Structural basis for CCR6 modulation by allosteric antagonists

David Jonathan Wasilko[1], Brian S. Gerstenberger[2], Kathleen A. Farley[1], Wei Li[3], Jennifer Alley[3], Mark E. Schnute[2], Ray J. Unwalla[2], Jorge Victorino[1], Kimberly K. Crouse[3], Ru Ding[3], Parag V. Sahasrabudhe[1], Fabien Vincent[1], Richard K. Frisbie[1], Alpay Dermenci[4], Andrew Flick[4], Chulho Choi[4], Gary Chinigo[4], James J. Mousseau[4], John I. Trujillo[4], Philippe Nuhant[2], Prolay Mondal[4], Vincent Lombardo[4], Daniel Lamb[5,6], Barbara J. Hogan[5,6], Gurdeep Singh Minhas[5], Elena Segala[5], Christine Oswald[5], Ian W. Windsor[1], Seungil Han[1], Mathieu Rappas[5,6], Robert M. Cooke[5], Matthew F. Calabrese[1], Gabriel Berstein[3], Atli Thorarensen[2], and Huixian Wu[1]*

[1]Discovery Sciences, Medicine Design, Pfizer Inc., Groton, CT 06340, USA

[2]Medicine Design, Pfizer Inc., Cambridge, MA 02139, USA

[3]Inflammation and Immunology Research, Pfizer Inc., Cambridge, MA 02139, USA

[4]Medicine Design, Pfizer Inc., Groton, CT 06340, USA

[5]Sosei Heptares, Steinmetz Building, Granta Park, Great Abington, Cambridge CB21 6DG, United Kingdom

[6]Current address: Nxera Pharma UK Limited, Steinmetz Building, Granta Park, Great Abington, Cambridge CB21 6DG, United Kingdom

*Correspondence to: huixian.wu@pfizer.com

Abstract

The CC chemokine receptor 6 (CCR6) is a potential target for chronic inflammatory diseases such as psoriasis and inflammatory bowel disease. Previously, we reported an active CCR6 structure in complex with its cognate chemokine CCL20, revealing the molecular basis of CCR6 activation mediated by CCL20. Here, we present two inactive CCR6 structures determined by cryo-EM in ternary complexes with different allosteric antagonists, CCR6/SQA1/OXM1 and CCR6/SQA1/OXM2. OXM1 and OXM2 are oxomorpholine (OXM) analogues which are highly selective for CCR6 and disrupt the molecular network critical for receptor activation by binding to an extracellular allosteric pocket within the transmembrane domain. A U-shaped conformation stabilized by intramolecular interactions was revealed by structural and NMR studies of active OXM analogues. SQA1 is a squaramide (SQA) derivative with close-in analogues that were previously reported to be antagonists of CCR6 and other chemokine receptors. Our structures reveal an intracellular pocket occupied by SQA1 that overlaps with the G protein binding site. In addition, SQA1 stabilizes a closed conformation of the intracellular pocket, a hallmark of the inactive state of GPCRs. Minimal communication was found between the two allosteric pockets. Overall, our work provides new evidence of the versatility of GPCR antagonism by small molecules, complementing previous knowledge on CCR6 activation, and sheds light on drug discovery approaches to target CCR6 for autoimmune disorders.

Introduction

The chemokine receptors (CKRs) are a group of class A G protein-coupled receptors (GPCRs) that respond to a family of cytokines known as chemokines[1]. More than twenty distinct CKRs have been discovered in humans, including 18 conventional CKRs that are typically G_i-coupled and 5 atypical CKRs which are structurally related to the conventional receptors but are non-signaling[2]. CKRs can also be classified by their primary type of chemokine ligands as CC, CXC, CX_3C, and C receptors, where C refers to the conserved N-terminal cysteines and X represents the amino acids between the two N-terminal cysteines of the chemokine[1]. Upon activation by chemokines, CKRs mediate a wide range of physiological processes including embryonic development, immune cell proliferation, activation, differentiation, and cell death. Consequently, CKRs play a significant role in the development and homeostasis of the immune system, and are important drug targets for cancer, infectious diseases, inflammation, and autoimmune disorders[3,4].

The CC chemokine receptor 6 (CCR6)[5] is a member of the family expressed on a variety of lymphocytes including T cells, particularly the T helper 17 (Th17) cells[6], B cells, and dendritic cells. CCR6 is activated by its specific ligand CCL20[7], which stimulates the migration of CCR6[+] lymphocytes to sites of inflammation. The expression of both CCL20 and CCR6 is upregulated in the epithelial tissues of patients with chronic inflammatory conditions such as psoriasis[8] and inflammatory bowel disease (IBD)[9]. Therefore, inhibiting CCL20-mediated CCR6 signaling is a promising strategy for anti-inflammatory drug discovery[10]. Indeed, a monoclonal antibody targeting CCL20 has resulted in the effective alleviation of inflammation in preclinical models[11]. In addition, CCR6 knockout mice are protected from induced psoriasis-like inflammation[12]. Accordingly, a small molecule antagonist of CCR6 could be useful as an oral treatment for autoimmune disorders.

Previously, we reported an active CCR6 structure in complex with CCL20 and G protein[13].

This structure revealed a shallow CCL20 pocket formed mainly by extracellular loops and the N-

terminus of CCR6, leaving the orthosteric pocket within the seven-transmembrane (7TM) domain

largely unoccupied. The CCL20-CCR6 recognition mode as well as the large extracellular pocket

of this protein receptor pose significant challenges to the development of small molecule

antagonists targeting the orthosteric site. Alternatively, modulating CKRs via allosteric ligands,

notable for its versatility and diversity of mechanisms-of-action (MoA), has been an attractive and

promising approach in CKR drug discovery[14-18].

In this work, we present two inactive CCR6 structures in ternary complexes with different

allosteric antagonists from squaramide (SQA) and oxomorpholine (OXM) series. The structures

were determined using single-particle cryogenic electron microscopy (cryo-EM) facilitated by an

engineered fiducial marker. The SQA and OXM analogues occupy two different pockets, binding

of which can occur simultaneously on CCR6. Different mechanisms to allosterically antagonize

CCR6 are revealed for SQA and OXM analogues, highlighting the diversity of GPCR allosteric

modulation by small molecules.

Results and Discussion

Additive stabilization of CCR6 by SQA and OXM

SQA1 (**Fig. 1a**) is a SQA derivative that was initially discovered as a CXCR2 inhibitor and later

found to also inhibit CCR6 activity[19]. Conversely, the OXM analogues OXM1 and OXM2 (**Fig.**

1a) were discovered in a ligand-based virtual screen using SQA1 as the seed and subsequent

structure-activity relationship (SAR) studies. All three small molecules demonstrate robust

inhibition of CCL20-mediated chemotaxis in primary human T cells expressing endogenous CCR6

(**Fig. 1b**). Significant CCR6 stabilization was observed in a thermal shift assay (**Fig. 1c**) upon

incubation with the small molecules, confirming direct target engagement for the SQA and OXM

molecules. Surprisingly, the stabilization of CCR6 by analogues from the two series is additive, as

a significantly larger thermal shift was observed when analogues from both series were added to

the receptor. Such additive stabilization was not dependent on the order of ligand addition between

SQA1 and OXM analogues but was not observed when CCR6 was co-incubated with the two

different OXM molecules. These results suggest that despite the discovery of the OXM series via

a SQA-based virtual screen, the two series bind to different pockets on CCR6 and can bind

independently and simultaneously to the receptor.

Construct engineering for cryo-EM studies

To reveal how the SQA and OXM analogues bind to CCR6, we conducted cryo-EM studies using

an engineered CCR6 construct Nα7.1 which contains seven thermostabilizing alanine-

substitutions[20,21] in the transmembrane domain and a point mutation L134$^{3.41}$W (superscripts

indicate Ballesteros–Weinstein numbering for GPCRs[22]) to increase receptor expression[23].

Binding of the SQA and OXM analogues to the thermostabilized CCR6 construct was confirmed

with comparable potency to the wild-type (WT) receptor (**Fig. S1**). Additive stabilization between

the SQA and OXM analogues was also observed with the stabilized construct (**Fig. 1d**). For

structural studies, twenty-seven residues at both N and C-termini of CCR6 were truncated to

improve protein homogeneity. Furthermore, the residues between L241 and K251 of intracellular

loop 3 (ICL3) were replaced by a thermostabilized apocytochrome b562 (BRIL)[24]. Two amino

acid linkers were grafted from the A$_{2A}$ adenosine receptor[25] to stabilize continuous alpha-helices

as the junctions between TM5 (CCR6) and helix 1 (BRIL), as well as TM6 (CCR6) and helix 4

(BRIL). The engineered CCR6-BRIL chimeric construct (**Fig. S2**) was co-purified with analogues

from both SQA and OXM series. An anti-BRIL Fab[26] with an anti-Fab nanobody (Nb)[27] was

employed to complex with the BRIL in the chimeric CCR6 construct. The BRIL/Fab/Nb module

was introduced as a fiducial marker in the single-particle cryo-EM analysis to facilitate particle

alignment[28]. This approach led to the successful determination of two inactive CCR6 structures

bound by ligands from the SQA and OXM series.

The complex structures of CCR6/SQA1/OXM1 and CCR6/SQA1/OXM2 were determined

at overall resolutions of 2.63 Å and 3.02 Å, respectively (**Figs. S3-4, Table S1**). Cryo-EM

reconstructions of both complexes reveal robust density, allowing the unambiguous fitting of

BRIL, Fab, Nb, and the 7TM of CCR6 (**Figs. 2a, b**). Most extracellular loops (ECLs) and

intracellular loops (ICLs) are unresolved or ambiguous in the reconstruction except for the bottom

of ECL2, which adopts a β-hairpin structure like that reported in the active CCR6 and other CKR

structures[13]. The disordered loops in the final reconstructions are not unexpected based on the

known intrinsic flexibility of these regions in GPCRs. In both structures, the densities of the SQA

and OXM analogues are robustly resolved, with each series binding to a distinct but well-defined

pocket within the 7TM bundle, allowing confident modeling of ligand binding for subsequent

structural analysis (**Figs. 2, S3-4**).

The OXM analogues bind to an extracellular allosteric pocket

U-shaped density is resolved in an extracellular pocket for both reconstructions and can be fitted

by the two different OXM analogues, OXM1 and OXM2 (**Figs. 2a, b**). Binding of the OXM

analogues to CCR6 is mainly mediated by hydrophobic interactions with side chains from TMs 3,

4, 5, 6, and ECL2 (**Figs. 2c, 3a-d**). In particular, the *p*-chlorophenyl group of OXM1 and the *p*-

trifluoromethylphenyl moiety of OXM2 are sandwiched between TMs 5 and 6 with their *para*-

substitutions pointing towards the hydrophobic membrane, while making hydrophobic interactions

with F129$^{3.36}$, L219$^{5.43}$, F223$^{5.47}$, I268$^{6.49}$, M272$^{6.53}$, and L275$^{6.56}$. On the other end of the bent

molecules are hydrophobic substitutions on the amide nitrogen of the oxomorpholine core, i.e., an

indane moiety in OXM1 and a bicyclo[1.1.1]pentane in OXM2. These two groups bind in a

hydrophobic pocket capped by the C-terminal tip of ECL2 that connects to TM5. Due to the

compact size of this subpocket, the larger indane group of OXM1 bends down towards the *p*-

chlorophenyl group, forming an intramolecular T-shaped π stacking interaction which likely

further stabilizes the observed U-shaped conformation of OXM1. Other than the hydrophobic

interactions, several H-bonds are mediated by the two amide groups of the OXM analogues with

polar side chains such as N130$^{3.37}$, K122$^{3.29}$, T182$^{4.60}$, and N271$^{6.52}$. Remarkably, a majority of the

OXM binding pocket side chains are CCR6-specific (**Fig. S5**). Consequently, OXM1 was found

to be highly selective for CCR6 and is not active against other CKRs (**Table S2**).

Conformational restraints required for OXM activity

The observed bent pose of OXM1 in the ligand-bound structure prompted us to study the solution

conformation of this molecule using Nuclear Magnetic Resonance (NMR) experiments to

understand the energy landscape of this conformation. Rotating-frame nuclear Overhauser

Enhancement signals (ROEs) were collected using previously reported methods[29]. Robust ROEs

were obtained between the indane C(sp^3)-*H* (H-18) and the aromatic hydrogens from the *p*-

chlorophenyl ring (Hs-3,4,6,7) (**Fig. S6**), indicating intramolecular interactions between the indane

and the *p*-chlorophenyl moieties are also maintained in solution. Additionally, we performed

proton and carbon assignments for OXM1 (**Figs. S7-13**, **Table S3**) in a compressed dimethyl

sulfoxide (DMSO) gel[30,31] which were used to calculate the residual dipolar coupling (RDC) values

of the proton-carbon and proton-nitrogen J-couplings. The RDC data was fit to a conformational

ensemble that consisted of low energy structures generated by molecular dynamics simulations in

a solvent-dependent manner (see methods). This exercise output several U-shaped conformations

of OXM1 as the best fits. Ninety-two percent of the low energy conformers greatly resemble the

bound pose observed in the cryo-EM structure (**Fig. 3e**), suggesting a low energy conformation

for the bound state.

157   We rationalize the conformational restraints conferred by the cyclopropyl group as well as

the T-shaped π stacking interactions between the indane and the *p*-chlorophenyl groups are critical

factors in stabilizing the observed U-shaped conformation of OXM1 in both bound and free forms.

To test this hypothesis, we synthesized and analyzed two additional OXM analogues, OXM3

which bears a *gem*-dimethyl replacement at the cyclopropyl position, and OXM4 that lacks the

para-Cl substitution on the phenyl ring (**Fig. 3e**). As expected, extended conformations in solution

are observed for the two new OXM analogues and no long-range ROEs or nuclear overhauser

enhancement signals (NOEs) were observed (**Figs. 3e, S14-27, Table S3**). Consequently, neither

OXM3 nor OXM4 are active on CCR6. Finally, we also characterized the solution conformation

of OXM2 used in the cryo-EM studies. Despite the absence of the indane moiety, a curved

conformation that largely resembles the bound conformation was revealed in its solution state (**Fig.**

3e, S28-34, Table S3). The curved conformation of OXM2 in solution was further confirmed by

long-range NOEs observed between the bicyclo[1.1.1]pentane (Hs-21,24,25) and the aromatic

hydrogens from the *p*-trifluoromethyl phenyl ring (Hs-16,17,19,20) (**Fig. S35**). In sum, the NMR

studies support the hypothesis that a low-energy curved conformation in the free state is critical

for OXM analogues to be active on CCR6.

SQA1 binds to an intracellular pocket

In contrast to the OXM analogues, an intracellular binding site is revealed for the SQA molecule

SQA1 (**Figs. 2d, 4**). The binding poses of SQA1 from both structures are nearly identical.

Therefore, the CCR6/SQA1/OXM1 complex that led to a higher-resolution reconstruction was

used for analysis hereafter. Side chains interacting with SQA1 are from TMs 1, 2, 3, 6, 7, and helix

8 (**Figs. 4a, b**). CCR6 has a glycine residue $G320^{8.47}$ as the TM7-helix 8 linker. The absence of a

side chain at the 8.47 position provides a pocket that allows the squaramide core to bind. Indeed,

a CCR6 mutant bearing $G320^{8.47}V$ completely abolished SQA1 binding (**Figs. 4c, d**). Extensive

polar interactions are observed between the squaramide-picolinamide core of SQA1 and CCR6,

including the salt bridges between $D82^{2.40}$ and the squaramide NHs, as well as H-bonds mediated

by side chains of $S79^{2.37}$, $T81^{2.39}$, $R143^{3.50}$, and $K322^{8.49}$ with the 3'-hydroxyl-picolinamide moiety.

Consequently, SQA1 binding was abolished on mutants bearing single point mutations such as

$D82^{2.40}N$, $S79^{2.37}A$, $T81^{2.39}L$, $R143^{3.50}A$, or $K322^{8.49}A$ (**Figs. 4c, d**). On the other hand, binding of

the 1-(4-isopropyl furan-2-yl)propyl moiety is largely mediated by hydrophobic interactions with

side chains from TMs 1, 2, 7, and helix 8. Particularly, the 1-propyl binds in a hydrophobic pocket

formed by the side chains of $L85^{2.43}$, $Y316^{7.53}$, and $V67^{1.53}$. CCR6 mutants bearing mutations in

this pocket such as $L85^{2.43}A$ or $Y316^{7.53}A$ no longer respond to SQA1 (**Figs. 4c, d**).

A co-crystal structure of CXCR2 in complex with SQA1 was reported previously[32].

Comparison of the bound structures of SQA1 on CCR6 and CXCR2 reveal a highly similar mode

adopted by the molecule in binding to a mostly conserved pocket shared by the two receptors (**Figs.**

4e, S36). As observed in CCR6, extensive polar interactions are also seen in the CXCR2 structure,

mediated by the squaramide-picolinamide core with conserved side chains shared by the two

receptors. The most striking difference in the ligand binding pose is for the 1-(4-isopropyl furan-

2-yl)propyl moiety (**Fig. 4f**). In CXCR2, this moiety is pushed about 1 Å downward towards the

intracellular side compared to the bound position in CCR6. Movement of the 1-(4-isopropyl furan-

2-yl)propyl moiety is likely caused by the difference of side chains at the 1.57 position, which is

$F71^{1.57}$ in CCR6 and is replaced by $I73^{1.57}$ in CXCR2. The aliphatic isoleucine side chain reaches

deeper into the 1-propyl binding pocket, resulting in a smaller subpocket in CXCR2 that pushes

the 1-propyl group down. Furthermore, to avoid clashing into the isoleucine side chain, the 4-

isopropyl furan of SQA1 in CXCR2 undergoes a rotation away from TM1 compared to the bound

structure in CCR6.

SQA and OXM antagonize CCR6 via different allosteric mechanisms

The bound CCR6 structures are resolved in inactive conformations (**Fig. 5a**). A comparison of the

SQA- and OXM-bound CCR6 structures to the previous CCL20-bound structure reveals that the

binding sites of the OXM and SQA analogues do not overlap with the CCL20 binding site (**Fig.**

5b), indicating both OXM and SQA molecules are allosteric antagonists of CCR6. Compared to

the active structure, a significant rearrangement of the CCR6 7TM bundle is observed in the

inactive structures, which includes a 4-5 Å inward movement of the entire TM6 towards the center

of the TM bundle and a 4.6-Å outward movement at the IC tip of TM7 (**Figs. 5c, d**). These changes

result in a closed IC pocket that is not compatible with G protein coupling. Furthermore, the

conserved functional motifs such as $P226^{5.50}$-$M133^{3.40}$-$F263^{6.44}$, $D142^{3.49}$-$R143^{3.50}$-$Y144^{3.51}$, and

$N312^{7.49}$-$P313^{7.50}$-x-x-$Y316^{7.53}$ are now resolved in conformations resembling those observed in

most other inactive class A GPCR structures (**Figs. 5e-g**)[33]. In the inactive structures of CCR7[18]

and CCR9[16] reported previously, a conserved $N^{6.52}$-$Y^{3.32}$-$Q^{6.48}$ H-bond network was observed as a

hallmark of the inactive state[13]. In the inactive CCR6, a similar H-bond network mediated by

$Y125^{3.32}$, $Q267^{6.48}$, and $N271^{6.52}$ is also observed (**Fig. 5h**). Notably, both $Y125^{3.32}$ and $N271^{6.52}$

are residues located in the OXM binding pocket, and the side chains of these residues mediate

specific protein-ligand interactions via hydrophobic packing and a H-bond interaction with the

oxomorpholine core, respectively. Therefore, binding of OXM analogues likely further stabilizes

the $Y125^{3.32}$-$Q267^{6.48}$-$N271^{6.52}$ H-bond network that is important for the inactive state of CCR6.

Bound to an inactive IC pocket, SQA1 conformationally and sterically hinders the coupling

of the G protein (**Fig. 5b**). Furthermore, SQA1 engages the side chains of R143$^{3.50}$ and Y316$^{7.53}$,

which are part of the D$^{3.49}$R$^{3.50}$Y$^{3.51}$ and the N$^{7.49}$P$^{7.50}$xxY$^{7.53}$ motifs involved in G protein binding

and CCR6 activation. In the SQA1 co-structures, R143$^{3.50}$ and Y316$^{7.53}$ are found in inactive

conformations (**Figs. 5f, g**). Therefore, SQA1 binding spatially competes with G protein binding

and prevents conformational changes of the TMs and the critical activation motifs, which

altogether inhibits CCL20-mediated CCR6 activation. In addition to SQA1, a few other

intracellular antagonists have been reported for other CKRs, such as CXCR1/2[15,34], CCR2[17],

CCR7[18], and CCR9[16], likely all utilizing a highly similar molecular mechanism to regulate their

corresponding receptors.

Located in the extracellular half of the 7TM, the OXM pocket is about 9 Å away from the

CCL20 binding site (**Fig. 5g**). Binding of the OXM analogues, however, reshapes the orthosteric

pocket of CCR6 to a conformation that is vastly different from that observed in the CCL20-bound

form (**Figs. 5b, c**). Particularly, by interacting with N186 from ECL2 and the C-terminal tip of the

loop, OXM1 or OXM2 binding brings the β-hairpin of ECL2 4 Å closer to the center of the EC

pocket, a conformation that would clash with the N-terminus of CCL20 resolved in the active

structure. Moreover, sandwiched in between TMs 5 and 6, the *para*-substituted phenyl groups of

OXM1 and OXM2 wedge TM5 8 Å away from the center of the 7TM bundle, which makes room

to accommodate the 4-5 Å inward movement of the entire TM6 associated with receptor

inactivation. ECL3 provides an important docking site for the N-loop and the 30s-loop of CCL20

during receptor activation. The large movement of TM6 observed in the OXM analogue-bound

structures results in an EC tip conformation of the helix that is now no longer compatible with

CCL20 docking. Altogether, OXM binding stabilizes a distinct receptor conformation of the EC

region from that utilized by CCL20, thereby allosterically inhibiting CCL20-mediated CCR6

activation.

Positive cooperativity between orthosteric agonists and G protein is a hallmark of GPCR

activation. Cooperative stabilization of inactive CCR2 between orthosteric and allosteric

antagonists has also been reported[17]. However, crosstalk between different allosteric ligands has

not been well characterized to date. To understand the communication between the extra- and

intracellular allosteric pockets discovered on CCR6, we examined binding of [^{3}H]SQA1 to WT

CCR6 with and without OXM1. To our surprise, the presence of OXM1 did not enhance the

binding of [^{3}H]SQA1 (Fig. S37), in contrast to the report that the presence of an orthosteric CCR2

antagonist increased binding of the intracellular antagonist, CCR2-RA[17]. We also studied the

impact of SQA1 on the binding of OXM1 using SPR with a stabilized CCR6. Despite a slight

increase in on-rate and a slight decrease in off-rate of OXM1 to CCR6 preincubated with SQA1,

the overall change of the dissociation constant (K_D) was within noise (Figs. S38). Taken together,

these binding data suggest minimal communication occurred between the two allosteric pockets

occupied by the OXM and SQA molecules, differing from the cooperativity observed between

orthosteric agonists and intracellular signaling transducers.

Discussion

Despite the significant advancement of cryo-EM and its tremendous success in active GPCR

structure determination[35], limited examples have been reported on the application of cryo-EM to

small molecule-bound inactive GPCR structure determination. The challenges are twofold: the

small size of the receptors alone, and the highly dynamic nature of this protein family. Here we

present a successful example of the application of cryo-EM in tackling inactive GPCR structure

determination for small molecule drug discovery. A BRIL fusion was introduced to replace the

ICL3 with a carefully designed junction to achieve optimal rigidity. The BRIL/Fab/Nb module is

proven to be an effective fiducial marker that allows robust particle alignment in cryo-EM[28].

Thermostabilizing mutations together with a ternary complex strategy of incorporating two

different allosteric antagonists led to a highly stabilized CCR6. As a result, two high-resolution

structures in complex with analogues from the SQA and OXM series were determined in this

study, providing the first known examples of inactive CCR6 structures bound by small molecules.

The structures reveal that SQA and OXM utilize different allosteric mechanisms to inhibit CCR6

activation. Importantly, the allosteric pockets of CCR6 discovered in our work are also likely

present in other CKRs. Therefore, the lessons described herein are useful for structure-based drug

discovery of small molecules targeting CCR6 and other related CKRs.

Figures

Figure 1. SQA and OXM analogues inhibit CCL20-mediated activation of CCR6 and display

additive binding stabilization to the receptor. a, Chemical structures of SQA1, OXM1 and

OXM2. **b**, Inhibition of CCL20-mediated CCR6$^+$ human T-cell chemotaxis demonstrated by

SQA1 (IC$_{50}$ = 199 nM or pIC$_{50}$ = 6.7 ± 0.8, mean ± s.d., n = 6), OXM1 (IC$_{50}$ = 146 nM, or pIC$_{50}$

= 6.8 ± 0.2, mean ± s.d., n = 4), and OXM2 (IC$_{50}$ = 163 nM, pIC$_{50}$ = 6.78 ± 0.11, mean ± s.d., n =

2). **c-d**, Thermal shifts of **c**, WT (n = 4) and **d**, thermostabilized (n = 6) CCR6_Nα7.1 from Apo

to liganded conditions as indicated. Statistical analysis was done by unpaired t test (*ns*, not

significant; ** p < 0.01; *** p < 0.001; **** p < 0.0001). Data for graphs **b-d** are available as

source data.

Figure 2. Overall structures of CCR6 in complex with OXM and SQA analogues. a-b, Cryo-

EM map (left) and model (right) of CCR6 complex with anti-BRIL Fab and anti-Fab Nb bound to

a, OXM1 (orange carbon spheres) and SQA1 (yellow carbon spheres), or **b**, OXM2 (slate carbon

spheres) and SQA1 (yellow carbon spheres). Protein is colored by subunit as follows: Fab heavy

chain, magenta; Fab light chain, salmon; Nb, yellow; CCR6, aquamarine in **a** and deep teal in **b**.

c, Extracellular (EC) view of CCR6 bound to OXM analogues, OXM1 (left) and OXM2 (right).

d, Intracellular (IC) view of CCR6 bound to the SQA analogue, SQA1. Both **c** and **d** use the same

color code as **a** and **b**. **e**, SQA1 and OXM1 fit into the cryo-EM density map. **f**, EM density for

SQA1 and OXM2 fit into the cryo-EM density map.

Figure 3. Structures of OXM analogues bound to CCR6 and comparison to solution conformations. a, Binding pocket of OXM1 (orange carbon sticks). **b**, Protein-ligand interaction

diagram for OXM1 prepared with Molecular Operating Environment (MOE)[36]. **c**, Binding pocket

of OXM2 (slate carbon sticks). **d**, Protein-ligand interaction diagram for OXM2 prepared with

MOE[36]. In **a** and **c**, CCR6 protein is shown in aquamarine and deep teal ribbons, respectively. Side

chains interacting with the small molecules are shown as white carbon sticks. Hydrogen bonds are

highlighted by yellow dashed lines. In **b** and **d**, OXM pocket residues are presented as follows:

polar residues in pink, hydrophobic residues in green, acidic residues with a red contour ring, basic

residues with a blue contour ring. Green dotted arrows indicate hydrogen and halogen bonds

mediated by side chains that contribute to ligand binding. Ligand atoms exposed to environment

are shaded in blue according to degrees of exposure, scaled by size. Light-blue halos around

residues indicate the degree of interaction with ligand, scaled by size. The dotted contour around

the ligand reflects steric room for methyl substitution. **e**, Chemical structures of different OXM

analogues and their solution conformations were determined by NMR using residual dipolar

couplings (RDC)[30]. Three most populated conformations are illustrated by dark green, light green,

and orange carbon sticks, respectively. The bound conformations of OXM1 and OXM2 are shown

as purple carbon sticks in the overlays for comparison. The IC_{50} value of each OXM analogue

determined in the CCL20-mediated human CCR6[+] T-cell chemotaxis assay is shown under the

corresponding chemical structure.

Figure 4. SQA1 binds to an intracellular pocket. a, Structure of SQA1 (yellow carbon sticks) bound to CCR6 (aquamarine ribbon) from the higher-resolution CCR6/SQA1/OXM1 cryo-EM structure. Side chains interacting with SQA1 are shown by white carbon sticks. Hydrogen bonds are highlighted by yellow dashed lines. **b**, Protein-ligand interaction diagram for SQA1 prepared

with MOE[36]. Definitions of the schematic representation are same as that in Fig. 3b and d. **c**,

Saturation binding curves of [^{3}H]-SQA1 on CCR6 WT and mutants as indicated. Binding of [^{3}H]-

SQA1 on WT provided a K_D of 250 ± 22 nM. Data shown as mean ± s.d. are representative of

three independent experiments ($n = 3$). No significant binding was observed at all doses measured

on the CCR6 mutants. **d**, Normalized surface expression (dark grey bars) and CPM (counts per

minute) of [^{3}H]-SQA1 measured at 1,000 nM ligand concentration (blue bars) for CCR6 mutants

transiently expressed in HEK293 cells. Data of WT was used as the baseline for normalization

calculation. **e**, Overlay of CCR6 and CXCR2 (PDB ID 6LFL) inactive structures bound by SQA1.

f, Major differences of the binding pockets in CCR6/- and CXCR2/SQA1-bound structures. Side

chains of pocket residues are shown in sticks. In **e** and **f**, CXCR2 and SQA1 from 6LFL are shown

by dark grey ribbon and dark grey sticks. CCR6 structure is shown by the same cartoon

representations as in **a**. Data for graphs **c-d** are available as source data.

Figure 5. Different mechanisms of allosteric antagonism of CCR6 by OXM and SQA. a, Overall structures of the active CCR6 (magenta, PDB ID 6WWZ, left) in complex with its cognate chemokine ligand CCL20 (yellow) and a heterotrimeric G protein colored by subunit (Gαo, green, Gβ, cyan, and Gγ, light grey), and the inactive CCR6 (aquamarine, right) in complex with SQA1 (yellow carbon spheres) and OXM1 (orange carbon spheres). The light grey box highlights the location of cell membranes. **b**, Overlays of CCL20, OXM1, and SQA1 bound structures onto the

active (left) and the inactive (right) CCR6 structures. Structures are presented in the same color

codes as in **a**. Blue lines depict pockets revealed by the structures. **c-d**, Structural overlays of the

active (magenta) and inactive (aquamarine) CCR6 structures from **c**, EC and **d**, IC views. **e-h**,

Conformational changes of conserved functional motifs, **e**, P226$^{5.50}$-M133$^{3.40}$-F263$^{6.44}$, **f**, D142$^{3.49}$-

R143$^{3.50}$-Y144$^{3.51}$, **g**, N312$^{7.49}$-P313$^{7.50}$-x-x-Y316$^{7.53}$, and **h**, Y125$^{3.32}$-Q267$^{6.48}$-N271$^{6.52}$ revealed

by structural overlay of active (magenta) and inactive (aquamarine) CCR6. In **c-h**, significant TM

movements and side chain conformational changes from active to inactive states are highlighted

by black arrows.

References

References

1   Hughes, C. E. & Nibbs, R. J. B. A guide to chemokines and their receptors. *FEBS J* **285**,
2944-2971 (2018). https://doi.org:10.1111/febs.14466
2   Bonecchi, R. & Graham, G. J. Atypical Chemokine Receptors and Their Roles in the
Resolution of the Inflammatory Response. *Front Immunol* **7**, 224 (2016).
https://doi.org:10.3389/fimmu.2016.00224
3   Murdoch, C. & Finn, A. Chemokine receptors and their role in inflammation and
infectious diseases. *Blood* **95**, 3032-3043 (2000).
4   Chow, M. T. & Luster, A. D. Chemokines in cancer. *Cancer Immunol Res* **2**, 1125-1131
(2014). https://doi.org:10.1158/2326-6066.CIR-14-0160
5   Murphy, P. M. *et al.* International union of pharmacology. XXII. Nomenclature for
chemokine receptors. *Pharmacol Rev* **52**, 145-176 (2000).
6   Paulissen, S. M., van Hamburg, J. P., Dankers, W. & Lubberts, E. The role and
modulation of CCR6+ Th17 cell populations in rheumatoid arthritis. *Cytokine* **74**, 43-
53 (2015). https://doi.org:10.1016/j.cyto.2015.02.002
7   Schutyser, E., Struyf, S. & Van Damme, J. The CC chemokine CCL20 and its receptor
CCR6. *Cytokine Growth Factor Rev* **14**, 409-426 (2003).
8   Homey, B. *et al.* Up-regulation of macrophage inflammatory protein-3 alpha/CCL20
and CC chemokine receptor 6 in psoriasis. *J Immunol* **164**, 6621-6632 (2000).
https://doi.org:10.4049/jimmunol.164.12.6621
9   Skovdahl, H. K. *et al.* Expression of CCL20 and Its Corresponding Receptor CCR6 Is
Enhanced in Active Inflammatory Bowel Disease, and TLR3 Mediates CCL20
Expression in Colonic Epithelial Cells. *PLoS One* **10**, e0141710 (2015).
https://doi.org:10.1371/journal.pone.0141710
10   Ranasinghe, R. & Eri, R. Modulation of the CCR6-CCL20 Axis: A Potential Therapeutic
Target in Inflammation and Cancer. *Medicina (Kaunas)* **54** (2018).
https://doi.org:10.3390/medicina54050088
11   Katchar, K., Kelly, C. P., Keates, S., O'Brien M, J. & Keates, A. C. MIP-3alpha
neutralizing monoclonal antibody protects against TNBS-induced colonic injury and
inflammation in mice. *Am J Physiol Gastrointest Liver Physiol* **292**, G1263-1271
(2007). https://doi.org:10.1152/ajpgi.00409.2006
12   Hedrick, M. N. *et al.* CCR6 is required for IL-23-induced psoriasis-like inflammation
in mice. *J Clin Invest* **119**, 2317-2329 (2009). https://doi.org:10.1172/jci37378
13   Wasilko, D. J. *et al.* Structural basis for chemokine receptor CCR6 activation by the
endogenous protein ligand CCL20. *Nat Commun* **11**, 3031 (2020).
https://doi.org:10.1038/s41467-020-16820-6
14   Rennard, S. I. *et al.* CXCR2 Antagonist MK-7123. A Phase 2 Proof-of-Concept Trial for
Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med* **191**, 1001-1011
(2015). https://doi.org:10.1164/rccm.201405-0992OC
15   Gonsiorek, W. *et al.* Pharmacological characterization of Sch527123, a potent
allosteric CXCR1/CXCR2 antagonist. *J Pharmacol Exp Ther* **322**, 477-485 (2007).
https://doi.org:10.1124/jpet.106.118927
16   Oswald, C. *et al.* Intracellular allosteric antagonism of the CCR9 receptor. *Nature*
540, 462-465 (2016). https://doi.org:10.1038/nature20606

17   Zheng, Y. *et al.* Structure of CC chemokine receptor 2 with orthosteric and allosteric
antagonists. *Nature* **540**, 458-461 (2016). https://doi.org:10.1038/nature20605
18   Jaeger, K. *et al.* Structural Basis for Allosteric Ligand Recognition in the Human CC
Chemokine Receptor 7. *Cell* **178**, 1222-1230 e1210 (2019).
https://doi.org:10.1016/j.cell.2019.07.028
19   Li, W. *et al.* A Novel C-C Chemoattractant Cytokine (Chemokine) Receptor 6 (CCR6)
Antagonist (PF-07054894) Distinguishes between Homologous Chemokine
Receptors, Increases Basal Circulating CCR6(+) T Cells, and Ameliorates Interleukin-
23-Induced Skin Inflammation. *J Pharmacol Exp Ther* **386**, 80-92 (2023).
https://doi.org:10.1124/jpet.122.001452
20   Magnani, F. *et al.* A mutagenesis and screening strategy to generate optimally
thermostabilized membrane proteins for structural studies. *Nat Protoc* **11**, 1554-
1571 (2016). https://doi.org:10.1038/nprot.2016.088
21   Robertson, N. *et al.* The properties of thermostabilised G protein-coupled receptors
(StaRs) and their use in drug discovery. *Neuropharmacology* **60**, 36-44 (2011).
https://doi.org:10.1016/j.neuropharm.2010.07.001
22   Ballesteros, J. A. & Weinstein, H. Integrated methods for the construction of three-
dimensional models and computational probing of structure-function relations in G
protein-coupled receptors. *Methods in Neurosciences* **25**, 63 (1995).
https://doi.org:https://doi.org/10.1016/S1043-9471(05)80049-7
23   Roth, C. B., Hanson, M. A. & Stevens, R. C. Stabilization of the human beta2-
adrenergic receptor TM4-TM3-TM5 helix interface by mutagenesis of Glu122(3.41),
a critical residue in GPCR structure. *J Mol Biol* **376**, 1305-1319 (2008).
https://doi.org:10.1016/j.jmb.2007.12.028
24   Chun, E. *et al.* Fusion partner toolchest for the stabilization and crystallization of G
protein-coupled receptors. *Structure* **20**, 967-976 (2012).
https://doi.org:10.1016/j.str.2012.04.010
25   Liu, W. *et al.* Structural basis for allosteric regulation of GPCRs by sodium ions.
Science **337**, 232-236 (2012). https://doi.org:10.1126/science.1219218
26   Mukherjee, S. *et al.* Synthetic antibodies against BRIL as universal fiducial marks for
single-particle cryoEM structure determination of membrane proteins. *Nat Commun*
11, 1598 (2020). https://doi.org:10.1038/s41467-020-15363-0
27   Ereno-Orbea, J. *et al.* Structural Basis of Enhanced Crystallizability Induced by a
Molecular Chaperone for Antibody Antigen-Binding Fragments. *J Mol Biol* **430**, 322-
336 (2018). https://doi.org:10.1016/j.jmb.2017.12.010
28   Tsutsumi, N. *et al.* Structure of human Frizzled5 by fiducial-assisted cryo-EM
supports a heterodimeric mechanism of canonical Wnt signaling. *Elife* **9** (2020).
https://doi.org:10.7554/eLife.58464
29   Butts, C. P. *et al.* Interproton distance determinations by NOE--surprising accuracy
and precision in a rigid organic molecule. *Org Biomol Chem* **9**, 177-184 (2011).
https://doi.org:10.1039/c0ob00479k
30   Farley, K. A. *et al.* Cyclic Peptide Design Guided by Residual Dipolar Couplings, J-
Couplings, and Intramolecular Hydrogen Bond Analysis. *J Org Chem* **84**, 4803-4813
(2019). https://doi.org:10.1021/acs.joc.8b02811

31 Horst, R., Farley, K. A., Kormos, B. L. & Withka, J. M. NMR spectroscopy: the swiss
army knife of drug discovery. *J Biomol NMR* **74**, 509-519 (2020).
https://doi.org:10.1007/s10858-020-00330-0
32 Liu, K. *et al.* Structural basis of CXC chemokine receptor 2 activation and signalling.
Nature (2020). https://doi.org:10.1038/s41586-020-2492-5
33 Wu, H. *et al.* Structure of the human kappa-opioid receptor in complex with JDTic.
Nature **485**, 327-332 (2012). https://doi.org:10.1038/nature10939
34 Salchow, K. *et al.* A common intracellular allosteric binding site for antagonists of
the CXCR2 receptor. *Br J Pharmacol* **159**, 1429-1439 (2010).
https://doi.org:10.1111/j.1476-5381.2009.00623.x
35 Danev, R. *et al.* Routine sub-2.5 A cryo-EM structure determination of GPCRs. *Nat*
Commun **12**, 4333 (2021). https://doi.org:10.1038/s41467-021-24650-3
36 Molecular Operating Environment (MOE), 2022.02 Chemical Computing Group ULC,
910-1010 Sherbrooke St. W., Montreal, QC H3A 2R7, Canada. (2023).
37 Jumper, J. & Hassabis, D. Protein structure predictions to atomic accuracy with
AlphaFold. *Nat Methods* **19**, 11-12 (2022). https://doi.org:10.1038/s41592-021-
01362-6
38 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501 (2010).
https://doi.org:10.1107/S0907444910007493
39 Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons
and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861-
877 (2019). https://doi.org:10.1107/S2059798319011471
40 Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
https://doi.org:10.1107/S0907444909042073
41 Schrödinger, L. & DeLano, W. *PyMoL*, <http://www.pymol.org/pymol> (2020).
42 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research
and analysis. *J Comput Chem* **25**, 1605-1612 (2004).
https://doi.org:10.1002/jcc.20084
43 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82 (2021).
https://doi.org:10.1002/pro.3943
44 Alexandrov, A. I., Mileni, M., Chien, E. Y., Hanson, M. A. & Stevens, R. C. Microscale
fluorescent thermal stability assay for membrane proteins. *Structure* **16**, 351-359
(2008). https://doi.org:10.1016/j.str.2008.02.004
45 *Blaze, version, Cresset, Litlington, Cambridgeshire, UK;*, <https://www.cresset-
group.com/software/blaze/> (
46 Cheeseright, T. J., Mackey, M. D., Melville, J. L. & Vinter, J. G. FieldScreen: virtual
screening using molecular fields. Application to the DUD data set. *J Chem Inf Model*
48, 2108-2117 (2008). https://doi.org:10.1021/ci800110p
47 Cheeseright, T., Mackey, M., Rose, S. & Vinter, A. Molecular field extrema as
descriptors of biological activity: definition and validation. *J Chem Inf Model* **46**, 665-
676 (2006). https://doi.org:10.1021/ci050357s

48 Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. The Cambridge Structural
Database. *Acta Crystallogr B Struct Sci Cryst Eng Mater* **72**, 171-179 (2016).
https://doi.org:10.1107/S2052520616003954
49 Gil-Silva, L. F., Santamaria-Fernandez, R., Navarro-Vazquez, A. & Gil, R. R. Collection
of NMR Scalar and Residual Dipolar Couplings Using a Single Experiment. *Chemistry*
22, 472-476 (2016). https://doi.org:10.1002/chem.201503449
50 Navarro-Vazquez, A. MSpin-RDC. A program for the use of residual dipolar couplings
for structure elucidation of small molecules. *Magn Reson Chem* **50 Suppl 1**, S73-79
(2012). https://doi.org:10.1002/mrc.3905
51 Cornilescu, G., Delaglio, F. & Bax, A. Protein backbone angle restraints from
searching a database for chemical shift and sequence homology. *J Biomol NMR* **13**,
289-302 (1999). https://doi.org:10.1023/a:1008392405740
52 *Macromodel, Schrodinger, LLC, New York*, <http://www.schrodinger.com> (2011).
499

Methods

No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

CCR6 thermostabilization and characterization. HEK293T cells used for transient transfection for thermostabilization experiments were cultured in DMEM + GlutaMAX™-I with 4.5 g/L D-Glucose and Pyruvate (Gibco 31966-021), supplemented with 10 % (v/v) of fetal calf serum (FCS). Transient transfections were carried out using GeneJuice (Merck) in accordance with manufacturer's protocols. Full-length WT CCR6 used as a template for thermostabilization was cloned into pcDNA3.1-based plasmid vector to allow transient expression with C-terminal GFP-TEV-myc-His×10 tags. Thermostability of transiently expressed CCR6 constructs in HEK293T cells was assayed by solubilized [^{3}H]-SQA1 binding. Cells were solubilized in detergent (1 % (w/v) *n*-Dodecyl-*β*-D-Maltopyranoside (DDM, Anatrace), *n*-Decyl-*β*-D-Maltopyranoside (DM, Anatrace), or 2 % (w/v) *n*-Nonyl-*β*-D-Glucopyranoside (NG, Anatrace) with cholesteryl hemisuccinate (CHS, Sigma) added at 0.012 % (w/v) as appropriate) prior to incubation with radioligand and heating, and unbound radioligand was subsequently removed using microscale immobilized metal affinity chromatography (IMAC). Melting temperature (T_m) was defined as the temperature at which 50 % of maximal radioligand binding was measured. Stabilizing mutants were made, identified, and combined as described previously[20,21]. The thermostabilized construct CCR6_Nα7.1 used in the cryo-EM study contains the following seven mutations: L47$^{1.33}$A, L86$^{2.44}$A, V96$^{2.54}$A, S140$^{3.47}$A, R159A, F236$^{5.60}$A, G295$^{7.32}$A (superscripts indicate Ballesteros–Weinstein numbering for GPCRs[22]). The thermostabilized construct CCR6_Nβ10.2 used in the

crosstalk SPR study contains F73$^{1.59}$A, V276$^{6.57}$A, and A279$^{6.60}$L in addition to the seven

mutations of Nα7.1.

The affinity of [^{3}H]-SQA1 binding to CCR6 constructs was determined by saturation

binding. Increasing concentrations of [^{3}H]-SQA1 were incubated for 2 h at 25 °C with aliquots of

HEK293T cells transiently expressing CCR6 (or mock transfected control) resuspended in 50 mM

HEPES pH 7.5, 150 mM NaCl, 5 mM MgCl$_2$ and protease inhibitors. For the crosstalk study, the

cells were prebound by 10 μM of OXM1 for 1 h before addition of [^{3}H]-SQA1. The cell suspension

with ligand mix was then chilled to 4 °C prior to receptor solubilization in 1 % DDM and 0.012 %

CHS, and unbound radioligand was separated using microscale IMAC. Non-specific binding to

mock-transfected controls was subtracted from binding to CCR6-containing samples prior to

calculation of K$_d$ in GraphPad Prism 9.0.0, using one site specific curve fitting.

Expression, membrane preparation, and protein purification. The thermostabilized

CCR6_Nα7.1-BRIL chimeric construct was cloned into a modified pFastBac1 vector (Invitrogen)

with a haemagglutinin (HA) signal sequence followed by a cleavable FLAG-10×His tags. Protein

expression was done in Spodoptera frugiperda (*Sf9*) insect cells (ATCC) using the Bac-to-Bac

Baculovirus Expression System (Invitrogen). The insect cells were infected at a density of 2-3 ×

10^6 cells/ml with high titer viral stock multiplicity of infection (MOI) of 5.0. At 48 h post infection,

cells were collected. Cell pellet was resuspended by dounce homogenization in buffer containing

10 mM HEPES pH 7.5, 150 mM NaCl, 10 mM MgCl$_2$, 20 mM KCl, nuclease (Pierce) and EDTA-

free cOmplete^{TM} protease inhibitor cocktail (Roche). Cell debris were removed by centrifugation

at 1000×g for 20 min. The supernatant was then subjected to ultracentrifugation at 158,420×g for

30 min and the membrane pellet was resuspended by dounce homogenization in 50 mM HEPES

pH 7.5, 150 mM NaCl, 10% glycerol, and EDTA-free cOmplete^{TM} protease inhibitor cocktail

(Roche). The resuspension was solubilized at 4 °C for 4 h with 2% (w/v) Lauryl Maltose Neopentyl

Glycol (LMNG, Anatrace) and 0.2% (w/v) CHS in the presence of 2 mg/ml iodoacetamide and 50

μM of the compounds. The sample was clarified by ultracentrifugation at 340,252×g for 30 min at

4 °C, and the supernatant was batch bound to TALON cobalt-affinity resin (Takara) overnight at

4 °C. The resin was washed with successively lower concentrations of detergent and eluted in

buffer containing 50 mM HEPES pH 7.5, 500 mM NaCl, 0.003% LMNG, 0.0003% CHS, 300 mM

imidazole, 50 μM OXM1 or OXM2, and 50 μM SQA1. The elution fractions were concentrated

and incubated with anti-BRIL Fab and Fab-Nb in a molar ratio of 1 : 1.2 : 1.5 on ice for 1 hour.

The complex was then subjected to size-exclusion chromatography (SEC) using a Superose 6 GL

Increase 10/300 (Cytiva) in buffer containing 50 mM HEPES pH 7.5, 150 mM NaCl, 0.003%

LMNG, 0.0003% CHS, 50 μM of OXM1 or OXM2 and 50 μM of SQA1. The SEC peak fraction

was collected and concentrated to 1 mg/ml for cryo-EM grid preparation.

Cryo-EM sample preparation and data acquisition. For both samples, cryo-EM grids were

prepared with a Vitrobot Mark IV (Thermo Scientific) by applying 3 μl of sample to freshly glow-

discharged Quantifoil Au R1.2/1.3 200-mesh grids and plunge freezing into liquid ethane. The

grids were prepared using a blot force of +3, a blot time of 2 s, and an absorption time of 20 s. The

sample chamber was kept at 4°C and 100% humidity.

The CCR6/SQA1/OXM1 dataset was collected using EPU (Thermo Scientific) on a Titan

Krios operating at 300 KeV and equipped with a Selectris energy filter (Thermo Scientific) at a

slit width of 10 eV and a Falcon 4 direct electron detector (Thermo Scientific) at a magnification

of 215,000× for a magnified pixel size of 0.575 Å per pixel. Data were collected using a defocus

range of -0.8 to -2.4 μm and a dose of 7.93 electrons per pixel per s for a total exposure time of

1.76 s divided into 60 subframes for a total accumulated dose of 40 electrons per Å^2.

The CCR6/SQA1/OXM2 dataset was collected using EPU (Thermo Scientific) on a Titan

Krios G3i operating at 300 KeV and equipped with a Bioquantum imaging filter at a slit width of

20 eV and a K3 direct electron detector (Gatan) at a magnification of 130,000× for a magnified

pixel size of 0.654 Å per pixel. Data were collected using a defocus range of -0.4 to -1.8 μm and

a dose of 18.27 electrons per pixel per s for a total exposure time of 1.06 divided into 45 subframes

for a total accumulated dose of 45 electrons per $Å^2$.

Cryo-EM data processing. Data processing was carried out using cryoSPARC (Structura

Biotechnology). For the CCR6/SQA1/OXM1 dataset, 12,852 movies were imported into

cryoSPARC and subjected to patch motion correction and patch CTF estimation. Exposures with

CTF fits better than 5 Å were selected (12,776 movies) and blob picking was used to select and

extract 5,332,464 particles. These particles were subjected to multiple rounds of 2D classification.

Ab initio model generation was carried out with 476,361 particles and 3 classes. The best model

was used as a template for heterogeneous refinement starting with 2,168,721 particles using one

good class model and 3 junk models. When most of the particles were sorted to the good class

(487,232 particles), the particles were re-extracted using the full box size of 520 pix. These

particles (482,484 particles) were subjected to nonuniform refinement, resulting in a map of 2.63

Å.

For the CCR6/SQA1/OXM2 dataset, 15,650 movies were subjected to patch motion

correction and patch CTF estimation. Exposures with CTF fits better than 5 Å were selected

(14,790 movies) and blob picking was used to select and extract 10,786,639 particles. These

particles were subjected to multiple rounds of 2D classification. Ab initio model generation for

three classes was carried out with 129,004 particles selected from 2D classification. The best

model was used as a template for heterogeneous refinement starting with 1,127,335 particles using

one good class model and 3 junk models. After three rounds of heterogeneous refinement, most of

the particles sorted to the good class (371,301 particles), the particles were re-extracted using the

full box size of 520 pix (368,745 particles). These particles were subjected to nonuniform

refinement, resulting in a map of 3.02 Å.

Model building and refinement. For each of the CCR6 structures presented, an atomic model

predicted by AlphaFold2[37] (thermostabilized CCR6_Nα7.1$_{ICL3}$BRIL fusion construct sequence)

was rigid-body fit into the map density. Starting models of the anti-BRIL Fab and the hinge-

binding nanobody were obtained from PDB entry 6WW2. Following the docking of each model

and ligands into the EM density map, iterative manual adjustment and real-space refinement was

carried out using Coot[38]. Subsequently, global refinement and minimization in real space was

performed using *phenix.real_space_refine* in Phenix[39]. The final models were visually inspected

for fit to the map, and Molprobity[40] was used for geometry validation. All structure figures were

generated by using PyMOL[41], UCSF Chimera[42], and UCSF ChimeraX[43].

Intrinsic fluorescence-based thermal shift assay. Purified CCR6 ICL3-BRIL fusion protein (9

µM) was incubated with 10 µM of the specified compounds in buffer containing 50 mM HEPES

pH 7.5, 500 mM NaCl, and 0.003% LMNG/0.0003% CHS. After a 5 min incubation at 25 °C, the

samples were loaded into capillary tubes for Tycho NT.6 instrument (NanoTemper). Tryptophan

fluorescence at emission wavelengths of 350 nm and 330 nm were measured as the sample

temperature was ramped from 35 °C to 95 °C according to the preprogrammed instrument protocol.

The ratio of the emission fluorescence was used to determine the T_i for the melting curves reported

by the instrument software.

CPM-based thermal shift assay. CPM-based thermal shift assay was conducted for CCR6 WT

by a differential scanning fluorimetry assay adapted from a previous publication[44] using UNit

(Unchained Labs). In brief, 5 μM of protein was incubated for 5 min at 25 °C with the presence of

10 μM of testing compound or 1% DMSO as control in buffer containing 50 mM HEPES pH 7.5,

500 mM NaCl, 0.05% DDM, and 2 μM of the thiol-specific fluorochrome *N*-[4-(7-diethylamino-

4-methyl-3-coumarinyl)phenyl]-maleimide (CPM) dye. After incubation, the samples were heated

from 20 °C to 80 °C at a temperature ramping rate of 2 °C/min, during which the CPM fluorescence

(excitation 384 nm, emission 470 nm) was recorded. Melting temperature (T_m) is defined as the

temperature at which 50% of the protein was denatured based on Boltzmann sigmoidal fitting.

OXM discovery by SQA-based virtual screen. A field-based 3D similarity search as

implemented in the Blaze module within the Cresset software[45-47] was performed using the small

molecule x-ray conformation[48], of an SQA analogue as a query. Briefly the method uses

electrostatics and shape of the query ligand to rapidly search large chemical collections for

molecules with similar properties. The basis of electrostatic potential calculations is the polarizable

XED force field which allows an accurate description of the charges around atoms distribution.

Each molecule is described as a Field Point using the local extrema of four molecular fields:

positive electrostatic, negative electrostatic, surface (van der Waals interactions), and

hydrophobic. Unlike traditional electrostatic field methods that place single point partial charges

at atom centers, this field redefines charges towards a multipole electron distribution around the

atom. The molecular fields developed from this representation are condensed to their local field

points and used to rapidly search and score scaffolds with similar electronic properties for

molecules in our internal database. The OXM analogue OXM1 with a similarity score of 0.74 was

identified from this approach.

NMR experiments using ROE techniques. Rotating-frame nuclear Overhauser enhancements

(ROEs) or nuclear Overhauser effect (NOE) experiments were collected using previously reported

methods[29] in CDCl$_3$ or DMSO-d_6. The absolute values for the ROE/NOE intensities were

measured relative to the irradiated peak which was arbitrarily assigned a value of -100. Calibration

for this experiment used a ROE/NOE intensity for the adjacent aromatic proton at a distance of 2.8

Å.

RDC NMR solution conformations. The proton and carbon assignments for OXM1, OXM2,

OXM3, and OXM4 were completed using a combination of 1D proton, 2D COSY, 2D HSQC, 2D

HMBC and 1D carbon experiments using standard Bruker pulse sequences and a concentration of

7 mg/200 μl. The assignments are shown in Supplementary Information (Figs. S1-S5, S8-12, S15-

19, S22-26). All NMR experiments were collected on a Bruker AVANCE III spectrometer

operating at a ^{1}H-Larmor frequency of 600.1 MHz that was equipped with a 5 mm TCI helium

cryoprobe using Topspin 3.2. The chemical shifts are referenced to residual solvent (deuterated

DMSO). Both $^1J_{CH}$ and $^1J_{NH}$'s[49] were collected for the four OXM analogues using compressed

DMSO-d$_6$ swollen poly-HEMA gels in New Era Enterprises compression devices. For the proton-

carbon and proton-nitrogen J-couplings, two J-scaled Bird (JSB) HC-HSQC spectra were

collected for each compound, see Supplementary Information (Figs. S6-S7, S13-14, S20-21, S27-

28). The first JSB HC-HSQC was collected on the compound in DMSO-d$_6$ alone (isotropic media)

while the second JSB HC-HSQC was collected on the compound in a poly-HEMA gel (anisotropic

media). The residual dipolar coupling (RDC) values (^1D) for proton-carbon and proton-nitrogen

J-couplings were determined using the formula: $^1D = {}^1T - {}^1J$ as reported previously[30]. Overall,

RDCs ($^1J_{CH}$ and $^1J_{NH}$) in a range of -8.9 to +13.3 Hz were measured for each compound in DMSO-

d$_6$ as shown in Supplementary Information (Table S3). The RDC data was fit to each member of a

conformational ensemble that consisted of 80-160 low energy structures that were generated in a

solvent dependent manner based on molecular dynamic simulations. Two files were generated for

each compound: a linear conformer file and a U- shaped conformer file. The RDC restraints were

fit to each file independently and the best results (lowest Q) are reported. The RDC data was fit

using Mestre MSpin/Stereofitter software using the input files provided (see Supplementary

Information)[50]. With this software, the Cornilescu Q-factor (Q)[51] was used to judge the goodness

of the fit for each conformation. The conformations with the best fit had $Q = 0.005$ for OXM1, Q

$= 0.097$ for OXM2, $Q = 0.001$ for OXM3, and $Q = 0.035$ for OXM4. In addition, the experimental

RDC's correlated well with the back-calculated values as shown in Table S3.

Conformer generation for RDC fitting experiments. All structures were first subject to

geometry optimization in Macromodel[52] using the OPLS3e forcefield and water as solvent, with

default parameters and convergence criteria. A conformational search was performed on each

structure using the mixed MCMM/low-mode search as implemented in the Macromodel program

with 1000 steps per rotatable bond and an energy window of 21 kJ mol−1 for retention of

conformers. The conformer files (OXM1, OXM2, OXM3, and OXM4) are provided in a

supplementary file for each compound.

OXM binding measurement by SPR. Binding affinity and kinetics of OXM1 were measured

with the CCR6_Nα7.1 construct using a Surface Plasmon Resonance (SPR)-based binding assay

on Biacore T200 instrument (Cytiva). The steps of sensor preparation and protein capture were

carried out at 25 °C and the binding interactions were measured at 10 °C. The CCR6 protein was

initially captured via immobilized metal affinity chromatography (IMAC) followed by amine

coupling onto an activated Ni-NTA sensor chip treated by a 1:1 mixture of 0.05 M *n*-

hydroxysuccinimide (NHS) and 0.2 M 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC).

The experiments were carried out in 50 mM HEPES buffer (pH 7.5) containing 150 mM NaCl,

0.1% DDM, and 3% DMSO. OXM1 was injected in a concentration series consisting of 6-point,

3-fold dilutions from 10 µM to 0.014 µM. The data were processed and analyzed using Biacore

Biaeval and Scrubber 2.0 software. The binding affinities and on and off rates were calculated by

globally fitting the sensorgrams to a 1:1 binding model.

Crosstalk study by SPR. The crosstalk SPR experiments were performed using a Biacore T200

Instrument (Cytiva) equipped with a NiHC 200M sensor chip (Xantec) at 25 °C. The run buffer

for all experiments, unless otherwise stated, was 50 mM HEPES pH 7.5, 500 mM NaCl, 1 mM

$MgCl_2$, 0.03% DDM, 0.003% CHS, 5% DMSO. The CCR6_Nβ10.2 protein was immobilized on

the sensor chip by Ni-NTA pull down and amine coupling. 0.5 M EDTA pH 8.0 was injected for

60 s at 5 µL/min followed by a 60 s injection of run buffer containing 0.5 mM $NiCl_2$. The surface

was activated with a 1:1 mixture of 0.1 M NHS and 0.5 M EDC at a flow rate of 10 µL/min. The

receptors were injected at a concentration of 0.5 µM at 5 µL/min until a surface response of 2000

RU was achieved. The surface was equilibrated for 10 h in run buffer before commencing

experiments. OXM1 binding affinity was measured in a 5-point, 2-fold dilution series in single

cycle kinetics format with a top concentration of 1 µM in run buffer with and without 100 nM

SQA1. The flow rate of binding measurement was 50 µL/min. The association phase was 120 s

and dissociation phase was between 1800 s. All experiments were run in triplicate. The data were

processed using Biacore T200 Evaluation Software (Version 2.0, Cytiva). Responses were

corrected for volume exclusion effects. Sensorgrams were reference and blank subtracted before

fitting with 1:1 binding model which accounted for drift, bulk shift and mass transport. Sensorgram

images were prepared using GraphPad Prism software.

Protein surface expression measurement. HEK293 cells transiently expressing CCR6 constructs

or mock transfected control after 72 h post transfection were detached using an enzyme-free

dissociation buffer. Cells were blocked with 0.1 mg/ml mouse IgG prior to staining with BV-421

labeled anti-human CCR6 antibody (Clone 11A9, BD Biosciences) for 1 h at 4 °C. After washing,

cells were analyzed on a BD LSRFortessa cytometer (BD Biosciences) for cell surface expression

of CCR6 by determining the mean fluorescence intensity (MFI) of BV-421 using FlowJo™

Software (Becton, Dickinson and Company).

Radioligand saturation binding in HEK293 or CHO-K1 cells. Saturation binding experiments

were performed on HEK293 cells transiently expressing human CCR6 48 h post transfection, or

CHO-K1 cells stably expressing human CCR6 (DiscoverX). HEK293 cells were plated at

80,000/well in growth media (DMEM + 10% Serum + 20 mM HEPES) in white/clear bottom PDL

coated 96 well plates (Corning); CHO-K1 cells were plated at 50,000/well in white/clear bottom

96 well Isoplates (PerkinElmer) in growth media a day before the experiments. On the day of

assay, cells in the assay well were treated with either 0.3-0.4 % DMSO (for total binding) or 90-

120 μM of unlabeled SQA1 (for non-specific binding), followed by addition of a serial dilution of

[^{3}H]-SQA1 in growth media. For each dose of [^{3}H]-SQA1, triplicate reactions were prepared for

both total and non-specific binding conditions. After 3 hours of incubation at room temperature,

cells were washed 3 times with wash buffer (PBS + 1 mM $CaCl_2$ + 0.1% BSA) before adding

Microscint-20 (PerkinElmer) for tritium concentration measurement using a TriLux MicroBeta2

plate reader (PerkinElmer). Data was analyzed with GraphPad Prism software using a one-site

saturation binding equation.

CCR6+ T cell chemotaxis assay. Human CD4$^+$CCR6$^+$CXCR3$^-$ T cells were isolated from human

donor leukopaks using EasySep™ Human Th17 Cell Enrichment Kit (StemCell Technologies,

18162). To obtain large quantities of cells, CCR6+ T cells were activated with Dynabeads Human

T-activator (Gibco, 11132D) at a density of 1×10^6 cell/ml in growth media (RPMI1640 media

with 10% serum, 4 ng/ml IL-2) with a 1:1.5 cell to bead ratio. On day 4 post activation, Dynabeads

were removed from the culture. The activated T cells were maintained at 1–2×10^6 cells/ml for 15

days by feeding fresh growth media when needed. The CCR6+ T cell chemotaxis assay was carried

out on day 12 to 15 post T cell activation using the 96 well ChemoTx® disposable chemotaxis

system (Neuroprobe 101-5) according to the manufacturer's protocol. After one wash with assay

buffer (1 × HBSS containing 20 mM HEPES and 0.25% BSA), cells were incubated with test

compounds for 30 minutes at room temperature prior to initiation of chemotaxis. CCL20 induced

$CCR6^+$ human T cell chemotaxis with an $EC_{50} = 0.19$ nM (pEC50 = 9.7 ± 0.9, mean $\pm$ s.d., $n = 5$).

CCL20 was used at 0.5 nM in all measurements. For IC50 determination, the top and bottom of

the chemotaxis chamber contained the same concentration of compound. DMSO was kept constant

at 0.1% (v/v) in all wells. The final concentration of CCL20 (Peprotech, 300-29A) in the bottom

chamber is 0.5 nM. The fully assembled chemotaxis plate was placed in a cell culture incubator at

37 °C, 5% CO_2 for 1 h. After incubation, the top filter was removed, followed by a quick freeze of

the bottom chamber at -80 °C for 1 h. The migrated cells in the bottom chamber were stained with

CyQUANT dye (Life Technologies, C7026) for cell number determination. Data was analyzed

with GraphPad Prism Software using a non-linear regression analysis of dose response curves for

IC50 determination.

GPCR β-arrestin assay panel. OXM1 was tested at 1 μM for agonistic and antagonistic activity

on a panel of GPCRs in *gpcr*MAX℠ GPCR assay panel using PathHunter® β-arrestin enzyme

complementation technology as described in Eurofins Discovery website (gpcrMAX-

(discoverx.com)).

Preparation of [^{3}H]-SQA1 and Synthesis of different OXM analogues, OXM1, OXM2,

OXM3, and OXM4 are in supplementary information.

Data availability statement. The cryo-EM density maps for the complexes are deposited in the

Electron Microscopy Data Bank (EMDB) under accession codes EMD-xxxxx

(CCR6/SQA1/OXM1) and EMD-xxxxx (CCR6/SQA1/OXM2). The coordinates for the models

are deposited in the Worldwide Protein Data Bank (wwPDB) under accession codes XXXX

(CCR6/SQA1/OXM1) and XXXX (CCR6/SQA1/OXM2). Solution conformer library files of the

OXM analogues are provided as source data in .sdf format. Other source data are also provided

with this paper. All the other data supporting the findings of this study are available within the

article and its Supplementary Information files and from the corresponding author upon reasonable

request.

Acknowledgements

Cryo-EM data for the CCR6/SQA1/OXM2 complex was collected at the Cryo-EM Facility at

MIT.nano. We thank K. Fennell and H. Zhao for recombinant protein expression support; K.

Schildknegt, T. Kenakin, and E.J. Corey for consultancy support for the CCR6 antagonist project;

C. Tang, Q. Jin, and N. Fadeyi for synthesis support; X. Qiu and D. Hepworth, for project

management support. We thank all members of Pfizer CCR6 research project team for suggestions

and comments.

Author contributions

D.J.W. performed protein purification, thermal shift experiments, prepared cryo-EM grids,

collected and processed cryo-EM data, determined the structures, and assisted with preparing the

manuscript; B.S.G. supervised discovery and design of the OXM analogues; K.A.F. performed

NMR studies; W.L. designed and supervised all pharmacology experiments; J.A. performed

radioligand binding and CCR6 mutagenesis studies; M.E.S. supervised discovery of the SQA

analogue; R.J.U. performed ligand-based virtual screening and generated conformer libraries for

the OXM analogues; J.V. performed protein purification; K.K.C. performed T cell chemotaxis experiments; R.D. performed protein surface expression measurements; P.V.S performed SPR experiments; F.V. and R.K.F. performed T cell chemotaxis experiments; A.D., A.F., C.C., G.C., J.J.M., J.I.T., P.N., P.M., and V.L. performed synthesis; D.L., E.S., and C.O. performed StaR engineering, characterization, and saturation bindings; B.J.H. performed crosstalk SPR studies; G.S.M. purified CCR6 protein for crosstalk SPR studies. I.W.W. collected cryo-EM data of the SQA1/OXM2 complex; S.H. assisted with cryo-EM data collection of the SQA1/OXM1 complex; M.R. and R.M.C. supervised StaR engineering and characterization; M.F.C. managed structural and biophysical studies and reviewed the data; G.B. analyzed and interpreted pharmacology data; A.T. supervised and managed the project; H.W. designed and supervised all structural and biophysical studies, determined the structures, analyzed the results and wrote the manuscript.

Competing interests

All authors were either employees of Pfizer Inc. or were employees of Sosei Heptares at the time the work was performed.

Additional information

Extended data is available for this paper at (web).

Supplemental information is available for this paper at (web).

Correspondence and requests for materials should be addressed to huixian.wu@pfizer.com.

Reprints and permissions information is available at (web).