Title: Predation by a ciliate community mediates temperature and nutrient effects

on a peatland prey prokaryotic community.

Katrina DeWitt[1*], Alyssa A. Carrell[3], Jennifer D. Rocca[1], Samantha Votzke[1], Andrea Yammine[1],

Ariane L. Peralta[2], David J. Weston[3], Dale A. Pelletier[3], Jean P. Gibert[1]

[1]Department of Biology, Duke University, Durham, NC, USA

[2]Department of Biology, East Carolina University, Greenville, NC, USA

[3]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

Running Title: Predation and temperature affect bacterial communities.

#Address correspondence to Katrina DeWitt, katrina.mariedewitt@gmail.com

*Present address: Samantha Votzke, Department of Earth and Planetary Sciences, Johns Hopkins

University, Baltimore, MD, USA

Abstract word count: 222

Importance word count: 111

Main text word count: 4940

Keywords: Ciliates, Protoctists, Climate Change, Synthetic Communities, Microbial Food Webs

Abstract:

Temperature significantly impacts microbial communities' composition and function, which

subsequently plays a vital role in the global carbon cycle that ultimately fuels climate change.

Interactions between different microorganisms might be critical in shaping how these

communities react to both temperature changes. Additionally, rising temperatures are occurring

in the context of increasingly nutrient-rich ecosystems due to human activity. Nonetheless, we

lack a comprehensive understanding of how predation influences microbial communities in

future climate scenarios and an increasingly nutrient-rich world. Here, we assess whether

predation by key bacterial consumers—ciliates—influences a microbial community's freshwater

temperature and nutrient response regarding biomass, diversity, structure, and function. In a

three-week microcosm experiment, we exposed mostly prokaryotic microbial communities to a

community of ciliate predators at two different temperature scenarios (ambient and +3°C, i.e., a

conservative projection of climate change by 2050) and nutrient levels (low and elevated).

Nutrients, temperature, and ciliate presence influenced microbial biomass and function

separately, but their interaction had the largest explanatory power over the observed changes in

microbial community biomass, structure, and function. Our study supports previous findings that

temperature and nutrients are essential drivers of microbial community structure and function but

also demonstrates that the presence of predators can mediate these effects, indicating that the

biotic context is as important as the abiotic context to understand microbial responses to novel

climates.

Importance:

While the importance of the abiotic environment in microbial communities has long been

studied, how prevalent ecological interactions, like predation and the broader abiotic context,

may influence these responses is largely unknown. Our study disentangles the complex interplay

between temperature, nutrients, and predation and their joint effects on microbial community

diversity and function. The findings suggest that while temperature and nutrients are

fundamental drivers of microbial community dynamics, the presence of predators significantly

mediates these responses. Our study underscores the profound impact of abiotic factors on

microbial communities, but how to properly understand, let alone predict, these responses, we

need to account for the biotic context in which these are occurring.

Introduction:

Rising global temperatures are currently affecting populations [1,2], communities [3], and ecosystems [4] through an increase in metabolic rates [5,6]. Increased metabolic rates then lead to increased energetic demands for organisms within populations [7–9], and this has cascading effects across ecological scales [5,10–12]. In addition to temperature-driven ecological change, ecosystems face increasingly large nutrient loads from anthropogenic sources (e.g., agricultural runoff) and increased mineralization rates [13], which have long been known to impact the structure and dynamics of communities and ecosystems [14–19]. While the independent effects of rising temperature and nutrients on ecological processes are relatively well understood [20–22], joint increases in temperature and nutrients have been shown to have interactive effects that can be hard to predict and are thus poorly understood [23–28]. Yet, this understanding is central to assessing whether and how ecological communities will respond to an increasingly warm, nutrient-rich, and human-dominated world.

Some of the most consequential biological communities worldwide are also some of the least well-understood –prokaryotic communities– which comprise upwards of 14% of all existing biomass on Earth [29]. These communities are present on all continents and ecosystems [30,31], and their central role in the global carbon and nutrient cycles –which ultimately determine the pace of global climate change– is increasingly recognized [32,33]. For example, prokaryotic decomposition recycles organic material into food webs [34,35], and prokaryotic respiration is one of the atmosphere's most significant sources of carbon dioxide (CO_2) [36,37]. These respiration rates can change dramatically across temperatures via shifts in individual metabolic rates [5–7,38–41], which could result in a "warming begets warming" scenario. Therefore, how prokaryotic

communities respond to joint changes in temperature and nutrients is central to understanding

and better predicting the global effects of climate change.

In addition to rapid abiotic change, prokaryotic communities worldwide will likely also

respond to a rapidly changing biotic environment as food webs rewire under climate change [42–44].

These shifting ecological interactions often determine community and ecosystem-level processes

[45]. For example, predation is a crucial factor leading to decreased biomass in prokaryotic

communities[46–49], and bacterivores are predicted to increase foraging rates to offset metabolic

costs [50]. This increase in bacterivory –and predation, in general– may impact the composition

and function of many prokaryotic communities in a rapidly changing world [49,51–53]. However,

how these prokaryotic communities may respond to both temperature increases and nutrient

additions in the presence of predators is an active area of research [54].

Ciliates are among the planet's most abundant predators of prokaryotes [29,30,55,56]. These

microbial predators show strong, well-known temperature [41,57–59] and nutrient responses [28,60–62],

as well as temperature-dependent population dynamics [41,59] and feeding interactions [6,58]. Ciliates

have been shown to affect prokaryotic community structure, dynamics, and function strongly [63]

and can affect species interactions and ecological dynamics within microbial food webs [28,58,64].

Under warmer conditions, predation by ciliates could, therefore, significantly impact prokaryotic

respiration and overall function. Whether this would result in a "warming begets warming"

scenario or a "warming dampens warming" scenario, however, depends on the specific

temperature response of the predators and the prey and how this ecological interaction constrains

the rising temperature and nutrient effects.

We thus examine whether and how predation of prokaryotes by a ciliate community

mediates the combined effects of nutrient additions and temperature on community biomass,

diversity, composition, and function (measured as total respiration rates). We used experimental

microcosms containing a synthetic community of ciliates preying on a natural peatland

prokaryotic community to address the questions: 1) How do temperature, nutrients, and the

presence of a ciliate community affect prokaryotic community biomass, diversity, and

composition? 2) How do these factors affect total community respiration? 3) How does the

ciliate community, in turn, respond to imposed changes in temperature and nutrients?

We hypothesize that prokaryotic biomass and respiration rates should increase under

warming temperatures and high nutrient conditions [65,66]. We should also observe that ciliate

predation decreases prokaryotic biomass [67,68], reducing total respiration rates [46], which also

should result in a shift in the composition of the prokaryotic community due to differential

consumption by ciliates on different species[67,68]. As predation by a single ciliate species has been

observed to result in an interactive effect of predation, temperature, and nutrients on prokaryotic

community respiration and biomass (Rocca et al. 2022), we hypothesize that an entire ciliate

community should also have an interactive effect of temperature and nutrients on prokaryotic

community respiration and biomass. Last, a recent study has shown that natural ciliate

communities can respond interactively to temperature and nutrients[69,70], so we expect that the

predator ciliate community should also respond differently to temperature depending on nutrient

levels.

Results

Effects of temperature, nutrients, and ciliate presence on total prey community biomass

and respiration

Preliminary data analysis suggested that ciliate presence, temperature, and nutrient levels, as well

as their interactions, likely influenced OD600, while nutrient levels were the most important

variable explaining respiration, followed by the presence of ciliates and temperature

(Supplementary Figures S4, S5; Appendix 3).

140   Additional analyses showed that temperature, nutrients, and the presence of a ciliate

community all had independent effects on total community biomass (OD600, Figure 1). Higher

temperature (Figure 1A) and low nutrient levels (Figure 1B) resulted in less total biomass (i.e.,

smaller OD600, temp effect = -0.008 $\pm$ 0.003 SE, t-value = -2.788, df = 118, p-value = 0.006;

nutrient effect = 0.007 $\pm$ 0.003 SE, t-value = 2.524, df = 118, p-value = 0.012), while the

presence of ciliates resulted in more overall biomass (Figure 1C, effect = 0.009 $\pm$ 0.003 SE, t-

value = 3.025, df = 118, p-value = 0.003). Temperature, nutrients, and the presence of ciliates

had a significant interactive effect on total community biomass (Figure 1D, 1E). Under high

nutrient levels, the presence of ciliates increased OD600 (effect = 0.016 $\pm$ 0.007 SE, t-value =

2.041, df = 112, p-value = 0.043, Figure 1D, 1E) compared to the treatments that do not have

ciliates. Moreover, how nutrients affected total biomass in the presence of ciliates varied across

temperatures (i.e., three-way interaction, effect = -0.035 $\pm$ 0.011, t-value = -3.215, p-value =

0.001, Figure 1D, E). Specifically, under low nutrient conditions, the microbial biomass

increased in response to warmer temperatures when ciliates were added compared to a decrease

in the absence of ciliates. Under high nutrient conditions, there was a decrease in the response to

warming of the microbial OD600 in the presence of ciliates but no change in the absence of

ciliates.

157   Respiration rates were not independently influenced by temperature (Figure 2A) or ciliate

presence (Figure 2C) but were affected significantly by nutrient level (Figure 2B, effect = 2.213

$\times 10^{-4} \pm 4.84$ SE, t-value = 4.564, df = 118, p-value = 1.24×10^{-5}). However, respiration rates

increased with temperature only under high nutrient levels for treatments that did not have

ciliates, suggestive an interactive effect of nutrient levels and temperature (Figure 2D, 2E, effect

= $3.313 \times 10^{-4} \pm 1.654 \times 10^{-4}$, t-value = 2.004, df = 112, p-value = 0.047).

Effects of temperature, nutrients, and ciliate presence on the prey community structure

Across all treatments, Proteobacteria were the most abundant group in the prey prokaryotic

community, followed by Firmicutes and Cyanobacteria (Figure 3A). Temperature, nutrients, and

ciliate presence all had a significant interactive effect on Shannon diversity (ANOVA, F =

13.449, p-value = 3.76×10^{-4}, Figure 3B), while temperature and ciliate presence also had a

significant interactive effect (ANOVA, F = 7.3073, p-value = 0.007). Shannon diversity

increased in the presence of ciliates under high nutrient levels and low temperatures (22°C,

Figure 3B). In the high temperature (25°C), however, Shannon diversity increased in the absence

of the ciliates and high nutrients but decreased in the presence of the ciliates and high nutrients

(Figure 3B). This three-way interaction is likely the result of an increase in the dominance of

Firmicutes relative to Proteobacteria in conditions of low nutrients, low temperature, and ciliate

presence (Figure 3A).

Beta diversity was significantly affected by temperature, nutrient levels, and the presence

of ciliates (Temp: PERMANOVA, F = 4.919, p-value = 0.001. Nutrients: PERMANOVA, F =

4.625, p-value = 0.001. Pro: PERMANOVA, F = 15.913, p-value = 0.001. Figure 3C). In

addition, all two-way interactions and the triple interaction between temperature, nutrients, and

ciliates were significant (Temp:Nutrient, PERMANOVA, F = 2.551, p-value = 0.002,

Temp:Ciliate, PERMANOVA, F = 2.218, p-value = 0.008, Nutrient:Ciliate, PERMANOVA, F =

1.939, p-value = 0.011, Temp:Nutrient:Ciliate, PERMANOVA, F = 2.253, p-value = 0.003,

Figure 3C). These results suggest that while prey alpha diversity remained constant across

treatments, the composition responded strongly to all variables and their interaction, indicating

substantial species turnover among treatments.

Effects of temperature and nutrients on the predatory ciliate community

Across all treatments, *Blepharisma sp.* was the most abundant ciliate species, followed by

Colpidium sp. and *Paramecium bursaria* (Figure 4A); two species, *Tetrahymena pyriformis* and

Cyclidium glaucoma, were not present in any samples. This is most likely due to being

outcompeted and eaten by other ciliates within the system. Nutrient levels significantly impacted

predator ciliate community alpha diversity, while temperature or temperature and nutrient

interactions did not. Shannon diversity was higher in low nutrient conditions (ANOVA, F =

6.370, df = 1, p-value = 0.013 Figure 4B). All population counts were similar across

experimental treatments except for *Blepharisma sp.* counts, which were significantly affected by

temperature, nutrients, and their interaction (temperature effect = -205.533 ± 68.833 SE, t-value

= 2.986, p-value = 0.003; nutrient effect = -277.450 ± 97.902 SE, t-value = -2.834, p-value =

0.005; nutrient:temp effect = 10.650 ± 4.161 SE, t-value = 2.559, p-value = 0.012).

Interestingly, the imposed treatments seem to have shaped the composition of the ciliate

communities in such a way that three distinct community configurations were possible, each

corresponding to a principal component: a *Colpidium sp.*, *Paramecium bursaria, Paramecium*

aurelia, and *Glaucoma sp.* dominated community (PC1, Figure 4D), a *Blepharisma sp.* and

Colpoda steinii dominated community (PC2, Figure 4D), and a *Halteria grandinella*, and

Strombidium sp. dominated community (PC3, Figure 4D). The overall PCA was significantly

affected by nutrient level and its interaction with temperature (Nutrient: PERMANOVA, F =

2.144, df = 1, p-value = 0.029; Nutrient:Temp: PERMANOVA, F = 2.022, df = 1, p-value =

0.034). This suggests that a combination of direct environmental effects and indirect responses of

the prey community to the abiotic manipulations likely influence the structure and composition

of the predator community.

Joint direct and indirect effects of temperature and nutrients on the prokaryotic and ciliate

communities

The SEM model fitted the data well (Fisher.C = 5.017, p-value = 0.286) and confirmed that

temperature alone influences prey biomass (0.72, p-value = 0.024) and the Shannon diversity of

the prey community (1.328, p-value = 0.0001). However, the SEM suggested that neither

temperature, nutrients, nor their interaction influence the predator's community structure (Fig 5),

which does not align with our earlier findings (Fig 4). Nutrients alone affected total microbial

biomass (5.31, p-value = 0.0009), diversity of the prey community (5.839, p-value = 0.0006),

and the biomass of the ciliate predators (4.13, p-value = 0.0133), but not the diversity of ciliates

or the overall function (Fig 5). The interaction between temperature and nutrients directly

affected ciliate biomass (standardized effect size (ses) = -3.7601, p-value = 0.0240), the biomass

of the prey (ses = -5.36, p-value = 0.0011), and prey community diversity (ses = -5.930, p-value

= 0.0007). Total functioning (respiration) was directly affected by the biomass of the predators

(ses = 0.44, p-value = 0.003) and prey (ses = -0.28, p-value = 0.028) and ciliate diversity (ses =

0.32, p-value = 0.006), and was indirectly affected by nutrients through the biomass of ciliates

(Fig 5).

Additionally, the SEM uncovered critical indirect effects of temperature and nutrients,

notably, an indirect negative effect of temperature on total respiration through prey microbial

biomass (Fig 5). Nutrients, in turn, had two distinct indirect pathways: one positive indirect

effect through ciliate biomass and one negative indirect effect through ciliate biomass's effect on

prey microbial biomass (Fig 5). Last, temperature and nutrient interaction also indirectly affected

respiration through their effect on ciliate biomass, and the effects of ciliate biomass directly on

respiration and indirectly on respiration through total prey microbial biomass (Fig 5).

Discussion

Shedding light on how changes in environmental temperature and nutrient levels may impact

prokaryotic biomass and function (e.g., total respiration) is crucial to our ability to recognize and

predict future warming trends [33]. First, our study shows how temperature and nutrient levels

determine a prokaryotic community's biomass, composition, and respiration rates (Figures 1, 2,

3). Second, results reveal that a ciliate predator community influences these responses (Figures

1A, 1B, and 1C). Last, we observe that the predatory community's biomass, diversity, and

composition respond to temperature and nutrient levels directly, to shifts in the prey prokaryotic

communities they prey upon, or a combination of both (Figure 4, Figure 5). These results

indicate that accounting for ecological interactions between prokaryotic prey and eukaryotic

predators is important to understanding prokaryotic communities' responses to compound

environmental changes.

Increased temperatures and low nutrient levels reduced microbial biomass (Figures 1A

and B), consistent with results in non-prokaryotic taxa across ecosystems[46,87]. Interestingly, joint

increases in temperature and nutrient levels have been predicted to increase microbial biomass

by intensifying eutrophication [27,88,89]. We, however, do not see an increase in biomass in the high

nutrient-high temperature treatment, perhaps due to the resilience of prokaryotic biomass to

changes in nutrient loads through compositional shifts [90]. While under elevated temperatures,

prokaryotic communities are known to shift to be mesophilic and perform anaerobic respiration;

thus, the prevalence of Firmicutes tends to be higher [91,92]. Proteobacteria are found in higher

relative abundances in warmer temperatures; perhaps this is most likely due to the presence of

ciliates (also observed in Rocca *et al.* 2022) and could represent prey switching by the ciliates.

The composition of the prokaryotic community, however, responded strongly to all treatments

and their interaction, as beta diversity was significantly affected by temperature, nutrient levels,

and the presence of ciliates (Figure 3C).

Total community respiration rates increased with temperature, but only under high

nutrient levels (Figures 2D and 2E), regardless of ciliate presence. These results are partly

consistent with theoretical expectations of increased metabolic demands with temperature [5,7,47]

and, hence, respiration rates [40,93–95]. However, they also suggest that changes in metabolic

demands with temperature depend on nutrient availability: metabolic rates cannot increase unless

enough nutrients fuel that increases. Further, this result implies that under low nutrients, the

function of these microbial communities is less likely to change with warming unless there is a

concomitant increase in nutrient levels to allow for increased function [96]. Alternatively, increased

respiration rates due to nutrient enrichment could indicate increased metabolic activity in

organisms due to increased turnover [97,98].

Moreover, while temperature, nutrient levels, and ciliate presence interactively influenced

prokaryotic biomass (Figure 1), only nutrient levels significantly impacted total respiration

(Figure 2). In other words, despite the effects of ciliate presence on prey prokaryotic community

structure and biomass, there seems to be no effect on its function. Yet, recent studies have argued

that direct and indirect effects of ecological interactions can influence the function of prokaryotic

communities in shifting abiotic conditions [49,56,99–101]. Indeed, previous work has shown that

individual ciliate species can influence the biomass and the temperature response of respiration

of a prokaryotic community [53], while one study even found that predation could reverse the

effects of warming in a tundra community [102], which seems at odds with our results. On the other

hand, the SEM did reveal that temperature, nutrients, and their interaction did indirectly affect –

albeit weakly– total microbial respiration (Fig 5) through ciliate biomass, suggesting that direct

effects of temperature and nutrients on the predatory ciliate community mediate their effect on

total respiration rates and ecosystem function of these microbial communities. This seeming

inconsistency in our results likely means that any effects on total microbial function may only be

observed in simpler communities, but perhaps not speciose, more complex ones. Additionally, all

indirect pathways are accounted for, highlighting the need for a deeper understanding of how

predation by ciliates and other Eukaryotes impacts the global carbon cycle through its influence

on the function of prokaryotic communities.

The predatory ciliate communities strongly responded to nutrient levels and their

interaction with temperature (Figure 4). We showed that the alpha diversity of the predator

ciliate community was higher under low nutrient levels (Figure 4B) and that nutrient levels

affected community composition significantly differently across temperatures (Figure 4C).

However, temperature can magnify or dampen this nutrient effect. Ciliate community

composition (and function) can change across temperatures in both tightly controlled laboratory

conditions [41] and whole-ecosystem warming experiments in boreal peatlands [70]. These recent

results are consistent with ours and suggest that neither nutrients nor temperature alone can

explain observed complex shifts in these ciliate predator communities.

Caveats. We did not track the responses or dynamics of the prey prokaryotic and ciliate

communities over time. Consequently, our results only represent a snapshot in time, and some of

the effects of predation reported here may be transient. Also, our temperature manipulations only

represent the effects of mean temperature in relatively stable conditions. So, our results should

not be used to infer the broader consequences of seasonal temperature shifts, as we do not

consider diurnal or seasonal temperature variation. As global temperatures rise, seasonality, in

particular, is undoubtedly also changing [103,104]; it remains an open question whether the effects of

temperature variability are distinct from those of changes in mean temperatures shown here.

While the results presented are strong and clear enough to suggest that the processes at play may

also be acting in nature, the short-term nature of our experimental manipulations and our

experimental setup in tightly controlled experimental conditions inevitably limit the possible

reach of our inference.

Conclusions. Our results shed some light on how warming may affect carbon

sequestration in peatlands. Indeed, peatlands are exceptionally susceptible to future climate

change impacts [105–107]. Despite covering less than 3% of the Earth's surface, these ecosystems

store approximately 25% -30 % of the world's soil carbon as recalcitrant peat [108]. However, peat

moss growth in nutrient-poor peatlands depends on symbiotic interactions with prokaryotic

associates [109,110]. Shifts in prokaryotic community composition due to warming [111] may impact

the activity of moss symbionts and other important microbes, potentially leading to reductions in

carbon sequestration in these peatlands, through changes in moss growth [112]. Our results

highlight that to fully understand the breadth and consequences of these changes, we need to

consider the interactive effects of rising temperatures and increasing nutrient loads in these

rapidly changing ecosystems. Moreover, our results suggest that essential but largely overlooked

predatory interactions between these organisms and their predators may be important to

understanding and predicting how climate change may affect carbon dynamics in peatlands[13].

We highlighted critical temperature and nutrient effects on prey prokaryotic community

structure and function. We also show that these effects are often interactive, suggesting that we

cannot predict these microbial communities' response to climate change without considering

possible eutrophication or other anthropogenic nutrient additions. Last, we show that ciliate

predation mediates the effects of temperature, nutrients, and their interactions on prokaryotic

communities. The predators themselves respond to these abiotic conditions, possibly through a

combination of direct and indirect effects. Our study suggests interesting mechanisms for

predation to influence prokaryotic function in a changing world.

Materials and Methods:

Prey Prokaryotic Community

We isolated a prey –mostly prokaryotic– community from a peatland bog in the Marcell

experimental forest (Minnesota, USA) right next to the SPRUCE (Spruce and Peatland

Responses Under Changing Environments) whole-ecosystem warming experiment [71]. These

communities were isolated from 5 cm core samples containing a top layer of living *Sphagnum*

moss tissue and a lower layer of peat. To isolate the prokaryotes, we filtered out larger fungi and

ciliates using autoclaved filters (11 μm pore size, Whatman), then removed smaller flagellates

and fungal spores [49] using sterile GF/A filters (1.6 μm pore size, Whatman). While the

community is not guaranteed only to have prokaryotes (e.g., eukaryotic nanoflagellates cannot be

filtered out using this approach), it should be composed in its majority by Prokaryotes and

Archaea [53] and will henceforth be called the prokaryote community.

Experimental Treatments

The resultant community was homogenized and incubated in 120, 200 mL acid-washed, and

autoclaved borosilicate jars filled in 150 mL of Carolina Biological Ciliate culture medium [72].

Jars were assigned to one of two possible nutrient treatments: low nutrients (75 mL of media and

75 mL of distilled water, 60 jars) and high nutrients (150mL of media, 60 jars), and one of two

possible temperature treatments (60 jars at 22°C, representing ambient temperature, and 60 jars

at 25°C, representing an increase in the average temperature of 3ºC). All microcosms were

incubated under controlled environmental conditions in Percival AL-22 L2 growth chambers

(Percival Scientific, Perry, Iowa) at 10% light intensity (1700 lux), 75% humidity, and a 16:8 hr.

day-night cycle for three days before further manipulation.

We added the synthetic ciliate communities after the three-day pre-incubation period to

acclimate the prokaryotic communities. To this end, we assigned each experimental microcosm

to one of two possible ciliate treatments (40 jars had no ciliates, and 80 jars were inoculated with

ciliates). The asymmetrical design for the ciliate treatment was to ensure that even small or

nuanced effects of predation could be detected. These "ciliate" treatment microcosms were

inoculated with the ciliates *Paramecium aurelia, Paramecium bursaria, Halteria grandinella,*

Tillina magna, Tetrahymena pyriformis, Cyclidium glaucoma, Colpoda steinii, Strombidium sp.,

Blepharisma sp., Glaucoma sp., and *Colpidium sp.,* all of which naturally occur in *Sphagnum*

peatlands [30]. The ciliates came from long-term (> five years) lab cultures [41], sourced from

Carolina Biological (Burlington, NC, US) or Sciento (Manchester, UK), except for *P. bursaria,*

which was cultured from SPRUCE samples. Ciliates were introduced by pipetting well-mixed

stock cultures at carrying capacity into experimental microcosms. Each species was introduced at

a density of at least 17 individuals per jar (see Supplementary Table 1, ST1; Appendix 1 for

initial densities). To control for possible effects of other microbes already occurring in the ciliate

cultures –notably, prokaryotes– we added the same volume of filtered (11 μm pore size,

Whatman), homogenized, ciliate stock media from the original cultures into all "non-ciliate"

treatment microcosms in the identical amounts these were introduced in the ciliate treatments, as

done elsewhere [53,63,73]. This inoculation of filtered media through filters allows the prokaryotes to

pass through but filters out the ciliates to ensure that all experimental jars start with almost

identical prey prokaryotic communities. Following standard ciliate culturing procedures [72], we

also added one wheat seed (~35 mg ea.) to high-nutrient treatments and half of a seed (~17.5 mg

ea.) to the low-nutrient treatments to serve as a slow-release carbon source and prevent resource

limitation issues [72]. The wheat seeds were pre-autoclaved and sourced from the same batch to

control additional wheat seed-associated microorganisms.

Microcosms were assigned to one of eight possible treatments: (1) 22°C, no ciliates, low

nutrient; (2) 22°C, no ciliates, high nutrient; (3) 25°C, no ciliates, low nutrient; (4) 25°C, no

ciliates, high nutrient; (5) 22°C, ciliates, low nutrient; (6) 22°C, ciliates, high nutrient; (7) 25°C,

ciliates, low nutrient; and (8) 25°C, ciliates, high nutrient.

Measurement of Prey Community Biomass, Diversity, and Respiration Rate

After 21 days, we quantified the prey community's biomass, diversity, and total community

respiration rate to assess the individual and joint impacts of temperature, nutrient levels, and

ciliate predation. We measured the optical density at 600 nm wavelength (or OD600) [74] as a

proxy for the total biomass of each community microcosm using a BioTEK Epoch-2 microplate

spectrophotometer (Winooski, VT, USA). Greater OD600 measurements (indicating increased

absorbance and scattering) signify elevated overall biomass [74]. Protists scatter light to a lesser

extent due to their larger size and reduced densities [64]. Community respiration was determined

using an optode-based real-time OXY-4 SMA respirometer (PreSens, Regensburg, Germany)

[72,75]. The respiration rate was measured on 150 ml samples for at least 30 minutes, with a

collection rate of one measurement every three seconds, after a 30-min acclimation period (n =

120) at their original experimental temperature and in the dark. Respiration rates were estimated

as the rate of change (slope) of the estimated oxygen concentration over time (in μmol O_2/min;

Supplementary Figures S1, S2, and S3; Appendix 2) following standard optode-based closed-

system respirometry procedures [76].

Analysis of Prey Community Biomass, Diversity, and Respiration Rate Data

To understand the joint effects of temperature, time, and ciliate presence on biomass (OD600)

and respiration, we did exploratory data analysis using the R package "MuMIn." In doing so, we

fitted all possible models containing OD600 and respiration as the response variable and all

combinations of temperature, nutrients, ciliate presence, and their interactions as explanatory

variables. To disentangle the joint effect of nutrients, temperature, and ciliates on the response

variables, we used multi-model inference to measure the relative importance of each model term

using the package MuMIn version 1.47.1 [77]. This approach fits all possible models –i.e., models

considering all possible combinations of explanatory variables and their interactions–then ranks

them all by AIC to calculate the relative importance of each model term across all possible

models [77]. To assess whether total community biomass, diversity, and respiration rate changed

with experimental treatment, we used linear models with ciliate presence, nutrient levels (high or

low), and temperature (22 or 25°C) (and their possible interactions) as explanatory variables, and

either biomass, diversity, or respiration rates as response variables. Response variables were log-

transformed, while all explanatory variables were treated as discrete. All analyses were done in R

version 4.2.2 [78]. One of the 80 ciliate jars had no ciliates at the end of the experiment and was

discarded.

417   Prey Community Structure: Amplicon Sequencing and Processing

We used 16S rRNA amplicon sequencing to examine the impacts of temperature, nutrient level,

and ciliates on the community structure of the prey prokaryotic community. After 21 days of

incubation, we collected the prokaryotic communities by pelleting 1 ml from each microcosm

into 1.5 mL Eppendorf tubes and storing them at -20°C until DNA extraction. Genomic

microbiome DNA was isolated with an Omega Mag-Bind Environmental kit. Amplification and

preparation followed the Illumina 16S rRNA amplicon sequencing protocol with a custom

mixture of 515F and 806R primers [79] for archaea/bacteria and 18SV4F and 18SV4R primers [80]

targeting the 18S region. Prokaryotic sequences were processed with the QIIME 2 v 2021.2

platform [81]. Paired sequences were demultiplexed with the plugin demux and quality filtered

(denoised, dereplicated, chimera filtered, and pair-end merged) and processed in Amplicon

Sequence Variants (ASVs) with the DADA2 plugin [82]. Taxonomy was assigned using a pre-

trained Naive Bayes classifier based on the Silva database (version 138) trimmed to the

515F/806R primer pair (16S rRNA). Sequence variant-based alpha diversity (Shannon) and beta

diversity (Bray Curtis distance) were calculated with the phyloseq package [83]. Alpha diversity

was calculated from the analysis of variance tests of the relationship between alpha diversity and

moss species. Beta diversity was calculated from a linear model test of the relationship between

Bray Curtis distance and site.

Ciliate Responses

The ciliate community may itself respond to the imposed nutrients and temperature treatments

and understanding that response is crucial to shedding light on the direct and indirect effects of

temperature and nutrients on the prokaryotic community. To understand how the predator-ciliate

communities may have been affected by the treatments, we estimated ciliate population sizes at

the end of the experiment through fluid imaging of 1 mL subsamples of all microcosms using a

FlowCam (Fluid Imaging Technologies, Scarborough, ME, United States). We used linear

models to assess whether ciliate population counts changed with either temperature or nutrient

level. Shannon diversity was calculated using the diversity() function in the R package "vegan"

(v2.6-4;[84]). Using principal components analysis on abundance data, we assessed whether the

overall ciliate community structure responded to temperature and nutrient levels. We tested for

individual and interactive treatment effects with a perMANOVA with the adonis2() function in

the R package "vegan" (v2.6-4;[84]).

Direct and indirect influences on microbial communities using Structural Equation

Modeling

Last, we assessed the possibility of joint direct effects of temperature and nutrients on the

prokaryote community and indirect effects of temperature and nutrients through their effects on

the ciliates, using Structural Equation Models we fitted piecewiseSEM version 2.3.0 [85].

Compared with the traditional variance covariance-based SEM, the piecewise SEM (1) pieces

together multiple separate (generalized) linear models into a single causal network, (2) uses

Shipley's test of d-separation to test whether any paths are missing from the model, and (3) uses

Akaike information criterion (AIC) to compare nested models. We then automatically calculated

direct, indirect, and total effects for piecewise structural equation models using the package

semEff version 0.6.1 [86] in R. We checked the model fit using Fisher's C test at the 0.05 level

where the p-value needs to be larger than 0.05 for the model to describe the data appropriately.

Data Availability

The sequence data has been deposited in GenBank SRA under accession PRJNA1095004;

https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1095004. All other data and code have been

deposited in GitHub: https://github.com/kmd304/Predation-Mediates-TempXNut.git.

Acknowledgments

We thank Ze-Yi Han, Christopher Kilner, and Daniel J. Wieczynski for their support and

thoughtful discussions. This work was supported by the US Department of Energy, Office of

Science, Office of Biological and Environmental Research, Genomic Science Program Grant

award number DE-SC0020362, NSF DEB award number 2224819, and a Simons Foundation

Early Career Fellowship in Aquatic Microbial Ecology and Evolution number LS-ECIAMEE-

00001588 to JPG. Additional support for prokaryotic and a ciliate collection from SPRUCE, was

supported by US Department of Energy (DOE), Grant/Award Number: DE-AC05–00OR22725

to DJW.

References:

1. Alto, B. W. & Juliano, S. A. Precipitation and Temperature Effects on Populations of Aedes albopictus (Diptera: Culicidae): Implications for Range Expansion. *J Med Entomol* **38**, 646–656 (2001).

2. Bærum, K. M., Finstad, A. G., Ulvan, E. M. & Haugen, T. O. Population consequences of climate change through effects on functional traits of lentic brown trout in the sub-Arctic. *Sci Rep* **11**, 15246 (2021).

3. Zander, A., Bersier, L.-F. & Gray, S. M. Effects of temperature variability on community structure in a natural microbial food web. *Glob Chang Biol* **23**, 56–67 (2017).

4. Baron, J. S. *et al.* Effects of Land Cover, Water Redistribution, and Temperature on Ecosystem Processes in the South Platte Basin. *Ecological Applications* **8**, 1037–1051 (1998).

5. Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. & Charnov, E. L. Effects of size and temperature on metabolic rate. *Science* **293**, 2248–2251 (2001).

6. DeLong, J. P. *et al.* The combined effects of reactant kinetics and enzyme stability explain the temperature dependence of metabolic rates. *Ecology and Evolution* **7**, 3940–3950 (2017).

7. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a Metabolic Theory of Ecology. *Ecology* **85**, 1771–1789 (2004).

8. West, G. B. & Brown, J. H. The origin of allometric scaling laws in biology from genomes to ecosystems: towards a quantitative unifying theory of biological structure and organization. *J Exp Biol* **208**, 1575–1592 (2005).

9. Price, C. A., Gilooly, J. F., Allen, A. P., Weitz, J. S. & Niklas, K. J. The metabolic theory of ecology: prospects and challenges for plant biology. *New Phytologist* **188**, 696–710 (2010).

10. Anderson, K. J., Allen, A. P., Gillooly, J. F. & Brown, J. H. Temperature-dependence of biomass accumulation rates during secondary succession. *Ecology Letters* **9**, 673–682 (2006).

11.   O'Connor, M. I. *et al.* Temperature control of larval dispersal and the implications for marine

ecology, evolution, and conservation. *Proc Natl Acad Sci U S A* **104**, 1266–1271 (2007).

12.   Padfield, D. *et al.* Metabolic compensation constrains the temperature dependence of gross

primary production. *Ecology Letters* **20**, 1250–1260 (2017).

13.   Iversen, C. M. *et al.* Whole-Ecosystem Warming Increases Plant-Available Nitrogen and Phosphorus

in an Ombrotrophic Bog. *Ecosystems* **26**, 86–113 (2023).

14.   Rosenzweig, M. L. Paradox of Enrichment: Destabilization of Exploitation Ecosystems in Ecological

Time. *Science* **171**, 385–387 (1971).

15.   Vitousek, P. M. *et al.* Human Alteration of the Global Nitrogen Cycle: Sources and Consequences.

Ecological Applications **7**, 737–750 (1997).

16.   Fussmann, G. F., Ellner, S. P., Shertzer, K. W. & Hairston Jr., N. G. Crossing the Hopf Bifurcation in a

Live Predator-Prey System. *Science* **290**, 1358–1360 (2000).

17.   Morris, R. J. Anthropogenic impacts on tropical forest biodiversity: a network structure and

ecosystem functioning perspective. *Philos Trans R Soc Lond B Biol Sci* **365**, 3709–3718 (2010).

18.   Hautier, Y. *et al.* Eutrophication weakens stabilizing effects of diversity in natural grasslands.

Nature **508**, 521–525 (2014).

19.   Guignard, M. S. *et al.* Impacts of Nitrogen and Phosphorus: From Genomes to Natural Ecosystems

and Agriculture. *Frontiers in Ecology and Evolution* **5**, (2017).

20.   McClelland, J. W. & Valiela, I. Changes in food web structure under the influence of increased

anthropogenic nitrogen inputs to estuaries. *Marine Ecology Progress Series* **168**, 259–271 (1998).

21.   Carlier, A. *et al.* Food web structure of two Mediterranean lagoons under varying degree of

eutrophication. *Journal of Sea Research* **60**, 264–275 (2008).

22.   Gibert, J. P. Temperature directly and indirectly influences food web structure. *Scientific Reports* **9**,

5312 (2019).

23.   Binzer, A., Guill, C., Brose, U. & Rall, B. C. The dynamics of food chains under climate change and

nutrient enrichment. *Philos Trans R Soc Lond B Biol Sci* **367**, 2935–2944 (2012).

24.   Cross, W. F., Hood, J. M., Benstead, J. P., Huryn, A. D. & Nelson, D. Interactions between

temperature and nutrients across levels of ecological organization. *Glob Change Biol* **21**, 1025–

1040 (2015).

25.   Binzer, A., Guill, C., Rall, B. C. & Brose, U. Interactive effects of warming, eutrophication and size

structure: impacts on biodiversity and food-web structure. *Global Change Biology* **22**, 220–227

(2016).

26.   Tatters, A. O. *et al.* Interactive effects of temperature, CO2 and nitrogen source on a coastal

California diatom assemblage. *Journal of Plankton Research* **40**, 151–164 (2018).

27.   Tabi, A., Petchey, O. L. & Pennekamp, F. Warming reduces the effects of enrichment on stability

and functioning across levels of organisation in an aquatic microbial ecosystem. *Ecology Letters* **22**,

1061–1071 (2019).

28.   Han, Z.-Y., Wieczynski, D. J., Yammine, A. & Gibert, J. P. Temperature and nutrients drive eco-

phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society B: Biological*

Sciences **290**, 20222263 (2023).

29.   Bar-On, Y. M., Phillips, R. & Milo, R. The biomass distribution on Earth. *Proceedings of the National*

Academy of Sciences **115**, 6506–6511 (2018).

30.   Foissner, W. & Berger, H. A user-friendly guide to the ciliates (Protozoa, Ciliophora) commonly

used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their

ecology. *Freshwater Biology* **35**, 375–482 (1996).

31.   Oliverio, A. M. *et al.* The global-scale distributions of soil protists and their contributions to

belowground systems. *Science Advances* **6**, eaax8787 (2020).

32.   Wang, L. Microbial control of the carbon cycle in the ocean. *National Science Review* **5**, 287–291

(2018).

33.   Jansson, J. K. & Hofmockel, K. S. Soil microbiomes and climate change. *Nat Rev Microbiol* **18**, 35–46

(2020).

34.   Cordone, G. *et al.* Green vs brown food web: Effects of habitat type on multidimensional stability

proxies for a highly-resolved Antarctic food web. *Food Webs* **25**, e00166 (2020).

35.   Mougi, A. Coupling of green and brown food webs and ecosystem stability. *Ecology and Evolution*

10, 9192–9199 (2020).

36.   Falkowski, P. *et al.* The global carbon cycle: a test of our knowledge of earth as a system. *Science*

290, 291–296 (2000).

37.   Guo, X. *et al.* Gene-informed decomposition model predicts lower soil carbon loss due to

persistent microbial adaptation to warming. *Nat Commun* **11**, 4897 (2020).

38.   Zhou, J. *et al.* Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate*

Change **2**, 106–110 (2012).

39.   Bradford, M. A. *et al.* Cross-biome patterns in soil microbial respiration predictable from

evolutionary theory on thermal adaptation. *Nat Ecol Evol* **3**, 223–231 (2019).

40.   Smith, T. P. *et al.* Community-level respiration of prokaryotic microbes may rise with global

warming. *Nat Commun* **10**, 5124 (2019).

41.   Wieczynski, D. J. *et al.* Linking species traits and demography to explain complex temperature

responses across levels of organization. *Proceedings of the National Academy of Sciences* **118**,

e2104863118 (2021).

42.   Montoya, J. M. & Raffaelli, D. Climate change, biotic interactions and ecosystem services. *Philos*

Trans R Soc Lond B Biol Sci **365**, 2013–2018 (2010).

43.   Bartley, T. J. *et al.* Food web rewiring in a changing world. *Nat Ecol Evol* **3**, 345–354 (2019).

44.   Barbour, M. A. & Gibert, J. P. Genetic and plastic rewiring of food webs under climate change.

Journal of Animal Ecology **90**, 1814–1830 (2021).

45.   Collins, C. G. *et al.* Global change re-structures alpine plant communities through interacting

abiotic and biotic effects. *Ecology Letters* **25**, 1813–1826 (2022).

46.   O'Connor, M. I., Piehler, M. F., Leech, D. M., Anton, A. & Bruno, J. F. Warming and Resource

Availability Shift Food Web Structure and Metabolism. *PLOS Biology* **7**, e1000178 (2009).

47.   Yvon-Durocher, G. & Allen, A. P. Linking community size structure and ecosystem functioning using

metabolic theory. *Philos Trans R Soc Lond B Biol Sci* **367**, 2998–3007 (2012).

48.   Baltar, F. *et al.* Marine bacterial community structure resilience to changes in protist predation

under phytoplankton bloom conditions. *ISME J* **10**, 568–581 (2016).

49.   Geisen, S., Hu, S., dela Cruz, T. E. E. & Veen, G. F. (Ciska). Protists as catalyzers of microbial litter

breakdown and carbon cycling at different temperature regimes. *ISME J* **15**, 618–621 (2021).

50.   Dell, A. I., Pawar, S. & Savage, V. M. Systematic variation in the temperature dependence of

physiological and ecological traits. *Proceedings of the National Academy of Sciences* **108**, 10591–

10596 (2011).

51.   Trap, J., Bonkowski, M., Plassard, C., Villenave, C. & Blanchart, E. Ecological importance of soil

bacterivores for ecosystem functions. *Plant Soil* **398**, 1–24 (2016).

52.   Geisen, S. *et al.* Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology Reviews*

42, 293–323 (2018).

53.   Rocca, J. D., Yammine, A., Simonin, M. & Gibert, J. P. Protist Predation Influences the Temperature

Response of Bacterial Communities. *Frontiers in Microbiology* **13**, (2022).

54.   Van der Putten, W. H., Macel, M. & Visser, M. E. Predicting species distribution and abundance

responses to climate change: why it is essential to include biotic interactions across trophic levels.

Philos Trans R Soc Lond B Biol Sci **365**, 2025–2034 (2010).

55.   Gao, Z., Karlsson, I., Geisen, S., Kowalchuk, G. & Jousset, A. Protists: Puppet Masters of the

Rhizosphere Microbiome. *Trends in Plant Science* **24**, 165–176 (2019).

56.   Wieczynski, D. J. *et al.* Viral infections likely mediate microbial controls on ecosystem responses to

global warming. *FEMS Microbiol Ecol* **99**, fiad016 (2023).

57.   Atkinson, D., Ciotti, B. J. & Montagnes, D. J. S. Protists decrease in size linearly with temperature:

ca. 2.5% degrees C(-1). *Proc Biol Sci* **270**, 2605–2611 (2003).

58.   DeLong, J. P. & Lyon, S. Temperature alters the shape of predator–prey cycles through effects on

underlying mechanisms. *PeerJ* **8**, e9377 (2020).

59.   Gibert, J. P., Wieczynski, D. J., Han, Z.-Y. & Yammine, A. Rapid eco-phenotypic feedback and the

temperature response of biomass dynamics. *Ecology and Evolution* **13**, e9685 (2023).

60.   Irwin, A. J., Finkel, Z. V., Schofield, O. M. E. & Falkowski, P. G. Scaling-up from nutrient physiology

to the size-structure of phytoplankton communities. *Journal of Plankton Research* **28**, 459–471

(2006).

61.   Marañón, E. *et al.* Unimodal size scaling of phytoplankton growth and the size dependence of

nutrient uptake and use. *Ecology Letters* **16**, 371–379 (2013).

62.   Gibert, J. P., Allen, R. L., Hruska, R. J. & DeLong, J. P. The ecological consequences of

environmentally induced phenotypic changes. *Ecol Lett* **20**, 997–1003 (2017).

63.   Pernthaler, J. Predation on prokaryotes in the water column and its ecological implications. *Nat

Rev Microbiol* **3**, 537–546 (2005).

64.   Thakur, M. P. *et al.* Resilience of rhizosphere microbial predators and their prey communities after

an extreme heat event. *Functional Ecology* **35**, 216–225 (2021).

65.   Crowther, T. W. & Bradford, M. A. Thermal acclimation in widespread heterotrophic soil microbes.

Ecol Lett **16**, 469–477 (2013).

66.   Ye, J.-S., Bradford, M. A., Maestre, F. T., Li, F.-M. & García-Palacios, P. Compensatory Thermal

Adaptation of Soil Microbial Respiration Rates in Global Croplands. *Global Biogeochemical Cycles*

34, e2019GB006507 (2020).

67.   Hahn, M. W. & Höfle, M. G. Grazing of protozoa and its effect on populations of aquatic bacteria.

FEMS Microbiology Ecology **35**, 113–121 (2001).

68.   Glücksman, E., Bell, T., Griffiths, R. I. & Bass, D. Closely related protist strains have different grazing

impacts on natural bacterial communities. *Environ Microbiol* **12**, 3105–3113 (2010).

69.   Han, Z.-Y., Wieczynski, D. J., Yammine, A. & Gibert, J. P. Temperature and nutrients drive eco-

phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society B: Biological*

Sciences **290**, 20222263 (2023).

70.   Kilner, C. L. *et al. Climate Change Factors Interact to Shift Functional Microbial Composition in*

Peatlands. http://biorxiv.org/lookup/doi/10.1101/2023.03.06.531192 (2023)

doi:10.1101/2023.03.06.531192.

71.   Hanson, P. J. *et al.* Attaining whole-ecosystem warming using air and deep-soil heating methods

with an elevated CO_2 atmosphere. *Biogeosciences* **14**, 861–883 (2017).

72.   Altermatt, F. *et al.* Big answers from small worlds: a user's guide for protist microcosms as a model

system in ecology and evolution. *Methods in Ecology and Evolution* **6**, 218–231 (2015).

73.   Moreno, A. M., Matz, C., Kjelleberg, S. & Manefield, M. Identification of Ciliate Grazers of

Autotrophic Bacteria in Ammonia-Oxidizing Activated Sludge by RNA Stable Isotope Probing.

Applied and Environmental Microbiology **76**, 2203–2211 (2010).

74.   Beal, J. *et al.* Robust estimation of bacterial cell count from optical density. *Commun Biol* **3**, 1–29

(2020).

75.   DeLong, J. P. & Vasseur, D. A. A dynamic explanation of size–density scaling in carnivores. *Ecology*

93, 470–476 (2012).

76.   Tomlinson, S. *et al.* Measuring metabolic rates of small terrestrial organisms by fluorescence-based

closed-system respirometry. *Journal of Experimental Biology* **221**, jeb172874 (2018).

77.   Barton, K. MuMIn: Multi-model inference. Version 1.43.17. (2020).

78.   R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing (2022).

79.   Dove, N. C. *et al.* Microbiome Variation Across Two Hemlock Species With Hemlock Woolly Adelgid

Infestation. *Frontiers in Microbiology* **11**, (2020).

80.   Bradley, I. M., Pinto, A. J. & Guest, J. S. Design and Evaluation of Illumina MiSeq-Compatible, 18S

rRNA Gene-Specific Primers for Improved Characterization of Mixed Phototrophic Communities.

Applied and Environmental Microbiology **82**, 5878–5891 (2016).

81.   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using

QIIME 2. *Nat Biotechnol* **37**, 852–857 (2019).

82.   Callahan, B. J. *et al.* DADA2: High resolution sample inference from Illumina amplicon data. *Nat*

Methods **13**, 581–583 (2016).

83.   McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLOS ONE* **8**, e61217 (2013).

84.   Oksanen, J. *et al.* vegan: Community Ecology Package. (2022).

85.   Lefcheck, J. S. piecewiseSEM: Piecewise structural equation modeling in R for ecology, evolution,

and systematics. *Methods in Ecology and Evolution* **7**, 573–579 (2016).

86.   Murphy, M. V. semEff: Automatic Calculation of Effects for Piecewise Structural Equation Models.

(2022).

87.   Carr, L. A., Gittman, R. K. & Bruno, J. F. Temperature Influences Herbivory and Algal Biomass in the

Galápagos Islands. *Frontiers in Marine Science* **5**, (2018).

88.   Moss, B. *et al.* Allied attack: climate change and eutrophication. *Inland Waters* **1**, 101–105 (2011).

89.   Trolle, D. *et al.* Projecting the future ecological state of lakes in Denmark in a 6 degree warming

scenario. *Climate Research* **64**, 55–72 (2015).

90.   Lürling, M., van Oosterhout, F. & Faassen, E. Eutrophication and Warming Boost Cyanobacterial

Biomass and Microcystins. *Toxins (Basel)* **9**, 64 (2017).

91.   Girkin, N. T. *et al.* Peat Properties, Dominant Vegetation Type and Microbial Community Structure

in a Tropical Peatland. *Wetlands* **40**, 1367–1377 (2020).

92.   AminiTabrizi, R. *et al.* Microbial sensitivity to temperature and sulfate deposition modulates

greenhouse gas emissions from peat soils. *Global Change Biology* **29**, 1951–1970 (2023).

93.   Petchey, O. L., McPhearson, P. T., Casey, T. M. & Morin, P. J. Environmental warming alters food-

web structure and ecosystem function. *Nature* **402**, 69–72 (1999).

94.   Gulis, V. & Suberkropp, K. Leaf litter decomposition and microbial activity in nutrient-enriched and

unaltered reaches of a headwater stream. *Freshwater Biology* **48**, 123–134 (2003).

95.   Tant, C. J. & Rosemond, A. D. Stream nutrient enrichment has a greater effect on coarse than on

fine benthic organic matter. *jnbs.1* **32**, 1111–1121 (2013).

96.   Brookes, J. D. & Carey, C. C. Ecology. Resilience to blooms. *Science* **334**, 46–47 (2011).

97.   Ostertag, R. Effects of Nitrogen and Phosphorus Availability on Fine-Root Dynamics in Hawaiian

Montane Forests. *Ecology* **82**, 485–499 (2001).

98.   Watson, E. B. *et al.* High nutrient loads amplify carbon cycling across California and New York

coastal wetlands but with ambiguous effects on marsh integrity and sustainability. *PLOS ONE* **17**,

e0273260 (2022).

99.   Kuzyakov, Y. & Mason-Jones, K. Viruses in soil: Nano-scale undead drivers of microbial life,

biogeochemical turnover and ecosystem functions. *Soil Biology and Biochemistry* **127**, 305–317

(2018).

100. Trubl, G. *et al.* Soil Viruses Are Underexplored Players in Ecosystem Carbon Processing. *mSystems*

3, e00076-18 (2018).

101. Jansson, J. K. & Wu, R. Soil viral diversity, ecology and climate change. *Nat Rev Microbiol* **21**, 296–

311 (2023).

102. Kaarlejärvi, E., Eskelinen, A. & Olofsson, J. Herbivores rescue diversity in warming tundra by

modulating trait-dependent species losses and gains. *Nat Commun* **8**, 419 (2017).

103. Easterling, D. R. *et al.* Climate Extremes: Observations, Modeling, and Impacts. *Science* **289**, 2068–

2074 (2000).

104. Rummukainen, M. Changes in climate and weather extremes in the 21st century. *WIREs Climate*

Change **3**, 115–129 (2012).

105. Dorrepaal, E. *et al.* Carbon respiration from subsurface peat accelerated by climate warming in the

subarctic. *Nature* **460**, 616–619 (2009).

106. Page, S. E. & Baird, A. J. Peatlands and Global Change: Response and Resilience. *Annual Review of*

Environment and Resources **41**, 35–57 (2016).

107. Hugelius, G. *et al.* Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw.

Proceedings of the National Academy of Sciences **117**, 20438–20446 (2020).

108. Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W. & Hunt, S. J. Global peatland dynamics since the

Last Glacial Maximum. *Geophysical Research Letters* **37**, (2010).

109. Kostka, J. E. *et al.* The Sphagnum microbiome: new insights from an ancient plant lineage. *New*

Phytologist **211**, 57–64 (2016).

110. Carrell, A. A. *et al.* Habitat-adapted microbial communities mediate Sphagnum peatmoss resilience

to warming. *New Phytologist* **234**, 2111–2125 (2022).

111. Carrell, A. A. *et al.* Experimental warming alters the community composition, diversity, and N2

fixation activity of peat moss (Sphagnum fallax) microbiomes. *Glob Chang Biol* **25**, 2993–3004

(2019).

112. Gill, A. L., Giasson, M.-A., Yu, R. & Finzi, A. C. Deep peat warming increases surface methane and

carbon dioxide emissions in a black spruce-dominated ombrotrophic bog. *Global Change Biology*

23, 5398–5411 (2017).

Figures

Figure 1. The independent and interactive effects of temperature, nutrients, and ciliate presence

on prokaryotic biomass. (A) Boxplots of the temperature, (B) the nutrient level, (C) presence of

ciliate community effects on prokaryotic biomass, (D) interaction between temperature, low

nutrient level, and ciliate presence on prokaryotic biomass, and (E) interaction between

temperature, high nutrient level, and ciliate presence on prokaryotic biomass. Lines within the

boxplots represent median, 25th, and 75th percentile values, while whiskers are defined by the

largest value not greater than 1.5× the interquartile range (IQR). Individual open circles represent

raw data points. Pink colors represent treatments with no ciliates, while purple represents those

that have the addition of ciliates.

761
Figure 2. The independent and interactive effects of temperature, nutrients, and ciliate presence

on prokaryotic respiration. (A) Boxplots of the temperature, (B) the nutrient level, (C) presence

of ciliate community effects on prokaryotic respiration, (D) interaction between temperature, low

nutrient level, and ciliate presence on prokaryotic respiration, and (E) interaction between

temperature, high nutrient level, and ciliate presence on prokaryotic respiration. Lines within the

boxplots represent median, 25th, and 75th percentile values, while whiskers are defined by the

largest value not greater than 1.5× the interquartile range (IQR). Individual open circles represent

raw data points. Pink colors represent treatments with no ciliates, while purple represents those

that have the addition of ciliates.

772
773

Figure 3. (A) Phyla relative abundances of prokaryotic (16S rRNA) prokaryotic communities

across treatments. (B) Boxplot of alpha diversity of prokaryotic communities across temperature,

nutrient level, and presence of ciliate community. Lines within the boxplots represent median,

25th, and 75th percentile values, while whiskers are defined by the largest value not greater than

1.5× the interquartile range (IQR). (C) NMDS ordination plot of Bray-Curtis dissimilarity

of prokaryotic community composition (16S rRNA). Pink colors represent treatments with no

ciliates, while purple represents those that have the addition of ciliates. Squares represent high

nutrient conditions, while triangles represent low nutrient conditions. If the shape is filled in with

color, it is experiencing warming (25°C), whereas those that are not filled in are at base

conditions (22°C).

785
Figure 4. (A) Final density data of ciliate communities for each sample across treatments. (B)

Boxplot of alpha diversity of ciliate communities across temperature and nutrient level. Lines

within the boxplots represent the median. (C) The ordination plot is based on a principal

components analysis of ciliate community composition. Arrows show dominant communities'

variable loadings within the functional trait space, while individual points show individual

samples and are colored by treatment. Individual open circles represent raw data points. Lines

within the boxplots represent median, 25th, and 75th percentile values, while whiskers are

defined by the largest value not greater than 1.5× the interquartile range (IQR).

Figure 5. Structural equation model showing proposed relationships between latent variables

temperature, nutrients, and their interaction on diversity and biomass of microbes and their

function. All numbers correspond to the standardized path coefficients. Solid arrows shown are

significant direct effects ($p < 0.05$). Red arrows represent a negative relationship, while green

shows positive.