Title: Predation by a ciliate community mediates temperature and nutrient effects on a peatland

prey prokaryotic community.

Katrina DeWitt[1*], Alyssa A. Carrell[3], Jennifer D. Rocca[1], Samantha Votzke[1], Andrea Yammine[1],

Ariane L. Peralta[2], David J. Weston[3], Dale A. Pelletier[3], Jean P. Gibert[1]

[1]Department of Biology, Duke University, Durham, NC, USA

[2]Department of Biology, East Carolina University, Greenville, NC, USA

[3]Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

Running Title: Predation and temperature affect bacterial communities.

#Address correspondence to Katrina DeWitt, katrina.mariedewitt@gmail.com

*Present address: Samantha Votzke, Department of Earth and Planetary Sciences, Johns Hopkins

University, Baltimore, MD, USA

Abstract word count: 173

Importance word count: 108

Main text word count: 4966

Keywords: Ciliates, Protoctists, Climate Change, Synthetic Communities, Microbial Food Webs

Abstract:

Temperature significantly impacts microbial communities' composition and function, which plays a vital role in the global carbon cycle that determines climate change. Nutrient influxes often accompany rising temperatures due to human activity. While ecological interactions between different microorganisms could shape their response to environmental change, we do not understand how predation may influence these responses in a warmer and increasingly nutrient-rich world. Here, we assess whether predation by a ciliate community of bacterial consumers influences changes in the diversity, biomass, and function of a freshwater prokaryotic community under different temperature and nutrient conditions. We found that predator presence mediates the effects of temperature and nutrients on total prokaryotic community biomass and composition through various mechanisms, including direct and indirect effects. However, the total community function was resilient. Our study supports previous findings that temperature and nutrients are essential drivers of microbial community composition and function but also demonstrates how predation can mediate these effects, indicating that the biotic context is as important as the abiotic context to understanding microbial responses to novel climates.

Importance:

While the importance of the abiotic environment in microbial communities has long been acknowledged, how prevalent ecological interactions like predation may influence these microbial community responses to shifting abiotic conditions is largely unknown. Our study addresses the complex interplay between temperature, nutrients, predation, and their joint effects on microbial community diversity and function. Our findings suggest that while temperature and nutrients are fundamental drivers of microbial community dynamics, the presence of predators significantly alters these responses. Our study underscores the impact of abiotic factors on microbial communities and the importance of accounting for the biotic context in which these occur to understand, let alone predict, these responses properly.

Introduction:

Rising global temperatures are currently affecting populations[1,2], communities[3], and ecosystems[4]

rising organismal metabolic rates[5,6], leading to higher energetic demands from populations[7–9],

and cascading effects within communities and ecosystems[5,10–12]. Additionally, human activity is

increasing nutrient loads and mineralization rates worldwide[13], which also affects communities

and ecosystems[14–19]. While the independent effects of temperature and nutrients are relatively

well-understood[20–22], their joint increase can interactively influence communities[23–28], and these

interactions can be hard to both understand and predict. Communities of prokaryotes, in

particular, comprise upwards of 14% of all existing biomass on Earth[29] are present on all

continents and ecosystems[30,31], and play a central role in the global carbon and nutrient cycles[32–35]. As with other organisms, prokaryotic respiration often increases with temperature[5–7,36–39],

possibly leading to a "warming begets warming" scenario. Increasing nutrient loads can also

increase heterotrophy and respiration[14,15] in these communities. However, how rising

temperature and nutrients might jointly influence prokaryotes in an increasingly warm, nutrient-

rich, and human-dominated world is unclear.

Prokaryotic communities might also respond to the rapid rewiring[40–42] of the broader food

webs they are a part of[43]. Indeed, bacterivores are predicted to increase foraging rates to offset

increased metabolic costs[44], possibly resulting in decreased prokaryotic biomass with warming[45–48], which could, in turn, impact the composition and function of prokaryotic communities in

future climates[48–52]. Ciliate microbes are among the most abundant predators of prokaryotes

worldwide[29,30,53,54]. These microbial predators show strong temperature[39,55–57] and nutrient

responses[28,58–60], temperature-dependent population dynamics[39,57] and feeding interactions[6,56],

and predation by a single ciliate has been shown to influence prokaryotic community structure,

93   dynamics, and function[61,56,62,28]. Prokaryotic communities will interact with ciliate communities

94   in nature –not just a single species– but how the presence of a ciliate community might influence

95   the response to joint changes in temperature and nutrients is not known.

96   Here, we examine whether and how the presence of a ciliate community on a mostly

prokaryotic prey community influences how nutrient additions and temperature affect total

microbial biomass, diversity, composition, and function. To do so, we used experimental

microcosms containing a synthetic community of naturally co-occurring ciliates[63] and a peatland

prokaryotic community to address: 1) How do temperature, nutrients, and the presence of a

predatory ciliate community jointly affect prokaryotic community biomass, diversity, and

composition? 2) What are the consequences for total microbial community respiration (i.e.,

function)? However, changes in the prey prokaryotic community might give feedback to the

ciliates, and the ciliates might also directly respond to temperature and nutrients. So, we also ask:

3) How do temperature and nutrients affect the ciliates? We combined these answers to provide

a perspective on how prokaryotes respond directly to temperature and nutrients and indirectly

through ciliate responses to those same factors and their resulting impacts on the prokaryotes.

108   We hypothesize that prokaryotic biomass and total community respiration rates should

increase with temperatures and nutrients[64,65]. Predation by a single ciliate decreases prokaryotic

biomass[66,67], so predation by the ciliate community should also lead to lower prokaryotic

biomass, resulting in lower total respiration rates (prokaryotes + ciliates)[45]. However, ciliate

presence can sometimes facilitate prokaryotic growth by mobilizing otherwise inaccessible

resources (e.g., fertilization)[68,69]. Alternatively, the presence of a ciliate community could

increase prokaryotic biomass and total community respiration rates. Ciliate presence should

result in a compositional shift among the prokaryotes due to the differential consumption or

facilitation[66,67]. Last, predation by a single ciliate species can interact with temperature and

nutrients, to co-determine total respiration and biomass responses of the prokaryotes[52], and so we

expect that an entire community of ciliates might have similar effects. Because, the ciliates can

themselves show ecological[28] and phenotypic[70,57] responses to temperature and nutrients[71,72], we

hypothesize that the prokaryotic community likely will respond to both direct, and indirect

effects of temperature and nutrients, the later mediated by the ciliate direct responses to

temperature and nutrient change.

Results

Direct effects of temperature, nutrients, and ciliate community on prokaryotic biomass and

total community respiration

We conducted a fully factorial microcosm experiment manipulating temperature (22°C/25°C),

nutrient availability (low/high), and the presence of an 8-species synthetic ciliate community

known to co-occur in nature. After three weeks, we quantified prokaryotic biomass via

spectrophotometry (OD600) and total community respiration using real-time respirometry (see

Methods).

Contrary to what was observed in mono-ciliate treatments[52], the same approach we used,

we observed an increase in prokaryotic biomass in the presence of the ciliate community (effect

$= 0.151 \pm 0.056$ SE, $t = 2.677$, $df = 111$, $p = 0.009$; Figure 1A). This effect was temperature-

dependent, as indicated by a significant negative interaction between temperature and ciliate

presence (interaction $= -0.0067 \pm 0.0024$ SE, $t = -2.783$, $df = 111$, $p = 0.006$; Figure 1B, 1C).

Additionally, ciliates and nutrients interacted: under low nutrient conditions, the presence of

ciliates reduced prokaryotic biomass (interaction $= -0.206 \pm 0.080$ SE, $t = -2.577$, $df = 111$, $p =$

0.011; Figure 1D, 1E). Finally, we detected a significant three-way interaction between

temperature, nutrient level, and ciliate presence (interaction = 0.0092 ± 0.0034 SE, t = 2.700, df

= 111, p = 0.008), suggesting that the temperature effect on biomass depends on both nutrient

availability and protist presence (Figure 1B, 1C). In contrast, the main effects of temperature (p =

0.986) and nutrient treatment (p = 0.787) were not significant.

The ciliate community did not influence total community respiration (effect = 9.601×10^{-5}

$\pm 1.013 \times 10^{-4}$ SE, t-value = 0.948, df = 112, p-value = 0.345, Fig 2). However, a weak two-

way interaction between temperature and nutrients led to decreased respiration under high

nutrient conditions at 25°C relative to other treatments (effect = 0.000331 ± 0.000165 SE, t-value

= 2.004, df = 112, p-value = 0.048, Fig 2B, 2C), and no significant effect of ciliates presence

(Fig 2B, 2C).

Direct effects of temperature, nutrients, and ciliate presence on the prokaryotic prey

community diversity and composition

We quantified shifts in the Shannon diversity and composition of the prokaryotic community

across treatments through 16S rRNA amplicon sequencing.

Proteobacteria were the most abundant group, followed by Firmicutes and Cyanobacteria

(Figure 3A). Shannon diversity responded to a two-way interaction between temperature and

ciliates (ANOVA, F = 7.3073, p-value = 0.007), as well as a three-way interaction between

temperature, nutrients, and ciliates (ANOVA, F = 13.449, p-value = 3.76×10^{-4}, Figure 3B) such

that Shannon diversity increased in the presence of ciliates, but only under low nutrients and high

temperatures (Figure 3B), leading to significant reduction in Firmicutes under those conditions

relative to all other scenarios (Figure 3A).

The composition of the prokaryotic prey community changed significantly due to two-

way and three-way interactions between temperature, nutrients, and ciliate presence (Fig 3C).

First, temperature increases species turnover under low nutrients but stabilizes it under high

nutrient conditions (PERMANOVA, F = 2.491, p-value = 0.002, Fig 3C). Second, temperature

amplified ciliate-driven effects on prokaryotic prey community composition (PERMANOVA, F

= 2.056, p-value = 0.010, Fig 3C). Third, ciliates promoted higher species turnover under high

nutrients than low ones (PERMANOVA, F = 1.970, p-value = 0.010, Fig 3C). Last, the ciliates

strongly modulated the combined effects of temperature and nutrients on prokaryotic prey

community composition (PERMANOVA, F = 2.336, p-value = 0.004, Figure 3C). For example,

ciliates enhanced species turnover under high nutrient conditions at elevated temperatures,

whereas their influence was weaker under low nutrient conditions regardless of temperature

(Figure 3C).

Direct effects of temperature and nutrients on the ciliate community

While temperature and nutrients directly influenced the prokaryotic community, ciliate presence

also influenced their effect on prokaryotic biomass, diversity, and composition, but not total

respiration. However, if the ciliates themselves respond to temperature and nutrients, the

prokaryotic community could be responding indirectly to these ciliate shifts instead of simply

showing different temperature and nutrient responses in their presence. To understand this, we

first ask whether the ciliate communities respond to nutrients and temperatures by quantifying

changes in ciliate density, diversity, and composition (see Methods).

Nutrients led to higher ciliate densities across all species, but this effect decreased with

temperature (temperature: effect = 0.389 ± 0.192 SE, t-value = 2.025, p-value = 0.046; nutrient:

effect = 0.865 ± 0.192 SE, t-value = 4.050, p-value = 2.4×10^{-5}; two-way interaction effect = -

0.710 ± 0.271 SE, t-value = -2.613, df = 74, p-value = 0.010, Fig 4A).

Blepharisma sp. was the dominant ciliate species by density, followed by *Colpidium* sp.

(Figure 4B); two species, *Tetrahymena pyriformis,* and *Cyclidium glaucoma,* were not present in

any samples, most likely due to competition –or predation– by other ciliates (e.g., *Blepharisma*

sp.). Ciliate Shannon diversity decreased with nutrients (effect = -0.229 ± 0.090 SE, t-value = -

2.542, df = 75, p-value = 0.013), and so did evenness (effect = -0.104 ± 0.041 SE, t-value = -

2.542, df = 75, p-value = 0.013), regardless of temperature (Fig 4C).

Ciliate communities shifted significantly in composition across temperature and nutrient

treatments such that four distinct community configurations were possible—each corresponding

to a principal component in PCA space: *Colpidium* sp.–*Paramecium bursaria*–*Paramecium*

aurelia–*Glaucoma* sp. (PC1, Figure 4D), *Blepharisma* sp.–*Colpoda steinii* (PC2, Figure 4D),

Halteria grandinella–*Strombidium* sp. (PC3, Figure 4D), *Tillina magna* (PC4, Figure 4D).

Composition was significantly affected by nutrients (PERMANOVA, F = 2.144, p-value =

0.021) and the interaction between nutrients and temperature (PERMANOVA, F = 2.022, p-

value = 0.039).

Direct and indirect effects of temperature and nutrients on the total microbial community

structure and function

Having shown that temperature and nutrients directly affect the prokaryotic prey and ciliate

predator community and that the presence of the ciliate community seems to mediate how the

prey prokaryotic community responds to temperature and nutrients, we used a Structural

Equation Modeling approach to understand the direct and indirect effects of temperature on the

entirety of the microbial community. The best SEM model (model selection described in

Appendix 1) converged after 11 iterations, including 25 parameters, and showed good alignment

with our previous analyses while uncovering additional effects, both direct and indirect (Fig 5).

All measures of goodness-of-fit suggested that the model correctly described the data ($\chi^2(10) =$

4.61, p-value = 0.990; CFI = 1.000; TLI = 1.194; RMSEA = 0.000, SRMR = 0.041).

In the best model, multiple regression paths were statistically significant, including a

positive effect of temperature on the Shannon diversity of the prey prokaryotic community ($\beta =$

0.213, p = 0.036), a negative effect of nutrients on the Shannon Diversity of the predator ciliate

community ($\beta = -0.279$, p = 0.009), and a negative effect of the interaction between nutrients and

temperature on the prey prokaryotic diversity ($\beta = -0.373$, p < 0.001, Fig 5). Their diversity

significantly influenced the composition of the predator ciliate and prey prokaryotic

communities: prokaryotic diversity negatively influenced prokaryotic compositional change ($\beta =$

-0.414, p < 0.001), while ciliate diversity positively influenced ciliate community compositional

change ($\beta = 0.58$, p < 0.001). Nutrients led to ciliate community compositional change ($\beta =$

0.244, p = 0.024), while ciliate diversity declined with temperature ($\beta = -0.210$, p = 0.052). Last,

the ciliate community composition drove prokaryotic composition ($\beta = 0.375$, p = 0.022),

supporting all other analyses presented thus far (Fig 5).

Total microbial biomass –prokaryotic and ciliate– declined with temperature ($\beta = -0.51$, p

= 0.001) and increased with nutrients ($\beta = 0.50$, p < 0.001). Prokaryotic diversity positively

influenced total biomass ($\beta = 0.34$, p = 0.022), as did ciliate community composition ($\beta = 0.464$,

p = 0.002) and prokaryotic community composition ($\beta = 0.471$, p = 0.003). A significant

Temperature and Nutrient interaction ($\beta = -0.58$, p < 0.001) revealed that nutrient availability

moderates the negative effect of temperature on total microbial biomass.

As opposed to biomass, total community respiration was influenced by fewer factors:

nutrients led to an increase in total respiration ($\beta = 0.382$, $p < 0.001$), and so did ciliate diversity

($\beta = 0.300$, $p = 0.004$), suggesting that total respiration is more robust to shifts in the biotic and

abiotic environment than composition or biomass. Last, total respiration was also negatively

affected by nutrients indirectly through the diversity of the ciliates ($\beta = -0.084$, $p = 0.052$).

Discussion

Understanding how temperature and nutrients influence microbial structure, biomass, and

function is important in a warming world[35]. Our study shows how temperature and nutrients

determine prokaryotic community biomass, composition, and respiration rates in the presence

and absence of a ciliate community (Figures 1, 2, 3) and that the ciliate community also responds

to these abiotic conditions (Figure 4). We then disentangle the direct and indirect effects of

temperature and nutrients on the structure and function of the overall microbial community

through their differential impacts on the diversity and composition of prokaryotes and ciliates,

their biomass, and total respiration rates (Figure 5). Our results highlight the importance of

ecological interactions in shaping prokaryotic community responses to environmental change.

We showed that elevated temperatures and nutrients reduced prokaryotic biomass

(Figures 1B, 1C), aligning with observations in non-prokaryotic taxa across various

ecosystems[45,73]. The presence of ciliates mediated these effects, likely due to selective predation

on certain bacterial taxa[52]. While rising nutrient levels are expected to increase microbial

biomass under high temperatures[27,74,75], our findings do not support that trend, maybe owing to

enhanced prokaryotic communities' ability to maintain biomass despite nutrient fluctuations[76], or

compositional shifts, which were driven by interactive effects of temperature, nutrients, and

ciliate presence (Figure 3C). Interestingly, ciliate presence drove species turnover in nutrient-rich

environments (Figure 3C), which might be required for prokaryotic communities to maintain

biomass production and function under shifting abiotic conditions, maybe resulting in a ciliate

"rescue" effect of production and diversity under adverse conditions.

Total community respiration rates increased with temperature –consistent with theoretical

expectations[77,5,7,46,78,38]– but only under high nutrient levels (Figures 2B and 2C), and showed no

effect of ciliate presence (Figure 2A). This is striking, considering that predation by a single

ciliate species was shown to decrease respiration under elevated temperatures[52]. Thus, the effects

of a few species compared to entire ciliate communities can drastically affect their prokaryotic

prey and their response to environmental change. While rising temperatures are expected to

increase microbial respiration through increased heterotrophy[5,64,54], our results suggest that may

not always be true. This issue compounds with the fact that while ciliates showed density (Figure

4A, 4B), diversity (Figure 4C), compositional (Figure 4D), and biomass (Appendix 2) shifts with

temperature and nutrients interactions consistent with laboratory[39] and whole-ecosystem

warming studies[72], so neither nutrients nor temperature alone can fully explain the complex shifts

in ciliate predator communities, possibly obscuring our ability to predict the fate of the entire

microbial community under novel climates.

Our SEM approach indicated the direct and indirect effects of temperature and nutrients

on the structure and function of the different components of the microbial community (Figure 5).

For example, temperature directly reduces total microbial biomass. However, it also increases

prokaryotic diversity, which negatively affects prokaryotic compositional variation from

microcosm to microcosm, which increases total microbial biomass, resulting in an indirect effect

that, while synergistic with its direct effect on total biomass, occurs through a completely

different –indirect– mechanism. Interestingly, strong positive effects of ciliate community

composition on prokaryotic compositional change highlight a possible role of top-down control

in these microbial dynamics. In non-microbial food webs, the loss of top predators often

mediates how the food web responds to abiotic shift conditions[79–81], which our results suggest

could also occur in our study system.

Accounting for both direct and indirect effects of abiotic conditions revealed that

nutrients, but not temperature, influenced total microbial respiration directly and indirectly,

through ciliate diversity and the effect of this diversity on respiration (Fig. 5). This highlights the

importance of accounting for indirect effects when understanding joint biotic and abiotic effects

on microbial function: indirect effects are likely to manifest as strong collinearity among

explanatory model variables in a linear model framework, which often results in one variable's

effect masking the other's (e.g., Fig. 2). Last, the conspicuous lack of explanatory variables for

total community respiration –e.g., compared to total community biomass (Fig. 5)– suggests some

level of robustness of respiration to shifts in community composition in response to

environmental change[64,65,52] and highlighting that functioning may be less vulnerable to

environmental change than composition[82,83].

Caveats. We did not track the dynamics of either community over time. Consequently,

our results represent a snapshot in time, and some effects reported here may be transient, limiting

our understanding of the underlying mechanisms and processes. As global temperatures rise,

seasonality and variability are also changing[84,85], but because we only manipulate mean

temperature, our results do not inform how microbial communities may respond to shifts in

temperature seasonality or variability and whether the effects of temperature variability are

distinct from those of mean temperature, remains an open question. While the results presented

are clear enough to suggest that these processes may also be at play in nature, the short-term

nature of our experimental manipulations and tightly controlled experimental setup inevitably

limit the possible scope of our inference.

Conclusions. Our results shed some light on how warming may affect carbon

sequestration in peatlands. Indeed, peatlands are exceptionally susceptible to future climate

change impacts[86–88]. Despite covering less than 3% of the Earth's surface, these ecosystems store

approximately 25%-30 % of the world's soil carbon as recalcitrant peat[89]. However, peat moss

growth in nutrient-poor peatlands depends on symbiotic interactions with prokaryotic associates

[90,91]. Shifts in prokaryotic community composition due to warming[92] may impact the activity of

moss symbionts and other important microbes, potentially leading to reductions in carbon

sequestration in these peatlands through changes in moss growth[93]. Our results highlight that to

fully understand the breadth and consequences of these changes, we need to consider the

interactive effects of rising temperatures and increasing nutrient loads in these rapidly changing

ecosystems. Moreover, our results suggest that essential but largely overlooked predatory

interactions between these organisms and their predators may be important to understanding and

predicting how climate change may affect the responses of these microbial communities in

peatlands[13]. Still, total respiration levels might be influenced by a much narrower set of biotic

and abiotic variables.

Materials and Methods:

Prey Prokaryotic Community

We isolated a prokaryotic prey community from a peatland bog in the Marcell experimental

forest (Minnesota, USA), next to the Spruce and Peatland Responses Under Changing

Environment whole-ecosystem warming experiment[94] from 5 cm core samples containing a top

layer of living *Sphagnum* moss tissue and a lower layer of peat. We filtered out larger fungi and

ciliates using 11μm pore size Whatman autoclaved filters, then removed smaller flagellates and

fungal spores[48] using sterile 1.6 μm pore size Whatman GF/A filters. While the community is not

guaranteed to only have prokaryotes (e.g., eukaryotic nanoflagellates cannot be filtered out using

this approach), it should be composed in its majority by Prokaryotes and Archaea[52]. From now

on, we call it the *prokaryotic* community.

Experimental Treatments

The prokaryotic community was homogenized and incubated in 120, 200mL acid-washed and

autoclaved borosilicate jars filled with 150 mL of Carolina Biological Ciliate culture medium

plus one autoclaved wheat seed as a carbon source[95]. Jars were assigned to nutrient to possible

nutrient treatments –low nutrients (75 mL of media and 75 mL of distilled water, 60 jars) and

high nutrients (150mL of media, 60 jars)– and one of two possible temperature treatments (60

jars at 22°C, representing ambient temperature, and 60 jars at 25°C, representing an increase in

the average temperature of 3ºC. There was no observable difference in pH between low and high

nutrient treatments (pH measured using a Mettler Toledo SevenGo Duo SG68 pH/Ion/DO

Meter). We added one autoclaved wheat seed (~35 mg ea.) to high-nutrient treatments and half a

seed (~17.5 mg ea.) to the low-nutrient treatments[95,60], both from the same batch to control for

seed-associated microbes. All microcosms were incubated for three days in Percival AL-22 L2

growth chambers (Percival Scientific, Perry, Iowa) at 10% light intensity (1700 lux), 75%

humidity, and a 16:8 hr. day-night cycle before further manipulation.

After three days, we added a synthetic ciliate community to 80 jars across all nutrient and

temperature treatments, leaving 40 no-ciliate jars equally distributed among temperature and

nutrient manipulations. This asymmetric design ensures the detection of even small effects in the

jars with ciliates. We included the following ciliates: *Tillina magna, Tetrahymena pyriformis,*

Cyclidium glaucoma, Colpoda steinii, Strombidium sp.*, Blepharisma* sp.*, Glaucoma* sp.*,*

Colpidium sp.*, Halteria grandinella, Paramecium aurelia,* and *Paramecium bursaria* all of

which naturally occur in *Sphagnum* peatlands[30], and came from long-term (> five years) lab

cultures[39]. Ciliates were introduced by pipetting well-mixed stock cultures at carrying capacity

into experimental microcosms. Each species was introduced at a density of at least 17 individuals

per jar (see Supplementary Table S6; Appendix 3 for initial densities). To control for possible

effects of other prokaryotes in the ciliate cultures, we added filtered, homogenized ciliate growth

medium to all "non-ciliate" jars, matching the amounts used in ciliate jars, as in previous

studies[52,61,96]. The filtered media was inspected under the microscope to confirm the absence of

ciliates before use, then confirmed again through microscopy and 18S sequencing post-

experiment (no contamination was found). However, 18S sequencing revealed that one of the 80

ciliate jars had no ciliates by day 21 and was discarded for analysis.

Every jar was thus assigned to one of eight possible treatments –with 10 replicates for no

ciliate jars, and 20 replicates for jars with ciliates: (1) 22°C, no ciliates, low nutrient; (2) 22°C,

no ciliates, high nutrient; (3) 25°C, no ciliates, low nutrient; (4) 25°C, no ciliates, high nutrient;

(5) 22°C, ciliates, low nutrient; (6) 22°C, ciliates, high nutrient; (7) 25°C, ciliates, low nutrient;

and (8) 25°C, ciliates, high nutrient.

Measurement and Analysis of Prokaryotic Biomass, Prey Community Diversity, and Total

Community Respiration Rate

Prokaryotic Biomass

After 21 days, we quantified prokaryotic biomass, prey community diversity, and total

community respiration rates. We measured the optical density at 600 nm wavelength (or OD600)

as a proxy for the prokaryotic biomass[97] using a BioTEK Epoch-2 spectrophotometer (Winooski,

VT, USA). Higher OD600 indicates greater prokaryotic biomass[97]. Ciliates can scatter some

light, but their contribution relative to that of the prokaryotes is much smaller[98,62], due to their

larger size and lower densities. Yet, OD600 could still reflect changes in ciliate biomass, even if

it mostly reflects changes in prokaryotic biomass[98,62]. Our analyses show that ciliate biomass

does a poor job at predicting OD600 ($R^2 = 0.14$; Appendix 4), with a substantial spread away

from the 1:1 line (SF1 Appendix 4), suggesting only a weak relationship between the two. To

further control this minor influence, we used the residuals from a linear model fit to raw OD600

values as a function of ciliate biomass as a corrected estimate of prokaryotic biomass. These

residuals (hereafter, prokaryotic biomass --OD600 residuals) were used in downstream models

testing treatment effects. Based on these analyses and consistent with prior work[98,62], we

interpret OD600 as a reliable proxy for prokaryotic biomass rather than total microbial biomass.

Prey community diversity and composition

To estimate prokaryotic community diversity and composition, we transferred 1mL from all

microcosms into 1.5mL Eppendorf tubes, pelleting and storing them at -80°C until DNA

extraction. Genomic microbiome DNA was isolated with an Omega Mag-Bind Environmental

kit. Amplification and preparation followed the Illumina 16S rRNA amplicon sequencing

protocol with a custom mixture of 515F and 806R primers[99] for archaea/bacteria and 18SV4F

and 18SV4R primers[100] targeting the 18S region. Prokaryotic sequences were processed with the

QIIME 2 v 2021.2 platform[101]. Paired sequences were demultiplexed with the plugin demux and

quality filtered (denoised, dereplicated, chimera filtered, and pair-end merged) and processed in

Amplicon Sequence Variants (ASVs) with the DADA2 plugin[102]. Taxonomy was assigned using

a pre-trained Naive Bayes classifier based on the Silva database (version 138) trimmed to the

515F/806R primer pair (16S rRNA). Sequence variant-based alpha (Shannon) and beta diversity

(Bray-Curtis distance) were calculated with the *phyloseq* package[103]. Shannon diversity was

calculated using the *diversity*() function in the R package "*vegan*" (v2.6-4;[104]). Beta diversity –

which measures compositional change across jars– was calculated from a linear model test of the

relationship between Bray-Curtis distance and jar.

Total Respiration rates

Total community respiration by day 21 was determined using optode-based real-time

respirometry (OXY-4 SMA, PreSens, Regensburg, Germany)[95,105] on the entire jar microcosm

(150 ml) for 30 minutes and a collection rate of one measurement every three seconds, after a 30-

minute acclimation period (n = 120) at their original experimental temperature and in the dark.

Respiration rates were estimated as the slope of the oxygen concentration over time (in μmol

O_2/min; Supplementary Figures S2, S3, and S4; Appendix 5)[106].

Analyses of Prokaryotic Biomass, Community Diversity and Composition, and Total

Respiration Rates

Brute-force exploratory data analysis was performed by fitting all possible models with

prokaryotic biomass (OD600) and respiration as response variables and all combinations of

temperature, nutrients, ciliate presence, and their interactions as explanatory variables, then did

multi-model inference to disentangle their joint effects –measured as the relative importance of

each model term across all possible models– using R package MuMIn (v1.47.1)[107]. This analysis

suggested that ciliate presence, temperature, and nutrient levels all interactively influenced

prokaryotic biomass, while nutrients were the most important predictor of total microbial

respiration, followed by the presence of ciliates and temperature (Supplementary Figures S6, S7;

Appendix 6).

To quantify the magnitude and direction of these effects, we used separate linear models

with ciliate presence, nutrient levels, temperature, and their two- and three-way interactions as

explanatory variables, for either prokaryotic biomass (OD600), prokaryotic diversity (richness

and Shannon diversity), or total respiration rates, as response variables. Additionally, we

assessed how temperature, nutrients, and ciliate presence influenced the composition of the

prokaryotic community using PerMANOVA on compositional data from Bray-Curtis. All

response variables were log-transformed, and explanatory variables were treated as categorical.

All analyses were done in R version 4.2.2[108].

Changes in the Ciliate Density, Biomass, Diversity, and Composition: Estimation and

Analyses

Because the ciliate community could potentially respond to the imposed treatments, we

tracked all ciliate species densities through fluid imaging of 1 mL subsamples of all microcosms

(FlowCam, Fluid Imaging Technologies, Scarborough, ME, United States) at day 21. The

FlowCam takes pictures of all censused cells and can use these pictures to estimate length, width,

area, volume and other properties, thus providing estimates of density, and allowing us to

calculate cell volume, mass, and hence, biomass. Ciliate biomass was calculated as the average

body mass of each ciliate species times its density (i.e., $\sum_{Species} N_i M_i$, where N is the density of

species i and M is its average mass). The average mass was estimated by using the volume of

each censused individual –i.e., volume of an ellipsoid, $V = \frac{4}{3}\pi \left(\frac{length}{2}\right)\left(\frac{width}{2}\right)^2$, in µm³, and

assuming that the two minor axes of each cell are the same. Individual cell volume was then

converted into cubic centimeters (cm³) and then multiplied by the density of water (1 g/cm³) to

individual cell mass estimates. These estimates were averaged within each species to obtain

average mass, then multiplied by each species' density and added across species to obtain ciliate

biomass density in g/mL.

Classification of individual cell images into different species was done manually, for

accuracy, but measurements were taken automatically by the FlowCam's proprietary software.

Shannon ciliate diversity was calculated using the *diversity*() function in the R package "*vegan*"

(v2.6-4;[104]). Mean dissimilarity, a measure of community composition, was quantified using the

Bray-Curtis distances among jars. Bray-Curtis dissimilarity was calculated using *vegdist* function

in R package "*vegan*". We then calculated mean dissimilarity row-wise: the resulting mean

dissimilarity values were appended to the original dataset and treated as a measure of community

average differences in community composition (or beta diversity), for each sample[109,110].

We used linear models to evaluate how ciliate density and diversity changed with

temperature, nutrients, or their interaction. We used Principal Components Analysis (PCA) on

the ciliate community density data across all species, then tested for individual and interactive

treatment effects on the first two principal components of such compositional data (PC1= 27%,

PC2= 15%) using perMANOVA as implemented the adonis2() function in "*vegan*".

Direct and Indirect Effects of Temperature and Nutrients

So far, we have separately addressed how prokaryotic communities respond to direct effects of

temperature and nutrients with and without ciliates, and how ciliates may also respond to direct

effects of these abiotic variables. However, it is unclear whether temperature and nutrients

indirectly affect prokaryotes via ciliate responses or whether and how microbial community

function (prokaryotes + ciliates) is driven by responses to temperature and nutrients by either

prokaryotes, ciliates, or both.

We address these questions by fitting alternative Structural Equation Models (SEMs) in R

package *lavaan* (version 0.6-18)[111]. The most complex model included the effects of

temperature, nutrients, and their interaction, on biotic variables, Shannon diversity and Bray-

Curtis composition for both prokaryotic and ciliate communities. The models also included the

effects of these abiotic and biotic variables on (1) total microbial biomass—defined as a

composite of prokaryotic biomass (OD600) and ciliate biomass—and (2) the joint effects of the

biotic and abiotic variables and total microbial biomass, on total community respiration (Table

S3, Appendix 1). We retained the best model by AIC and BIC (Appendix 1, Table S1-4).

Data Availability

477   The sequence data have been deposited in GenBank SRA under accession PRJNA1095004;

478   https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1095004. All other data and code have been

479   deposited in GitHub: https://github.com/kmd304/Predation-Mediates-TempXNut.git.

Acknowledgments

We thank Ze-Yi Han, Christopher Kilner, and Daniel J. Wieczynski for their feedback. This

work was supported by the US Department of Energy, Office of Science, Office of Biological

and Environmental Research, Genomic Science Program Grant award number DE-SC0020362,

NSF DEB award number 2224819, and a Simons Foundation Early Career Fellowship in Aquatic

Microbial Ecology and Evolution number LS-ECIAMEE-00001588 to JPG. Additional support

for prokaryotic and ciliate collection from SPRUCE, was supported by US Department of

Energy (DOE), Grant/Award Number: DE-AC05–00OR22725 to DJW.

References:

1. Alto, B. W. & Juliano, S. A. Precipitation and Temperature Effects on Populations of Aedes albopictus (Diptera: Culicidae): Implications for Range Expansion. *J Med Entomol* **38**, 646–656 (2001).

2. Bærum, K. M., Finstad, A. G., Ulvan, E. M. & Haugen, T. O. Population consequences of climate change through effects on functional traits of lentic brown trout in the sub-Arctic. *Sci Rep* **11**, 15246 (2021).

3. Zander, A., Bersier, L.-F. & Gray, S. M. Effects of temperature variability on community structure in a natural microbial food web. *Glob Chang Biol* **23**, 56–67 (2017).

4. Baron, J. S. *et al.* Effects of Land Cover, Water Redistribution, and Temperature on Ecosystem Processes in the South Platte Basin. *Ecological Applications* **8**, 1037–1051 (1998).

5. Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. & Charnov, E. L. Effects of size and temperature on metabolic rate. *Science* **293**, 2248–2251 (2001).

6.   DeLong, J. P. *et al.* The combined effects of reactant kinetics and enzyme stability explain the

temperature dependence of metabolic rates. *Ecology and Evolution* **7**, 3940–3950 (2017).

7.   Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a Metabolic Theory of

Ecology. *Ecology* **85**, 1771–1789 (2004).

8.   West, G. B. & Brown, J. H. The origin of allometric scaling laws in biology from genomes to

ecosystems: towards a quantitative unifying theory of biological structure and organization. *J Exp*

Biol **208**, 1575–1592 (2005).

9.   Price, C. A., Gilooly, J. F., Allen, A. P., Weitz, J. S. & Niklas, K. J. The metabolic theory of ecology:

prospects and challenges for plant biology. *New Phytologist* **188**, 696–710 (2010).

10.   Anderson, K. J., Allen, A. P., Gillooly, J. F. & Brown, J. H. Temperature-dependence of biomass

accumulation rates during secondary succession. *Ecology Letters* **9**, 673–682 (2006).

11.   O'Connor, M. I. *et al.* Temperature control of larval dispersal and the implications for marine

ecology, evolution, and conservation. *Proc Natl Acad Sci U S A* **104**, 1266–1271 (2007).

12.   Padfield, D. *et al.* Metabolic compensation constrains the temperature dependence of gross

primary production. *Ecology Letters* **20**, 1250–1260 (2017).

13.   Iversen, C. M. *et al.* Whole-Ecosystem Warming Increases Plant-Available Nitrogen and Phosphorus

in an Ombrotrophic Bog. *Ecosystems* **26**, 86–113 (2023).

14.   Rosenzweig, M. L. Paradox of Enrichment: Destabilization of Exploitation Ecosystems in Ecological

Time. *Science* **171**, 385–387 (1971).

15.   Vitousek, P. M. *et al.* Human Alteration of the Global Nitrogen Cycle: Sources and Consequences.

Ecological Applications **7**, 737–750 (1997).

16.   Fussmann, G. F., Ellner, S. P., Shertzer, K. W. & Hairston Jr., N. G. Crossing the Hopf Bifurcation in a

Live Predator-Prey System. *Science* **290**, 1358–1360 (2000).

17.   Morris, R. J. Anthropogenic impacts on tropical forest biodiversity: a network structure and

527   ecosystem functioning perspective. *Philos Trans R Soc Lond B Biol Sci* **365**, 3709–3718 (2010).

528   18.   Hautier, Y. *et al.* Eutrophication weakens stabilizing effects of diversity in natural grasslands.

529   *Nature* **508**, 521–525 (2014).

530   19.   Guignard, M. S. *et al.* Impacts of Nitrogen and Phosphorus: From Genomes to Natural Ecosystems

531   and Agriculture. *Frontiers in Ecology and Evolution* **5**, (2017).

532   20.   McClelland, J. W. & Valiela, I. Changes in food web structure under the influence of increased

533   anthropogenic nitrogen inputs to estuaries. *Marine Ecology Progress Series* **168**, 259–271 (1998).

534   21.   Carlier, A. *et al.* Food web structure of two Mediterranean lagoons under varying degree of

535   eutrophication. *Journal of Sea Research* **60**, 264–275 (2008).

536   22.   Gibert, J. P. Temperature directly and indirectly influences food web structure. *Scientific Reports* **9**,

537   5312 (2019).

538   23.   Binzer, A., Guill, C., Brose, U. & Rall, B. C. The dynamics of food chains under climate change and

539   nutrient enrichment. *Philos Trans R Soc Lond B Biol Sci* **367**, 2935–2944 (2012).

540   24.   Cross, W. F., Hood, J. M., Benstead, J. P., Huryn, A. D. & Nelson, D. Interactions between

541   temperature and nutrients across levels of ecological organization. *Glob Change Biol* **21**, 1025–

542   1040 (2015).

543   25.   Binzer, A., Guill, C., Rall, B. C. & Brose, U. Interactive effects of warming, eutrophication and size

544   structure: impacts on biodiversity and food-web structure. *Global Change Biology* **22**, 220–227

545   (2016).

546   26.   Tatters, A. O. *et al.* Interactive effects of temperature, CO_2 and nitrogen source on a coastal

547   California diatom assemblage. *Journal of Plankton Research* **40**, 151–164 (2018).

27. Tabi, A., Petchey, O. L. & Pennekamp, F. Warming reduces the effects of enrichment on stability

and functioning across levels of organisation in an aquatic microbial ecosystem. *Ecology Letters* **22**,

1061–1071 (2019).

28. Han, Z.-Y., Wieczynski, D. J., Yammine, A. & Gibert, J. P. Temperature and nutrients drive eco-

phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society B: Biological*

Sciences **290**, 20222263 (2023).

29. Bar-On, Y. M., Phillips, R. & Milo, R. The biomass distribution on Earth. *Proceedings of the National*

Academy of Sciences **115**, 6506–6511 (2018).

30. Foissner, W. & Berger, H. A user-friendly guide to the ciliates (Protozoa, Ciliophora) commonly

used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their

ecology. *Freshwater Biology* **35**, 375–482 (1996).

31. Oliverio, A. M. *et al.* The global-scale distributions of soil protists and their contributions to

belowground systems. *Science Advances* **6**, eaax8787 (2020).

32. Cordone, G. *et al.* Green vs brown food web: Effects of habitat type on multidimensional stability

proxies for a highly-resolved Antarctic food web. *Food Webs* **25**, e00166 (2020).

33. Mougi, A. Coupling of green and brown food webs and ecosystem stability. *Ecology and Evolution*

10, 9192–9199 (2020).

34. Wang, L. Microbial control of the carbon cycle in the ocean. *National Science Review* **5**, 287–291

(2018).

35. Jansson, J. K. & Hofmockel, K. S. Soil microbiomes and climate change. *Nat Rev Microbiol* **18**, 35–46

(2020).

36. Zhou, J. *et al.* Microbial mediation of carbon-cycle feedbacks to climate warming. *Nature Climate*

Change **2**, 106–110 (2012).

37.   Bradford, M. A. *et al.* Cross-biome patterns in soil microbial respiration predictable from

evolutionary theory on thermal adaptation. *Nat Ecol Evol* **3**, 223–231 (2019).

38.   Smith, T. P. *et al.* Community-level respiration of prokaryotic microbes may rise with global

warming. *Nat Commun* **10**, 5124 (2019).

39.   Wieczynski, D. J. *et al.* Linking species traits and demography to explain complex temperature

responses across levels of organization. *Proceedings of the National Academy of Sciences* **118**,

e2104863118 (2021).

40.   Montoya, J. M. & Raffaelli, D. Climate change, biotic interactions and ecosystem services. *Philos*

Trans R Soc Lond B Biol Sci **365**, 2013–2018 (2010).

41.   Bartley, T. J. *et al.* Food web rewiring in a changing world. *Nat Ecol Evol* **3**, 345–354 (2019).

42.   Barbour, M. A. & Gibert, J. P. Genetic and plastic rewiring of food webs under climate change.

Journal of Animal Ecology **90**, 1814–1830 (2021).

43.   Collins, C. G. *et al.* Global change re-structures alpine plant communities through interacting

abiotic and biotic effects. *Ecology Letters* **25**, 1813–1826 (2022).

44.   Dell, A. I., Pawar, S. & Savage, V. M. Systematic variation in the temperature dependence of

physiological and ecological traits. *Proceedings of the National Academy of Sciences* **108**, 10591–

10596 (2011).

45.   O'Connor, M. I., Piehler, M. F., Leech, D. M., Anton, A. & Bruno, J. F. Warming and Resource

Availability Shift Food Web Structure and Metabolism. *PLOS Biology* **7**, e1000178 (2009).

46.   Yvon-Durocher, G. & Allen, A. P. Linking community size structure and ecosystem functioning using

metabolic theory. *Philos Trans R Soc Lond B Biol Sci* **367**, 2998–3007 (2012).

47.   Baltar, F. *et al.* Marine bacterial community structure resilience to changes in protist predation

under phytoplankton bloom conditions. *ISME J* **10**, 568–581 (2016).

48.   Geisen, S., Hu, S., dela Cruz, T. E. E. & Veen, G. F. (Ciska). Protists as catalyzers of microbial litter

595   breakdown and carbon cycling at different temperature regimes. *ISME J* **15**, 618–621 (2021).

596   49.   Van der Putten, W. H., Macel, M. & Visser, M. E. Predicting species distribution and abundance

597   responses to climate change: why it is essential to include biotic interactions across trophic levels.

598   *Philos Trans R Soc Lond B Biol Sci* **365**, 2025–2034 (2010).

599   50.   Trap, J., Bonkowski, M., Plassard, C., Villenave, C. & Blanchart, E. Ecological importance of soil

600   bacterivores for ecosystem functions. *Plant Soil* **398**, 1–24 (2016).

601   51.   Geisen, S. *et al.* Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology Reviews*

602   **42**, 293–323 (2018).

603   52.   Rocca, J. D., Yammine, A., Simonin, M. & Gibert, J. P. Protist Predation Influences the Temperature

604   Response of Bacterial Communities. *Frontiers in Microbiology* **13**, (2022).

605   53.   Gao, Z., Karlsson, I., Geisen, S., Kowalchuk, G. & Jousset, A. Protists: Puppet Masters of the

606   Rhizosphere Microbiome. *Trends in Plant Science* **24**, 165–176 (2019).

607   54.   Wieczynski, D. J. *et al.* Viral infections likely mediate microbial controls on ecosystem responses to

608   global warming. *FEMS Microbiol Ecol* **99**, fiad016 (2023).

609   55.   Atkinson, D., Ciotti, B. J. & Montagnes, D. J. S. Protists decrease in size linearly with temperature:

610   ca. 2.5% degrees C(-1). *Proc Biol Sci* **270**, 2605–2611 (2003).

611   56.   DeLong, J. P. & Lyon, S. Temperature alters the shape of predator–prey cycles through effects on

612   underlying mechanisms. *PeerJ* **8**, e9377 (2020).

613   57.   Gibert, J. P., Wieczynski, D. J., Han, Z.-Y. & Yammine, A. Rapid eco-phenotypic feedback and the

614   temperature response of biomass dynamics. *Ecology and Evolution* **13**, e9685 (2023).

615   58.   Irwin, A. J., Finkel, Z. V., Schofield, O. M. E. & Falkowski, P. G. Scaling-up from nutrient physiology

616   to the size-structure of phytoplankton communities. *Journal of Plankton Research* **28**, 459–471

617   (2006).

59. Marañón, E. *et al.* Unimodal size scaling of phytoplankton growth and the size dependence of

nutrient uptake and use. *Ecology Letters* **16**, 371–379 (2013).

60. Gibert, J. P., Allen, R. L., Hruska, R. J. & DeLong, J. P. The ecological consequences of

environmentally induced phenotypic changes. *Ecol Lett* **20**, 997–1003 (2017).

61. Pernthaler, J. Predation on prokaryotes in the water column and its ecological implications. *Nat*

Rev Microbiol **3**, 537–546 (2005).

62. Thakur, M. P. *et al.* Resilience of rhizosphere microbial predators and their prey communities after

an extreme heat event. *Functional Ecology* **35**, 216–225 (2021).

63. *Protist Diversity and Geographical Distribution*. vol. 8 (Springer Netherlands, Dordrecht, 2009).

64. Crowther, T. W. & Bradford, M. A. Thermal acclimation in widespread heterotrophic soil microbes.

Ecol Lett **16**, 469–477 (2013).

65. Ye, J.-S., Bradford, M. A., Maestre, F. T., Li, F.-M. & García-Palacios, P. Compensatory Thermal

Adaptation of Soil Microbial Respiration Rates in Global Croplands. *Global Biogeochemical Cycles*

34, e2019GB006507 (2020).

66. Hahn, M. W. & Höfle, M. G. Grazing of protozoa and its effect on populations of aquatic bacteria.

FEMS Microbiology Ecology **35**, 113–121 (2001).

67. Glücksman, E., Bell, T., Griffiths, R. I. & Bass, D. Closely related protist strains have different grazing

impacts on natural bacterial communities. *Environ Microbiol* **12**, 3105–3113 (2010).

68. Jousset, A., Scheu, S. & Bonkowski, M. Secondary metabolite production facilitates establishment

of rhizobacteria by reducing both protozoan predation and the competitive effects of indigenous

bacteria. *Functional Ecology* **22**, 714–719 (2008).

69. Weidner, S., Latz, E., Agaras, B., Valverde, C. & Jousset, A. Protozoa stimulate the plant beneficial

activity of rhizospheric pseudomonads. *Plant Soil* **410**, 509–515 (2017).

70.   Gibert, J. P., Han, Z.-Y., Wieczynski, D. J., Votzke, S. & Yammine, A. Feedbacks between size and

642   density determine rapid eco-phenotypic dynamics. *Functional Ecology* **36**, 1668–1680 (2022).

643   71.   Han, Z.-Y., Wieczynski, D. J., Yammine, A. & Gibert, J. P. Temperature and nutrients drive eco-

644   phenotypic dynamics in a microbial food web. *Proceedings of the Royal Society B: Biological*

645   *Sciences* **290**, 20222263 (2023).

646   72.   Kilner, C. L. *et al. Climate Change Factors Interact to Shift Functional Microbial Composition in*

647   *Peatlands.* http://biorxiv.org/lookup/doi/10.1101/2023.03.06.531192 (2023)

648   doi:10.1101/2023.03.06.531192.

649   73.   Carr, L. A., Gittman, R. K. & Bruno, J. F. Temperature Influences Herbivory and Algal Biomass in the

650   Galápagos Islands. *Frontiers in Marine Science* **5**, (2018).

651   74.   Moss, B. *et al.* Allied attack: climate change and eutrophication. *Inland Waters* **1**, 101–105 (2011).

652   75.   Trolle, D. *et al.* Projecting the future ecological state of lakes in Denmark in a 6 degree warming

653   scenario. *Climate Research* **64**, 55–72 (2015).

654   76.   Lürling, M., van Oosterhout, F. & Faassen, E. Eutrophication and Warming Boost Cyanobacterial

655   Biomass and Microcystins. *Toxins (Basel)* **9**, 64 (2017).

656   77.   Petchey, O. L., McPhearson, P. T., Casey, T. M. & Morin, P. J. Environmental warming alters food-

657   web structure and ecosystem function. *Nature* **402**, 69–72 (1999).

658   78.   Tant, C. J. & Rosemond, A. D. Stream nutrient enrichment has a greater effect on coarse than on

659   fine benthic organic matter. *jnbs.1* **32**, 1111–1121 (2013).

660   79.   Rooney, N., McCann, K., Gellner, G. & Moore, J. C. Structural asymmetry and the stability of

661   diverse food webs. *Nature* **442**, 265–269 (2006).

662   80.   McCann, K. S. & Rooney, N. The more food webs change, the more they stay the same.

663   *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**, 1789–1801 (2009).

664   81.   Trzcinski, M. K. *et al.* The effects of food web structure on ecosystem function exceeds those of

665   precipitation. *Journal of Animal Ecology* **85**, 1147–1160 (2016).

666   82.   Louca, S. *et al.* Functional structure of the bromeliad tank microbiome is strongly shaped by local

667   geochemical conditions. *Environ. Microbiol.* **19**, 3132–3151 (2017).

668   83.   Hillebrand, H. & Kunze, C. Meta-analysis on pulse disturbances reveals differences in functional

669   and compositional recovery across ecosystems. *Ecology Letters* **23**, 575–585 (2020).

670   84.   Easterling, D. R. *et al.* Climate Extremes: Observations, Modeling, and Impacts. *Science* **289**, 2068–

671   2074 (2000).

672   85.   Rummukainen, M. Changes in climate and weather extremes in the 21st century. *WIREs Climate*

673   *Change* **3**, 115–129 (2012).

674   86.   Dorrepaal, E. *et al.* Carbon respiration from subsurface peat accelerated by climate warming in the

675   subarctic. *Nature* **460**, 616–619 (2009).

676   87.   Page, S. E. & Baird, A. J. Peatlands and Global Change: Response and Resilience. *Annual Review of*

677   *Environment and Resources* **41**, 35–57 (2016).

678   88.   Hugelius, G. *et al.* Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw.

679   *Proceedings of the National Academy of Sciences* **117**, 20438–20446 (2020).

680   89.   Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W. & Hunt, S. J. Global peatland dynamics since the

681   Last Glacial Maximum. *Geophysical Research Letters* **37**, (2010).

682   90.   Kostka, J. E. *et al.* The Sphagnum microbiome: new insights from an ancient plant lineage. *New*

683   *Phytologist* **211**, 57–64 (2016).

684   91.   Carrell, A. A. *et al.* Habitat-adapted microbial communities mediate Sphagnum peatmoss resilience

685   to warming. *New Phytologist* **234**, 2111–2125 (2022).

92.　Carrell, A. A. *et al.* Experimental warming alters the community composition, diversity, and N2

fixation activity of peat moss (Sphagnum fallax) microbiomes. *Glob Chang Biol* **25**, 2993–3004

(2019).

93.　Gill, A. L., Giasson, M.-A., Yu, R. & Finzi, A. C. Deep peat warming increases surface methane and

carbon dioxide emissions in a black spruce-dominated ombrotrophic bog. *Global Change Biology*

23, 5398–5411 (2017).

94.　Hanson, P. J. *et al.* Attaining whole-ecosystem warming using air and deep-soil heating methods

with an elevated CO_2 atmosphere. *Biogeosciences* **14**, 861–883 (2017).

95.　Altermatt, F. *et al.* Big answers from small worlds: a user's guide for protist microcosms as a model

system in ecology and evolution. *Methods in Ecology and Evolution* **6**, 218–231 (2015).

96.　Moreno, A. M., Matz, C., Kjelleberg, S. & Manefield, M. Identification of Ciliate Grazers of

Autotrophic Bacteria in Ammonia-Oxidizing Activated Sludge by RNA Stable Isotope Probing.

Applied and Environmental Microbiology **76**, 2203–2211 (2010).

97.　Beal, J. *et al.* Robust estimation of bacterial cell count from optical density. *Commun Biol* **3**, 1–29

(2020).

98.　Amacker, N. *et al.* Biocontrol Traits Correlate With Resistance to Predation by Protists in Soil

Pseudomonads. *Front. Microbiol.* **11**, (2020).

99.　Dove, N. C. *et al.* Microbiome Variation Across Two Hemlock Species With Hemlock Woolly Adelgid

Infestation. *Frontiers in Microbiology* **11**, (2020).

100.　Bradley, I. M., Pinto, A. J. & Guest, J. S. Design and Evaluation of Illumina MiSeq-Compatible, 18S

rRNA Gene-Specific Primers for Improved Characterization of Mixed Phototrophic Communities.

Applied and Environmental Microbiology **82**, 5878–5891 (2016).

101.　Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using

QIIME 2. *Nat Biotechnol* **37**, 852–857 (2019).

102. Callahan, B. J. *et al.* DADA2: High resolution sample inference from Illumina amplicon data. *Nat*

Methods **13**, 581–583 (2016).

103. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLOS ONE* **8**, e61217 (2013).

104. Oksanen, J. *et al.* vegan: Community Ecology Package. (2022).

105. DeLong, J. P. & Vasseur, D. A. A dynamic explanation of size–density scaling in carnivores. *Ecology*

93, 470–476 (2012).

106. Tomlinson, S. *et al.* Measuring metabolic rates of small terrestrial organisms by fluorescence-based

closed-system respirometry. *Journal of Experimental Biology* **221**, jeb172874 (2018).

107. Barton, K. MuMIn: Multi-model inference. Version 1.43.17. (2020).

108. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing (2022).

109. Collins, S. L. Fire Frequency and Community Heterogeneity in Tallgrass Prairie Vegetation. *Ecology*

73, 2001–2006 (1992).

110. Grace, J. B. & Keeley, J. E. A structural equation model analysis of postfire plant diversity in

California shrublands. *Ecol Appl* **16**, 503–514 (2006).

111. Rosseel, Y. lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical Software*

48, 1–36 (2012).

Figures

Figure 1. The independent and interactive effects of temperature, nutrients, and ciliate presence

on prokaryotic biomass. (A) The presence of a ciliate community affects prokaryotic biomass.

(B) Interaction between temperature, low nutrient levels, and ciliate presence. (C) Interaction

between temperature, high nutrient levels, and ciliate presence. Each boxplot displays the mean

(horizontal line), the 25th and 75th percentiles (box edges), and whiskers extending to the largest

value within 1.5× the interquartile range (IQR). Open circles represent individual raw data

points. Pink indicates treatments without ciliates; purple indicates treatments with ciliates.

Figure 2. The independent and interactive effects of temperature, nutrients, and ciliate presence

on total community respiration. (A) the presence of ciliate community effects on total

respiration, (B) interaction between temperature, low nutrient level, and ciliate presence on total

respiration, and (C) interaction between temperature, high nutrient level, and ciliate presence on

total respiration. Lines within the boxplots represent median, 25th, and 75th percentile values,

while whiskers are defined by the largest value not greater than 1.5× the interquartile range

(IQR). Individual open circles represent raw data points. Pink colors represent treatments with no

ciliates, while purple represents those that have the addition of ciliates.

754
755

Figure 3. (A) Phyla relative abundances of prokaryotic (16S rRNA) prokaryotic communities

across treatments. (B) Boxplot of alpha diversity of prokaryotic communities across temperature,

nutrient level, and presence of ciliate community. Lines within the boxplots represent median,

25th, and 75th percentile values, while whiskers are defined by the largest value not greater than

1.5× the interquartile range (IQR). (C) NMDS ordination plot of Bray-Curtis dissimilarity

of prokaryotic community composition (16S rRNA). Pink colors represent treatments with no

ciliates, while purple represents those that have the addition of ciliates. Squares represent high

nutrient conditions, while triangles represent low nutrient conditions. If the shape is filled in with

color, it is experiencing warming (25☐), whereas those that are not filled in are at base

conditions (22☐).

Figure 4. (A) Boxplot of ciliate density across temperature and nutrient level. Lines within the

boxplots represent the median. (B) Final density data of ciliate communities for each sample

across treatments. (C) Boxplot of alpha diversity of ciliate communities across temperature and

nutrient level. Lines within the boxplots represent the median. (D) The ordination plot is based

on a principal components analysis of ciliate community composition. Arrows show dominant

communities' variable loadings within the functional trait space, while individual points show

individual samples and are colored by treatment. Individual open circles represent raw data

points. Lines within the boxplots represent median, 25th, and 75th percentile values, while

whiskers are defined by the largest value not greater than 1.5× the interquartile range (IQR).

Figure 5. A structural equation model shows proposed relationships between latent variables,

such as temperature and nutrients, and their interaction with the diversity, composition, and total

biomass of microbes and their function. All numbers correspond to the standardized path

coefficients. Solid arrows shown are significant direct effects (p < 0.05). Red arrows represent a

negative relationship, while blue shows a positive.