Genome-wide profiling of the hierarchical control of autophagy dynamics using deep learning

Nathalia Chica[1-2,†,*], Aram N. Andersen[1-3,†], Sara Orellana-Muñoz[1,2], Ignacio Garcia[1,2], Aurélie Nguéa P[1,2], Pilar Ayuda-Durán[1,2], Linda Håkensbakken[1,2], Eline Rødningen[1,2], Christopher D. Putnam[4,5], Manuela Zucknick[6,7], Tor Erik Rusten[1-2,7] and Jorrit M. Enserink[1-3,*]

[1] Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital, 0379 Oslo, Norway
[2] Centre for Cancer Cell Reprogramming, Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, 0318 Oslo, Norway
[3] Section for Biochemistry and Molecular Biology, Faculty of Mathematics and Natural Sciences, University of Oslo, 0371 Oslo, Norway
[4] Ludwig Institute for Cancer Research, 9500 Gilman Drive, La Jolla, CA 92093-0660, USA
[5] Department of Medicine University of California School of Medicine, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0660, USA
[6] Oslo Centre for Biostatistics and Epidemiology, University of Oslo, 0317 Oslo, Norway
[7] Faculty of Medicine, University of Oslo, 0318 Oslo, Norway
† Equal contribution
* Corresponding authors: nathac@uio.no or j.m.enserink@ibv.uio.no

Abstract:

Recycling of nutrients through autophagy is a crucial mechanism for cells to sustain internal stability in a fluctuating environment. Dysregulation of the process has been associated with a range of human disorders, and the core components of autophagy have been comprehensively outlined. However, limited insight into its systems-wide dynamical control has hampered predictive modeling and effective *in vivo* manipulation. Here, we mapped yeast genomic influences on autophagy dynamics in response to changes in nitrogen levels. Using time-resolved high-content imaging coupled with deep learning, we examined the kinetics of autophagy activation and inactivation in 5919 gene deletion mutants, and classified their profiles based on temporal responsiveness and activation potential, as well as their contribution to autophagosome formation and clearance. By integrating these profiles with functional and genetic network data, we unveiled a hierarchical and multi-layered control of autophagy dynamics and exposed novel regulatory features of the core components and well-established nutrient-sensing pathways. Furthermore, by leveraging multi-omics resources and explainable machine learning to model genetic perturbation effects, we identified the retrograde pathway as a central time-varying transcriptional modulator of autophagy execution. These findings offer valuable insights into the systems-wide tuning of autophagy, and advance our understanding of the dynamical control by providing genome-wide quantitative data under concurrent genetic and environmental interventions. We further anticipate that our study can serve as a blueprint for high-content, deep learning-driven exploration of complex dynamical processes in any organism.

Introduction

Living organisms dynamically tune their metabolism to changes in nutrient availability. Macroautophagy, hereafter autophagy, sustains cellular homeostasis by adaptively degrading and recycling cellular components in response to nutrient scarcity, as well as other environmental cues, and its fine-tuning is crucial for a healthy lifespan[1,2]. This is evidenced by tight multi-layered control of the autophagy core regulatory network, and the association of several autophagy-related genes (ATGs) with human disorders[2-4]. Thus, characterizing autophagy execution has been the subject of intense research and several regulatory pathways have been identified[5-9]. However, a generalized understanding of how living organisms dynamically control autophagic activity in the context of a broader genetic landscape is lacking. Along these lines, pharmacological manipulation of autophagy remains rudimentary and involves either complete inhibition of the process or strong activation through the use of TORC1 inhibitors, which have substantial side effects on growth and cell proliferation[10]. A systems-level understanding of the tuning of autophagic activity may help develop predictive models and support the development of effective treatments for autophagy-related conditions.

While emerging studies utilizing multi-omics approaches, mathematical modeling and network analyses have begun to elucidate the complex landscape of potential autophagy regulators, systematic genome-wide data on causal influences on the autophagic nutrient response are limited[11-15]. To date, the study of autophagy has largely been based on gain- or loss-of-function discoveries from autophagic flux assays, which measure the cumulative degradation or re-distribution of cargo over a fixed time window[16,17]. This approach has been successful in elucidating the central mechanisms orchestrating autophagy, as well as proximal upstream switches acting in response to starvation. However, these assays do not fully capture the dynamic nature of autophagy regulation and may overlook important genomic factors influencing activation thresholds, sensitivity to nutrient shifts, and more distant homeostatic tuning[17-21]. These shortcomings also extend to a limited focus on the deactivation of autophagy and return to homeostasis upon nutrient replenishment. Moreover, systematic information on the regulation of different stages of autophagy, such as the formation and clearance of autophagosomes, is lacking. To address these limitations, more sensitive real-time recordings of changes in autophagy dynamics during nutrient depletion and repletion are needed, along with an analysis of divergent features associated with distinct stages of the process.

In this study, we performed a functional analysis of the genome-wide regulation of autophagy dynamics in *Saccharomyces cerevisiae*. By utilizing high-content imaging combined with deep learning, we quantified the nutrient response kinetics and differential influences on autophagy execution in 5919 yeast gene mutants (90% of the yeast genome) with high precision. By assembling quantitative regulatory profiles, and integrating these with the backdrop of genome-wide resources that exist in yeast, we constructed a comprehensive map of the autophagy network response to shifts in nitrogen supply. Our approach reveals previously unexplored gene and systems-level regulators of autophagy kinetics, including phase-dependent control of autophagic flux, and enables genome-scale predictive modeling of autophagy dynamics.

Results

Genome-wide influences on autophagy kinetics. To identify genomic influences on autophagy response kinetics, an arrayed library of 4760 deletion mutants and 1159 DAmP mutants expressing the autophagosome marker mNeonGreen-Atg8 and the vacuole-resident protease Pep4-mCherry was grown to log-phase in a nitrogen-containing medium (+N). Subsequently, it was imaged every hour for 12 hours during nitrogen starvation (-N) and for 7 hours during

nitrogen replenishment (+N) using high-content fluorescence microscopy (Fig. 1A). Each array contained triplicate wells of wild-type (WT) cells and autophagy induction-deficient $atg1\Delta$ mutant cells. We also included triplicates of $vam6\Delta$ mutant cells, which exhibit defects in Gtr1-dependent TORC1 activation and in homotypic fusion with the vacuole, resulting in an abundance of uncleared autophagosomes[22–24].

To robustly quantify the dynamic autophagy response across the genome-wide library with minimal technical confounders and observation bias, we first assembled a large training dataset of extracted image features from 1.34M automatically annotated WT and $atg1\Delta$ cells spanning 70 independent experiments and we used a deep learning approach for classifying autophagic activity (Fig. 1A and S1A-K). The annotation was based on experimental control conditions where autophagic (WT cells over 7 hours in synthetic defined medium lacking nitrogen, -N) and non-autophagic features (either $atg1\Delta$ across all time points, or WT cells over 4 hours in synthetic defined medium containing nitrogen, +N) were maximized in the cell populations, thus avoiding the limitations of human annotation. A hyperparameter search revealed that multiple deep neural network (DNN) models could not only reproducibly predict the activation rate of autophagy with high accuracy (>98%) and dynamic robustness (Fig. S1B-F), but also represent a temporal continuity of latent autophagic features in WT cells and intermediary phenotypes ($vam6\Delta$) relative to non-autophagic noise (variance in $atg1\Delta$; Fig. S1G-I). This allowed for reliable profiling of nutritional responses across plates, resulting in 5678 unique starvation responses and 5442 replenishment responses after removing mutants with low-confidence predictions and low cell counts (Fig. S2A-E).

For WT cells and unperturbed phenotypes, both the activation and inactivation phases of the nutrient response displayed sigmoidal kinetics, with the inactivation phase exhibiting a much steeper decline (Fig. 1B and S2A). Using a double-sigmoidal model to extract 15 response parameters (Fig. 1C and S3A-B)[25], we performed multiple significance tests for mutant perturbations using an expected error model (H_0) derived from replicate WT and negative control measurements (Fig. S3C-E and S4A). The global statistical significance of the response of each mutant was scored by combining test results using a harmonic mean p-value (HMP) weighted by the genomic rate of rejection at a 5% Benjamini-Hochberg (BH) correction cutoff for each test (Fig. 1D and S4B)[26]. A 1% HMP cutoff identified 1613 mutants with perturbations in multiple parameters, and a slight overrepresentation of essential genes (Fig. 1D, E and S4C; Data File S2). Strains with changes in starvation responses were more numerous and distinct compared to those with changes in replenishment responses, indicating a greater regulatory complexity in the activation phase of autophagy (Fig. 1F and S4D). Clustering the response dynamics of the significant mutants revealed six distinct autophagy perturbation profiles (Fig. 1G and S4E), with the three major groups ('ultrasensitive', 'hyposensitive', and 'hyperactive') being characterized by changes in temporal sensitivity and activation rates, and three minor groups ('insufficient activation', 'failed response', and 'null response') exhibiting impaired response potential (activity; Fig. 1H and S4E)[21].

Accuracy and robustness of autophagy perturbation profiling. To evaluate the capacity of the perturbation statistics to retrieve true autophagy-associated genes, we defined two 'gold-standard' reference sets, one set consisting of ATG genes essential for the induction of autophagy ('ATG' set, n=16), and one set comprising genes involved in the maturation or clearance of autophagosomes ('Fusion' set, n=19), identified through literature curation and manual inspection of images (Fig. S5A; see Methods 4.1). Furthemore, we defined a negative control set that comprised deletions of dubious ORFs that did not overlap with verified genes ('control' set, n=23).

Fig. 1. Deep learning-based profiling of autophagy nutritional response kinetics. a, Study workflow overview: 1. An autophagic reporter integration into a yeast gene deletion library; hourly image capture during starvation (-N) and nitrogen replenishment (+N). 2. Cell segmentation and extraction of 30 image features (p). 3. Neural network training to predict autophagy probability (P0 non-autophagy; P1 autophagy) from control data. 4. DNN output and latent space are used to predict autophagy response and states in mutants. 5-6. Identification of outliers, and computation of perturbation differentials. **b,** Predicted autophagy response as a percentage of cells in autophagy for KO and DAmP collection strains. Median (solid line), negative plate control means (dashed line) and standard deviation (ribbon). **c,** Double sigmoidal model fitting was used to measure kinetic perturbations and interpolate the autophagy response dynamics. **d,** HMP correlates with the minimum FDR over the set of hypothesis tests performed and is more sensitive than FDR for mutants with multiple significant perturbations. Distributions and percentages indicate the proportions of DAmP collection mutants; Fisher's exact test. **e,** Distribution of mutants with significant parameter perturbations (p-value ≤0.01). Inset indicating the cumulative probability. Significant mutants (HMP ≤0.01) have multiple significant perturbations, with 50% having four or more. **f,** Distribution of the different parameter perturbations among significant mutants. The color coding indicates groups of related parameters based on clustering (see S4D). **g,** 30% of the yeast genome has a significant autophagy perturbation phenotype, which clustered into six dynamic profiles. **h,** Autophagy response dynamics for the six profiles. Insets indicate log2-fold enrichment of significant perturbations (p-value ≤0.01) in the respective profile over the entire genome. Parameter order and class color coding are the same as in (**f**).

For the combined 'ATG' and 'Fusion' reference sets, the HMP resulted in a precision-recall area under the curve (PRC-AUC) > 0.97 (Fig. S5B-C), and the individual kinetic parameter p-values

yielded PRC-AUCs that were in the 0.7-1 range when compared with genome-wide random samples and in the 0.8-1 range when compared with the control set (Fig. S5D-F). While the 'ATG' and 'Fusion' mutants presented a severe autophagic deficiency in starvation, many 'Fusion' mutants with delays in autophagosome clearance displayed slow deactivation (see T50 in +N) in response to nitrogen replenishment (Fig. S5A,G).

We also evaluated the reproducibility of the measured perturbation phenotypes. Repeating the starvation-induced autophagy protocol for the top mutants (selected based on HMP) revealed a positive association between autophagy perturbation magnitudes and perturbation dynamics correlation between experiments (Fig. S6A-G). In addition, we observed global concordance between measured and predicted phenotypes based on nearest neighbor network interactions, which was subsequently used to identify outliers (potential false positives or negatives) that were retested (Fig. S6E-G). Moreover, to test the reliability of the genome-wide analysis, 33 strongly significant (HMP ≤ 0.001) deletion mutants corresponding to different dynamic profiles and biological functions were reconstructed in triplicate and subjected to a set of validation experiments with variations in the growth protocol (Fig. S7A-B, see Methods 7.1-7.3). The reproducibility of phenotypes across independent clones and different protocols was similar to that across all experiments, with a minor strain background effect and elevated noise in autophagy response caused by growing the cultures to stationary phase (Fig. S7C-G). Moreover, autophagy perturbations, as well as estimated kinetic parameters, correlated in a screen-wide manner, with response potential metrics (rank correlation > 0.8) and starvation slope (rank correlation > 0.7) being the most robust (Fig. S8A-F).

Cellular processes that impact autophagy dynamics. To identify pathways affecting autophagy dynamics, we first performed gene-set enrichment analysis (GSEA) on each of the individual response parameter statistics (signed $-\log_{10}$ p-values) for sets defined by GO biological process terms. Significantly enriched terms (at least one test p-value < 0.05) were analyzed using principal component analysis (PCA) which revealed that response potential and sensitivity to starvation represented the most dominant axes of enriched GO biological processes (Fig. 2A), with terms related to macroautophagy and vacuolar activity yielding the strongest association (Fig. 2B). In contrast, sensitivity to nitrogen replenishment was the second most dominant component (Fig. 2A), and was enriched for processes related to membrane trafficking and fusion (Fig. 2B). Moreover, perturbations in initial and final autophagy levels, which were the least correlated with the other parameters (Fig. S4D and 2A), were enriched for terms related to amino acid and nucleoside metabolism, nitrogen utilization, and mitochondrial activity (Fig. S9A).

We then identified enriched GO terms by GSEA using the combined HMP statistic (Fig. 2C). An enrichment map provided an integrated overview of the primary functional categories influencing autophagy with three large, interconnected clusters representing 1) autophagy and membrane trafficking, 2) RNA metabolism and translation, and 3) gene expression regulation. This map highlighted a graded distribution of perturbation profiles with homogenous enrichment of severe autophagy deficiencies ('null response' and 'failed response') close to autophagy-essential GO clusters, and less severe deficiencies in neighboring clusters (Fig. 2C). Interestingly, the gain-of-function profiles 'ultrasensitive' and 'hyperactive' phenotypes co-occurred more frequently with each other, as well as co-occurring more frequently with the loss-of-function profiles 'hyposensitive' and 'insufficient activation' respectively, indicating consistency in regulatory phenotype pairing along the dimensions of sensitivity and response potential (Fig. 2D).

Fig. 2. Kinetic perturbation profiling is highly sensitive and potentiates deeper annotation of regulatory modules. a, PCA loadings for kinetic perturbation parameters using signed -log$_{10}$ p-values from GSEA of GO-BP. Parameter class colors coding are the same as in (**1f**). **b,** PCA of GO-BP terms from (**a**). Colors indicate the perturbation parameter class (see **1f**) with the most significant enrichment. **c,** Enrichment map based on Jaccard similarity of enriched GO terms from GSEA over the HMP. Edges are drawn for terms with 10% overlap. The pie charts represent the composition of perturbation profiles among enriched genes (leading edge). **d,** Similarity in functional distribution of perturbation profiles across enriched GO terms, assessed by Spearman correlation of GSEA-leading edge counts from (**c**).

Causal structure and network architecture of autophagy regulators. A phenotypic change in response to a genetic intervention does not necessarily imply that a gene is a direct regulator of the measured phenotype. We therefore more closely examined the organization of autophagy-influencing genes and their relation to the autophagy core-machinery in functional networks. Previous work in yeast has shown that genome-wide networks can accurately capture the functional organization of the cell when constructed using similarities in genetic interactions, which are calculated using the Pearson Correlation Coefficients of interactions from Synthetic Genetic Array data (SGA-PCC), or when constructed using protein-protein interactions (PPIs)[27–29]. Using spatial analysis of functional enrichment (SAFE) for the different perturbation profiles, we found distinct genome-wide enrichment patterns for all the profiles except

'ultrasensitive' in both genetic similarity and STRING-based PPI networks (Fig. 3A and S10A)[30]. Noteworthy. the 'ATG set' did not form a distinct cluster in the genetic similarity network but did so in the PPI network, where its spatial enrichment overlapped with that of 'null responses', and it co-localized with processes related to membrane fusion (Fig. S10A).

Fig. 3. Genome-wide regulatory network architecture of autophagy influencing genes. a, SAFE analysis of the global genetic similarity network. Reference map of bioprocesses (left) and of ATG and fusion set (middle) for interpreting the spatial enrichment of autophagy perturbation profiles (right). **b,** Quantification of shortest paths from the ATG core machinery. **c,** Average HMP (-$\log_{10}$) per shortest path distances in different functional interaction networks, computed as indicated in **(b)**. The dashed line indicates HMP = 0.05. **d,** Shortest causal pathlength per autophagy perturbation profile relative to the non-significant genes (HMP > 0.01). **e,** Correlations in temporal autophagy perturbations or sets of perturbation parameters (signed -$\log_{10}$ p-values) between genes that share network interactions or complexome association or lack any interaction altogether (none). Correlations were considered for different HMP cutoffs (1, 0.01, or 0.001). **f,** Enrichment of positive or negative genetic interaction similarities (SGA-PCCs) per correlation intervals of autophagy perturbations between pairs of strongly significant genes (HMP ≤ 0.001). *p-value ≤0.05 (Fisher's exact test). **g,** Direction preferential enrichment of perturbation profiles along the SGA-PCC spectra for individual genes of the core autophagy machinery. *p-value ≤0.05 (permutation test). Screen phenotypes were manually categorized according to the expected or degree of perturbation. NA refers to gene mutants not available in the screen. **h,** Illustration depicting hypothetical regulatory entry points within the modules

of the autophagy core machinery, derived from the significant enrichment of different perturbation profiles along SGA-PCC spectra in (**g**).

Interestingly, more severe autophagy phenotypes were enriched closer to the ATG core sub-graph ('ATG set', including *ATG1* and *ATG8*). The proximity of autophagy-influencing genes to the ATG core in these networks was confirmed in two ways when analyzing the shortest path lengths (Fig. 3B). First, the HMP was more significant for genes that were closer to the ATG core than for those were further away (Fig. 3C). Second, the average shortest path length for each autophagy perturbation profile, except 'ultrasensitive', were significantly closer to the ATG core than the average shortest path length for genes that were not in any perturbation profile (HMP > 0.01; Fig. S10B-F), suggesting that these phenotypes on average represent more direct influences on autophagy. This relationship became even stronger when pathlengths were calculated using the shortest causal path (requiring all intermediate nodes to have a significant influence on autophagy in response to a gene deletion, HMP ≤ 0.01; Fig. 3B, D). In contrast, the 11% of mutations in the 'ultrasensitive' profile, which only affect the response sensitivity, might involve more distant fine-tuning of the nutrient response, spanning through several intermediates as indicated by the observed path length.

Consistent with the clustering of autophagy-influencing genes in these networks, gene pairs with reported functional relations (BioGrid PPIs, SGA-PCC, and STRING)[27,31,32] or complexome associations[33] exhibited correlated autophagy perturbations (Fig. 3E and S11A-B), suggesting that the dynamic profiles were consistent with gene functional relations on a genome-wide scale. The signs of the pairwise SGA-PCC scores can distinguish between inhibitory and activating gene relations[34–36]. We examined whether these relations would also be consistent with the signs of autophagy correlations by assessing whether gene sets with different SGA-PCC cutoffs were enriched in the perturbation dynamics correlations using Fisher's exact test. Indeed, divergences in correlations between strongly significant genes caused a shift in the representation of positive and negative SGA-PCC (Fig. 3F). Moreover, when testing for parametric enrichment of perturbation profiles over the SGA-PCC of core autophagy genes by GSEA, 'hyperactive' phenotypes (in contrast to loss-of-function phenotypes) were preferentially associated with negative SGA-PCCs for both ATG and fusion core genes, suggesting an inhibitory relation (Fig. S11C-D). Using this approach, we inferred differential regulation of the various sub-systems executing autophagy, which cannot be measured from their null or deficient individual deletion phenotypes (Fig. 3G and S11E). Here, 'ultrasensitive' responses were negatively related to genes involved in the early stage of autophagosome formation, particularly Atg1, suggesting that this may be the main limiting factor for the starvation response rate. Interestingly, 'failed response', which was positively related to the fusion core genes, was negatively related to some genes involved in autophagy induction (Fig. 3H). This could indicate the existence of negative feedback circuits from stages involved in autophagosome clearance[37], or pleiotropic functions of genes involved in amino acid liberation and TORC1 re-activation (such as Vps41 and Vam6)[22,38].

To explore the regulatory relations between specific autophagy-influencing genes, we mapped the autophagy perturbation phenotypes across curated complexome networks representing the three major hierarchies of the cell with system-wide influences on the autophagy nutrient response (Fig. S10G-J). This allowed us to observe the penetrance and distribution of phenotypes across well-characterized functional modules. For instance, co-aggregation of multiple significant phenotypes (see chromatin remodeling sub-module), or consistency in the direction of autophagy perturbation for regulatory sub-graphs (Ure2 and Gln3, Gal80 and Gal3, or the CURI complex)[39–41]. For RNA metabolism and translation in particular, we observed a

systemic enrichment of stronger phenotypes along modules related to mRNA processing and translation decoding (binding of mRNA by the 40S small ribosomal subunit), where perturbations in mRNA maturation and binding resulted in a reduced autophagic response, while perturbations in mRNA decay caused the opposite phenotype.

Dynamic latent-space analysis of autophagosome formation and clearance. We reasoned that the latent autophagic features generated by the DNN (Fig. 1A, see autophagy prediction) potentially provided more information about cellular states that could be captured by the binary classifier used to determine the autophagy response (Fig. S1I). We therefore projected these latent autophagic features onto two dimensions using UMAP (Fig. 4A-B and S12A-B), where information about autophagy was conserved together with cell-to-cell variation and dynamic data from the DNN latent space (Fig. 12C-F). WT cells progressed from a UMAP region characterized by no autophagosomes (-N, 0 hours), through a region characterized by free autophagosomes (-N, early time points, 3 to 5 hours), and finally to a region characterized by cleared autophagosomes, where autophagosomes fused with the vacuole (-N, late time points, 5 to 11 hours); (Fig 4A-B and S12A-B). Upon nitrogen replenishment, the WT cell trajectory reversed. As expected, the *atg1Δ* trajectory never progressed out of the no autophagosome UMAP region. In contrast, the *vam6Δ* cell trajectory overlapped with the early WT cell trajectory but never advanced to the cleared autophagosome UMAP region. Instead, it exhibited a distinct UMAP flux characterized by the progressive accumulation of free autophagosomes (Fig. 4A-B), which was statistically separable from that of WT cells in the UMAP within similar intervals of low confidence DNN predictions (classification probabilities in the range $0.1<P_1<0.9$; Fig. S12G-H). Thus, with substantial genome-wide variation in UMAP dynamics and many mutants exhibiting elevated classification uncertainty (Fig. S12E-F), we tested the use of the *vam6Δ* latent space as an "out-of-distribution" intermediate reference state in a two-step model framework for autophagy execution (Fig. 4C and S13A-E). Here, Bayes factors (BFs) were used to score the UMAP latent space evidence of mutant cells executing autophagosome formation (VAM6:ATG1) or clearance (WT:VAM6; details in Methods 6.2). We employed a time-independent BF to classify the overall mutant behavior between the three reference phenotypes (Fig. S13B, D). Additionally, we scored the relative dynamics of the mutants using fixed reference distributions as an indicator of their activity (BF_t) (Fig. S13C, E; Data File S7).

The BFs accurately scored selective defects in autophagy execution, while also quantifying gradual increases in the severity of the mutant phenotypes (Fig. 4C-D, S13F-I, and S14A). For instance, proteins involved in vacuole tethering and the HOPS complex showed evidence of a strong clearance defect and presented a high autophagosome burden, while retromer subunits and several other membrane trafficking proteins were more moderate, with autophagosomes being slowly docked and gradually resolved (Fig. 4C-E, and S14B). Analyzing temporal correlations between genome-wide BF_t variations revealed highly nitrogen-dependent regulation of autophagosome formation, with a rapid decay in correlations between time-lags, indicating transient states of free autophagosomes over brief time intervals (Fig. 4F). In contrast, variations in clearance activity remained relatively constant, with clearance in -N being consequential for the autophagosome burden in +N (Fig. 4F, *lower right*). This distinction highlights the dynamic sensitivity of autophagosome formation to nitrogen availability, while underscoring the steady and robustly regulated activity that characterizes the clearance process.

Fig. 4. Dynamic latent-space analysis of autophagy execution. a, Latent-space model for autophagy execution. *atg1Δ* distribution overlaps with the region for no autophagosomes, *vam6Δ* distribution overlaps with the region for free autophagosomes, and the arrow represents the dynamic of WT cells as they execute autophagy. **b,** Comparison of UMAP-embedded latent-space distributions for WT (ORF) controls, *vam6Δ*, and *atg1Δ* cells with representative micrographs for the indicated time points. The densities represent distributions from 6 independent plates, grouped per time point and colored with the average autophagy prediction (%). Arrows indicate the position and direction of the average latent-space flux per plate. **c,** BF classification of autophagy execution phenotypes across the genome. Densities represent WT (blue), *vam6Δ* (orange), and *atg1Δ* (gray) cells. Reference ATG and fusion mutants are indicated and labeled. **d,** Color-coded functional overview of fusion mutants within each complex, ranging from red to orange, based on severity as determined by BF WT:VAM6. **e,** UMAP-embedded latent-space distributions and autophagy response dynamics for fusion mutants with representative micrographs from indicated time points. **f,** Correlations in genome-wide BF$_t$ variations for different time-lags (indicated by black arrows from BF$_t$ to BF$_{t+l}$). **g,** UMAP-embedded latent-space distributions and autophagy response dynamics for *rtg1Δ*, *rtg2Δ*, and *rtg3Δ* with representative micrographs from indicated time points. **h,** Correlations of average BF$_t$ with autophagic flux measured as the average GFP-Atg8 cleavage (%) from 0, 2, and 4 hours in -N. **i,** Relative contribution of log BF$_t$ scores in predicting autophagic flux per time point, estimated using mixed model regression (with time as grouping factor) based on a univariate model (WT:ATG1) or a bivariate model (VAM6.ATG1 + WT:VAM6). For *i* and *h* points and bars in indicate mean and standard error (see methods for replicas and quantifications).

Parametric gene-set enrichment revealed that perturbations in amino acid synthesis or sensing via TORC1 were distinctly associated with elevated autophagosome formation, while perturbations in proteasomal activity and glucose uptake increased autophagosome clearance (Fig. S14B). Furthermore, using time-independent BFs grouped over each treatment phase, we were able to identify nitrogen-status dependent roles of some of these groups in autophagy execution (Fig. S14C-D). Interestingly, the deletion of genes in the RTG pathway[42] resulted in excessive autophagosome formation in +N, and delayed clearance overall (Fig. 4G and S14E). In particular, *rtg1Δ* exhibited a severe hyperactive phenotype that seemed to be decoupled from

variations in TORC1 signaling (with higher RpS6-p in +N, Fig S15A). Moreover, all RTG mutants exhibited exhausted autophagic flux with a moderate increase in autophagy in response to starvation as quantified by the delivery of distinct cargos, GFP-Atg8 and the modified phosphatase precursor Pho8Δ60, to the vacuole (Fig. S15A-B). This was in contrast to spermidine synthesis mutants, which had a delay in autophagosome formation and clearance (Fig. S14E and S15C), and presented elevated RpS6-p levels in -N (Fig. S15D).

When measuring autophagic flux (GFP-Atg8 cleavage percent) over a set of selected mutants, the time-averaged BF_t WT:ATG1 yielded the best correlation out of any screen measurement (95%), supporting its use as a proxy for autophagic activity (Fig. 4H and S16A-B). Since this measure is the sum of the BF_t metrics with VAM6 as an intermediate step, we quantified the progressive contribution of formation (VAM6:ATG1) and clearance (WT:VAM6) activity on autophagic flux over time using a bivariate regression model (Fig. S16C-D). This finding underscores the importance of clearance regulation, which is the dominant factor for predicting Atg8 flux variation deep into starvation conditions and immediately upon replenishment (Fig. 4I).

Fig. 5. Multi-stage control of autophagy by the TORC1 pathway. a, Average autophagy (%) and BF_t levels in response to starvation, rapamycin or both. Points and bars indicate mean and standard deviation for the plate controls. **b,** Overview of components of the TORC1 pathway. **c,** Autophagy perturbation (%) and log BF_t change from WT ORF controls for different mutants of the TORC1 signaling pathway, the RTG pathway, and the spermidine synthesis pathway in response to starvation, rapamycin or both. **d,** Correlations of RpS6 phosphorylation with GFP-Atg8 cleavage and average BF_t scores for mutants in (**S15a,d;** see methods for replicas and quantifications).

The TORC1 pathway controls multiple stages of autophagy. Defects in the TORC1 signaling pathway caused nitrogen-dependent imbalances between formation and clearance that varied depending on the sub-complex (Fig. S14D-E, S15D-E). Therefore, we performed a secondary screen of selected mutants involved in TORC1 signaling, as well as the RTG pathway and spermidine synthesis. Mutants were subjected to rapamycin treatment in combination with -N or

+N for 10 hours and analyzed using our deep learning approaches (Fig. S17 and Fig. S18). Overall, rapamycin boosted autophagy induction and synergized with genetic disruptions of TORC1 signaling (Fig. 5A-C). Remarkably, rapamycin reduced autophagosome clearance activity in -N as measured by the reduction in the BF_t WT:VAM6 values (Fig. 5A), which was decorrelated from the effect in +N (Fig. S17F). This suggests a possible role of nutrient-regulated vacuole fusion, through TORC1 and other mechanisms. Notably, deleting the TORC1-independent branch of Gtr1/2 involving Ltv1[43] selectively compromised clearance without causing hyperactive autophagosome formation in response to starvation and rapamycin (Fig. 5B-C and Fig. S18). DAmP disruption of Tor2, Kog1 and Lst8, or non-essential deletion of Tor1 or the vacuole-specific component Tco89[44], led to an elevated autophagosome formation under all conditions, and rapamycin-independent suppression of clearance in all the mutants except Tor1 (Fig. 5C and Fig. S18). This autophagosome clearance defect was confirmed in *tco89Δ* cells, exhibiting severely compromised autophagic flux in -N (lower GFP-Atg8 cleavage; Fig. S15D). Moreover, comparing RpS6 phosphorylation with the BF_s revealed a negative correlation with autophagosome formation in -N, underscoring the role of TORC1 as a suppressor of autophagy induction, while yielding a slight positive correlation with clearance in +N (Fig. 5D), suggesting that TORC1 phosphorylation activity might be positively related to fusion activity.

Cross-omics inference of transcriptional mediators of autophagy regulation. Genome-wide quantification of dynamic signatures enables the construction of comprehensive systems-level models that can predict and simulate processes under different scenarios without sacrificing structural complexity or causal sufficiency. This has been demonstrated with biologically informed machine learning models for growth and metabolism in yeast, where multi-omics data resources are plentiful[45,46]. As a proof-of-principle, we used random forest to predict different autophagy perturbation metrics from distinct yeast gene deletion datasets containing transcriptomic, proteomic, and amino acid metabolomic features[47–49], as well as from genetic (SGA) and STRING genome-wide correlation networks[27,32]. All datasets could predict autophagy perturbations with performances varying with the experimental reproducibility of the different perturbation parameters (Fig. S19A and S8F), and the genome-wide networks performing the best overall with test predictions yielding over 40% median correlations between predicted and true outcomes for BF_t metrics in -N (Fig. S19A).

Given the abundance of transcription factor (TF) deletions found to strongly influence autophagy kinetics and the general challenge of establishing the exact causal mediators through which TFs act on a cellular process, we performed a mediation analysis by searching for gene expression (GE) changes predicting the initial perturbations in autophagy dynamics. For this, we used the GE induction profiles for 186 overlapping TFs from the Induction Dynamics gene Expression Atlas (Fig. 6A)[50].

Fig. 6. Cross-omics mediation analysis reveals transcriptional regulatory mechanisms of autophagy dynamics. a, Transcription factor (TF) gene expression induction profiles from Hackett et al. (*50*) were used to predict the corresponding autophagy perturbation dynamics of TF gene deletions, using random forest (RF). Gene targets (model features) with high importance for predicting autophagy perturbations were evaluated as potential causal mediators of TF influence on autophagy. **b,** Adjusted SHAP values for differential regulation of gene expression comparing feature classes across models and autophagy metrics. F: formation genes; C: clearance genes with negative BF$_t$ z-scores beyond two standard deviations from the mean. Points and bars indicate mean and standard error. **c,** Gene-wise SHAP values for differential regulation of gene expression predicting autophagy responses (%). **d,** Hierarchical clustering of average TF-induced gene expression changes (z-scores) for gene targets that were selected as important model features (using sumGain). **e,** qRT-PCR for expression of *ATG1*, *ATG13*, and *ATG14* in *rtg1Δ*, *rtg2Δ*, *rtg3Δ*, and WT cells subjected to 4 hours of -N followed by 1 hour of +N. +N t=0h refers to cells before shifting to -N. Graphs represent mean and standard error. **f,** Differential expression of various target genes in *rtg1Δ*, *rtg2Δ*, *rtg3Δ* cells per phase from (**e**) and **(S19f)**. n=3. *p ≤0.05;**p ≤0.01;***p ≤0.001; t-test in (**e**) and regression in (**f**). **g,** Immunoblot analysis of GFP-Atg8 cleavage and RpS6 phosphorylation for WT, *rtg1Δ*, *gcn4Δ*, and *gcn4Δ rtg1Δ* cells (*upper panels*), and for WT, *rtg1Δ*, *gln3Δ*, and *gcn3Δ rtg1Δ* cells (*lower panels*) in -N for the indicated time points. +N t=0h refers to cells before shifting to -N. *p ≤0.05**p ≤0.01;***p ≤0.001; mixed model regression (n=4 for WT/*rtg1Δ*; n=2 for *gln3Δ*/*gcn4Δ*/double deletions).

Models built with only potential causal mediators (significant genes, HMP ≤ 0.01, and excluding TFs) were able to sustain the predictivity, exhibiting only a moderate drop from the genome-wide models (Fig. S19B). Moreover, gene targets with severe autophagy perturbations were estimated to mediate stronger influences across all models (Fig. S19C), with targets involved in autophagosome formation and clearance predicted as positive mediators of autophagy regulation overall (Fig. 6B and S19D). Interestingly, only targets involved in clearance were predicted to mediate a positive influence on clearance dynamics, whereas targets involved in autophagosome formation were predicted to exert a negative influence (Fig. 6B and S19D, *BF WT:VAM6 panels*). This suggests that transcriptional suppression of certain autophagy induction genes could play a role in promoting resolution. *ATG1* and *ATG14* were among the major TF targets predicted to mediate autophagy initiation control (Fig. 6C), while *VAM10* and the ESCRT-0 subunit-encoding *VPS27* were predicted to mediate autophagosome clearance regulation (Fig. S19E). By clustering the average induction response of predicted mediators for

select autophagy-regulating TFs, we found that RTG-mediated suppression of autophagy induction was likely mediated through Gcn4-regulated expression of *ATG1* (Fig. 6D)[50,51]. Indeed, RT-qPCR experiments revealed that *ATG1* expression was substantially greater in *rtg1Δ* cells than in WT control under both -N and +N conditions, with the effects of the RTG subunits on expression of different ATGs and upstream TORC1 regulators, decoupling during starvation (Fig. 6E-F and S19F-G). When *rtg1Δ* was crossed with *gcn4Δ* strains, the hyperactive autophagy phenotype was readily rescued, but this effect was not observed in *rtg1Δgln3Δ* strains (Fig. 6G). This epistatic effect of *rtg1Δ* on *gln3Δ* was consistent with *rtg1Δ* supressing *GLN3* expression in +N, leading to decreased expression of Gln3-regulated gene targets in -N (i.e *ATG14;* Fig. 6F)[39].

Discussion

Studying the kinetics of dynamical cell behavior is becoming increasingly relevant as we strive to achieve predictability and treatment guidance from our scientific models. This may be particularly important for autophagy, which is intrinsically coupled to an organism's metabolism and growth, and continuously fluctuates with environmental changes such as feeding cycles and circadian rhythms. For instance, manipulating the sensitivity of autophagy activation in phases of declining nutrient levels will require knowledge about how kinetic influences manifest through functional interactions with rate-limiting factors (Fig. 3G). Since genetic perturbations of the core machinery severely compromise autophagy execution, high-quality temporal data from associated genes and multi-omics strategies to model the observed dynamics are warranted. Our observation that there is a consistency of autophagy perturbation profiles and the direction of functional similarities with the core machinery (Fig. 3G and S11B), and the concordance in phenotypes between TF deletions and the predicted GE mediators (Fig. 6B), strongly supports such a utility of the presented data.

By demonstrating the predictive power of our approach, we identified a novel phase-dependent tuning mechanism through the transcriptional repression of autophagy by the RTG signaling pathway. With Gcn4 positively regulating RTG gene expression (Fig. S19H)[50], the retrograde pathway could work as a negative feedback, buffering a positive loop between Gcn4 and Gln3 to support temporally coordinated *ATG* expression, limiting uncontrolled autophagy induction and promoting autophagosome clearance[50,51]. Recent discoveries suggest that Atg1 plays a role in activity-dependent disassembly of several ATG sub-complexes involved in autophagosome formation[52]. Thus, modulating the relative levels of ATG proteins may play an important role in tuning the strength and persistence of autophagy induction, and the imbalanced expression observed in RTG deletion cells could explain the diminishing returns in autophagic flux from starvation (Fig. 4G and S15A-B). This adaptive control of autophagy dynamics may be evolutionarily conserved, as the yeast retrograde response and analogous pathways in higher organisms support cellular survival in response to environmental perturbations or to cellular challenges associated with aging[53].

The versatility of our expectation-based deep learning approach offered us a richer characterization of the autophagic feature space with high accuracy. Human annotation of heterogeneous data is a fundamental challenge for the utility of machine learning in cell biology[54], while experimental generation of high-content reference data is becoming increasingly easy. Using interventional control conditions to instruct the expected outcome space could easily be extended to a range of other statistical learning tasks[54]. When combined with a systematic analysis across functionally related gene sets, this approach revealed discrete phenotypes that otherwise would likely go unnoticed. An unexpected finding was the connection between the TORC1 pathway and autophagosome clearance. While this requires further investigation, one possibility is that TORC1 inhibition leads to a flux imbalance between autophagosome formation

and clearance (Fig. 4H and 5), or that nutrient-reactivation of TORC1 promotes clearance of residual autophagosomes[37]. Alternatively, Gtr1/2 regulation of Ltv1, which has previously been connected to endosomal membrane trafficking[43], could play a role in supporting vacuolar fusion[24,55]. Treatment with rapamycin also revealed further levels of variation, and otherwise unobserved nutrient-gene interactions (Fig. 5C and S17E-F). Hence, further manipulations of the culture environment and metabolic growth history can potentially unveil an even broader spectrum of unexplored autophagy regulatory mechanisms.

Due to the sensitivity of our approach we were able to detect a vast number of genes with a significant influence on autophagy kinetics—approximately 30% of the investigated genome (Fig. 1). These genes were distributed across multiple hierarchies of cell functions, with varying connectivity to the autophagy core machinery that correlated with the observed tuning strength (Fig. 2-3). In conclusion, our study provides a time-resolved genome-wide map of the activation and inactivation of autophagy, providing comprehensive insight into the dynamic regulation of the cellular nutrient response.

Data and material availability: Genome-wide high-content images will be shared by the lead contact upon request, considering storage constraints. Any additional information required for the reanalysis of the data reported in this study can be obtained by contacting corresponding authors.

Supplementary Materials: Materials and Methods, Figs. S1 to S19, Table S1 and References (*56–95*).

Acknowledgments: We thank Dr. Anne Simonsen and Dr. Harald Stenmark for critical comments on the manuscript. We also thank Dr. Eduardo Cebollero and Dr. Fulvio Reggiori for providing valuable guidance on the experimental protocol for alkaline phosphatase (ALP) assay. We acknowledge Dr. Anna Lång and Dr. Vigdis Sørensen and Advanced Light Microscopy Core Facilities at Oslo University Hospital Gaustad and Montebello for technical support. We thank members of the Enserink group for feedback and suggestions.This work was funded by Norwegian Cancer Society (project numbers 182524 and 208012), Norwegian Health Authority South-East (2017064, 2017072, 2018012, 2019096), Research Council of Norway (261936, 301268), Research Council of Norway through its Centres of Excellence funding scheme, project number 262652, and Fundación Alfonso Martín Escudero (SMO).

Author contributions: Conceived the study and acquired funding: J.M.E. Conceptualization: N.C., A.N.A. and J.M.E. Methodology – construction genome-wide libraries: I.G., A.N.P., P.A.D. and C.D.P. Methodology – optimizing cell-growth and starvation protocols: N.C. Methodology – developing and optimizing high-content screening protocols: N.C., S.O.M. and A.N.P. Methodology – construction of validation libraries: N.C., S.O.M. and L.H. Methodology – image segmentation and feature extraction analysis: I.G. Methodology – developing and optimizing image analysis and deep learning pipeline: A.N.A. and I.G. Investigation: N.C., L.H., S.O.M., P.A.D., A.N.P., and E.R. Investigation – performing and validating high-content screening: N.C. and L.H. Data curation and analysis: N.C. and A.N.A.; Formal analysis: A.N.A. Network analysis: N.C. and A.N.A. Visualization: N.C. and A.N.A. Writing – original draft: N.C. and A.N.A. Writing – review & editing: N.C., A.N.A., C.D.P., M.Z., T.E.R., and J.M.E. Project coordination: N.C. and J.M.E. Supervision: N.C., M.Z., T.E.R., and J.M.E. All authors discussed, revised, and approved the manuscript.

Competing interests: Authors declare that they have no competing interests.

References

1. He, C. Balancing nutrient and energy demand and supply via autophagy. *Curr. Biol.* **32**, R684–R696 (2022).
2. Hansen, M., Rubinsztein, D. C. & Walker, D. W. Autophagy as a promoter of longevity: insights from model organisms. *Nat. Rev. Mol. Cell Biol.* **19**, 579–593 (2018).
3. Galluzzi, L., Pietrocola, F., Levine, B. & Kroemer, G. Metabolic control of autophagy. *Cell* **159**, 1263–1276 (2014).
4. Aman, Y. *et al.* Autophagy in healthy aging and disease. *Nat Aging* **1**, 634–650 (2021).
5. Feng, Y., He, D., Yao, Z. & Klionsky, D. J. The machinery of macroautophagy. *Cell Res.* **24**, 24–41 (2014).
6. González, A. & Hall, M. N. Nutrient sensing and TOR signaling in yeast and mammals. *EMBO J.* **36**, 397–408 (2017).
7. Tsukada, M. & Ohsumi, Y. Isolation and characterization of autophagy-defective mutants of Saccharomyces cerevisiae. *FEBS Lett.* **333**, 169–174 (1993).
8. Mizushima, N. *et al.* A protein conjugation system essential for autophagy. *Nature* **395**, 395–398 (1998).
9. Kirisako, T. *et al.* Formation process of autophagosome is traced with Apg8/Aut7p in yeast. *J. Cell Biol.* **147**, 435–446 (1999).
10. Galluzzi, L., Bravo-San Pedro, J. M., Levine, B., Green, D. R. & Kroemer, G. Pharmacological modulation of autophagy: therapeutic potential and persisting obstacles. *Nat. Rev. Drug Discov.* **16**, 487–511 (2017).
11. Behrends, C., Sowa, M. E., Gygi, S. P. & Harper, J. W. Network organization of the human autophagy system. *Nature* **466**, 68–76 (2010).
12. Kramer, M. H. *et al.* Active Interaction Mapping Reveals the Hierarchical Organization of Autophagy. *Mol. Cell* **65**, 761–774.e5 (2017).
13. Hu, Z. *et al.* Multilayered Control of Protein Turnover by TORC1 and Atg1. *Cell Rep.* **28**, 3486–3496.e6 (2019).
14. Diehl, V. *et al.* Minimized combinatorial CRISPR screens identify genetic interactions in autophagy. *Nucleic Acids Res.* **49**, 5684–5704 (2021).
15. Türei, D. *et al.* Autophagy Regulatory Network - a systems-level bioinformatics resource for studying the mechanism and regulation of autophagy. *Autophagy* **11**, 155–165 (2015).
16. Mizushima, N. & Murphy, L. O. Autophagy Assays for Biological Discovery and Therapeutic Development. *Trends Biochem. Sci.* **45**, 1080–1093 (2020).
17. Loos, B., du Toit, A. & Hofmeyr, J.-H. S. Defining and measuring autophagosome flux—concept and reality. *Autophagy* **10**, 2087–2096 (2014).
18. Beesabathuni, N. S., Park, S. & Shah, P. S. Quantitative and temporal measurement of dynamic autophagy rates. *Autophagy* **19**, 1164–1183 (2023).
19. Martin, K. R. *et al.* Computational model for autophagic vesicle dynamics in single cells. *Autophagy* **9**, 74–92 (2013).
20. Kholodenko, B. N. Cell-signalling dynamics in time and space. *Nat. Rev. Mol. Cell Biol.* **7**, 165–176 (2006).
21. Di-Bella, J. P., Colman-Lerner, A. & Ventura, A. C. Properties of cell signaling pathways and gene expression systems operating far from steady-state. *Sci. Rep.* **8**, 17035 (2018).
22. Binda, M. *et al.* The Vam6 GEF controls TORC1 by activating the EGO complex. *Mol. Cell* **35**, 563–573 (2009).
23. Ostrowicz, C. W. *et al.* Defined subunit arrangement and rab interactions are required for functionality of the HOPS tethering complex. *Traffic* **11**, 1334–1346 (2010).
24. Chen, Y. *et al.* A Vps21 endocytic module regulates autophagy. *Mol. Biol. Cell* **25**, 3166–3177

(2014).

25. Caglar, M. U., Teufel, A. I. & Wilke, C. O. Sicegar: R package for sigmoidal and double-sigmoidal curve fitting. *PeerJ* **6**, e4251 (2018).

26. Wilson, D. J. The harmonic mean -value for combining dependent tests. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 1195–1200 (2019).

27. Costanzo, M. *et al.* A global genetic interaction network maps a wiring diagram of cellular function. *Science* **353**, (2016).

28. Hao, B. & Kovács, I. A. A positive statistical benchmark to assess network agreement. *Nat. Commun.* **14**, 2988 (2023).

29. Huang, J. K. *et al.* Systematic Evaluation of Molecular Networks for Discovery of Disease Genes. *Cell Syst* **6**, 484–495.e5 (2018).

30. Baryshnikova, A. Systematic Functional Annotation and Visualization of Biological Networks. *Cell Syst* **2**, 412–421 (2016).

31. Oughtred, R. *et al.* BioGRID: A Resource for Studying Biological Interactions in Yeast. *Cold Spring Harb. Protoc.* **2016**, db.top080754 (2016).

32. Szklarczyk, D. *et al.* STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res.* **43**, D447–52 (2015).

33. Meldal, B. H. M. *et al.* Analysing the yeast complexome-the Complex Portal rising to the challenge. *Nucleic Acids Res.* **49**, 3156–3167 (2021).

34. Collins, S. R. *et al.* Functional dissection of protein complexes involved in yeast chromosome biology using a genetic interaction map. *Nature* **446**, 806–810 (2007).

35. Ulitsky, I., Shlomi, T., Kupiec, M. & Shamir, R. From E-MAPs to module maps: dissecting quantitative genetic interactions using physical interactions. *Mol. Syst. Biol.* **4**, 209 (2008).

36. Schuldiner, M. *et al.* Exploration of the function and organization of the yeast early secretory pathway through an epistatic miniarray profile. *Cell* **123**, 507–519 (2005).

37. Yu, L. *et al.* Termination of autophagy and reformation of lysosomes regulated by mTOR. *Nature* **465**, 942–946 (2010).

38. Takeda, E. *et al.* Vacuole-mediated selective regulation of TORC1-Sch9 signaling following oxidative stress. *Mol. Biol. Cell* **29**, 510–522 (2018).

39. Chan, T. F., Bertram, P. G., Ai, W. & Zheng, X. F. Regulation of APG14 expression by the GATA-type transcription factor Gln3p. *J. Biol. Chem.* **276**, 6463–6467 (2001).

40. Pilauri, V., Bewley, M., Diep, C. & Hopper, J. Gal80 dimerization and the yeast GAL gene switch. *Genetics* **169**, 1903–1914 (2005).

41. Albert, B. *et al.* A Molecular Titration System Coordinates Ribosomal Protein Gene Transcription with Ribosomal RNA Synthesis. *Mol. Cell* **64**, 720–733 (2016).

42. Butow, R. A. & Avadhani, N. G. Mitochondrial signaling: the retrograde response. *Mol. Cell* **14**, 1–15 (2004).

43. MacDonald, C. & Piper, R. C. Genetic dissection of early endosomal recycling highlights a TORC1-independent role for Rag GTPases. *J. Cell Biol.* **216**, 3275–3290 (2017).

44. Hatakeyama, R. *et al.* Spatially Distinct Pools of TORC1 Balance Protein Homeostasis. *Mol. Cell* **73**, 325–338.e8 (2019).

45. Ma, J. *et al.* Using deep learning to model the hierarchical structure and function of a cell. *Nat. Methods* **15**, 290–298 (2018).

46. Culley, C., Vijayakumar, S., Zampieri, G. & Angione, C. A mechanism-aware and multiomic machine-learning pipeline characterizes yeast cell growth. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 18869–18879 (2020).

47. Kemmeren, P. *et al.* Large-scale genetic perturbations reveal regulatory networks and an abundance of gene-specific repressors. *Cell* **157**, 740–752 (2014).

48. Messner, C. B. *et al.* The proteomic landscape of genome-wide genetic perturbations. *Cell* **186**, 2018–2034.e21 (2023).

49. Mülleder, M. *et al.* Functional Metabolomics Describes the Yeast Biosynthetic Regulome. *Cell* **167**, 553–565.e12 (2016).

50. Hackett, S. R. *et al.* Learning causal networks using inducible transcription factors and transcriptome-wide time series. *Mol. Syst. Biol.* **16**, e9174 (2020).

51. Natarajan, K. *et al.* Transcriptional profiling shows that Gcn4p is a master regulator of gene expression during amino acid starvation in yeast. *Mol. Cell. Biol.* **21**, 4347–4368 (2001).

52. Schreiber, A. *et al.* Multilayered regulation of autophagy by the Atg1 kinase orchestrates spatial and temporal control of autophagosome formation. *Mol. Cell* **81**, 5066–5081.e10 (2021).

53. Jazwinski, S. M. & Kriete, A. The Yeast Retrograde Response as a Model of Intracellular Signaling of Mitochondrial Dysfunction. *Front. Physiol.* **3**, 26575 (2012).

54. Mattiazzi Usaj, M. *et al.* Systematic genetics and single-cell imaging reveal widespread morphological pleiotropy and cell-to-cell variability. *Mol. Syst. Biol.* **16**, e9243 (2020).

55. Nair, U. *et al.* SNARE proteins are required for macroautophagy. *Cell* **146**, 290–302 (2011).

Supplementary Materials for

Genome-wide profiling of the hierarchical control of autophagy dynamics using deep learning

Nathalia Chica[1-2,†,*], Aram N. Andersen[1-3,†], Sara Orellana-Muñoz[1,2], Ignacio Garcia[1,2], Aurélie Nguéa P[1,2], Pilar Ayuda-Durán[1,2], Linda Håkensbakken[1,2], Eline Rødningen[1,2], Christopher D. Putnam[4,5], Manuela Zucknick[6,7], Tor Erik Rusten[1-2,7] and Jorrit M. Enserink[1-3,*]

* Corresponding authors: nathac@uio.no or j.m.enserink@ibv.uio.no

The PDF file includes:

Table of contents

Materials and Methods

1. High-content screening

1.1 Genome-wide libraries details

The yeast mutant libraries harboring the autophagy reporter (autophagy library KO-A and DAmP-A) were constructed based on the collection of non-essential deletion mutants (YKO collection; Open Biosystems), and the collection of essential genes with a decreased abundance by mRNA perturbation (DAmP collection; Open Biosystems). The DamP mutations were chosen over conditional temperature-sensitive (ts) mutations to eliminate temperature as a variable that could impact the autophagy response[55]. The mutant strains were originally distributed across 56 KO-A and 12 DAmP-A 96-well plates. To introduce control strains into each plate, the mutant strains in row A were transferred to new plates designated as Mixed plates. The mutants with significantly reduced growth rates, that may affect the cell count for imaging, were re-inoculated with higher density into independent plates that were annotated as Recovery plates. A subset of mutants was rearranged into Repetition plates, which were used for validation and reproducibility experiments. Rapamycin plates were prepared for the analysis of rapamycin response.

Each plate included three control strains in triplicate, specifically the SGA starting strain WT (Y7092)[56], the deletion mutant *VAM6Δ::kanMX4* (KO-A), and the deletion mutant *ATG1Δ::kanMX4* (KO-A). The libraries Mix, Recovery, Repetition and Rapamycin included the deletions of three dubious ORFs *YDR535CΔ::kanMX4* (KO-A), *YCR102W-AΔ::kanMX4* (KO-A), and *YEL074WΔ::his3MX6* (this study), in addition to *ATG1Δ* and *VAM6Δ,* to control the influence of the kanamycin and histidine selectable markers on the autophagy response. The complete list of mutant strains and controls distributed into the arrays KO-A, DAmP-A, Recovery, Mixed, Repetition, and Rapamycin is provided in Data File S1.

1.2 Genome-wide libraries preparations

To prepare genome-wide libraries for high-throughput screening, duplicates of the yeast libraries were created in 96-well plates (Nunc™ Microwell™, Thermo Fisher). In each well, 1 µl of both control strains and mutant strains was inoculated into 150 µl of synthetic medium containing amino acids (SC: 6.9 g yeast nitrogen base with ammonium sulfate and without amino acids, 20 g/ glucose anhydrous, and 0.790 g Complete Supplement Mixture Formedium™ per liter). To promote optimal growth and low cell death rate, the duplicate plates were incubated at 25 °C for 72 h, before being stored at -80 °C with 20% glycerol.

    A working batch of 16 plates from duplicates was prepared five days before an imaging

round. In each plate, 1 µl of both mutant and control strains was inoculated into 200 µl of SC medium. Following a 48-hour incubation at 25°C, these plates were stored at 4°C with a cover film. The batch was divided into four sets, and each set underwent pre-culturing and imaging on consecutive days.

1.3 Cell growth and starvation protocol for high-content screening

The growth protocol was optimized to enable high-throughput cell growth while inducing an optimal autophagy response in yeast cells. This involved periodic medium replacement, which kept cells metabolically active and prevented premature autophagy initiation due to nutrient unavailability upon prolonged incubations without shaking. It ensured that yeast cells were in the log phase and minimized population response variability before the onset of starvation.

After a 48-hour incubation of the previous batch, 1 µl of strains with a normal growth rate and 4 µl of strains with a reduced growth were transferred into 200 µl of fresh SC medium. These plates were then incubated at 25°C for an additional 48 h, followed by reinoculation and another 24-hour incubation at 25°C. For time-lapse imaging, 4-6 µl of cells were inoculated into 96-well black plates (Corning® Black microplate, Sigma-Aldrich) coated with Con A (0.25 mg/ml) and incubated for 8 hours at 25 °C to ensure cells were in exponential phase. To induce autophagy, cells were washed twice with sterile MilliQ water and transferred to a nitrogen-starvation medium for 12 h (SD-N: 1.9 g yeast nitrogen base without ammonium sulfate and amino acids, and 20 g glucose anhydrous Formedium™ per liter). Subsequently, the same cells were shifted to SC medium (at t=12 h) to record autophagy inactivation upon nitrogen replenishment for 7 hours. Live imaging was performed at 30 °C, capturing images every hour for 19 hours. The cells were maintained at room temperature during the intervals between each time point of image acquisition.

1.4 Autophagy reporter details

To assess autophagy activity, a fluorescent protein-tagged Atg8 was examined, and its translocation from the cytoplasm to the vacuole was monitored using fluorescent microscopy[57]. To ensure photostability throughout time-lapse experiments, the optimized mNG fluorescent protein was fused into the N-terminus of Atg8, under the control of the ScTDH3 (GPD) promoter. Furthermore, to visualize the vacuole, the mCherry fluorescent protein was fused into the C-terminus of the vacuolar proteinase A Pep4.

1.5 Plasmid construction

The pYM-N natNT2-GPD1pr-mNeongreen plasmid was generated by replacing the yeGFP coding sequence from pYM-N17[58] by the codon-optimized monomeric NeonGreen (mNG) DNA sequence[59]. The mNG sequence was amplified from the pFA6-mNG-his3MX6 plasmid using primers mNG-Fwd-XbaI and mNG-Rv-EcoRI, which included *XbaI* and *EcoRI* sites for the tag replacement (see Data File S1, *primer sequences*).

1.6 Integration of autophagy reporter into libraries and SGA query construction

The autophagy reporter was crossed with the collections of non-essential and the essential deletion mutants to create YKO-A and DAmP-A collections. Triple allele strains harboring the autophagic reporter and the deletion of non-essential or DAmP alleles were obtained by adapting the Synthetic Genetic Array (SGA) analysis method[60,61].

The query bait containing autophagy reporter was created using the SGA host strain *MATα* haploid Δ*can1*::*STE2*pr-Sp_*his5* Δ*lyp1* (Y7092; kind gift from C. Boone, University of Toronto, Canada). Endogenous *ATG8* and *PEP4* were tagged with mNG and mCherry using a standard PCR-based approach [58]. In brief, *natNT2:GPD1pr-mNG-ATG8* was constructed by PCR amplification of a *natNT2-GPD1pr-mNeongreen* fragment from the pYM-N17-mNeonGreen plasmid using the primer pair ATG8-S1 and ATG8-S4, whereas *PEP4-mCherry:hphMX6* was constructed by integrating an *mCherry:hphMX6* PCR fragment amplified from plasmid pBS35 using primer pair PEP4-S3 and PEP4-S2 at the 3' end of the *PEP4* locus (see Data File S1, *primer sequences*). This resulted in the construction of JEY11511 (*MATα can1Δ::STE2pr-Sp_his5 lyp1Δ his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 natNT2:GPDpr-mNeongreen-ATG8 PEP4-mCherry:hphMX6*).

The SGA bait strain JEY11511 was crossed with a target array of 4760 viable MATa deletion mutants and 1159 conditional alleles of essential genes (KO and DAmP mutants respectively) in the BY4741 background (MAT**a** *his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*), wherein each ORF or 3' UTR was disrupted by a kanMX4 resistance cassette marker. Crossing of JEY11511 was performed in quadruplicate by pinning onto fresh YPD agar plates using a Singer RoToR (Singer Instruments) followed by growth for two days. Cells were subjected to two rounds of pinning onto diploid selection medium (YPD-agar containing 200 µg/ml geneticin (G418, ThermoFisher) and 200 µg/ml hygromycin (Gibco)) and grown for 1-2 days at 30˚C. The cells were then pinned onto pre-sporulation medium (15 g Difco nutrient broth (BD Biosciences), 5 g Bacto-yeast extract (ThermoFisher), 10 g Bacto-agar (ThermoFisher), and 62.5 ml 40% glucose (ThermoFisher) per 500 ml) and grown for 3 days at 30˚C. Cells from the pre-sporulation medium were then pinned onto sporulation medium (10 g potassium acetate, 0.05 g zinc acetate, 20 g Bacto-agar per liter, containing a final concentration of 50 µg/ml G418 and 50 µg/ml hygromycin) and incubated for 7 days at 30˚C. The resulting spore-containing cells were then subjected to two rounds of pinning onto diploid killing medium (1.7 g yeast nitrogen base without amino acids and without ammonium sulfate (BD Biosciences), 1 g L-glutamic acid monosodium salt, 2 g SC dropout mix without lysine, canavanine, and histidine (US Biological), 20 g Bacto-agar, 50 ml of 40% glucose per liter, containing a final concentration of 50 µg/ml thialysine (Sigma Aldrich), 60 µg/ml canavanine (Sigma Aldrich), 200 µg/ml G418 (Gibco), 200 µg/ml hygromycin (Invitrogen), and 100 µg/ml nourseothricin (Jena Bioscience)) followed by growth for 5 days at 30˚C for the first pinning and 2 days at 30˚C for the second pinning. Cells were then subjected to two rounds of pinning and growth on haploid selection medium (1.7 g yeast nitrogen base without amino acids and without ammonium sulfate, 1 g L-glutamic acid monosodium salt, 2 g SC dropout mix without histidine, 20 g Bacto-agar, 50 ml of 40% glucose per liter, containing a final concentration of 200 µg/ml G418, 200 µg/ml hygromycin, and 100 µg/ml nourseothricin) and grown for 2 days at 30˚C. Then the cells were pinned and grown on YPD-agar followed by storage at -80˚C in YPD media containing 20% (v/v) glycerol. The resulting MATa haploid triple mutants were used for high-content screening.

1.7 High-content fluorescent imaging

The high-content screening (HCS) for autophagy analysis was carried out in a 96-well plate format using an ImageXpress® Micro Confocal microscope equipped with a temperature control unit, LED light source, and specific filter sets (Molecular Devices). The images were acquired using automated acquisition in wide-field, utilizing the Nikon Plan Apochromat 60x/0.95 Air objective coupled to a 16-bit 4MPix scientific CMOS camera and a laser-based autofocusing unit. Image acquisition was controlled by the ImageXpress Analysis Software (Molecular

Devices). Live cell imaging was conducted at 30 °C ± 2 °C. Fluorescence signals were captured using specific filter sets for FITC and TexRed, with exposure times of 200 ms and 800 ms, respectively. For transmitted light (TL) images, an exposure time of 10 ms was used. The same cells were imaged every hour for 19 hours with 2-3 images per well by selecting adjacent imaging sites near the center of the well and employing modus focus on the plate and bottom to maintain the same plane.

1.8 Automatic image analysis using Fiji/ImageJ

We developed an automated image analysis tool using Fiji/ImageJ to extract features from the genome-wide dataset and evaluate changes in the autophagy reporter along the time-lapse experiments. The tool employed a macro to extract parameters for each image categorized by well (row+col), time point (starvation/replenishment), image within well (s1, s2, s3), and channel (w1=BF, w2=green, w3=red). The extracted features included parameters related to both fluorescent signals and cell morphology. Parameters related to fluorescent signals were mean and standard deviation of signal intensity, modal and minimum signal values, and integrated median signal intensity. Parameters related to cell morphology included cell area, centroid and center coordinates, perimeter, bounding box, best-fit ellipse dimensions, shape descriptors, ferret dimensions, skewness, and kurtosis.

For image segmentation, we first enhanced contrast using a saturation value of 0.3 and then sharpened it to improve clarity and pixel contrast. We further enhanced contrast with the same saturation value to refine the image after sharpening. We then applied the "Unsharp Mask" plugin with a radius of 5 and a mask value of 0.70. We used the "Auto Threshold" plugin with the default method and set the image to white. To finalize the segmentation and make objects white on a black background, we inverted the image and applied the "Minimum" filter with a radius of 0.5. To eliminate small holes or gaps in the image, we used the "Close-" plugin and followed it with another inversion of the image.

To identify and count the objects in the image, we used the "Analyze Particles" plugin with the size range of 400-2500 and circularity range of 0.50-1.00. After correcting the background using a rolling value of 50px and measuring the signal intensity of ROIs in both channels, we quantified colocalization by multiplying the pixel values of the two channels with the 'multiply create 32-bit' command.

2. Deep learning analysis

2.1 Data preparation for model training

All neural net models were trained using the *Keras* API (version 2.2.5.0) for *Tensorflow* (version 2.00) in R (version 3.6.3)[62]. The training data with associated state labels was compiled from *atg1Δ* or WT cells under the experimental conditions where autophagic or non-autophagic features were uniformly maximized across the cell culture populations (Fig. S1A). *atg1Δ* cells across the entire experimental protocol or WT cells in T0 or from 5 hours in nitrogen replenishment were assigned state 0 (non-autophagic), while WT cells from 8 hours to 12 hours of nitrogen starvation were assigned state 1 (autophagic). Although autophagic features, including free GFP-Atg8 puncta, and colocalization of GFP-Atg8 and Pep4-mCherry, could be observed earlier in WT cells, 8 hours of starvation represented a demarcation where GFP-Atg8 and Pep4-mCherry were clearly overlapping homogeneously across all WT cells, representing that autophagosomes had been delivered to the vacuole. Experimental outliers visibly deviating

from these standards were manually removed from the training data. Also, cells with low GFP-Atg8 signal (mean intensity < 150) or low (mean intensity < 120) or high (mean intensity > 5000) Pep4-mCherry signal were removed from the training data.

Single-cell parameters for variations in signal distributions from both color channels and the co-occurrence were used as input features. Prior to training, we computed the mean and standard deviation feature vectors from the entire WT dataset, including all timepoints, which were used to transform the training data features and transforming all future new data. The resulting training set included 1 073 981 observations in state 0 and 269 655 observations in state 1.

2.2 Model training, hyperparameter search and model selection

To train the models binary cross-entropy was used as loss and the models were trained using the AdaGrad algorithm, with a batch size of 50 000, learning rate of 0.01, and patience of 5 epochs. In the hyperparameter search, 10% of the data was kept out for testing. Of the remaining dataset 80% of the data was used for training the models and 20% was used as validation data for early stoppage. To find the optimal model architecture a grid search was performed over all combinations of activation functions (*relu, elu, tanh, sigmoid*) with networks of different sizes (Fig. S1A). The networks were grown automatically by defining a number of layers (1-5 and 6-8, respectively) and neurons in the first layer (60-120 and 80-120, respectively, with increments of 5) and adding the layers with a decay rate in the number of neurons from the former layer (0.5-1 and 0.7-1, respectively, with increments of 0.05), with a minimum layer size of 10 neurons. This resulted in 3096 models that were trained for a maximum of 150 epochs (with early stoppage). The models were evaluated based on the test set prediction averages compared respective state labels, training and validation set loss and accuracy, and model overfitting (Fig. S1B-D). With exception of *sigmoid*-based models, which underperformed compared to the other models, most architectures resulted in similar test performances (Fig. S1B). Thus, the *sigmoid*-based models were excluded from further analyses.

Model complexities were quantified from the number of model weights. Since model test and validation performances were asymptotically scaling with model complexity, while more complex models also tended to have more overfitting for some architectures (*relu*), we defined a model complexity cutoff at the smallest model in the top 1% (validation loss; Fig. S1D). Moreover, network size may affect the rate of convergence and generalization of a model for complex function approximations, where 150 epochs may not be enough. Subsequently, we selected the top-performing 5 models (validation loss) among the least and most complex models (given the complexity cutoff) for *relu*, *elu*, and *tanh* activation functions, respectively (Fig. S1E). These 30 models were re-evaluated by re-training them for a maximum of 3000 epochs (with early stoppage) using all of the data in a 80/20 training and validation split. The models were evaluated based on training and validation set loss and accuracy, as well as the between-experimental correlation of predicted autophagy dynamics (plate averages for WT, *atg1Δ*, and *vam6Δ* data points combined). On average, the less complex models slightly outperformed the more complex ones for architectures using *tanh* and *elu*, but not *relu* activation functions (Fig. S1E).

Additionally, the predicted last latent layers (autophagic feature vectors) for the top 2 models (validation loss) for each activation function, were extracted for 500 000 randomly sampled WT, *atg1Δ*, or *vam6Δ* cells from any time point and embedded with uniform manifold approximation and projection (UMAP; Fig. S1A), using the *umap* package (version 0.2.7.0) in R[63]. The UMAP was used to force a low-dimensional and regularized representation of the

predictive information in the neural networks and to investigate the relative contribution of noise and extracted autophagic features between three distinct autophagy phenotypes. The UMAP distance metric was set to Pearson correlation, and n_neighbors were set to 10, 15, or 30. The between-experimental correlation of predicted UMAP dynamics (coordinate-wise plate averages for WT and *vam6Δ* data points combined) was used to test latent space robustness (Fig. S1E). The two highest performing and most robust models across multiple metrics (*elu* model 30 - 7 layers, 90 neurons in layer 1 and 66 neurons in layer 7; *tanh* model 22 - 8 layers, 115 neurons in layer 1 and 37 neurons in layer 8) were deployed for GW predictions, and used to corroborate each other (Fig. S1H). Statistical analyses of percent autophagy response dynamics (Fig. 1-3 and 'Autophagy %' results of Fig. 5) were based on predictions from model 30. For latent space analyses, model 22 had less noise and were primarily used to visualize the UMAP dynamics (Fig. 4), but all statistical analyses and inferences from the UMAP distributions were averaged or corroborated by predictions from both models (Fig. 4-5).

2.3 Parametric UMAP

Since transforming new data with UMAP is slow, we implemented a fast parametric transformation of the inputs (autophagic feature vectors) to the embedded coordinates using a neural net (pUMAP: fully connected, 8 layers, 120 neurons/layer, *tanh* activation functions, with bivariate linear output, and trained using the adam optimizer with MAE loss; Fig. S1A). The specific implementation here was conceived before the publication by Sainburg et al. 2020[64], but follows the same essence.

2.4 Genome-wide predictions and quality control

New screen data were filtered for cells with low (mean intensity < 120) or high (mean intensity > 2000) Pep4-mCherry signal. A more stringent upper bound was used for the red channel in order to ensure the removal of dead autofluorescent cells. The input features were z-transformed using the feature mean and standard deviation from the training data, and the probability for cells activating autophagy (P_a) was predicted using the neural nets. Subsequently, the expected probability of autophagy (predicted autophagy %), $A\%_t = \frac{100}{n} \sum_{i=1}^{n} P_{a,i,t}$, was computed for the mutant cell population at every time point. Alternatively, we also computed the autophagy classification %, $A_{binary}\%_t = \frac{100}{n} \sum_{i=1}^{n} I(P_{a,i,t} > 0.5)$, which strongly correlated with $A\%_t$, but was not used as it was less robust for smaller cell populations. The expected standard deviation per cell was computed as $\sigma_{A,t} = \frac{100}{n} \sum_{i=1}^{n} \sqrt{P_{a,i,t}(1 - P_{a,i,t})}$, to assess the prediction uncertainty. We also computed the classification uncertainty using the entropy, $S_A = \frac{1}{n} \sum_{i=1}^{n} [P_{a,i} log(P_{a,i}) + (1 - P_{a,i}) log(1 - P_{a,i})]$ (See Data File S2, *autophagy % DNN predictions*).

      Outlier plates were identified by computing the IQR from all plate median responses and finding plates that had a mean absolute deviation greater than $2IQR$ over the time series. These plates were subsequently repeated. We also identified mutant outliers based on the 99% high

density region of average cell numbers and expected standard errors over the time course (Fig. S2E), which were subsequently repeated in new experiments with higher initial cell seeding concentration. Four DAmP plates that did not have the replenishment response, were manually removed from genome-wide analyses and clustering procedures.

3. Statistical analyses

3.1 Double-sigmoidal model fitting

Double-sigmoidal models were fitted to the data using the *sicegar* package (version 0.2.4) in R (version 4.2.1)[25]. This model includes six parameters (maximum autophagy, final autophagy, two transition midpoints/T50s, and two slope parameters) (Fig. 1C). Tangent lines crossing the transition midpoints (T50s) were computed to extract additional parameters describing the dynamic part of the curve. To de-noise, interpolate potential missing time points, and standardize responses with minor shifts in imaging times, the models were also used to predict the response for all mutants for the entire time series with 30-minutes increments. The goodness-of-fit was assessed using the log-likelihood (on normalized data) and RMSE (See Data File S2, *autophagy % curvefits* and *goodness-of-fit measures*).

3.2 Hypothesis testing and perturbation parameter statistics

Using the double-sigmoidal model predictions, autophagy perturbations were computed from the difference between the interpolated curve fits ($A_f\%$) for all mutants (m) and the corresponding plate medians (ctr), $\Delta_f\%_{m,t} = A_f\%_{m,t} - A_f\%_{ctr,t}$, and subsequently averaged per phase (Perturbation -N or +N) or over the entire time series (Perturbation overall; Fig. S3D). Equivalently, all the mutant parameters were centered by subtracting the corresponding plate medians to compute parameter perturbations, $\Delta parm_m = parm_m - parm_{ctr}$ (Fig. S3E).

Autophagy perturbation/parameter errors were computed for each of the plate controls (WT, *atg1Δ*, and *vam6Δ*), by subtracting their respective plate-wise average. Then, the WT errors were used as H_0-references for each parameter by fitting the error distributions using the *locfit.raw* function of the *locfit* package (version 1.5-9.8) in R[65], and assessing significance of the parameters by computing p-values from the re-sampled fitted H_0-densities. To control for the false discovery rate (FDR) the parameter p-values were adjusted using the Benjamini-Hochberg method (see Data File S2, *parameter statistics and parameter controls*)

3.3 p-value aggregation and mutant statistics

Due to the high correlation between the different perturbation parameters, causing a lack of independence between the individual hypothesis tests (non-independent p-values), we used a harmonic mean p-values (HMP) technique to aggregate the p-values and score the overall significance of the mutants, while controlling for the false discovery rate[26]. The HMPs were computed as $HMP = \sum_j w_j / \sum_j \frac{w_j}{p-value_j}$, where weights corresponded to the H_0-rejection rate at a 5% Benjamini-Hochberg (BH) correction cutoff, $w_j = F_j(FDR_{BH} \leq 0.05)$, in order to favor the

more likely alternative hypotheses based on the genome-wide data. This grouping of hypothesis tests maintained sensitivity of the least stringent BH-adjusted p-value (FDR min) or improved the sensitivity for mutants with multiple significant tests. HMPs based on equal weights were also computed, but in comparison did improve the power over individual BH-FDR corrections per mutant (see Fig. 1D and S4B and Data File S2, *mutant profiles and parameter groups*).

3.4 Mutant profiling and clustering

Statistically significant mutants, HMP ≤ 0.01, were extracted for clustering. The parameter perturbations were clustered and grouped by computing the log_{10}-transformed p-values and performing a hierarchical clustering using Pearson correlation as a similarity metric and ward.D2 clustering. The parameters were split into five groups (Fig. S4D) using the *cutreeDynamic* function from the *dynamicTreeCut* package in R[66].

To group the mutants into distinct dynamic profiles, the response predictions from the double sigmoidal models for mutants with HMP ≤ 0.01 were subjected to hierarchical clustering using Euclidean distances and ward.D2 (Fig. S4E). For mutants with multiple significant independent experimental measurements the median responses were computed before the analysis. Groups were identified with the *cutreeDynamic* function.

4. Screen quality assessments

4.1 Recall of reference mutants

'Gold-standard' test sets for evaluating the sensitivity and accuracy for retrieving true autophagy perturbation phenotypes, were defined through manual inspection of images and literature curation (Table S1)[5,23,24,55,67–74]. This included two true positive test sets, including genes involved in autophagosome formation ('ATG set', n=16) and clearance ('Fusion set', n=19), respectively. A negative control set was also developed from several independent replicates of two dubious ORFs[75] that produced WT autophagy responses (n = 23) (Table S1). None of the mutants included in the test sets were part of the training data or used as reference data for the UMAPs.

Precision-recall curves (PRC) and receiver operating characteristic (ROC) curves were computed from the p-value rank orders for the different perturbation parameters, comparing the true positive sets ('ATG set', 'Fusion set', or combined) with the negative control set (Fig. S5C, E), and area under the curves (AUCs) were computed to score the precision and accuracy of recalling true positives (Fig. S5B,G). Alternatively, negative sets with equal size to the true positive sets were bootstrapped by repeated random sampling with replacement from the GW population (b=1000) for a more statistical assessment of the different perturbation parameters with respect to the average deletion mutant phenotype (Fig. S5A, B and S5D, F, G). For mutants with multiple independent experimental measurements the median p-values or HMP were computed before the analysis. To use ROC curves for a direction-specific separation of the test sets from the GW population average, the p-values were log-transformed and signed by the direction of the perturbation (Fig. S5F, G).

4.2 Gene set enrichment analysis (GSEA)

GSEA was performed using the *gseGO* function of the *clusterProfiler* package (version 4.4.4) and *org.Sc.sgd.db* (version 3.15.0) in R (version 4.2.1)[76], with 10^5 permutations, and min and

max gene set size set to 5 and 200, respectively. To analyze the directional association of GO-BP (gene ontology - biological processes) terms for the individual parameter perturbations, we used the log-transformed p-values signed by the direction of the perturbation. For mutants with multiple independent experimental measurements the median p-values were computed before the analysis. Subsequently, enriched GO-BP terms with at least one test p-value < 0.005, were analyzed across the multivariate parameter space using PCA (Fig. 2A, B) and hierarchical clustering (Fig. S9A). In both cases, we used a matrix with the signed log-transformed enrichment p-values for the selected GO-BP terms for each parameter (see Data File S3). Hierarchical clustering was performed using Euclidean distances and ward.D2.

To generate an enrichment map (Fig. 2C and S9A), we performed the same GSEA over the -log-transformed HMP values that were zero-centered around -log(0.01). For mutants with multiple independent experimental measurements the median HMP was computed before the analysis. Here, all GO categories BP, MF (molecular function) and CC (cellular compartment) were included. GO terms with p-values < 0.05 and NES > 0 were graphed based on the Jaccard correlation coefficient (JC) for scoring the GO term similarities and a minimum JC > 0.01 for each edge, using the *enrichplot* (version 1.16.2) and *clusterProfiler* packages in R. The Spearman correlation over the profile count matrix for the GO terms was used to quantify the functional co-occurrence and purity of the different profiles in the enrichment map (Fig. 2D).

5. Network analyses

5.1 Spatial analysis of functional enrichment (SAFE)

SAFE was performed on the SGA-PCC network as published by Costanzo et al. 2016 (Fig. 3A)[27]. The SGA-PCC network was projected as in Costanzo et al. using the edge-weighted, spring embedded implementing Kamada-Kawai force-directed algorithm and loaded in Cytoscape (v.3.9.1)[77]. We also prepared a STRING network, based on a combined_score ≥ 990 of the STRING v10 database, and projected using the unweighted spring embedded layout algorithm (Fig. S10A, see Data File S4, *Costanzo and STRING node-edge*). Both networks were annotated with the GO-BP reference data from Costanzo et al. using SAFE[27,30]. The SGA-PCC network displayed a coverage of 84% and an attribute coverage of 56% and the STRING ≥ 990 network displayed a coverage of 99% and an attribute coverage of 66%. The enrichment landscape was built with a Jaccard correlation value of 0.75 and the resulting reference map labels were manually curated according to the functional domains identified. We also used SAFE to annotate the ATG core sub-graph using the 'ATG set' including ATG1 and ATG8, and the fusion core sub-graph using the entire tethering complex (including VTI1 and YKT6), the HOPS complex, and VPS17, VPS29, VPS5, VPS35, VAM10, VPS1, VSP45, VPS3, VPS36 and VPS26. The profiles were enriched using SAFE and the enrichment colors were scaled with the SAFE score ranging from a min=0.35 to a max=0.35 (See Data File S4, *Costanzo and STRING SAFE*)

5.2 Network distance analyses

The shortest network path from any gene to the ATG core machinery (ATG set) was computed by finding the set of genes one degree away from any gene within the ATG set, and iteratively finding the set of genes one degree away from any gene within the previous set, until all possible paths had been found (Fig. 3B). For the shortest causal network path, connector genes within the

previous set were required to have a significant phenotype given by HMP < 0.01 or an autophagy perturbation p-value$_{min}$ < 0.01.

5.3 Gene-pair correlations

Correlations in phenotypes between genes were computed using the Pearson correlations of autophagy perturbation dynamics, $PCC_{\Delta,\{m1,m2\}} = cor(\Delta\%_{m1,t}, \Delta\%_{m2,t})$, or equivalently using signed -log$_{10}$-transformed p-values over the set of kinetic parameter hypothesis tests performed. The correlations were subsequently grouped based on network interaction or complexome association status of the gene pairs as well as significance status (HMP ≤ 1, 0.01, or 0.001).

For strongly significant gene pairs (HMP ≤ 0.001) the correlations in perturbation dynamics were stratified into intervals for SGA-PCC concordance testing. The three intervals considered positive correlations (PCC$_\Delta$ > 0.25) representing similar phenotypes, negative correlations (PCC$_\Delta$ < -0.25) representing opposite phenotypes, and no correlations. Enrichment of SGA-PCCs was computed for each interval given different positive or negative SGA-PCC thresholds (0.1, 0.125, 0.15. 0.175, 0.2). Enrichment was calculated as $E_{k,l} = \frac{N(PCC_\Delta=k, SGA\cdot PCC>l)}{N(PCC_\Delta=k)} \times \frac{N}{N(SGA\cdot PCC>l)}$, for a PCC$_\Delta$ interval $k = (lower, upper]$ and a SGA-PCC threshold l (here indicated as greater than, but lesser than for $-l$). A Fisher's exact test was used to compute the p-values for over- or underrepresentation.

5.4 Directional GSEA along SGA-PCC spectra

Directional, positive or negative associations of autophagy perturbation profiles with SGA-PCC spectra for specific core machinery genes were computed using a modified GSEA algorithm (Fig. S11C). A positive or negative enrichment score (ES) was computed as:

$$ES_+ = max(F_c^{GS} - F_c^{\overline{GS}}) \text{ and } ES_- = min(F_c^{GS} - F_c^{\overline{GS}}).$$

Here the F represent the cumulative distribution functions for the ranked SGA-PCC scores (x)

$$F_c^{GS} = \sum_{i=1}^{c} abs(x_i)1_{\{m_i \in GS\}} / \sum_{i=1}^{N} abs(x_i)1_{\{m_i \in GS\}} \text{ and } F_c^{\overline{GS}} = \sum_{i=1}^{c} 1_{\{m_i \in \overline{GS}\}} / \sum_{i=1}^{N} 1_{\{m_i \in \overline{GS}\}},$$

where 1 represent a indicator function for whether a mutant m_i is present in the perturbation profile gene set (GS) or among the non-gene set mutants ($\overline{GS}$). Direction preferential enrichment was computed as $\Delta ES = ES_+ + ES_-$. For statistical evaluation we used a permutation test where the distribution of GS membership along x were randomized 500 times to generate respective H0-distributions of rES_+, rES_-, and ΔrES, which were used to compute respective p-values, as well as the normalized enrichment scores $NES_+ = \frac{ES_+}{\langle rES_+ \rangle}$, $NES_- = \frac{ES_-}{\langle rES_- \rangle}$, and $\Delta NES = NES_+ - NES_-$. NES values were grouped for genes belonging to the ATG core and the fusion core machinery respectively, and profile enrichment per group was tested using a one-sample t-test.

5.5 Module networks

The module networks were generated by identifying complexes through the Complex Portal[33] or applying the MCODE algorithm[78] to the STRING network (combined_score $\geq$ 990)[32], with multiple significant genes. These complexes were then manually curated for protein-protein interactions using cross-references from SGD, STRING and BioGRID databases[31,32,79]. The networks were embedded in Cytoscape using the prefuse force-directed open layout algorithm and subsequently manually organized with node colors scaling with the signed -$\log_{10}$ p-values for the overall perturbation and size of the nodes scaling with their HMP (-$\log_{10}$) (see Data File S5).

6. Latent-space and Bayes factors analysis

6.1 Statistical analysis of the UMAP latent space

pUMAP predictions were performed on the latent autophagy feature vectors for all cells. The embedded cell densities were plotted by grouping populations per time points and color coding either for time or predicted autophagy % ($A\%_t$). Moreover, the population means ($\mu_{U1,t}$, $\mu_{U2,t}$) and standard deviations ($\sigma_{U1,t}$, $\sigma_{U2,t}$) were computed for each UMAP coordinate (x_{U1} and x_{U2}) per time point.

The UMAP embeddings were analyzed for representation of predictive information, population dynamics and noise. To test whether the single-cell UMAP embeddings conserved the information about autophagy from the DNNs, we trained a bag of regression trees to predict the DNN outputs, either represented as the probability of autophagy P_a or its log-odds, $LO = log\frac{P_a}{(1-P_a)}$. For this we used the *XGBoost* package (version 1.7.5.1) in R to train 50 parallel trees with subsampling = 0.63, depth = 10 and min_child_weight = 3[80]. The regression was performed per plate, where the data was split to 70% training data and 30% test data, and a PCC between the predicted and true values was reported. For comparison, the same procedure was used to test representation of time. We also compared these results with the ability of the UMAPs to separate WT from vam6Δ cells, by using bags of classification trees (with the same hyperparameter settings) subjected to 10-fold training and testing over the entire dataset, and scoring the concordance between the predicted class and the true label using an F1 score. To assess whether the information distinguishing these phenotypes were due to latent information about intermediate autophagic states in the UMAP or simply due to distributional differences based on the overall DNN prediction for the two strains, we stratified the UMAP distributions into percentiles based on the predicted autophagy P_1 or LO, and then performed the same training and testing procedure within each interval.

To measure UMAP latent space noise and flux, we first calculated the population bivariate standard deviation, $\sigma_{U,t} = \sqrt{\sigma^2_{U1,t} + \sigma^2_{U2,t}}$, and the average population displacement, $\Delta_{U,t} = \sqrt{\Delta_{U1,t}^2 + \Delta_{U2,t}^2}$, per time point. Here, the population displacement per coordinate, for instance $\Delta_{U1,t}$, is given as $\Delta_{U1,t} = \mu_{U1,t+1} - \mu_{U1,t}$. $\Delta_{U,t}$ and $\sigma_{U,t}$ were compared with equivalent metrics computed from the unembedded latent feature vectors directly for populations of greater than 100 cells (Fig. S12D). For this comparison outliers for the latent

space metrics beyond the range given by the quartiles Q1/Q3 -/+ 1.5 IQR (interquartile range) were filtered out.

As a measure of signal-to-noise we also computed the normalized flux $n\Delta_{U,t} = \sqrt{(\frac{\Delta_{U1,t}}{\sigma_{U1,t}})^2 + (\frac{\Delta_{U2,t}}{\sigma_{U2,t}})^2}$. The overall population variability and flux was assessed by averaging $\Delta_{U,t}$, $n\Delta_{U,t}$ and $\sigma_{U,t}$ for each mutant time series per treatment phase. For this comparison only observations with an average of 50 cells were evaluated (see Data File S6).

6.2 Bayes factor computation and assessment

Reference UMAP states of WT, *atg1*Δ, and *vam6*Δ cell populations were collected from 21 independent time series across 7 plates (Mix8, Mix10, and Rec1-5), where the WT cells comprised the dubious ORF negative controls in order for the latent space distributions to match that of unperturbed H_0-outcomes of the KO collection. Kernel density estimators (KDE) were fitted using the *kde* function of the *ks* (Kernel Smoothing) package (version 1.13.2) in R (version 4.2.1)[81]. The KDEs were computed for the reference states based on their overall distributions independent of treatment phase (K_{WT}, $K_{atg1\Delta}$, $K_{vam6\Delta}$), or their distributions stratified per time-point ($K_{WT,t}$, $K_{atg1\Delta,t}$, $K_{vam6\Delta,t}$) in order to perform timewise phenotype comparisons. By using the *dkde* function we computed the probabilities $P(D|K_S)$, which represented the marginal likelihoods for UMAP observations D (U1, U2) given a reference state S. Thus, ratios of the marginal likelihoods comparing any two given states represented Bayes factors, $BF(WT:atg1\Delta)$, $BF(WT:vam6\Delta)$ and $BF(vam6\Delta:atg1\Delta)$ that summarized the evidence favoring a given autophagic reference state in a specific UMAP (Fig. S14A,C). For time-independent phenotype classification of mutants between reference states A and B, the time-wise KDEs were paired with data for the corresponding time points to compute time-wise likelihoods. The individual KDEs were computed with equal matrix bandwidths generated by taking the medians of the bandwidth selection results using the *Hpi.diag* function on each time point. Subsequently, the log-likelihood ratios were averaged per mutant over all cells and all times points:

$$log\ BF(B:A) = \frac{1}{n}\sum_{i=1}^{n}\sum_{t=1}^{T}log\ \frac{P(D_{i,t}|K_{B,t})}{P(D_{i,t}|K_{A,t})}.$$

This population average or expected log Bayes factor can be viewed as a time-independent overall phenotype comparison per mutant, where dynamical effects have been 'integrated' out with the moving KDEs. This averaging was done for the entire experimental time course (overall), as well as for each treatment phase individually. On the other hand, to quantify the execution 'activity' as a timewise transition between temporally fixed reference states, the expected log-likelihood ratio was computed with the time-invariant KDEs over a cell population per time point:

$$log\ BF_t(B:A) = \frac{1}{n_t}\sum_{i=1}^{n_t}log\ \frac{P(D_{i,t}|K_B)}{P(D_{i,t}|K_A)}.$$

Within this Bayes factor framework, given the execution model $\{A \to B \to C\}$ ($ATG1 \to VAM6$: autophagosome formation; $VAM6 \to WT$: autophagosome clearance), the overall activity for the transition $\{A \to C\}$ can be factorized into the intermediate transitions ($\{A \to B\}$ and $\{B \to C\}$) as

$$log\,BF_t(C\colon A) \;=\; log\,BF_t(C\colon B) \;+\; log\,BF_t(B\colon C).$$

For comparison with the time-independent Bayes factors, and to score the expected activity over a given time frame, these timewise Bayes factors were averaged per mutant for all cells over a given time frame as well. This was done for the entire experimental time course (overall), as well as for each treatment phase individually (see Data File S7). Because these Bayes factors represent comparisons between fixed reference states, they are more sensitive to dynamical variation among the mutants, and hence indicated with the t-subscript. All time-averaged Bayes factors were corrected for distributional biases across plates by subtracting the plate-wise median difference from the screen median:

$$log\,BF(B\colon A)_m \;=\; log\,BF(B\colon A)_m \;-\; [log\,BF(B\colon A)_{plate} \;-\; log\,BF(B\colon A)_{screen}].$$

Analysis was restricted to mutants with at least n = 100 cells for time-averaged Bayes factors overall, and n = 50 cells when averaging per treatment phase. For mutants with multiple independent experimental measurements the median Bayes factors were computed. ROC curves and AUCs for analyzing retrieval of test gene sets ('ATG set' and 'Fusion set') were conducted as described above. Potential overfitting of the density kernels to the reference distributions was assessed by comparing the Bayes factors between *atg1Δ* and the 'ATG set', and between *vam6Δ* and the 'Fusion set'. To compare the performance (ROC analysis) of the Bayes factors directly with the predicted autophagy perturbations (not subjected to sigmoidal curve fitting), $\Delta\%_{m,t} = A\%_{m,t} - A\%_{ctr,t}$ and $\Delta_{binary}\%_{m,t} = A_{binary}\%_{m,t} - A_{binary}\%_{ctr,t}$ were computed per mutant and averaged over the entire time series. Comparison with the fluorescent signal co-occurrence per experimental measurement was done by computing the mean co-occurrence early (0-1 hour) and late (> 7 hours) of the starvation protocol (Fig. S14A). Functional analyses of Bayes factors were performed over the KO library only. We removed Bayes factors for experimental measurements where the mean red or green fluorescent signals early (0-1 hour of starvation) were below the 25th percentile - 1.5IQR of the GW screen distribution. For functional analyses we also computed the mean of the Bayes factor from the two neural latent space models (model 22 and model 30). To assess the concordance in phenotype classification between treatment phases, we paired the treatment phase specific Bayes factors and computed the fraction of deletion mutants with with substantial evidence for the same reference state (either log $BF > 5$ or log $BF < -5$) across both phases (Fig. S14C). Analysis of lagged correlations for time-wise Bayes factors ($log\,BF_t$) were computed using the deletion library only.

6.3 Parametric gene-set enrichment analysis

Parametric GO (BP and CC) enrichment analyses of Bayes factors were performed using the *piano* package (version 2.8.0) in R (version 4.2.1)[82], using the gene sampling permutation test (nPerm = 10^5) of the mean gene set statistics. The genesets were retrieved and annotated using the *GOdb* package (version 2.9) and org.Sc.sgd.db (version 3.15.0). Only the KO library phenotypes were analyzed, including the median of all *atg1Δ* and *vam6Δ* phenotypes observed. The Bayes factors or their respective treatment phase differentials were standardized over the

GW population (to compute log-BF z-scores) prior to the analyses (see Data File S8, *z-scores*).

For analyzing the BF_t enrichment, the directional statistics (representing average log BF_t z-score per gene set) were visualized for gene sets with a p-value < 0.05 for at least one Bayes factor. For analyzing the BF differential enrichment, the directional statistics were visualized for gene sets with a differential enrichment p-value < 0.05. For the heatmap the gene-sets with a differential enrichment p-value < 0.01 for BF VAM6:ATG1 or BF WT:VAM6 and a minimum absolute statistic of one. Euclidean distance and ward.D2 were used for the clustering.

7. Experimental validation of autophagy predictions

7.1 Yeast and growth condition details

S. cerevisiae strains were grown at 30 °C until mid-log phase in standard yeast extract peptone dextrose (YPD) medium or in synthetic medium supplemented with relevant amino acids (SC) before proceeding with genetic manipulation and experimental assays. See *strain*s tab and *purpose* in Data File S1 for strains used in this section.

7.2 Reproducibility library and strains construction for autophagy assays

All strains used in this section were derived from the S288c strain BY4741[83] using either PCR-targeted gene replacement methods or crossing. Standard cell culture and genetic techniques were employed according to protocols[84,85]. Unless, otherwise noted, all genetic modifications were carried out chromosomally. Deletions and promoter replacements were performed as described[58].

For creation of the "reproducibility" set, *ATG8* was first N-terminal tagged and expressed under the ScTDH3 (GDP) promoter using the plasmid pYM-N17-mNGreen. PEP4 was then c-terminal tagged with mCherry using the plasmid pBS35. Finally, all deletion strains and ORF control *YEL074W* were deleted using the plasmid pFA6a-his3MX6 in the resultant strain that contained aforementioned alleles (See Data File S1; *strains* tab and *reproducibility* purpose).

For the quantitative Pho8Δ60 assay, the PYM-N15 plasmid was used to delete the first 60 amino acid residues from the amino terminus of the phosphatase Pho8, and to express the truncated ORF under the control of the ScTDH3 (GPD) promoter (See Data File S1; *strains* tab and *autophagy assays* purpose)[86,87].

For the GFP-Atg8 processing assay, yeast cells were transformed with a plasmid expressing GFP-Atg8 under the control of the endogenous ATG8 promoter (pRS416 GFP-ATG8/AUT7; See Data File S1; *strains* tab and *autophagy assays* purpose)[88,89]. All strains were confirmed by PCR and fusion proteins were additionally validated by fluorescent microscopy and Western blot analysis. See Data File S1 for primer sequences.

7.3 Growth and starvation protocols for screen reproducibility analysis

The screen reproducibility analysis comprised two approaches as described (See section Screen reproducibility analyses). In the first, aimed at replicating starvation observations in the initial screening, strains were grown sequentially to logarithmic phase, following an identical protocol as in the initial screening (See section 1.3- cell growth protocol for high-content screening).

In the second approach, focused on evaluating the uniformity of autophagy phenotypes across various experiments and strain backgrounds, the growth protocol was adjusted to induce

autophagy with cells in stationary phase. Cells were grown in SC at 25 °C for three consecutive periods of 48 h each, followed by 8 h at 25 °C in the imaging plates, before being transferred to SD-N to induce nitrogen starvation for 12 h and replenishment for 7 h. Live imaging was performed at 30 °C, capturing images every hour for 19 h.

7.4 Screen reproducibility analyses

7.4.1 Nearest neighbor analysis and repetition set selection. Autophagy perturbation gene sets of different sizes were generated by setting various HMP cutoffs ($\leq$ 0.01, 0.005, 0.001, or 0.0005; see Fig. S6B). Subsequently, we performed a Fisher's exact test for overrepresentation of autophagy perturbations among the nearest neighbors using different GW yeast network data, including BioGrid PPIs, SGA-PCCs (top 1%), as well as all STRING interactions or only STRING-PPIs with a combined score > 500. These overrepresentation tests resulted in eight p-values, from which a harmonic mean was computed to yield an overall network enrichment significance (network p-value). Potential false positives (FPs) were defined as mutants with significant autophagy perturbations (perturbation FDR_{min} < 0.01; min referring to the lowest value between -N and +N), but non-significant network enrichment (network p-value > 0.1). Potential false negatives (FNs) were defined as mutants with non-significant autophagy perturbations (perturbation FDR_{min} > 0.1), and significant network enrichment (network p-value < 0.01). Potential FNs were also selected based on genes with key autophagy GO-term annotations (see Fig. S6C) and non-significant phenotype (FDR_{min} > 0.1 and HMP > 0.01). Top hits were selected by rank-ordering mutants based on the HMP (See Data File S9 and Data File S10).

7.4.2 *Replication of repetition set and outliers.* Only mutants from the KO library were used for the validation screen, due to the lack of WT DAmP controls. This included 109 potential FNs, 56 potential FPs, and 75 top hits (See Data File S1, *repetition library* and Data File S10, *repetition responses*). For the validation screen perturbations in predicted autophagy ($\Delta\%_{m,t}$) were computed from the mean WT ORFs included in each plate (since plate medians would not be valid WT references due to the enrichment of mutant phenotypes). The replicability was tested by repeating the GW screen protocol up to 8 hours of starvation, and computing the correlation in predicted autophagy dynamics ($PCC_{A,m} = cor(A\%_{m,t,R1}, A\%_{m,t,R2})$) and autophagy perturbation dynamics ($PCC_{\Delta,m} = cor(\Delta\%_{m,t,R1}, \Delta\%_{m,t,R2})$) for each mutant between replicates R. The dynamics correlations were compared with the mean absolute autophagy perturbations, $|\Delta A\%|_m = \frac{1}{T}\sum_{t=1}^{T} abs(A\%_{m,t} - A\%_{ctr,t})$. For mutants with multiple independent experimental measurements in the GW screen the median responses were computed before the comparisons.

7.4.3 *Reproducibility across experiments, strain background and conditions.* To assess the reproducibility of autophagy phenotypes across strains-background, independently generated clones, and growth conditions, 33 mutants covering different cellular functions and with different autophagy dynamic profiles were picked manually. All mutants were strongly significant (HMP $\leq$ 0.001). Three independently generated new clones were included for each mutant, along with the collection clone, and three new screens were done independently (See Data File S10, *validation responses*; see Fig. S7A-B).

For evaluation of mutant replication error, the mean absolute perturbations were computed per mutant and compared with the mean absolute errors,

$$\epsilon_{A,m} = \frac{1}{T}\sum_{t=1}^{T} abs(A\%_{m,t,R1} - A\%_{m,t,R2}),$$ between different types of replicates R. These included technical replicates (same mutant clone, different wells within the same plate), experimental replicates (same mutant clone, different plates/experiments), replication of mutant phenotypes generated in a new background strain, and replication between independent clones generated within the same background strain. Correlations in autophagy dynamics (PCC_A) and autophagy perturbation dynamics (PCC_Δ) were also computed per mutant across experimental replicates and between collection clones and clones generated in a new background strain. Mutant perturbations in autophagy on the new screen plates were computed with respect to the average from 12 independent WT controls, 6 based on the collection background strain and 6 based on the new background strain.

For assessing screen-level experimental reproducibility of autophagy perturbations and kinetic parameters, Pearson and Spearman correlations (due to large outliers for some parameters) were computed from global pairwise comparisons of the screen metrics across all mutant clones. Comparisons were done between the GW screen and the new screens for autophagy perturbations, $\Delta\%_t$, grouped on treatment phase (-N, +N, or both) and strain background (collection clones, new clones, or both), or between the new screens across all clones for the parameters (See Data File S10, *validation statistics*).

7.5 Rapamycin-dependent autophagy response and analysis

7.5.1 Experimental cell growth and conditions. Cells were grown in SC at 25 °C for two consecutive periods of 48 h each, followed by a period of 24 h and a last for 8h at 25 °C in the imaging plates. Cells were then washed twice with sterile MilliQ water and transferred to independent plates containing 200ul of SC+Vehicle, SC+Rapamycin, SD-N+Vehicle and SD-N+Rapamycin. Live imaging was performed at 30 °C, capturing images every hour for 10 h. Rapamycin was dissolved in ethanol and Triton-X in a 9:1 ratio in a concentration of 1 mg/ml. The final concentration of rapamycin used was 440 nM. As a vehicle control, the solvent of rapamycin was added in the same proportions. Each plate included duplicate experimental conditions of every strain (triplicates for the controls), and the complete experiment was repeated three times.

7.5.2 Data Analysis. Autophagy ($A\%_t$) and UMAP Bayes factors (BF_t) were computed for each well (see Data File S11). Autophagy perturbations ($\Delta\%_t$) were computed with respect to the mean of ORF plate controls, and for all metrics the time series averages were computed. Reproducibility between experiments was assessed by computing PCC for each metric.

7.6 Yeast growth conditions for protein and RNA analysis

To correspond to library growth conditions, cells were serially diluted in a fresh medium to maintain them in logarithmic phase before inducing autophagy. Cells were first grown in YPD O/N and transferred to grow in three sequential cultures in SC-URA at 30 °C to mid log-phase for 24 hours. Cells were then washed five times with nitrogen-starvation medium (SD-N: 1.9 g/ml yeast nitrogen base without amino acids and ammonium sulfate and 20 g/L glucose) using membrane filtration and transferred to SD-N at an $OD_{600} = 0.25$. Cells were then incubated for 4 or 8 h at 30 °C according to the experiment. In the experiments with prolonged starvation (8 h),

cells were shifted to SC-URA medium and further incubated for 4 h. Cells were collected at specified time points during nitrogen starvation and replenishment and subjected to protein and RNA analysis. For RNA analysis, samples at t=0 were collected from a portion of the culture that was shifted to fresh SC once more and incubated at 30°C for an additional hour.

7.7 GFP-Atg8 processing assay

The GFP-Atg8 processing assay was performed and interpreted as described[57,89]. According to the experiment, cells were collected at indicated times, and protein extracts were prepared and resolved by SDS-PAGE. Full-length GFP-Atg8 and free GFP were detected by Western blot using an anti-GFP antibody (see below).

7.8 Protein extraction, immunoblotting and antibodies

Cell lysis and immunoblotting were performed as previously described[90]. For protein analysis, 2 x 10^8 cells were collected by filtration, washed with STOP buffer (150 mM NaCl, 50 mM NaF, 10 mM EDTA, 1 mM sodium azide) and frozen on dry ice before extraction. Total protein extracts were prepared by precipitation with trichloroacetic acid (TCA) as previously described [91]. Following gel electrophoresis in 4–20% Criterion™ TGX™ Precast Midi (BioRad), proteins were transferred onto a PVDF membrane using a semi-dry blotting system (Trans-Blot® Turbo™ from BioRad). The membrane was blocked in 5% skimmed milk with TBS-0.1% Tween and incubated with primary antibodies in 5% BSA with TBS-0.1% Tween (for antibody against phosphorylated residues) or BSA plus 5% milk (in all other instances). Western blots were developed using Pierce™ and SuperSignal® Dura ECL detection reagents (Thermo Fisher) and imaged using Chemidoc MP Imaging system (Bio-Rad laboratories). The following primary antibodies were used in this study: rabbit anti-phospho-(Ser/Thr) Akt Substrate Antibody to detect phosphorylated RpS6 (1:1000, Cell Signaling Technology 9611), goat anti-GFP (1:1000, Abcam 5440) to detect GFP-Atg8 and mouse anti-beta Actin (1:1000, Abcam 8224). As secondaries, HRP-conjugated Goat anti rabbit IgG (1: 5000 Sigma), HRP-conjugated Bovine anti goat IgG (1:5000 JIR) and HRP-conjugated Goat anti mouse IgG (1:10000 JIR). The Western blot experiments were performed in replicates as indicated in Data File S12 and representative images were chosen for publishing.

7.9 Quantifications of Western blot with Bio-Rad Lab software

The degradation of GFP-fused Atg8 into the vacuole and the phosphorylation of RpS6 in relation to actin as loading control were estimated by calculating the densitometric values of the bands in the raw digital WB images using Image Lab software from Bio-Rad Laboratories 6.0.1. The WB experiments were performed for multiple independent replicates as indicated in Data File S12 (*see GFP quant. and Rps6 quant.*). See Data File S12 for all quantifications and numerical transformations.

7.10 Autophagy flux and Rps-6-p comparison with BFs

WB quantifications were averaged over two exposure times for every experiment before computing the mean and SEM from the independent biological replicates (see Data File S12, *Bayes Factor comparison*). GFP cleavage was computed from the adjusted band volume (intensity) as the percentage of free GFP signal divided by the total GFP signal (free GFP +

GFP-Atg8) within each lane using densitometric analysis (Bio-Rad Image Lab Software). The relative RpS6-phosphorylation (RpS6-p) was computed from the absolute band volume by dividing the RpS6-p signal by the β-actin signal and normalizing each experiment to their respective WT T0 value.

The time-averaged BFs, kinetic parameters and average screen metrics were compared with the autophagic flux by computing the mean GFP cleavage from 8 hours of starvation (0, 2, 4, 6 and 8 hours). The similarity in autophagy phenotypes measured across mutants were assessed using the Pearson correlation. For time-series comparison of BF_t with GFP cleavage, we matched the time-wise measurements for -N (0, 2, 4, 6 and 8 hours) and +N (0, 1, and 4 hours) for the two time series experiments. Note that +N for the BF_t occurred from 12 hours in -N, while for GFP cleavage +N occurred from 8 hours in -N. Moreover, the 5 min time point in +N for GFP cleavage were used to match with T0 replenishment for BF_t. The time-wise similarity were then assessed using a mixed model based on BF_t WT:ATG1 alone (model 1) or BF_t WT:VAM6 and BF_t VAM6:ATG1 (model 2), with random intercept and slope for each time point:

$$\text{model 1: } y_{GFP\%,t} = \beta_0 + b_{0,t} + (\beta_{BF1_t} + b_{BF1_t,t})x_{BF1_t,t} + \epsilon$$

$$\text{model 2: } y_{GFP\%,t} = \beta_0 + b_{0,t} + (\beta_{BF2_t} + b_{BF2_t,t})x_{BF2_t,t} + (\beta_{BF3_t} + b_{BF3_t,t})x_{BF3_t,t} + \epsilon$$

Here, β represent fixed effects, b represent random effects per time point t, ϵ is the residual error, x_{BF1_t} represent log BF_t WT:ATG1 values, x_{BF2_t} represent log BF_t WT:VAM6 values, and x_{BF3_t} represent log BF_t VAM6:ATG1 values. Goodness-of-fit was assessed by measuring the Pearson correlations between predicted and observed values across mutants for each time point and across time points for each mutant. The time-wise variable (BF_t) contribution to autophagic flux point was computed as $(\beta_{BF_t} + b_{BF_t,t})\sigma_{BF_t,t}$, where $\sigma_{BF_t,t}$ is the standard deviation of the log BF_t sample space at time point t.

For comparing with the Bayes factors with the RpS6-p status, the time-averaged BF_ts were correlated with RpS6-p at T0, average RpS6-p in -N (2, 4, 6 and 8 hours), and average RpS6-p in +N (15 and 30 min, 1, 2 and 4 hours).

Statistical evaluation of genetic interactions effects on GFP cleavage between *rtg1Δ* and *gcn4Δ* or *gln3Δ*, were done using a mixed effects model with random intercept on experiment identity, *rtg1Δ* as baseline and interaction between the mutants and time (see Data File S12, *GFP stats*).

7.11 Quantitative Pho8Δ60 assay

Pho8Δ60 assay was performed as previously described[87,92]. For the analysis of alkaline phosphatase (ALP), cells were first grown to mid-log phase (5 x 10^6 cells/ml) in SC at 30 °C and then diluted to 2.5 x 10^6 cells/ml and shifted to SC for 1h and SD-N for 4h. For sampling, 1 to 1.25 x 10^6 cells were collected by filtration, washed out with 1 ml of PBS pH 7.2 and pellets were frozen on dry ice. For protein extraction, cell pellets were dissolved in 500 μl of lysis buffer at pH 6.8 containing 1 M PIPES, 10% Triton-100X, 1 M KCl, 1M KOAc, 1M MgS04, 10 mM ZnSo4 and 2mM of PMSF. Cell disruption was performed by agitation and homogenization with a cryogenic bead beating grinder (FastPrep®-24 5G). For the phosphatase assay, 0.1-0.5 mg of total protein was used as the phosphatase source, with 125 mM p-nitrophenyl phosphate serving as the substrate in the reaction buffer. The reaction buffer was prepared with 1M Tris-HCl at pH

8.6, 10% Triton-100X, 1M MgSO4, and 10 mM ZnSO4. The mixture was incubated at 37°C for 5 and 7 min, allowing the phosphatase present in the total protein to dephosphorylate the p-nitrophenyl phosphate substrate. The enzymatic reaction was then stopped by the addition of 1M glycine at pH 11. The ALP activity was calculated as the amount of p-nitrophenol (nmol) produced per minute per milligram of total protein. Both incubation times were measured using independent reactions. The Pho8Δ60 assay experiments were performed for multiple independent replicates as indicated in Data File S12 (*see pho8d60 measures*). Data was presented as mean ± SEM. Significant differences were calculated using a one-way ANOVA with a TukeyHSD post-hoc test. Significance is reported for mutant comparisons with the negative control within each treatment phase (see Data File S12, *pho8d60 stats*).

7.12 RNA extraction and quantitative PCR (qPCR)

RNA extraction and quantitative PCR (qPCR) were performed as previously described[90]. For qPCR analysis, 8 x 10^7 cells were collected by centrifugation, washed in DEPC-treated water and frozen in dry ice. Total RNA preparation was performed with MasterPure™ Yeast RNA Purification Kit (Epicentre) following the manufacturer instructions. 1 µg of RNA was used for cDNA synthesis using SuperScript® III Reverse Transcriptase (Invitrogen). qPCR was performed with SYBR® Select Master Mix (Applied Biosystems). Primer sequences (fwd and rv) are listed in Data File S1. RNA analysis experiments were performed in triplicate as indicated in Data File S13 (see *input data*).

7.13 qPCR analysis

Three independent experiments were performed, each with two technical replicates for all targets per treatment condition. The mean C_t value was calculated for each target per treatment condition and subsequently normalized by subtracting the median value of the actin control to compute a ΔC_t value, which was transformed to the signal $S = 2^{-\Delta Ct}$. The complete dataset was batch-corrected with the *ComBat* algorithm from the *sva* package (version 3.44.0) in R[93], using parametric adjustments with experimental replicate as batch variable and genotype as model variable. S was log transformed before the batch correction, and exponentiated after to generate S_{adj}. Finally, S_{adj} was normalized to the mean S_{adj} of the WT controls at T0 and $\log_2$-transformed (See Data File S13, *input data*).

Significance for individual time points was assessed by comparing the mutants with the WT using a t-test. For statistical evaluation of the overall differential regulation effect by the mutants on every target, a regression was performed per treatment phase (T0 +N, -N, and +N), with WT as baseline and genotype and time point as independent variables. The reported values are the coefficients for the genotype effects (See Data File S13, *statistics*)

8. Cross-omics inference for autophagy predictions

8.1 Cross-omics prediction benchmarking

Cross-omics modeling and prediction benchmarking performed against the time-averaged BFs and kinetic parameter statistics (signed -$\log_{10}$-transformed p-values) for the KO library only. Three omics datasets from a genome-wide deletion library were tested, as well as the SGA-PCC dataset and a STRING protein similarity network (STRING-PCC). The individual datasets were

matched for overlapping deletion mutants with autophagy phenotypes as the output and subsequently tested with random forest and 10-fold cross-validation using all features as model inputs.

The proteomics dataset was retrieved from Messner et al. 2023[48], and we used the complete imputed dataset. Each protein abundance value was normalized by the median value of the WT controls and subsequently $\log_2$-transformed. The dataset overlapped with 4189 deletion mutants and included 1850 proteins as model features.

The transcriptomic dataset was retrieved from Kemmeren et al. 2014[47], and represented microarray-based $\log_2$-fold changes from the WT controls. The dataset overlapped with 1419 deletion mutants and included 6112 transcripts as model features. The expression profiles for each mutant were standardized to have zero mean and unit variance.

The metabolomics dataset was retrieved from Mülleder et al. 2016[49], and represented concentration changes in deletion mutants for 19 amino acids. The dataset overlapped with 4501 deletion mutants.

The SGA-PCC was retrieved from Costanzo et al. 2016[27]. Because this dataset includes some correlation with multiple deletions for the same gene, we averaged the correlations per ORF over the columns and rows, subsequently. The resulting dataset overlapped with 4565 deletion mutants and included 5707 gene correlations as model features.

The STRING-PCC dataset was retrieved from the STRING database version 10^{32}. A binary interactome matrix for combined_sore ≥ 200 was generated and subsequently used to compute pairwise PCC between each column. The resulting correlation matrix overlapped with 4684 deletion mutants and had 6385 gene correlations as model features.

The response variables were standardized before training. Random forest was executed using the regression implementation of the *XGBoost* package (version 1.7.5.1) in R (version 4.2.1)[80] using squared error as loss and learning_rate = 1. Number of parallel trees were set to 1000, subsample = 0.63 and colsample_bynode = $\sqrt{p}/p$, where p indicated the number of features and varied per dataset. The number of splits were set to max_depth = 20, and we used a min_child_weight = 3. The r^2 was computed for each test fold and the median value was reported.

8.2 Random forest mediation analysis

Discovery of transcriptional mediators of regulatory effects of transcription factor (TF) on autophagy responses where executed by matching the time series of the Induction Dynamics gene Expression Atlas (IDEA) by Hackett et al. 2020[50] with the initial starvation responses dynamics of 186 overlapping TFs, using a 1 hour time shift. The rationale for using the IDEA dataset, is that the gene expression profiles presented in IDEA are directly causally driven by the acute activation of the TFs. Moreover, because of the abundance of autophagy perturbations caused by TF deletions, and the relatively low variation in autophagy at the onset of the starvation protocol, it is likely that a lot of the dynamic variation is related to loss-of-functions of crucial gene expression changes acutely involved in the starvation response. Gene target induction profiles that strongly predict the autophagy perturbation responses generated across multiple TF deletions, where the targets themselves are genes with a measured influence on autophagy, represent probable candidates as targets for transcriptional modulation of autophagy.

In the IDEA datasets we used 'log2_cleaned_ratio' variable for differential gene expression time series (X), which are normalized to the T0 baseline of each induction protocol. To compensate for varied efficacies on gene expression regulation from induction for different

TFs, as well as batch effects, we performed the harmonization procedure $X_{TF,adj} = X_{TF} \cdot (\sigma_X / \sigma_{X_{TF}})$, where σ_X is the standard deviation of X for the entire screen and $\sigma_{X_{TF}}$ is the standard deviation of X_{TF} for a specific TF. The reported log fold change (log FC) values in the study are based on these adjusted values. Missing values in the time series were imputed using the na_interpolation function with the stine method from the *imputeTS* package (version 3.3) in R[94], and time series values from 30, 45, 60 and 90 minutes after TF induction were extracted for the remaining analysis. These were matched with 90, 105, 120 and 150 minutes after starvation for $A\%_t$ and the BF_t time series, where the intermediate missing values were imputed using the na_interpolation function with the spline method. Thus, the resulting dataset had 744 samples (186 TFs x 4 time points). Before the regression the response variables were centered around the median of each time point, to compute the perturbations, and then standardized. The dataset had 6175 gene targets as features, 5989 after removal of the 186 TFs, and 4508 that were present in the autophagy dataset, where 1354 targets also had a significant autophagy phenotype (HMP ≤ 0.01).

Random forest was executed as described above using the *XGBoost* package (version 1.7.5.1) in R (version 4.2.1), with number of parallel trees set to 5000. For model evaluation 10-fold cross-validation was performed, where each fold was defined by withholding a random sample of individual data points or a random sample of TF time series. The r^2 was computed for each test fold and the median value was reported. To score the feature importance we used the sumGain, which was computed using the importance function in the *XGBoost* package. For clustering selected TFs with potential mediators, target genes with a sumGain > 2200 against one of the response variables and a HMP < 0.001, were considered (See Data File S14, *importance scores*). The sumGain cutoff was estimated as 2 times the average sumGain of features from control models trained with randomized sample IDs. SHAP (SHapley Additive exPlanations) values were used to assess the predictive effect from gene expression changes on autophagy perturbations. To compute SHAP values we used the SHAPforxgboost package (version 0.1.3)[95]. The individual SHAP values were grouped on whether the corresponding rfvalue (raw feature value; corresponding to the log fold changes in expression, $X_{TF}.adj$) were positive or negative, and then computing the mean per group for each feature (see Data File S14, *SHAP scores*) . Thus, this yielded average SHAP values for upregulated and downregulated gene expression, respectively. Differential SHAP values were computed by subtracting the SHAP value for upregulation with the SHAP value for downregulation. To compare the SHAP values for models trained with different number of features they were multiplied with p/p_{max}, where p indicate the number of features for the specific model and p_{max} indicate the maximum number of features.

References

56. Baryshnikova, A. *et al.* Quantitative analysis of fitness and genetic interactions in yeast on a genome scale. *Nat. Methods* **7**, 1017–1024 (2010).
57. Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition). *Autophagy* **17**, 1–382 (2021).
58. Janke, C. *et al.* A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947–962 (2004).
59. Shaner, N. C. *et al.* A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. *Nat. Methods* **10**, 407–409 (2013).
60. Kuzmin, E. *et al.* Systematic analysis of complex genetic interactions. *Science* **360**, (2018).
61. Tong, A. H. *et al.* Systematic genetic analysis with ordered arrays of yeast deletion mutants.

Science **294**, 2364–2368 (2001).

62. Chollet, F. & j. Allaire, J. *Deep Learning with R*. (Pearson Professional, 2018).

63. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. *arXiv [stat.ML]* (2018).

64. Sainburg, T., McInnes, L. & Gentner, T. Q. Parametric UMAP: learning embeddings with deep neural networks for representation and semi-supervised learning. arXiv. *Preprint] arXiv2009*.

65. Loader, C. & Liaw, M. A. Package 'locfit'. https://mirror.linux.duke.edu/cran/web/packages/locfit/locfit.pdf.

66. Langfelder, P., Zhang, B. & Horvath, S. dynamicTreeCut: Methods for detection of clusters in hierarchical clustering dendrograms. *R package version*.

67. Darsow, T., Rieder, S. E. & Emr, S. D. A multispecificity syntaxin homologue, Vam3p, essential for autophagic and biosynthetic protein transport to the vacuole. *J. Cell Biol.* **138**, 517–529 (1997).

68. Sato, T. K., Darsow, T. & Emr, S. D. Vam7p, a SNAP-25-like molecule, and Vam3p, a syntaxin homolog, function together in yeast vacuolar protein trafficking. *Mol. Cell. Biol.* **18**, 5308–5319 (1998).

69. Wang, C.-W., Stromhaug, P. E., Shima, J. & Klionsky, D. J. The Ccz1-Mon1 protein complex is required for the late step of multiple vacuole delivery pathways. *J. Biol. Chem.* **277**, 47917–47927 (2002).

70. Yang, S. & Rosenwald, A. A High Copy Suppressor Screen for Autophagy Defects in Δ and Δ Strains. *G3* **7**, 333–341 (2017).

71. Ohashi, Y. & Munro, S. Membrane delivery to the yeast autophagosome from the Golgi-endosomal system. *Mol. Biol. Cell* **21**, 3998–4008 (2010).

72. Arlt, H. *et al.* The dynamin Vps1 mediates Atg9 transport to the sites of autophagosome formation. *J. Biol. Chem.* **299**, 104712 (2023).

73. Shimobayashi, M., Takematsu, H., Eiho, K., Yamane, Y. & Kozutsumi, Y. Identification of Ypk1 as a novel selective substrate for nitrogen starvation-triggered proteolysis requiring autophagy system and endosomal sorting complex required for transport (ESCRT) machinery components. *J. Biol. Chem.* **285**, 36984–36994 (2010).

74. Kato, M. & Wickner, W. Vam10p defines a Sec18p-independent step of priming that allows yeast vacuole tethering. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 6398–6403 (2003).

75. Fisk, D. G. *et al.* Saccharomyces cerevisiae S288C genome annotation: a working hypothesis. *Yeast* **23**, 857–865 (2006).

76. Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

77. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

78. Bader, G. D. & Hogue, C. W. V. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* **4**, 2 (2003).

79. Cherry, J. M. *et al.* Saccharomyces Genome Database: the genomics resource of budding yeast. *Nucleic Acids Res.* **40**, D700–5 (2012).

80. Chen, T. *et al.* Xgboost: extreme gradient boosting. *R package version 0. 4-2* **1**, 1–4 (2015).

81. Duong, T. ks: Kernel Density Estimation and Kernel Discriminant Analysis for Multivariate Data in R. *J. Stat. Softw.* **21**, 1–16 (2007).

82. Väremo, L., Nielsen, J. & Nookaew, I. Enriching the gene set analysis of genome-wide data by incorporating directionality of gene expression and combining statistical hypotheses and methods. *Nucleic Acids Res.* **41**, 4378–4391 (2013).

83. Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**, 115–132 (1998).

84. Guthrie, C. & Fink, G. R. *Guide to Yeast Genetics and Molecular and Cell Biology: Guide to Yeast Genetics and Molecular Biology*. (Gulf Professional Publishing, 2004).

85. Gietz, R. D. & Woods, R. A. Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* **350**, 87–96 (2002).

86. Araki, Y., Kira, S. & Noda, T. Quantitative Assay of Macroautophagy Using Pho8△60 Assay and GFP-Cleavage Assay in Yeast. *Methods Enzymol.* **588**, 307–321 (2017).

87. Noda, T. & Klionsky, D. J. The quantitative Pho8Delta60 assay of nonspecific autophagy. *Methods Enzymol.* **451**, 33–42 (2008).

88. Suzuki, K. *et al.* The pre-autophagosomal structure organized by concerted functions of APG genes is essential for autophagosome formation. *EMBO J.* **20**, 5971–5981 (2001).

89. Cheong, H. & Klionsky, D. J. Biochemical methods to monitor autophagy-related processes in yeast. *Methods Enzymol.* **451**, 1–26 (2008).

90. Chica, N., Portantier, M., Nyquist-Andersen, M., Espada-Burriel, S. & Lopez-Aviles, S. Uncoupling of Mitosis and Cytokinesis Upon a Prolonged Arrest in Metaphase Is Influenced by Protein Phosphatases and Mitotic Transcription in Fission Yeast. *Front Cell Dev Biol* **10**, 876810 (2022).

91. Foiani, M., Marini, F., Gamba, D., Lucchini, G. & Plevani, P. The B subunit of the DNA polymerase alpha-primase complex in Saccharomyces cerevisiae executes an essential function at the initial stage of DNA replication. *Mol. Cell. Biol.* **14**, 923–933 (1994).

92. Mitchell, J. K., Fonzi, W. A., Wilkerson, J. & Opheim, D. J. A particulate form of alkaline phosphatase in the yeast, Saccharomyces cerevisiae. *Biochim. Biophys. Acta* **657**, 482–494 (1981).

93. Zhang, Y., Jenkins, D. F., Manimaran, S. & Johnson, W. E. Alternative empirical Bayes models for adjusting for batch effects in genomic studies. *BMC Bioinformatics* **19**, 262 (2018).

94. Moritz, S. & Bartz-Beielstein, T. ImputeTS: Time series missing value imputation in R. *R J.* **9**, 207 (2017).

95. Liu, Y., Just, A. & Mayer, M. SHAPforxgboost: SHAP plots for 'XGBoost'. *R package version 0.1. 0*.

Fig. S1.

Fig. S1. Robust quantification of autophagy using deep learning. a, Overview of deep learning approach. *Top left*: Causal control conditions for training data. Non-autophagic states (0) were defined as WT cells at T0 of starvation and after 4 hours of replenishment and as *atg1Δ* cells for any time points. Autophagic states (1) were defined as WT cells after 7 hours of starvation until T0 replenishment. *Top right*: Neural nets were trained on

extracted single-cell signal distribution features to classify autophagic cells. *Bottom left*: A hyperparameter grid search was initially performed over different neural net models with varying activation functions, hidden layers, number of initial neurons, and decay in the number of neurons per layer. The initial collection of models was trained for 150 epochs with early stoppage on validation loss and subsequently evaluated for test predictions, validation accuracy loss, and overfitting. Top models with an *elu*, *relu*, or *tanh* architecture were selected based on the validation loss and retrained for 3000 epochs with early stoppage on validation loss. *Bottom right*: A UMAP embedding of the last latent layer containing predictive autophagic features was used to evaluate the continuity of single-cell distributions between non-autophagic and autophagic states. For fast embedding, we used a parametric UMAP (pUMAP) approach where neural nets were trained to map the high-dimensional latent space features to a two-dimensional UMAP embedding. **b,** The average autophagy % in test data for neural nets, trained with various hyperparameters controlling the size and shape of the architecture, ranked by validation loss, consistently shows high separation between reference autophagy states for models with *elu*, *relu*, and *tanh* activation functions. **c,** Similar training and validation accuracies indicate low overfitting across most models. **d,** Sufficient model complexity is required to predict autophagy confidently. Only *relu*-based models show a positive association between increased model complexity and overfitting. Dashed lines indicate a complexity cutoff based on the model with the lowest complexity in top 1% (validation loss). Top 5 high and low complexity models (based on validation loss) were re-trained and examined further. **e,** Top model architectures trained for a greater number of epochs show similar performances (accuracy and loss) and robustness (correlation of predicted autophagy dynamics of WT and *vam6Δ* between experiments). Dashed lines indicate separation between low complexity (*left*) and high complexity (*right*) models. For *tanh* and *elu* architectures, low complexity models perform slightly better on average. **f,** Different model architectures result in very similar predicted autophagy dynamics. **g,** Top model architectures show high robustness in dynamics of embedded latent autophagic features using UMAP (correlation of population mean for WT and *vam6Δ* between experiments), with higher experimental correlation for models based on *elu* and *tanh*. **h,** UMAP dynamics of latent autophagic features are similar and consistent with reference phenotypes across different model architectures, with some activation function-dependent distributional differences in mean and variance. Points and bars indicate the mean and standard deviation at given time points across all experiments performed. The two models with the least variation in non-autophagic features are indicated by the boxes. **i,** Latent space UMAP embedding of control cells (model 30 and model 22). Points and bars indicate the population mean and standard deviation of an experiment at a given time point. **j,** Independent models yield highly consistent autophagy predictions across the genome-wide yeast library *(left panel)*, with a similar correlation to the reference controls color-coded as in (**h**) *(right panel)*. **k,** Association between input features and predicted autophagy log-odds.

Fig. S2.

Fig. S2. Quality control of genome-wide data. a, Predicted autophagy response dynamics for plate median and controls (model 30). The ribbons indicate the upper and lower boundary for plate and control filtering given by the 1.5xIQR +/- the 25%/75% quartiles, respectively. Plate medians with an average deviation from the screen median greater than 2xIQR were repeated. **b,** Prediction uncertainty, given by the average standard deviation per well for the genome-wide library and reference controls. **c,** Average number of analyzed cells per well for the genome-wide library and reference controls. **d,** Prediction uncertainty given by the average standard error per well for the genome-wide library and reference controls. **e,** Filtering of high uncertainty predictions due to low cell count, indicated by red dots. These mutants were repeated with higher seeding density.

Fig. S3.

Fig. S3. Double sigmoidal curve-fit results. a, Goodness-of-fit distributions for the genome-wide library and the reference controls. The RMSE distributions *(left panels)* show that all mutants have similar and low absolute curve-fit errors, while the log-likelihood score on scaled responses *(right panels)* indicates that *atg1Δ* and *vam6Δ* have lower compatibility with a sigmoidal model, consistent with the observed response curves. **b,** Sigmoidal model parameter distributions highlight differences between the KO collection strains, DAmP collection strains, and the WT control strains. **c,** Workflow for statistical analysis of autophagy perturbations. **d,** Distribution of autophagy perturbations from the plate medians for the KO collection strains, DAmP collection strains, and the control strains compared with the error distributions for the control strains. **e,** Distributions of sigmoidal model parameter differentials with the plate median for the KO collection strains, DAmP collection strains, and the control strains compared with the error distributions for the control strains.

Fig. S4.

Fig. S4. Statistics of parameter perturbations, and autophagy profile clustering. a, Frequency of mutants from the KO and DAmP collections having perturbations given a 1% p-value cutoff (*left*) or 5% FDR cutoff (*right*). **b,** Correlation between unweighted HMP (uHMP) and the minimum FDR over the set of hypothesis tests performed. **c,** Distribution of mutants with significant parameter perturbations (FDR ≤ 0.05), with inset indicating the cumulative probability. **d,** Hierarchical clustering of perturbation parameters using Pearson's correlations of signed significance scores for the significant mutants (HMP ≤ 0.01), with ward.D2 clustering and dynamic tree cutting. **e,** Hierarchical clustering of autophagy response dynamics for the significant mutants (HMP ≤ 0.01), with Euclidean distances, ward.D2 clustering and dynamic tree cutting.

Fig. S5.

Fig. S5. Tests for detecting autophagy reference genes. a, Model-fitted autophagy response curves for reference mutants. Repeated starvation measurements from a repetition screen are indicated in red for some genes. **b,** Area under the curve for PRC curves over ranked p-values for kinetic perturbation parameters comparing autophagy reference test sets versus equisized randomly sampled negative sets or ORF negative controls (squares). **c,** PRC curves for perturbation parameters and HMP ranked by p-values comparing a combined autophagy reference test set (n=35) versus equisized randomly sampled negative sets. The dashed lines represent the p-values and HMP for the recalled genes. **d,** PRC curves for perturbation parameters ranked by p-values comparing a combined autophagy reference test set (ATG: n=16; Fusion: n=19) versus an ORF negative control set (n=23). **e,** ROC curves for perturbation parameters ranked by p-values comparing a combined autophagy reference test set (ATG: n=16; Fusion: n=19) versus an ORF negative control set (n=23). **f,** ROC curve for perturbation parameters ranked by their signed p-values comparing ATG genes reference test set (n=16) and fusion genes reference test set (n=19) versus equisized randomly sampled negative sets. **g,** Area under the curve for ROC curves over ranked p-values (*top panel*) and over ranked signed p-values (*bottom panel*) for kinetic perturbation parameters comparing autophagy reference test sets versus equisized randomly sampled negative sets or ORF negative controls (squares).

Fig. S6.

Fig. S6. Reproducibility of top hits, and potential errors. a, The first stage of reproducibility testing was done by selecting the top-ranked mutants based on the HMP along with predicted false positives and negatives and repeating the starvation protocol (see methods 1.3). Subsequently, the mutant-wise correlation in autophagy dynamics and perturbation dynamics was compared between the two screens. **b,** Prediction of false positives and negatives based on enrichment tests of significant phenotypes (generated by different HMP cutoffs) in the set of first-degree neighbors for different network sources. The enrichment test p-values were aggregated by taking their harmonic mean to an overall network enrichment p-value and compared with the autophagy perturbation statistics of the respective genes. **c,** Non-significant genes annotated with the indicated autophagy GO terms were predicted as false negatives. **d,** Network enrichment p-value correlates with autophagy perturbation FDR_{min} (minimum perturbation BH-FDR value). Potential false positives or negatives were predicted in the off-diagonal areas as indicated. **e,**

Network enrichment p-value is concordant with both autophagy perturbation FDR_{min} and HMP. **f,** Mutant replicate correlation of autophagy perturbations for top hits and predicted false positives and negatives versus average absolute autophagy perturbation for GW data *(top panel)* and validation data *(bottom panel)*. The vertical lines represent the 2x median perturbation of the screen. **g,** Distribution of mutant replicate correlation of autophagy *(left)* and autophagy perturbation *(right)* for top hits and predicted false positives and negatives.

Fig. S7.

Fig. S7. Reproducibility across experimental replicates, strain background, and experimental protocols. a, The second stage of reproducibility testing was performed by selecting a sub-library of strongly significant (HMP ≤ 0.001) verified gene deletions with diverse perturbation phenotypes (dynamic variability and covering multiple gene classes for biological variability). For each mutant three new clones were made in a new background strain, and the entire experimental protocol was repeated three times under two distinct growth preparation protocols (see methods 7.1 and 7.3). Reproducibility was assessed in a mutant-wise manner by comparing autophagy dynamics and perturbation dynamics, and in a screen-wise manner by comparing computed kinetic parameters. **b,** Workflow of experimental organization for comparing reproducibility. P1 refers to a growth preparation protocol where the cells were grown to log phase before being transferred to fresh media 8 hours before the experimental procedure. P2 refers to a growth preparation protocol where the cells were grown to stationary phase before being transferred to fresh media 8 hours before the experimental procedure. The genome-wide screen R1 and replicate screen R2 were subjected to P1 and replicate screens R3 and R4 were subjected to P2. Colored arrows indicate the different levels of replication tested. **c,** Comparison of autophagy perturbation and mutant-wise replication errors with the same color-coding as in (**b**). The horizontal dashed line represents the 2x median perturbation of the screen. **d,** Mutant-wise replicate correlation of autophagy (*left panels*) and autophagy perturbation (*right panels*) for collection clones and new clones across biological replicates. **e,** Distribution of autophagy (*top panels*) and autophagy perturbation (*bottom panels*) correlation between collection clones and new clones for each experimental replicate. The distributions are stratified based on the ranked clone correlation from best to worst for each gene. The dashed horizontal lines represent the best, median, and worst correlation between WT control strains of the collection and new mutants. The median of all WT replicates was used as a plate control for computing autophagy perturbations, and a negative correlation for R2 indicates a distinct WT strain effect. **f,** Autophagy response dynamics for collection (Y7092) and new (BY4741) WT control strains. Dashed lines indicate the common median across all strains and experimental replicates for reference. **g,** Autophagy perturbation correlation between collection clones and new clones versus average absolute autophagy perturbation for new clones and collection clones across protocols. The red and black dots represent the best and median clone correlation for each gene, respectively. The dashed horizontal lines represent the best and median correlation between WT control strains of the collection and new mutants.

Fig. S8.

Fig. S8. Between-screen reproducibility. a, b Correlation scatter plots of autophagy perturbations for collection clones across experimental replicates. **c,** Correlation scatter plots of autophagy perturbations for new clones across experimental replicates. The inset values in (**a-c**) represent the Pearson correlations and associated p-values. **d,** Pearson correlations (*left*) and Spearman correlations (*right*) summarizing the between-screen similarities in autophagy perturbations depending on phase or mutant source. **e,** Correlation scatter plots of kinetic parameters from both collection clones and new clones across the experimental replicates. The inset values represent the Spearman correlations and associated p-values. **f,** Pearson correlations (*left two panels*) and Spearman correlations (*right two panels*) summarizing the between-screen similarities in autophagy perturbations and kinetic parameters.

Fig. S9.

Fig. S9. GSEA of kinetic perturbations. a, Hierarchical clustering of signed GSEA p-values for GO-BP terms and different kinetic perturbation parameters.

Fig. S10.

Fig. S10. Distribution of autophagy regulating genes in genome-wide networks. a, SAFE analysis of the genome-wide STRING network (version 10), with a combined score cutoff of 990 for edges. Reference map of bioprocesses (left) and of ATG and fusion set (middle) for interpreting the spatial enrichment of autophagy perturbation profiles (right). **b,** Shortest pathlength and shortest causal pathlength per autophagy perturbation profile. Causal paths were determined by requiring HMP ≤ 0.01 for all intermediate nodes. **c,** Shortest path length per autophagy perturbation profiles relative to the non-significant genes (ns). **d,** Cumulative count of significant genes (HMP ≤ 0.01) per shortest path or shortest causal path distances. **e,** Cumulative enrichment ($\log_2$) of significant genes (HMP ≤ 0.01) per shortest path or shortest causal path distances. **f,** Enrichment of perturbation profiles per shortest causal path distances in the SGA-PCC network (Costanzo et al. 2016). **g-i,** Distribution of autophagy perturbations across gene modules spanning three regulatory network hierarchies. Node colors indicate the overall perturbation statistics (signed -$\log_{10}$ p-values) and the size of the nodes are proportional to their HMP (-$\log_{10}$). **j,** Hierarchy of autophagy regulatory network layers presented in (**g-i**).

Fig. S11.

Fig. S11. Network associations of autophagy regulatory genes. a, Overview of the two strategies for gene pair comparisons in **(b)**. **b,** Correlations in temporal autophagy (dynamics) perturbations or sets of parameter perturbations (signed -log$_{10}$ p-values) between genes that share network interactions or complexome associations. Correlations were considered for all possible gene pairs (HMP $\leq$ 1) or only between subsets of genes with different HMP cutoffs (HMP $\leq$ 0.01 or 0.001). **c,** Modified gene set enrichment analysis (GSEA) to account for directional enrichment along SGA-PCC scores. **d,** Directional normalized enrichment score (*left panels*) or direction preferential normalized enrichment score (*right panels*) per perturbation profile for core ATG genes (*upper panels*) and fusion genes (*lower panels*). *p-value $\leq$ 0.05 (one-sample t-test). **e,** Directional enrichment of perturbation profiles along the SGA-PCC spectra for individual genes of the core autophagy machinery. *p-value $\leq$ 0.05 (permutation test).

Fig. S12. Autophagy dynamics and noise in the latent-space. a, UMAP-embedded latent-space distributions for control cells from model 22 and model 30. The densities represent distributions from 6 independent plates, grouped per time point and colored with the average autophagy prediction. Arrows indicate the position and direction of the average latent-space flux per plate. Left panel is presented with representative micrographs in *4b*. **b,** Latent space UMAP flux for control cells from model 22 and model 30. Arrows indicate the position and direction of the average latent-space flux per plate, and color indicates the time for each population average. Note the slight greater variability of non-autophagic features (*atg1Δ*) in model 30. **c,** Training and test scores for UMAP predictions, showing that predictive information from the DNN models is fully conserved. The DNN outputs, autophagy probability (P_a) or log-odds (LO), or the sample time per cell were predicted from bagged regression trees and scored using the Pearson correlation coefficient (PCC). Classification of WT versus *vam6Δ* cells were performed with bagged classification trees and scored using an F1 score. **d,** Comparing multivariate properties of the UMAP embeddings and latent feature vectors from the DNN models of the control cells. Each point represents standard deviation (SD) and flux per cell population for a given experiment and time point. **e,** Comparing phase-averaged UMAP deviation and flux, as well as prediction uncertainty (binary entropy) across models and datasets. **f,** Comparing UMAP variance and flux between the two DNN models. Note that UMAP flux is more reproducible between the two models. **g,** Distribution of WT and *vam6Δ* for different DNN prediction intervals. **h,** Training and test F1 scores for UMAP classification of WT versus *vam6Δ* cells per DNN prediction intervals. The classifications were performed with bagged classification trees.

Fig. S13.

Fig. S13. Bayesian modeling of autophagy execution from "out-of-distribution" latent-space features. a, Causal control conditions for learning intermediate states of autophagy execution. WT cells can execute autophagy based on nitrogen status. *atg1Δ* cells cannot produce autophagosomes. *vam6Δ* cannot clear autophagosomes and has de-regulated inhibition of autophagosome formation due to diminished TORC1 activation. The consequence is empirical latent-space distributions of *vam6Δ* cells that are populating an area between the non-autophagic and maturely autophagic feature space of *atg1Δ* and WT cells. The latent-space distributions for the three mutants can be used as reference states in a Bayesian modeling framework using Bayes factors (BFs). **b,** BF scores for classifying mutant phenotypes where time is a marginalized variable by using timewise reference probability densities. **c,** BF_t's for quantifying latent-space movements between fixed reference states by using time-invariant probability densities. **d,** Time-independent log BFs from **b** for the reference mutants from 6 independent experiments. **e,** Time-wise log BF_t's from **c** for the reference mutants from 6 independent experiments, quantifying the dynamics for autophagy execution. **f,** Comparison of log BF and BF_t between model 22 and model 30. *R* indicates the Pearson correlations, and *p* the respective p-values. **g,** ROC-AUC scoring the separation of ATG and fusion set mutants versus equisized randomly sampled negative sets, versus ORF negative controls, or the test sets versus each other. The BF and BF_t scores are compared with the predicted average autophagy perturbation and average binarized autophagy perturbation from the same neural classifiers. **h,** log BF and BF_t z-scores for control cells and ATG and fusion set cells. Points and bars indicate the means and standard deviations from collection of mutants. **i,** log BF z-scores per autophagy perturbation profiles.

Fig. S14.

Fig. S14. Genome-wide Bayes factor relations and GO-enrichment. a Genome-wide associations between BF_t or average autophagy perturbation and average signal co-occurrence for (GxR) early (first two time points; *upper panels*) or (GxR) late (time points after 7 hours in -N; *lower panels*). Pearson correlations with their respective p-values are reported. Green and red indicate outliers with low mNeonGreen-Atg8 and Pep4-mCherry for early time points, where red outliers were filtered out from the genome-wide analyses of BFs. **b,** Average BF_t z-scores for significantly enriched GO terms. **c,** Genome-wide associations between treatment phase averages of BF and BF_t for the KO collection. Blue and orange points indicate the ATG and fusion mutants, respectively. Pearson correlations with their respective p-values are reported. The BF classification concordances between the treatment phases (-N: starvation and +N: replenishment) are reported above each plot. Note that time-independent BFs have higher concordances compared to the phase-averaged time-wise BF_t values. **d,** Comparing the average BF z-scores for GO terms per treatment phase, starvation and replenisment, that had a significant differential enrichment between the phases (p-value $\leq$ 0.05). **e,** BFs (*upper panels*) or BF z-scores (middle and *lower panels*) for different components of the TORC1 signaling pathway, the RTG pathway, and the spermidine synthesis pathway. The BFs have been computed per phase (-N and +N; *upper* and *middle panels*) and as a differential between phases (starvation-replenishment) (*lower panels*).

Fig. S15.

Fig. S15. Experimental validation of autophagic flux perturbations. a, Immunoblot analysis of GFP-Atg8 cleavage and RpS6 phosphorylation for *rtg1Δ, rtg2Δ, rtg3Δ,* and WT cells in +N and -N for the indicated time points. +N t=0h refers to cells before shifting to -N. β-actin is used as a loading control. *p ≤0.05**p ≤0.01;***p ≤0.001; mixed model regression (n=4). **b,** Pho8Δ60 activity (%) as a measure of autophagic flux in WT, *atg1Δ, rtg1Δ, rtg2Δ,* and *rtg3Δ* cells in +N (before starvation) and followed by -N for 4 hours. *p ≤0.05**p ≤0.01***p ≤0.001; ANOVA with TukeyHSD. **c,** UMAP-embedded latent-space distributions and autophagy response dynamics for gene deletions of the spermidine synthesis pathway with representative micrographs from indicated time points, **d,** Representative images of immunoblot analysis of GFP-Atg8 cleavage and Rps6 phosphorylation for WT, *spe1Δ, spe2Δ, spe4Δ, tco89Δ, gtr1Δ,* and *lst4Δ* in -N followed by +N for the indicated time points. □-actin is used as a loading control. See methods for replicas and quantifications. **e,** UMAP-embedded latent-space distributions and autophagy response dynamics for gene deletions of selected gene deletions related to TORC1 activation.

Fig. S16.

Fig. S16. Comparing Bayes factors with autophagic flux. a, GFP-Atg8 cleavage (%) correlations with average Bayes factors, autophagy predictions, and fluorescence signals. **b,** GFP-Atg8 cleavage (%) correlations with autophagy perturbation parameters. Red and blue indicate positive and negative correlations, respectively. **c,** Time-wise GFP-Atg8 cleavage (%) correlations with BF_t-based regression predictions, using either a univariate model (WT:ATG1) or a bivariate model (VAM6.ATG1 + WT:VAM6). Pearson correlations with their respective p-values are reported. **d,** Time-wise correlations of GFP-Atg8 cleavage % with regression model predictions based on BF_t from DNN model 22, model 30 or the average of the models combined. The colored area for regression model 2 represents the relative contribution of the two model covariates.

Fig. S17.

Fig. S17. Rapamycin response screen. a, Autophagy (%) response dynamic in response to starvation, rapamycin or both for WT, ORF control and selected components of the TORC1 signaling pathway, the RTG pathway, and the

spermidine synthesis pathway. **b,** Similarity in autophagy perturbations (%) between different treatment conditions. **c,** Similarity in BF responses between different treatment conditions. **d,** Reproducibility in autophagy % and BFs between models and between experimental conditions. **e,** Comparison of the BF response to starvation dependent on the rapamycin treatment status. **f,** Comparison of the BF response to rapamycin treatment dependent on the nitrogen status.

Fig. S18.

Fig. S18. Effect of rapamycin on autophagy in TORC1 mutants. a, Representative micrographs depicting control and TORC1 signaling pathway mutants in response to nitrogen presence (+N) and starvation (-N), both with and without rapamycin (R), at indicated time points.

Fig. S19.

Fig. S19. Cross-omics prediction of autophagy dynamics. a, Benchmarking prediction of different autophagy metrics using 10-fold training and testing. Model performance as median test correlation between measured and predicted values. Transcriptome: Kemmeren et al. 2014 (*47*); Proteome: Messner et al. 2023 (*48*); Metabolome: Mülleder et al. 2017 (*49*); SGA-PCC: Costanzo et al. 2016 (*27*). **b,** Benchmarking prediction of autophagy perturbation dynamics from TF expression induction profiles (*50*) using 10-fold training and testing, with random sampling of the entire TF time-series (*left panel*) or random sampling of individual measurements (*right panel*). Model performance as median test correlation between measured and predicted values. Color coding indicates different feature pre-selection criteria. **c,** Adjusted SHAP values for regulation of gene expression across models and

autophagy metrics comparing feature classes based on autophagy perturbation profiles. Points and bars indicate mean and standard error. **d,** Adjusted SHAP values for regulation of gene expression comparing feature classes across models and autophagy metrics. F: formation genes; C: clearance genes with negative BF_t z-scores beyond two standard deviations from the mean. Points and bars indicate mean and standard error. **e,** Gene-wise SHAP values for differential regulation of gene expression predicting BF responses. **f,** qRT-PCR for expression of *GCN4*, *ATG13*, *GTR1*, *GTR2*, *VAM6*, *VPS15*, *VPS34*, and *ATG9* in *rtg1Δ*, *rtg2Δ*, *rtg3Δ*, and WT cells subjected to 4 hours of -N followed by 1 hour of +N. Graphs represent mean and standard error from n=3. *$p \leq 0.05$;**$P \leq 0.01$;***$P \leq 0.001$; t-test. **g,** Correlations in differential expression between per time-point for the panel of genes targets analyzed in (**6e**) and (**S19f**). **h,** Median gene expression change in response to TF induction in Hackett et al. 2020 (*50*).

Table S1. List of reference mutants for test sets.

Gene	ORF	Complex/ module	Test set	Literature
ATG11	YGL180W	Atg1 complex	ATG set	PMID: 24366339
ATG13	YPR185W	Atg1 complex	ATG set	PMID: 24366339
ATG17	YLR423C	Atg1 complex	ATG set	PMID: 24366339
ATG31	YDR022C	Atg1 complex	ATG set	PMID: 24366339
ATG29	YPL166W	Atg1 complex	ATG set	PMID: 24366339
VPS30 (ATG6)	YPL120W	PI3K complex	ATG set	PMID: 24366339
ATG14	YBR128C	PI3K complex	ATG set	PMID: 24366339
ATG9	YDL149W	Atg9-Atg2 system	ATG set	PMID: 24366339
ATG18	YFR021W	Atg9-Atg2 system	ATG set	PMID: 24366339
ATG2	YNL242W	Atg9-Atg2 system	ATG set	PMID: 24366339
ATG16	YMR159C	Atg12 conjugation system	ATG set	PMID: 24366339
ATG12	YBR217W	Atg12 conjugation system	ATG set	PMID: 24366339
ATG5	YPL149W	Atg12 conjugation system	ATG set	PMID: 24366339
ATG10	YLL042C	Atg12 conjugation system	ATG set	PMID: 24366339
ATG7	YHR171W	Atg12/Atg8 conjugation system	ATG set	PMID: 24366339
ATG3	YNR007C	Atg8 conjugation system	ATG set	PMID: 24366339
ATG4	YNL223W	Atg8 conjugation system	ATG set	PMID: 24366339
VAM3	YOR106W	Vacuole tethering/SNARE	Fusion set	PMID: 9245783
VAM7	YGL212W	Vacuole tethering/SNARE	Fusion set	PMID: 9710615
YPT7	YML001W	Vacuole tethering/GTPase	Fusion set	PMID: 12364329
CCZ1	YBR131W	Vacuole tethering/GEF complex	Fusion set	PMID: 12364329
MON1	YGL124C	Vacuole tethering/GEF complex	Fusion set	PMID: 12364329
VAM6 (VPS39)	YDL077C	HOPS complex	Fusion set	PMID: 20604902; PMID: 25143401
VPS41	YDR080W	HOPS complex	Fusion set	PMID: 20604902; PMID: 25143401

PEP3 (VPS18)	YLR148W	HOPS complex	Fusion set	PMID: 20604902; PMID: 27974437
VPS45	YGL095C	Golgi SNARE complex	Fusion set	PMID: 25143401
TLG2	YOL018C	Golgi SNARE complex	Fusion set	PMID: 20861302; PMID: 21784249
GOS1	YHL031C	Golgi SNARE complex	Fusion set	PMID: 20861302; PMID: 21784249
VPS5	YOR069W	Retromer complex	Fusion set	PMID: 37060997
VPS17	YOR132W	Retromer complex	Fusion set	PMID: 37060997
VPS29	YHR012W	Retromer complex	Fusion set	PMID: 37060997
VPS35	YJL154C	Retromer complex	Fusion set	PMID: 37060997
PEP8 (VPS26)	YJL053W	Retromer complex	Fusion set	PMID: 37060997
VPS27	YNR006W	ESCRT-0 complex	Fusion set	PMID: 20855891
VPS1	YKR001C	Dynamin/GTPase	Fusion set	PMID: 37060997
VPS3	YDR495C	CORVET complex	Fusion set	PMID: 20604902; PMID: 25143401
VAM10	YOR068C	HOPS related	Fusion set	PMID: 12748377
	YCR102W-A		Negative control set	PMID: 17001629
	YDR535C		Negative control set	PMID: 17001629