***In vivo* properties of Arabidopsis FCA condensates involved in RNA 3' processing**

Geng-Jen Jang[1], Alex L. Payne-Dwyer[2], Robert Maple[1], Zhe Wu[1,3], Fuquan Liu[1,4], Sergio G. Lopez[1],
Yanning Wang[5], Xiaofeng Fang[5], Mark C. Leake[2,6] and Caroline Dean[1,*]

[1]Department of Cell and Developmental Biology, John Innes Centre, Norwich, UK

[2]School of Physics, Engineering and Technology, University of York, York, UK

[3]Institute of Plant and Food Science, Southern University of Science and Technology, Shenzhen,
China

[4]Institute of Global Food Security, Queen's University Belfast, Belfast, UK

[5]Center for Plant Biology, Tsinghua University, Beijing, China.

[6]Department of Biology, University of York, York, UK

*Corresponding author, Tel: +44 (0)1603 450000; E-mail: caroline.dean@jic.ac.uk

Abstract

Our understanding of the functional requirements underpinning biomolecular condensation *in vivo* is
still relatively poor. The Arabidopsis RNA binding protein FCA is found in liquid-like nuclear
condensates that function in transcription termination, promoting proximal polyadenylation at many
targets in the Arabidopsis genome. To further understand the properties of these condensates *in vivo* we
used single particle tracking experiments on FCA reporters stably expressed at endogenous levels in
plant nuclei. These revealed FCA forms a core oligomer of ~4 molecules that multimerizes into higher-
order particles. This assembles into macromolecular condensates through the function of the coiled-coil
protein FLL2, which is genetically required for FCA function. FLL2 predominately co-localizes with
FCA in larger sized condensates. A missense mutation in the FCA RRM domain, also genetically
required for FCA function, reduced both average FCA particle and condensate size but did not perturb
the core oligomer. Our work points to a modular structure for FCA condensates involving
multimerization of core oligomers that assemble into functional macromolecular condensates via
associated RNA and FLL2 interactions.

Keywords: Biomolecular condensate/*COOLAIR*/*FLC*/Non-coding RNA/RNA processing

31
Introduction

Cellular compartmentalization by biomolecular condensates is associated with diverse processes in
eukaryotic cells (Banani et al, 2017; Shin & Brangwynne, 2017). While *in vitro* studies have highlighted
their roles in increasing local concentration and residence time of interacting components, the functional
characteristics of heterogeneous condensates *in vivo* remain unclear. Plants are an excellent system in
which to study biomolecular condensates due to their response to external conditions (Field et al, 2023).

For example, Arabidopsis *FLOWERING LOCUS C* (*FLC*)*,* a gene encoding a key flowering repressor,

is regulated by co-transcriptional and antisense-mediated chromatin silencing mechanisms (Fang et al,

2020; Xu et al, 2021) that involve multiple nuclear biomolecular condensates with very different

properties (Berry et al, 2017; Fang et al, 2019; Zhu et al, 2021; Fiedler et al, 2022). One of those

involves the RNA-binding protein FCA, which forms liquid-like nuclear condensates (Fang et al, 2019).

Another involves FRIGIDA, a strong activator of *FLC* expression, which forms condensates at low

temperature, with turnover dynamics that are slow compared to FCA (Zhu et al, 2021). FCA directly

interacts with FY, a Pfs2p/WDR33 homologue and a central component of the polyadenylation

machinery in eukaryotes (Simpson et al, 2003). FCA contains two RRM (RNA Recognition Motif)

domains and a WW domain located between two predicted prion-like domains (Macknight et al, 1997).

FCA recruits RNA 3' processing factors to an R-loop, which is a DNA:RNA hybrid generated by the

antisense non-coding RNA, *COOLAIR*, to induce transcription-coupled chromatin repression (Xu et al,

2021; Mateo-Bonmati et al, 2024; Menon et al, 2024). This mechanism also involves an FCA-

interacting coiled-coil protein, FLL2, which is required for both formation of FCA nuclear condensates

and FCA function (Fang et al, 2019). *In vitro* assays revealed that FCA condensates were enhanced by

addition of low amounts of Arabidopsis RNA. However, the immunoprecipitation of *COOLAIR* nascent

transcripts by FCA was not dependent on FLL2 (Fang et al, 2019).

Knowledge of the properties and molecular structure of biomolecular condensates helps dissection of

the molecular mechanisms involved (Peran & Mittag, 2020). However, studies of biological

condensates have focused on *in vitro* analysis and transfection studies and if *in vivo* stable expression

fusions have been made, they are often not carefully matched to endogenous expression levels. This

will influence the *in vivo* biophysical properties (Bienz, 2020). Here, we therefore generated stable

transgenic plants with fusions expressed under their own regulatory regions and chose lines that

expressed the fusion at endogenous levels. Determining the biophysical properties of condensates *in*

vivo in plant systems is challenging so we exploited SlimVar microscopy, a single-molecule imaging

technique (Plank et al, 2009; Payne-Dwyer et al, 2024). This enables simultaneous tracking of

individual particles within a wide range of biological condensates providing quantitative measurements

of molecule number and molecular motion behaviour (Shen et al, 2023). This method has been used to

study biomolecular condensates in live cells including bacteria, yeast or mammalian cells (Kent et al,

2020; Izeddin et al, 2014; Ladouceur et al, 2020; Biswas et al, 2022) and we adapted it for living plant

tissues to specifically study nuclear proteins (Bayle et al, 2021).

Our findings indicate that FCA-mScarlet-I particles multimerize and then assemble in an FLL2-

dependent manner into the liquid-like condensates observed using confocal microscopy. FLL2 and FCA

predominantly co-localize in the larger condensates. The involvement of RNA in promoting FCA

condensation was shown through the identification of a missense mutation in the FCA second RRM

domain. This mutation reduced RNA-binding activity *in vitro*, significantly attenuated FCA particle

size and condensate formation, lowered RNA 3' processing efficiency and prevented *FLC* repression

but did not influence formation of the core FCA particle. Our work points to a modular structure for

RNA 3' processing FCA condensates, with a low-order oligomer core that multimerizes into higher

order particles via associated RNA and into larger functionally important condensates through FLL2

interaction.

Results

Single particle tracking reveals physical characteristics of FCA particles.

For the SlimVar microscopy we chose to generate an FCA-mScarlet-I translational fusion as a

fluorescent reporter. mScarlet-I is a bright red fluorescent protein known for its characteristics as a

rapidly maturing monomer (Bindels et al, 2016). A mScarlet-I coding sequence was inserted into an

FCA genomic clone and transgenic plant lines were generated, where the fusion was expressed at similar

level as the endogenous FCA gene (Fig. EV1A and B). The FCA-mScarlet-I fusion formed nuclear foci

(Fig. 1A and Fig. 3A) and rescued the *fca-9* mutant phenotype (Fig. EV1B), just like the FCA-eGFP

fusion (Fang et al, 2019). We exploited SlimVar microscopy with a narrow, oblique illumination angle

of $60 \pm 5°$ to enhance imaging contrast and dynamically track very small FCA particles in nuclei in

whole root tissue (Fig. 1A and Movie EV1). The single-molecule sensitivity of SlimVar *in vivo*, coupled

with stepwise photobleaching of fluorescent protein tags (Leake et al, 2006; Reyes-Lamothe et al, 2010;

Payne-Dwyer et al, 2024), allowed us to characterize single-molecule brightness of the FCA-mScarlet-

I. This enabled quantification of the number of FCA-mScarlet-I molecules (stoichiometry) within the

particles in intact tissue (Fig. EV2 and Movie EV2; Jin et al, 2021).

Single particle tracking was undertaken on homozygous transgenic plants carrying the FCA-mScarlet-

I transgene expressed at endogenous levels in three biological replicate experiments. Analysis of the

FCA-mScarlet-I stoichiometry distribution identified a distinct peak representing a low-order oligomer

of FCA-mScarlet-I (Fig. 1B) with other peaks in multiples of approximately 4, from approximately 16

to 56 molecules (Fig. 1C). This finding aligns with a previous study that described the prion-like domain

of FCA, which when fused to GFP in yeast formed small oligomers (Chakrabortee et al, 2016). To

further investigate the oligomerization properties of FCA molecules we analysed in vitro produced FCA

protein. Purified recombinant FCA proteins containing either the NH2-terminal half containing RRM

domains (MBP-RRM, Fang et al, 2019) or the C-terminal FCA half containing prion-like domains

(MBP-PrLD) -shown schematically in Fig. EV3 - were analysed using size exclusion chromatography

with multi-angle static light scattering (SEC-MALS). The FCA RRM fragment forms low-order

oligomers containing 4 or possibly 5 FCA proteins, consistent with the particle tracking assay *in vivo*.

The prion-like C-terminal domain formed very large aggregates with more than 100 molecules (Fig.

EV3), a characteristic frequently associated with *in vitro* expression of PrLDs at relatively high
concentrations.
113
114 SlimVar also enabled us to analyse the *in vivo* diffusion coefficient of individual particles with different
115 stoichiometry. Generally, particles with low stoichiometry displayed variable mobility, while those with
116 high stoichiometry showed low mobility (Fig. 1D). In particular, the diffusion coefficient of particles
117 with stoichiometry less than 10 or more than 10 was 0.54 ± 0.75 $\mu m^2/s$ (mean $\pm$ s.d.) and 0.27 ± 0.4
118 $\mu m^2/s$, respectively.
119
FCA and FLL2 show co-localization in a larger size class of condensates.
Our previous study using transient assays suggested that FCA condensates serve as the sites for the
interaction between FCA, FLL2 and 3'-end processing factors (Fang et al, 2019). However, the
condensates formed by FCA and FLL2 exhibited different patterns in low resolution images of
Arabidopsis roots (Fang et al, 2019). To investigate the potential co-localization of FCA and FLL2
within the same nuclear condensates *in vivo*, we crossed plants carrying an FCA-mTurquoise transgene
(Xu et al, 2021) with plants carrying the FLL2-eYFP transgene (Fang et al, 2019) and used F3 plants
expressing both reporters for imaging. 3D imaging and Airyscan confocal microscopy were used,
acknowledging that the Z-axis resolution of the confocal microscope and the limitations of the imaging
algorithm may underestimate the number and size of the condensates. We found that FCA-mTurquoise
and FLL2-eYFP foci generally reside in distinct condensates within the cell nucleus (Fig. 2A and
Appendix Fig. S1) but co-localize within a few larger condensates (Fig. 2B and D). We conducted a
comprehensive analysis to determine the size distributions and the frequency of FCA, FLL2 and
FCA/FLL2-co-localized condensates (Fig. 2C and Appendix Fig. S2). The *sof78* missense mutation in
FLL2 is predicted to influence a salt bridge connecting two coiled coils, and this attenuates FCA
condensation and R-loop resolution (Fig. EV4; Fang et al, 2019). We used single particle tracking to
analyse whether *sof78* affected the assembly of the low-order oligomers of FCA. The analysis reveals
that the periodicity and stoichiometry distribution of FCA particles in *sof78* shows no obvious difference
compared to the control (Fig. EV4B to D). Overall, the data suggest that association of FLL2 promotes
formation of the larger functional FCA condensates.
140
To address if these prominent FCA/FLL2 condensates directly associate with the *FLC* locus, we crossed
the FCA-mScarlet-I transgene into a line carrying a previously established *FLC-lacO*/eYFP-LacI
system enabling spatial localization of the *FLC* locus in the nucleus. This line also carries an active
FRIGIDA allele so *FLC* is in an active transcriptional state (Rosa et al, 2013). FCA condensates did not
frequently co-localize with the *FLC-lacO*/eYFP-LacI marked locus (Fig. EV5). This result suggests
that FCA condensates may only transiently associate with the *FLC* locus in this genotype and
environmental conditions. However, it will be interesting to repeat this analysis in a genotype where

FLC is transcriptionally repressed by FCA, but where the Polycomb silencing has not yet been

established (Menon et al, 2024).

FCA condensation is dependent on a functional RRM domain.

To further investigate the functional significance of the molecular composition in FCA condensates we

continued a genetic screen for factors essential for FCA function (Liu et al, 2007). We used transgenic

plants with overexpressed *35S::FCAγ* and conducted a forward genetic analysis to identify mutations

that suppressed FCA's ability to repress *FLC* expression. This identified a missense mutation on the

FCA overexpression transgene at amino acid 304 of the FCA protein, resulting in a leucine to

phenylalanine change (Fig. EV1C). This alteration is located within the second RRM domain of FCA,

a domain conserved across various species (Macknight et al, 1997; Fig. EV1D). That the mutant came

out of an *FLC* mis-expression screen and exhibited late flowering shows the RNA-binding activity is

important for FCA function (Fig. EV1E). The impact of this missense mutation on the reduction of

RNA-binding activity was established through *in vitro* RNA-binding assays (Fig. EV1F).

To examine whether this RRM mutation affects FCA condensation, we generated transgene constructs

expressing an FCA wild-type genomic DNA fragment, with (FCArrm) and without (FCAwt) the

mutation fused with mScarlet-I and introduced them into *fca-9* mutant (Fig. EV1A). After selecting for

single insertion transgenic plants and homozygotes, we analysed the FCAwt condensates within the

nuclei of young roots using an Airyscan confocal microscope. The FCArrm transgene produced smaller,

more diffuse condensates at lower numbers, compared to wild type (Fig. 3A and B).

We also conducted single particle tracking with the FCArrm lines and observed the major oligomer

peak was slightly higher than that for FCAwt (Fig. 3C; Appendix Fig. S3). FCArrm particles with more

than 16 molecules were scarce, however, the integrated nuclear intensity of fluorescent signal in

FCArrm and FCAwt showed no obvious change (Appendix Fig. S4). This result suggests that the

reduced RNA-binding activity of FCA proteins decreases their likelihood of forming the larger particles

in the range of 16 to 56 molecules. We also compared the diffusion coefficients of individual particles

with their stoichiometry (Fig. 3D). The mean stoichiometry of FCArrm was lower than that of FCAwt,

while the overall the diffusion coefficients were slightly higher (Fig 3E; Appendix Fig. S5), however,

the periodic unit was not significantly changed (Fig. 3F). Thus, the RNA-binding activity of FCA

proteins affects multimerization of the FCA particles, and formation of the larger condensates, but does

not affect core oligomer assembly.

The functional RNA-binding domain is required for *COOLAIR* R-loop resolution and FCA auto-

regulation.

We then used the FCAwt and FCArrm lines to further investigate the functional importance of the RNA-

binding domain. Compared to the FCAwt transgenic plants, FCArrm plants displayed higher levels of

FLC expression and therefore flowered late (Fig. 4A and B, EV1B). FCA associates with *COOLAIR*,

recruiting 3' processing factors to promote proximal *COOLAIR* polyadenylation, thus resolving a

COOLAIR-formed R-loop (Liu et al, 2010; Xu et al, 2021). The ratio of proximal-to-distal isoforms of

COOLAIR transcripts was reduced in the FCArrm lines compared to FCAwt (Fig. 4C). Consistent with

this result, S9.6-DNA/RNA immunoprecipitation followed by cDNA conversion (DRIPc)-qPCR

analysis showed higher R-loop level in FCArrm relative to FCAwt, indicating that FCArrm has reduced

ability to resolve the *COOLAIR*-induced R-loop (Fig. 4D).

FCA autoregulates itself promoting polyadenylation at the proximal sites in intron 3 (Quesada et al,

2003), so we conducted Quant-seq analysis (Moll et al, 2014) to capture polyadenylated *FCA* transcripts

in Col-0, FCAwt, FCArrm and *fca-9*. In FCArrm and *fca-9*, there were significantly fewer reads

corresponding to proximally polyadenylated *FCA* transcripts compared to FCAwt and Col-0.

Conversely, reads at the distally polyadenylated site of *FCA* increased (Fig. 5), all consistent with

reduced FCA function. RNA-binding activity is thus important for FCA particle multimerization and

formation of the condensates, and for the well-defined FCA functional roles; promotion of proximal

FCA and *COOLAIR* termination and R-loop resolution.

Discussion

Our study provides insight on the functional properties of the dynamic liquid-like RNA 3' processing

condensates containing the Arabidopsis nuclear RNA-binding protein FCA. We conducted single

particle tracking to characterize molecular stoichiometry and diffusivity and the influence of protein

and RNA partners that function in *FLC* repression. Our work points to a modular structure for FCA

condensates centred around a low-order core oligomer, which multimerize to oligomers and

macromolecular condensates via associated RNA and FLL2 interaction. The genetic data support a

requirement for these condensates in FCA function for RNA 3' processing (Fig. 3A, 4 and EV4A; Fang

et al, 2019) with RNA-protein and protein-protein interactions playing different roles in the

oligomerization and condensation processes.

The FCA protein has many domains including an extended prion-like domain (Chakrabortee et al, 2016),

two RRM domains (Macknight et al, 1997), interrupted by a WW domain (Simpson et al, 2003;

Henderson et al, 2005) and an extended NH2 terminus that initiates translation with a CTG (non-

methionine) start codon (Simpson et al, 2010). Further work will be required to fully determine how all

these domains influence FCA condensate dynamics. Notably, the FCA core modular structure is

reminiscent of ELAV, a well-characterized Drosophila RNA-binding protein. ELAV with three RRM

domains tetramerizes on its target RNAs (Soller & White, 2005). For FCA condensates, the role of RNA

may involve functionality of the hydrophobic core in the RRM domain. The missense mutation (L304F)

could strengthen the hydrophobic core in the FCA second RRM domain, an equivalent hydrophobic

region to that recently found to influence the protein stability of TDP-43, an amyotrophic lateral

sclerosis (ALS)-linked RNA binding protein (Mackness et al, 2024). RNA methylation may also be a

contributing factor to the condensation dynamics as m6A has been shown to function in the regulatory

mechanisms of *FLC* and *COOLAIR* (Xu et al, 2021; Wang et al, 2022; Sun et al, 2022; Amara et al,

2023).

COOLAIR transcription is enhanced upon cold exposure, and the transcripts form dense 'clouds' at each

FLC locus, as judged by single-molecule RNA fluorescence in situ hybridization (Rosa et al, 2016).

Whether the FCA condensates associate with these *COOLAIR* clouds will require RNA live imaging

analysis. It is interesting to speculate that *COOLAIR* molecules might undergo thermo-responsive phase

transitions as has been found recently for other RNAs (Wadsworth et al, 2023), with protein partners

tuning this phase behaviour (Ruff et al, 2020). The *FLC* biological system has a lot of potential to

uncover fundamental principles of condensate functionality *in vivo,* especially if we can successfully

image the condensates at the locus. In mammalian cells, Mediator condensates only sporadically co-

localize with *Sox2*, and this variability is attributed to a dynamic kissing model, where a Mediator

condensate interacts with the gene only at specific timepoints (Cho et al, 2018; Du et al, 2024). This

may be the case for *FLC,* which can exist in either a transcriptionally active (with functional FRIGIDA)

or inactive (without functional FRIGIDA) state; with switching between these states connected to co-

transcriptional RNA processing (Fang et al, 2020; Mateo-Bonmati et al, 2024; Menon et al, 2024).

Exploring whether the transient association of FCA condensates with *FLC* locus reflects the

transcriptional status of *FLC* will be the next important question.

In this study, we used SlimVar microscopy, which integrates Slimfield's technical capabilities with

enhanced imaging contrast from variable-angle laser excitation (Plank et al, 2009; Payne-Dwyer et al,

2024). In contrast to a previous approach using a combination of Airyscan and Slimfield to infer the

stoichiometry of FCA assemblies indirectly (Payne-Dwyer et al, 2022), SlimVar allowed for direct and

precise single particle tracking and stoichiometric quantification of molecular assemblies in plant root

tissue. The oblique excitation beam reduces out-of-focus background fluorescence, similar to HILO and

lightsheet microscopy, and the use of an oil immersion objective lens improves fluorescence signal

despite some optical aberration (Tokunaga et al, 2008; Stelzer et al, 2021). SlimVar calibration

techniques such as adjusting the objective lens correction collar helped minimize these aberrations,

enabling effective postprocessing "sifting" to collect robust signals (Payne-Dwyer et al, 2024). The

limitations of SlimVar microscopy when observing FCA particles included the typically small field of

view, which usually captures just one nucleus per acquisition. Additionally, the methodology suffered

from fewer or shortened tracks due to bleaching, diffusion of assemblies, and residual aberration and

scattering, necessitating large sample numbers. Furthermore, at higher assembly concentrations, careful
image analysis was required to avoid overlapping assemblies leading to incorrect stoichiometry
estimates. Despite these limitations, SlimVar provides the ability to functionally study particles
involved in biomolecular condensation in Arabidopsis. This opens up the possibility of combining a
genetically tractable system with dissection of condensate function and their dynamic responses to
various environmental exposures.

264
265 **Methods**
266
Plant materials and growth conditions
The wild type Col-0, *fca-9* mutant allele*, pFCA::FCA-mTurquoise* and *pFLL2::FLL2-eYFP* transgenic
plants were described previously (Fang et al, 2019; Xu et al, 2021). To generate the *pFCA::FCA-*
mScarlet-I transgenic plants, the FCA genomic DNA fragment was amplified and inserted into the
pENTR vector. The mScarlet-I coding sequence was inserted before the stop codon via In-Fusion
seamless cloning. The FCA genomic DNA fragment fused mScarlet-I was transferred to pSLJ755I5
vector (Jones et al, 1992) via gateway cloning. The RRM mutation construct is modified from wild-
type FCA genomic DNA constructs via In-Fusion seamless cloning. Primers were listed in Table EV1.
These constructs were transformed into the *fca-9* mutant. Arabidopsis seeds were surface sterilized and
sown on standard Murashige and Skoog (MS) medium plate without glucose and stratified at 4 °C for
3 days before transferred to long-day conditions (16-h light at 20 °C, 8-h darkness at 16 °C) for 4-7
days.

279
Confocal microscopy of FCA-mScarlet-I
Confocal images were acquired on a Zeiss LSM 880 upright microscope equipped with an Airyscan
detector. Samples were imaged through a C-Apochromat 63x/NA 1.2 water-immersion objective by
exciting them at 561 nm and using the Airyscan detector in super resolution mode. The fluorescence
emission arrived at the detector after passing through a filter that allowed wavelengths between 495-
550 nm and above 570 nm to pass through. Typically, the voxel size was $0.06 \times 0.06 \times 0.26$ μm (xyz)
and the pixel dwell time was 1.78 μs. Z-stacks typically comprised 28 slices and spanned a 7.13 μm
range for more than 70% of the nucleus. In Fig. 3A, the FCA-mScarlet-I signal was enhanced by shifting
its dynamic range to 0–160, and the images were processed using maximum intensity projection. After
Airyscan processing, the image analysis was performed in Arivis Vision4D ver. 4.1.0. (Zeiss). The blob
finder algorithm was applied to the mScarlet-I channel using a diameter value of 0.35 μm, a probability
threshold of 45%, and a split sensitivity of 50%. Blob finder is a segmentation algorithm that excels at
finding round or spherical objects in 2D or 3D, respectively. It uses a Gaussian scale to identify seeds
for the objects and subsequently applies a watershed algorithm to determine the boundaries of the
objects. To exclude non-specific identified condensates, a minimum size of 0.03 μm³ and a sphericity
(Mesh) of 0.6 were set during the analysis, and non-specific foci outside the nuclear region of interest

were removed. Violin plots reflecting the data distribution were produced with GraphPad Prism 10

software.

Co-localization analysis of FCA and FLL2

Confocal images were acquired either on the Zeiss LSM 880 microscope mentioned above or a Zeiss

LSM 980 confocal microscope. When using the LSM880, line by line sequential acquisition was

employed. Samples were imaged through a C-Apochromat 63x/NA 1.2 water-immersion objective and

were excited at 458 nm (mTurquoise) and 514 nm (eYFP). The fluorescence emission of the samples

passed through a filter that only allowed wavelengths in the 465-505 nm range and above 525 nm to

reach the Airyscan detector, which was operating in super resolution mode. The voxel size was 0.05 ×

0.05 × 0.21 µm (xyz) and the pixel dwell time was 0.72 µs. Z-stacks comprised 25 slices spanning a

4.96 µm range. 0.2 µm TetraSpeck microspheres (Thermo Fisher Scientific) were imaged using these

same settings and the resulting images were used to correct the sample images for chromatic aberration.

The correction was performed in Zen (Zeiss) using the channel alignment method. In Fig. 2A, FCA-

mTurquoise and FLL2-eYFP signal was enhanced by shifting the dynamic range of the signal to 0–

1650 and 0-750.

After Airyscan processing, channel alignment, and nuclei segmentation, the image analysis was

performed in Arivis Vision4D ver. 4.1.0. (Zeiss). Firstly, the blob finder algorithm was applied to the

mTurquoise channel using a diameter value of 0.3 µm, a probability threshold of 37%, and a split

sensitivity of 50%. Then, the blob finder algorithm was applied to the eYFP channel using a diameter

value of 0.35 µm, a probability threshold of 37%, and a split sensitivity of 50%. Afterwards, the

intersection between the output of the two blob finder operations was calculated. Finally, metrics such

as volume, mean intensity, and total intensity were computed for the objects generated by each of the

blob finder operations, as well as for their intersection. In general, a minimum size of 0.03 μm^3 and a

sphericity (Mesh) of 0.6 were set during the analysis, and non-specific foci outside the nuclear region

of interest were removed. Violin plots and x-y plots reflecting the data distribution were produced with

GraphPad Prism 10 software.

SlimVar microscopy

For all imaging modalities, seedlings' root tips were oriented horizontally on slides containing MS

media with 1% agarose, sealed with MS media under a No. 1.5 coverslip. The customized SlimVar

microscope was previously described (Payne-Dwyer et al, 2024). Continuous wave lasers (Coherent

OBIS) delivered a de-expanded Gaussian beam mode (TEM_{00}) to the back aperture of an NA 1.49 Apo

TIRF 100× oil immersion objective lens (Nikon) with a lateral displacement such that the beam

illuminated a sample area no greater than 9 × 16 µm at 60 ± 5º incidence. Single molecule sensitivity

was achieved using a fast sCMOS camera (Photometrics Prime95B) which triggered a continuous wave

laser for 10 ms exposure per frame at 80 Hz, providing optimal contrast for detection (Appendix Fig.

S6). The total magnification was approximately 200×, resulting in an oversampled image pixel edge

size of 53 nm. Excitation and emission filters were tailored to best contrast for mScarlet-I (594/25 nm).

Fluorescence acquisition settings in the red and complementary green channels were optimized to

prevent initial saturation and to ensure the detection of individual molecular brightness tracks for

mScarlet-I. To calibrate SlimVar, the beam displacement lens and objective lens correction collar were

adjusted in situ to minimise optical aberrations at a representative 20 μm depth. These settings were

then locked to prevent systematic variation in the characteristic single molecule brightness. Individual

nuclei in the outermost 3 cell layers were identified in brightfield mode to determine the best focus

without bleaching, and subsequently imaged in the SlimVar fluorescence mode until completely

photobleached. Videos were subject to a temporal bleach correction (Fiji, Schindelin et al, 2012).

Gaussian smoothing of 0.5 pixels (Fiji) were used on images, min/max adjusted for contrast, and the

black point set at the background level outside the nucleus.

Single particle tracking and intensity analysis

Fluorescent particles were identified from SlimVar image sequences and linked into tracks using

ADEMScode software (Plank et al, 2009) using a strict 'sifting' approach. This sets signal-to-noise and

distance thresholds to discard foci with low contrast, whose integrated intensity is influenced by

background or detection noise, or exhibit nonphysical mobility. Each of these particles were localised

to a super-resolved, subpixel precision, typically 40 nm. The random overlap probability was estimated

using a nearest neighbour distance approach as <5% for all observed tracks (Payne-Dwyer et al, 2024).

Analysis was spatially restricted to data lying within a region of effectively uniform laser illumination

(80% ± 9% s.d. peak intensity) by cropping images to an area of 240 × 184 pixels (12.7 × 9.8 μm).

Fluorescence image sequences were further cropped to specify individual nuclei, using manually

segmented masks from brightfield images.

The 2D diffusion coefficient of each track (D) was computed using the increase of mean-square

displacement across time interval according to a random walk model (Leake et al, 2008). Focus intensity

was calculated by summing the pixel values within a 5-pixel distance and subtracting an averaged

background level from the rest of a 17-pixel square (Leake et al, 2008). The track stoichiometry was

found by extrapolating the particle's intensity in the first 5 image frames s to account for photobleaching

(Syeda et al, 2019), then dividing this initial intensity by the characteristic single molecule brightness

(Wollman et al, 2017). The characteristic single molecule brightness linked to each reporter was

determined as 110 ± 15 (mean ± s.d.) for mScarlet-I based on the modal size of steps in track intensity

during late-stage photobleaching. The track intensities over time were first denoised using an edge

preserving filter (Chung and Kennedy, 1991). This characteristic single molecule brightness provides

an internal calibration for track stoichiometry, periodicity, and integrated nuclear intensity which is

refined to nuclear protein number.

To calculate periodicity, the stoichiometries of all tracks within each nucleus were represented as a

kernel density distribution (Leake et al, 2014), employing an empirical standard deviation of 0.6

molecules on the characteristic single molecule brightness. Peaks within this distribution were identified

using the MATLAB *findpeaks* function, and the intervals between nearest neighbour peaks were

computed (Payne-Dwyer et al, 2024). Raw estimations of the total number of molecules in each nucleus

were determined using ImageJ macros to integrate the pixel values inside segmented nuclei as in

previous work (Wollman & Leake, 2015; Payne-Dwyer et al, 2024). The nuclear protein number of

each reporter dataset was refined to exclude autofluorescence by calculating the difference between

mean integrated nuclear intensities of the labelled dataset and an unlabeled control, adjusted

proportionally to the ratio of mean nuclear segment areas.

In vitro protein expression and purification

MBP-RRM and MBP-PrLD proteins were expressed and purified from *Escherichia coli* (Rosetta) cells

using Ni-NTA resin as described previously (Fang et al, 2019). Briefly, protein expression was induced

by 0.4 mM isopropyl-β-D-1-thiogalactopyranoside (IPTG) at 18°C overnight. Cells were collected by

centrifugation and re-suspended in lysis buffer (40 mM Tris-HCl pH7.4, 500 mM NaCl, 10% glycerol).

The suspension was sonicated for 20 min (3 s ON, 3 s OFF, SCIENTZ) and centrifuged at 13000 *g* for

30 min at 4°C. The supernatant was incubated with Ni-NTA agarose for 20 min, washed and eluted with

buffer E (40 mM Tris-HCl pH7.4, 500 mM NaCl, 500 mM Imidazole, Sangon). Proteins were further

purified by gel filtration chromatography (Superdex-200; GE Healthcare) and stored in buffer S (40

mM Tris-HCl pH7.4, 500 mM NaCl, 1 mM DTT) at −80 °C.

Size-exclusion chromatography coupled with multi-angle laser light scattering (SEC-MALS)

Multi-angle light scattering experiments were performed in 40 mM Tris-HCl pH7.4 and 500 mM NaCl

using a Superdex-200 10/300 GL size exclusion column (GE Healthcare). The concentration of MBP-

RRM and MBP-PrLD used was 2 mg/mL. The chromatography system was connected to a Wyatt

DAWN HELEOS laser photometer and a Wyatt Optilab T-rEX differential refractometer. Wyatt ASTRA

7.3.2 software was used for data analysis.

Expression analysis

Total RNA was extracted from plant materials as described previously (Qüesta et al, 2016), treated with

TURBO DNase (Thermo Fisher Scientific) to eliminate DNA contamination, and reverse-transcribed

using SuperScript IV Reverse Transcriptase (Thermo Fisher Scientific) with gene-specific reverse

primers. Quantitative PCR (qPCR) analysis was conducted on a LightCycler480 II (Roche), and the

qPCR data were normalized to the reference genes *UBC*. To measure the proximal ratio of *COOLAIR*,

the level of the proximal component was normalized to the distal *COOLAIR*. Primers were listed in

Table EV1.

DNA−RNA hybrid immunoprecipitation followed by cDNA conversion (DRIPc)

The DRIP protocol is as described previously (Xu et al, 2021) with modifications. In general, 3 g 10-day-old seedlings were harvested and grounded into a fine powder. The powder was suspended in 35 mL of Honda buffer (20mM HEPES, 0.44M sucrose, 1.25% Ficoll, 2.5% dextran T40, 10mM MgCl$_2$, 0.5% Triton x-100, 5 mM DTT, 1x protease inhibitor cocktail (Roche), filtered through one layer of Miracloth, and centrifuged at 3500×g for 15 min. Nuclear pellets were resuspended in 1mL Honda buffer and centrifuged at 8000×g for 1 min. Pellets were then resuspended in the lysis buffer (50mM Tris-HCl pH 8.0, 10mM EDTA, 1% SDS) supplied with 0.1 mg/mL proteinase K (AM2546, Thermo Fisher Scientific) and digested at 55 °C overnight with gentle rotation. The mixture was phenol/chloroform extracted, followed by nucleic acid precipitation with NaOAc and ethanol. The nucleic acid pellet was dissolved in water and quantified with Qubit DNA quantification kit (Thermo Fisher Scientific). In general, 1 μg of nucleic acid was digested by restriction enzymes including DdeI, NdeI and XbaI (NEB) overnight. The digested nucleic acid was phenol/chloroform extracted, followed by nucleic acid precipitation with NaOAc and ethanol. Then the nucleic acid was treated with RNAse III (Thermo Fisher Scientific) and then extracted by phenol/chloroform and ethanol precipitation. The treated nucleic acid was then diluted ten times with dilution buffer (16.7mM Tris pH 7.5, 167mM NaCl, 2.2mM EDTA, 0.1% Triton X-100) and 10% was stored at −20 °C as input. In all, 5 μg of S9.6 antibody (1:100 dilution, ENH001, Kerafast) was added, then incubated overnight at 4 °C. The next day, 50 μl Dynabeads Protein G (Thermo Fisher Scientific) was added and incubated for another 2 h. The immunoprecipitants were washed five times with dilution buffer and twice with TE buffer, then were eluted in 100 μl elution buffer (10mM Tris pH 7.5, 2 mM EDTA, 0.2% SDS, 100 ng/μl tRNA) and 5 μl proteinase K at 55 °C for 2 hours, together with input samples. The nucleic acids were precipitated with NaOAc, isopropanol, and glycogen, dissolved in water. The DRIP nucleic acid samples were then treated with TURBO DNase (Thermo Fisher Scientific) to eliminate DNA contamination, and reverse-transcribed using SuperScript IV Reverse Transcriptase (Thermo Fisher Scientific) with gene-specific reverse primers. All the samples including inputs were subjected to qPCR analysis via LightCycler480 II (Roche). The data were normalized to 1% of input. Primers were listed in Table EV1.

Quant-seq

The Quant-seq protocol is as described previously (Mateo-Bonmatí et al, 2024). The total RNAs prepared for Quant-seq underwent purification using the Qiagen RNeasy miniprep kit (QIAGEN, 74106). Lexogen GmbH (Austria) conducted the library preparation, sequencing, and data analysis. Sequencing-ready libraries were created from 100 ng of input RNA using a QuantSeq 3' mRNA-Seq Library Prep Kit REV for Illumina (015UG009V0271), following standard procedures. The RNA integrity and quality of indexed libraries were assessed on a Fragment Analyzer device (Agilent Technologies) using a DNF-471 RNA Kit and HS-DNA assay, respectively. Library quantification was

carried out using a Qubit dsDNA HS assay (Thermo Fisher Scientific). A sequencing-ready pool of
indexed libraries was then sequenced on an Illumina NextSeq 2000 with a 100-cycle cartridge, using
the Custom Sequencing Primer (CSP). Verification of read quality utilized FastQC version v0.11.7, and
read adapter trimming employed cutadapt version 1.18 (Martin, 2011). Clean reads were mapped to the
latest version of the Arabidopsis genome (TAIR10) using the splice-aware aligner STAR version 2.6.1a
(Dobin et al, 2013).
450

Electrophoretic Mobility Shift Assay
FCA cDNA with L304F mutation was amplified from the mutant as described in Fig. EV1.
To generate RNA probes, dsDNA of the corresponding region was amplified through PCR (primers see
Table EV1) with the T7/T3 phage polymerase promoter added at the 5' end of the primer. *COOLAIR*
RNA was then generated by following a protocol provided by the MAXIscript® T7 Transcription Kit
(Thermo Fisher Scientific). The resulting product was DNA digested, denatured, and separated on 5%
TBE gel containing 8 M Urea. The band at the right size was cut out, sliced, and eluted in gel extraction
buffer (0.5 M NH_4Ac, 1 mM EDTA, 0.2% SDS, 1xRNAsecure (Thermo Fisher Scientific) at 37°C
overnight, followed by precipitation with isopropanol. 10 pmol purified RNA was 3'biotinylated by
following the instructions provided by the RNA 3' End Biotinylation Kit (Thermo Fisher Scientific).
The labelling efficiency was determined to be at least 90%. For the gel shift assay, a 20 µl reaction (25
ng/µl protein, 2 nM probe, 250 ng/µl tRNA, 50 ng/µl heparin, 10 mM HEPES (pH 7.3), 120 mM KCL,
1 mM $MgCl_2$, 1 mM DTT) was set up on ice. For the competition assay, 2x to 100x of unlabelled probe
was included in the reaction. The reaction with only GST was used as a negative control. The mixture
was incubated at room temperature for 5 min and resolved on 3.5% native TBE-acrylamide gel and
transferred onto a positively charged nylon membrane. The biotinylated RNA on the membrane was
detected by chemiluminescence according to a protocol provided with the Chemiluminescent Nucleic
Acid Detection Module (Thermo Fisher Scientific, 89880). For the binding reaction, 1 µg of different
recombinant proteins was incubated with 2 pmol of biotinylated probe in 20 µL of 1× buffer containing
20 mM Tris (pH 7.5), 100 mM NaCl, 10 mM $MgCl_2$, non-specific competitors (1µg/µl heparin and 5µg
tRNA) and 0.05% Triton X-100 at room temperature for 10 min. The RNA-protein mixture was
resolved on 6% 1× TBE acrylamide gel under 100 V for 50 min, followed by electrophoretic transfer
to positively charged nylon membranes (GE Healthcare). The biotinylated RNA on the membrane was
UV cross-linked and detected using chemiluminescent nucleic acid detection following the
manufacturer's instructions (Thermo Fisher Scientific).
476

Data availability
Raw image data of confocal microscopy and co-localization analysis have been deposited in the
BioImage Archive (https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD1154). The raw and
processed imaging data and analysed tracks of single particle tracking analysis are available at the

BioStudies: https://doi.org/10.6019/S-BIAD1234. The ADEMScode MATLAB tracking analysis

software can be found at https://github.com/alex-payne-dwyer/single-molecule-tools-alpd.

The raw reads of Quant-seq have been deposited in the Short Read Archive (SRA) under the reference

(PRJNA1087342 and PRJNA1076161).

Acknowledgements

We thank Shuqin Chen, Tina Zhang, JIC Horticultural Services and JIC Bioimaging facility for their

excellent technical assistance and Hsuan Pai (TSL) for her artwork assistance. We thank Simon

H. Stitzinger and Jose A. M. Lantero (MPI-IE) for helpful discussion. We also thank all members of

the Caroline Dean and Martin Howard groups. This work was funded by the European Research

Council Advanced Grant (EPISWITCH, 833254), Wellcome Trust (210654/Z/18/Z), and the Royal

Society Professorship (RSRP\R\231006) to Caroline Dean, and EPSRC grants (EP/T00214X/1 to

C.D. and EP/T002166/1 and EP/W024063/1 to M.L).

494
Disclosure and competing interest statement

The authors declare that they have no conflict of interest.

References

Amara U, Hu J, Cai J & Kang H (2023) FLK is an mRNA m6A reader that regulates floral transition
by modulating the stability and splicing of *FLC* in Arabidopsis. *Mol Plant* 16: 919–929

Banani SF, Lee HO, Hyman AA & Rosen MK (2017) Biomolecular condensates: Organizers of
cellular biochemistry. *Nat. Rev. Mol. Cell Biol.* 18: 285-298

Bienz M (2020) Head-to-tail polymerization in the assembly of biomolecular condensates. *Cell* 182:
799–811

Bayle V, Fiche JB, Burny C, Platre MP, Nollmann M, Martinière A & Jaillais Y (2021) Single-
particle tracking photoactivated localization microscopy of membrane proteins in living plant
tissues. *Nat Protoc* 16: 1600–1628

Berry S, Rosa S, Howard M, Bühler M & Dean C (2017) Disruption of an RNA-binding hinge region
abolishes LHP1-mediated epigenetic repression. *Genes Dev* 31: 2115–2120

Bindels DS, Haarbosch L, Van Weeren L, Postma M, Wiese KE, Mastop M, Aumonier S, Gotthard
G, Royant A, Hink MA *et al* (2016) mScarlet: A bright monomeric red fluorescent protein for
cellular imaging. *Nat Methods* 14: 53–56

Biswas S, Chen Z, Karslake JD, Farhat A, Ames A, Raiymbek G, Freddolino PL, Biteen JS &
Ragunathan K (2022) HP1 oligomerization compensates for low-affinity H3K9me recognition and
provides a tunable mechanism for heterochromatin-specific localization. *Sci Adv* 8

Chakrabortee S, Kayatekin C, Newby GA, Mendillo ML, Lancaster A & Lindquist S (2016)
Luminidependens (LD) is an Arabidopsis protein with prion behavior. *Proc Natl Acad Sci U S A*
113: 6065–6070

Cho WK, Spille JH, Hecht M, Lee C, Li C, Grube V & Cisse II (2018) Mediator and RNA
polymerase II clusters associate in transcription-dependent condensates. *Science* 361: 412–415

Chung SH & Kennedy RA (1991) Forward-backward non-linear filtering technique for extracting
small biological signals from noise. *J Neurosci Methods* 40: 71–86

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M & Gingeras TR
(2013) STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15–21

Du M, Stitzinger SH, Spille J, Hijaz M, Quintana A, Cisse II, Du M, Stitzinger SH, Spille J, Cho W *et*
al (2024) Direct observation of a condensate effect on super- enhancer controlled gene bursting.

    Cell 187: 331–344

Fang X, Wang L, Ishikawa R, Li Y, Fiedler M, Liu F, Calder G, Rowan B, Weigel D, Li P *et al*
    (2019) Arabidopsis FLL2 promotes liquid–liquid phase separation of polyadenylation complexes.
    Nature 569: 265–269

Fang X, Wu Z, Raitskin O, Webb K, Voigt P, Lu T, Howard M & Dean C (2020) The 3' processing of
    antisense RNAs physically links to chromatin-based transcriptional control. *Proc Natl Acad Sci U
    S A* 117: 15316–15321

Fiedler M, Franco-Echevarría E, Schulten A, Nielsen M, Rutherford TJ, Yeates A, Ahsan B, Dean C
    & Bienz M (2022) Head-to-tail polymerization by VEL proteins underpins cold-induced Polycomb
    silencing in flowering control. *Cell Rep* 41

Field S, Jang GJ, Dean C, Strader LC & Rhee SY (2023) Plants use molecular mechanisms mediated
    by biomolecular condensates to integrate environmental cues with development. *Plant Cell* 35:
    3173–3186

Henderson IR, Liu F, Drea S, Simpson GG & Dean C (2005) An allelic series reveals essential roles
    for FY in plant development in addition to flowering-time control. *Development* 132: 3597–3607

Izeddin I, Récamier V, Bosanac L, Cissé II, Boudarene L, Dugast-Darzacq C, Proux F, Bénichou O,
    Voituriez R, Bensaude O *et al* (2014) Single-molecule tracking in live cells reveals distinct target-
    search strategies of transcription factors in the nucleus. *Elife* 3: 1–27

Jin X, Lee JE, Schaefer C, Luo X, Wollman AJM, Payne-Dwyer AL, Tian T, Zhang X, Chen X, Li Y,
    et al (2021) Membraneless organelles formed by liquid-liquid phase separation increase bacterial
    fitness. *Sci Adv* 7

Jones JDG, Shlumukov L, Carland F, English J, Scofield SR, Bishop GJ & Harrison K (1992)
    Effective vectors for transformation, expression of heterologous genes, and assaying transposon
    excision in transgenic plants. *Transgenic Res.* 1: 285–297

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
    Žídek A, Potapenko A *et al* (2021) Highly accurate protein structure prediction with AlphaFold.
    Nature 596: 583–589

Kent S, Brown K, Yang C hsun, Alsaihati N, Tian C, Wang H & Ren X (2020) Phase-separated
    transcriptional condensates accelerate target-search process revealed by live-cell single-molecule
    imaging. *Cell Rep* 33: 108248

Ladouceur AM, Parmar BS, Biedzinski S, Wall J, Tope SG, Cohn D, Kim A, Soubry N, Reyes-
    Lamothe R & Weber SC (2020) Clusters of bacterial RNA polymerase are biomolecular
    condensates that assemble through liquid-liquid phase separation. *Proc Natl Acad Sci U S A* 117:
    18540–18549

Leake MC, Chandler JH, Wadhams GH, Bai F, Berry RM & Armitage JP (2006) Stoichiometry and
    turnover in single, functioning membrane protein complexes. *Nature* 443: 355–358

Leake MC, Greene NP, Godun RM, Granjon T, Buchanan G, Chen S, Berry RM, Palmer T & Berks
    BC (2008) Variable stoichiometry of the TatA component of the twin-arginine protein transport
    system observed by in vivo single-molecule imaging. *Proc Natl Acad Sci U S A* 105: 15376–15381

Leake MC (2014) Analytical tools for single-molecule fluorescence imaging in cellulo. *Phys Chem
    Chem Phys* 16: 12635–12647

Liu F, Quesada V, Crevillén P, Bäurle I, Swiezewski S & Dean C (2007) The Arabidopsis RNA-
    Binding Protein FCA Requires a Lysine-Specific Demethylase 1 Homolog to Downregulate *FLC*.
    Mol Cell 28: 398–407

Liu F, Marquardt S, Lister C, Swiezewski S & Dean C (2010) Targeted 3′ processing of antisense
    transcripts triggers Arabidopsis *FLC* chromatin silencing. *Science* 327: 94-7

Mackness BC, Morgan BR, Deveau LM, Kathuria S V., Zitzewitz JA & Massi F (2024) A
    hydrophobic core stabilizes the residual structure in the RRM2 intermediate state of the ALS-
    linked protein TDP-43. *bioRxiv* 2024.06.12.598648.

Macknight R, Bancroft I, Page T, Lister C, Schmidt R, Love K, Westphal L, Murphy G, Sherson S,
    Cobbett C *et al* (1997) FCA, a gene controlling flowering time in Arabidopsis, encodes a protein
    containing RNA-binding domains. *Cell* 89: 737–745

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
    EMBnet.journal 17: 10-12

Mateo-Bonmatí E, Montez M, Maple R, Fiedler M, Fang X, Saalbach G, Passmore LA & Dean C

(2024) A CPF-like phosphatase module links transcription termination to chromatin silencing. *Mol Cell* 84: 2272–2286

Menon G, Mateo-Bonmati E, Reeck S, Maple R, Wu Z, Ietswaart R, Dean C & Howard M (2024) Proximal termination generates a transcriptional state that determines the rate of establishment of Polycomb silencing. *Mol Cell* 84: 2255–2271

Moll P, Ante M, Seitz A & Reda T (2014) QuantSeq 3' mRNA sequencing for RNA quantification. *Nat Methods* 11: i–iii

Payne-Dwyer AL & Leake MC (2022) Single-molecular quantification of flowering control proteins within nuclear condensates in live whole Arabidopsis root. *Methods Mol Biol.* 2476:311–328.

Payne-Dwyer AL, Jang GJ, Dean C & Leake MC (2024) SlimVar : rapid in vivo single-molecule tracking of chromatin regulators in plants. bioRxiv 2024.05.17.594710.

Peran I & Mittag T (2020) Molecular structure in biomolecular condensates. *Curr Opin Struct Biol* 60: 17–26

Plank M, Wadhams GH & Leake MC (2009) Millisecond timescale slimfield imaging and automated quantification of single fluorescent protein molecules for use in probing complex biological processes. *Integr Biol* 1: 602–612

Quesada V, Macknight R, Dean C & Simpson GG (2003) Autoregulation of FCA pre-mRNA processing controls Arabidopsis flowering time. *EMBO J* 22: 3142–3152

Qüesta JI, Song J, Geraldo N, An H & Dean C (2016) Arabidopsis transcriptional repressor VAL1 triggers Polycomb silencing at *FLC* during vernalization. *Science* 353: 485–488

Reyes-Lamothe R, Sherratt DJ & Leake MC (2010) Stoichiometry and architecture of active DNA replication machinery in *Escherichia coli*. *Science* 328: 498–501

Rosa S, De Lucia F, Mylne JS, Zhu D, Ohmido N, Pendle A, Kato N, Shaw P & Dean C (2013) Physical clustering of *FLC* alleles during polycomb-mediated epigenetic silencing in vernalization. *Genes Dev* 27: 1845–1850

Rosa S, Duncan S & Dean C (2016) Mutually exclusive sense-antisense transcription at *FLC* facilitates environmentally induced gene repression. *Nat Commun* 7: 1–7

Ruff KM, Dar F & Pappu R V. (2021) Ligand effects on phase separation of multivalent macromolecules. *Proc Natl Acad Sci U S A* 118

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676–682

Shen Z, Jia B, Xu Y, Wessén J, Pal T, Chan HS, Du S & Zhang M (2023) Biological condensates form percolated networks with molecular motion properties distinctly different from dilute solutions. *Elife* 12: 1–32

Shin Y & Brangwynne CP (2017) Liquid phase condensation in cell physiology and disease. *Science* 357

Simpson GG, Dijkwel PP, Quesada V, Henderson I & Dean C (2003) FY is an RNA 3′ end-processing factor that interacts with FCA to control the Arabidopsis floral transition. *Cell* 113: 777–787

Simpson GG, Laurie RE, Dijkwel PP, Quesada V, Stockwell PA, Dean C & Macknight RC (2010) Noncanonical translation initiation of the Arabidopsis flowering time and alternative polyadenylation regulator FCA. *Plant Cell* 22: 3764–3777

Soller M & White K (2005) ELAV multimerizes on conserved AU4-6 motifs important for *ewg* splicing regulation. *Mol Cell Biol* 25: 7580–7591

Stelzer EHK, Strobl F, Chang B-J, Preusser F, Preibisch S, McDole K & Fiolka R (2021) Light sheet fluorescence microscopy. *Nat Rev Methods Prim* 1: 73

Syeda AH, Wollman AJM, Hargreaves AL, Howard JAL, Bruning JG, McGlynn P & Leake MC (2019) Single-molecule live cell imaging of Rep reveals the dynamic interplay between an accessory replicative helicase and the replisome. *Nucleic Acids Res* 47: 6287–6298

Sun B, Bhati KK, Song P, Edwards A, Petri L, Kruusvee V, Blaakmeer A, Dolde U, Rodrigues V, Straub D *et al* (2022) FIONA1-mediated methylation of the 3'UTR of *FLC* affects *FLC* transcript levels and flowering in Arabidopsis. *PLoS Genet* 18: 1–22

Tokunaga M, Imamoto N & Sakata-Sogawa K (2008) Highly inclined thin illumination enables clear single-molecule imaging in cells. *Nat Methods* 5: 159–161

Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G,
Laydon A *et al* (2022) AlphaFold Protein Structure Database: Massively expanding the structural
coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* 50: D439–
D444
Wang C, Yang J, Song P, Zhang W, Lu Q, Yu Q & Jia G (2022) FIONA1 is an RNA N 6-
methyladenosine methyltransferase affecting Arabidopsis photomorphogenesis and flowering.
Genome Biol 23: 1–29
Wadsworth GM, Zahurancik WJ, Zeng X, Pullara P, Lai LB, Sidharthan V, Pappu R V., Gopalan V &
Banerjee PR (2023) RNAs undergo phase transitions with lower critical solution temperatures. *Nat*
Chem 15: 1693–1704
Wollman AJM & Leake MC (2015) Millisecond single-molecule localization microscopy combined
with convolution analysis and automated image segmentation to determine protein concentrations
in complexly structured , functional cells , one cell at a time. *Faraday Discuss* 00: 1–24
Wollman AJM, Shashkova S, Hedlund EG, Friemann R, Hohmann S & Leake MC (2017)
Transcription factor clusters regulate genes in eukaryotic cells. *Elife* 6: 1–36
Xu C, Wu Z, Duan HC, Fang X, Jia G & Dean C (2021) R-loop resolution promotes co-
transcriptional chromatin silencing. *Nat Commun* 12, 1790
Zhu P, Lister C & Dean C (2021) Cold-induced Arabidopsis FRIGIDA nuclear condensates for *FLC*
repression. *Nature* 599: 657–661
656

657   **Table EV1: Primers**

Primer name	Primer sequence 5' -> 3'	Annotations
Primers for cloning		
Inf-XhoI-pFCA-F	CGGGCCCCCCCTCGACGAGCGGTGAAGATGTATTGT	Cloning of FCA genomic fragments with Infusion strategy
Inf-BamHI-FCA-3UTR-R	TAGAACTAGTGGATCACAACCTTGAGTTCCTCGAGT	
Inf-FCA-stop-linear-R	AGCTTTATTCTTCCACATGAGTTCTTGG	
Inf-FCA-stop-linear-F	TGAGGTTCATATCTACCCTTTCTCTCCT	
Inf-FCA-linker-mScarletI-F	TGGAAGAATAAAGCTTCTGCTGCCGCATCCGCGGCAGCTTCAGCCGTGAGCAAGGGCGAGGCA	Cloning of FCA-mScarlet-I with Infusion strategy
Inf-FCA-mScarletI-R	TAGATATGAACCTCACTTGTACAGCTCGTCCATGC	
Inf-FCA-stop-linear-R	AGCTTTATTCTTCCACATGAGTTCTTGG	Cloning of FCA fusion protein with Infusion strategy
Inf-FCA-stop-linear-F	TGAGGTTCATATCTACCCTTTCTCTCCT	
Inf-NdeI-FCA-In3-F	TAGGTGGTGCAAACATATGGA	Cloning of L304F mutation on FCA fragments with Infusion strategy
Inf-FCA-L304F-R	AAAAAACTTAAACTCTAGGGTGC	
Inf-FCA-L304F-F	CCCTAGAGTTTAAGTTTTTTGTTGG	
Inf-AfeI-FCA-In6-R	GTAGAATGTCCAAGCGCTT	

Primers for transcript analyses		
FCAγ-qPCR-F	TCAGTCGTGCAACTGCTCCTC	Spliced *FCAγ* qPCR
FCAγ-qPCR-R	GATCCAGCCCACTGTTGTTT	
FLC spliced F	AGCCAAGAAGACCGAACTCA	Spliced *FLC* qPCR
FLC spliced R	TTTGTCCAGCAGGTGACATC	
Set1 new LP	TGGTTGTTATTTGGTGGTGTG	RT for proximal *COOLAIR*
set2_new_2_LP	CCTGCTGGACAAATCTCCGA	Proximal *COOLAIR* qPCR
set2_new_2_RP	TCACACGAATAAGGTGGCTAATTAAG	
FLC -158_F	GCCCGACGAAGAAAAAGTAG	RT for distal *COOLAIR*
Set4-new-F-195	GTATCTCCGGCGACTTGAAC	Distal *COOLAIR* qPCR
Set4-new-R-195	GGATGCGTCACAGAGAACAG	
UBC_qPCR_F	CTGCGACTCAGGGAATCTTCTAA	*UBC* qPCR
UBC_qPCR_R	TTGTGCCATTGAATTGAACCC	
Primers for DRIPc-qPCR		
4319f	AGAACAACCGTGCTGCTTTT	A
4466r	TGTGTGCAAGCTCGTTAAGC	
5030F	CCGGTTGTTGGACATAACTAGG	B
5153R	CCAAACCCAGACTTAACCAGAC	
5327f	TTTTTGTTATGGTTAGGTTTGGA	C
5395r	AGTAGCACTACTTCTAGACACTTGGA	
5456f	GCTTCCAAACTTAAAAGCTTAAACA	D
5584r	TCTTTTTGTCTTCTATCCAAGGAAT	
5531F	TGGTTGTTATTTGGTGGTGTG	E

5652R	CTGCTCCCACATGATGATTA	
Primers for EMSA		
FCA-EMSA-F1-718-T7	AAGTAATACGACTCACTATAGGGATTTCAACCGCCG ATTTAAG	
FCA-EMSA-R-T3	AAGAATTAACCCTCACTAAAGGGTTCCGGTTGTTG GACATAACT	

Figure 1. Super-resolution microscopy and single particle tracking to determine the physical characteristics of FCA particles

(**A**) A schematic of SlimVar microscopy used for the imaging Arabidopsis root cells and an example of a SlimVar image of a nucleus showing FCA-mScarlet-I (magenta). Scale bar: 2 µm. (**B**) Distribution of the stoichiometry (molecule number) of individual FCA particles. (**C**) Periodicity of FCA particle stoichiometry estimated from the most common spacing between neighboring peaks in each stoichiometry distribution. (**D**) The diffusion coefficient and stoichiometry of each FCA particle. The single particle tracking analysis is from three biologically independent experiments. Results from individual replicate are provided in Appendix Figure S3.

Figure 2. The co-localization of FCA and FLL2 is associated with larger sized condensates in Arabidopsis nuclei

(**A**) Selected 3D images of FCA and FLL2 colocalization. FCA-mTurquoise signal (green) and FLL2-eYFP signal (magenta) were segmented using the blob finder algorithm of Arivis Vision4D for foci identification, as described in the Methods section. White arrows indicate the co-localized condensates. The enlarged images of co-localized foci are shown insets at the bottom. Scale bars: 2 μm. (**B**) Quantification of size of FCA (total FCA-mTurquoise identified foci), FLL2 (total FLL2-eYFP identified foci) and FCA identified foci that are co-localized with FLL2. The lines in the violin plots indicate the median and quartiles. (**C**) Size and frequency distribution of FCA, FLL2 foci and FCA foci that are co-localized with FLL2. (**D**) Number of FCA, FLL2 foci and FCA foci that are co-localized with FLL2 in each nucleus. The lines in the violin plots indicate the median and quartiles. The quantitative analysis was based on 18 nuclei from 5 independent samples. Statistical significance was determined using non-parametric Brunner-Munzel test. ****P-value < 0.001.

Figure 3. The RNA-binding activity of FCA influences FCA condensate size and number

(A) Selected Airyscan confocal images of wild type FCA fused to mScarlet-I (FCAwt) and FCA-mScarlet-I with the missense (L304F) mutation (FCArrm). Scale bars: 5 μm. (B) Quantification of size and number of FCAwt and FCArrm condensates using Vision4D software. The lines in the violin plots indicate the median and quartiles. The quantitative analysis is based on 43 FCAwt nuclei and 65 FCArrm nuclei from 3 independent experiments. (C) Distributions of stoichiometry of individual FCAwt (black) and FCArrm (red) particles. (D) Diffusion coefficient and stoichiometry of each FCAwt (black) and FCArrm (red) particle. (E) Statistic analysis of stoichiometry and diffusion coefficient of FCAwt and FCArrm particles. (F) Periodicity of FCAwt (black) and FCArrm (red) particle stoichiometry. The single particle tracking analysis is from 3 biologically independent experiments. The FCAwt data corresponds to that shown in Figure 1. Statistical significance was determined using non-parametric Brunner-Munzel test. **P-value < 0.01, ****P-value < 0.0001. Results from individual replicates are provided in Appendix Figure S3 and S5.

Figure 4. A missense mutation in the RRM of FCA only partially complements the *fca-9* mutant phenotype

(**A**) Flowering time of selected transgenic lines (#8 for FCAwt and #1 for FCArrm in Fig. EV1) grown in a long-day photoperiod. Data are presented as mean ± s.d. (n = 14, except for Col-0 which n = 13). (**B**) Relative values of *FLC* in the indicated lines. Values were normalized to the *UBC* gene and to Col-0. Data are presented as mean ± s.d. (n = 5, except for *fca-9* where n = 4). (**C**) The ratio of proximal: distal isoforms of *COOLAIR* transcripts relative to control (Col-0) in the indicated plants. Data are presented as mean ± s.d. (n = 4). (**D**) Schematic of *FLC* and *COOLAIR* transcripts at the *FLC* locus. Blue dots indicate the locations of primer sets for qPCR. Untranslated regions are indicated by grey boxes, and exons by black boxes. kb, kilobase. (**E**) DRIPc–qPCR analyzing *COOLAIR* R-loop in indicated genotypes. Data are presented as mean from four technical repeats. Three biological replicates show similar trends (Appendix Figure S7). TSS, transcription start site. The primer sets used for DRIPc-qPCR (A to E) corresponds to that shown in Fig. 4D. Statistical significance was calculated using the two-tailed t-test, with *P*-values indicated. ****P*-value < 0.001, *****P*-value < 0.0001.

Figure 5. The rrm mutation perturbs the autoregulatory function of FCA in promoting proximal polyadenylation

(**A**) Quant-seq experiments assaying polyA sites at the *FCA* locus in different genotypes. The vertical pale grey bars highlight previously described proximal and distal polyA sites. (**B**) Quantification of reads corresponding to proximal and distal polyadenylated *FCA* transcripts. Data are presented as mean ± s.d. (n = 3). Statistical significance was calculated using the two-tailed t-test, with *P*-values indicated. **P-value < 0.01, ***P-value < 0.001.

Figure EV1. A missense mutation (L304F) on the second RRM of FCA attenuates RNA-binding

(**A**) Schematic representation of the transgenic FCA-mScarlet-I fusion within the FCA locus - the position of the missense L304F mutation is indicated. Untranslated regions are indicated by grey boxes, exon regions by black boxes, and mScarlet-I coding sequence by the red box. (**B**) Expression of *FLC* and *FCA* relative to *UBC* in plants at the T3 generation following transformation. Data represent mean values from technical repeats. The FCAwt #8 and FCArrm #1 lines are used for further analysis. (**C**) FCA protein structure predicted by Alphafold Protein Structure Database (AF-Q5I5A2-F1-v4, Jumper et al, 2021; Varadi et al, 2022). The arrow indicates the location of the amino acid change from the missense mutation (amino acid 213 in this protein structure database corresponds to amino acid 314 when FCA uses the non-canonical start CTG codon). (**D**) Peptide alignment of Arabidopsis FCA proteins from different plant species (https://phytozome-next.jgi.doe.gov). The arrow indicates the conserved Leucine 304. (**E**) Plants carrying transgenes with the L304F mutation show delayed flowering compared to the control. Data are mean ± s.d. (n = 23 and 24 for the control and L304F, respectively). Statistical significance was calculated using the two-tailed t-test, with *P*-value indicated. ****P*-value < 0.0001. (**F**) *In vitro* RNA binding assay showing that the Leu 304 and each RRM domain are important.

Figure EV2. Automated tracking analysis of FCA particles and determination of the characteristic single molecule brightness in living tissues.

(**A**) An example of FCA-mScarlet-I particle tracking at 10 ms/frame, which intermediate frame overlaid with tracks (white arrows) corresponding to the paths of achieves super-resolved localization precision and provides estimates of the corresponding stoichiometry and mobility. The left panel shows the initial frame of the acquisition, while the middle panels display the intermediate stage of the acquisition (after 75 ms), where contrast is greatest and assemblies are clearly visible. The nuclear boundary (white closed line) is obtained by blurring the initial frame and setting the Otsu intensity threshold. The third panel features the same ach detected assembly throughout the acquisition. Scale bar: 2 μm. (**B**) The consistent photobleaching steps yield an estimate of about 110 photoelectron counts for the characteristic single-molecule brightness of the FCA-mScarlet-I *in planta*. The intensity of particles near the end of five particle tracks in the FCA-mScarlet-I image sequences is represented by square dots, and the denoised signals by solid lines of the corresponding color. Here, zero intensity corresponds to the mean background level after complete photobleaching. (**C**) A subset of n = 35 particles was identified as having four photobleaching steps. This subset gives a characteristic single molecule brightness (black line) of 110 ± 31 counts (mean ± s.d., n=35). The initial intensities of these particles (grey line) are each greater than the characteristic single molecule brightness by a factor of ~ four (stoichiometry: 4.0 ± 0.6 molecules, mean ± s.d., n=35), suggesting that each particle is composed of ~ 4 FCA-mScarlet-I molecules. Data for individual particles are shown above the corresponding distribution in a jitter and box plot (showing median, interquartile range and min-max range). (**D**) A characteristic single molecule brightness of 110 ± 15 counts (mean ± s.d., n=504) is calculated from the time-averaged intensity levels of each of n=504 particles photobleached to a single molecule above background, sifted using the signal-to-noise threshold.

Figure EV3. SEC-MALS of purified FCA truncated proteins

SEC-MALS of purified MBP-RRM (**A**) or MBP-PrLD (**B**). Line traces indicate molar mass as determined from MALS. The predicted molecular mass of MBP fused FCA truncated protein MBP-RRM and MBP-PrLD is approximately 72 kDa and 89 kDa, respectively.

Figure EV4. The *COOLAIR* R-loop level and FCA-eGFP single particle tracking in the *sof78* mutant

(**A**) Two biological replicates of DRIPc–qPCR *COOLAIR* R-loop in the L*er*, *sof78* and *fca-1* genotypes. The *sof78* mutant carries a missense mutation in a region of FLL2 that is predicted to form a salt bridge, which influences FCA condensation (Fang *et al*., 2019). Data are means from three technical repeats. TSS, transcription start site. The primer sets used for DRIPc-qPCR (C and D) are the same as those shown in Figure 4D. (**B**) An example of FCA-eGFP particle tracking at 10 ms/frame. The left panel shows the initial frame of the acquisition. The third panel features the same intermediate frame overlaid with tracks (white arrows) corresponding to the paths of each detected assembly throughout the acquisition. The nuclear boundary (white closed line) is obtained by blurring the initial frame and setting the Otsu intensity threshold. Scale bar: 2 μm. (**C**) Distributions of stoichiometry of individual FCA-eGFP (black) and FCA-eGFP/*sof78* (blue) particles. (**D**) Periodicity of FCA-eGFP (black) and FCA-eGFP/*sof78* (blue) particle stoichiometry. The stoichiometry in the figure is capped at 100. The single particle tracking analysis is from three biologically independent experiments.

Figure EV5. Co-localization analysis of FCA condensates and *FLC-lacO*/eYFP-LacI

(A) An illustration of the *FLC-lacO*/eYFP-LacI transgenic plants. (B) Single slice of Arabidopsis root Airyscan image showing expression of FCA-mScarlet-I (red) and *FLC-lacO*/eYFP-LacI (yellow) in nuclei. Scale bars: 20 µm. (C) Subsets of Airyscan images from B showing nuclei co-expressing FCA condensates (red) and *FLC-lacO*/eYFP-LacI (yellow). Scale bars: 2 µm. The FCA-mScarlet-I and LacI-eYFP signals were enhanced by shifting their dynamic ranges to 0–1099 and 0–4220 in EV5B, and to 0–1000 and 0–6000 in EV5C. The images were acquired on a Zeiss LSM980 using LD C-Apochromat 40x/NA 1.1 water-immersion objective. The fluorescent proteins, eYFP and mScarlet-I, were excited at 488 and 561 nm, respectively.

Appendix Figure S1.
The co-localization of 3D images is achieved after object identification using the blob finder algorithm
Front view (A) and side view (B) of the 3D image with grid lines, consistent with Figure 2A, illustrating the FCA-mTurquoise, FLL2-eYFP, and the objects identified using the blob finder algorithm, as described in the Methods. In the identified objects, green dots denote FCA-mTurquoise identified foci, magenta dots indicate FLL2-eYFP foci identified, and the white region highlights the co-localised area.

Appendix Figure S2. An additional experiment analysing the co-localization of FCA and FLL2
(A) Quantification of size of FCA (total FCA-mTurquoise identified foci), FLL2 (total FLL2-eYFP identified foci) and FCA identified foci that are co-localized with FLL2. The lines in the violin plots indicate the median and quartiles. (B) Size and frequency distribution of FCA, FLL2 foci and FCA identified foci that are co-localized with FLL2. (C) Number of FCA, FLL2 foci and FCA foci that are co-localized with FLL2 in each nuclei. The lines in the violin plots indicate the median and quartiles. The images for the quantification were acquired on a Zeiss LSM980 using LD C-Apochromat C-Apochromat 63x/NA 1.2 water-immersion objective. The fluorescent probes, mTurquoise2 and eYFP, were excited at 458 and 514 nm, respectively. The voxel size of images for quantification analysis was 0.05 µm x 0.05 µm x 0.21 µm (xyz). The quantitative analysis was based on 7 nuclei.

Appendix Figure S3. Stoichiometry distribution of individual FCA particles in FCAwt and FCArrm genotypes in different experiments

Appendix Figure S4. Integrated nuclear intensity in FCAwt and FCArrm genotypes from different experiments

The violin plots represent the integrated nuclear intensity (in molecules) of FCA reporters in FCAwt and FCArrm from each biological replicate. Col-0 is negative control.

Tracking number				
	Replicate 1	Replicate 2	Replicate 3	Total
FCAwt	504	202	302	1008
FCArrm	560	171	182	913

Appendix Figure S5. Quantification of FCA particle stoichiometry and diffusion coefficient
The violin plots represent the FCA reporter stoichiometry and diffusion coefficient in FCAwt and FCArrm from each biological replicate. The table displays the sample statistics used for quantitative SlimVar analysis, including the numbers of tracks.

FCA-mScarlet-I

FCA-eGFP

Exposure time (ms) 0.1*   1*   10   100*   1000*

Noise   Motion blur

Appendix Figure S6. SlimVar at 10 ms/frame provides optimal contrast for detection of FCA-mScarlet-I and FCA-eGFP particles
A single acquisition at 10 ms/frame exposure was used to simulate a range of exposure times (labelled with *). Longer exposure times were generated by summing 10 or 100 frames directly, while shorter exposures were created by dividing a frame's pixel intensity by 10. This was done either with FCA-mScarlet-I or FCA-eGFP. Frames are displayed with individual min/max contrast stretching. Below 1 ms/frame, shot noise reduces the contrast to a level insufficient for observing single molecules, whereas at 100 ms/frame, motion blur and photobleaching degrade the contrast for all but the brightest and least mobile assemblies. Scale bar: 2 μm.

Replicate 2

Replicate 3

Appendix Figure S7. Two additional independent DRIPc–qPCR experiments were performed to analyze the *COOLAIR* R-loop in the indicated genotypes

Data are presented as mean from four technical repeats for Replicate 2 and three technical repeats for Replicate 3. TSS, transcription start site. The primer sets used for DRIPc-qPCR (A to E) corresponds to that shown in Figure 4D.