Title

Negative selection allows human primary fibroblasts to tolerate high somatic mutation loads induced by N-ethyl-N-nitrosourea

Authors

Johanna Heid[1, =], Ronald Cutler[1, 2, =], Shixiang Sun[1], Moonsook Lee[1], Alexander Y. Maslov[1, 3], Xiao Dong[4,5], Simone Sidoli[2] and Jan Vijg[1, 6, *]

Affiliations

[1] Department of Genetics, Albert Einstein College of Medicine, Bronx, NY 10461, USA

[2] Department of Biochemistry, Albert Einstein College of Medicine, Bronx, NY 10461, USA

[3] Laboratory of Applied Genomic Technologies, Voronezh State University of Engineering Technology, Voronezh 394000, Russia

[4] Institute on the Biology of Aging and Metabolism, University of Minnesota, Minneapolis, MN 55455, USA

[5] Department of Genetics, Cell Biology and Development, University of Minnesota, Minneapolis, MN 55455, USA

[6] Center for Single-Cell Omics, School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

[=] Authors contributed equally to this work

[*] Correspondence to J.V. (jan.vijg@einsteinmed.edu)

Summary

Single-cell sequencing has shown that thousands of mutations accumulate with age in most human tissues. While there is ample evidence that some mutations can clonally amplify and lead to disease, the total burden of mutations a cell tolerates without functional decline remains unknown. Here we addressed this question by exposing human primary fibroblasts to multiple, low doses of N-ethyl-N-nitrosourea (ENU) and analyzed somatic mutation burden using single-cell whole genome sequencing. The results indicate that individual cells can sustain ~60,000 single-nucleotide variants (SNVs) with only a slight adverse effect on growth rate. We provide evidence that such high levels of mutations are only tolerated through negative selection against variants in gene coding regions, and in sequences associated with genetic pathways for maintaining basic

cellular function and growth. Since most tissues in adults are non-dividing, these results suggest
that somatic mutations in the absence of negative selection may have functionally adverse effects.

Key words

Aging; Somatic mutations; Mutation burden; N-ethyl-N-nitrosourea; Single-cell whole-genome
sequencing; Negative selection; Coding regions; Cell growth pathways; Human lung fibroblasts

Introduction

Aging is a complex process of continuous decline in cellular and tissue function that is
accompanied with increased disease risk [1]. DNA damage is thought to play a central role in this
process and may be a universal cause of aging [2]. Among the major molecular consequences of
DNA damage during aging are somatic DNA mutations. Somatic mutations are the result of errors
during repair or replication of a damaged DNA template and cannot be corrected except by the
death of the cell or organism. The accumulation of somatic mutations has since long been
implicated as a major cause of aging [3, 4], which is strongly supported by the age-related
exponential increase of cancer, a disease known to be caused by mutations [5]. Somatic mutations
are difficult to analyze because of their random nature, with very low variant allele frequencies.
With the recent emergence of single-cell and single-molecule assays this situation has changed.
There is now ample evidence that somatic mutation rate is far higher than the germline mutation
rate and that thousands of somatic mutations accumulate with age in most if not all human tissues
[6-9] [10], including postmitotic tissues such as the brain [11]. Evidence has now also been
provided that postzygotic, somatic mutations are a cause of a wide variety of human diseases other
than cancer [12, 13]. Like cancer, many of these age-related effects may be due to clonally
amplified mutations [14-16]. Indeed, in blood, somatic mutations lead to clonal hematopoiesis,
which has been associated with increased risk of age-related diseases, such as cardiovascular
disease, as well as with overall mortality [17].

What thus far remains unknown is the possible functional impact, collectively, of thousands of
random mutations accumulating in a cell, for example, by adversely affecting gene-coding and
gene-regulatory sequences relevant for the cell type studied. If so, one would expect a limit to the
burden of random mutations a cell can tolerate. To directly test this, we treated IMR-90 human
primary fibroblasts with low, repeated doses of N-ethyl-N-nitrosourea (ENU), a powerful mutagen
that induces almost exclusively single-nucleotide variants (SNVs). Our results show that repeated
ENU treatment linearly increased mutation frequency up to ~60,000 SNVs per cell, without a
major effect on growth rate. We show that such a high tolerance for somatic mutations can be
explained by negative selection against potentially deleterious mutations in gene coding regions
and in functional genomic regions associated with cell growth. These results suggest that

accumulating somatic mutations may have adverse effects on function when negative selection is
not possible, as is the case in most cells and tissues in adult organisms [18].

Results

<u>Repeated Mutagen Treatment Only Slightly Affects Cell Growth</u>

To determine the impact of increased somatic mutation burden on primary human cells *in vitro*,
we repeatedly treated human fetal lung fibroblasts (IMR-90) a total of 9 times with a sublethal
dose of 50 µg/mL of N-ethyl-N-nitrosourea (ENU), followed by a recovery period (Methods). To
monitor cell growth and toxicity, we measured cell number and apoptosis rate at 3 and 7 days after
each treatment to assess both short- and long-term effects of the treatment (Methods). The study
design is schematically depicted in Figure 1A. We found that repeated treatment with ENU
gradually decreased cell growth rate over the course of the experiment, resulting in a total of
28.25±1.87 (mean ± standard deviation) population doublings (PDs) for control cells and
24.1±1.34 PDs for ENU-treated cells at cycle 9 (Figure 1B). The effect of ENU treatment on cell
growth was more pronounced at the 3-day time point (Figure S1A) compared to the 7-day time
point (Figure S1B), which explains the decrease in overall growth rate (Figure 1B). This would be
expected due to adverse effects of a DNA damage response directly following mutagen treatment
and subsiding over time. Indeed, we found slight increases in early or late apoptosis at the 3- and
7-day time points, albeit neither was statistically significant (Figures 1C-D, and S1C-D).

To analyze possible effects of a DNA damage response on the control and ENU-treated cycling
cells in more detail, we performed bulk mRNA-sequencing on samples collected at cycle 1 and 9
for the control group and cycle 9 for the ENU-treated group (Figures S2A-K; Table S1; Methods).
We first compared the control cells at cycle 1 to the control cells at cycle 9 which resulted in 979
significantly differentially expressed genes (Figure 1E; Table S2) that were significantly enriched
in genes related to senescence and DNA repair (Figures 1F and S2L), not unexpected during cell
passaging, even under the low oxygen levels the cells were cultured (Methods). Interestingly, there
were no differentially expressed genes due to ENU treatment at cycle 9 (Figures 1G and S2M-N,
Supplemental note 1), although senescence genes were still found to be significantly enriched
(Figure 1H). Therefore, both the slightly increased rates of apoptosis and cellular senescence could
explain the observed phenotypes of slowed cellular growth (Figures 1B-D). Altogether, these

results indicate high resilience of proliferating cells in culture to low doses of ENU, which is a
very powerful mutagen, but not very toxic [19].

<u>Repeated Mutagen Treatment Results in Extremely High Mutation Burden</u>

Since somatic DNA mutations are random, they cannot be distinguished from sequencing artifacts
after direct whole genome sequencing. Hence, we used our established and highly accurate single-
cell whole genome sequencing assay [20, 21] to analyze mutation burden in 21 single-cells from
control and ENU-treated groups collected in 3 independent batches at cycles 3, 6, and 9 (Figure
1A, Table S3). Mutation frequencies were assessed at the 7-day time point after each treatment, as
DNA damage responses should be abated by that time and all mutations fixed (Figures S1B-D).
After correction for genome coverage and sensitivity (Figures S3A-G; Methods), we found the
number of estimated single nucleotide variants (SNVs) per cell in the control group increased from
2,434±406 at cycle 1 to 19,064±823 in the ENU-treated group at cycle 3, i.e. a 7.8-fold increase
(Figure 2A and S3H). At cycle 9, there was a ~32-fold increase in estimated SNVs per cell in the
ENU-treated group relative to the controls, where the total amount of estimated SNVs per cell in
the ENU-treated group reached a maximum of 55,954±4,066 (Figure 2A and S3H). The increase
in SNVs over the course of the experiment was best fit by a linear model (Figure 2B). Importantly,
there was no significant difference between control cells at cycle 1 and 9, ruling out that the
increase in SNVs in ENU-treated cells was due to passaging (Figure 2A and S3H). In addition,
these results were orthogonally validated with our recently developed single-molecule assay [22]
(Figures S4A-D; Table S4; Supplementary note 2).

While ENU is known to mainly induce point mutations through alkylation damage, the estimated
number of small insertions and deletions (INDELs) per cell was also observed to increase due to
ENU treatment, from 123±49 in the control group to 246±49 in the ENU-treated group at cycle 3,
without further increase, i.e. a 1.8-fold increase (Figures 2C and S3I). While rare, INDELS are
induced by ENU, albeit at low frequencies [23], and could be caused by DNA polymerase slippage
during DNA replication at sites of alkylation damage. As with SNVs, there was no significant
difference in the amount of estimated INDELs between control cells from cycle 1 and cycle 9
(Figures 2C and S3I). However, as opposed to the linear increase in SNVs with increasing cycles
(Figure 2B), there was no further increase in the estimated INDELs per cell in the ENU-treated
cells at cycle 6 or cycle 9 relative to cycle 3 (Figures 2C and S3I), suggestive of negative selection
acting to prevent a further increase. This can also be seen by the downward trend of the
INDEL/SNV ratio in the ENU-treated cells from cycle 3 to cycle 9 (Figure 2D and S3J).

We then characterized the underlying SNV mutational processes by combining cells from all
cycles within the control and ENU-treated groups and found increases in the relative contribution
of thymine-to-adenine (T>A) and thymine-to-cytosine (T>C) base transversions (Figures 2E and
S5A). This is in keeping with previously published ENU-induced mutational spectra by us and
others [19, 22]. We then inferred SNV mutational signatures *de novo* using non-negative matrix

factorization [24] (Figure S5B) and identified two signatures, named SBSA and SBSB, which were
dominant in the cells of the control and ENU-treated groups, respectively (Figure 2F). SBSA
resembles COSMIC human signature SBS40 (Figure 2G), while SBSB resembles COSMIC
human signature SBS5 (Figure 2H). Both mutational signatures are known to correlate with aging
[24]. Importantly, we also found that SBSB resembled the SIGNAL human signature associated
with ENU-treatment, validating the source of the induced mutations (Figure 2I) [25]. Though
relatively few INDELs were observed (369 in total), we also provide a characterization of their
mutational spectra (Figure S5E, Supplementary note 3).

Cells With Extremely High Mutation Burden Show Signs of Negative Selection Against
Damaging Variants

Based on our results showing that the mutation burden of primary cells could reach an estimated
~60,000 SNVs per cell (Figure 2A), i.e., ~10 times the SNV burden typically found in cells from
aged tissues [6], without a substantial effect on cell growth (Figures 1B and 2A), we hypothesized
that the sustained cell survival could be explained in part by negative selection. This was tested by
analyzing the 108,600 SNVs that were observed collectively in all cycles of both controls and
ENU-treated cells (Figure S3H). From these SNVs, we first annotated the different types of coding
region variants, (i.e., synonymous, stop retained, missense, start loss, stop gain, and stop loss)
using the Ensemble Variant Effect Predictor program [26] and then determined their frequency
(Methods). This revealed significant cycle-dependent increases in both synonymous and non-
synonymous variants in the ENU-treated group relative to the controls, with non-synonymous
variants reaching up to 80±35.3 variants per cell (73%±5.6% of all coding variants per cell) at
cycle 9 in the ENU-treated cells (Figure 3A, right panel). We then assessed for selection by
calculating the Log_2 observed/expected (O/E) ratios of the variants observed in all cells of each
cycle-condition group, which allows one to compare the observed frequency of variants to the
frequency expected by chance, i.e., neutral selection (Methods). Using this approach, O/E ratios
of synonymous variants were not significantly different than what was expected by chance at all
cycles in the ENU-treated groups, indicative of neutral selection as would be expected (Figure 3B,
left panel). O/E ratios of non-synonymous variants at cycle 1 and 9 of the control group were also
indicative of neutral selection (Figure 3B, right panel). However, it should be noted that the
numbers of mutations in the control group were very low (Figure 3A, left panel), which does not
allow for concrete conclusions. By contrast, the O/E ratios at cycle 6 and 9 in the ENU-treated
group were significantly less than what would be expected by chance, indicative of negative
selection (Figures 3B, right panel and S6A-B). From the O/E ratios of non-synonymous variants
in the ENU-treated group, it is estimated that these correspond to protection against 14.5±9.1, and
19.9±12.5 non-synonymous variants per cell at cycles 6 and 9, respectively (Figure 3C).

The signal of negative selection for non-synonymous variants in the ENU-treated group could
mostly be attributed to a significant depletion of missense variants (Figures 3D, bottom panel, and

S6C), which accounted for the majority of the coding region variants (Figure 3D, top panel). The
same trend (although not significant due to the limited number of observed variants) could be seen
for stop-loss variants, but not for start-lost or stop-gain variants (Figures 3D, bottom panel, and
S6C). The finding of negative selection against non-synonymous mutations in the ENU-treated
group was confirmed by examining the ratio of non-synonymous variants to synonymous variants
(N/S). While there was no difference of the observed N/S ratio (Figures 3E and S6D), when
analyzing this ratio using the O/E method, it showed a cycle-dependent decrease in the ENU-
treated group that was significantly less than what would be expected from chance alone at cycles
6 and 9 (Figure 3F).

We then used the same Variant Effect Predictor program [26] to annotate the different types of
non-coding region variants (accounting for >99% of all variants) to also test for selection (e.g.
introns, splice sites, etc.). As with the coding variants, non-coding variants showed cycle-
dependent increases in the ENU-treated group relative to the controls, with variants in intronic and
intergenic regions accounting for 49.9% and 35.1% of all non-coding variants, respectively (Figure
3G, top panel). For both control and ENU-treated groups at all cycles, O/E ratios of intronic region
variants were significantly less than what would be expected by chance, while intergenic region
variants were significantly greater than what would be expected by random chance (Figure 3G,
bottom panel, See supplementary note 4 for discussion of intergenic region results). The O/E ratio
result for the intronic regions was an expected finding due to transcription coupled repair and was
used to normalize the O/E ratios of coding and select non-coding variants (Methods). Out of all
the non-coding variants, only O/E ratios of variants occurring in 3' untranslated regions (UTR),
and splice regions in the ENU-treated group at cycle 9 were significantly less than would be
expected by chance alone (Figures 3G, bottom panel and S6F), which indicates negative selection
for mutations in these functional regions despite being outside of the coding region. The O/E ratios
of 3' UTRs and splice sites in the ENU-treated group at cycle 9 corresponded to protection against
19.3 ± 12.5 and 6.2 ± 2.7 variants, respectively (Figure 3H).

Taken together, these results indicate that damaging variants in both coding and a subset of non-
coding regions do not accumulate to levels that would be expected in the context of the extremely
high mutation burden induced by ENU and instead were subject to negative selection, likely to
protect critical genes supporting essential cellular functions *in vitro*. These results may explain
how a population of cells exposed to multiple mutagen treatments is able to avoid a dramatic
decline in growth rate (Figure 1B), even as overall SNV burden continued to increase linearly
throughout the experiment (Figure 2B). Of note, since in our selection studies we could only
analyzed the mutations that were actually observed, which are a fraction (~20%) of the total
estimate of ~60,000 mutations at cycle 9 in the ENU-treated group (Figure S3H), these results
have relevance for the much smaller numbers of SNVs that are estimated to accumulate during
normal aging, i.e., from ~3,000 in human B lymphocytes to ~ 9,000 in human hepatocytes [6].

Pathways Supporting Lung Fibroblast Growth are Protected from Mutation Accumulation

To investigate if negative selection may prevent mutations to adversely affect functional pathways as a whole, we analyzed the distribution of SNVs across functional genomic regions relevant to the cells under study, i.e. exons, upstream promoters, and enhancers of all expressed genes (13,849 genes) and non-expressed genes (12,697 genes) derived from our bulk mRNA-sequencing data (Figures S2 and S7A-B; Table S5; Methods) [27-29]. We then calculated the frequency and O/E ratios of SNVs within these regions (Methods). This revealed significant cycle-dependent increases of SNV frequency in the ENU-treated group relative to the controls, as would be expected (Figure 4A). O/E ratios in functional regions associated with expressed genes were significantly less than what would be expected by chance at all cycles in the ENU-treated group and at cycle 9 in the control group (Figures 4B right panel, and S7C-D, right panels), while they did not deviate from expected by chance alone in functional regions associated with non-expressed genes at all cycles in both groups (Figures 4B left panel, S7C-D, left panels). This is indicative of negative selection against mutations in the functional regions of expressed genes, which appears to be sufficient for cell survival regardless of the mutation burden.

Moving to the pathway level, we identified 399 gene ontology (GO) biological processes as significantly enriched in the set of expressed genes from our bulk RNA-seq data (Figure 4C; Table S6; Methods). The mean frequency of SNVs within each pathway, although low, showed significant cycle-dependent increases of SNV frequency in the ENU-treated group relative to the controls (Figure 4D; Table S7). At a global level, O/E ratios of SNVs across all pathways showed a cycle-dependent decrease in the ENU-treated group, but not in the controls, indicative of increasing negative selection strength at the pathway level (Figures 4E and S7E-F; Table S7). At cycle 9 in the ENU-treated group, 6 pathways showed O/E ratios significantly below zero, which is indicative of a depletion of SNVs within the functional sequences corresponding to these pathways (Figures 4E-F and S7E-F; Table S7). The pathways under negative selection were characterized as those related to basic cellular processes supporting lung fibroblast cell growth *in vitro*, such as 'response to oxidative stress' (GO:0006979; protected against 9.3±6.7 SNVs), 'respiratory system development' (GO:0006979; protected against 4.3±6.9 SNVs), 'cellular response to chemical stress' (GO:0062197; protected against 6.3±6.9 SNVs), 'respiratory tube development' (GO:0030323; protected against 3.8±6.7 SNVs), 'proteasome-mediated ubiquitin-dependent protein catabolic process' (GO:0043161; protected against 6.2±6.9 SNVs), and 'regulation of response to DNA damage stimulus' (GO:2001020; protected against 3±6.3 SNVs). As a negative control, we employed bootstrapping to randomly shuffle the genes amongst the identified pathways before calculating the O/E ratios, which validated our findings of the pathways under negative selection at cycle 9 in the ENU-treated group (Figures S8A-D; Supplementary note 5; Methods).

251 In summary, these results suggest that in the context of very high mutation burden, pathways
252 supporting stress response, cell proliferation, protein homeostasis, and metabolism *in vitro* are
253 protected from the accumulation of SNVs by negative selection. This highlights a mechanism by
254 which dividing cells can survive high mutation burden and underscores the functional impact of
255 random somatic SNVs on genetic pathways active in proliferating lung fibroblasts.

Discussion

Somatic mutations are known since the 1970s to be the main cause of cancer [30]. Mainly because
of the exponential increase in cancer risk with age [31], aging itself has been suggested to be
caused by the accumulation of somatic mutations [3, 8, 32]. Testing this hypothesis has long been
constrained by a lack of methods to quantitatively analyze somatic mutations in tissues and organs
during aging. However, both single-cell and single-molecule methods have recently been
developed to accurately quantify SNVs and INDELs. Using these methods, we and others have
demonstrated that somatic mutation burden is far higher than expected based on the known
germline mutation frequency and increases with age up to several thousands of mutations per cell
in aged subjects depending on the tissue [6, 9]. However, a causal relationship of increased
mutation burden and age-related functional decline and disease risk remains untested. Instead,
attention has been focused on clonally amplified somatic mutations and their causal relationship
to non-cancerous and age-related diseases [14, 33].

In the present study we tested for a possible limit to somatic mutation accumulation in a cell culture
model in which high levels of somatic mutations were induced by multiple low doses of the
powerful mutagen ENU (Figure 1A). In this system we were able to linearly increase the somatic
mutation burden to ~60,000 SNVs per cell, over 30-fold higher than untreated control cells (Figure
2B). Remarkably, this supraphysiological mutation load only slightly impacted cell growth rate
(Figure 1B), suggesting that most cells can sustain an astounding level of genetic damage. The
most straightforward explanation for these findings is that the mammalian genome is highly robust,
and that even very high numbers of point mutations are unlikely to adversely affect cellular
function. Indeed, this would be supported by results from a recent study in which as many as
~30,000 SNVs were observed in intestinal crypts of elderly patients with genetic defects in DNA
mismatch repair [34].

An alternative explanation for the high tolerance of actively proliferating fibroblasts for mutation
accumulation is negative selection against mutations affecting growth rate and cell viability.
Evidence for negative selection has been found for somatic mutations in the cancer genome [35].
However, in cancer, positive selection is dominant resulting in many driver mutations [36].
Evidence for negative selection of somatic mutations in normal cells is scarce because of the much
lower numbers of somatic mutations available. Our present analysis of untreated, control cells
suffered from this limitation, and we were unable to convincingly demonstrate negative selection
of mutations in these cells. However, our ENU-treated cells yielded a sufficiently high number of

SNVs to test for negative selection (Figure S3H) (positive selection does not occur over the limited number of population doublings of the experiment). We found multiple lines of evidence for negative selection acting on potentially deleterious coding and non-coding region variants. While the frequency of both synonymous and non-synonymous variants increased linearly with ENU treatment (Figure 3A), deeper analysis using observed over expected (O/E) ratios revealed a significantly smaller number of high impact non-synonymous variants, especially missense, than would be expected by chance alone in the ENU-treated cells (Figures 3B-C, 3D, bottom panel). This suggests that an actively proliferating population of fibroblasts have the capacity to selectively purge the most harmful coding mutations from their genomes, preserving cellular fitness despite the extremely high number of random mutations accumulating genome wide. Negative selection was also detected in variants occurring in 3' untranslated regions or splice sites, which made up ~1.1% of all the observed non-coding variants (Figures 3G, bottom panel and 3H).

Evidence for negative selection of somatic mutations was also found at the pathway level, which revealed significantly less mutations in functional genomic regions associated with genes actively expressed in pathways relevant to growth and metabolism of the primary lung fibroblasts treated with ENU (Figure 4E), such as response to oxidative stress, respiratory system development, and regulation of response to DNA damage stimulus (Figure 4F). This suggests that negative selection acts to preferentially maintain the integrity of pathways critical for cellular survival and proliferation, despite the massive somatic mutational burden across the genome overall.

These results provide the first evidence that very high numbers of somatic mutations, collectively, have a functional impact on cells above and beyond clonally amplified mutations. However, they also indicate that negative selection is active on the much lower numbers of somatic mutations accumulating during normal aging. Indeed, while after 9 treatments with ENU we estimated as many as 55,954±4,066 SNVs per cell (Figure 2A), the actual number of observed SNVs that could be assessed was only as much as 11,623±4,820 SNVs per cell (Figure S3H), not that much higher than what has been found for cells from aged human subjects, i.e., between 3,000 and 9,000 SNVs per cell, depending on the cell type [6]. These numbers are not solely determined by mitotic activity, which would lead to SNVs as replication errors. Indeed, even postmitotic cells as human neurons have been demonstrated to accumulate up to ~4,000 SNVs per cell on average in the dentate gyrus [11], while reversed postmitotic liver hepatocytes had as many as ~9,000 in people in their 70 [37].

Hence, it seems likely that negative selection is a general mechanism to eliminate somatic mutations that adversely impact cell function, similar to purifying selection in the germline. However, such a mechanism would only work in actively proliferating cells when negative selection is possible based on growth, similar to the situation in tumors. In normal adult subjects, cell division rate is very low with few exceptions, most notably the lymphoid and intestinal systems [15]. In human B lymphocytes, we previously demonstrated that SNVs accumulate with age much slower in the functional part of the genome than in the genome overall, indicating negative selection of mutations arising during the process of B cell development [38]. By contrast,

somatic mutations accumulating in most adult cells will inevitably affect coding or regulatory sequences associated with the particular function these cells are programmed to carry out. However, such mutations cannot be cleared through negative selection in the absence of a relevant cell fitness advantage. Therefore, our present results suggest that somatic mutations can functionally impact critical physiological functions at old age, especially as the force of negative selection has been shown to decline with aging[39].

In this study we only analyzed SNVs, and the few small insertions and deletions induced by ENU. However, SNVs are much less likely to have a functional impact than genome structural variants, including large deletions, translocations, and copy number variants. Thus far, there is very little information about somatic SVs, which cannot be detected very well, either using single-cell or single-molecule methods. While their relative frequency is much lower than that of SNVs (~100X), SVs can have a major phenotypic impact, as indicated by germline SVs in human diseases [40]. SVs are much more likely to adversely affect cell function than SNVs, but unless they result in cell death, a mechanism to select against them *in vivo* during aging is difficult to imagine.

In summary, our present findings demonstrate that actively proliferating human cells possess a remarkable capacity to tolerate extremely high levels of random somatic mutations, specifically SNVs, primarily through negative selection against the most damaging variants in coding regions and preservation of genomic integrity of critical cellular pathways. These results have important implications for our understanding of the role of somatic mutations in aging and age-related diseases, where the ability to effectively select against deleterious mutations may be limited compared to the *in vitro* culture conditions examined here.

Data Availability

Raw sequencing data has been deposited in the NCBI Sequence Read Archive under BioProject accession PRJNA1129131. Processed gene expression data has been deposited in the NCBI Gene Expression Omnibus database under accession GSE271867. Processed variant call files have been deposited in the NCBI dbSNP database and will be made available upon acceptance.

Code Availability

Scripts used for processing and analyzing data have been deposited at: figshare.com/articles/software/Negative_selection_allows_human_primary_fibroblasts_to_tolera te_high_somatic_mutation_loads_induced_by_N-ethyl-N-nitrosourea/26110453

Acknowledgements

This study was supported by National Institute of Health grants: T32GM007491 (R.C.) T32AG023475 (R.C.), U19AG056278 (J.V.), U01HL145560 (J.V.), U01ES029519 (J.V.), P01AG017242 (J.V.), P01AG047200 (J.V.), RF1AG068908 (J.V.), and P30AG038072 (J.V.). Other funding sources: US Department of Defense grant BC180689P1 (J.V.), Glenn Foundation for Medical Research Postdoctoral Fellowships in Aging Research (J.H.), The Michael J. Fox Foundation, and The Paul F. Glenn Center for the Biology of Human Aging at Albert Einstein College of Medicine.

Author Contributions

A.Y.M., and J.V. conceived this study and designed the experiments. J.H., M.S., and A.Y.M performed the experiments. J.H., R.C., and S.S. analyzed the data. X.D. provided guidance on the data analysis. J.H., R.C., and J.V. wrote the manuscript.

Declaration of interests

M.L., A.Y.M., X.D., and J.V. are co-founders and shareholders of SingulOmics Corp. A.Y.M., and J.V. are co-founders of MutagenTech Corp. Other authors declare no conflict of interest.

Declaration of generative AI and AI-assisted technologies in the writing process

The authors employed ChatGPT for spelling and grammar checks during this work's preparation. They have reviewed and edited the content accordingly, assuming full responsibility for the publication's content.

Materials and methods

Cell culture and treatment

Human IMR-90 primary fibroblasts were obtained from the American Tissue Culture Collection (ATCC; LOT number 70020198). They were cultured in DMEM (Dulbecco's Modified Eagle Medium, from ATCC) supplemented with 10% fetal bovine serum (FBS, Gibco/Life Technologies) and 1% Penicillin-Streptomycin solution (Gibco/Life Technologies) at 3% oxygen and 37°C. Cells obtained from ATCC were at passage 11 upon arrival and were cultured for an additional 5 passages before the start of the experiment. Cells were then treated for a total of 9 times over the course of several months with ENU. Cells were treated with medium containing 50 µg/ml ENU 24 hours after seeding 1 million cells in a 10 cm dish. After 3 days cells were counted,

396	1 million cells re-seeded, and any remaining cells were collected and stored at -80°C. The same
397	procedure was repeated 7 days after treatment. All downstream analysis was performed on samples
398	collected 7 days after the last treatment unless stated otherwise.

| 399 | |

| 400 | Cell growth and apoptosis assays |

401	Cells were counted using the ViaCount assay (Cytek) on a Guava EasyCyte flow cytometer to
402	assess cell number and viability at each time point. Apoptosis rates were determined utilizing the
403	Nexin assay (Cytek) on the same machine.

| 404 | |

| 405 | Bulk mRNA transcriptome analysis |

406	RNA was extracted from frozen cell pellets using the Zymo DNA/RNA extraction kit (Microprep
407	plus D7005) according to the manufacturer's instructions and quantified using a Qubit kit (Thermo
408	Fisher Scientific, USA). Libraries for mRNA analysis were prepared and sequenced using 150
409	paired-end mode on an Illumina Novaseq platform by Novogene Corp Inc., CA. Raw sequence
410	reads were processed using the nf-core RNA-Seq pipeline (https://nf-co.re/rnaseq/3.10) to obtain
411	gene expression levels [41]. Gene filtering was performed using the zPKM program (Figure S2A)
412	[42]. Count normalization and differential expression testing was performed using DESeq2 [43]
413	where the design was set to test for the effect of the cycle-condition variable while controlling for
414	the batch variable (Supplemental note 1). Gene set enrichment analysis (GSEA) was performed
415	using the clusterProfiler program [44] along with the SenMayo gene set [45] and KEGG DNA
416	repair gene sets [46].

| 417 | |

| 418 | Single-cell isolation, whole-genome amplification, library preparation, and sequencing |

419	Single cells were isolated using the CellRaft AIR system (Cell Microsystems). The CellRaft array
420	was prepared following the manufacturer's instruction and washed three times for 3 minutes with
421	warm PBS before addition of cell suspension. 1,000 cells were seeded in 1ml for one CellRaft
422	array. Cells were allowed to settle for 3-4 hours at 37°C before isolation. Then, the raft was
423	carefully washed with warm medium to remove potential debris and dead cells. Individual rafts
424	containing one cell were deposited in PCR tubes with 2.5 µl PBS and stored at -80°C until further
425	processing. Single-cell whole-genome amplification and library preparation for WGS was
426	performed as previously described [12]. In brief, single-cells were whole genome-amplified using
427	SCMDA and subject to quality control using a locus dropout test [47]. Libraries were prepared
428	using a NEBNext Ultra II FS kit (NEB). Library quantity and size were assessed with a Qubit kit
429	and TapeStation, respectively. WGS libraries were sequenced using the 150 paired-end mode on
430	an Illumina Novoseq platform by Novogene Corp Inc., CA.

| 431 | |

<u>Single-cell whole-genome read alignment and mutation calling</u>

Data analysis was performed as previously described [21]. Briefly, raw sequencing data underwent
quality control using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and
Trim Galore (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/), followed by
alignment to the human hg19 reference genome using the GATK best practices pipeline. This
included marking adapters, sequence alignment, base quality score recalibration, and variant
calling for germline variants using GATK HaplotypeCaller (https://gatk.broadinstitute.org/hc/en-
us/articles/360037225632-HaplotypeCaller).   SCcaller   2.0
(https://github.com/biosinodx/SCcaller) was then employed to call somatic single-nucleotide
variations (SNVs) and small insertions and deletions (INDELs), adjusting for allelic amplification
bias using known germline variants (Figures S3A-C). Estimated mutation burden was calculated
as follows:

445
$$\text{Estimated somatic SNVs|INDELs per cell} = \frac{\text{observed somatic SNVs|INDELs}}{\frac{surveyed\ genome\ (bp) * sensitivity}{size\ of\ genome\ (bp)}}$$

<u>Mutation filtering and identification of technical artifacts in single-cell whole-genome mutation</u>
<u>data</u>

Overlapping SNVs and INDELs were observed amongst the single-cells from batches 2 and 3
(Figures S3D-E; Table S3). Since this was not observed in the first batch, it is unlikely that these
mutations are the result of clonal amplification. Mutations that were shared between all cells within
a batch were thus considered to be germline and were filtered out prior to downstream analysis
(Figures S3F-G).

Technical artifacts inherent to single-cell MDA library preparation steps, such as overheating
during cell lysis or whole genome amplification, can be identified by their specific mutational
signatures scE and scF [48]. We checked for the presence of these signatures and found scF to be
present in the signature extracted from control cells at cycle 1 (Figure S5A, C-D). The scF technical
artifact is characterized by higher than expected cytosine-to-thymine (C>T) transversions at G<u>C</u>N
contexts. Thus, control cells from cycle 1 were excluded from the downstream mutational
signature analyses.

<u>Mutation signature analysis</u>

The MutationalPatterns program [49] use used to analyze mutational spectra and signature analysis
at the level of single-cells cycle-condition groups, respectively. To determine the number of
mutational signatures to be extracted from the data, a non-negative matrix factorization rank

survey was employed which resulted in 2 factors (Figure S5B). *De novo* signatures were extracted
and then compared to those from the COSMIC and SIGNAL databases using cosine similarity [50,
51].

Variant annotation

Observed mutations in each single-cell and simulations of single-cells (where mutations were
randomly re-distributed as described in the section below) were annotated using the Ensembl
Variant Effect Prediction program and the given 'Consequences' annotation was used for
downstream analysis [26]. In the case where mutations were predicted to lead to more than 1
unique consequence (i.e. multiple entries of the same mutation), the highest ranking annotation
that occurred in the resulting output was kept which resulted in a 1:1 mapping of mutations and
consequences. In the case where mutations occurred in a region that overlapped multiple possible
consequences (e.g. splice region variant and intron variant), we classified this mutation to be
consistent with the consequence that occurred first in this annotation (e.g. splice region, if multiple
annotation was splice region variant and intron variant). For mutations annotated as splice variants
(i.e.   splice_acceptor_variant,   splice_donor_variant,   splice_donor_5th_base_variant,
splice_region_variant, splice_donor_region_variant, splice_polypyrimidine_tract_variant), we
classified these mutations as 'splice.' Synonymous variants were classified as mutations annotated
as synonymous_variant and stop_retained_variant. Non-synonymous variants were classified as
mutations annotated as missense_variant, start_lost, stop_gained, and stop_lost

Selection analysis of annotated variants

Selection analysis of mutations must account for the fact that deleterious mutations will result in
lower cellular fitness, which will in turn lead to lower frequencies of these mutations being
observed when sampling from a population of proliferating cells in culture (i.e. survivor bias).
Thus, selection analysis was performed by first creating 10,000 simulations of each single-cell in
our dataset to be as the null expectation (i.e. neutral selection) where mutations are randomly re-
distributed throughout the genome. This was done using the SigProfilerSimulator program [52],
where for each single-cell, observed mutations were randomly re-distributed across the genome
while accounting for cell-specific sequencing coverage, mutational signatures, and nucleotide
contexts.

Observed/expected (O/E) ratios were calculated for each single-cell as well as cycle-condition
groups by first summing the frequency of annotated consequences of all single-cells and their
respective simulations within each group. The O/E ratio was then calculated with respect to each
simulation (resulting in 10,000 O/E ratios per cell/group) as follows:

$$O/E = Log_2 \frac{(observed\ consequence\ freq. + 1)}{(simulated\ consequence\ freq. + 1)}$$

A permutation test was then used to determine the p-value of the observed consequence frequency
where the simulated consequence frequency served as the null distribution and is as follows:

505
$$p = \frac{1}{n}\sum_{i=1}^{n} I\,(simulated\ consequence\ freq._{\cdot i} \geq observed\ consequence\ freq.)$$

Where p is the permutation test p-value, n is the number of permutations (i.e. 10,000), i is the i-th
permutation, and I is the indicator function, which equals 1 if the condition inside is true, and 0
otherwise.

When assessing for mutation selection, it is also necessary to account for the cell-intrinsic bias in
mutation frequency due to DNA repair mechanisms (i.e. transcription coupled repair), which is
known to reduce mutation frequency in transcribed regions and is agnostic to damage that would
result in synonymous or non-synonymous variants [53]. This bias in mutation frequency was
indeed observed in our data and consistent with a depletion of mutations due to DNA repair (Figure
3G, bottom panel). Thus, we normalized the O/E ratio of coding variants (e.g. synonymous,
missense, etc.) by the O/E ratio of mutations which occurred within intronic regions and not in
splice regions (Figure 3G, bottom panel). We chose to use this normalization because the
occurrence of both exonic and intronic mutations is dependent on transcription coupled repair [54]
however, since intronic mutations do not alter the translated protein (with the exception of splice
site variants), they are not subject to selective mechanisms. In addition, the majority of variants in
our data occur within introns (~46.7%, Figure 3G, top panel), allowing us to make more confident
O/E ratio for these mutations as compared to using the O/E ratio of synonymous mutations which
only made up ~0.23% of all variants. A comparison of unnormalized and normalized O/E ratio for
non-synonymous and synonymous variants is shown by comparing Figured 3B to S6A, where it
can be seen to slightly increase the O/E ratio.

To estimate the number of variants that are protected against resulting from negative selection, we
subtracted the average number of variants from simulated samples from the number of variants in
the respective observed sample.

<u>Selection analysis of pathways</u>

Bulk mRNA transcriptome data was processed as described above. Samples from cycle 9 were
used to determine expressed and non-expressed genes which was performed using the zPKM
program (Figure S2A) [42]. Genomic coordinates of exons, introns (used for normalization),
promoters (5KB upstream of transcription start site), and annotated enhancers [27] were retrieved
for expressed and non-expressed genes. Over representation analysis for gene ontology biological
processes was then performed on the list of expressed genes using the clusterProfiler program [44].
For the enriched biological process, we randomly shuffled the genes associated between all other
identified biological processes which resulted in 10,000 permutations of each biological process
that contained a random set of genes. O/E ratios and p-values were then calculated for each cell as
well as cycle-condition groups in the same manner as described above. P-values of O/E ratio for

each pathway were adjusted for multiple hypothesis tests using the Benjamini-Hochberg method.
A comparison of unnormalized and normalized O/E ratio for non-synonymous and synonymous
variants is shown by comparing Figures 4E to S7C, where it can be seen to slightly increase the
O/E ratio. Average O/E ratios were calculated for 10,000 permutations of each biological process
and served as a negative control to ensure that the signal of negative selection within a pathway
was not due to random chance (Supplemental note 5).

<u>Single-molecule mutation analysis</u>

DNA was extracted from frozen cell pellets using the Zymo DNA/RNA extraction kit (Microprep
plus D7005) according to the manufacturer's instructions and quantified using a Qubit kit (Thermo
Fisher Scientific, USA). Library preparation for single-molecule mutation sequencing (SMM-Seq)
was performed as previously described [22]. In brief, 200 ng gDNA was used as input for library
construction and then fragmented using restriction endonucleases Alu I and Mlu CI, followed by
purification with AMPure XP beads. The fragmented DNA was then processed with the NEBNext
Ultra II DNA Library Prep Kit for Illumina, utilizing a custom adapter. Post size selection with
AMPure XP beads, the product was quantified with a Qubit kit and analyzed on a TapeStation
(Agilent). For library preparation, samples were diluted based on molar concentration, and 1 µl of
the diluted sample was used as a template in a pulse-RCA reaction with a strand displacement
polymerase. The reaction product was purified, PCR-amplified, and further purified before
quantification and pooling for sequencing. Conventional whole-genome sequencing (WGS)
libraries that are used to identify germline variants were similarly prepared by PCR amplification
of adapter-ligated samples, followed by purification, quantification, pooling, and sequencing as
described above. WGS libraries were sequenced using 150 paired-end mode on an Illumina
Novoseq platform by Novogene Corp Inc., CA.

<u>Single-molecule whole-genome read alignment and mutation calling</u>

Data analysis for SMM-Seq data was performed as previously described [22]. In brief, raw
sequence reads were adapter- and quality-trimmed, then aligned to the hg19 reference genome.
Realignment and recalibration were performed based on known indels. For variant calling, filters
were applied to ensure only high-quality reads (mapping quality ≥ 60, in proper pairs, no secondary
alignments) were considered. Positions covered by at least seven reads from each strand within a
UMI family and covered at least 20× in regular sequencing were deemed qualified. Variants were
identified if all reads within a UMI family showed the same base, differing from the reference
genome. Germline variants were filtered out using a list of SNPs from the sample and dbSNP
[55]Variants were rejected if detected in different UMI families or conventional WGS data
containing germline variants. Somatic SNV frequency was calculated as the ratio of identified
variants to the total number of qualified positions. See supplemental note 2 for details of analysis.

References

1.   López-Otín, C., et al., *Hallmarks of aging: An expanding universe.* Cell, 2023. **186**(2): p.
243-278.
2.   Schumacher, B., et al., *The central role of DNA damage in the ageing process.* Nature,
2021. **592**(7856): p. 695-703.
3.   Failla, G., *The aging process and cancerogenesis.* Ann N Y Acad Sci, 1958. **71**(6): p. 1124-
40.
4.   Szilard, L., *ON THE NATURE OF THE AGING PROCESS.* Proc Natl Acad Sci U S A,
1959. **45**(1): p. 30-45.
5.   Laconi, E., F. Marongiu, and J. DeGregori, *Cancer as a disease of old age: changing*
mutational and microenvironmental landscapes. Br J Cancer, 2020. **122**(7): p. 943-952.
6.   Ren, P., X. Dong, and J. Vijg, *Age-related somatic mutation burden in human tissues.* Front
Aging, 2022. **3**: p. 1018119.
7.   Franco, I. and M. Eriksson, *Reverting to old theories of ageing with new evidence for the*
role of somatic mutations. Nat Rev Genet, 2022. **23**(11): p. 645-646.
8.   Franco, I., G. Revêchon, and M. Eriksson, *Challenges of proving a causal role of somatic*
mutations in the aging process. Aging Cell, 2022. **21**(5): p. e13613.
9.   Franco, I., et al., *Whole genome DNA sequencing provides an atlas of somatic mutagenesis*
in healthy human cells and identifies a tumor-prone cell type. Genome Biol, 2019. **20**(1):
p. 285.
10.   Franco, I., et al., *Somatic mutagenesis in satellite cells associates with human skeletal*
muscle aging. Nat Commun, 2018. **9**(1): p. 800.
11.   Lodato, M.A., et al., *Aging and neurodegeneration are associated with increased*
mutations in single human neurons. Science, 2018. **359**(6375): p. 555-559.
12.   Erickson, R.P., *Somatic gene mutation and human disease other than cancer: an update.*
Mutat Res, 2010. **705**(2): p. 96-106.
13.   Li, C. and S.M. Williams, *Human Somatic Variation: It's Not Just for Cancer Anymore.*
Current Genetic Medicine Reports, 2013. **1**(4): p. 212-218.
14.   Abyzov, A., et al., *One thousand somatic SNVs per skin fibroblast cell set baseline of*
mosaic mutational load with patterns that suggest proliferative origin. Genome Res, 2017.
27(4): p. 512-523.
15.   Laurie, C.C., et al., *Detectable clonal mosaicism from birth to old age and its relationship*
to cancer. Nat Genet, 2012. **44**(6): p. 642-50.
16.   Abyzov, A., et al., *Somatic copy number mosaicism in human skin revealed by induced*
pluripotent stem cells. Nature, 2012. **492**(7429): p. 438-42.
17.   Jaiswal, S., et al., *Age-related clonal hematopoiesis associated with adverse outcomes.* N
Engl J Med, 2014. **371**(26): p. 2488-98.
18.   Sender, R. and R. Milo, *The distribution of cellular turnover in the human body.* Nat Med,
2021. **27**(1): p. 45-48.
19.   Barbaric, I., et al., *Spectrum of ENU-induced mutations in phenotype-driven and gene-*
driven screens in the mouse. Environ Mol Mutagen, 2007. **48**(2): p. 124-42.
20.   Dong, X., et al., *Accurate identification of single-nucleotide variants in whole-genome-*
amplified single cells. Nat Methods, 2017. **14**(5): p. 491-493.
21.   Zhang, L., et al., *Analyzing somatic mutations by single-cell whole-genome sequencing.*
Nat Protoc, 2024. **19**(2): p. 487-516.
22.   Maslov, A.Y., et al., *Single-molecule, quantitative detection of low-abundance somatic*
mutations by high-throughput sequencing. Sci Adv, 2022. **8**(14): p. eabm3259.

23.   Watson, D.E., M.L. Cunningham, and K.R. Tindall, *Spontaneous and ENU-induced mutation spectra at the cII locus in Big Blue Rat2 embryonic fibroblasts.* Mutagenesis, 1998. **13**(5): p. 487-97.

24.   Alexandrov, L.B., et al., *The repertoire of mutational signatures in human cancer.* Nature, 2020. **578**(7793): p. 94-101.

25.   Kucab, J.E., et al., *A Compendium of Mutational Signatures of Environmental Agents.* Cell, 2019. **177**(4): p. 821-836 e16.

26.   McLaren, W., et al., *The Ensembl Variant Effect Predictor.* Genome Biol, 2016. **17**(1): p. 122.

27.   Boix, C.A., et al., *Regulatory genomic circuitry of human disease loci by integrative epigenomics.* Nature, 2021. **590**(7845): p. 300-307.

28.   Carlson M, M.B. *TxDb.Hsapiens.UCSC.hg19.knownGene: Annotation package for TxDb object(s).* 2015; Available from: https://bioconductor.org/packages/release/data/annotation/html/TxDb.Hsapiens.UCSC.hg19.knownGene.html.

29.   Boix, C.A. *EpiMap Repository.* 2023; Available from: https://compbio.mit.edu/epimap/.

30.   Nowell, P.C., *The clonal evolution of tumor cell populations.* Science, 1976. **194**(4260): p. 23-8.

31.   Group, U.S.C.S.W., *US cancer statistics: 1999–2009 incidence and mortality web-based report.* 2013, USDHHS, CDC: Atlanta GA.

32.   Burnet, F.M., *Intrinsic mutagenesis: a genetic basis of ageing.* Pathology, 1974. **6**(1): p. 1-11.

33.   Ng, S.W.K., et al., *Convergent somatic mutations in metabolism genes in chronic liver disease.* Nature, 2021. **598**(7881): p. 473-478.

34.   Robinson, P.S., et al., *Increased somatic mutation burdens in normal human cells due to defective DNA polymerases.* Nat Genet, 2021. **53**(10): p. 1434-1442.

35.   Zapata, L., et al., *Negative selection in tumor genome evolution acts on essential cellular functions and the immunopeptidome.* Genome Biol, 2018. **19**(1): p. 67.

36.   Martincorena, I., et al., *Universal Patterns of Selection in Cancer and Somatic Tissues.* Cell, 2017. **171**(5): p. 1029-1041.e21.

37.   Brazhnik, K., et al., *Single-cell analysis reveals different age-related somatic mutation profiles between stem and differentiated cells in human liver.* Sci Adv, 2020. **6**(5): p. eaax2659.

38.   Zhang, L., et al., *Single-cell whole-genome sequencing reveals the functional landscape of somatic mutations in B lymphocytes across the human lifespan.* Proc Natl Acad Sci U S A, 2019. **116**(18): p. 9014-9019.

39.   Cheng, C. and M. Kirkpatrick, *Molecular evolution and the decline of purifying selection with age.* Nat Commun, 2021. **12**(1): p. 2657.

40.   Yang, L., *A Practical Guide for Structural Variation Detection in the Human Genome.* Curr Protoc Hum Genet, 2020. **107**(1): p. e103.

41.   Ewels, P.A., et al., *The nf-core framework for community-curated bioinformatics pipelines.* Nat Biotechnol, 2020. **38**(3): p. 276-278.

42.   Hart, T., et al., *Finding the active genes in deep RNA-seq gene expression studies.* BMC Genomics, 2013. **14**: p. 778.

43.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

670 44. Wu, T., et al., *clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.*
671   Innovation (Camb), 2021. **2**(3): p. 100141.
672 45. Saul, D., et al., *A new gene set identifies senescent cells and predicts senescence-associated*
673   *pathways across tissues.* Nat Commun, 2022. **13**(1): p. 4827.
674 46. Kanehisa, M. and S. Goto, *KEGG: kyoto encyclopedia of genes and genomes.* Nucleic
675   Acids Res, 2000. **28**(1): p. 27-30.
676 47. Gundry, M., et al., *Direct, genome-wide assessment of DNA mutations in single cells.*
677   Nucleic Acids Res, 2012. **40**(5): p. 2032-40.
678 48. Petljak, M., et al., *Characterizing Mutational Signatures in Human Cancer Cell Lines*
679   *Reveals Episodic APOBEC Mutagenesis.* Cell, 2019. **176**(6): p. 1282-1294.e20.
680 49. Blokzijl, F., et al., *MutationalPatterns: comprehensive genome-wide analysis of*
681   *mutational processes.* Genome Med, 2018. **10**(1): p. 33.
682 50. Sondka, Z., et al., *COSMIC: a curated database of somatic variants and clinical data for*
683   *cancer.* Nucleic Acids Res, 2024. **52**(D1): p. D1210-d1217.
684 51. Degasperi, A., et al., *A practical framework and online tool for mutational signature*
685   *analyses show inter-tissue variation and driver dependencies.* Nat Cancer, 2020. **1**(2): p.
686   249-263.
687 52. Bergstrom, E.N., et al., *Generating realistic null hypothesis of cancer mutational*
688   *landscapes using SigProfilerSimulator.* BMC Bioinformatics, 2020. **21**(1): p. 438.
689 53. Selby, C.P., et al., *Molecular Mechanisms of Transcription-Coupled Repair.* Annu Rev
690   Biochem, 2023. **92**: p. 115-144.
691 54. Heilbrun, E.E., M. Merav, and S. Adar, *Exons and introns exhibit transcriptional strand*
692   *asymmetry of dinucleotide distribution, damage formation and DNA repair.* NAR Genom
693   Bioinform, 2021. **3**(1): p. lqab020.
694 55. Sherry, S.T., et al., *dbSNP: the NCBI database of genetic variation.* Nucleic Acids Res,
695   2001. **29**(1): p. 308-11.

Figure 1 | Repeated Mutagen Treatment Only Slightly Affects Cell Growth.
A) Schematic of experimental design. 1 million human primary lung fibroblasts (IMR-90) cells were plated at the beginning of each cycle and allowed to expand for 24 hours. Cells were then treated with 50 µg/mL N-ethyl-N-nitrosourea (ENU) and allowed to recover for 3 days. Cells were then collected and assayed for cell number and cell viability. Of the collected cells, 1 million cells were re-plated and then allowed another 4 days to recover and expand until they were collected to assess cell number and viability for the 7-day time point. This procedure was repeated 9 consecutive times over the course of ~10 weeks. Cells were isolated at the 7-day time point at cycles 1, 3, 6, and 9 for somatic mutation analysis and at cycles 1 and 9 for transcriptome analysis.
B) Population doublings (PD) of the control (CTRL) and ENU-treated (ENU) groups throughout the experiment. Statistics: n = 3, Data represent mean and S.D.; two-way repeated measures ANOVA. See figures S1A-B for results separated at the 3- and 7-day time points.
C-D) Percentage of cells detected in early (C) and late (D) apoptosis at the 3-day time point throughout the experiment. See figures S1C-D for results at 7-day time point. Statistics: n = 2, mean and S.D; Two-way repeated measures ANOVA.
E-F) Differential gene expression (DGE) of control cycle 9 vs control cycle 1 (E) and gene set enrichment analysis (GSEA) (F) to test for the effect of cell passaging. Red points in (E) are Log_2 fold change > 2 and adjusted p-value < 0.05. See figure S2L for GSEA results on DNA repair pathways. Statistics: n = 3; DGE results obtained using Wald test followed by Benjamini-Hochberg correction for multiple hypothesis testing; GSEA p-values obtained using a permutation test.
G-H) DGE of ENU-treated cycle 9 vs control cycle 9 (G) and GSEA (H) to test for the effect of ENU treatment. Red points in (G) are Log_2 fold change > 2 and adjusted p-value < 0.05. See figures S2M-N and supplemental note 1 for DGE results obtained using only cycle 9 samples. Statistics: n = 3; DGE results obtained using Wald test followed by Benjamini-Hochberg correction for multiple hypothesis testing; GSEA p-values obtained using a permutation test.

Figure 2 | Repeated Mutagen Treatment Results in Extremely High Mutation Burden.
A) Estimated burden of somatic single nucleotide variants (SNVs) per cell corrected for genome coverage and sensitivity. See figure S3H for observed SNVs per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; P-values of comparisons between cycles within the same condition were obtained with a two-sided linear model with estimated marginal means, Tukey's HSD. P-values of comparisons of control and ENU-treated groups are estimated using a two-way ANOVA and shown at top of plots.
B) Fit of linear model to estimated burden of SNVs per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; Shown at top is Pearson R correlation coefficient and p-value of the correlation that was obtained using a T-test.
C) Estimated burden of somatic short insertions/deletions and (INDELs) per cell corrected for genome coverage and sensitivity. See figure S3I for observed INDELs per cell. Statistics: n = 3-6; Data represent mean ± S.D.; p-values of comparisons between cycles within the same condition were obtained with a two-sided linear model with estimated marginal means, Tukey's HSD. P-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA and are shown at top of plots.
D) Ratio of estimated INDELs to SNVs per cell using data shown in Figures 2A and 2C. See figure S3L for observed INDELs to SNVs per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; P-values of comparisons between cycles within the same condition were obtained with a two-sided linear model with estimated marginal means, Tukey's HSD. P-values of comparisons of control and ENU-treated groups are estimated using a two-way ANOVA and shown at top of plots.
E) Mutation spectra of the relative contribution of SNV types per cell grouped by condition. See figure S5A for each group separately. Note that cells from the control cycle 1 group were excluded from this analysis due to abnormally high C>T frequency suggestive of a sample preparation artifact (Figures S1C-D; Methods). Statistics: n = 3-6; Data represent mean ± S.D.
F) Relative contribution of *de novo* mutational signatures SBSA and SBSB to individual cells (shown in rows) grouped by condition and cycle number. Each individual cell is shown as a row. Note that cells from the control cycle 1 group were excluded from this analysis (Methods). See figure S5B for why 2 signatures were chosen to be extracted.
G) Comparison of *de novo* mutational signature SBSA to most similar fitted signature SBS40 from the COSMIC database. Note that cells from the control cycle 1 group were excluded from this analysis (Methods). Statistics: cosine similarity=0.88, residual sum of squares=$4.03*10^{-3}$
H) Comparison of *de novo* mutational signature SBSB to most similar fitted signature SBS5 from the COSMIC database. Note that cells from the control cycle 1 group were excluded from this analysis (Methods). Statistics: cosine similarity=0.71, residual sum of squares=$1.1*10^{-2}$
I) Comparison of *de novo* mutational signature SBSA to most similar fitted signature SBS40 from the COSMIC database. Note that cells from the control cycle 1 group were excluded from this analysis (Methods). Statistics: cosine similarity 0.86, residual sum of squares=$7.54*10^{-3}$
P-value legend: ns: p > 0.05; ***: p <= 0.001; ****: p <= 0.0001

Figure 3 | Negative Selection Against Damaging Coding and Non-Coding Variants.
A) Observed frequency of synonymous (left panel) and non-synonymous (right panel) variants per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; p-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA.
B) Log_2 observed vs expected (O/E) ratio of synonymous and non-synonymous variants grouped by cycle-condition. O/E ratios were normalized for preferential DNA repair by the O/E ratio of variants in intronic regions (Figure 3G, bottom panel; Methods). See figure S6A for results without normalization. See figure S6B for results per cell. Statistics: n = 3-6; Data represent mean ± S.D.; p-values for comparing the frequency of observed variants to that expected by chance alone (Log_2 O/E = 0) were estimated using a permutation test (n = 10,000).
C) Percent of non-synonymous variants out of total coding variants that are protected against in the ENU-treated group due to negative selection (Methods). Statistics: n = 3-6; Data represent mean ± S.D.
D) Top panel: Observed frequency of specific coding variants per cell. O/E ratios were normalized by the O/E ratio of variants in intronic regions (Figure G, bottom panel; Methods). See figure S6C for results per cell. Statistics: n = 3-6; Data represent mean ± S.D.; p-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA. Bottom panel: Log_2 O/E of specific coding variants grouped by cycle-condition. Statistics: n = 3-6 cells; Data represent mean ± S.D.; p-values for comparing the frequency of observed variants to that expected by chance alone (Log_2 O/E = 0) were estimated using a permutation test (n = 10,000).
E) Ratio of observed non-synonymous to synonymous (N/S) variants per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; p-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA.
F) Log_2 O/E of the N/S ratio grouped by cycle-condition. See figure S6D for results per cell. Statistics: n = 3-6 cells; Data represent mean ± S.D.; p-values for comparing the frequency of observed variants to that expected by chance alone (Log_2 O/E = 0) were estimated using a permutation test (n = 10,000).
G) Same as in D, but for specific non-coding variants. O/E ratios only of splice region variants were normalized by the O/E ratio of variants in intronic regions (Figure G, bottom panel; Methods). See figure S6F for results per cell. See supplementary note 4 for discussion on intergenic region O/E results.
H) Reduction of variants in 3' untranslated region (UTR) and splice region variants that are protected against in the ENU-treated group due to negative selection (Methods). Statistics: n = 3-6; Data represent mean ± S.D.
P-value legend: ns: p > 0.05; *: p <= 0.05; **: p <= 0.01; ***: p <= 0.001; ****: p <= 0.0001

Figure 4 | Protection Against ENU-induced Mutation Accumulation in Functional Pathways.
A) Observed frequency of SNVs per cell in functional regions associated with genes that were determined to be non-expressed or expressed from bulk mRNA transcriptome data from control and ENU-treated cells at cycle 9. Statistics: n = 3-6 cells; Data represent mean ± S.D.; p-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA.
B) Log_2 observed vs expected (O/E) ratio of SNVs grouped by cycle-condition in functional regions associated with genes determined to be non-expressed or expressed from bulk mRNA transcriptome data from control and ENU-treated cells at cycle 9. O/E ratios were normalized by the O/E ratio of variants in intronic regions (Figure 3G, bottom panel; Methods). See figure S7C for results without intron O/E normalization. See figure S7D for results per cell. Statistics: n = 3-6 cells; data represent mean ± S.D.; p-values for comparing the frequency of observed variants to that expected by chance alone (Log_2 O/E = 0) were estimated using a permutation test (n = 10,000).
C) Top 20 gene ontology biological processes resulting from gene set over representation analysis of expressed genes from bulk mRNA transcriptome data from control and ENU-treated cells at cycle 9. See figure S2A-I for detailed analysis. Statistics: n = 6 cells; Adjusted p-values obtained using a hypergeometric distribution test followed by Benjamini-Hochberg correction for multiple hypothesis testing.
D) Mean observed frequency of SNVs grouped by cycle-condition within the functional regions of expressed pathways. Statistics: n = 3-6 cells; p-values of comparisons of control and ENU-treated groups were estimated using a two-way ANOVA.
E) Log_2 O/E ratio of SNVs grouped by cycle-condition within the functional regions of expressed pathways. O/E ratios were normalized by the O/E ratio of variants in intronic regions (Figure 3G, bottom panel; Methods). See figure S7E for results without intron O/E normalization. See figure S7F for results plotted as volcano. Statistics: n = 3-6 cells; p-values for comparing the frequency of observed variants to that expected by chance alone (Log_2 O/E = 0) for each pathway are highlighted in red and were estimated using a permutation test (n = 10,000) followed by Benjamini-Hochberg correction for multiple hypothesis testing.
F) Reduction of SNVs in functional regions of pathways found to be under significant negative selection at cycle 9 in the ENU-treated group. This was calculated by subtracting the frequency of variants in the observed by the average frequency of variants in the simulated samples (Methods). Statistics: n = 3-6; Data represent mean ± S.D.
P-value legend: ****: p <= 0.0001

Figure 1

Figure 2

Figure 3

Figure 4