Population size rescaling significantly biases outcomes of forward-in-time population genetic simulations

Amjad Dabi and Daniel Schrider[*]

Department of Genetics, University of North Carolina, Chapel Hill, North Carolina, USA

Address: 120 Mason Farm Rd, Chapel Hill, NC 27599

[*]Corresponding author; e-mail: drs@unc.edu

ABSTRACT

Simulations are an essential tool in all areas of population genetic research, used in tasks such as the validation of theoretical analysis and the study of complex evolutionary models. Forward-in-time simulations are especially flexible, allowing for various types of natural selection, complex genetic architectures, and non-Wright-Fisher dynamics. However, their intense computational requirements can be prohibitive to simulating large populations and genomes. A popular method to alleviate this burden is to scale down the population size by some scaling factor while scaling up the mutation rate, selection coefficients, and recombination rate by the same factor. However, this rescaling approach may in some cases bias simulation results. To investigate the manner and degree to which rescaling impacts simulation outcomes, we carried out simulations with different demographic histories and distributions of fitness effects using several values of the rescaling factor, Q, and compared the deviation of key outcomes (fixation times, fixation probabilities, allele frequencies, and linkage disequilibrium) between the scaled and unscaled simulations. Our results indicate that scaling introduces substantial biases to each of these measured outcomes, even at small values of Q. Moreover, the nature of these effects depends on the evolutionary model and scaling factor being examined. While increasing the scaling factor tends to increase the observed biases, this relationship is not always straightforward, thus it may be difficult to know the impact of scaling on simulation outcomes *a priori*. However, it appears that for most models, only a small number of replicates was needed to accurately quantify the bias produced by rescaling for a given Q. In summary, while rescaling forward-in-time simulations may be necessary in many cases, researchers should be aware of the rescaling effect's impact on simulation outcomes and consider investigating its magnitude in smaller scale simulations of the desired model(s) before selecting an appropriate value of Q.

AUTHOR SUMMARY

Computer simulations have become an indispensable tool for researchers studying the dynamics of evolution in nature, since the underlying processes are often complex and cannot always be described using purely theoretical approaches. Unfortunately, these simulations often require large computing resources and long run times. To overcome this, researchers often employ an approximation which allows them to run the desired simulation faster by decreasing the size of the simulated population by a factor of

their choosing while increasing other parameters of the simulation by the same factor to compensate. While it is generally believed that such an approximation should not greatly affect the outcomes of the simulation, this has only been rigorously tested in a small set of scenarios. Our study tests this rescaling approximation across a larger set of scenarios by simulating each scenario with a full population size, using the approximation to simulate the same scenario with smaller populations, and investigating whether this affects important simulation outcomes. Our results indicate that the rescaling approximation does introduce significant biases in our measured outcomes which generally increase with further downscaling of the population size, and that caution must be taken when it is employed to decrease the computation burden of simulations.

INTRODUCTION

Stochastic simulations are widely used in population genetics [1–3]. Their uses include the study of the behavior of a wide variety of population genetic models, either when used on their own in simulation studies [4–9] or when used to validate theoretical predictions [10–15], the comparison of simulations to data to infer model parameters or choose between competing models [16–20], assessing the accuracy of inference methods [21–23] and the generation of training datasets for machine learning methods [24–34]. Given their wide range of research applications, considerable effort has been made to improve the accessibility, computational speed, and flexibility of population genetic simulations [35–40].

Two main modes of population genetic simulation are available to researchers: coalescent simulations, which trace the genealogy of a sample of individuals back in time until all segments of the chromosome have reached the sample's MRCA, and forward-in-time simulations, which explicitly simulate an entire population of chromosomes evolving under the desired model. Coalescent simulations have been a mainstay of population genetics for the last few decades because of their computational efficiency [41–43], as they are often used to simulate only a relatively small sample of genomes rather than the entire population [44], and jump backward from one event (e.g. coalescence and recombination) rather than simulating each generation of the evolutionary scenario.

While computationally efficient, coalescent simulations are relatively limited in the evolutionary scenarios that they can model, especially with respect to natural selection, and are therefore more commonly used for modeling demographic histories (see Ewing and Hermisson [36]; Kern and Schrider [37]; Baumdicker et al. [45]; Teshima and Innan [46] for notable exceptions). Forward-in-time simulations, on the other hand, provide much greater flexibility [47–51], giving users the ability to simulate more complex models involving not only demographic changes, but several types of natural selection, and scenarios that fall outside the bounds of the Wright-Fisher model [52]. Unfortunately, forward-in-time simulations can require a large amount of computational power and memory because the entire population is simulated for each generation of the desired scenario, rendering simulations of large populations and/or very long histories infeasible. One popular approach to alleviate this computational burden is to rescale forward-in-time simulations such that a smaller population is used to capture the desired evolutionary dynamics that would be expected for the larger population [53]. Typically, this is done by scaling down the population by some factor (Q) while scaling up the mutation and recombination rates by the same factor. Theoretically, this scaling approximation would preserve important population-level measures and allow easy recovery of the times at which events occurred during the simulation. For instance, the population-scaled mutation rate, θ, for a population of size N and with a mutation rate of μ,

is equal to $4N\mu$. After rescaling, our new population size is N/Q and our new mutation rate is $\mu \times Q$, and thus θ is unchanged:

$$\theta_{scaled} = \frac{4N}{Q}\mu Q = 4N\mu$$

Meanwhile, under the standard neutral model, the expected time to the most recent common ancestor (TMRCA) between any two individuals in the unscaled simulation is $2N$ generations. Because the population size is decreased by a factor of Q in the scaled simulations, the expected TMRCA becomes $2N/Q$. Consequently, to produce genealogies of the desired shape and size, the evolutionary model being simulated must be modified such that the time intervals between events specified in the simulated model (e.g. population splits, migration events, etc.) are divided by Q. Following the simulation, if one wishes to investigate the timescale of any stochastic events occurring during the simulation such as fixations, coalescent events between particular pairs of individuals, etc., the times of these events can simply be multiplied by Q to produce results that approximate those expected under the original unscaled model.

This population-size "rescaling trick" has been used in forward simulations for decades, with its first use dating all the way back to Hill and Robertson [54]). However, the consequences of rescaling have rarely been evaluated, and with the exception of Hill and Robertson's initial comparison of the effects of changing N from 16 to 8, those few studies that have examined the impact of rescaling have noticed a significant effect. For example, although the approximations underlying rescaling appear to be sound at least for simulations lacking natural selection, Adrion et al. [2] found that in some cases, rescaling by modest values of Q can bias the outcomes of purely neutral simulations, resulting in downwardly biased estimates of nucleotide diversity (π [55], Tajima's D [56], and linkage disequilibrium (LD) as measured by r^2). Moreover, when natural selection is included, rescaling may introduce further biases. The selection coefficient is often rescaled by multiplying it by Q, such that Ns, a commonly used measure of the efficacy of selection, is unchanged [12,53,57]. However, the time to fixation of a selected mutation does not scale linearly with N. Indeed, Uricchio and Hernandez [53] showed that, under a recurrent hitchhiking model, rescaling blunts the impact of selection on diversity levels and the site frequency spectrum. Furthermore, Comeron and Kreitman [58] showed that rescaling alters the dynamics of interference selection for weakly selected mutations, affecting important measures such as Tajima's D. Although these studies have examined a very limited set of scenarios, their findings suggest that undesirable effects of rescaling could be commonplace. This possibility, combined with the increasing importance and accessibility of forward simulations, underscores the need to investigate the impact of rescaling across a larger array of simulated models. We still know little about the manner in which simulations are affected (i.e. which simulation outcomes are upwardly or downwardly biased by rescaling), how widespread these effects are across simulated models, and how the severity of these effects varies with Q.

In this study, we carry out both scaled and unscaled simulations under a variety of demographic and selective scenarios and investigate the impact of rescaling on important simulation outcomes such as fixation times, site frequency spectra, and linkage disequilibrium. Our simulations explore various selection models containing different combinations of neutral, deleterious, and beneficial mutations, including simulations with a distribution of fitness effects estimated from human population genomic data. We also examine scaling effects in constant, expanding, and contracting populations. To quantify the effect of scaling, we train classification models that use a set of simulation outcomes as features to determine whether the results of scaled and unscaled simulations are distinguishable across increasing values of Q. Furthermore, for each feature we characterize the direction and magnitude of the deviation

between scaled and unscaled simulations to provide a detailed picture of the effects of scaling across our scenarios. Our results show that rescaling simulations can result in a substantial deviation of numerous simulation outcomes, and that the magnitude of these effects usually increases with the value of Q. We also find that the choice of demographic model can have an impact on the severity of the rescaling effect. Overall, the results indicate the need to exercise caution in making inferences from rescaled simulations, especially when large values of Q are used.

METHODS

Simulation and rescaling strategy

We used `SLiM 4.0` [35] and `stdpopsim 0.2.0` [2,59] to carry out our scaled and unscaled simulations, all of which conform to Wright-Fisher assumptions: generations are non-overlapping, mating is random, and all individuals are monoecious, and the population size is solely determined by the specified demographic model and is not affected by the population's fitness. For our main simulations, the mutation and recombination rates are both $Q \times 1.2 \times 10^{-8}$, the genome size is 25 Mb, and the scaled population size was set to N/Q, with $N = 10,000$. For mutations with a constant population size, a burn-in was performed for a duration of $10 \times N \times Q$ generations, after which statistics were recorded during and after a simulation run of $4 \times N \times Q$ generations. We also performed simulations of the full model in the presence of a single-event population expansion or contraction. In the expanding simulation, the unscaled population size expands from $N_{anc} = 10,000/Q$ individuals to $N_0 = 20,000/Q$ individuals after the burn-in, while in the contraction simulations the size decreases from $N_{anc} = 10,000/Q$ individuals to $N_0 = 5,000/Q$ individuals after the burn-in. In both demographic scenarios, the burn-in runs for $10 \times N_{anc}$ generations, followed by N_0 generations after the population size change. For example, an unscaled simulation under the expansion scenario would run for 100,000 generations of burn-in followed by 20,000 generations, during and after which statistics summarizing the simulation results are recorded. For each simulated scenario, we generated 1000 replicates for each scaling factor (Q) except for the simulations of larger populations described below, where the computational demand necessitated a smaller sample of 100 replicates.

The fitness effects of mutations in our simulations were as specified in Table 1, and used the gamma distribution of deleterious fitness effects estimated from African American genomes by Boyko et al. [60]. Using this general structure, we performed scaled and unscaled simulations of several different scenarios each with different mixes of neutral, deleterious, and beneficial mutations (Table 2). Note that the parameter values listed in the second column of Table 2 specify the probability that any new mutation will be of a given mutation type as follows: f_n is the probability that a new mutation will be neutral, f_b is the probability it will be beneficial, and f_d is the probability it will be deleterious.

Table 1: Mutation parameters for simulations.

Mutation Type	Fitness Effect (s)
Neutral	0
Beneficial	$0.01 \times Q$
Deleterious	$\sim Gamma(-0.0294 \times Q, 0.184)$

Table 2: Proportions of neutral, deleterious, and beneficial mutations in each simulation model.

Simulation Model Name	Model parameter values
Full model	$f_n = 0.25, f_b = 0.75 \times 0.05, f_d = 0.75 \times 0.95$
No-beneficials model	$f_n = 0.25, f_d = 0.75$
Sparse model	$f_n = 0.95, f_d = 0.05$
Neutral-only model	$f_n = 1$

We also tested the effect of scaling in several additional models as well. To test the effect of starting with a larger unscaled population size, for example, we simulated the no-beneficials model with a constant unscaled population size $N = 20{,}000$. The no-beneficials model was selected since it exhibited the most dramatic effects due to scaling. We also tested a single sweep model with a population size of $100{,}000/Q$, where a single beneficial mutation ($s{=}0.05 \times Q$) was introduced in the middle of the genome and then swept to fixation, whereupon the simulation was terminated. If the sweeping mutation was lost, the simulation was reset and run again until fixation occurred. Finally, we tested two *Drosophila melanogaster* constant population size models based on the `stdpopsim` catalog parameters for the species [2] which specifies a population size of $1{,}720{,}600/Q$ [61], a mutation rate of $5.49 \times 10^{-9} \times Q$ [62], and a recombination rate of $2.40463 \times 10^{-8} \times Q$ (based on the estimate from Comeron et. al [63]). For these simulations, we chose a 10 kb genome size and only conducted simulations at Q values of 20, 50, 100, 150, 200 due to computational limitations. For the first model, we used the distribution of fitness effects from Ragsdale et al. [64]. Since this DFE is expressed in units of $2N_e$ `stdpopsim` converts the selection coefficients to units of s using a population size estimate of 2.8×10^6 [65]. This yielded a lognormal distribution with a mean of $9.679 \times 10^{-7} + \log(Q)$ and a standard deviation of 3.36 for the selection coefficient of deleterious mutations, and a fixed value of $7.125 \times 10^{-6} \times Q$ for the selection coefficient of beneficial mutations. The second *Drosophila* model was identical to the first except that the selection coefficient for beneficial mutations was increased to $0.01 \times Q$. All code for performing the simulations and downstream analyses in this study are available at https://github.com/SchriderLab/simscale-snakemake.

Statistics summarizing simulation outcomes

For each simulation replicate, we recorded the following statistics summarizing the outcome:

- The site frequency spectrum (SFS) for each mutation type, each estimated from the same sample of 100 randomly selected individuals taken at the end of the simulation.
- Linkage disequilibrium (LD), as measured by the r^2 statistic calculated from the same sample of 100 individuals for all pairs of mutation or a set of 5000 randomly chosen mutations (whichever was smaller). These r^2 values were then binned by distance to produce 50 equal distance bins across the genome and the average r^2 value in each bin was recorded.
- The fixation probability for each mutation type, recorded only for mutations occurring after the burn-in. These probabilities were estimated by taking the fraction of mutations of a given type that reached fixation and rescaled by dividing this fraction by Q.
- The distribution of fixation times, recorded separately for each mutation type and only for mutations occurring after the burn-in. These times were all rescaled by multiplying by Q and binned to produce around 20-30 bins for each simulation model.

Assessing the deviation between scaled and unscaled simulations

To assess whether scaled and unscaled simulations could be distinguished from one another for a given simulation model and a given value of Q, we trained classifiers to label a given simulation as unscaled or scaled, with the input features being the SFS, the binned data for fixation times and LD, and the fixation probabilities as described above. 80% of replicates for a given model and value of Q were used to train the classifier, and the remaining 20% of the replicates were set aside for testing. For each dataset, we trained both a logistic regression model and a random forest model. Model parameters were the default values of the Python package scikit-learn, version 1.1.3 [66]. The input features were grouped into the following four separate feature sets to separately test their ability to distinguish between scaled and unscaled simulations: 1) the SFS across all mutation types (199 features per mutation type) represented as a 1-dimensional vector per replicate, 2) the average r^2 values at different (binned) inter-mutation distances (50 features, all mutation types), 3) the fixation probabilities (one feature per mutation type), and 4) the histograms of fixation times (20-30 features per mutation type) concatenated along the time axis to produce a 1-dimensinoal vector per replicate. The varying number of bins was due to our goal of keeping the width of the fixation time bins fixed across all replicates, such that they map correctly to each other. Therefore, a random unscaled replicate was chosen to decide the width of the fixation time bin (in generations) to produce 20 bins, and then propagated to all replicates across all scaling factors. For instance, if the maximum fixation time for the randomly chosen unscaled replicated was 10000 generations, then the bin width would correspond to 500 generations. Since some replicates will have a different maximum fixation time, the number of bins would vary across mutation types and simulation scenarios, but each bin would correspond to the same fixation time range in all replicates and for all scaling factors. After training, classification accuracy (fraction of correct predictions on the test data) was recorded.

To further characterize the differences between scaled and unscaled simulations, the Kullback-Leibler (KL) Divergence between the scaled and unscaled fixation times and site frequency spectra were calculated. For these calculations, the distributions for each outcome were first averaged for each bin across all replicates for each scaling factor such that:

$$y_{bin} = \frac{1}{n_{replicates}} \Sigma_{n=i}^{n_{replicates}} y_{bin_i}$$

KL Divergence of the scaled distribution for each outcome was then calculated against the corresponding unscaled distribution. For LD, we opted to use the RMSE instead of KL Divergence since we summarized LD by the average r^2 values within genomic distance bins rather than probability distributions of r^2 values. This was obtained by averaging the squared error values for each bin in the LD distribution for each scaling factor such that:

$$RMSE_{LD} = \sqrt{\frac{\Sigma_{bin=1}^{50} \left(r^2_{bin_{scaled}} - r^2_{bin_{unscaled}} \right)^2}{50}}$$

Additionally, we calculated the mean percent error (MPE) in between the scaled and unscaled fixation times, mutation frequency, LD r^2 values, and fixation probabilities by first obtaining the mean of each outcome for each replicate, averaging the means across all replicates, and calculating the MPE as:

$$MPE = 100 \times \frac{\overline{\mu_{unscaled}} - \overline{\mu_{scaled}}}{\overline{u_{scaled}}}$$

Where μ is the mean of the outcome for each replicate.

Assessing the impact of sample size on estimates of deviation between scaled and unscaled simulations

To investigate the impact of the number of replicates on the estimation of deviation due to scaling, we recalculated our MPEs, KLD, and LD average RMSE values for our simulation outcomes between scaled and unscaled simulations using smaller sample sizes of 100 and 10 for the full model, the no-beneficials model, and the *Drosophila* model with stronger selection on beneficials. For each of those deviation measures, we sampled the appropriate number of replicates randomly for each scaling factor and recalculated our estimates as described above.

RESULTS

Rescaling alters the SFS, LD, and fixation times/probabilities in a simulated population with a gene-dense genome

To characterize the impact of population size rescaling on simulations, we simulated a variety of scenarios and examined how rescaling alters allele frequencies, LD, and the probability of and time to fixation (Methods). We began by examining simulations of a constant-sized population with a genome densely packed with selected mutations (both deleterious and beneficial, with 5% of selected mutations belonging to the latter category); because this is our richest model in terms the amount of natural selection, we refer to it as the "full model" (Methods).

When examining the distribution of mutation fixation times for neutral mutations under the full model (Fig 1A), the differences between the unscaled and scaled distributions are subtle and non-monotonic. For instance, fixation times at Q of 2 are slightly shifted towards longer sojourns, while those at Q of 20 are shifted in the opposite direction. The distributions of fixation times for beneficial and deleterious mutations (S1 Fig) also show this non-monotonic trend, with longer sojourns at Q of 2, 5 and 10 and shorter sojourns at Q of 20. The site frequency spectrum for neutral mutations (Fig 1B), again shows a non-monotonic trend across values of Q. The fraction of mutations that are singletons, for which differences are most pronounced, decreases with increasing values of Q and slightly increases at Q of 20 compared to Q of 10. The SFS for deleterious mutations follows a very similar pattern to neutral mutations, while for beneficial mutations the fraction of singletons increases more dramatically at Q=20 (S1 Fig). The amount of LD across the genome markedly changes with Q (Fig 1C) as it appears that rescaling increases LD between mutations regardless of the distance between them. Furthermore, the fixation probabilities for all mutation types also change with Q (Fig 1D), with the largest shift occurring for beneficial mutations; we note that this comparison was done after rescaling all fixation probability estimates themselves to account for the impact of rescaling on the neutral fixation probability (Methods). For example, for beneficial mutations, the fixation probability was around 0.20% for unscaled simulations and gradually decreased to around 0.05% for $Q = 20$. On the other hand, the fixation probabilities for neutral and deleterious mutations remain steady at ~ 0.005% and 0.003%, respectively, for all values of Q.

Fig 1. Select simulation results for the full model across scaling factors. (**A**) Fixation times for neutral mutations. (**B**) The site frequency spectra for neutral mutations. (**C**) Linkage disequilibrium across 50 genomic bins for all mutations. (**D**) Fixation probabilities for all mutations. As higher values of Q are used, all statistics show larger deviations from values observed in unscaled simulations. Fixation probabilities for beneficial mutations are impacted markedly more than for other mutation types.

Next, we trained classifiers to determine whether scaled and unscaled simulations were distinguishable from one another based on these simulation outcomes. To assess this, we trained a set of binary classifiers, each using a different set of input features (fixation times, the SFS, fixation probabilities, and LD) to discriminate between unscaled simulations at a given value of Q. As Q increases, classification accuracy increases for the SFS- and LD-based classifiers, starting at 67% for the SFS at $Q = 2$ and increasing to 100% at $Q = 10$ for the logistic regression classifier. For LD, the accuracy of logistic regression increases from 61% at $Q = 2$ to 99.5% at $Q = 10$. For fixation times and fixation probabilities, accuracy is already at 100% when $Q=2$ (Fig 2A), consistent with a larger deviation between scaled and unscaled simulation outcomes at higher values of Q. Similar results were obtained by the random forest classifiers (Fig 2B). Most importantly, we found that all feature sets could be used to distinguish between scaled and unscaled simulation replicates, even at relatively modest values of Q. For example, the minimum accuracy observed at $Q=5$ was 89% for the random forest classifier and 88% for the logistic regression classifier (both examining LD). The high accuracy obtained by classifiers discriminating between unscaled and rescaled simulations demonstrates that the rescaling effects summarized in (Fig 1) are significant and systematic.

Fig 2. Classifier accuracy for models trained to distinguish between scaled and unscaled replicates of the full model. **(A)** Accuracy of the Random Forest classifier trained to classify a given simulation replicate as scaled or unscaled. **(B)** Accuracy of the Logistic Regression classifier trained to classify a given simulation replicate as scaled or unscaled. A separate classifier was trained for each combination of Q value and set of features (those summarizing the SFS, fixation probabilities, LD, and the distribution of fixation times). Accuracy represents fraction of simulation replicates in the test dataset correctly identified as scaled or unscaled.

Having established that rescaling significantly influences simulation outcomes, we sought to quantify the magnitude of this effect. We first examined the mean percent deviation between scaled and unscaled simulations (hereafter referred to as mean percent error, or MPE) for each feature examined above (Fig 3). MPE for fixation time and allele frequency shows a non-monotonic rescaling effect across values of Q. For fixation times, MPE for all mutation types increases between Q values of 2 and 5, but decreases with Q=10, and then flips sign at Q=20 where it exhibits the largest magnitude across scaling factors. The decrease in error between Q=5 and Q=10 occurs due to an initial increase in the mean fixation time as Q increases. For example, in the case of neutral mutations the mean fixation time increases from 622 generations at Q=1 to 629 and 637 generations at Q of 2 and 5, respectively. This is followed by a decrease in mean fixation time to 634 and 600 generations at Q of 10 and 20, respectively. This causes the mean fixation time to be closer to that of the unscaled simulation at Q=10 compared to Q=5, before it decreases below the unscaled simulation mean at Q=20. When comparing the different mutation types within each Q, fixation time MPEs appear to be the lowest for beneficial mutations across all values of Q while neutral and deleterious mutations both exhibit slightly larger MPEs that are similar to one another. For allele frequencies, we observed a considerably higher MPE, which increases monotonically in magnitude from Q=2 to Q=20 for all mutation types. Here, neutral mutations have the highest MPE across Q values, reaching ~44% for Q=20 while beneficial mutations have the lowest error at −6.2% for Q=20 and are the only type of mutation to experience a sign change in MPE (occurring at Q=20). MPE was also large for LD and for the fixation probability of beneficial mutations, and again

increased in magnitude with Q. However, deleterious and neutral mutations exhibit miniscule MPEs for their fixation probabilities.

Fig 3. Mean percentage error of various statistics for the full model by values of Q. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage error for average linkage disequilibrium as measured by the values of r^2. For each statistic, the mean value was calculated for each replicate, followed by averaging the means across all replicates in each value of Q. The mean percentage error was then calculated between the resulting value from each Q and the corresponding value at Q of 1.

As an alternative way of quantifying the degree of deviation between scaled and unscaled simulations, we examined KL divergence for fixation times and SFS distributions and RMSE for LD (S2 Fig). For the distribution of fixation times, all mutation types show similar KL divergence within each scaling factor, with KL divergence increasing steadily for each mutation type as Q increases. This trend changes at $Q = 20$, where beneficial mutations display substantially higher KL divergence than other mutation types. Similarly, when examining the SFS, KL divergence for each mutation type increases with Q, although in this case, while beneficial mutations again have the highest KL divergence at $Q=20$, neutral mutations have markedly higher KL divergence at $Q=10$. For LD decay, the root mean squared

error (RMSE) between scaled and unscaled simulations a similar trend as MPE, increasing with Q from around 0.00006 at $Q=2$ to around 0.02 at $Q=20$.

Rescaling effects persist in simulations with deleterious mutations only

Our full model serves as a useful stress test of the robustness of forward simulations to rescaling but contains a larger number of beneficial mutations than will be realistic in most cases (5% of non-neutral mutations). To see how rescaled simulations would behave without such a pervasive impact of positive selection, we altered this model by removing all beneficial mutations (the "no-beneficials" model). Under this model, we again observe a significant scaling effect, as our classifiers were able to discriminate between scaled and unscaled simulations with higher accuracy as obtained for the full model (S4 Fig). One notable exception to this was the random forest using fixation probabilities at Q of 2, where accuracy was ~50%, no better than random guessing.

We next examined the magnitude of the rescaling effect for the no-beneficials model as measured by MPE. We observed much larger MPEs for fixation times for the no-beneficials model (Fig 4) than for the full model for fixation times, LD, and fixation probabilities. However, we observed smaller MPEs for allele frequencies than under the full model (MPE is approximately 5% at $Q=10$ compared to 28% for neutral mutations in the full model and increase to over 20% at $Q=20$ compared to 44% for neutral mutations in the full model), with both mutation types having similar errors across values of Q. Of note here is that unlike our full model, the magnitude of error increases monotonically and in the same direction for all measured statistics, with fixation times and allele frequencies exhibiting negative MPE. This indicates that rescaling in this model lowers average fixation times and allele frequencies, rather than increasing them as was the case for most values of Q under the full model. LD and fixation probabilities, on the other hand, exhibit positive MPE, with the MPE for LD increasing with Q in a similar manner to the full model, with slightly smaller values at all values of Q except 20, where the error is around 37% compared to the full model's 33%. The MPE for fixation probabilities was quite high, especially for deleterious mutations, with values peaking at around 142% for deleterious mutations and 64% for neutral mutations at $Q=20$. Interestingly, under the full model, the corresponding MPEs for these mutation types were negligible, as that model instead showed large MPEs for beneficial mutations, suggesting a substantial change in the dynamics of rescaling effects depending on the presence or absence of large numbers of beneficial mutations.

KL divergence for the fixation time distribution in the no-beneficials simulations (S4 Fig) again increases monotonically with Q, with similar values for both mutation types, but shows a much greater magnitude than observed for the full model across Q values. For the SFS, KL divergence is lower than that of the full model for lower values of Q, but suddenly increases to higher values than observed under the full model at $Q=20$. Consistent with the full model, however, neutral mutations show larger KL divergence than deleterious ones. For LD, KL divergence increases monotonically with Q and exhibits higher values at all values of Q compared to the full model. Overall, the comparison between the full model and no-beneficials model suggests that the presence of large amounts of positive selection may, at least in some cases, actually diminish the impact of rescaling on simulation outcomes.

Fig 4. Mean percentage error of various statistics for the model with deleterious mutations only by values of Q. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for fixation probabilities. **(D)** Mean percentage error for linkage disequilibrium as measured by the values of r^2.

Given the large scaling effects present in this model, we sought to test the impact of increasing the population size on these effects. To this end, we carried out simulations of this same model containing only deleterious and neutral mutations but with double the starting population size (N=20000). Results indicate that MPE for fixation times (Fig 5A) and fixation probabilities (Fig 5C) are nearly identical to their counterparts in a smaller population (compare to Fig 4). MPE for LD and the SFS are slightly diminished (Fig 5D and 5B), on the other hand, compared to their counterparts in the smaller starting population model. We note that this change is not uniform across all values of Q. For instance, increasing the unscaled population size decreases the MPE for allele frequency by around ~3% at Q=20 for both neutral and deleterious mutations, but only by ~1% at Q=10. For LD, increasing the population size reduces the MPE by ~10% at Q=20 and by ~5% for Q=10.

Fig 5. Mean percentage error of various statistics for the model with deleterious mutations only but a larger starting population by values of Q. (A) Mean percentage error for average mutation fixation times. (B) Mean percentage error for average allele frequencies. (C) Mean percentage error for average fixation probabilities. (D) Mean percentage error for linkage disequilibrium as measured by the values of r^2.

Sparse genome model

Next, we examined a model with a much smaller fraction of deleterious mutations (5% of all mutations), as might be expected in a large genome where functional elements are sparse. As in the previous section, we did not include beneficial mutations. Under this model, our classifiers were not as effective at discriminating between unscaled and rescaled simulations at lower values of Q (S8 Fig), especially for fixation times and fixation probabilities where the random forest model accuracy is approximately 53% for both features at Q=2 and 76% and 69% respectively at Q=5. However, we note that at Q=20 our classifiers were still quite accurate, reaching 100% on all feature sets for both classifiers. Thus, while simulation outcomes are not as impacted at lower Q values compared to the full model or a denser genome lacking beneficial mutations, they still appear to be significantly affected by rescaling under our sparse genome model, particularly at higher Q values.

Concordant with our decreased classifier accuracy, the MPE of fixation times for this model is lower across all statistics than that of the dense model without beneficial mutations (Fig 5). Again, the MPE for allele frequencies, fixation times and probabilities, and LD increases monotonically with Q. However, one notable difference is that the direction of the bias caused by rescaling, as indicated by the sign of the MPE, is reversed for allele frequencies compared to the dense model with deleterious mutations only—here we typically observe a slight increase in the average frequencies of neutral and deleterious mutations, as was observed under the full model including beneficial mutations rather than the dense genome model with deleterious mutations only. However, we again stress that the magnitude of this bias is much smaller under this sparse genome model than either of the two dense genome models (MPE <1% for allele frequencies at Q=20). The MPE is still appreciable for the other simulation outcomes at Q=20, however (~8% for fixation times, ~17.5% for LD, and ~35% for fixation probabilities).

Fig 6. Mean percentage error of various statistics for the sparse genome model by values of Q. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage error for average linkage disequilibrium as measured by the values of r^2.

KL Divergence for fixation times and the SFS (S8 Fig) reveals that for simulations under the sparse model, the distributions of outcomes for deleterious mutations are generally more impacted by rescaling than are neutral mutations, except for fixation times at $Q=20$ where the divergence is similar for both mutation types. When compared with the dense-deleterious model and, consistent with the smaller MPE values observed in this sparse model, KL Divergence for fixation times and the SFS are also smaller under the sparse model, with KL divergence for fixation times declining from ~14 for the dense-deleterious model for both mutation times at Q of 20 to ~0.04 at this Q for the sparse model. SFS KL divergence for neutral and deleterious mutations is around 0.0003 and 0.0005 respectively at Q of 20, compared to 0.046 and 0.034 for the corresponding values in the dense-deleterious model. Also consistent with the MPE trends for the sparse model is the observation that KL divergence for fixation times is similar for both mutation types and is higher for deleterious mutations in the case of the SFS. RMSE for LD is also lower compared to the dense-deleterious model, measured at 0.001 at Q of 20 compared to the corresponding dense-deleterious value of 0.002.

Fig 7. Mean percentage error of various statistics for the strictly neutral model by values of Q. (A) Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage error for average linkage disequilibrium as measured by the values of r^2.

Strictly neutral model

Next, to investigate the degree to which natural selection (whether linked or direct) is responsible for the rescaling effects described above, we performed simulations identical to the previous models but with neutral mutations only. In this case, classification of simulations as scaled or unscaled becomes more challenging for both models (S10 Fig) and fixation times and probabilities are uninformative features for all values of Q. However, some impacts of scaling on the SFS and LD appear to persist, as the random forest model achieves an accuracy of approximately 90% and 100% on those features, respectively, at Q=20. This indicates that rescaling can still impact allele frequencies and LD even in the absence of selection, as previously shown by [2]. These results are corroborated by the MPE values for these statistics (Fig 6) which show very small values for fixation times and probabilities but larger values for allele frequencies and LD. For LD, the MPE at Q=20 is around 16% under the strictly neutral model, which is comparable to that of the full model at Q=10, while the allele frequency MPE at Q=20 is around 1.1% under the strictly neutral model which is smaller than the MPE for neutral allele frequencies under the full model (the MPE for the neutral model at Q=20 is comparable to that of the full model at Q=2). Fixation times and SFS KL divergence and LD RMSE values for this model (S10 Fig) are also smaller than for the full model, but the LD RMSE is the closest with an RMSE of around 0.001 at Q=20 under the strictly neutral model, compared to 0.002 for the corresponding value in the full model.

The rescaling effect is visible in both expanding and contracting populations

Researchers are often interested in observing evolutionary dynamics in populations experiencing demographic changes. We therefore next sought to examine the effect of population size change on the rescaling effect. First, we simulated a variant of our full model where the population experienced a recent two-fold expansion (Methods). Our classifiers for distinguishing between rescaled and unscaled simulations performed similarly to those for the full model with a constant-size population (S12 Fig), although accuracy was generally slightly lower under the expansion model for LD features across values of Q below 20. MPEs for fixation times (Fig 8A) for all mutations are higher across all values of Q except for 20, where the MPE becomes smaller in magnitude compared to the constant sized model. For instance, the MPE for neutral mutations at Q values of 2, 5, and 10 were around 1.3%, 3%, and 3.8% respectively for this model compared to around 1.1%, 2.4%, and 2% respectively for the constant-size full model. At Q of 20 however, the error here decreases to 1.6% compared to the constant-size model's -3.4% (note the sign change, which does not occur under the expansion model). The MPEs for all other statistics (Fig 8B-8D) are very similar to those under the constant-size model but are smaller for LD across all Q values. KL divergence values (S12 Fig) for fixation times are lower in this model compared to the constant-size model, with beneficial mutations having the lowest KL divergence at Q=10 and Q=20 in contrast to the constant-size model (where beneficial mutations exhibited the largest KL divergence at these Q values). For the SFS KL divergence here is very similar to the constant-size model, while the LD RMSE values are lower across all values of Q. Overall, simulations of a population experience a recent expansion seem to produce a rescaling effect that is fairly similar to that of a simulated constant-size population, with the biggest observed difference being in the magnitude of divergence for LD and the magnitude and direction of divergence for fixation times at Q of 20.

Fig 8. Mean percentage error of various statistics for the population expansion model by values of
Q. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average
allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage
error for average linkage disequilibrium as measured by the values of r^2.

 Our next exploration of the effect of population size change was the simulation of a two-fold
decrease in population size. Again, our classifiers for this model perform similarly to those for the
constant-size model, although the classifiers examining LD were slightly more accurate (S14 Fig).
Comparing MPE between the contraction and constant-size models, we observe fairly similar results for
allele frequencies, fixation probabilities, and LD (Fig 9). However, for fixation times, we observe smaller
MPE values for Q=2 through Q=10, and a sign change from positive to negative MPE occurs at Q=10
whereas this is not observed until Q=20 under the constant-size model. More notably, the magnitude of
MPE for fixation times suddenly increases to values much larger than those observed under the constant-
size model (~8–10%, depending on the mutation type, for the contraction model vs. ~3% for the constant-
size model). Concordantly, fixation times show similar KL divergence under the contraction model (S14
Fig) compared to the constant-size model except for Q=20 where KL divergence increases markedly
compared to the constant-size model, with values of 0.057, 0.041, and 0.057 for neutral, beneficial, and
deleterious mutations at Q=20 compared to the constant-size model's values of 0.015, 0.022, and 0.016
respectively. Beneficial mutations are also no longer showing the highest KL divergence for fixation time

across any Q. The KL divergence for the SFS is very similar, with slightly higher KL divergence for beneficial mutations especially at $Q=20$, where the KL divergence is around 0.35 compared to the constant-size model's 0.03. For LD, RMSE values are all higher than the constant-size model, with a value of 0.0027 at $Q=20$ compared to the constant-size model's 0.002.

Fig 9. Mean percentage error of various statistics for the population contraction model by values of Q. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage error for average linkage disequilibrium as measured by the values of r^2.

Scaling effects are greatly diminished in a larger simulated population modeled after *Drosophila* populations

Next, to investigate the impact of rescaling in a larger population, we performed simulations of a *Drosophila melanogaster* population (Methods). When applying the classifiers to this model, we observe the worst performance across all values of Q than seen for any of the other simulation models examined above (S16 Fig). While accuracy hovers around 50% (i.e. random guessing) for all features across almost

all Q values, the random forest classifier using the SFS can achieve 63% accuracy at Q=20, suggesting that the SFS is the simulation outcome most affected by rescaling in this model. Note that the lower accuracy here may be in part due to the smaller chromosomes we simulated for this model because of its greater computational burden (Methods). Thus, there are fewer mutations within each replicate, providing less information to the classifiers. However, this would not affect KL divergence and MPE as much since these measures aggregate information across replicates (although those metrics could still be inflated by noise in sparse datasets; see below).

When examining MPE (S16 Fig) and KL divergence (S17 Fig) we observe substantial bias in allele frequencies and fixation times and probabilities for deleterious mutations, but comparatively subtle effects observed for neutral and beneficial mutations. For instance, the average fixation time MPE is ~70% at Q=200 for deleterious mutations, but only around -0.7% and -0.1% for neutral and beneficial mutations, respectively. The MPE for the average fixation probability is also very high for deleterious mutations at Q=200 (~200%) and low for beneficial and neutral mutations (~-0.6% and ~1.5% respectively). While the MPE for fixation probability is higher at Q=200 for deleterious mutations, it is not monotonic for neutral and beneficial mutations, since the highest fixation time MPE in magnitude is at Q=150 for neutral mutations (~-0.84%) and at Q=50 for beneficial mutations (~0.36%) and the highest fixation probability MPE is at Q=50 for beneficial mutations (~2.5%). While the fixation probability MPE is at its highest at Q=200 for neutral mutations, it only increases in magnitude monotonically, but abruptly flips sign at Q=100. For allele frequency MPE, the trend is non-monotonic for all three mutation types and is higher at Q=150 for deleterious mutations (~17%), at Q=100 for beneficial mutations (~-2%) and at Q=50 for neutral mutations (~-0.8%). However, it is important to note here that the probability of the fixation of deleterious mutations is extremely small, ~1.5×10^{-12} for Q=20 and no such fixations were observed for Q values of 50, 100, and 150 (yielding a fixation probability estimate of zero). This also means that the distribution of fixation times for deleterious mutations was based on very sparse information and in some cases could not be estimated at all. This was not a concern for neutral and beneficial mutations, however, and we observed that KL divergence for fixation times of these mutations (S16 Fig) shows a weak and non-monotonic trend with increasing Q. KL divergence for the SFS and RMSE for the LD were also small and non-monotonic. Overall, the degree of the rescaling effects under this model was fairly small for beneficial and neutral mutations. Although our error measures were much higher for deleterious mutations, these probably reflect the sparsity of our data rather than a genuinely strong rescaling effect.

Scaling effects are substantial in a population modeled after *Drosophila* but with stronger selection on beneficial mutations

In the previous section we examined the behavior of a *D. melanogaster* population model with weak selection acting on beneficial mutations (s=7.125×10⁻⁶ [59,64]). We therefore sought to examine whether the scaling effects remain diminished in our *Drosophila* modeled population when a higher selection coefficient for beneficial mutations is used (s=0.01); we refer to this model as the "strong beneficial" model. Classifier results (S19 Fig) indicate that scaling effects are much more drastic when strongly beneficial mutations are present. Accuracy for the random forest model for fixation times and probabilities are 100% for all values of Q. Accuracy for LD and SFS features are close to 50% for Q of 50 but increase to around 63% and 66% respectively for Q of 100 and increase to around 69% and 75% respectively at Q of 200.

MPE of fixation times (Fig 10A) shows a monotonic increase in error values for all mutation types with increasing Q. While deleterious mutations have errors that are smaller and in the opposite direction than the model with weaker beneficial mutations across all values of Q, errors for beneficial and neutral mutations are much larger. For instance, neutral mutations exhibit an MPE for average fixation time of ~12% at Q=50 which increases to 70% for Q=200. In contrast the fixation time MPEs for neutral mutations under the weak beneficial model at those same values of Q are -0.4% and -0.6%. MPE for beneficial mutations' average fixation time in the strong beneficial model is ~13% at Q=50 and increasing to around 70% at Q=200, compared to the corresponding values of 0.4% and -0.09% in the weak beneficial model. Deleterious mutations exhibit errors of ~10% at Q=1 and 54% at Q=20. We note that unlike under the weak beneficial model, deleterious mutations do occasionally fix at all values of Q under the strong beneficial model, likely due to hitchhiking with beneficial mutations.

The trend for mutation frequency MPE (Fig 10B) is more complex and non-monotonic. For neutral mutations, MPE alternates its sign at each successive Q, increasing in magnitude from ~-0.2% at Q of 50 to ~13% for Q of 200. Deleterious mutations exhibit the highest error at Q of 50 (around 48%) which then declines at Q of 100, increases again at Q of 150, and then decreases to its lowest value of ~14% at Q of 200. In contrast, beneficial mutation frequency MPE increases monotonically from ~4% at Q of 50 to ~14% at Q of 200. Of note here is that while all mutation types have drastically different values of MPE for allele frequency at Q=50, 100, 150, they eventually exhibit similar errors at Q of 200. Furthermore, all error values for this model are higher compared to their corresponding mutations type in the weak beneficial model, even for deleterious mutations which have a maximum error of around 17% for the weak beneficial model. For fixation probabilities (Fig 10C), MPE is lowest for neutral mutations, where the highest magnitude MPE is at Q of 50 (around -0.5%) which flips sign at Q of 100 to 0.11% and increases to around 0.4% at Qs of 150 and 200. Beneficial and deleterious mutations have similar errors, which are negative and increase in magnitude monotonically with Q to -0.30% for both mutation types. This starkly different than the corresponding values for the lower selection *Drosophila* model, where neutral and beneficial mutations have maximum errors of 1.2% and 2.5%, respectively and again where deleterious mutations fixed very rarely, preventing the reliable estimation of MPEs. MPE values for LD (Fig 12C) are also non-monotonic with Q as was the case for the weak beneficial model, but are much higher in magnitude, with the highest value being around 4.7% at Q of 100, compared to the highest value of around 0.34% at Q of 200 for the lower selection model.

KL divergence for fixation times (S19 Fig) increases monotonically with Q from values of around 0.05, 0.16, and 0.08 for neutral, deleterious, and beneficial mutations respectively at Q of 50 to around 1.09, 1.7, and 1.51, respectively, at Q of 200. On the other hand, KL divergence values for the SFS exhibit different trends for each mutation type; for beneficial mutations, KL divergence increase monotonically from around 0.02 at Q=50 to around 0.06 at Q=200. For neutral mutations, the trend is actually a monotonic decrease from a value of around 0.3 at Q=50 to around 0.21 at Q of 200. KL divergence for deleterious mutations is non monotonic, initially increasing from a value of about 0.53 at Q=50 to around 0.6 at Q=100, decreasing 0.52 and 0.54 and the two subsequent Q values, respectively. RMSE for LD also does not show a clear trend, with all values hovering around 0.005.

Figure 10. Mean percentage error of various statistics for the higher selection *Drosophila* model by values of *Q*. **(A)** Mean percentage error for average mutation fixation times. **(B)** Mean percentage error for average allele frequencies. **(C)** Mean percentage error for average fixation probabilities. **(D)** Mean percentage error for average linkage disequilibrium as measured by the values of r^2.

Scaling effects are present in a single selective sweep model

Finally, we sought to exam the effects of scaling in a simple hitchhiking model with only neutral mutations and a single beneficial mutation sweeping through the population. We focus here on the distribution of fixation times for the beneficial mutation across all replicates, and on the SFS and LD for neutral mutations. Classifier accuracy results (S21 Fig) show that scaling also affects these simulations with the intensity of the effects increasing with *Q*. While classifier accuracy is low at *Q*=2, with the best accuracy being 56% for LD, accuracy increases with *Q*, especially for LD where accuracy is at 85% for *Q*=5 and reaches 100% for *Q*=20. The highest classification accuracy for fixation times and the SFS are also observed at *Q*=20 (80% and 91%, respectively). MPE for fixation times of the beneficial mutation (Fig 11A) are substantial and increase monotonically with *Q* from 1.8% at *Q*=2 to 35% at *Q*=20. MPEs for the neutral mutation allele frequency also increase with *Q* but are smaller, reaching only as high as 1.2% at *Q*=20. For LD, we also observe a monotonic increase in MPE from 0.5% at *Q*=2 to 16% at *Q*=20.

KL divergence for the distribution of fixation times (S21 Fig) of the beneficial mutation decreases from 0.05 at Q=2 to 0.03 at Q=5 and then increases monotonically to 0.31 at Q=20. For neutral mutations, the SFS KL divergence and LD RMSE for both increase monotonically with Q, with the highest KL divergence and RMSE values being 0.0005 and 0.001 respectively at Q=20. These results demonstrate that even with a scenario involving a single sweep, scaling can bias simulation outcomes, especially fixation times for the sweeping mutation and LD between pairs of linked neutral mutations.

Fig 11. Mean percentage error of various statistics for the single sweep model by values of Q. **(A)** Mean percentage error for average beneficial mutation fixation times. **(B)** Mean percentage error for average allele frequencies for neutral mutations. **(C)** Mean percentage error for average linkage disequilibrium for neutral mutations as measured by the values of r^2.

Scaling effects can be often recapitulated from a small number of replicates

Although some general conclusions can be drawn from our experiments above (e.g. rescaling effects are common and generally increase with Q), behavior differs substantially across models and thus it may be difficult for a researcher to know what the impact of rescaling may be on their simulations without conducting similar experiments themselves on the model of interest. Since it is often unfeasible to acquire many replicates of minimally-scaled (ideally unscaled) simulations under some models due to computational limitations, we sought to investigate whether statistical measures of the impact of scaling such as MPE, KL divergence, and RMSE can be reasonably approximated with a smaller number of

simulation replicates for a subset of our simulations. We therefore randomly sampled smaller sets of replicates for the full model, the no-beneficials model, and the *Drosophila* model with strong beneficials for each value of Q. Our results indicate that when information about the outcome of interest is sufficiently dense in each replicate, measures of deviations gleaned from a smaller number of replicates are generally close to those obtained using all 1000 replicates. For example, in a sample of 100 replicates simulated under the full model (S22 Fig), all measures of deviation are extremely similar to those obtained using 1000 replicates, with the exception of KL divergence for the SFS of beneficial mutations, which is lower for all values of Q compared to the results from 1000 replicates. When using only 10 replicates (S23 Fig), deviation outcomes for the full model are still informative but less well preserved. For instance, the magnitude of the MPE of average fixation times is slightly higher for neutral and beneficial mutations at Q=5 and slightly lower at Q=20. KL divergence for fixation times and the SFS is more severely affected, especially for neutral and deleterious mutations, whereas the average RMSE for LD is still well-preserved.

For the no-beneficials model, using 100 replicates (S24 Fig) preserves all measures of deviation rather well, with the exception of a lower KL divergence for fixation times of neutral mutations at Q=20 compared to that estimated from 1000 replicates. Using 10 replicates only was sufficient to recover similar estimates of MPE for average fixation times, probabilities, and LD, but induced a more marked change in the average allele frequency MPE (S25 Fig). Changes in KL divergence for fixation times and the SFS are also more pronounced for this small number of replicates, with lower estimated KL divergence for both mutation types for both outcomes. The average RMSE for LD, however, remains well-preserved.

Unfortunately, down-sampling the number of replicates for the *Drosophila* model does not recapitulate the measured errors for some outcomes as well. While down-sampling to 100 replicates (S26 Fig) does conserve some error measurements, such as the average fixation time MPEs for neutral and beneficial mutations, the average fixation probability MPE for beneficial mutations, and the average allele frequency for beneficial mutations, we observed substantial shifts for all other measurements, including KL divergence for fixation times and the SFS for all mutation types and the average LD RMSE. Further down-sampling to 10 replicates (S27 Fig) causes further loss of ability to capture all measures of error. This result is not unexpected, as we simulated much smaller regions in the *Drosophila* genome to examine smaller scaling factors (Methods), and this resulted in a much smaller number of mutations available for analysis (e.g. the estimated SFS was based on fewer than 20 mutations present in a sample for the *Drosophila* higher selection model on average).

In summary, using a smaller number of replicates can be informative for characterizing deviation between scaled and unscaled simulation when replicates are sufficiently information dense (e.g. there are enough polymorphisms in the sample taken at the end of the simulation to estimate the SFS, or there were enough fixations during the simulation to accurately estimate the fixation probability). Using a much smaller number of replicates can impair these estimates, especially for KL divergence, and most severely for simulation models whose replicates contain sparse information such as a low number of segregating or fixed polymorphisms (e.g. our simulations of 10 kb regions in the *Drosophila* genome).

DISCUSSION

Population genetic simulation software is increasingly powerful and flexible [43,45]. This is especially so for forward-in-time simulators that allow for complex scenarios involving different types of natural

selection [47,48] departures from Wright-Fisher assumptions [52], the modeling of multispecies dynamics [35], and more. Unfortunately, because forward simulations require the entire population to be modeled, they can be too computationally costly when the population size is large. Rescaling the simulation parameters to simulate a smaller population to model a larger one is therefore necessary in many cases. Unfortunately, it has been shown that this rescaling trick can bias simulation outcomes [2,53]. However, the rescaling effect has only been examined for a small number of simulation scenarios, and thus we still do not know the manner and magnitude with which it will affect simulation results.

Here, we sought to investigate the rescaling effect on a wider array of simulation models, including those of varying genome densities, population sizes, and strengths of positive and negative selection. We used $\texttt{stdpopsim}$ and $\texttt{SLiM}$ to generate these simulations under varying values of the rescaling factor Q, and record a set of simulation outcomes summarizing the SFS, LD, and the probability of and time to fixation. Ideally, the rescaling strategy would produce results that are difficult to distinguish from unscaled simulations. However, this was not the case—for most of the models that we examined, binary classifiers were able to discriminate between rescaled and unscaled simulations quite easily. For example, for our full model, classifiers examining either the estimated fixation probability and the distribution of fixation times could label simulation replicates as unscaled or rescaled by a factor of only $Q=2$ with 100% accuracy (Fig 2). Classifiers using the SFS or LD also performed well under this model with only a modest degree of rescaling (e.g. nearly 100% accuracy at $Q=5$). Our classifiers generally performed well for the other models examined here, but there were two notable exceptions: 1) for the neutral model, only the LD-based classifier could discriminate between unscaled and scaled simulations with 100% accuracy at $Q=20$ (the highest rescaling factor examined for this model; see S10 Fig); 2) for the *Drosophila* model with weaker selection on beneficial mutations, none of the classifiers performed well at any value of Q examined (although this was not the case for the *Drosophila* model with strong beneficials).

Even if rescaling alters simulations sufficiently to be detected by classifiers, the size of this rescaling effect may still in some cases be small enough to be tolerable by researchers. We therefore sought to more directly quantify the magnitude by measuring the mean percent error between scaled and unscaled simulations for each of our simulation outcomes. We found that MPE between scaled and unscaled outcomes was often substantial. For example, the fixation probability of deleterious mutations under our no-beneficials model is overestimated by ~140% when $Q=20$ (Fig 4C). However, MPE did vary considerably across simulation outcomes and models. Consider again the MPE of fixation probability under the no-beneficials model. When we add beneficial mutations into this model (yielding the "full model"; Methods), the MPE for deleterious mutations is only 0.2% at $Q=20$ (Fig 3C). We note that classification accuracy did not always correspond with the observed trends in MPE. For instance, in the no-beneficials model, the random forest classifier achieves higher accuracy on the SFS features than fixation times, despite allele frequencies having an MPE that is an order of magnitude lower than that of fixation times. This is probably a result of higher variance on our fixation time estimates, resulting in more overlap between scaled and unscaled simulations despite a comparatively large shift in the mean. Thus, one should bear in mind that our classification results are indicative of the degree to which the scaled and unscaled simulations are separable—which could be the case for two distributions that have a slight difference in mean value but low variance—while our estimates of MPE and KL divergence/RMSE more directly speak to the magnitude of the rescaling effect.

Our analysis of the variety of simulation models considered here does yield some conclusions that are likely to generalize to other models. For one, we might expect the rescaling effect to be more severe in smaller populations than in larger ones. For example, the magnitude of the rescaling effect observed in

our no-beneficials model (Fig 4) was reduced somewhat when we re-ran these simulations with a population that was twice as large (Fig 5). We also observed a weaker rescaling effect in the simulated *Drosophila* population, although this was in part a result of the sparsity of mutation data in these simulations which modeled smaller chromosomes. However, the rescaling effect may still be appreciable in larger populations when positive selection plays a more prominent role and when Q is large: the rescaling effect in the *Drosophila* model with stronger beneficial mutations is far greater than that under the model with weaker beneficials (compare Fig 10 to S16 Fig).

Intuitively, it is not unexpected that the rescaling effect would be greater in magnitude in the presence of selection. For example, in the presence of positive selection, the rationale for rescaling the selection coefficient by Q is not supported by theory, as neither the time to fixation nor the fixation probability are expected to be a linear function of the population-scaled selection coefficient $2Ns$ [67,68]—our results from simple simulations of a single selective sweep underscore this (Fig 11). Similarly, Uricchio and Hernandez [53] showed that the theoretical prediction that the impact of a sweep on a linked variation is determined by $2Ns$ [69] is not valid when s is large, which will be the case when s is multiple by Q during rescaling. Moreover, we would expect there to be a stronger rescaling effect in the presence of large amounts of natural selection because rescaling will cause each simulated genome to receive a larger number of selected mutations each generation, which may alter short-term dynamics before the increased recombination rate can break up this extra linkage. Finally, rescaling divides the total number of generations by Q, thereby reducing the temporal resolution of the simulation when it comes to events that occur on short timescales. For example, all estimates of selective sweep sojourn times will necessarily be rounded up to the nearest multiple of Q, resulting in underestimates. A corollary of all these lines of reasoning is that the rescaling effect should be less severe under neutral simulations, perhaps especially those of large populations. However, although the effect is more subtle, we do observe one here (Fig 7). We note that this finding is consistent with results from Adrion et al. [2] who observed biased estimates of several summary statistics when performing purely neutral rescaled forward simulations of the demographic model from Kamm et al. [70].

Although we find that the consequences of population-size rescaling for forward simulations are somewhat predictable, when examining the totality of our results one cannot help but be struck by numerous exceptions to the general trends described above. Indeed, as noted in numerous places throughout the Results we did not always find that the magnitude of MPE increased monotonically with Q. Thus, it is not as of yet possible to predict a priori precisely how, for a population genetic model of interest, which simulation outcomes will be most affected by rescaling, and whether these outcomes will be upwardly or downwardly biased for a given value of Q. Thankfully, we have found that even a relatively small number of simulation replicates (e.g. 10, for the cases shown in (S25 Fig, S26 Fig, and S27 Fig) can be enough to accurately describe the impact of rescaling both qualitatively and quantitatively (via MPE) for a given Q. Thus, a small-scale test of varying Q values may be sufficient for a researcher wishing to determine which value of Q will provide the best balance between computational efficiency and accuracy of estimated outcomes. For researchers wishing to simulate very large numbers of simulation replicates each of reasonably small chromosome lengths (e.g. for ABC inference of models involving natural selection [57]), this will mean just simulating a small number of replicates for each of a range of rescaling factors. For researchers instead hoping to generate chromosome-scale simulation replicates, it may be possible to first do a small-scale test on shorter chromosomal regions to select a value of Q before conducting the full-scale study. For simpler scenarios, more theoretically motivated algorithmic approaches to selecting an optimal Q value may be possible [53]. Thus, although population rescaling may in many cases be unavoidable for the time being, researchers may be able to mitigate the undesirable effects of rescaling by carefully considering and investigating them before performing

simulations at scale. Still, we suggest that continued study of the manner in which population rescaling effects simulation results and the models affected will be necessary to ensure that the continuing advances in simulation power and flexibility are being used to produce accurate results.

ACKNOWLEDGMENTS

The authors thank members of the Popsim Consortium for feedback on preliminary results from this study. This work was funded by NIH awards R01HG010774 and R35GM138286.

References

1. Peng B, Chen HS, Mechanic LE, Racine B, Clarke J, Gillanders E, et al. Genetic data simulators and their applications: an overview. Genet Epidemiol. 2015 Jan;39(1):2–10.

2. Adrion JR, Cole CB, Dukler N, Galloway JG, Gladstein AL, Gower G, et al. A community-maintained standard library of population genetic models. Coop G, Wittkopp PJ, Novembre J, Sethuraman A, Mathieson S, editors. eLife. 2020 Jun 23;9:e54967.

3. Hoban S, Bertorelle G, Gaggiotti OE. Computer simulations: tools for population and evolutionary genetics. Nat Rev Genet. 2012 Feb;13(2):110–22.

4. Hudson RR. Properties of a neutral allele model with intragenic recombination. Theor Popul Biol. 1983 Apr 1;23(2):183–201.

5. Charlesworth B, Morgan MT, Charlesworth D. The Effect of Deleterious Mutations on Neutral Molecular Variation. Genetics. 1993 Aug;134(4):1289–303.

6. Hermisson J, Pennings PS. Soft Sweeps. Genetics. 2005 Apr;169(4):2335–52.

7. Orr HA, Unckless RL. Population extinction and the genetics of adaptation. Am Nat. 2008 Aug;172(2):160–9.

8. Zeng K. A coalescent model of background selection with recombination, demography and variation in selection coefficients. Heredity. 2013 Apr;110(4):363–71.

9. Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of deleterious mutations during range expansions. Mol Ecol. 2013;22(24):5972–82.

10. Morgan K, Strobeck C. Is intragenic recombination a factor in the maintenance of genetic variation in natural populations? Nature. 1979 Feb;277(5695):383–4.

11. Braverman JM, Hudson RR, Kaplan NL, Langley CH, Stephan W. The hitchhiking effect on the site frequency spectrum of DNA polymorphisms. Genetics. 1995 Jun;140(2):783–96.

12. Schrider DR. Background Selection Does Not Mimic the Patterns of Genetic Diversity Produced by Selective Sweeps. Genetics. 2020 Oct;216(2):499–519.

13. Battey CJ, Ralph PL, Kern AD. Space is the Place: Effects of Continuous Spatial Structure on Analysis of Population Genetic Data. Genetics. 2020 May 1;215(1):193–214.

14. Torres R, Stetter MG, Hernandez RD, Ross-Ibarra J. The Temporal Dynamics of Background Selection in Nonequilibrium Populations. Genetics. 2020 Apr 1;214(4):1019–30.

15. Thornton KR. Polygenic Adaptation to an Environmental Shift: Temporal Dynamics of Variation Under Gaussian Stabilizing Selection and Additive Effects on a Single Trait. Genetics. 2019 Dec 1;213(4):1513–30.

16. Csilléry K, Blum MGB, Gaggiotti OE, François O. Approximate Bayesian Computation (ABC) in practice. Trends Ecol Evol. 2010 Jul 1;25(7):410–8.

17. Beaumont MA. Approximate Bayesian Computation in Evolution and Ecology. Annu Rev Ecol Evol Syst. 2010;41(1):379–406.

18. Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population genetics. Genetics. 2002 Dec;162(4):2025–35.

19. Tavare S, Balding DJ, Griffiths RC, Donnelly P. Inferring Coalescence Times from DNA Sequence Data. Genetics. 1997 Feb;145(2):505–18.

20. Pritchard JK, Seielstad MT, Perez-Lezaun A, Feldman MW. Population growth of human Y chromosomes: a study of Y chromosome microsatellites. Mol Biol Evol. 1999 Dec 1;16(12):1791–8.

21. Galimberti M, Leuenberger C, Wolf B, Szilágyi SM, Foll M, Wegmann D. Detecting Selection from Linked Sites Using an F-Model. Genetics. 2020 Dec 1;216(4):1205–15.

22. Alachiotis N, Pavlidis P. RAiSD detects positive selection based on multiple signatures of a selective sweep and SNP vectors. Commun Biol. 2018 Jun 27;1(1):1–11.

23. Pavlidis P, Živković D, Stamatakis A, Alachiotis N. SweeD: Likelihood-Based Detection of Selective Sweeps in Thousands of Genomes. Mol Biol Evol. 2013 Sep 1;30(9):2224–34.

24. Caravagna G, Heide T, Williams MJ, Zapata L, Nichol D, Chkhaidze K, et al. Subclonal reconstruction of tumors by using machine learning and population genetics. Nat Genet. 2020 Sep;52(9):898–907.

25. Sheehan S, Song YS. Deep Learning for Population Genetic Inference. PLOS Comput Biol. 2016 Mar 28;12(3):e1004845.

26. Schrider DR, Kern AD. Supervised Machine Learning for Population Genetics: A New Paradigm. Trends Genet. 2018 Apr 1;34(4):301–12.

27. Mughal MR, DeGiorgio M. Localizing and Classifying Adaptive Targets with Trend Filtered Regression. Mol Biol Evol. 2019 Feb 1;36(2):252–70.

28. Pavlidis P, Jensen JD, Stephan W. Searching for footprints of positive selection in whole-genome SNP data from nonequilibrium populations. Genetics. 2010 Jul;185(3):907–22.

29. Ronen R, Udpa N, Halperin E, Bafna V. Learning Natural Selection from the Site Frequency Spectrum. Genetics. 2013 Sep;195(1):181–93.

30. Flagel L, Brandvain Y, Schrider DR. The Unreasonable Effectiveness of Convolutional Neural Networks in Population Genetic Inference. Mol Biol Evol. 2019 Feb 1;36(2):220–38.

31. Chan J, Perrone V, Spence JP, Jenkins PA, Mathieson S, Song YS. A Likelihood-Free Inference Framework for Population Genetic Data using Exchangeable Neural Networks. Adv Neural Inf Process Syst. 2018 Dec;31:8594–605.

32. Gao F, Ming C, Hu W, Li H. New Software for the Fast Estimation of Population Recombination Rates (FastEPRR) in the Genomic Era. G3 GenesGenomesGenetics. 2016 Jun 1;6(6):1563–71.

33. Smith CCR, Tittes S, Ralph PL, Kern AD. Dispersal inference from population genetic variation using a convolutional neural network. Genetics. 2023 May 26;224(2):iyad068.

34. Sanchez T, Cury J, Charpiat G, Jay F. Deep learning for population size history inference: Design, comparison and combination with approximate Bayesian computation. Mol Ecol Resour. 2021 Nov;21(8):2645–60.

35. Haller BC, Messer PW. SLiM 4: Multispecies Eco-Evolutionary Modeling. Am Nat. 2023 May;201(5):E127–39.

36. Ewing G, Hermisson J. MSMS: a coalescent simulation program including recombination, demographic structure and selection at a single locus. Bioinformatics. 2010 Aug 15;26(16):2064–5.

37. Kern AD, Schrider DR. Discoal: flexible coalescent simulations with selection. Bioinformatics. 2016 Dec 15;32(24):3839–41.

38. Hudson RR. Generating samples under a Wright–Fisher neutral model of genetic variation. Bioinformatics. 2002 Feb 1;18(2):337–8.

39. Kelleher J, Thornton KR, Ashander J, Ralph PL. Efficient pedigree recording for fast population genetics simulation. PLOS Comput Biol. 2018 Nov 1;14(11):e1006581.

40. Messer PW. SLiM: Simulating Evolution with Selection and Linkage. Genetics. 2013 Aug 1;194(4):1037–9.

41. Liang L, Zöllner S, Abecasis GR. GENOME: a rapid coalescent-based whole genome simulator. Bioinformatics. 2007 Jun 15;23(12):1565–7.

42. Carvajal-Rodriguez A. Simulation of Genomes: A Review. Curr Genomics. 2008 May 1;9(3):155–9.

43. Kelleher J, Etheridge AM, McVean G. Efficient Coalescent Simulation and Genealogical Analysis for Large Sample Sizes. PLOS Comput Biol. 2016 May 4;12(5):e1004842.

44. Hudson RR. TESTING THE CONSTANT-RATE NEUTRAL ALLELE MODEL WITH PROTEIN SEQUENCE DATA. Evol Int J Org Evol. 1983 Jan;37(1):203–17.

45. Baumdicker F, Bisschop G, Goldstein D, Gower G, Ragsdale AP, Tsambos G, et al. Efficient ancestry and mutation simulation with msprime 1.0. Genetics. 2022 Mar 1;220(3):iyab229.

46. Teshima KM, Innan H. mbs: modifying Hudson's ms software to generate samples of DNA sequences with a biallelic site under selection. BMC Bioinformatics. 2009 May 30;10(1):166.

47. Haller BC, Messer PW. SLiM 2: Flexible, Interactive Forward Genetic Simulations. Mol Biol Evol. 2017 Jan 1;34(1):230–40.

48. Thornton KR. A C++ template library for efficient forward-time population genetic simulation of large populations. Genetics. 2014 Sep;198(1):157–66.

49. Hernandez RD. A flexible forward simulator for populations subject to selection and demography. Bioinformatics. 2008 Dec 1;24(23):2786–7.

50. Peng B, Kimmel M. simuPOP: a forward-time population genetics simulation environment. Bioinformatics. 2005 Sep 1;21(18):3686–7.

51. Neuenschwander S, Hospital F, Guillaume F, Goudet J. quantiNemo: an individual-based program to simulate quantitative traits with explicit genetic architecture in a dynamic metapopulation. Bioinforma Oxf Engl. 2008 Jul 1;24(13):1552–3.

52. Haller BC, Messer PW. SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model. Mol Biol Evol. 2019 Mar 1;36(3):632–7.

53. Uricchio LH, Hernandez RD. Robust Forward Simulations of Recurrent Hitchhiking. Genetics. 2014 May 1;197(1):221–36.

54. Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genet Res. 1966 Dec;8(3):269–94.

55. Nei M, Li WH. Mathematical model for studying genetic variation in terms of restriction endonucleases. Proc Natl Acad Sci U S A. 1979 Oct;76(10):5269–73.

56. Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. Genetics. 1989 Nov;123(3):585–95.

57. Johri P, Charlesworth B, Jensen JD. Toward an Evolutionarily Appropriate Null Model: Jointly Inferring Demography and Purifying Selection. Genetics. 2020 May;215(1):173–92.

58. Comeron JM, Kreitman M. Population, Evolutionary and Genomic Consequences of Interference Selection. Genetics. 2002 May 1;161(1):389–410.

59. Lauterbur ME, Cavassim MIA, Gladstein AL, Gower G, Pope NS, Tsambos G, et al. Expanding the stdpopsim species catalog, and lessons learned for realistic genome simulations. Gao Z, Przeworski M, editors. eLife. 2023 Jun 21;12:RP84874.

60. Boyko AR, Williamson SH, Indap AR, Degenhardt JD, Hernandez RD, Lohmueller KE, et al. Assessing the Evolutionary Impact of Amino Acid Mutations in the Human Genome. PLOS Genet. 2008 May 30;4(5):e1000083.

61. Li H, Stephan W. Inferring the Demographic History and Rate of Adaptive Substitution in Drosophila. PLOS Genet. 2006 Oct 13;2(10):e166.

62. Schrider DR, Houle D, Lynch M, Hahn MW. Rates and Genomic Consequences of Spontaneous Mutational Events in Drosophila melanogaster. Genetics. 2013 Aug;194(4):937–54.

63.   Comeron JM, Ratnappan R, Bailin S. The Many Landscapes of Recombination in Drosophila melanogaster. PLOS Genet. 2012 Oct 11;8(10):e1002905.

64.   Ragsdale AP, Coffman AJ, Hsieh P, Struck TJ, Gutenkunst RN. Triallelic Population Genomics for Inferring Correlated Fitness Effects of Same Site Nonsynonymous Mutations. Genetics. 2016 May 1;203(1):513–23.

65.   Huber CD, Kim BY, Marsden CD, Lohmueller KE. Determining the factors driving selective effects of new nonsynonymous mutations. Proc Natl Acad Sci. 2017 Apr 25;114(17):4465–70.

66.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine Learning in Python. J Mach Learn Res. 2011;12(85):2825–30.

67.   Charlesworth B. How Long Does It Take to Fix a Favorable Mutation, and Why Should We Care? Am Nat. 2020 May;195(5):753–71.

68.   Kimura M. On the Probability of Fixation of Mutant Genes in a Population. Genetics. 1962 Jun;47(6):713–9.

69.   Wiehe TH, Stephan W. Analysis of a genetic hitchhiking model, and its application to DNA polymorphism data from Drosophila melanogaster. Mol Biol Evol. 1993 Jul 1;10(4):842–54.

70.   Kamm J, Terhorst J, Durbin R, Song YS. Efficiently Inferring the Demographic History of Many Populations With Allele Count Data. J Am Stat Assoc. 2020 Jul 2;115(531):1472–87.