Loss of precise auditory sampling as a sign of value-driven visual attentional capture

For correspondence:
fcervantes@psico.edu.uy

Present address: Centro de Investigación Básica en Psicología, Facultad de Psicología, Universidad de la República, Montevideo, 11.200 Uruguay

Keywords: value-driven attentional capture | reward-driven distraction | auditory encoding | phase-locking value

Rodrigo Caramés Harcevnicow[1,*], Thaiz Sánchez-Costa[1,*], Alejandra Carboni[1], and Francisco Cervantes Constantino[1,2]

[1]Facultad de Psicología, Universidad de la República;

[2]Instituto de Investigaciones Biológicas Clemente Estable

[*]These authors contributed equally

Abstract   Conditioning sensory signals endows persistent salience that influences attention, even when their presence no longer connects with rewards. These irrelevant distractors also shape multisensory processing, but how do effects sustain throughout continuous encoding remains poorly understood. We test if value-driven capture interferes with the ability of cortical activity to reliably phase-lock to temporal modulations in audiovisual (AV) signals. Listening to periodically-modulated sound, observers discriminated between two visual object streams flickering at different rates, in order to identify an AV match. Peripheral color cues were shown simultaneously, and evaluated for their ability to modulate participant responses due to a color-reward associative training taken beforehand. Behavioral sensitivity and electroencephalography recordings show that performance impoverished in presence of colors previously associated with reward, as did the phase locking of AV responses. Decreased temporal precision predicted participants' reward-driven distraction, evidencing the attentional shift away from temporal representations of the multimodal target. Consistency loss was also confirmed in auditory responses, identifying a previously unknown effect of value-driven attentional capture to withdraw from unimodal tracking fidelity. The findings offer understanding on the effective inter-modal competition set out when tracking incentive salience cues overtakes goal-oriented tracking of multisensory streams connected in time.

Value-driven attentional capture (VDAC) encapsulates the experience where rewarded aspects of our recent past involuntarily influence selection of sensory information (*Anderson, 2019*; *Anderson et al., 2011*, *2021*). It describes situations in which previously tangible benefits are no longer provided in relation to current goals, illustrating the

constraints that rewarded backgrounds place upon how we engage with the sensory environment (*Todd & Manaligod, 2018*). VDAC transiently adjusts the share of neural processing dedicated to current targets, biasing the balance towards 'has-been' cues (*Rusz et al., 2020*), and penetrates through different sensory modalities than current demands require. One of the few studies that specifically investigated the latter aspect demonstrated the reshaping of visual cortex representations in presence of sounds with a rewarded history, predicting observers' visual sensitivity changes (*Pooresmaeili et al., 2014*). Using brief stimulation designs, this and newer findings left open the question of how multimodal streams that unfold over time through common temporal dynamics could be robust to or instead molded by persisting salience signals. Importantly, time features in most if not all natural associations promoting binding of multisensory streams into stable objects (*Bizley et al., 2016*).

To build multimodal associations from dynamic streams, the brain seems best served if it precisely entrains cortical sensory and associative networks to multiple component features. A prime form of entrainment of cortical populations takes place in the tracking responses of specialized auditory primary and secondary areas to slow (< 12 Hz) amplitude modulations of sound. With its relatively high temporal precision, human auditory cortical entrainment may be nonetheless molded by the dynamics of voluntarily attended visuals that are temporally coherent, turning them to greater multimodal consistency (*Atilgan et al., 2018*; *Maddox et al., 2015*). To date, it is unclear if bottom-up influences such as VDAC similarly modulate entrainment or are prevented from doing so, e.g., via reactive suppression mechanisms (*Klink et al., 2023*). Importantly, the same region that relays cross- and supramodal input to auditory cortex, the temporo-parieto-occipital junction (*Rolls et al., 2022*), has been directly involved in AV manifestations of VDAC (*Pooresmaeili et al., 2014*; *Antono et al., 2023*).

Switching attention from one sense to the other brings in variability in neurally entrained activity across time: when visual input is attended, local sub-threshold excitability patterns in auditory cortex become aligned to the visual event (*Lakatos et al., 2009*). Responses to ignored auditory elements also appear relatively less sharp and stable when competing elements are being attended (*Lakatos et al., 2013*). These variations suggest that ongoing vision of reward-associated cues could set a recurring context unfavorable to steady tracking of auditory stimuli. If the latter become less salient by virtue of competing cues, temporal variability in associated phase-locked auditory responses may be expected to increase (*Kayser, 2009*). The precision of temporal locking patterns of activity within auditory networks moreover appears as a relevant index in perceptual AV synchrony judgment tasks (*Kösem et al., 2014*).

The current study aims to demonstrate the impact of visual VDAC on the precision of cortical responses locked to temporal modulations in AV stimuli. Specifically, we evaluate changes on phase variability of audiovisual and unimodal auditory responses during AV perceptual judgments. In a two-alternative forced choice task, subjects matched one of two central visual flicker options with a single amplitude-modulated (AM) sound. Conditioned visual distractors, shown at the periphery, conveyed past reward associations. Typically set in probabilistic designs (*Anderson et al., 2011*; *Hickey & Peelen, 2015*; *Pooresmaeili et al., 2014*; *Sanz et al., 2018*; *Cho & Cho, 2021*; *Koenig et al., 2017*), but see (*Anderson, 2016*; *Tankelevitch et al., 2020*), reward associative learning was addressed through probabilistic

and non-probabilistic (fixed) conditioning schemes as the latter are expected to be less effective (*Rouhani et al., 2018*; *Monosov, 2020*; *Hart et al., 2015*). Our results provide first evidence of the ability of both visual conditioning setups to disrupt steady AV processing, decreasing the temporal precision of sound-locked responses through time.

Materials and Methods

Participants

Thirty-four subjects (15 female; mean age 26.4 ± 4.0 SD; 4 left-handed) with no history of neurological or psychiatric disorders volunteered in the study. All participants provided formal written informed consent. They reported normal hearing and normal or corrected to normal visual acuity. The experiments were performed in accordance with the World Medical Association Declaration of Helsinki guidelines. The Ethics in Research Committee of the School of Psychology at the Universidad de la República approved the experimental procedures. For two subjects, the second half of EEG recordings were not saved in error, reducing analyses to 32 participants. Specific participant data affected by an incorrect color display (1 participant) or missing sound output feedback recordings (1 participant) were not included in analyses based on that information, as indicated in the relevant Statistical analysis sections, below.

Experimental equipment and setup

Participants were seated at 50 cm in front of a 40 cm CRT monitor (E. Systems, Inc., CA) delivering visual presentations with an 83 dpi resolution and an 85 Hz refresh rate. Audio presentations were delivered with a Sound Blaster Z sound card (Creative Labs, Singapore) in combination with a Scarlett 4i4 hub interface (Focusrite Plc, UK) and high-quality Sennheiser HD 25 headphones (Sennheiser, Germany). Neural recordings were performed using a BioSemi ActiveTwo 64-channel EEG system (BioSemi, The Netherlands) with a 10/20 CMS/DRL (ground) layout at a 2048 Hz digitization rate. Audio signal sampling was obtained from sound interface output fed back to the ActiveTwo system through an optical link cable (ERGO Optical, BioSemi). A fifth-order cascaded integrator-comb low-pass filter with a −3 dB point at 410 Hz was applied to sensor channels online, after which all signals were decimated to 1024 Hz. Online high-pass response was fully DC-coupled. Participants adjusted the sound volume presentation to a comfortable listening level prior to starting the computerized tasks. Full experimental sessions lasted ~2.5 h.

Experimental protocol overview

To generate reward associations, participants were trained on the Eriksen flanker test, a response inhibition task designed to maximize gains by suppressing objects outside of the focus of spatial attention (*Eriksen & Eriksen, 1974*). At this Training stage, target and flankers were presented along with peripheral color displays which could predict earnings via controlled color-reward mappings. Unlike visual training paradigms where success is contingent on a color search display (e.g., *Anderson et al., 2011*), here participants discriminated within a constant, narrow spatial domain. The purpose of this was two-fold: to ensure that the target space remained comparable to that at the subsequent testing stage (see below), and that color-signaling spaces remained similar across stages.

A main AV task was subsequently used as a Testing stage to investigate subject performance and neural responses in the presence of reward-associated but irrelevant color distractors. From electroencephalography (EEG) recordings of testing stage data, modulations of temporal consistency (phase coherence) were probed via frequency tagging methods (*Grahek et al., 2021*). Prior to the main experiment, participants also performed a 2-3 minute auditory-only listening session where they were presented with long versions of the experiment sound stimuli (see Auditory stimuli, below), and instructed to listen attentively with their eyes closed. Recordings from the unimodal presentations served to inform the spatial distribution associated to auditory-evoked activity across listeners.

Main experimental sessions consisted of an initial Training task (Tr1) followed by an initial AV Testing task (Tt1), both of which represented baseline conditions without reward learning (**Figure 1A**). To address reward learning effects, participants thereafter performed a second round of Training (Tr2) and Testing (Tt2) tasks with reward feedback. Presentation and response logging were performed with PsychoPy (*Peirce, 2007*) for the Training tasks and with MATLAB R2010a (Natick, United States) for the Testing tasks.

Training tasks (Tr1 & Tr2)

Protocol and stimuli. Participants were asked to quickly and accurately report the direction of a central target (**Figure 1B**). Each task consisted of 216 trials with balanced congruent, incongruent, and neutral flanker conditions. A static colored ring display was shown at the start of a trial over a black background, with an internal[external] diameter subtending 21[27] degrees of visual angle (dva). Color displays were either solid red (#FF0000), green (#00FF00), blue (#0000FF), cyan (#00FFFF), magenta (#FF00FF), or yellow (#FFFF00), distributed by 36 trials per color. After an initial 200 ms of ring presentation, the task's target and flanker symbols appeared in white, subtending 6 dva horizontally. All elements were shown for a further 940 ms, after which the ring disappeared. The symbols remained for an additional 50 ms, and there was a final 1.5 s response timeout period. Participants responded to the direction of the central symbol by pressing a keyboard button. Correct and incorrect responses were treated as valid trials whereas missed/late responses were treated as invalid.

Reward structure. Points were awarded for valid and correct trials, by amounts determined by ring color as follows: a flat 80[20] points reward was given for valid responses in trials representing High[Low] reward and fixed conditions. In High[Low] reward and probabilistic conditions, valid responses earned a number of points that was pseudo-randomly sampled from a normal distribution (mean = 80[20] points; standard deviation = 7 points) and rounded to the nearest integer. Finally, 'No Reward' conditions always yielded 0 points and served as a common baseline. Colors were paired with conditions following a four cohort schedule (**Figure 1C**). Invalid (but not incorrect) trials were deducted 50 points from the accumulated score. Trial conditions and colors were presented in random order. An ad lib pause was made every 72 trials, indicating the number of accumulated points so far. The order of trials was randomized for each task. Tr1 and Tr2 were identical except on feedback presentation.

Feedback. After entering a valid response participants received text feedback, but it was intelligible only at Tr2. For Tr1 (baseline), all characters appeared in scrambled order into uninterpretable strings. In consequence, reward-color mappings were not available for participants at the first stage. Pause-related feedback, and feedback from invalid trials which

resulted in point deductions, were clearly presented in both stages. In both cases participants were instructed to accumulate as many points as possible. They were informed that the points would be converted into tokens at the end of each Testing task, to be exchanged for a prize at the end of the session. Other than through feedback, participants were not explicitly debriefed about reward-color mappings.

Testing tasks (Tt1 & Tt2)

In a two-alternative forced choice (2AFC) AV task, participants were presented with trials consisting of a dual visual object flicker display in tandem with a single amplitude-modulated (AM) noise sound. The goal was to correctly associate one of the two flicker object rates with the AM rate.

Visual flicker display. The visual dual visual object flicker display consisted of a combination of two non-overlapping central (target and non-target) figure sets with peripheral distractor presentations. Central targets and nontargets. On each trial, two symbols were taken at random from a standard chess pool, i.e., king, queen, bishop, knight, rook, or pawn figures. Two replicates of each figure were shown, and flickered in phase at 9 Hz in one figure and 11 Hz in the other. Target/non-target figure designations were determined by the auditory stream (see below). The flicker rates were selected for the following technical reasons: *i*) to elicit reliable steady-state responses from visual delivery adjusted to the screen refresh rate (85 Hz) via frequency approximation (*Nakanishi et al., 2014*; Supplementary Figure 1); *ii*) for both rates to remain relatively close to the optimal low-frequency domain where auditory EEG cortical responses also lock; and *iii*) for them to be separate enough from to each other to avoid excessively difficult discrimination. The four figure locations were scattered over a circular surface extending 8 dva in diameter and shown with a steady central fixation cross (**Figure 1C**). Peripheral distractors. A peripheral ring-shaped aperture was presented with similar dimensions as in the Training task. This domain was filled with a mixture of grey and colored dots; the specific color varied per trial and was from the same color set as in Training. The ratio of grey to colored dots was fixed at 4:3 (320 grey dots), and the diameter of each dot was 0.53 dva with a density of 2.6 dots/dva^2. Critically, to prevent the potential confound that auditory activity is differentially entrained because of mismatching dynamics between targets and distractors, peripheral dot sets flickered with colored dots exclusively synced to the target set and gray dots to the non-target set (Video 1, Supplementary Figure 1). The dots changed position every 1 s, making central figures sole reliable visual cues to perform the task. Color, figure and rates were equally distributed and balanced across trials. Dots and symbols did not overlay. Chess figures remained in their same location for 1 s before changing to different locations within the delimited central area. This resulted in a highly challenging task and then for half of the participants chess figures remained at the same initial positions throughout the trial, which was accounted for as a nuisance variable (see Data and statistical analysis). Trial stimuli displays lasted 6 seconds in total.

Auditory stream. Two 6 s AM white noise stimuli were synthesized with 9 and 11 Hz sinusoidal modulation rates, respectively, at a 75% modulation depth and sampled at 44.1 KHz. For listening-only pre-experimental sessions (see Experimental protocol, above), two additional stimuli were created with the same conditions except that they were 54 s in duration. The same auditory stimuli were used across trials and participants.

Audiovisual presentation. Each AV trial presentation consisted of the visual flicker display described above plus either the 9 or 11 Hz AM noise sound, and lasted 6 s. The AM rate presented at a trial defined which of the two figure flickers was the target. The onset of the noise relative to the first video frame included a uniformly distributed delay in the $(0,1/f)$ s range, where f is the sound AM rate, and a random phase was sampled at every trial. This manipulation allowed to discard visually-locked contributions from phase-locked activity at the inter-trial level.

AV task and feedback. Participants were asked to detect in each trial the chess symbol whose flicker matched the presented sound, and explicitly instructed to ignore the colors. Subjects were encouraged to minimize eye movement by maintaining central fixation to maximize hit chances and reduce noise; eye movement was free in the task nevertheless. After the AV presentation, the two symbols presented in the trial were shown again on the screen and a click sound prompted the observer to respond by pressing a keyboard button selecting one option (**Figure 1D**). After a choice, a high- or low-pitched feedback tone then indicated whether the response was correct or incorrect. In this task, accuracy was explicitly emphasized to participants over response speed, and responses made before the click were not accepted. Symbol pairs, colors, and auditory AM rates were equally distributed over 120 trials. In each task, trial order was randomized, and the percent correct performance was calculated and shown as feedback at the end. This value was multiplied by the number of points awarded at the immediately preceding Training task, resulting in the amount of tokens earned; for instance, 2000 points gained from Tr1 could be turned into a maximum of 2 tokens at a 100% correct rate at Tt1. Participants were informed of the number of tokens accumulated so far at the end of each Testing task. Testing tasks Tt1 and Tt2 were identical except that, as an incentive to improve their performance, at the start of Tt2 subjects were offered the possibility to pick (rather than be given) from an unspecified set of award options if they over-performed Tt1. At the end of the session participants either selected, or were assigned, a chocolate bar among different styles.

Figure 1. *Design*. **(A)** The full experiment consists of two sequential, interleaved rounds of *Training* and AV *Testing* tasks designed to evaluate effects of reward learning. **(B)** At the Training stages (Tr1 and Tr2), participants accumulate points over correct trials in a standard flanker task. The colored peripheral ring may predict the range of points to be earned from correct trials. At Tr1 this information is not available (baseline) and becomes clear at Tr2 (conditioning). **(C)** Color-reward associations determine mean reward amounts and range, and are determined by cohort assigned to participants (see Materials and Methods). For instance, cohort 3 participants receive on average 80 points on correct responses if a cyan ring is also shown, and the realized amount varies across such trials. Participants do not earn points from baseline trials. **(D)** Participants are tested in a 2AFC AV task before and after conditioning, (Tt1 and Tt2, respectively), which contains a dynamic 6-second AV display consisting of an AM noise stream plus two visual symbol sets flickering at different rates. The goal is to determine which of the two sets flickers to the same sound modulation rate (9 or 11 Hz). Also included is a peripheral ringed dot pattern with flickering grey and colored dots. At Tt2, colored dots may signal reward associations learned from Tr2. See Video 1 for a full AV trial example or Supplementary Figure 1 for a frame-by-frame video sequence.

Data preprocessing

Data analysis was implemented in MATLAB 2018b. EEG recordings from pre-experiment (auditory-only) sessions were downsampled to 1024 Hz, and to 512 Hz from main sessions. Head and reference sensor datasets were common average-referenced, after which DC offset was removed per channel. Signals were bandpass-filtered between 1 and 40 Hz with a 10-

order elliptic IIR filter of 60 dB attenuation and 0.5 dB passband ripple. Single channel data were rejected in a blind manner according to a channel variance-based criterion (*Junghöfer et al., 2000*) with confidence coefficient $\lambda_p=4$, and the same procedure was performed separately for the external reference channels. To reduce artifact contamination, EEG data were decomposed into independent components using FastICA (*Hyvärinen, 1999*). The two independent components with maximal broadband power were automatically selected and projected out of the data. A time-shifted principal component analysis (*de Cheveigné & Simon, 2007*) was applied to discard environmental signals recorded on the external reference sensors (±4 ms shifts). A sensor noise suppression algorithm (*de Cheveigné & Simon, 2008*) was applied in order to attenuate artifacts specific to any single channel (63 neighbors). After epoching, the variance-based rejection procedure was again applied across single channels and epochs' timeseries per participant, resulting in less than 2.9% rejected channel-epoch timeseries form main sessions on average (subject range 2.7%–3.2%).

Data and statistical analysis

Behavioral data. In Tr1 and Tr2 blocks, hit rates and response times were obtained per participant. In Tt1 and Tt2 blocks, d-prime (*d'*) scores indexing sensitivity to the audiovisual discrimination task were computed as the *z*-transform of the hit rates minus the *z*-transform of false positives to 9 Hz auditory stimulus presentations. *z*-scores were obtained by the normal cumulative distribution inverse function approximation, and *d'* values were obtained per condition, per subject. To ensure convergence, all hit rates and false alarm scores were approximated by the log-linear approach (*Stanislaw & Todorov, 1999*).

Statistical analysis of behavioral data. All statistical analyses were performed with MATLAB (R2018a) (The MathWorks, Natick MA), both with built-in functions as well as custom-written code. Effects were statistically assessed for confirmatory testing using linear mixed-effect (LME) models with the MATLAB function `fitlme`. The omnibus LME analyses evaluated the effects of independent ordinal variable "Stage" (Testing task 1 and Testing task 2), numeric variable "Reward" level (High, +80 points average; Low, +20 points; and No Reward, +0 points), and numeric variable "Uncertainty" (Probabilistic, 7 points standard deviation; Fixed, 0 points s.d.) on the dependent numeric variable (I: hit rate at Training; II: reaction time at Training; III: *d'* sensitivity at Testing). For Training tasks, the relevant "Flanker" variable (congruent, neutral, incongruent) was also included.

For Training data (*N*=32), the full model includes 16 fixed effects (Stage, Reward, Uncertainty, Flanker, and all higher-order interactions). To account for repeated measures within participants, the model includes random effects (intercept and slope) for each participant. In Wilkinson notation, the full general model is expressed as: *y* ~ Stage × Reward × Uncertainty × Flanker + (Stage × Reward × Uncertainty × Flanker | participant).

For Testing data (*N*=31), the full model includes 8 fixed effects (same as above except Flanker, interactions are therefore restricted to Stage:Reward, Stage:Uncertainty, Reward:Uncertainty, Stage:Reward:Uncertainty). Similarly to Training data, the model includes random effects (intercept and slope) for each participant in the between-subjects design involving two task difficulty levels as nuisance variable. The full general model is

expressed as: y ~ Stage × Reward × Uncertainty + (Stage × Reward × Uncertainty | participant/difficulty).

Where a significant interaction was found, post hoc tests were carried out on with a Bonferroni-Holm correction to account for multiple comparisons. For reward learning effects observed at Tr2 or Tt2, subject-level association strength analyses were additionally performed. These analyses served to examine and illustrate the individual dependence of observers' behavior with reward level as predictor. Beta (slope) parameters obtained by linear regression on individual subject data were then submitted to Student's *t*-testing to evaluate the hypothesis that reward associations are established, i.e., association strength is significantly different from zero.

Audiovisual locking data analysis. We probe whether reward learning impacts the stability of locking responses to audiovisual temporal at the single trial level, under both probabilistic and fixed modes of learning. EEG channel data from Tt1 and Tt2 blocks were epoched by 6 s AV presentation segments, sub-epoched into 1 s partitions, then transformed into the frequency domain via a Fast Fourier Transform, resulting in six sub-epochs spectra per trial with 1 Hz resolution. Phase coherence analysis (*Nozaradan et al., 2012*; *van Diepen & Mazaheri, 2018*) was used to measure temporal and spectral phase synchronization in EEG signals at the single trial level, via circular averaging of the corresponding phase angle timeseries across sub-epochs spectra. Resulting intra-trial phase locking values (PLV) were used to evaluate the stability of steady state responses in each trial as they become modulated by temporal lapses of attention (*Keitel et al., 2019*). PLVs were obtained by estimating peak sizes of phase coherence spectra at the target auditory stimulation rate plus the first and second harmonics (i.e., 9, 18, and 27 Hz in 9 Hz trials; alternatively, 11, 22, and 33 Hz). Harmonics were included (**Figure 2A**) as they may contain non-redundant neural representations that complement those directly linked to the base rate (*Jenkins et al., 2011*; *Keitel & Müller, 2016*). Each PLV peak size was expressed as the ratio of the coherence at the target frequency bin divided over the average coherence at the two immediate neighbor bins. The PLV for a trial was defined as the average of the three sound-related peak frequency ratios across nine auditory-dominant (Fz, FC1, FCz, FC2, C3, C1, Cz, C2, C4, see **Figure 2B**), or when indicated visual-dominant (PO7, PO3, O1, POz, Oz, Iz, PO4, O2, PO8), EEG channels and expressed in decibels.

Figure 2. *Single-trial estimation of temporal locking precision during AV scene presentations.* **(A)** Intra-trial phase coherence spectra averaged across trials, conditions and channels, by sound modulation rate. AV presentations elicit characteristic frequency patterns in participants' EEG responses, with significant PLV differences at the sound modulation rate and two harmonics (adjusted *p*-values ≤ 0.004). **(B)** Precision of temporal locking measured by the size of PLV peaks, estimated by referencing the crest phase coherence value relative to the immediate neighboring bin frequencies. Expressed in dB, the average of these peak-to-baseline ratio estimates appears specific to the auditory stimulation rate (left). Error bars indicate ±1 SEM. Topographically, peak sizes show relatively precise temporal locking over the frontocentral scalp of the EEG array, in addition to occipital areas (right). The highlight indicates channels that were used to index AV locking to the auditory rate in subsequent analyses, unless otherwise indicated.

Auditory locking data and statistical analysis. Finding of reward learning impacting precise temporal locking of audiovisual responses in auditory sites at Tt2 (see Results) prompted further investigation of the impact of learned reward effects on unimodal responses. In particular, we aimed to evaluate whether reward learning and performance effects similarly hold for auditory responses to the sound stream at this stage. Importantly, recordings of intra-trial coherent activity at the auditory channels include contributions to PLV from cross-modal activity locked to both visual flicker rates ([Figure 2A](#)). This is a consequence of visual and auditory streams maintaining a consistent phase lag within each trial, overall contributing to consistency across trial sub-epochs; AV lag is however variable between trials. Next, we evaluated phase consistency of auditory-locked activity devoid of visual contributions by using an inter-trial approach in combination with spatial source separation.

First, a spatial filter was constructed to emphasize reproducible auditory processing across subjects using an implementation of joint decorrelation (JD, *de Cheveigné and Parra, 2014*; *de Cheveigné and Simon, 2008*). The spatial filter weights were estimated simultaneously from all listeners' recordings of the 'auditory-only' pre-experiment session, each organized in two 54 s epochs corresponding to 9 and 11 Hz presentations. Data were submitted to a first round of JD using as bias data the average across listeners in the two separate presentations. From the resulting spatial decomposition, the subspace resulting with the highest stimulus-evoked activity ratio across the two stimuli was extracted. This reproducible subspace, composed of the top 3 components, was projected back into sensor data, treating less reproducible activity across listeners as noise. Next, denoised unisensory data were filtered using an order-2 Butterworth filter in the 8–10 (for 9 Hz stimuli) or 10–12 Hz (11 Hz stimuli) bandpass region and averaged across participants. Denoised sensor data were submitted to a second round of JD using as bias data the filtered average across listeners in the two separate presentations. The most repeatable (top) component was consistent with auditory source activity and used together with the first step to decompose. The procedure provided two transformations: one to remove the spatial components least likely to reproduce across listeners, and another to optimize the signal to noise ratio of isolated stimulus-evoked narrowband activity (*de Cheveigné and Parra, 2014*) from the auditory system.

Second, the above transformations obtained from listening-only data were then applied to AV data. Specifically, 1) the first JD matrix was applied to single-trial Tt1 and Tt2 data; 2) the top 3 components were kept and projected back into sensor space; 3) the second JD matrix was applied to the denoised data; and 4) the top component was used for auditory tracking analyses. As only linear weightings of sensor data are applied, narrow-band spectral filtering of the main experiment epochs was not imposed. After obtaining auditory source estimates, inter-trial phase coherence (ITPC) measures were applied across trials. ITPC measures are sensitive to sample size difference biases, and were thus computed for each subject at 20 trials per reward level condition and subject. Because visual flicker phase lags varied from trial to trial relative to sound AM phase, the procedure effectively discarded visual oscillatory contributions to phase variability. In this case, only the first harmonic of each fundamental rate resulted in an obvious peak across listeners and conditions, phase variability analyses of auditory locking therefore did not include the second harmonic. For each reward level and subject, ITPC spectrum data (N=30) were collected at the sound presentation AM rate and its first harmonic, across both AM conditions in the Tt2 stage. ITPC data were regressed against reward level and resulting beta parameters were submitted to Student's *t*-testing.

Results

We examined participants on a two-alternative forced choice (2AFC) AV coherence task before and after a reward conditioning visual procedure (**Figure 1**). The multistage process was used to determine whether the temporal precision tracking AV modulations during the 2AFC task (**Figure 2**) may be influenced by reward associations established through conditioning.

Reward manipulation effectively incentivizes observer behavior during conditioning

To first determine whether the reward association was established, we inspected subject behavior before and during conditioning. Both training stages, consisting of a standard flanker's task with color cues (**Figure 1**), resulted in performance that was assessed by linear mixed effects modeling. Hit rates and, separately, mean reaction times analyzed by the factors 'Flanker' (standard congruent, neutral, incongruent conditions), 'Stage', 'Reward', 'Uncertainty', and their interactions, first showed an effect of 'Flanker' condition in both hit rate ($F_{2,936} = 9.83$; $p < 0.001$) and reaction time data ($F_{2,996} = 67.6$; $p < 0.001$), as is expected for this task. While hit rates were not further significantly modulated by any other factor or interaction ($p > 0.09$; Supplementary Table 1), mean reaction times showed a significant main effect of 'Stage' ($F_{1,996} = 11.5$; $p < 0.001$). This main effect was qualified by (*i*) a significant interaction with 'Flanker' ($F_{2,996}=4.59$; $p = 0.010$); (*ii*) a significant interaction with 'Reward' ($F_{1,996} = 5.08$; $p = 0.024$; Supplementary Table 2). Addressing the significant 'Stage' by 'Flanker' interaction, post-hoc analyses revealed that responses to neutral and incongruent flanker trials were faster in the second training stage (during conditioning, Tr2) compared to the first (before, Tr1; Supplementary Table 3). Post-hoc analyses conducted to address the significant 'Stage' by 'Reward' interaction additionally revealed that responses to Low (20 point average reward) and High reward trials (80 point average) were faster in Tr2 compared to Tr1 (Supplementary Table 4). Mean reaction times were not significantly modulated by any other factor or interaction in the omnibus test ($p > 0.42$).

The results suggest that participants speed up inhibitory responses as a function of learning the color payoff context, relevant to increasing task gains (**Figure 3A**), indistinctly of varying and fixed conditioning. Therefore, we expect that if reward-driven learning has been substantiated at conditioning (Tr2), then reaction times at that point may be expected to vary as a function of reward. For each participant, mean reaction time was regressed against expected reward indicated by color, evaluated at 0, 20 or 80 points, and individual association slope parameters were obtained. Faster responses to higher expected reward predict negative slopes, and a one-sided *t*-test confirmed that at this reward learning stage the average of participants' slopes was significantly lower than zero ($t(31) = −2.574$, $p = 0.008$).

Cue-dependent sensitivity changes in AV perception consistent with reward-driven distraction

Given an effective color-reward association, behavioral measures of AV discrimination sensitivity performance (*d'*) were next evaluated at the pre- and post-conditioning stages of AV testing (Tt1 & Tt2, respectively). Rather than aligning with task goals (as in the training tasks), conditioned peripheral color cues here are likely to interfere and lead observers to suboptimal performance. Participants observed the single-auditory/dual-visual streams conforming the target AV display before committing to a single visual option that matched the sound. Hence, performance measurements at Testing were not based on reaction time and were limited to sensitivity. Linear mixed effects analysis of *d'* by the factors 'Stage', 'Reward' associated with the peripheral dots, 'Uncertainty' associated with the same, and their interactions, first confirmed a significant main effect of 'Stage' on AV sensitivity ($F_{1,33} = 7.50$; $p = 0.010$), which increased from Tt1 to Tt2 (**Figure 3B**) consistent again with practice effects. This main effect was qualified by a significant interaction with 'Reward' ($F_{1,35} = 4.54$; $p = 0.040$). *d'* values were not significantly affected further by any other factor or interaction

(*p* > 0.19; Supplementary Table 5). To address the significant interaction, post-hoc analyses were conducted to evaluate the effect of stage across each reward condition, revealing a significant increase in sensitivity in Tt2 compared to Tt1 that was limited to No reward trials ($F_{1,31}$ = 9.94; adj. *p* = 0.011; **Figure 3B**); for Low and High reward trials there was no significant change (*p* > 0.9; Supplementary Table 6). To clarify whether these findings may be interpreted as reward cues interfering with optimal performance, Tt2 sensitivity data were evaluated as a function of expected reward magnitude. For each participant, mean *d'* was regressed against expected reward signaled through color distractors, and individual association slope parameters were obtained (**Figure 3C**). Weaker sensitivity under higher expected reward associations at this testing stage also predict negative slopes, and one-sided *t*-testing confirmed that the mean of participants' slopes was significantly below zero ($t(30)$ = −2.324, *p* = 0.013).

The results thus are consistent with implicit reward learning transferred over the testing stage, and overall suggest that practice benefits at the second round may be offset or not available in the presence of rewarded-associated signals. To facilitate comparison with reward-driven distraction studies (*Rusz et al., 2020*) we additionally computed the standardized mean change effect size of d' sensitivity measures at Tt2, contrasted between the High reward (mean *d'* = 0.80) and No Reward (mean *d'* = 1.05) conditions. Expressed in standard deviations, the parameter quantified how much of potential performance appears hindered by the reward cue. The estimated effect size of 0.192 result is moderate for visual reports but in line with cross-modal studies (*Rusz et al., 2020*).

Figure 3. *Behaviorally-relevant precise of locking responses to AV stimuli have stability interfered by reward-driven visual distraction.* **(A)** In a standard flankers task subjects speed their inhibitory responses as a function of training and learned color-reward mappings, indicating consolidation of reward conditioning at the Tr2 stage. **(B)** Before and after conditioning subjects perform an independent 2AFC AV correspondence task. Improved sensitivity performance at the latter Tt2 stage is contingent on lack of reward-associated color distractors. Dots denote subjects per condition (boxplot centroid: median; thick line: inter-quartile range). **(C)** Sensitivity data regressed against reward level shows decreased sensitivity at Tt2 to AV correspondence. **(D)** Neural results before and after conditioning show decreased intra-trial phase-locking to the common AV temporal modulation rate in the presence of high-reward-associated distractors, even as the latter's temporal flicker is synced with visual targets. **(E)** Phase locking data regressed against reward level shows the retreat in stable temporal encoding as a function of prior reward

context cued by distractors at Tt2. **(F)** The loss in temporal precision predicts subjects' weaker behavioral sensitivity in the AV task (reward-driven distraction). Contour indicates distance from bivariate mean; data with $D_s \geq 6$ squared z-units are defined as outliers (open circles). Shaded region: 95% confidence intervals (CI) of linear regression model prediction. **(G)** 95% CI of the empirical cumulative distribution function of single-trial PLVs show stable, precise locking as a relevant decision variable in the AV task. All single-trial data combined show that, across all conditions and stages, decreased stability of responses locked to the common temporal AV rate predict incorrect outcomes, measured at the frontocentral locations (left) but also at occipital sensors (right).

Contextual reward cues disengage stable tracking of AV objects

Next we studied how induction of reward associations changed the intra-trial stability (temporal precision) of locking responses to key modulations of the AV scenes. Scalp-wide EEG activity during these presentations responded as expected to the stimulation design (**Figure 2**). To assess the temporal precision of locking responses, we estimated the peak-to-baseline ratios (peak sizes) of EEG channels' PLV spectra at frequencies determined by auditory/target modulation rates (9, 18, and 27 Hz; alternatively, 11, 22, and 33 Hz). These estimates were found to map topographically to a distribution predominantly consistent with auditory activity (*Nozaradan et al., 2012*) and including occipital locking. To study cross-modal VDAC effects on the temporal precision of tracking responses, PLV peak size data from frontocentral auditory channels were averaged (**Figure 2B**) and analyzed, on a trial-by-trial basis, by a full linear mixed effects model as in behavioral sensitivity analyses. Analysis of PLV peak size estimates by the factors 'Stage', 'Reward', 'Uncertainty', and their interactions confirmed a significant interaction between 'Stage' and 'Reward' ($F_{1,7431} = 4.30$; $p = 0.038$) in these neural data. No other significant interaction or main effect was found (Supplementary Table 7). The significant interaction was analyzed in post hoc tests, revealing a significant decrease in PLV by 'Stage' at the High reward condition ($F_{1,2478} = 13.63$; adj. $p = 6.8\times10^{-4}$), with no evidence of change at the other reward conditions (Supplementary Table 8). The results suggest that when visual distractors cue previous reward to observers in a sustained fashion, the ability of the brain to precisely sustain temporal representations of target AV dynamics becomes suboptimal (**Figure 3D**).

As with behavioral data, we addressed whether the neural changes that were transferred via conditioning to the AV task may reflect reward magnitude. Single-trial Tt2 phase locking data were investigated as a function of expected reward signaled by color distractors, and individual association slope parameters were obtained (**Figure 3E**). Weaker phase locking by increasing expected reward associations similarly predict negative slopes in this case. One-sided *t*-testing confirmed that the average of participants' slopes was significantly below zero ($t(30) = -3.763$, $p = 3.7\times10^{-4}$). The results indicate that visual distractors decreased the temporal precision of phase locked AV responses at frontocentral scalp, via their past associations with reward level.

Cue-dependent loss of stable tracking indexes observers' reward-driven distraction

The finding of reward association effects at both behavioral and neural levels motivated the hypothesis that both may be related. We examined whether, at a subject level, the scale of changes to temporal encoding evidenced by PLV predicted the behavioral performance differences. Effect sizes (in *d'* and PLV) were defined by the separation between the High

and No Reward conditions. At the Tt2 stage, participants' PLV effects were significantly correlated with their behavioral differences (Spearman's $\rho = 0.473$, $[0.113 - 0.735$ C.I.$]$; $p = 0.007$). Steeper losses of precise temporal locking in a participant predicted their greater loss of sensitivity in the task (**Figure 3F**). The correlation was robust to potential outliers (Shepherd's Pi = 0.529; $p = 0.010$), and a robust regression analysis additionally confirmed its significance ($t=2.48$; d.f.=29; $p = 0.019$). By contrast, Tt1 stage (pre-conditioning) data failed to reveal a significant correlation between neural and behavioral effects ($\rho = 0.228$; $[-0.171 - 0.556$ C.I.$]$; $p=0.22$) in full, or after the removal of potential outliers (Pi = 0.159; $p = 0.82$).

Temporally stable selective tracking is a neural index of correct AV stream integration

The previous result suggests that visual reward-driven distraction may interfere with the tracking processes used to establish reliable temporal associations within the AV scene. VDAC deficits have been shown to reflect adverse sensory encoding or weak sensory evidence integration in at least sensory cortical regions (*Hickey & Peelen, 2015*; *Pearson et al., 2022*). While our findings are consistent with spontaneous out of phase fluctuations within a trial that negatively impact perceptual decisions, they do not exclude the contribution by dedicated attention to the non-target flicker into less reliable decision-making. Such scenario may tie the loss of temporal precision to varying top-down selection under different reward cue contexts, rather than to reward-driven distraction directly, with selectivity being a primary predictor of outcome. To dissect the link of precise temporal locking to perceptual decisions, we examined whether this neural index determined trial outcomes in relation to reward learning. For this, we assessed whether reward learning, i.e, the statistical interaction between 'Stage' and 'Reward', varies according to the 'Outcome' of a trial (correct or incorrect) through an exploratory factorial design (see Materials and Methods). The test involved a LME model run on the same single-trial PLV data which included the trial factors 'Stage', 'Reward', and 'Outcome' of the result (correct versus incorrect response). If a trial's successful outcome is supported by precise tracking of target modulations, a main effect of 'Outcome' may be expected. Temporal precision that is particularly affected in misses consistent with VDAC rather than misses generally, including those from attending nontargets, may be additionally shown through a triple interaction.

Confirming the expected interaction between 'Stage' and 'Reward' in the PLV data ($F_{1,7431} = 10.15$; $p = 0.001$), the analysis further revealed a significant main effect of 'Outcome' ($F_{1,7431} = 6.69$; $p = 0.010$). The effect was due to trials leading to correct responses showing greater PLVs than those leading to incorrect outcomes, across all experimental conditions, (**Figure 3G**). Nevertheless, no other significant interaction or main effect was observed (Supplementary Table 9). The findings indicate that down-modulations on PLV may predict incorrect trial outcomes generally, including but not limited those linked to VDAC. Through reward learning VDAC may effectively decrease PLV, regardless of whether this drop eventually leads to a correct or incorrect trial. The results therefore appear consistent with bottom-up interference forming temporally precise bimodal neural representations during the AV perceptual decision-making.

Auditory and supramodal activity is associated with loss of stable tracking elicited through VDAC

So far, stable locking to the target rate in frontocentral scalp within a trial may predict decision outcome. Because (*i*) locking changes by the visually cued VDAC may be present even in successful trials; and (*ii*) color distractors may not necessarily undermine visual temporal locking because they are strictly target-synced, we then evaluated activity from occipital regions. Such activity is also expected to reflect more directly the arbitration of the observer's time-varying top-down selection across the dual flicker of central objects (*Kim et al., 2007*). Importantly, querying visual activity may inform us on the source of VDAC effects on temporal locking as the modality of origin conveying the reward-associated signals. Single-trial visual selection PLVs from occipital electrodes (see Methods), were submitted to the same statistical model as above. The analysis similarly showed a significant main effect of 'Outcome' ($F_{1,7431} = 25.4$; $p = 4.7 \times 10^{-7}$), indicating greater temporal locking to target rates in correct versus incorrect trials. This result confirms the impact that precise temporal representations of the target rate have on task performance across conditions and stages. Unlike frontocentral activity nevertheless, a significant interaction between 'Stage' and 'Reward' was not observed in these responses ($F_{1,7431} = 1.54$; $p = 0.21$; Supplementary Table 10). The lack of evidence may appear consistent with observers' unchanged visual locking when spatial attention deflects away from targets to peripheral reward-associated color distractors, which are in phase with the central target. Moreover, it raises the question that negative VDAC influences on reliability may appear within supra-modal locking or even at unimodal auditory locking.

To illustrate the spatial profile associated with the temporal locking effects associated with reward learning and outcomes, we performed single channel analyses of single-trial PLV data from which the spatial topographies of associated *F*-values could be derived. Using these maps, the analysis confirmed that the signal-to-noise ratio of 'Outcome' effects was optimal at the occipital areas (**Figure 4A**). This is consistent with the ability of participants to successfully localize and focus their visual responses, entrained themselves to the single target rate that. Remarkably, 'Stage' by 'Reward' interaction effects were shown to peak instead only within a limited subset of electrodes at frontocentral scalp locations (**Figure 4B**), consistent with in supramodal and/or auditory locking responses (**Figure 2B**). This result may suggest that sound-entrained, rather than flicker-entrained responses were primarily susceptible to modulation by VDAC.

Figure 4. *Reward-associated visual distractors decrease the temporal precision of auditory phase-locked activity.* (A) Topography of repeated measures ANOVA *F*-values on single channel data, illustrating the localized strength of two relevant effects, reward learning and trial outcome. Largest effects of Outcome on single-trial sound/target rate PLVs are shown on occipital sensors (left), underscoring the relevance of selection of these rates for successful behavior. By contrast, the largest reward learning effect (Stage by Reward interaction) features a cluster of frontocentral electrodes (right), suggesting direct influence on auditory or supramodal phase locking. **(B)** Topography of the spatial component used for source

separation of AV data, trained from reproducible and narrowband activity that was obtained from additional auditory-only recordings of the same participants. **(C)** Unimodal auditory locking activity from Tt1 & Tt2 AV presentations, extracted via spatial source separation in combination with the inter-trial phase coherence method. Phase locking is clearly apparent at the sound AM rate plus a harmonic (dashed lines) exclusively. **(D)** To evaluate the influence of reward-associated visual distractors, auditory ITPCs were regressed against reward level, demonstrating the decreasing effect by expected reward from color cues on the inter-trial temporal precision of sound tracking responses. **(E)** Conceptual illustration. *Left*: Target flicker stimuli match the AM in rate but have random phase. Because of this, associated unimodal visual responses will decohere when aligning responses by the acoustic phase. Noisy physiological responses, represented in phase diagrams, result in a relatively compact phase-locking distribution for unimodal auditory but which is absent in unimodal visual responses. *Right*: The observer's behavior accommodates shifts of attentional sampling among the two candidate flickers in the dual visual stream. Visual attention to the correct flicker rate involves either the central object, in line with task requirements (top), or the periphery through reward-driven distraction (bottom). In the latter case, AV tracking supporting cross-modal integration decreases, suggesting intermodal competition (highlight).

Cue-dependent loss of temporal precision occurs within unimodal auditory tracking

In evaluating the hypothesis that auditory locking may lose temporal precision under VDAC, we acquired unimodal recordings of AM noise stimulation in the same listeners, which was used to train a series of spatial source separation filter (see Materials and Methods). Moreover, in each AV presentation during Tt1 and Tt2, the distractor flicker systematically varied in phase relative to the sound stream. Across-trial phase comparisons offer therefore an additional means to remove activity evoked by the dual visual flicker. Submitting unpartitioned (six-second) single-trial responses to source separation, in combination with inter-trial phase coherence (ITPC) estimation, first confirmed the absence of dual visual flicker responses (**Figure 4C**). Unimodal auditory locking, represented by peak ITPC values at the relevant AM stimulation frequencies (base plus first harmonic rates), was evaluated for the strength of reward associations by using linear regression. Here, loss of precise auditory locking under higher expected reward associations similarly predict negative slopes. Individual association slope parameters were obtained per participant (**Figure 4D**), and one-sided *t*-testing confirmed that the mean of participants' slopes was significantly below zero ($t(29) = -2.467$, $p = 0.010$). The result is consistent with the notion that continuous cross-modal signaling of reward cue associations weaken the reliability with which temporal patterns are encoded in unconditioned unimodal cortex. Specifically here, visually carried reward signals operating as distractors directly decrease the quality of auditory entrainment to stimuli. This inter-modal suppressive effect appears to compromise the operation of low-level AV binding processes that are based on detecting common temporal dynamics.

Discussion

We interact with our surroundings more effectively when we integrate the incoming flux of visual and auditory signals into veridically coherent audiovisual representations. How do past reward associations reach and tap into the reliable AV processing has been unknown. Here, we demonstrate visual dynamic contexts with conditioned signals that affect temporally reliable cortical responses. Taking the chances to sample reward cues means the brain reduces the precision with which cortical networks track temporal modulations in the non-

cuing sense. The re-balanced representation impacts the sensitivity observers use to detect AV correspondences in busy scenes.

Aside from the reward system, visual VDAC engages the frontal eye field and posterior parietal cortical areas, where plasticity effects underpin spatial priority mappings (*Anderson, 2019*; *Anderson et al., 2021*; *Tankelevitch et al., 2020*; *Bisley & Goldberg, 2010*; *Markovic et al., 2014*; *Kim et al., 2021*; *Peck et al., 2009*). These parietal regions receive projections from norepinephrine networks which encode stimulus significance and mediate shifts in the transmission of information according to behavioral relevance (*Sara, 2009*; *Todd & Manaligod, 2018*; *Mather et al., 2016*). Transmission of reward salience across modalities is conveyed under temporal priming (*Anderson, 2016*), spatial co-localization (*Pooresmaeili et al., 2014*), or semantic coincidence (*Sanz et al., 2018*). Our present results now implicate cross-modal associations based on common temporal structure. Reward salience triggered less stable audiovisual tracking activity, both in correct (audiovisually integrated) and incorrect trials, consistent with systematic lapses in temporal sampling of multimodal targets. After excluding unimodal visual contributions, our results furthermore identified the weakened precision of auditory locking. This was striking as the auditory stream shared the same periodicity as the rewarded distractors. The finding therefore suggests that recurrent lapses in attentional sampling originating from VDAC induce a form of inter-modal competition which recurs over time.

Learned value associations modulate sensory cortical responses, altering the precision of target input representations from the same or different modalities as the value-associated distractor (*Itthipuripat et al., 2019*; *Kim et al., 2021*; *Pooresmaeili et al., 2014*; *Antono et al., 2023*; *Stănişor et al., 2013*). When distractors are from a different modality, these bottom-up effects are observed relatively later (*Vakhrushev et al., 2023*; *MacLean & Giesbrecht, 2015*; but see *Tankelevitch et al., 2020*), yet earlier than some cross-modal influences operated through top-down attention (*Cervantes Constantino et al., 2023*). Longer latencies when multiple modalities are involved reflect less direct connectivity pathways in place between early sensory and frontal valuation areas. Specifically, both intra- and cross-modal VDAC modes appear mediated by the anterior intraparietal sulcus (IPS), but only in the cross-modal case communication between IPS and early sensory areas appears intermediately relayed through association areas of the superior temporal sulcus (STS, *Antono et al., 2023*).

Cross-modal VDAC manipulations have been typically studied through spatial associations. STS-IPS link findings from spatial association paradigms also appear of interest here due to its hypothesized role in the spatiotemporal transformations required to establish AV correspondences in the brain (*Miller and D'Esposito, 2005*). The IPS is involved in cross-modal interactions and auditory-to-visual entrainment (*Bauer et al., 2020*, *2021*; *Lakatos et al., 2009*), as well as estimates of audiovisual congruency based on normative spatial inference (*Rohe and Noppeney, 2015*). STS rather appears highly sensitive to AV temporal congruency, and often feeds back into early sensory areas (*Noesselt et al., 2007*; *Lewis and Noppeney, 2010*). Temporal synchrony effects observed in STS similarly feature in sensory areas, likely through bottom-up coherence detection (*Atilgan et al., 2018*). The comparatively visually dominant parietal versus auditory-dominant temporal association areas (*Werner and Noppeney, 2010*) is consistent with specialized STS populations to process AV temporal associations in a relatively automatic manner, and IPS to compute

mappings based on behaviorally relevant AV spatial priority (*Noppeney, 2021*; *Rohe and Noppeney, 2016*). Given these differences it is plausible that reward-associated signals in one sense use more than one pathways to reach another unisensory area. Among minimally necessary pathways in AV scenes, we speculate that STS-mediated interactions are required in cross-modal VDAC when target streams are causally connected through temporal correspondences alone. Streams with spatiotemporally common causes may result in reward bias transfer requiring IPS-mediated spatial transformations or inference. Future neuroimaging work using conjunction analyses will be required to confirm that dynamic re-priorization distinctively operates in these regions and the underlying mechanisms.

When reward-associated distractors are placed apart from task targets, they increase task demands with interference effects (*Tankelevitch et al., 2020*). By contrast, sharing the same target location typically boosts performance (*Pooresmaeili et al., 2014*; *Antono et al., 2023*; *Grahek et al., 2021*; see reviewed in *Rusz et al., 2020*). In the present design, deflecting overt or covert spatial attention from targets to the periphery met colored signals which continued the visuotemporal dynamics of target objects. Our findings of reduced (rather than sustained) locking appear consistent with 'de-sharpening' of neural representations of sensory target input when salient reward cues are present (*Hickey & Peelen, 2015*). Neural representations of distractor input signaling value do not appear to be systematically enhanced, however. Rather, VDAC biases may reflect the dedication of task resources to compute relative value (*Anderson, 2019*; *Kim & Beck, 2020*; *Liao & Anderson, 2020*; *Anderson et al., 2021*). Doing so may release from inhibition some visual cortical regions encoding salient distractor representations, with the effect that spatial and object-based target selectivity decreases (*Anderson et al., 2021*; *Wang et al., 2015*; *Hickey & Peelen, 2015*; *Stănişor et al., 2013*).

Gaining salience also modulates spectral components of neural activity. Cortical representations of visual stimuli exhibit more gamma-band power when stimuli appear more physically salient (*Fries, 2015*). In the STS, such activity increases are associated with a greater likelikood of perceptual binding of AV stimuli (*Mishra et al., 2007*; *Balz et al., 2016*), where in this case salience operates through redundancy gain. Conditioning is another means to boost perceptual salience of stimuli (*Qin et al., 2021*). Salience is more generally hypothesized in relation to upkeeping distractors' persistent gamma activity power and, possibly, temporal coding over attentional engagement of variable strength elsewhere (*Jensen et al., 2012*). The notion suggests implications for temporal phase coding during attentional reorienting guided by incentive salience. Outbound engagement from the target to the salient reward cue plausibly increases temporal uncertainty in target-following temporal responses at unimodal regions, or within supramodal areas, as target-specific coding moves to a subordinate temporal rank in the attentional cycle (e.g., *Jensen et al., 2012*). In the STS, such changes may be expected at unimodal visual dominant member populations, less so at auditory dominant or AV bimodal dominant populations (*Beauchamp et al., 2004*). The variable sub-threshold excitability dynamics driven by salience and attentional orienting, in combination with suprathreshold activity driven by stimulus-following responses across these sub-populations, may determine propagation of integrated or else segregated feedback to sensory networks (e.g. *Bauer et al., 2020*; *Fries, 2015*; *Keil & Senkowski, 2018*). We suggest that the interplay prompted by incentive salience promotes inter-modal competition: it creates recurrent instances of routing lower reliability to unimodal representations of unconditioned streams despite temporal redundancy cues (**Figure 4E**).

The breakup of cross-modal binding in favor of inter-modal competition opens the question of the natural mechanisms by which incentive salience engages dynamic attentional sampling. Sustained top-down visuospatial attention is hypothesized to fluctuate rhythmically on regular sampling periods committed to a target, interleaved with disengagement that allows exploratory shifts (*Fiebelkorn & Kastner, 2019*; *Lakatos et al., 2016*; *Landau & Fries, 2012*; *Todd & Manaligod, 2018*; *Chen & Gong, 2022*). The latter periods represent temporary blind spots, referred to as 'down' phases of the attentional cycle, and may be harnessed by the visual system to scan potentially relevant parts of the scene. The dynamic sampling framework appears to accommodate findings that reward cue signaling increases the likelihood of exploratory oculomotor behavior (*Rusz et al., 2020*). Recurrent exploitation of down phases may be warranted by incentive salient distractors and their regular position changes. Securing these periods consistently therefore offer a regular means to sustain reward-driven distraction across time. The cost is a reliable inference of the scene's temporal causal structure, compromised by recurring diminished sensitivity to central objects and the auditory stream, facing relatively expedited peripheral processing (*Jovanovic & Mamassian, 2020*). The tradeoff may expose limitations to low-level continuous audiovisual integration as both visual and auditory time in the brain accord with dynamic attentional and physiological balances.

Conclusion & impact statement

To see and hear things playing along means connecting them with confidence if they belong together. Many aspects of this function are tied with learning. Reward experience can instruct us valuable ways to learn but also creates bias in others. This study presents a type of learning bias, value driven attentional capture, that diverts from making reliable and timely linking of audiovisual information in the brain useful for observer success. The effect persistence puts into question the neural resources and organization set to track and bind multimodal cues that we attend to. These basic perceptual processes appear downgraded in the sensitized brain, a state which favors unimportant but salient cues that repeatedly sign a prominent incentive history.

Data and code availability

Data and code supporting the findings of this study are openly available at the Open Science Framework at http://doi.org/10.17605/OSF.IO/SPJ9F.

Author contributions

Rodrigo Caramés Harcevnicow: Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization **Thaiz Sánchez-Costa**: Investigation, Data Curation, Writing - Review & Editing **Alejandra Carboni**: Software, Resources, Writing - Review & Editing, Supervision, Funding acquisition **Francisco Cervantes Constantino**: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration, Funding acquisition

Funding

The research presented in this manuscript received funding from the Agencia Nacional de Investigación e Innovación, Uruguay, under the code PD_NAC_2018_1_150365.

Declaration of Competing Interests

None.

Acknowledgments

We thank Alex L. White and Germán A. Cipriani for code sharing, and Franco Caballero and Cecilia Biurrun for assistance with data collection.

References

Anderson, B. A. (2016). Value-driven attentional capture in the auditory domain. *Attention, Perception & Psychophysics*, *78*(1), 242–250. https://doi.org/10.3758/s13414-015-1001-7

Anderson, B. A. (2019). Neurobiology of value-driven attention. *Current Opinion in Psychology*, *29*, 27–33. https://doi.org/10.1016/j.copsyc.2018.11.004

Anderson, B. A., Kim, H., Kim, A. J., Liao, M.-R., Mrkonja, L., Clement, A., & Grégoire, L. (2021). The past, present, and future of selection history. *Neuroscience & Biobehavioral Reviews*, *130*, 326–350. https://doi.org/10.1016/j.neubiorev.2021.09.004

Anderson, B. A., Laurent, P. A., & Yantis, S. (2011). Value-driven attentional capture. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(25), 10367–10371. https://doi.org/10.1073/pnas.1104047108

Antono, J. E., Dang, S., Auksztulewicz, R., & Pooresmaeili, A. (2023). Distinct patterns of connectivity between brain regions underlie the intra-modal and cross-modal value-driven modulations of the visual cortex. *Journal of Neuroscience*, *43*(44), 7361–7375. https://doi.org/10.1523/JNEUROSCI.0355-23.2023

Atilgan, H., Town, S. M., Wood, K. C., Jones, G. P., Maddox, R. K., Lee, A. K. C., & Bizley, J. K. (2018). Integration of Visual Information in Auditory Cortex Promotes Auditory Scene Analysis through Multisensory Binding. *Neuron*, *97*(3), 640–655.e4. https://doi.org/10.1016/j.neuron.2017.12.034

Balz, J., Keil, J., Roa Romero, Y., Mekle, R., Schubert, F., Aydin, S., Ittermann, B., Gallinat, J., & Senkowski, D. (2016). GABA concentration in superior temporal sulcus predicts gamma power and perception in the sound-induced flash illusion. *Neuroimage*, *125*, 724–730. https://doi.org/10.1016/j.neuroimage.2015.10.087

Bauer, A.-K. R., Debener, S., & Nobre, A. C. (2020). Synchronisation of Neural Oscillations and Cross-modal Influences. *Trends in Cognitive Sciences*, *24*(6), 481–495. https://doi.org/10.1016/j.tics.2020.03.003

Bauer, A.-K. R., Ede, F. van, Quinn, A. J., & Nobre, A. C. (2021). Rhythmic Modulation of Visual Perception by Continuous Rhythmic Auditory Stimulation. *Journal of Neuroscience*, *41*(33), 7065–7075. https://doi.org/10.1523/JNEUROSCI.2980-20.2021

Beauchamp, M., Argall, B., Bodurka, J. Duyn, J. H., & Martin, A. (2004). Unraveling multisensory integration: patchy organization within human STS multisensory cortex. *Nature Neuroscience*, 7, 1190–1192. https://doi.org/10.1038/nn1333

Bisley, J. W., & Goldberg, M. E. (2010). Attention, intention, and priority in the parietal lobe. *Annual Review of Neuroscience*, *33*, 1–21. https://doi.org/10.1146/annurev-neuro-060909-152823

Bizley, J. K., Maddox, R. K., & Lee, A. K. C. (2016). Defining Auditory-Visual Objects: Behavioral Tests and Physiological Mechanisms. *Trends in Neurosciences*, *39*(2), 74–85. https://doi.org/10.1016/j.tins.2015.12.007

Cervantes Constantino, F., Sánchez-Costa, T., Cipriani, G. A., & Carboni, A. (2023). Visuospatial attention revamps cortical processing of sound amid audiovisual uncertainty. *Psychophysiology*, e14329. https://doi.org/10.1111/psyp.14329

Chen, G., & Gong, P. (2022). A spatiotemporal mechanism of visual attention: Superdiffusive motion and theta oscillations of neural population activity patterns. *Science Advances*, *8*(16), eabl4995. https://doi.org/10.1126/sciadv.abl4995

Cho, S. A., & Cho, Y. S. (2021). Uncertainty modulates value-driven attentional capture. *Attention, Perception, & Psychophysics*, *83*(1), 142–155. https://doi.org/10.3758/s13414-020-02171-3

de Cheveigné, A., & Parra, L. C. (2014). Joint decorrelation, a versatile tool for multichannel data analysis. *NeuroImage*, *98*, 487–505. https://doi.org/10.1016/j.neuroimage.2014.05.068

de Cheveigné, A., & Simon, J. Z. (2007). Denoising based on time-shift PCA. *Journal of Neuroscience Methods*, *165*(2), 297–305. https://doi.org/10.1016/j.jneumeth.2007.06.003

de Cheveigné, A., & Simon, J. Z. (2008). Sensor noise suppression. *Journal of Neuroscience Methods*, *168*(1), 195–202. https://doi.org/10.1016/j.jneumeth.2007.09.012

Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics*, *16*(1), 143–149. https://doi.org/10.3758/BF03203267

Fiebelkorn, I. C., & Kastner, S. (2019). A Rhythmic Theory of Attention. *Trends in Cognitive Sciences*, *23*(2), 87–101. https://doi.org/10.1016/j.tics.2018.11.009

Fries, P. (2015). Rhythms For Cognition: Communication Through Coherence. *Neuron*, *88*(1), 220–235. https://doi.org/10.1016/j.neuron.2015.09.034

Grahek, I., Schettino, A., Koster, E. H. W., & Andersen, S. K. (2021). Dynamic Interplay between Reward and Voluntary Attention Determines Stimulus Processing in Visual Cortex. *Journal of Cognitive Neuroscience*, *33*(11), 2357–2371. https://doi.org/10.1162/jocn_a_01762

Hart, A. S., Clark, J. J., & Phillips, P. E. M. (2015). Dynamic shaping of dopamine signals during probabilistic Pavlovian conditioning. *Neurobiology of Learning and Memory*, *117*, 84–92. https://doi.org/10.1016/j.nlm.2014.07.010

Hickey, C., & Peelen, M. V. (2015). Neural mechanisms of incentive salience in naturalistic human vision. *Neuron*, *85*(3), 512–518. https://doi.org/10.1016/j.neuron.2014.12.049

Hyvärinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis. *IEEE Transactions on Neural Networks*, *10*(3), 626–634. https://doi.org/10.1109/72.761722

Itthipuripat, S., Vo, V. A., Sprague, T. C., & Serences, J. T. (2019). Value-driven attentional capture enhances distractor representations in early visual cortex. *PLOS Biology*, *17*(8), e3000186. https://doi.org/10.1371/journal.pbio.3000186

Jenkins, J., Rhone, A. E., Idsardi, W. J., Simon, J. Z., & Poeppel, D. (2011). The Elicitation of Audiovisual Steady-State Responses: Multi-Sensory Signal Congruity and Phase Effects. *Brain Topography*, *24*(2), 134–148. https://doi.org/10.1007/s10548-011-0174-1

Jensen, O., Bonnefond, M., & VanRullen, R. (2012). An oscillatory mechanism for prioritizing salient unattended stimuli. *Trends in Cognitive Sciences*, *16*(4), 200–206. https://doi.org/10.1016/j.tics.2012.03.002

Jovanovic, L, & Mamassian, P. (2020). Events are perceived earlier in peripheral vision. *Current Biology*, *30*(21), R1299–R1300. https://doi.org/10.1016/j.cub.2020.08.096

Junghöfer, M., Elbert, T., Tucker, D. M., & Rockstroh, B. (2000). Statistical control of artifacts in dense array EEG/MEG studies. *Psychophysiology*, *37*(04), 523–532. https://doi.org/10.1111/1469-8986.3740523

Kayser, C. (2009). Phase Resetting as a Mechanism for Supramodal Attentional Control. *Neuron*, *64*(3), 300–302. https://doi.org/10.1016/j.neuron.2009.10.022

Keil, J., & Senkowski, D. (2018). Neural Oscillations Orchestrate Multisensory Processing. *The Neuroscientist*, *24*(6), 609–626. https://doi.org/10.1177/1073858418755352

Keitel, C., Keitel, A., Benwell, C. S. Y., Daube, C., Thut, G., & Gross, J. (2019). Stimulus-Driven Brain Rhythms within the Alpha Band: The Attentional-Modulation Conundrum. *The Journal of Neuroscience*, *39*(16), 3119–3129. https://doi.org/10.1523/JNEUROSCI.1633-18.2019

Keitel, C., & Müller, M. M. (2016). Audio-visual synchrony and feature-selective attention co-amplify early visual processing. *Experimental Brain Research*, *234*(5), 1221–1231. https://doi.org/10.1007/s00221-015-4392-8

Kim, A. J., Grégoire, L., & Anderson, B. A. (2021). Value-Biased Competition in the Auditory System of the Brain. *Journal of Cognitive Neuroscience*, *34*(1), 180–191. https://doi.org/10.1162/jocn_a_01785

Kim, S., & Beck, M. R. (2020). Impact of relative and absolute values on selective attention. *Psychonomic Bulletin & Review*, *27*(4), 735–741. https://doi.org/10.3758/s13423-020-01729-4

Kim, Y. J., Grabowecky, M., Paller, K. A., Muthu, K., & Suzuki, S. (2007). Attention induces synchronization-based response gain in steady-state visual evoked potentials. *Nature Neuroscience*, *10*(1), 117–125. https://doi.org/10.1038/nn1821

Klink, P. C., Teeuwen, R. R. M., Lorteije, J. A. M., & Roelfsema, P. R. (2023). Inversion of pop-out for a distracting feature dimension in monkey visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *120*(9), e2210839120. https://doi.org/10.1073/pnas.2210839120

Koenig, S., Kadel, H., Uengoer, M., Schubö, A., & Lachnit, H. (2017). Reward Draws the Eye, Uncertainty Holds the Eye: Associative Learning Modulates Distractor Interference in Visual Search. *Frontiers in Behavioral Neuroscience*, *11*. https://www.frontiersin.org/articles/10.3389/fnbeh.2017.00128

Kösem, A., Gramfort, A., & van Wassenhove, V. (2014). Encoding of event timing in the phase of neural oscillations. *NeuroImage*, *92*, 274–284. https://doi.org/10.1016/j.neuroimage.2014.02.010

Lakatos, P., Barczak, A., Neymotin, S. A., McGinnis, T., Ross, D., Javitt, D. C., & O'Connell, M. N. (2016). Global dynamics of selective attention and its lapses in primary auditory cortex. *Nature Neuroscience*, *19*(12), 1707–1717. https://doi.org/10.1038/nn.4386

Lakatos, P., Musacchia, G., O'Connell, M. N., Falchier, A. Y., Javitt, D. C., & Schroeder, C. E. (2013). The spectrotemporal filter mechanism of auditory selective attention. *Neuron*, *77*(4), 750–761. https://doi.org/10.1016/j.neuron.2012.11.034

Lakatos, P., O'Connell, M. N., Barczak, A., Mills, A., Javitt, D. C., & Schroeder, C. E. (2009). The leading sense: Supramodal control of neurophysiological context by attention. *Neuron*, *64*(3), 419–430. https://doi.org/10.1016/j.neuron.2009.10.014

Landau, A. N., & Fries, P. (2012). Attention samples stimuli rhythmically. *Current Biology: CB*, *22*(11), 1000–1004. https://doi.org/10.1016/j.cub.2012.03.054

Lewis, R., & Noppeney, U. (2010). Audiovisual synchrony improves motion discrimination via enhanced connectivity between early visual and auditory areas. Journal of Neuroscience, 30(37),12329–12339. https://doi.org/10.1523/JNEUROSCI.5745-09.2010

Liao, M.-R., & Anderson, B. A. (2020). Inertia in value-driven attention. *Learning & Memory*, *27*(12), 488–492. https://doi.org/10.1101/lm.052027.120

MacLean, M. H., & Giesbrecht, B. (2015). Neural evidence reveals the rapid effects of reward history on selective attention. *Brain Research*, *1606*, 86–94. https://doi.org/10.1016/j.brainres.2015.02.016

Maddox, R. K., Atilgan, H., Bizley, J. K., & Lee, A. K. C. (2015). Auditory selective attention is enhanced by a task-irrelevant temporally coherent visual stimulus in human listeners. *eLife*, *4*. https://doi.org/10.7554/eLife.04995

Markovic, J., Anderson, A. K., & Todd, R. M. (2014). Tuning to the significant: Neural and genetic processes underlying affective enhancement of visual perception and memory. *Behavioural Brain Research*, *259*, 229–241. https://doi.org/10.1016/j.bbr.2013.11.018

Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and memory. *Behavioral and Brain Sciences*, *39*, e200. https://doi.org/10.1017/S0140525X15000667

Miller, L. M., & D'Esposito, M. (2005). Perceptual fusion and stimulus coincidence in the cross-modal integration of speech. *Journal of Neuroscience*, *25*(25), 5884–5893. https://doi.org/10.1523/JNEUROSCI.0896-05.2005

Mishra, J., Martinez, A., Sejnowski, T. J., & Hillyard, S. A. (2007). Early cross-modal interactions in auditory and visual cortex underlie a sound-induced visual illusion. *Journal of Neuroscience*, *27*(15), 4120–4131. https://doi.org/10.1523/JNEUROSCI.4912-06.2007

Monosov, I. E. (2020). How Outcome Uncertainty Mediates Attention, Learning, and Decision-Making. *Trends in Neurosciences*, *43*(10), 795–809. https://doi.org/10.1016/j.tins.2020.06.009

Nakanishi, M., Wang, Y., Wang, Y.-T., Mitsukura, Y., & Jung, T.-P. (2014). Generating Visual Flickers for Eliciting Robust Steady-State Visual Evoked Potentials at Flexible Frequencies Using Monitor Refresh Rate. *PLOS ONE*, *9*(6), e99235. https://doi.org/10.1371/journal.pone.0099235

Noesselt, T., Rieger, J. W., Schoenfeld, M. A., Kanowski, M., Hinrichs, H., Heinze, H. J., & Driver, J. (2007). Audiovisual temporal correspondence modulates human multisensory superior temporal sulcus plus primary sensory cortices. *The Journal of Neuroscience*, *27*(42), 11431–11441. https://doi.org/10.1523/JNEUROSCI.2252-07.2007

Noppeney, U. (2021). Perceptual Inference, Learning, and Attention in a Multisensory World. *Annual Review of Neuroscience*, *44*, 449–473. https://doi.org/10.1146/annurev-neuro-100120-085519

Nozaradan, S., Peretz, I., & Mouraux, A. (2012). Steady-state evoked potentials as an index of multisensory temporal binding. *NeuroImage*, *60*(1), 21–28. https://doi.org/10.1016/j.neuroimage.2011.11.065

Pearson, D., Watson, P., Albertella, L., & Le Pelley, M. E. (2022). Attentional economics links value-modulated attentional capture and decision-making. *Nature Reviews Psychology*, *1*(6), Article 6. https://doi.org/10.1038/s44159-022-00053-z

Peck, C. J., Jangraw, D. C., Suzuki, M., Efem, R., & Gottlieb, J. (2009). Reward modulates attention independently of action value in posterior parietal cortex. *The Journal of Neuroscience*, *29*(36), 11182–11191. https://doi.org/10.1523/JNEUROSCI.1929-09.2009

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience Methods*, *162*(1), 8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Pernet, C., Wilcox, R., & Rousselet, G. (2013). Robust Correlation Analyses: False Positive and Power Validation Using a New Open Source Matlab Toolbox. *Frontiers in Psychology*, *3*. https://www.frontiersin.org/articles/10.3389/fpsyg.2012.00606

Pooresmaeili, A., FitzGerald, T. H. B., Bach, D. R., Toelch, U., Ostendorf, F., & Dolan, R. J. (2014). Cross-modal effects of value on perceptual acuity and stimulus encoding. *Proceedings of the National Academy of Sciences*, *111*(42), 15244–15249. https://doi.org/10.1073/pnas.1408873111

Qin, N., Gu, R., Xue, J., Chen, C., & Zhang, M. (2021). Reward-driven attention alters perceived salience. *Journal of Vision*, *21*(1):7, 1–13. https://doi.org/10.1167/jov.21.1.7

Rolls, E. T., Deco, G., Huang, C.-C., & Feng, J. (2022). The human language effective connectome. *NeuroImage*, *258*, 119352. https://doi.org/10.1016/j.neuroimage.2022.119352

Rohe, T., & Noppeney, U. (2015). Cortical hierarchies perform Bayesian causal inference in multisensory perception. *PLoS Biology, 13*(2), e1002073. https://doi.org/10.1371/journal.pbio.1002073

Rohe, T., & Noppeney, U. (2016). Distinct Computational Principles Govern Multisensory Integration in Primary Sensory and Association Cortices. *Current Biology*, *26*(4), 509–514. https://doi.org/10.1016/j.cub.2015.12.056

Rouhani, N., Norman, K. A., & Niv, Y. (2018). Dissociable effects of surprising rewards on learning and memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *44*(9), 1430–1443. https://doi.org/10.1037/xlm0000518

Rusz, D., Le Pelley, M. E., Kompier, M. A. J., Mait, L., & Bijleveld, E. (2020). Reward-driven distraction: A meta-analysis. *Psychological Bulletin, 146*(10), 872–899. https://doi.org/10.1037/bul0000296

Sanz, L. R. D., Vuilleumier, P., & Bourgeois, A. (2018). Cross-modal integration during value-driven attentional capture. *Neuropsychologia, 120*, 105–112. https://doi.org/10.1016/j.neuropsychologia.2018.10.014

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience*, *10*(3), Article 3. https://doi.org/10.1038/nrn2573

Schwarzkopf, D., de Haas, B., & Rees, G. (2012). Better Ways to Improve Standards in Brain-Behavior Correlation Analysis. *Frontiers in Human Neuroscience*, *6*. https://www.frontiersin.org/articles/10.3389/fnhum.2012.00200

Stanislaw, H., & Todorov, N. (1999). Calculation of signal detection theory measures. *Behavior Research Methods, Instruments, & Computers*, *31*(1), 137–149. https://doi.org/10.3758/BF03207704

Stănișor, L., van der Togt, C., Pennartz, C. M. A., & Roelfsema, P. R. (2013). A unified selection signal for attention and reward in primary visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(22), 9136–9141. https://doi.org/10.1073/pnas.1300117110

Tankelevitch, L., Spaak, E., Rushworth, M. F. S., & Stokes, M. G. (2020). Previously Reward-Associated Stimuli Capture Spatial Attention in the Absence of Changes in the Corresponding Sensory Representations as Measured with MEG. *Journal of Neuroscience*, *40*(26), 5033–5050. https://doi.org/10.1523/JNEUROSCI.1172-19.2020

Todd, R. M., & Manaligod, M. G. M. (2018). Implicit guidance of attention: The priority state space framework. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *102*, 121–138. https://doi.org/10.1016/j.cortex.2017.08.001

Vakhrushev, R., Cheng, F. P.-H., Schacht, A., & Pooresmaeili, A. (2023). Differential effects of intra-modal and cross-modal reward value on perception: ERP evidence. *PLOS ONE*, *18*(6), e0287900. https://doi.org/10.1371/journal.pone.0287900

van Diepen, R. M., & Mazaheri, A. (2018). The Caveats of observing Inter-Trial Phase-Coherence in Cognitive Neuroscience. *Scientific Reports*, *8*(1), 2990. https://doi.org/10.1038/s41598-018-20423-z

Wang, L., Yu, H., Hu, J., Theeuwes, J., Gong, X., Xiang, Y., Jiang, C., & Zhou, X. (2015). Reward breaks through center-surround inhibition via anterior insula. *Human Brain Mapping*, *36*(12), 5233–5251. https://doi.org/10.1002/hbm.23004

Werner, S., & Noppeney, U. (2010). Distinct functional contributions of primary sensory and association areas to audiovisual integration in object categorization. *Journal of Neuroscience*, *30*(7), 2662–2675. https://doi.org/10.1523/JNEUROSCI.5091-09.2010