Title: **Pannexin 1 and Pannexin 3 differentially regulate the tumorigenic properties of cutaneous squamous cell carcinoma**

Authors: **Brooke L. O'Donnell[1], Danielle Johnston[1], Ayushi Bhatt[2], Zahra Kardan[3], Dan Stefan[1], Andrew Bysice[4], Samar Sayedyahossein[5], Lina Dagnino[5,9], Matthew Cecchini[6], Sampath Kumar Loganathan[3,7], Kathryn Roth[4,8] and Silvia Penuela[1,9]**

Affiliations:

[1] Department of Anatomy and Cell Biology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1.

[2] Faculty of Medicine, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1

[3] Department of Surgical and Interventional Sciences, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec. Canada, H4A 3J1.

[4] Department of Otolaryngology—Head and Neck Surgery, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1.

[5] Department of Physiology and Pharmacology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1.

[6] Department of Pathology and Laboratory Medicine, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1.

[7] Departments of Otolaryngology—Head and Neck Surgery, Biochemistry and Experimental Medicine, Rosalind and Moris Goodman Cancer Research Institute, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec. Canada, H4A 3J1.

[8] London Regional Cancer Program, London Health Sciences Centre, London, Ontario. Canada, N6A 5W9.

[9] Department of Oncology, Division of Experimental Oncology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, N6A 5C1.

*Corresponding author: Silvia Penuela, Associate Professor, Department of Anatomy and Cell Biology, University of Western Ontario, London, Ontario, Canada. Tel. 519-661-2111 ext. 84735; E-mail: spenuela@uwo.ca

Running title: PANX1 and PANX3 regulate cSCC malignancy.

Key points summary

- Pannexin 1 and pannexin 3 are channel-forming proteins which are critical in the normal maintenance and function of keratinocytes in the skin but may become altered in cutaneous squamous cell carcinoma (cSCC) tumours.

- In this study, using a combination of culture models, mouse models and patient-derived tissues, we found pannexin 1 levels are increased in cSCC tumours and present in all tumour cell types, functioning to increase cSCC cell growth and migration.

- On the other hand, pannexin 3 levels are decreased in cSCC tumours and this protein reduces the incidence and growth of pre-cancerous lesions.

- Taken together, our data indicates that in cSCC these pannexin family members seem to have opposite effects, where pannexin 1 is pro-tumorigenic and pannexin 3 is anti-tumorigenic.

- These results help us to better understand the mechanisms of malignant transformation of keratinocytes and offer a new potential therapeutic target for the treatment of advanced cSCC.

Abstract

Pannexin (PANX) channels are present in skin and facilitate the movement of signalling molecules during cellular communication. PANX1 and PANX3 function in skin homeostasis and keratinocyte differentiation but were previously reduced in a small cohort of human cutaneous squamous cell carcinoma (cSCC) tumours compared to normal epidermis. In our study, we used SCC-13 cells, limited publicly available RNA-seq data and a larger cohort of cSCC patient-matched samples to analyze PANX1 and PANX3 expression and determine the association between their dysregulation and the malignant properties of cSCC. In a bioinformatics analysis, *PANX1* transcripts were increased in both cSCC and head and neck SCC tumours compared to normal tissues, but *PANX3* mRNA showed no differences. However, in our own patient cohort, *PANX3* transcripts were decreased in cSCC tissue compared to patient-matched aged skin, whereas PANX1 was upregulated in cSCC. PANX1 localized to all cell types within the cSCC tumour microenvironment and increased levels were associated with larger tumour dimensions. To investigate PANX1 channel function, we treated SCC-13 cells with PANX1 inhibitors which markedly reduced cell growth and migration. To assess PANX3 function in cutaneous

carcinogenesis, we employed the DMBA/TPA carcinoma model using our global *Panx3* knockout (KO) mice, where 60% of wildtype and 100% of KO mice formed pre-cancerous papillomas. Average papilloma volumes at endpoint were significantly increased in KO mice and showed moderate evidence of increases in KO mice over time. Collectively, these findings suggest PANX1 and PANX3 dysregulation may have potential tumour promoting and suppressive effects for keratinocyte transformation, respectively.

Introduction

Cutaneous squamous cell carcinoma (cSCC) is the second-most common form of skin cancer arising from squamous keratinocytes in the upper layers of the epidermis or stem cell populations in cutaneous glands and hair follicles (Firnhaber, 2012; Kallini et al., 2015). Despite only accounting for 20% of non-melanoma skin cancer (NMSC) cases (Madan et al., 2010; Firnhaber, 2012), cSCC is responsible for most NMSC-related deaths often due to tumour metastasis (Nissinen et al., 2016), and the incidence of cSCC is projected to continue increasing worldwide as the population ages (Lomas et al., 2012; Neagu et al., 2019). Generally, surgical resection is a viable treatment option for small and localized tumours, but approximately 5% of cSCC progress to advanced disease, becoming highly invasive (Schmults et al., 2013). This contributes to a very high recurrence rate and requires additional treatment methods such as chemotherapy, radiation, targeted therapies and immunotherapy, which have lower success rates that vary depending on the patient (Fahradyan et al., 2017; Aboul-Fettouh et al., 2021). Despite these considerations, the mechanisms of keratinocyte malignant transformation into cSCC are still poorly understood (Ratushny et al., 2012).

Within the skin, cells express pannexins (PANX1, PANX2, PANX3) which are canonically known for their roles as channel-forming glycoproteins (Panchin et al., 2000) that facilitate cellular communication by metabolite and ion conductance (Bao et al., 2004; Iwamoto et al., 2010; Ishikawa et al., 2011; O'Donnell & Penuela, 2023). PANX1 has been demonstrated to play a role in murine skin development and the establishment of proper skin architecture, with protein levels being high in young skin and decreasing with age in the absence of challenges such as wound repair (Penuela et al., 2014). Conversely, PANX3 is undetectable in young mouse dorsal skin but becomes upregulated with age and is highly abundant in aged mouse skin, functioning to maintain skin homeostasis during the process of aging (O'Donnell et al., 2023).

Both PANX1 and PANX3 have been localized to all living layers of keratinocytes, where they act in keratinocyte differentiation, adhesion and/or inflammation. During calcium-induced keratinocyte differentiation, PANX1 levels decrease (Penuela et al., 2014) whereas PANX3 levels were found to either be induced or remain unchanged (Zhang et al., 2019; Zhang et al., 2021; O'Donnell et al., 2023), indicating distinct expression patterns. Interestingly, PANX1 localization changes from a cell membrane profile in basal, undifferentiated keratinocytes, to a more intracellular profile in supra-basal, differentiated keratinocytes (Penuela et al., 2014), whereas PANX3 exhibits a solely cytoplasmic localization in both undifferentiated and differentiated cells (Cowan et al., 2012). Additionally, overexpression of each protein in keratinocytic cell lines reduced proliferation, but only PANX1 overexpression increased keratin 14 levels (basal keratinocyte marker) and disrupted terminal differentiation which normally facilitates the formation of epidermal architecture (Celetti et al., 2010). However, little is known about pannexin expression and function in keratinocytic cancers where the normal keratinocyte differentiation program is disrupted.

PANX1 has previously been implicated in melanoma tumorigenesis. Our group showed PANX1 levels are increased in melanoma cells compared to normal melanocytes, and *PANX1* mRNA is upregulated in patient melanoma tumour samples compared to control skin, with levels remaining high throughout all stages of melanoma disease progression (Penuela et al., 2012; Freeman et al., 2019). In human A375-P and A375-MA2 melanoma cells, decreased cellular growth rate, migration, invasion and mitochondrial metabolic activity, as well as increased melanin production were observed when PANX1 channel activity was inhibited or PANX1 levels were reduced via shRNA knockdown (Freeman et al., 2019). The malignant melanoma marker β-catenin, which was found to bind directly to the C-terminal tail of PANX1, was also significantly decreased (Freeman et al., 2019; Sayedyahossein et al., 2021). Altogether, this suggests a role for PANX1 throughout melanoma carcinogenesis, but little is known about PANX1 or PANX3 in NMSCs.

In a study using the 7,12-dimethylbenz(a)anthracene/12-Otetradecanoylphorbol-13-acetate (DMBA/TPA) mouse cutaneous carcinoma model, which mimics many aspects of human SCC (Abel et al., 2009), *Panx3* was found genetically linked to body mass index and tumorigenesis by quantitative trait loci analysis in male but not female mice. It was also noted that pre-treatment *Panx3* transcript levels in tail skin were positively associated with tumour

susceptibility, where the higher the expression, the higher the risk of papilloma and carcinoma formation. Despite this, *Panx3* transcript levels were markedly reduced in papillomas and carcinomas compared to untreated tail skin, suggesting that *Panx3* may not influence tumour maintenance or progression (Halliwill et al., 2016). In human cSCC patient-derived tumours and aged epidermis, immunofluorescence microscopy analysis using a small cohort of samples indicated that both PANX1 and PANX3 were visibly reduced in cSCC tumours (Cowan et al., 2012). However, data from the Human Protein Atlas shows conflicting results for PANX1 (PANX3 is not annotated), reporting that PANX1 is present at moderate levels in immunohistochemically stained cSCC tumour cores, where cells showed both membranous and cytoplasmic PANX1 localization. Collectively, these results suggest PANX1 and PANX3 levels are altered in cSCC tumorigenesis, but further investigation is needed to navigate the conflicting results and understand the effects of pannexin dysregulation.

In this study we utilized the SCC-13 cell line, publicly available cSCC and head and neck SCC (HNSCC) patient expression data, patient-derived skin and cSCC tissues, and a mouse model of cutaneous carcinogenesis to evaluate PANX1 and PANX3 expression in cSCC compared to normal skin and their role in cSCC malignancy. In our patient sample cohort, we found PANX1 protein is upregulated whereas *PANX3* mRNA is downregulated in human cSCC tumour fragments compared to patient-matched normal skin. Within SCC-13 cells, PANX1 exhibits both an intracellular and cell surface localization, whereas PANX3 is found predominantly in the cytosol. We also demonstrated that PANX1 promotes cancer cell properties such as growth and migration through its channel activity and is present in all cell types within the cSCC tumour microenvironment and adjacent skin. On the contrary, PANX3 has a tumour suppressive role where it reduces the incidence and growth of pre-cancerous papillomas.

Methods

Ethical approval

Animal experiments were approved by the Animal Care Committee at the University of Western Ontario (London, ON, Canada, Protocol #2022-025). For collection of human samples, the study was conducted in accordance with the Declaration of Helsinki, and the protocol was approved by the Health Science Research Ethics Board (HSREB) of Western University and London Health

Sciences Centre (London, ON, Canada, HSREB#103381). Subjects provided informed consent to participate in this study and tumour identity was determined by a pathologist.

Cell culture

SCC-13 cells (RRID:CVCL_4029) were cultured in DMEM medium (Gibco 12430062, Grand Island, NY) supplemented with 10% fetal bovine serum (WISENT 098-150, Saint-Jean-Baptiste, QC, Canada). U-2 OS (ATCC HTB-96, Manassas, VA; RRID:CVCL_0042), GBM17 (gifted from Dr. Matthew Hebb, University of Western Ontario), Hs578T (ATCC HTB-126, RRID:CVCL_0332), and Hs578T *PANX1* KO (Nouri-Nejad et al., 2021) cells used for positive and negative controls for Western blotting were cultured in DMEM medium supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin (Gibco 15140-122). Cells were maintained at 37°C and 5% CO_2 and dissociated using 0.25% trypsin/1 mM EDTA (WISENT 325-043-EL) for 7 min after one wash with 1x DPBS without Ca^{2+} and Mg^{2+} (Gibco 14190250) for SCC-13 cells or 3 min for all other cells. Cells were transfected using Lipofectamine™ 3000 (Invitrogen L3000015, Carlsbad, CA) in Opti-MEM medium (Life Technologies 51985-034, Carlsbad, CA) according to the manufacturer's protocol with 1 µg of PANX1 (InvivoGen puno1-hpanx1, San Diego, CA) or PANX3 plasmid in a 6 well plate. The PANX3 construct was modified from the original Origene plasmid (PS100001, Rockville, MD) by NorClone Biotech Laboratories (London, ON, Canada) to replace the FLAG tag with an HA tag. Protein extracts were prepared 48 hr after transfection for Western blot positive controls.

Protein isolation and Western blotting

Protein lysates were prepared from cells using 1x RIPA buffer, and from human tissue using 2x RIPA buffer after pulverizing in liquid nitrogen. Both extraction buffers also contained 1 mM sodium fluoride, 1 mM sodium orthovanadate and a half of a Pierce Protease Inhibitor EDTA-free Mini Tablet in 10 mL of buffer (Thermo Fisher Scientific A32955). Western blots were performed as reported (O'Donnell et al., 2023), using 70 µg per sample for SCC-13 cells and 40 µg per sample for human tissue and separated on 10% acrylamide SDS-PAGE gels. Gels were transferred to nitrocellulose membranes using an iBlot Gel Transfer Device (Invitrogen) on P3 and blocked for 1 hr at room temperature in 1x PBS with 3% bovine serum albumin (BioShop ALB001.100, Burlington, Canada). Primary antibody dilutions included: anti-PANX1 CT-412

1:1000, anti-PANX3 EL1-84 1:250, anti-PANX3 CT-379 1:1000 (Penuela et al., 2007; Penuela et al., 2009), anti-PANX3 (Thermo Fisher Scientific 433270) and anti-GAPDH 1:5000 (Millipore Sigma G8795, RRID:AB_1078991, Burlington, MA). Dilutions of 1:10,000 were used for all secondary antibodies, including IRDye-800CW goat anti-rabbit (926-32211, RRID:AB_621843) and IRDye-680RD goat anti-mouse (926-68070, RRID:AB_10956588) IgG secondary antibodies from LI-COR Biosciences (Lincoln, NE). All antibodies were dissolved in 3% bovine serum albumin in 0.05% Tween-20 PBS. PANX1 and PANX3 antibodies were incubated overnight at 4°C, whereas GAPDH and secondary antibodies were incubated for 45 min at room temperature. The LI-COR Odyssey Infrared Imaging System and ImageStudio software (LI-COR Biosciences) were used to image blots and perform protein level and molecular weight quantifications. Protein levels were normalized to GAPDH and presented relative to the control mean value.

Immunocytochemistry

For analysis of PANX1 localization in SCC-13 cells and SCC-13 cells ectopically expressing a PANX1 plasmid, 300,000 cells were plated on glass coverslips (Fisher Scientific 1254580, Pittsburgh, PA) in a 6 well plate. SCC-13 cells were transfected with 1 µg of PANX1 plasmid 24 hr after seeding. For analysis of PANX3-HA localization in SCC-13 cells, 500,000 cells were seeded on glass coverslips in a 6 well plate and transfected with 1 µg of PANX3-HA plasmid 24 hr after seeding. Forty-eight hours after plating SCC-13 cells (endogenous PANX1) or 48 hr post-transfection for PANX1 or PANX3-HA expressing SCC-13 cells, cells were washed twice with 1x DPBS without Ca^{2+} and Mg^{2+} and fixed with ice-cold methanol:acetone (5:1, v/v) for 15 min at 4°C. For 1 hr at room temperature, coverslips were blocked with 10% goat serum (Life Technologies 50062Z), and then incubated overnight at 4°C with PANX1 CT-412 (Penuela et al., 2007; Penuela et al., 2009) or HA (Cell Signaling 3724, RRID:AB_1549585, Danvers, MA) primary antibody in a humidity chamber. After three 5 min washes with 1x DPBS without Ca^{2+} and Mg^{2+}, coverslips were incubated with 1:700 Alexa Fluor™ 488 goat anti-rabbit (Invitrogen #A11008, RRID:AB_143165) or 1:700 Alexa Fluor™ 488 goat anti-mouse secondary antibody (Invitrogen A11029, RRID:AB_2534088) for 1 hr at room temperature. All antibodies were diluted in 1x DPBS without Ca^{2+} and Mg^{2+} with 1% goat serum. After one 1x DPBS without Ca^{2+} and Mg^{2+} and one double-distilled water wash for 5 min each, coverslips were incubated for

5 min at room temperature with Hoechst 33342 (Life Technologies H3570) diluted 1:1000 in double-distilled water to stain cell nuclei. ProLong™ Gold Antifade Mountant (Thermo Fisher Scientific P36934) was used to mount coverslips. A ZEISS LSM 800 AiryScan Confocal Microscope from the Schulich Imaging Core Facility (University of Western Ontario) was used to image cells at 63x oil magnification with 405 nm (Hoechst 33342) and 488 nm (Alexa Fluor™ 488) laser lines.

PANX1 inhibition

Probenecid (PBN, Invitrogen P36400) and spironolactone (SPL, Selleckchem S4054, Houston, TX) dissolved in water and 100% ethanol (EtOH), respectively, were used to block PANX1 channels. For all cell experiments a final concentration of 1 mM PBN or 20 µM SPL was used and treatments or corresponding vehicle controls were added to culture medium at day 0 and replaced every 2 days. A 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay (Abcam ab211091, Cambridge, UK) was used to confirm treatments did not affect cell viability. For the MTT assay, 50,000 SCC-13 cells were seeded in 6 well plates and treated with vehicle, PANX1 inhibitor or 1 µM staurosporine (Sigma-Aldrich S6942, Burlington, MA) dissolved in culture media 6 days later. Staurosporine was used as a control for reduced cell viability. Twenty-four hours after treatment, MTT was diluted 1:40 in culture medium and incubated with the cells for 1 hr at 37°C and 5% CO_2. After aspirating growth media, 500 µL of DMSO was added to each well and shaken for 5 min at room temperature to dissolve the formazan products. In a 96 well plate, triplicates of each condition were diluted 1:2 in DMSO and the OD570 was measured using the VICTOR^{3}™ Multilabel Counter plate reader (PerkinElmer, Inc., Shelton, CT). Triplicate values were averaged for each experiment and calculated as a fold change relative to vehicle averages.

Growth and migration assays

For growth assays, 50,000 SCC-13 cells were seeded in 12 well plates and treated with either vehicle or inhibitor treatment 48 hr after seeding. Phase contrast images were taken at 10x magnification every 6 hr for 5 days using the Incucyte® S3 Live-Cell Analysis Instrument (Sartorius, Goettingen, Germany), and cell counts per image were analyzed using the Incucyte® Cell-By-Cell Analysis Software Module (Sartorius BA-04871). For migration assays, 40,000

SCC-13 cells were seeded on Incucyte® Imagelock 96 well plates (Sartorius BA-04856) and grown to confluency. Scratch wounds were created in each well using the Incucyte® Woundmaker Tool (Sartorius BA-04858), and after one wash with DPBS without Ca^{2+} and Mg^{2+} to remove cell debris, cells were treated with serum free 1x DMEM medium supplemented with either vehicle or inhibitor treatment. Cells were imaged using the Incucyte® S3 Live-Cell Analysis Instrument and Incucyte® Scratch Wound Analysis Software Module (Sartorius 9600-0012), with phase contrast images taken every 2 hr for 46 hr at 10x magnification. Analysis of % wound confluence (% wound closure) was performed using the same software.

DMBA/TPA mouse model

The DMBA/TPA mouse carcinoma protocol (Sundberg et al., 1997; Filler et al., 2007; Halliwill et al., 2016) was employed to investigate the effects of PANX3 on papilloma and carcinoma formation and progression. Eight-week-old male C57BL/6N wildtype (WT) and congenic global *Panx3* knockout (KO) mice (Moon et al., 2015) (*N*=5) were treated with one dose of 25 µg DMBA (dissolved in acetone), followed by twice weekly treatments of 4 µg TPA (dissolved in acetone) until endpoint at 28 weeks. Treatments were administered topically to the first centimetre of the tail at the tail base. Throughout treatment, mice were monitored for their body condition, weight, and papilloma formation (volume/growth over time). Papillomas were imaged using an iPhone SE (Apple, Inc., Cupertino, CA) and measured once weekly with a digital caliper according to the University of Western Ontario's CLN-100 Monitoring and Humane Endpoints for Rodent Cancer Models SOP until endpoint. Papilloma volumes were calculated using the following equation: $V = 4/3\pi \times (Length/2) \times (Width/2)^2$ where V is tumour volume, length is the dimension along the longest axis, width is the dimension along the shortest axis. At endpoint, mice were sacrificed using CO_2 inhalation. Mice were genotyped as described previously (O'Donnell et al., 2023), fed Teklad 2018 chow (Envigo, Indianapolis, IN) *ad libitum* and maintained in a 12-hr-light to 12-hr-dark cycle.

Bioinformatics analysis of PANX1 and PANX3 expression in cSCC and HNSCC

We utilized the Gene Expression Omnibus (GEO) database to download the RNA-seq FASTQ files of two separate cSCC studies for further validation (Edgar et al., 2002). The GSE191334 database contained 8 bulk RNA-Seq samples of cSCC with paired normal controls of human skin

organoids, and GSE139505 consisted of the whole transcriptome profile of 9 cSCC and 7 unmatched healthy skin as controls. The TCGA biolinks package (Colaprico et al., 2016) was employed to retrieve mRNA expression generated from The Cancer Genome Atlas (TCGA) HNSCC RNA-sequencing data which contains 502 HNSCC tissues and 44 adjacent normal tissues, including clinical information and gene expression from TCGA. Incomplete data was deleted before analysis. Background correction, normalization, and expression value calculation were performed on the original datasets using the package Limma of R software (version x64 4.2.1). Fold change (FC) and adjusted p-values (p adj.) were used to screen differentially expressed genes (DEGs) using DESeq2. $|\log(FC)| \geq 1$ and p adj.<0.05 were defined as the screening criteria for DEGs. Scatter plots were produced using the Ggplot2 package (https://ggplot2.tidyverse.org/authors.html).

Patient samples

Patient-matched cSCC tumour fragments and normal aged skin (containing only the epidermis and dermis) samples from 20 patients as well as four additional full thickness normal aged skin samples were analyzed in this study. Patient demographics are outlined in Table 2 and 3 in the results section. Images of patient tumours were taken with consent using a password protected iPhone X (Apple. Inc.).

RNA isolation and RT-qPCR

RNA isolation and RT-qPCR for human tissue were performed as described previously (O'Donnell et al., 2023) using a hybrid TRIzol (Life Technologies 15596018) and RNeasy Plus Mini kit protocol after liquid nitrogen pulverization. The High-Capacity cDNA Reverse Transcription Kit with RNase Inhibitor (Thermo Fisher Scientific 4374966, Waltham, MA) was used to reverse transcribe 50 ng of RNA as outlined in the manufacturer's protocol in a T100TM Thermal Cycler (Bio-Rad Laboratories, Hercules, CA). SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Laboratories 1725274) was used to perform RT-qPCR with samples run in duplicate. Transcript levels were normalized to β-2 microglobulin (*β2M)*, presented as relative to the aged skin mean value and analyzed with the ΔΔCT method in Excel (Microsoft 365, Redmond, WA). See Table 1 for list of primers used. Logarithmic means were used to perform statistics, but geometric means were used to create figure graphs.

Immunohistochemistry

cSCC tumour and adjacent skin tissues were fixed in 10% formalin, with routine processing in surgical pathology. Parallel sections (4 µm thickness) were used for hematoxylin and eosin (H&E) and 3,3'-diaminobenzidine (DAB) staining. For DAB staining, tissue slides were baked at 40°C overnight before xylene deparaffinization (two 5 min and one 3 min incubations) and descending EtOH series rehydration (100% EtOH for 2 min, 100% EtOH for 1 min, 95% EtOH for 2 min, 95% EtOH for 1 min, 70% EtOH for 1 min, distilled water for 2 min). Endogenous peroxidases and alkaline phosphatases were quenched using BLOXALL® Endogenous Blocking Solution (Vector Laboratories SP-6000, Newark, CA) for 10 min. Sections were washed with 1x PBS for 5 min and antigen retrieval was performed with Antigen Unmasking Solution Citrate-based pH 6.0 (Vector Laboratories H-3300) in a Decloaking Chamber™ (Biocare Medical, LLC, Pacheco, CA) at 112.5°C for 1.5 min, followed by 90°C for 10 s. Slides were cooled with a tap water rinse and washed with 1x PBS for 5 min. Samples were blocked with 2.5% normal horse serum (Vector Laboratories S-2012-50) for 30 min at room temperature and incubated with 1:250 PANX1 CT-412 diluted in 2.5% normal horse serum at 4°C overnight in a humidity chamber. Secondary antibody control only samples were left in the blocking solution. The next day, slides were washed twice with 1x PBS and incubated with ImmPRESS® HRP Horse Anti-Rabbit IgG Polymer Reagent (Vector Laboratories MP-7401) for 30 min at room temperature. After two 5 min 1x PBS washes, slides were stained with DAB substrate (Vector Laboratories SK-4100, using reagents 1-3) for 3 min at room temperature. After rinsing in tap water, slides were counterstained with Harris hematoxylin (Leica 3801561, Wetzlar, Germany) for 1 min and rinsed again in tap water. Two to three quick dips in 0.6% acid alcohol and 2% ammonium alcohol with water rinses in between were used to change the Harris hematoxylin to a blue colour. Lastly, slides were cleared with an ascending ethanol series (70% EtOH 1 min, 95% EtOH 1 min, 95% EtOH 2 min, 100% EtOH 1 min, 100% EtOH 1 min, 100% EtOH 2 min) and xylene (two 5 min incubations) and mounted using Fisher Chemical Permount Mounting Medium (Fisher Scientific SP15) and microscope cover glass (Fisher Scientific 12545F). Slides were scanned using a Grundium Ocus®40 (Grundium Ltd., Tampere, Finland) and images were processed using QuPath version 0.5.0 (Bankhead et al., 2017). Objective magnifications of images are outlined in the figure legends. Adobe Photoshop CS6 (Adobe, San Jose, CA) was

used to white balance, sharpen and enhance contrast of images which were annotated by a pathologist.

Statistics

Statistical analyses were performed using GraphPad Prism 10 (GraphPad Software, San Diego, CA). Outliers were removed from datasets using the outliers test from the same software. Biological and technical replicates and statistical tests used are reported in each figure legend. Based on recommendations from the editorial by Wasserstein et al. to move beyond *p* values as the sole indicator of biological validity to observed differences (Wasserstein et al., 2019), data was referred to in terms of the strength of the evidence when *p* values do not reach significance. The following *p* value ranges were used in this study: strong evidence when $p<0.05$, moderate evidence when $0.05< p <0.15$ and weak evidence when $0.15< p <0.25$. Any *p* values > 0.25 were considered to show no evidence for differences.

Results

PANX1 and PANX3 are present in SCC-13 cells and exhibit distinct localization patterns
To assess PANX1 and PANX3 levels in cSCC, we first analyzed PANX1 and PANX3 protein levels in SCC-13 cells. Western blot analysis revealed both PANX1 and PANX3 were detectable in SCC-13 cell lysates (Fig. 1*A*,*B*). Immunocytochemistry using untransfected and PANX1-expressing SCC-13 cells (Fig. 1*C*) revealed that endogenous PANX1 showed a variety of presentations, where some cell clusters had a predominantly diffuse intracellular localization, whereas other clusters exhibited both cytosolic and prominent plasma membrane localization. In SCC-13 cells exogenously expressing PANX1, the localization pattern was consistent with endogenous PANX1 found in the cytosol and plasma membrane, but overexpression seemed to increase the prevalence of intracellular species. Conversely, SCC-13 cells exogenously expressing PANX3-HA showed a predominantly punctate intracellular localization although some PANX3-HA could be seen at the plasma membrane (Fig. 1*D*). We were unable to evaluate endogenous PANX3 localization in SCC-13 cells due to a lack of PANX3 antibodies which were suitable for immunocytochemistry.

Blocking PANX1 channel function decreases cancer cell properties of SCC-13 cells in vitro
We previously demonstrated that inhibiting PANX1 channels in another skin cancer reduced the malignant properties of melanoma cells such as growth and migration (Freeman et al., 2019). Since PANX1 was found to be abundant in SCC-13 cells, we evaluated the effects of previously established PANX1 channel blockers PBN and SPL on the corresponding cell properties in this cell line (Fig. 2). With both PBN (p=0.0045) and SPL (p=0.0066) treatment of SCC-13 cells, PANX1 inhibition significantly reduced cell growth over time compared to water and EtOH vehicle controls, where significant cell growth perturbations were more severe with SPL and evident after only 48 hr (p=0.0307) of incubation (Fig. 2*A,B*). Doses of 1 mM PBN and 20 µM SPL were selected based on doses previously reported (Good et al., 2018; Freeman et al., 2019) and confirmed to have no detectable effects on cell viability in comparison to vehicle controls (p=0.605 and p=0.578, respectively) via MTT assay (Supplementary Fig. 1). Next, using scratch assays we investigated SCC-13 migration, determining PANX1 channel inhibition also decreased SCC-13 motility (Fig. 2*C-H*). With PBN, although there was no significant treatment effect due to variability (p=0.0505), there was moderate evidence to suggest reduced percent wound closure over time compared to water treatment (Fig. 2*C*). This was also evident based on representative images of scratch areas (Fig. 2*E*) and quantifications (Fig. 2*G*) at 24 hr (p=0.0757) and 46 hr (p=0.0066), indicating blocking PANX1 channels with PBN reduces SCC-13 migration. The same trend with a more pronounced effect was observed with SPL treatment. There was a significant treatment effect with SPL treatment over time (p=0.0157) compared to vehicle control (Fig. 2*D*), where significant reductions in percent wound closure were evident as early as 16 hr (p=0.0192) after initial treatment. Similar to PBN, representative images (Fig. 2*F*) and quantifications (Fig. 2*H*) at 24 hr (p=0.0233) and 46 hr (p=0.0025) also indicated reduced SCC-13 migration with SPL-induced PANX1 channel inhibition. Thus, PANX1 promotes the cancerous properties of SCC-13 cells such as growth and migration.

Panx3 deletion increases papilloma formation and growth in mouse cutaneous carcinoma model
Due to a lack of PANX3-specific channel blockers, we utilized the DMBA/TPA mouse cutaneous carcinoma protocol (Sundberg et al., 1997; Filler et al., 2007; Halliwill et al., 2016) on a small subset of male WT and congenic global *Panx3* KO mice (*N*=5) to investigate the effects of PANX3 on papilloma and carcinoma formation and progression *in vivo*. Eight-week-old mice

were treated topically with one dose of 25 µg DMBA, followed by 28 weeks of twice weekly 4 µg topical TPA treatments at the tail base until endpoint at 28 weeks (Fig. 3*A*). Unfortunately, one WT mouse (which developed a papilloma and was included in any analyses of papilloma incidence) died during the treatment period before reaching endpoint of causes independent of the study and was therefore removed from all papilloma volume analyses (WT *N*=4, *Panx3* KO *N*=5). Most papillomas developed at the site of chemical administration on top of the tail base (Fig. 3*B*, left panel) and were well-demarcated, symmetrical, and dome-shaped without any ulceration, but few mice also developed papillomas on the dorsal skin immediately above the tail base, with some papillomas becoming keratinized (Fig. 3*B*, right panel). Of the mice in the pilot study, 3/5 WT and 5/5 *Panx3* KO mice developed papillomas (Fig. 3*C*), but none developed carcinomas. Generally, *Panx3* KO mice exhibited a slightly higher papilloma incidence, with KO mice developing an average of 2.2 papillomas per mouse compared to the average of 1 papilloma per WT control. Survival curves for time to papilloma onset showed weak evidence for differences between genotypes (Fig. 3*D*, p=0.176); however, the average response latency to first papilloma formation (for mice that developed papillomas) was similar, with WT mice developing papillomas after 25.7 weeks and *Panx3* KO mice developing papillomas after 25.2 weeks. Observation of growth curves for average papilloma volumes in individual mice showed some overlap between genotypes, but most KO mice tended to have increased papilloma growth over time (Fig. 3*E*). Additionally, although no significant genotype effect was observed due to high variability between mice (p=0.0974), there was moderate evidence to suggest the combined genotype averages for average papilloma volume over time and the corresponding area under the curve measurements (p=0.1106) were increased in KO mice compared to controls (Fig. 3*F,G*). When analyzed at endpoint (week 28), the average papilloma volume was found to be significantly increased in *Panx3* KO mice compared to WT mice (Fig. 3*H*, p=0.0491). A power analysis using the above results with an α value of 0.05 and power of 0.8 indicated a sample size of 5 animals per genotype, indicating our findings were within power. Taken together, this illustrates *Panx3* KO mice tend to have increased papilloma incidence, volumes and growth over time when subjected to a cutaneous carcinogenesis protocol.

Bioinformatic analysis of PANX1 and PANX3 expression in SCC tumours

To explore *PANX1* and *PANX3* transcript expression in patient-derived cSCC tumours compared to normal skin controls, we first performed a bioinformatics search by investigating two cSCC RNA-seq studies available on the GEO database (Fig. 4*A*). In the GSE191334 study, no significant differences were observed in *PANX1* (*p*=0.219) or *PANX3* (*p*=0.719) mRNA expression between tissue types, whereas *PANX1* transcript levels were significantly upregulated (*p*=2.35x10^{-10}) in cSCC tumours with a fold change of +2.65 compared to controls in the GSE139505 study. In the latter study, *PANX3* mRNA levels were either not detectable or not reported. Notably, both studies contained small N values and control tissues consisted of either paired human skin organoids or unpaired healthy skin, making it difficult to draw substantial conclusions about *PANX1* and *PANX3* expression in this cohort of patient-derived cSCC tumours. To address the issue of small N values, we next evaluated *PANX1* and *PANX3* transcript levels in HNSCC RNA-seq data obtained from TCGA (Fig. 4*B,C*). Although the tissue type of origin differs, with HNSCC arising from squamous mucosal surfaces, both cSCC and HNSCC tumours result from the malignant transformation of epithelial cells and share some key molecular markers (Yan et al., 2011). In this study, which contained a much larger N value of 502 HNSCC tumours and 44 adjacent normal tissues, *PANX1* mRNA was found to be significantly increased in HNSCC (*p*=6.54x10^{-14}), with a fold change of +2.20 compared to controls, similar to what was seen in cSCC. *PANX3* transcript levels were not found to be significantly different (*p*=0.655) between HNSCC and adjacent tissues but tended to be reduced in HNSCC tumours.

Patient demographics

Due to the limited cSCC samples present in online databases, we generated our own cohort of cSCC samples (patient demographics outlined in Table 2 and 3). In collaboration with local surgical oncologists, we accrued 20 different patients for our study cohort treated by the London Regional Cancer Program and collected patient-matched cSCC tumour fragments and normal aged skin. The normal skin was either excess tissue from the skin graft used in surgery or obtained via punch biopsy. Within our cohort, the average age was 75.1 years but ranged from adult (51 years) to geriatric (93 years) patients, with most samples coming from males. Seventy percent of patients had a history of previous skin cancer, with 30% being immunosuppressed and 35% exhibiting local recurrence. The cSCC tumours were localized to a variety of anatomic

locations with the highest percentage presenting on the scalp (30%), and a wide range of tumour sizes were observed. Most tumours were American Joint Committee on Cancer stage T3N0M0 (65%), with a moderate degree of differentiation (75%) and deep invasion (60%). Furthermore, the majority of patients included in this study were classified as having advanced cSCC, due to tumour size and invasion criteria (Migden et al., 2018). Lastly, the majority of the normal skin samples were obtained from the clavicle (25%), neck (25%) or leg (25%).

The patients included in this study had a variety of co-morbidities, with cardiovascular and diabetes mellitus being the most common at 80% and 45%, respectively. Consequently, common medications prescribed included those used to treat metabolic syndrome such as anti-hypertensives, statins, anti-platelets, anti-coagulants and anti-hyperglycemics. The PANX1 channel inhibitor SPL which was originally developed to treat hypertension (Koval et al., 2023) was not prescribed to any patient in the study. Other less common co-morbidities affected the gastrointestinal, genitourinary, musculoskeletal, nervous, hematologic and respiratory systems. In addition to cSCC tumour excision, some patients required adjuvant radiation (25%) and immunotherapy (15%). Patient outcomes included local or regional recurrence (25% for both) and metastasis (10%). Taken together, we believe that our cohort, although small, is representative of the patient population.

In human cSCC tumours, PANX3 mRNA is reduced compared to matched skin controls
We wanted to investigate whether the PANX1 and PANX3 expression patterns of our patient-derived cSCC and normal aged skin cohort were consistent with those seen previously (Cowan et al., 2012; Halliwill et al., 2016). Beginning with PANX3, we determined that in aged human full thickness skin samples, PANX3 was highly abundant (Fig. 5*A*), and *PANX3* transcripts were significantly reduced in cSCC tumour fragments compared to patient-matched aged skin samples containing the epidermal and dermal layers (Fig. 5*B*). When analyzing patient skin and cSCC for PANX3 protein levels, the 43 kDa PANX3 species present in the positive control was undetectable in both tissue types, but we identified shorter immunoreactive species ranging from ~32-38 kDa (Fig. 5*C*). We report the presence of PANX3-38K (37.9 kDa), PANX3-36K (35.7 kDa) and PANX3-32K (32.4 kDa) which showed variability in abundance between patients. Comparisons between tissue types showed that total PANX3 (p=0.0175) and PANX3-36K (p=0.0103) were significantly increased in cSCC tumour fragments compared to skin, but no

differences were seen in PANX3-38K (*p*=0.561) or PANX3-32K (*p*=0.971) levels (Fig. 5*D-G*). Further investigation into these immunoreactive species is required. Overall, this indicates that the canonical *PANX3* transcript is decreased in human cSCC tumours compared to patient-matched unaffected epidermis and dermis.

In human cSCC tumours, PANX1 is upregulated compared to matched skin controls and present within all the cell types of the tumour microenvironment

cSCC tumour fragments are three-dimensional structures that contain other cell types in addition to cancer cells such as cancer-associated fibroblasts, immune infiltrate and blood vessels. Thus, to localize PANX1 expression within the various cells of the cSCC tumour microenvironment we performed immunohistochemistry of tumours and adjacent skin tissue from the most common anatomical subtypes of the scalp, ear and cheek (Fig. 6, Supplementary Fig. 2). According to pathologist observations, PANX1 is present throughout all cell types in cSCC tumours and adjacent skin (Fig. 6*B*), with relative staining intensity slightly increased in the tumour. Within the tumour, PANX1 staining was present in both tumour nests and stromal regions, but seemed to be more prominent within the cSCC cancer cells as opposed to the tumour infiltrating lymphocytes and stromal fibroblasts present in the stroma surrounding tumour nests (Fig. 6*C*). However, PANX1 was visibly reduced in keratinized regions and necrotic tissue as expected (Supplementary Fig. 2*B*). In the adjacent skin, PANX1 was present throughout all the living layers of the epidermis, with relatively higher levels in basal keratinocytes (Fig. 6*C*). Visually increased PANX1 levels were also observed within blood vessels (Fig. 6*C*) and nerve bundles (Supplementary Fig. 2*B*).

To obtain a more quantitative understanding of the relative PANX1 levels in cSCC and normal skin controls, we analyzed our cohort of cSCC and non-adjacent aged skin samples for PANX1 transcript and protein expression. We used non-adjacent normal skin samples as our control for this analysis, since it has been previously shown that PANX1 levels transiently upregulate at the site of cutaneous wounds in mice (Penuela et al., 2014) and thus tumour-adjacent skin may have increased PANX1 levels in response to the cSCC tumour. In aged human full thickness skin samples, PANX1 was present at low levels (Fig. 7*A*). Despite a lack of differences seen in *PANX1* mRNA levels (Fig. 7*B*, *p*=0.925) between tissue types, immunoblotting showed a significant increase in cSCC tumour PANX1 levels compared to

patient-matched normal skin (containing epidermis and dermis only) controls (Fig. 7*C,D*, *p*=0.0011). Looking at our cohort of patient-matched samples (Fig. 7*E*), most sample pairs exhibited the pattern of higher PANX1 in cSCC tumours, but inter-patient variability was noticeable for PANX1 levels in both skin and cSCC samples. We also assessed how PANX1 levels may correlate with tumour recurrence and size. Since 70% of patients had a previous incidence of skin cancer, we analyzed PANX1 expression in tumours which arose at the site of previous skin cancer (local recurrence) or were present at a novel anatomic location (distant recurrence) but found no significant differences in PANX1 levels between sites (Fig. 7*F*, *p*=0.332). Patient tumours were also stratified based on tumour volume and the largest tumour dimension, dividing the 16 cSCC samples in half based on the median value (11.24 cm^2 and 3.7 cm, respectively) to create below median and above median groups. Although no differences were evident based on tumour volume (Fig. 7*G*, *p*=0.873), cSCC patient tumours with the largest dimension above the median value had significantly higher PANX1 levels compared to tumours with their largest dimension below the median (Fig. 7*H*, *p*=0.0377). Overall, this indicates PANX1 is increased in human cSCC tumours, where it localizes to virtually all cell types within the cSCC tumour microenvironment and is associated with larger tumour size.

Discussion

In this study, we found that PANX1 and PANX3 show opposite expression pattern changes in cSCC, with PANX1 increased and PANX3 reduced in this type of keratinocytic tumour compared to patient-matched healthy non-adjacent skin controls. We also determined that in full thickness normal aged skin, PANX1 was present in low abundance whereas PANX3 levels were found at high levels, consistent with reports of low PANX1 and high PANX3 expression in aged mouse dorsal skin (Penuela et al., 2014; O'Donnell et al., 2023). In each tissue type, both pannexins exhibit opposite expression patterns, most likely due to their distinct homeostatic actions in normal skin. Previous findings from our group suggest PANX1 plays more of a proliferative and developmental role in the skin, particularly in keratinocytes of which cSCC arises. PANX1 is prevalent in young murine skin, and basal keratinocytes where it traffics to the cell surface, but normally exhibits an age-related decline and reduction during keratinocyte differentiation where it changes its localization pattern to become more cytosolic (Penuela et al., 2014). This notion is consistent with the cSCC cancer condition where cells are aberrantly

proliferative and their ability to terminally differentiate is compromised. Conversely, PANX3 functions in maintaining epidermal structure and integrity and is present at high levels in aged skin and differentiated keratinocytes where it was found to have an intracellular localization (Cowan et al., 2012; Zhang et al., 2019; Zhang et al., 2021; O'Donnell et al., 2023). This indicates the protein is critical in homeostatic aging processes to ensure proper keratinocyte turnover within the epidermis, and therefore must be downregulated for cSCC carcinogenesis. Additionally, ablating *Panx3* in mice was found to increase inflammatory signalling in the neonatal epidermis and the incidence of dermatitis in an aged cohort (O'Donnell et al., 2023). Since an inflammatory microenvironment has been shown to facilitate cutaneous carcinogenesis (Neagu et al., 2019), reduced PANX3 levels in the skin could promote a pro-tumorigenic environment.

The PANX1 expression findings in our cSCC patient cohort conflict with earlier findings from our group (Cowan et al., 2012) but are consistent with the bioinformatics data from the Human Protein Atlas, and cSCC and HNSCC RNA-seq study data mined from GEO and TCGA, respectively. Inconsistencies were likely due to a smaller sample size in the previous study but may also be due to differences in the tumour stage, where most of the tumours included in this study were advanced. Since PANX1 is upregulated in most other cancer types in which it has been reported, including melanoma, breast cancer and leukemia (Laird & Penuela, 2021), it seems more likely that PANX1 is increased in cSCC compared to normal tissue. For PANX3, trends seen in cSCC corresponded to those observed in DMBA/TPA-induced murine papillomas and carcinomas (Halliwill et al., 2016), previous immunofluorescence microscopy findings from our group (Cowan et al., 2012) and HNSCC *in silico* findings, where *Panx3*/PANX3 levels were reduced in each case compared to controls.

We determined that PANX1 channel activity promoted cancer cell properties such as growth and migration in SCC-13 cells using established PANX1 channel inhibitors PBN and SPL. PANX1 channel activity in SCC-13 cells resembles that of human melanoma cells, whereby blocking PANX1 channels also reduced melanoma cell numbers and motility (Freeman et al., 2019). Although both channel blockers showed the same trends, reductions in SCC-13 tumorigenic properties were more prominent with SPL treatment, most likely due to the increased specificity and potency and decreased off-target effects when inhibiting PANX1 channels with SPL rather than PBN (Koval et al., 2023). Our finding that PANX1 levels are

increased in larger cSCC tumours also corresponds with our SCC-13 inhibition results and suggests that PANX1 channel function may influence cSCC cell proliferation, indicating a role for PANX1 in cSCC tumour progression. PANX1 was found to localize to both the cell surface and intracellularly in SCC-13 cells, indicating it could act as both an ATP release channel at the plasma membrane and/or a calcium leak channel in the endoplasmic reticulum—both of which would be sensitive to SPL inhibition—to promote tumorigenic properties like growth and migration. Additionally, PANX1 was prevalent in many tumour cell types in addition to the cSCC cancer cells such as in blood vessels and immune cells, meaning cross talk between the different cell types within the tumour microenvironment may be possible. Future studies could investigate the potential role of PANX1 in immune infiltration/evasion and angiogenesis in cSCC tumours. As for PANX3, we determined the protein localized predominantly intracellularly in SCC-13 cells, with a small pool trafficking to the cell surface, but we are unsure of the specific organelle compartments where PANX3 is present. Zhang et al. (2019) previously showed that in HaCaT keratinocytes, PANX3 overexpression increased extracellular ATP release and Ca^{2+} release from the endoplasic reticulum, so it is possible these homeostatic functions in keratinocytes are lost with reductions in PANX3 to promote cSCC formation. However, since our assessment on the effects of PANX3 on cSCC formation was performed using a global *Panx3* KO mouse model, it remains to be seen whether the reduced papilloma incidence and growth with *Panx3* ablation were due to reductions in PANX3 activity, signalling and/or protein interactions. Taken together, this indicates that in cSCC, PANX1 may exhibit pro-tumour activities and PANX3 may exhibit anti-tumour activities.

Despite cSCC being one of the most commonly diagnosed skin cancers with yearly increases in incidence estimated at 3-8% worldwide since the 1960s (Abbas & Kalia, 2015), there are very few tools to study this disease. In our study, experiments were performed using one of the only available cSCC cell lines (SCC-13), which we understand as a limitation since this singular cell line may not be representative of the patient population. Moreover, SCC-13 cells were derived from a carcinoma *in situ* (Rheinwald & Beckett, 1981) which is an early, more benign form of cSCC (Adalsteinsson et al., 2021) that more closely resembles a pre-cancerous papilloma lesion (as seen in the mouse carcinoma model) than an advanced cSCC (our cSCC patient sample tumours). Furthermore, there is a lack of epidemiological and expression data for NMSCs, as evidenced by the insufficient bioinformatics data available to be mined for this study,

which is most likely due to the low fatality with NMSCs and lack a NMSC surveillance system or any sort of record keeping of cSCC incidence in many countries (Abbas & Kalia, 2015). This reality has further impeded research into understanding this disease and was part of the rationale for performing this study.

For the chemical carcinoma model, we used our previously established *Panx3* KO mouse model (Moon et al., 2015) in order to assess the potential role of PANX3 in cSCC carcinogenesis since no there are currently no commercially available PANX3-specific channel blockers. Unfortunately, our mice were bred with a C57BL/6 background that has previously shown to be more tumour resistant with the DMBA/TPA protocol (Sundberg et al., 1997). This was evident in our study since none of the treated mice developed carcinomas, which we recognize as a limitation. We were only able to include *N*=5 mice per genotype, but opted to use male mice (instead of females which are normally suggested) (Filler et al., 2007) to be more representative of the human patient population, since men account for a much higher percentage of cSCC incidence than women (Urban et al., 2021). However, because our results were within power, we did not have reason to increase the number of mice used in this study. Lastly, we opted to administer DMBA/TPA treatments to the tail skin of mice to avoid shaving the dorsal skin with every treatment which would irritate or damage the epidermis and potentially affect results interpretation. Furthermore, murine tail skin is more similar to human thin skin since it contains less hair follicles and has a thicker epidermis than that of mouse dorsal skin, making it a more suitable selection to model human cSCC carcinogenesis (Xie & Chen, 2022).

Overall, through an *in vitro* investigation of PANX1 channel blocking, an *in vivo* characterization of a chemical carcinoma model in *Panx3* KO mice, and an analysis of patient-derived tissue, we observed that PANX1 and PANX3 dysregulation may have potential tumour promoting or suppressive effects for keratinocyte transformation into cSCC, respectively. These findings offer the potential for PANX1 and PANX3 as new therapeutic targets in cSCC treatment, specifically for advanced disease.

References

Abbas M. & Kalia S. (2015). Trends in non-melanoma skin cancer (basal cell carcinoma and squamous cell carcinoma) in canada: A descriptive analysis of available data. *Journal of Cutaneous Medicine and Surgery*, **20**(2), 166-175.

Abel E. L., Angel J. M., Kiguchi K. & DiGiovanni J. (2009). Multi-stage chemical carcinogenesis in mouse skin: Fundamentals and applications. *Nature Protocols*, **4**(9), 1350-1362.

Abitbol J., O'Donnell B., CB W., Jewlal E., Kelly J., Barr K., Willmore K., Allman B. & Penuela S. (2019). Double deletion of panx1 and panx3 affects skin and bone but not hearing. *Journal of Molecular Medicine*, **97**(5), 723-726.

Aboul-Fettouh N., Morse D., Patel J. & Migden M. R. (2021). Immunotherapy and systemic treatment of cutaneous squamous cell carcinoma. *Dermatology Practical & Conceptual*, **11**(Suppl 2), e2021169S.

Adalsteinsson J. A., Olafsdottir E., Ratner D., Waldman R., Feng H., Ungar J., Silverberg J. I., Kristjansson A. K., Jonasson J. G. & Tryggvadottir L. (2021). Invasive and in situ squamous cell carcinoma of the skin: A nationwide study in iceland. *British Journal of Dermatology*, **185**(3), 537-547.

Bankhead P., Loughrey M. B., Fernandez J. A., Dombrowski Y., McArt D. G., Dunne P. D., McQuaid S., Gray R. T., Murray L. J., Coleman H. G., James J. A., Salto-Tellez M. & Hamilton P. W. (2017). Qupath: Open source software for digital pathology image analysis. *Scientific Reports*, **7**(1), 16878.

Bao L., Locovei S. & Dahl G. (2004). Pannexin membrane channels are mechanosensitive conduits for atp. *FEBS Letters*, **572**(1-3), 65-68.

Celetti S. J., Cowan K. N., Penuela S., Shao Q., Churko J. & Laird D. W. (2010). Implications of pannexin 1 and pannexin 3 for keratinocyte differentiation. *Journal of Cell Science*, **123**(8), 1363-1372.

Colaprico A., Silva T. C., Olsen C., Garofano L., Cava C., Garolini D., Sabedot T. S., Malta T. M., Pagnotta S. M., Castiglioni I., Ceccarelli M., Bontempi G. & Noushmehr H. (2016). Tcgabiolinks: An r/bioconductor package for integrative analysis of tcga data. *Nucleic Acids Research*, **44**(8), e71.

Cowan K. N., Langlois S., Penuela S., Cowan B. J. & Laird D. W. (2012). Pannexin1 and pannexin3 exhibit distinct localization patterns in human skin appendages and are regulated

during keratinocyte differentiation and carcinogenesis. *Cellular Communication & Adhesion*, **19**(3-4), 45-53.

Edgar R., Domrachev M. & Lash A. E. (2002). Gene expression omnibus: Ncbi gene expression and hybridization array data repository. *Nucleic Acids Research*, **30**(1), 207-210.

Fahradyan A., Howell C., Wolfswinkel E., Tsuha M., Sheth P. & Wong A. (2017). Updates on the management of non-melanoma skin cancer (nmsc). *Healthcare (Basel)*, **5**(82).

Filler R. B., Roberts S. J. & Girardi M. (2007). Cutaneous two-stage chemical carcinogenesis. *Cold Spring Harbor Protocols*.

Firnhaber J. (2012). Diagnosis and treatment of basal cell and squamous cell carcinoma. *American Family Physician*, **86**, 161-168.

Freeman T. J., Sayedyahossein S., Johnston D., Sanchez-Pupo R. E., O'Donnell B., Huang K., Lakhani Z., Nouri-Nejad D., Barr K. J., Harland L., Latosinsky S., Grant A., Dagnino L. & Penuela S. (2019). Inhibition of pannexin 1 reduces the tumorigenic properties of human melanoma cells. *Cancers (Basel)*, **11**(102).

Good M. E., Chiu Y. H., Poon I. K. H., Medina C. B., Butcher J. T., Mendu S. K., DeLalio L. J., Lohman A. W., Leitinger N., Barrett E., Lorenz U. M., Desai B. N., Jaffe I. Z., Bayliss D. A., Isakson B. E. & Ravichandran K. S. (2018). Pannexin 1 channels as an unexpected new target of the anti-hypertensive drug spironolactone. *Circulation Research*, **122**(4), 606-615.

Halliwill K., Quigley D., Chung Kang H., Del Rosario R., Ginzinger D. & Balmain A. (2016). Panx3 links body mass index and tumorigenesis in a genetically heterogeneous mouse model of carcinogen-induced cancer. *Genome Medicine*, **8**, 83-100.

Ishikawa M., Iwamoto T., Nakamura T., Doyle A., Fukumoto S. & Yamada Y. (2011). Pannexin 3 functions as an er ca 2+ channel, hemichannel, and gap junction to promote osteoblast differentiation. *The Journal of Cell Biology*, **193**(7), 1257-1274.

Iwamoto T., Nakamura T., Doyle A., Ishikawa M., De Vega S., Fukumoto S. & Yamada Y. (2010). Pannexin 3 regulates intracellular atp/camp levels and promotes chondrocyte differentiation. *The Journal of Biological Chemistry*, **285**, 18948-18958.

Kallini J., Hamed N. & Khachemoune A. (2015). Squamous cell carcinoma of the skin: Epidemiology, classification, management, and novel trends. *International Journal of Dermatology*, **54**, 130–140.

Koval M., Schug W. J. & Isakson B. E. (2023). Pharmacology of pannexin channels. *Current Opinion in Pharmacology*, **69**, 102359.

Laird D. W. & Penuela S. (2021). Pannexin biology and emerging linkages to cancer. *Trends in Cancer*, **7**(12), 1119-1131.

Lomas A., Leonardi-Bee J. & Bath-Hextall F. (2012). A systematic review of worldwide incidence of nonmelanoma skin cancer. *British Journal of Dermatology*, **166**, 1069-1080.

Madan V., Lear T. & Szeimies R. (2010). Non-melanoma skin cancer. *The Lancet*, **375**, 637-685.

Migden M. R., Rischin D., Schmults C. D., Guminski A., Hauschild A., Lewis K. D., Chung C. H., Hernandez-Aya L., Lim A. M., Chang A. L. S., Rabinowits G., Thai A. A., Dunn L. A., Hughes B. G. M., Khushalani N. I., Modi B., Schadendorf D., Gao B., Seebach F., Li S., Li J., Mathias M., Booth J., Mohan K., Stankevich E., Babiker H. M., Brana I., Gil-Martin M., Homsi J., Johnson M. L., Moreno V., Niu J., Owonikoko T. K., Papadopoulos K. P., Yancopoulos G. D., Lowy I. & Fury M. G. (2018). Pd-1 blockade with cemiplimab in advanced cutaneous squamous-cell carcinoma. *New England Journal of Medicine*, **379**(4), 341-351.

Moon P. M., Penuela S., Barr K., Khan S., Pin C. L., Welch I., Attur M., Abramson S. B., Laird D. W. & Beier F. (2015). Deletion of panx3 prevents the development of surgically induced osteoarthritis. *Journal of Molecular Medicine*, **93**(8), 845-856.

Neagu M., Constantin C., Caruntu C., Dumitru C., Surcel M. & Zurac S. (2019). Inflammation: A key process in skin tumorigenesis. *Oncology Letters*, **17**, 4068-4084.

Nissinen L., Farshchian M., Riihilä P. & Kähäri V. (2016). New perspectives on role of tumor microenvironment in progression of cutaneous squamous cell carcinoma. *Cell and Tissue Research*, **365**, 691-702.

Nouri-Nejad D., O'Donnell B. L., Patil C. S., Sanchez-Pupo R. E., Johnston D., Sayedyahossein S., Jurcic K., Lau R., Gyenis L., Litchfield D. W., Jackson M. F., Gloor G. B. & Penuela S. (2021). Pannexin 1 mutation found in melanoma tumor reduces phosphorylation, glycosylation, and trafficking of the channel-forming protein. *Molecular Biology of the Cell*, **32**(5), 376-390.

O'Donnell B. L. & Penuela S. (2023). Skin in the game: Pannexin channels in healthy and cancerous skin. *Biochemical Journal*, **480**(23), 1929-1949.

O'Donnell B. L., Sanchez-Pupo R. E., Sayedyahossein S., Karimi M., Bahmani M., Zhang C., Johnston D., Kelly J. J., Wakefield C. B., Barr K., Dagnino L. & Penuela S. (2023). Panx3 channels regulate architecture, adhesion, barrier function, and inflammation in the skin. *Journal of Investigative Dermatology*, **143**, 1509-1519.

Panchin Y., Kelmanson I., Matz M., Lukyanov K., Usman N. & Lukyanov S. (2000). A ubiquitous family of putative gap junction molecules. *Current Biology*, **10**(13), 473-474.

Penuela S., Bhalla R., Gong X.-Q., Cowan K. N., Celetti S. J., Cowan B. J., Bai D., Shao Q. & Laird D. W. (2007). Pannexin 1 and pannexin 3 are glycoproteins that exhibit many distinct characteristics from the connexin family of gap junction proteins. *Journal of Cell Science*, **120**(21), 3772-3783.

Penuela S., Bhalla R., Nag K. & Laird D. (2009). Glycosylation regulates pannexin intermixing and cellular localization. *Molecular Biology of the Cell*, **20**, 4313–4323.

Penuela S., Gyenis L., Ablack A., Churko J. M., Berger A. C., Litchfield D. W., Lewis J. D. & Laird D. W. (2012). Loss of pannexin 1 attenuates melanoma progression by reversion to a melanocytic phenotype. *The Journal of Biological Chemistry*, **287**(34), 29184-29193.

Penuela S., Kelly J. J., Churko J. M., Barr K. J., Berger A. C. & Laird D. W. (2014). Panx1 regulates cellular properties of keratinocytes and dermal fibroblasts in skin development and wound healing. *Journal of Investigative Dermatology*, **134**(7), 2026-2035.

Ratushny V., Gober M., Hick R., Ridky T. & Seykora J. (2012). From keratinocyte to cancer: The pathogenesis and modeling of cutaneous squamous cell carcinoma. *The Journal of Clinical Investigation*, **122**, 464-472.

Rheinwald J. G. & Beckett M. A. (1981). Tumorigenic keratinocyte lines requiring anchorage and fibroblast support cultured from human squamous cell carcinomas. *Cancer Research*, **41**, 1657-1663.

Sayedyahossein S., Huang K., Li Z., Zhang C., Kozlov A. M., Johnston D., Nouri-Nejad D., Dagnino L., Betts D. H., Sacks D. B. & Penuela S. (2021). Pannexin 1 binds beta-catenin to modulate melanoma cell growth and metabolism. *Journal of Biological Chemistry*, 100478.

Schmults C. D., Karia P. S., Carter J. B., Han J. & Qureshi A. A. (2013). Factors predictive of recurrence and death from cutaneous squamous cell carcinoma. *JAMA Dermatology*, **149**(5), 541-547.

Sundberg J. P., Sundberg B. A. & Beamer W. G. (1997). Comparison of chemical carcinogen skin tumor induction efficacy in inbred, mutant, and hybrid strains of mice: Morphologic variations of induced tumors and absence of a papillomavirus cocarcinogen. *Molecular Carcinogenesis*, **20**, 19-32.

Urban K., Mehrmal S., Uppal P., Giesey R. L. & Delost G. R. (2021). The global burden of skin cancer: A longitudinal analysis from the global burden of disease study, 1990-2017. *JAAD International*, **2**, 98-108.

Wasserstein R. L., Schirm A. L. & Lazar N. A. (2019). Moving to a world beyond "p < 0.05". *The American Statistician*, **73**(sup1), 1-19.

Xie Y. & Chen T. (2022). Mouse tail skin wholemount staining and intravital calcium imaging. *STAR Protocols*, **3**(1), 101235.

Yan W., Wistuba I. I., Emmert-Buck M. R. & Erickson H. S. (2011). Squamous cell carcinoma - similarities and differences among anatomical sites. *Am J Cancer Res*, **1**(3), 275-300.

Ye J., Coulouris G., Zaretskaya I., Cutcutache I., Rozen S. & Madden T. L. (2012). Primer-blast: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, **13**, 134.

Zhang P., Ishikawa M., Doyle A., Nakamura T., He B. & Yamada Y. (2021). Pannexin 3 regulates skin development via epiprofin. *Scientific Reports*, **11**(1), 1779.

Zhang P., Ishikawa M., Rhodes C., Doyle A., Ikeuchi T., Nakamura K., Chiba Y., He B. & Yamada Y. (2019). Pannexin-3 deficiency delays skin wound healing in mice due to defects in channel functionality. *Journal of Investigative Dermatology*, **139**(4), 909-918.

Additional information

Data availability statement
All raw data is available upon request.

Competing interests
The authors have no competing interests to declare.

Author contributions
Conceptualization: B.O'D., D.J., L.D., S.L., K.R., S.P.; Data Curation: Z.K.; Formal Analysis: B.O'D., Z.K.; Funding Acquisition: S.P.; Investigation: B.O'D., D.J., A.Bh., Z.K., D.S., A.By. S.S.; Methodology: B.O'D., D.J., S.P.; Project Administration: B.O'D., D.J., S.P.; Resources: S.L., M.C., K.R., S.P.; Software: Z.K.; Supervision: L.D., S.L., K.R., S.P.; Visualization: B.O'D., A.Bh., Z.K., D.S., A.By. M.C.; Writing – original draft: B.O'D., S.P.; Writing – review and editing: all authors.

Funding
This work was supported by a Canadian Institutes of Health Research Project grant (FRN 153112) and a London Regional Cancer Program Catalyst Grant to S.P.

Acknowledgements
We would like to thank Dr. Rafael Sanchez-Pupo and Dr. Brent Wakefield (University of Western Ontario) for their assistance with statistical analyses. We would also like to thank the Pathology and Laboratory Medicine Translational Research Service at London Health Sciences Centre for providing tumour tissue slides and Linda Jackson-Boeters (University of Western Ontario) for her assistance with the DAB staining protocol. Lastly, we thank Dr. Kathryn Roth's patients who provided samples for this study.

Keywords
PANX1, PANX3, cutaneous squamous cell carcinoma, skin

Tables

Table 1. Genotyping and RT-qPCR primers

Primer	Forward sequence (5'–3')	Reverse sequence (5'–3')	Reference
Panx3 R1 (genotyping)	CGCAGCATTCCGGAACCTGAG	AACTAGCCGGAGGGGTCG	(Abitbol et al., 2019)
Panx3 RT (genotyping)	CGCAGCATTCCGGAACCTGAG	CTTGTCCTGCGTATGGTG	(Abitbol et al., 2019)
PANX1	AACCGTGCAATTAAGGCTG	GGCTTTCAGATACCTCCCAC	(Sayedyahossein et al., 2021)
PANX3	ATGTGAAAGCTGGAGCCGAA	TTGCCTCACTTGCTCTCTGG	(Ye et al., 2012)
β2M	GTATGCCTGCCGTGTGAACC	AAGCAAGCAAGCAGAATTTGGA	(Ye et al., 2012)

Table 2. Patient demographics and clinical information

Characteristic	N = 20
Average age (years)	75.1
Age range (years)	51 – 93
Male	17 (85%)
Female	3 (15%)
Immunosuppression	6 (30%)
Previous skin cancer	14 (70%)
Tumour location	
Scalp	6 (30%)
Ear	4 (20%)
Cheek	3 (15%)
Temple	2 (10%)
Lip	2 (10%)
Forehead	1 (5%)
Chin	1 (5%)
Neck	1 (5%)
Tumour characteristics	
Average size in largest dimension (cm)	3.2
Largest dimension range (cm)	0.6 – 5.7
Local recurrence	7 (35%)
Stage	
1	1 (5%)
2	1 (5%)
3	6 (30%)
4	12 (60%)
AJCC tumour stage	
T1N0M0	2 (10%)
T2N0M0	1 (5%)
T3N0M0	13 (65%)
T3N1M0	2 (10%)
T3N3aM0	1 (5%)
T3N0M1	1 (5%)
Degree of differentiation	
Poor	3 (15%)
Moderate	15 (75%)
Well	1 (5%)
In situ	1 (5%)
Degree of invasion	
Deep	12 (60%)
Perineural	7 (35%)
Lymphovascular	3 (15%)
Normal Skin Location	
Clavicle	7 (35%)
Neck	5 (25%)
Leg	5 (25%)
Arm	2 (10%)
Cheek	1 (5%)

AJCC, American Joint Committee on Cancer; T, tumour; N, node; M, metastasis.

Table 3. Additional patient demographics, including comorbidities, medications, and clinical outcomes

Characteristic	N = 20
Comorbidities	
Cardiovascular (HTN, dyslipidemia, MI, CAD, heart transplant, AAA, arrhythmia, valvular, CHF, angina)	16 (80%)
Endocrine (DM)	9 (45%)
GI (GERD, IBD, CRC, diverticulosis)	7 (35%)
GU (CKD, transplants, bladder cancer, BPH)	4 (20%)
MSK (OA, RA, gout)	4 (20%)
Neurologic (peripheral neuropathy, stroke, hearing loss, Parkinson's)	4 (20%)
Hematologic (polycythemia, anemia, previous PE, DVT)	4 (20%)
Respiratory (asthma, COPD, ILD)	3 (15%)
Previous cancers (prostate, bladder, colon)	2 (10%)
Immunosuppression	
Transplant	2 (10%)
Rheumatoid arthritis	1 (5%)
CNS vasculitis	1 (5%)
Crohn's disease	1 (5%)
Medications	
Metabolic syndrome (anti-hypertensives, statins, anti-platelets, anti-coagulants)	16 (80%)
Diabetes (anti-hyperglycemics)	8 (40%)
Immune suppressants (steroids, mycophenolate, tacrolimus, methotrexate, biologics)	6 (30%)
Pain control (pregabalin, tramadol, nortriptyline, diclofenac, acetaminophen, oxycocet)	5 (25%)
Anti-microbials (atovaquone, tobramycin)	1 (5%)
Clinical outcomes	
Adjuvant radiation	5 (25%)
Local recurrence	5 (25%)
Regional recurrence	4 (20%)
Immunotherapy	3 (15%)
Metastasis	2 (10%)
Mortality	0 (0%)

HTN, hypertension; MI, myocardial infarction; CAD, coronary artery disease; AAA, abdominal aortic aneurysm; CHF, congestive heart failure; DM, diabetes mellitus; GI, gastrointestinal; GERD, gastro-esophageal reflux; IBD, inflammatory bowel disease (Crohn's or ulcerative colitis); CRC, colorectal cancer; GU, genitourinary; CKD, chronic kidney disease; BPH, benign prostatic hypertrophy; MSK, musculoskeletal; OA, osteoarthritis; RA, rheumatoid arthritis; PE, pulmonary embolism; DVT, deep vein thrombosis; DBS, deep brain stimulation; COPD, chronic obstructive pulmonary disease; ILD, interstitial lung disease.

Figure Legends

Figure 1. SCC-13 cells express PANX1 and PANX3 with varying localizations

Immunoblotting assessed PANX1 (*A*, *N*=4) and PANX3 (*B*, *N*=3) protein levels in SCC-13 cells. U-2 OS cells ectopically expressing PANX3 (+) as positive control. GAPDH as protein loading control, protein sizes in kDa. *C*, confocal micrographs of PANX1 (turquoise) localization in SCC-13 and SCC-13+PANX1 cells (*N*=3). *D*, immunofluorescence of PANX3-HA (turquoise) localization in SCC-13 and SCC-13+PANX3-HA cells (N=3). Nuclei (blue) were counterstained with Hoechst 33342. Coverslips incubated with Alexa Fluor™ 488 goat anti-rabbit secondary antibody only as a secondary only control. Scale bar: 10μm.

Figure 2. PANX1 channel inhibition reduces SCC-13 growth and migration

SCC-13 cell growth was measured over a period of 4 days for cells treated with either 1 mM PBN (*A*), 20 µM SPL (*B*) or corresponding vehicle control (water and EtOH, respectively). $*p<0.05$; $**p<0.01$; $***p<0.001$; $****p<0.0001$. SCC-13 % wound closure was measured over a period of 2 days after treatment with either 1 mM PBN (*C*), 20 µM SPL (*D*) or corresponding vehicle control. Two-way repeated measures ANOVA followed by a Sidak's multiple comparisons test for both growth and migration. Bars indicate mean ± SD. Representative phase contrast images of scratch wounds in SCC-13 cells treated with either 1 mM PBN (*E*), 20 µM SPL (*F*) or corresponding vehicle control at 0 hr, 24 hr and 46 hr timepoints. Dotted white line outlines scratch. Scale bar: 200 µm. *G*, quantifications of 24 hr and 46 hr timepoints for SCC-13 cells treated with either water or 1 mM PBN showed a reduced trend at 24 hr ($p=0.0757$) and significant reduction for % wound closure at 46 hr ($p=0.0066$) with PBN treatment. *H*, SPL (20 µM) treatment of SCC-13 cells significantly reduced % wound closure at 24 hr ($p=0.0233$) and 46 hr ($p=0.0025$). Unpaired t test. Bars indicate mean ± SEM. *N*=3.

Figure 3. DMBA/TPA treated *Panx3* KO mice tend to have increased papilloma incidence, volumes and growth over time

A, timeline of experimental design. Eight-week-old WT and *Panx3* KO mice were subjected to one dose of 25 µg DMBA treatment (week 0), followed by 28 weeks of twice weekly treatments of 4 µg TPA. Twenty-eight weeks after the initial DMBA treatment, mice were sacrificed. *B*,

images of papillomas that developed on treated mice. Papilloma indicated by white arrow. *C*, KO mice had a slightly greater average papilloma incidence of 2.2 papillomas per mouse compared to the average of 1 papilloma per WT control, with 3/5 WT and 5/5 KO mice developing papillomas. *D*, survival curves for time to papilloma onset were similar between genotypes (*p*=0.176). Log-rank test. *N*=5 for WT, KO mice for panels *A-D*. *E*, papilloma growth curves for average papilloma volume over time. Individual lines depict measurements per mouse. *F*, although only significant at week 28 (**p*=0.0416), average papilloma volume over time showed moderate evidence to an increase in KO mice. Two-way repeated-measures ANOVA and multiple unpaired t tests. *G*, area under the curve (AUC) measurements for average papilloma volume over time showed moderate evidence to an increase in KO mice that was not statistically significant (*p*=0.1106). Unpaired t test. *H*, average papilloma volume at endpoint (week 28) was significantly increased in KO mice compared to controls (*p*=0.0491). Unpaired t test. *N*=4 for WT, *N*=5 for KO in panels *E-H*. Bars represent mean ± SEM.

Figure 4. *PANX1* mRNA is upregulated in cSCC and HNSCC tumours compared to corresponding control tissue

A, table showcases the normalized expression levels of *PANX1* and *PANX3* mRNA in cSCC samples in comparison to normal skin control samples (either human skin organoids or unmatched healthy controls). RNA-seq expression data was derived from the GEO gene expression studies GSE191334 and GSE139505. *p* adj, *p* value adjusted. *B*, table outlines the normalized expression levels of *PANX1* and *PANX3* in HNSCC tumour samples compared to normal adjacent tissue controls. Data was mined from TCGA database. FDR, false discovery rate. *C*, the z-score normalized transcript expression levels of *PANX1* and *PANX3* in TCGA_HNSCC RNA-seq samples compared to normal tissue show a marked increase in *PANX1* but a decrease in *PANX3* in HNSCC tumours compared to controls.

Figure 5. *PANX3* mRNA levels are reduced in cSCC tumour fragments compared to patient-matched aged skin

A, immunoblotting indicates PANX3 is present in aged full thickness human skin samples (*N*=4). *B*, *PANX3* transcripts were significantly reduced (*p*=0.0452, *N*=8) in cSCC tumour fragments compared to aged skin (minus hypodermis) via RT-qPCR. β-2-microglobulin (*β2M*) was used to

calculate normalized mRNA using the ΔΔCT method. *C*, PANX3 levels were assessed via Western blot in cSCC tumour fragments (T) compared to patient-matched aged skin (S, epidermis, and dermis only). Each # corresponds to a different patient. *N*=14. U-2 OS cells expressing a PANX3 plasmid as positive control for PANX3 (+). T_O, original tumour; T_R, recurrent tumour. GAPDH used as protein loading control, protein sizes in kDa. Quantifications of total PANX3 (*D*), PANX3-38K (*E*), PANX3-36K (*F*) and PANX3-32K (*G*) in patient-matched samples in panel *C*. Total PANX3 (*p*=0.0175) and PANX3-36K (*p*=0.0103) were significantly increased in cSCC samples, but PANX3-38K (*p*=0.5612) and PANX3-32K (*p*=0.9708) showed no differences between tissue types. Paired t tests. Bars represent mean ± SEM. Scatter plots with bars show mean values and before-after plots show patient sample matching.

Figure 6. PANX1 localizes to all cSCC tumour cell types

A, images of patient cSCC tumours from the most common anatomic subsites of scalp, ear and cheek (left to right). *B*, representative hematoxylin, and eosin (H&E) and immunohistochemical staining of patient-derived cSCC tumour (T) with adjacent skin (S). Brown DAB stain indicates PANX1 localization. Objective magnification 6.26x. Scale bar: 200 µm. *C*, H&E images and corresponding PANX1 immunohistochemistry of cSCC tumour and adjacent skin structures. A representative area of tumour is circled in a dotted line. Stroma region outlined in bracket, with arrowhead indicating a blood vessel. Epidermal layers bracketed, with asterisk (*) indicating basal layer keratinocytes. Objective magnification 22.37x. Scale bar: 50 µm.

Figure 7. PANX1 protein levels are upregulated in human cSCC tumour fragments compared to normal skin

A, Western blotting showed PANX1 is present at very low levels in aged full thickness normal human skin (*N*=4). Hs578T WT cells as positive control and Hs578T *PANX1* KO cells as negative control. *B*, RT-qPCR revealed no differences in *PANX1* transcripts between normal aged skin (minus hypodermis) and patient-matched cSCC tumour fragments (*p*=0.9254, *N*=9). β-2-microglobulin (*β2M*) was used to calculate normalized mRNA using the ΔΔCT method. *C*, immunoblotting assessed PANX1 levels in patient-matched normal aged skin (S, minus hypodermis) and cSCC tumour fragments (T) where each # corresponds to a different patient. *D*, quantification showed PANX1 levels are significantly increased in cSCC tumour fragments

compared to normal skin (N=16, p=0.0011). *E*, Western blots of all samples used for PANX1 quantification (*C*) show variability between patients. Quantifications excluded degraded tumour samples (and corresponding normal skin), evidenced by a lack of GAPDH expression (protein loading control). GBM17 cells as positive control for PANX1 (+). T_o, original tumour; T_R, recurrent tumour; T_S, tumour from scalp; T_C, tumour from cheek. Protein sizes in kDa. *F*, PANX1 levels were not significantly different (p=0.3323) in cSCC tumour fragments arising at local or distant recurrence sites from patients with a previous incidence of skin cancer. *G*, cSCC tumours above and below the median tumour volume of 11.24 cm^2 showed no differences in PANX1 levels (p=0.8734). *H*, cSCC tumours above the median largest tumour dimension of 3.7 cm had significantly increased PANX1 compared to those below the median value (p=0.0377). Paired t tests. Bars represent mean ± SEM. Scatter plots with bars show mean values and before-after plots show patient sample matching.

A cSCC

Study	Transcript	Log2 Fold Change	Fold Change	p value	p adj	N value Tumour	N value Control
GSE191334	*PANX1*	0.3187	+1.25	0.2190	0.7165	8	8
GSE191334	*PANX3*	1.0940	+2.13	0.7188	0.9952	8	8
GSE139505	*PANX1*	1.4062	+2.65	2.35×10^{-10}	3.23×10^{-9}	9	7
GSE139505	*PANX3*	-	-	-	-	9	7

B HNSCC

Transcript	Log2 Fold Change	Fold Change	FDR	p value	N value Tumour	N value Control
PANX1	1.1368	+2.20	1.54×10^{-12}	6.54×10^{-14}	502	44
PANX3	-0.3453	-1.27	0.6549	0.5752	502	44

C HNSCC

PANX1

PANX3