TITLE PAGE

TITLE: Histones are exosome membrane proteins regulated by cell stress

AUTHORS:

Birendra Singh[1][§], Marcus Fredriksson Sundbom[1][§], Uma Muthukrishnan[2][†], Balasubramanian

Natarajan[2][#], Stephanie Stransky[3], André Görgens[4,5,6], Joel Z. Nordin[4], Oscar P. B. Wiklander[4], Linda

Sandblad[7], Simone Sidoli[3,] Samir EL Andaloussi[4], Michael Haney[1*], and Jonathan D. Gilthorpe[2*]

[§]Contributed equally and should be considered as first authors.

*Considered as co-senior authors.

[1]Department of Diagnostics and Intervention, Anaesthesiology and Intensive Care, Umeå University,
Umeå, 90 185. Sweden.

[2]Department of Medical and Translational Biology, Umeå University, Umeå 90 187, Sweden.

[3]Department of Biochemistry, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

[4]Department of Laboratory Medicine, Division of Biomolecular and Cellular Medicine, Karolinska
Institute, Stockholm, Sweden, 141 86 Stockholm, Sweden.

[5]Department of Cellular Therapy and Allogeneic Stem Cell Transplantation (CAST), Karolinska
University Hospital Huddinge and Karolinska Comprehensive Cancer Center, Stockholm, Sweden
21
22 [6]Institute for Transfusion Medicine, University Hospital Essen, University of Duisburg-Essen, Essen,
23 Germany
24
25 [7]Department of Chemistry, Umeå University, Umeå 90 187, Sweden.

26 *Authors for correspondence: michael.haney@umu.se and jonathan.gilthorpe@umu.se

ABSTRACT

Histones are conserved nuclear proteins that function as part of the nucleosome in the regulation of chromatin structure and gene expression. Interestingly, extracellular histones populate biofluids from healthy individuals and when elevated may contribute to various acute and chronic diseases. It is generally assumed that most extracellular histones exist as nucleosomes, as components of extracellular chromatin. We analysed cell culture models under normal and stressed conditions to identify pathways of histone secretion. We report that core and linker histones localize to extracellular vesicles (EVs) and are secreted via the multivesicular body/exosome pathway. Upregulation of histone EV secretion occurs in response to cellular stress, with enhanced vesicle secretion and a shift towards a population of smaller EVs. Most histones were membrane associated with the outer surface of EVs. Degradation of EV-DNA did not impact significantly on EV-histone association. Individual histones or histone octamers bound strongly to liposomes and EVs, but nucleosomes did not, showing histones do not require DNA for EV binding. EV histones colocalized most frequently with the tetraspanin CD63 but using genetic or pharmacological intervention, we found that all known pathways of exosome biogenesis acted positively on histone secretion. Inhibition of autophagy and lysosomal degradation had a strong positive effect on EV histone release. Unexpectedly, EV-associated histones lacked the extensive post-translational modification of their nuclear counterparts, suggesting loss of PTMs may be involved in their trafficking or secretion. Our data does not support a significant role for EV-histones existing as nucleosomes. We show for the first time that histones are secreted from cells as membrane proteins via EVs/exosomes. This fundamental discovery provides support for further investigation of the biological activity of exosome associated histones and their role in disease.

280 words

KEY WORDS: Histone, extracellular vesicles, exosome, membrane associated proteins, cellular stress, post translational modification.

INTRODUCTION

Histones are basic proteins[1] abundant in the eukaryotic nucleus. The core histones (H2A, H2B, H3 and H4) associate with DNA to form the nucleosome[2, 3] and together with linker histone H1 they form chromatosomes[4]. Hence, the overwhelming bulk of investigation into the localization and function of histone proteins has focused on their roles in chromatin organization and gene regulation within the nuclear compartment. Functions for cytoplasmic and extracellular histones have also been described[5, 6]. Perhaps the best understood paradigm for histone release by activated cells is in association with modified chromatin during neutrophil extracellular trap (NET) formation, by a specific pathway of regulated cell death known as 'NETosis'[7-11]. However, our understanding of pathways for histone secretion in non-activated cells is lacking.

Elevated levels of extracellular histones are found in biofluids from critically ill patients with conditions such as acute lung injury/acute respiratory distress syndrome (ALI)/(ARDS), sepsis, trauma, severe COVID-19 infection and mediate inflammation through toll-like receptor (TLR)-4 and TLR-2 signaling[12-19]. Increased circulating histones have been implicated in poor survival as well as in organ dysfunction[13, 20-22], mechanical ventilation, secondary infection[23], thromboembolic events and acute kidney injury[13, 14, 20-24]. The toxic effects of extracellular histones can be meliorated, for example by degradation using activated protein C, or blocking using anti-histone monoclonal antibodies or C1 esterase inhibitors. These approaches have indicated therapeutic potential, with increased survival and reduced organ dysfunction[24, 25]. However, relatively low levels of circulating histones are also found in biofluids under normal physiological conditions[26-29]. This raises important question of how histones reach the extracellular environment and whether histones are normally secreted by healthy cells, as well as whether there is a function for extracellular/circulating histones in the non-disease state.

Histones are frequently identified as components of extracellular vesicle (EV) proteomes[30-32], including microvesicles/ectosomes or apoptotic vesicles (generated via outward budding of the plasma membrane), apoptotic bodies (released via membrane blebbing), amphisomes[33], and exosomes[30, 34, 35]. Exosomes are endosome-derived vesicles[36] formed as intraluminal vesicles from inward budding within multivesicular bodies (MVBs) and released from cells upon the fusion of an MVB with the plasma membrane[37]. They carry numerous proteins[34], bioactive lipids[38] RNAs[39] and DNA[40]. Two basic pathways for exosome biogenesis have

been defined; an endosomal sorting complex required for transport (ESCRT)-dependent and

an ESCRT-independent pathway requiring ceramide[41, 42]. It is unclear whether extracellular

histones are associated with a particular type of EV, whether they localize to the surface or

within EVs, or how they are trafficked to a secretory pathway.

The presence of histones in purified EVs has been suggested to be due to contamination with

apoptotic vesicles that co-purify[34], or fusion of EVs with free histones during

ultracentrifugation[33]. Another suggestion has been that histones are sorted into the lumen of

EVs with exosome DNA (exoDNA) in a nucleosome-like configuration[31, 43], with

micronuclei into exosomes[44], or are secreted via mitochondrial derived EVs[6]. Hence, the

dominant viewpoint is that EV associated histones are nucleosomes, or chromatin associated

particles that bind to EVs[30, 31]. Whether this is the case for all EV histones, or all cell types,

remains an important question.

We have previously identified histones as secretory components of injured rat brain tissue[45].

Significant progress has been made in understanding pathways of EV/exosome biogenesis

using rodent brain oligodendrocyte cell lines[46-48]. Oligodendrocyte EVs contain both core and

linker histones[47] and can be grown readily in 2-dimensional (2D) or 3D cell culture systems

under serum free conditions to facilitate the purification a sufficient quantity of EVs for the

biochemical characterization of cargo proteins, even for histones where there is a significant

risk of contamination with abundant cellular proteins such as histones[49]. Hence, we used this

as a starting point to define whether and where histones localize to EVs and to define how

they are secreted.

In this study, we used the rat oligodendrocyte progenitor cell line OLN-93[50] as a model to

elucidate the pathways for EV-mediated histone secretion. Using a combination of

experimental strategies, we show for the first time that core and linker histones are present in

MVBs and released via exosomes. Histone secretion was upregulated in response to different

stress paradigms. With stress there was a shift to a population of EVs of smaller diameter and

higher floating density. Surprisingly, most EV histones were strongly associated with the EV

membrane and interacted directly with membrane lipids in liposomes. In accordance with a

previous study[30], the majority EV-DNA associated with the EV surface. However, we found

that EV-DNA and EV-histones could be degraded independently. Both ESCRT-dependent

and –independent, as well as autophagosome-lysosomal pathways, were involved in the

routing of histones for secretion. Using imaging flow cytometry, we could show both H3 and

H4 colocalize to the surface of tetraspanin positive EVs, predominantly a CD63 positive

population. Intriguingly, secreted EV associated histones lacked post-translational

modifications, distinguishing them from their nuclear, chromatin associated, counterparts.

MATERIALS AND METHODS

All materials and reagents were obtained from Thermo Fisher Scientific or Merck/Sigma-

Aldrich unless otherwise stated.

Cell lines and culture

Cells were maintained in a humidified incubator at 37°C/5% (v/v) CO_2. OLN-93[50],

HEK/HEK293T (ATCC CRL-3216/CRL-1573TM) and NSC-34 (CEDARLANE) cells were

cultured in high glucose Dulbecco's modified Eagle's medium (DMEM). HeLa (ATCC CCL-

2) and Caco-2 (ATCC HTB-37) cells were maintained in in Eagle's Minimum Essential

Media, AC16 (ATCC CRL-3568) in DMEM:F12 and A549 (ATCC CCL-185) in F12K.

Complete media were routinely supplemented with 2 mM L-glutamine, 10% (v/v) heat

inactivated FBS with 100 IU/ml penicillin/streptomycin.

An OLN-93[EGFP-CD63] stable cell line (EGFP fused to the intraluminal N-terminus of human

CD63)[51] was established by selection in 1 mg/ml G418 sulphate[52]. A construct for mouse

H3.1 (Hist1h3a, P68433) was generated by RT-PCR from mouse cDNA and cloned (details

available on request) into a modified doxycycline-inducible lentiviral vector pCW57.1 (a gift

from David Root, Addgene plasmid # 41393). A HEK293–H3.1 stable cell line was

generated via transduction of lentiviral particles and selection with 2.5 µg/ml puromycin

(Sigma) for 4 days. Histone expression was induced by addition of 1 µg/ml doxycycline

(Sigma) for 24 h.

Antibodies

For most purposes, including WB and immunostaining, the primary antibodies used were:

Anti-histone H1 (clone AE-4, 05-457), anti-pan histone (clone), anti-H3-CT (07-690), anti-

H2A acidic patch (07-146), anti-H2B (07-371) and anti-H4 (05-858, clone 62-141-13)

purchased from Millipore. Anti-H2A (Novus NBP1-58095), Anti-H3 DyLight 488 (Novus,

NB21-1021G), Anti-H4 FITC (Novus, NB21-2044F), anti-H1 (orb10806, Biorbyt; 39708,

Active Motif), anti-human CD63 (clone 12,435, Fitzgerald), anti-human CD63

(Pierce/Thermo Fisher Scientific MA119281) anti-rat CD63 (NBP2-68077, Novusbio), anti-

CD9 (C-4, Santa Cruz; C9993 Sigma-Aldrich; Novus NBP2-67310), anti-GFP (Clones 7.1

and 13.1, Roche), anti-actin, beta (A1978, Sigma-Aldrich), anti-acetylcholine esterase

(AChE) and anti-TSG101 (ab97299 and ab83, Abcam), anti-GM130 (610823, BD

Transduction), anti-HSP60 (4869, Cell Signaling Technology), anti-hsp70 (SPA-812,

Stressgen Bioreagents), anti-annexin A1 (AF3770, R&D Systems), anti-MVP (JM74-73,

Invitrogen/Thermo Fisher Scientific), anti-Argonaute-2 (EPR10411, Abcam), anti-His tag

(2365S, Cell Signaling Technology), anti-HA tag (C29F4, Cell Signaling Technology).

Western blot

Protein concentrations were estimated using BCA Protein Assay Kit (Pierce/Thermo Fisher

Scientific) and Qubit (Thermo Fisher Scientific) proteins were separated by SDS-PAGE on

AnykD Criterion™ TGX™ precast gels (Bio-Rad) and transferred to 0.2 µm nitrocellulose

membranes by using a Trans-blot Turbo instrument (Bio-Rad). Blots were blocked with 2.5%

BSA in PBS and then incubated with primary antibodies as indicated above, prior to washing

(3x PBS containing 0.001% v/v Tween-20) and detection with HRP-conjugated secondary

antibodies (Vector Labs or Dako), followed by detection with ECL Prime Western Blot

Detection (GE Healthcare). Blots were imaged on an Odyssey®CLx system (LI-COR) or

ChemiDoc XRS+ (Biorad).

EV production and isolation

For 2D culture and EV production in flasks under normal and stress conditions, cells were

seeded in tissue culture flasks coated with 50 µg/ml poly-D-lysine and grown to confluence

in complete media. Cells were washed 3x with warm PBS and growth media was replaced

with serum-free Opti-MEM media, which was conditioned for 24 h. Where necessary, EVs

were concentrated from a large volume of conditioned media by tangential cross flow

diafiltration (100 kDa MWCO, Sartorius).

For 3D culture and EV production at scale under control conditions only, OLN-93 cells

grown in a hollow fibre bioreactor (HFBR; FiberCell Systems, USA). Cells were

seeded/maintained and harvested according to the manufacturer's protocol. Complete DMEM

was used as a reservoir feeding media. The cell compartment of the HFBR was supplemented

with OptiMEM supplemented with 10% (v/v) Exosome-Depleted FBS (Gibco/Thermo Fisher

Scientific). EV harvesting was performed at a regular 12 h interval and overgrowth of the

cells was controlled by flushing the cell compartment, as described in the manufacturer's

instructions. EVs produced using the HFBR were qualitatively indistinguishable from EVs

produced in cell culture flasks.

EVs were isolated from conditioned medium first cleared by sequential centrifugation[53] at

500 xg for 10 min, 2,000 xg for 20 min, 10,000 xg for 30 min (to remove cell debris and

larger microvesicles) followed by a 0.2 µm filtration step. EVs were pelleted by

ultracentrifugation at 100,000-110,000 xg for 90 min (Optima XPN-100 ultracentrifuge using

a SW28 or SW32Ti rotor, Beckman Coulter) at 4°C. EVs were subject to 2 washes in 50 mM

4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES)/150 mM NaCl (pH 7.4) with

recovery at 100,000 xg for 90 min. EVs (referred to as $EV^{100,000\ xg}$) were used directly after

resuspension, or stored in the same buffer at -80°C after snap freezing in liquid nitrogen.

Protein or DNA content were quantified with a Qubit 3.0 fluorometer using the appropriate

Qubit protein, or dsDNA HS assay kit (Life Technologies/Thermo Fisher Scientific).

OptiPrep equilibrium density gradient centrifugation

Iodixanol (OptiPrep) density gradients were used to fractionate $EV^{100,000\ xg}$. In brief, a 12-

36% (w/v) OptiPrep gradient was prepared in 25 mM HEPES (pH 7.4), 150 mM NaCl, 0.25

M sucrose, 1 mM EDTA and 10 mM Tris-HCl (pH 7.4). $EV^{100,000\ xg}$ (300-400 µg based on

protein content) were placed in the bottom 36% layer of a 6-step gradient in a 13.2 ml open

top polyallomer tube (331372, Beckman Coulter). Following ultracentrifugation at 100,000

xg for 18 h/4°C, (SW41 Ti rotor), fractions were collected, and density was estimated by

comparison to a reference gradient using the absorbance at 340 nm. OptiPrep was removed

by washing with 25 mM HEPES (pH 7.4), 150 mM NaCl for 4 hours at 100,000 xg. EV

pellets were resuspended in PBS or HEPES buffer. In some cases, fractions 4-7 were pooled

and in this case are referred to as called $EV^{OptiPrep}$.

Immunofluorescence staining and Duolink Proximity Ligation Assay (PLA)

OLN-93 or HEK293T were grown in complete media on poly-D-lysine coated glass

coverslips and switched into Opti-MEM for 16 h prior to fixation and staining. Formaldehyde

was added to a final concentration of 4% (w/v) and incubated for 15 min at room temperature

(RT). Cells were then washed three times with PBS and blocked in 10% FBS/0.1% (v/v)

Triton-X 100 for 1 h at RT. Primary antibodies were added in blocking solution and

incubated overnight at 4°C. Samples were washed 3x 5 min in PBS/0.1% (v/v) Triton-X 100

followed by addition of Alexa fluor conjugated secondary antibodies. After 1 hour at RT,

coverslips were again washed 3x with and then mounted on glass microscope slides with

Dako Fluorescence Mounting Medium and imaged using a confocal microscope (Nikon

Eclipse C1).

For PLA, permeabilization, blocking and primary antibody incubation was carried out as

described above. DuoLink PLA (Sigma Aldrich) was performed according to the

manufacturer's protocol. In brief, PLA probes were incubated with the samples for 1h at

37°C in a humidified chamber, followed by 30 min ligation and 100 min amplification. After

washing, samples were incubated with DAPI and briefly washed in ddH2O. Cover slips were

mounted and imaged as described above.

Cellular stress

Cells were seeded in tissue culture flasks (T-25 or T-175) or on coverslips coated with poly-

D-lysine and grown to confluence in complete DMEM. Cells were then washed 3x with

warm PBS and switched to serum-free conditions in Opti-MEM. For heat stress experiments

shown in Fig. 2A and S10, the conditioned medium was collected before heat stress

(Control/Before sample; 24 h conditioning period) before subjecting the cells to heat stress at

42°C. After a 3 h heat shock, the cells were returned to 37°C for a 4 h recovery period. EV-

conditioned medium was collected (Heat stress sample; 7 h conditioning period) and again

replaced with fresh Opti-MEM. Conditioned medium was then collected after a further 17 h

(After sample; 17 h conditioning period).

For other stress experiments, media was changed before exposure to the stressor and control

samples were maintained in parallel with the treated samples. Media was harvested after a 24

h conditioning period): Heat stress (42°C for 3 h), lipopolysaccharide (LPS) stress (1 µg/ml

LPS, *E. Coli* O111:B4 for 24 h), hypoxia (humidified 2% (v/v) O_2, 5% (v/v) CO_2, balance

N_2) for 24 h though O_2 equilibration occurs slowly in liquid media due to the slow absorption

of O_2 at a static interface water[54], oxidative stress (single application of 50 µM H_2O_2) which

provides a transient oxidative stress due to the rapid degradation of H_2O_2 in cell culture

media[55].

Cell viability assays

OLN-93 cells were seeded on to coverslips coated with poly-D-lysine and grown to 80-90%

confluency. Cells were washed 3x with warm PBS and switched to Opti-MEM. Following

exposure to stress conditions, media was removed, and cells were stained with Hoechst

333421 (5 µg/ml) and propidium iodide (50 µg/ml) in PBS for 20 min at RT. Cells were

washed 1x with PBS and coverslips were mounted and imaged by confocal microscopy

(Nikon Eclipse C1).

Apoptosis was quantified using the Incucyte S3 Live Cell Analysis platform (Sartorius).

OLN-93 cells were seeded (7000 cells/well) in 24-well plates (Corning #3526) coated with

poly-D-lysine. Cells were incubated overnight then washed 3x with PBS and switched to

Opti-MEM. Annexin V-red and Cytox-green reagents (Sartorius) were added according to the

manufacturer's recommendations. The cells were subjected to stress conditions and then

imaged under phase contrast and green/red fluorescence channels every 2 h over a 24 h

period. The number of single and doubled labelled cells were quantified in each channel and

using IncuCyte Base software (Sartorius).

StrataClean pull down of proteins

Cell culture media was cleared by sequential 500, 2000, and 10,000 xg centrifugation steps

followed by 0.2 µm filtration. StrataClean resin (Agilent) was added to the cleared media (5

µl/ml). Thereafter, samples were placed on a roller for 15 h at 4 ºC. The resin was collected

by centrifugation 5,000 xg for 10 min. Total proteins bound to the resin was resuspended in

1X SDS-PAGE loading buffer containing 2 mM DTT, heated to 95 ºC for 10 min and

separated by SDS-PAGE, prior to WB.

Nuclear and cytoplasmic fractionation

OLN-93 cells were pelleted by centrifugation and resuspended in 10 mM HEPES (pH 7.9),

1.5 mM $MgCl_2$, 10 mM KCl, 2 mM DTT with Protease Inhibitor Cocktail (Roche

Diagnostics) and incubated for 10 min on ice. Cells were pelleted at 2000 xg at 4ºC and

resuspended in the same buffer, followed by homogenization with a Dounce homogenizer.

After centrifugation at 10,000 xg for 10 min at 4ºC the cleared supernatant was collected as

the cytoplasmic extract and the pellet was resuspended in 20 mM HEPES (pH 7.9), 1.5 mM

$MgCl_2$, 500 mM NaCl, 25% (v/v) glycerol, 0.5 mM EDTA, 1 mM DTT and protease

inhibitors and incubated for 30 min at 4ºC with continuous mixing. The mixture was then

centrifuged, and the supernatant collected as the nuclear extract.

Histone extraction

301   Acid extraction of histones was performed with the addition of 0.4 M H_2SO_4 for 2 h at RT[56].

302   Following centrifugation, trichloroacetic acid (TCA) was added to the supernatant to a final

303   concentration of 20% (w/v). The mixture was incubated for 2 h at RT before centrifugation at

304   16,000 xg and 4 ºC for 30 min. The pellet was retrieved and washed 3x with acetone before

305   air drying. Histones were dissolved in ddH_2O.

Nanoparticle tracking analysis

Nanoparticle tracking analysis (NTA) was used to determine the size distribution and number

of particles[57] using a NanoSight NS300 instrument (Malvern Panalytical) equipped with NTA

3.1 software. For all our recordings, a camera level of 8-9 and a detection threshold of 3-4

were used. Samples were diluted in ddH_2O or buffer to achieve a particle concentration of

approximately 2×10^8-1×10^9 particles/ml. Five 60 s films were analysed for each sample and

measurements were repeated for 3-5 replicate samples.

NaCl or lithium 3,5-diiodosalicylate treatment of EVs

$NaCl$[58] and Li-diiodosalicylate[59] were used for extraction of proteins from purified EVs (10

µg in 20 mM Tris-HCl, pH 7.5) using NaCl (250 mM-750 mM) or Li-diiodosalicylate (5-50

mM). The final volume was adjusted to 25 µl using Tris-HCl. Samples were incubated for 12

h at 4ºC with gentle shaking. The next day, 5 µl of the sample was taken for TEM analysis

and 20 µl was subjected to ultracentrifugation at 150,000 xg in a TLA-100 rotor and

benchtop ultracentrifuge (Optima Max-XP, Beckman Coulter) for 1h at 4ºC. Supernatant and

pellet (resuspended in 20 µl Tris-HCl buffer) were quantified for DNA (dsDNA assay,

Qubit), or histone H3 by SDS-PAGE and WB. NaCl treated supernatants and pellet samples

were also analysed by dot blot, where 20 µl of supernatant/pellet samples were mixed with 30

µl of Tris-HCl buffer (pH 7.4) and blotted on to a 0.2 µm nitrocellulose membrane (Biorad)

using 96-well vacuum manifold (Biorad). After blotting, dot blots were treated as for WB

membranes for protein detection.

Nuclease or protease treatment of EVs

DNase I, S1 nuclease, and Exonuclease III (Thermo Fisher Scientific) were added to

EV^{OptiPrep}(10 µg; 2 U of DNase I, 100 U of S1 nuclease, and 20 U of Exonuclease III) in

combination. For combined enzyme digestion, S1 nuclease and Exonuclease III buffers were

both used at 1x concentration. Digestion mix was incubated at 37°C for 2 h. After digestion,

total DNA was quantified by Qubit or isolated using CeleCTEV tumour DNA enrichment kit

(Hansa Biomed). The concentration of histone H3 was estimated by WB. DNA extracted

from EVs before and after nuclease digestion was also analysed by TapeStation 4200

(Agilent), using reagents and screed Tape D5000 according to the manufacturer's

instructions.

For trypsin digestion, EV$^{100,000 \text{ xg}}$ from OLN-93 or HEK293-H3 were suspended in 50 mM

HEPES buffer (pH 7.5) and treated for 30 min. at 37°C with Trypsin Protease (MS Grade,

Pierce/Thermo Fisher Scientific) at ratios of 1:1500, 1:500 and 1:150 µg trypsin:µg protein

(determined in pilot experiments to span a suitable range for partial to complete digestion of

histones) in a total volume of 60 µl. Control EVs were incubated without trypsin. Trypsin

was inhibited using 10 µM phenylmethylsulfonyl fluoride (PMSF, Roche Diagnostics) in a

total volume of 1ml PBS for 10 min on ice. EVs were washed in PBS and pelleted at 110,000

g for 1 h prior to WB.

Lipid isolation and liposome preparation

Lipids from intact OLN-93 cells were extracted as described by Bittame *et al.*[60]. In brief, cells

were trypsinized from 5x T-175 culture flasks grown to 90 % confluency and collected by

centrifugation. Pellets were then resuspended in 3 ml of 1:2 chloroform:methanol and

incubated for 30 min with vortexing every 5 min. Chloroform (500 µL) and 900 µL deionized

water were added and samples were centrifuged at 1300 xg then incubated for 10 min at RT.

The organic bottom phase was collected reextracted with 500 µl chloroform and 900 µl

deionized water, after which the sample was centrifuged again at 1300 xg for 10 min at RT.

The organic phase was collected and dried with a gentle stream of nitrogen under constant

rotation. Samples were dried further under vacuum for 3 h. Dried lipids were rehydrated

overnight in 50 mM HEPES (pH 7.4), 150 mM NaCl and sonicated briefly in a sonicating

water bath. Following rehydration, liposomes were homogenized using a 100 nm membrane

mini extruder (Avanti Polar Lipids). The sample was heated to 65°C and then passed through

a 100 nm membrane 30 times. Liposome size distribution and morphology were confirmed

by NTA and TEM, respectively.

Liposome/EV binding assays

Human recombinant histones from New England Biolabs: H1 (M2501S), H2A (M2502S),

H2B (M2505S), H3.1 (M2503S) and H4 (M2504S), total histones extracted from calf thymus

(H9250, Type IIA Sigma-Aldrich), or recombinant His-tagged octamer (BPS-52037) or

nucleosomes (BPS-52038) were purchased from BPS/Nordic Biosite. For binding studies,10

µg of liposomes (in 50 mM HEPES (pH 7.4),150 mM NaCl,) were mixed with 1 µg histones

and incubated for 1 h at RT with gentle mixing. Histones without liposomes was used as

controls. Samples were loaded onto 100 kDa MWCO centrifugal filters with PES membrane

(Amicon Ultra, Millipore) and washed 7x with buffer by centrifugation. Samples retained by

the filter were analysed by WB. Liposome interaction with histones, nucleosomes, or

octamers was differentiated on sucrose gradients. Histones (5 µg) were mixed with 10 µg

liposomes. Nucleosomes and octamers (5 µg) were mixed with 20 µg liposomes. Total

reaction was performed as a 100 µl volume in a 50 mM HEPES (pH 7.4) containing150 mM

NaCl. Liposome-histone/octamer/nucleosome samples were loaded at the bottom of a 40%

sucrose layer, with a discontinuous 6-step gradient up to 0% sucrose. Samples were

centrifuged at 100,000 xg for 16 h/4°C using a TLS-55 swinging-bucket rotor and benchtop

ultracentrifuge (Optima Max-XP, Beckman Coulter). Fractions were collected and loaded

directly on SDS-PAGE prior to WB.

For dot blotting, EV^{OptiPrep} were diluted to 50 µg/ml in PBS. EVs were diluted in a 2-fold

series in a 96-well microtiter plate in triplicate. Solutions were then blotted on to a

nitrocellulose membrane using a 96-well vacuum manifold. After blotting, membranes were

blocked in PBS containing 2.5% (w/v) BSA, followed by overnight incubation of blots in 2

µg/ml octamer/nucleosome in the same buffer. Nucleosome/octamer binding were detected

by anti-His tag and HRP conjugated anti-rabbit secondary antibodies.

Genetic or pharmacological intervention in cell lines

All silencer RNA (siRNA) including Tsg101 (196580) Alix (s221953), Smpd2 (189939) and

negative control siRNA; 4390771 were purchased from Thermo Fisher Scientific. OLN-93

cells were grown to 90% confluency on poly-D-lysine coated 25 cm^2 flask. Cells were

washed with PBS and switched to OptiMEM. A total of 300 pmol siRNA was mixed with 6

µL Lipofectamine 3000 in Opti MEM media and incubated at room temperature for 15 min

prior to addition to cells. The siRNA-lipofectamine mix was incubated with cells at 37ºC for

6 h and then replaced with fresh OptiMEM and further incubated for 24 h before

harvest/analysis. Conditioned media was processed by StrataClean pull as described above.

For pharmacological intervention, drug concentrations were selected from previously

published protocols (Supplementary table S1). Cell viability was evaluated in OLN-93 cells

grown in 96-well plates treated with in poly-D-lysine using the MTT assay following

treatment with a range of drug concentrations, applied in OptiMEM to a set of 6 replicate

wells. After 24 h of incubation, 10 µl freshly prepared MTT (5 mg/ml in ddH$_2$O) was added

to each well. Cells were incubated for 3 h at 37ºC. Media was aspirated and 50 µl DMSO was

added to each well to dissolve formazan crystals with shaking at 200 r.p.m. Absorbance at

560 nm was read in a plate reader (Biotech Synergy 2). Wells without cells were used as

controls.

Analysis of histone secretion was analysed in OLN-93 cells grown in poly-D-lysine coated T-

25 flasks (80-90% confluency). Cells were washed with PBS and supplemented with 2.5 ml

OptiMEM per well containing compound or vehicle (control): 10 mM 3-methyl adenine (3-

MA, Sigma), 5 µM neutral sphingomyelinase inhibitor GW4869 (SelleckChem), 200 mM

trehalose (Merck), 50 µM imipramine (SelleckChem), 25 µM rapamycin (SelleckChem), 100

nM Bafilomycin A1 (SelleckChem), 15 µM Dyngo (SelleckChem), 10 mM valproic acid

(Sigma-Aldrich), 10 mM sodium butyrate (Sigma-Aldrich), 200 nM Trichostatin A (Sigma

Sigma-Aldrich), 20 nM romidepsin (SelleckChem), 50 µM YF-2 (SelleckChem), 15 µM

C646 (SelleckChem), 10 µM A-485 (SelleckChem). After 24 h treatment, supernatant

(conditioned media) was removed from the flasks. Conditioned media (2 mL) was taken and

processed by centrifugation at 500 xg, 2,000 xg and10,000 xg followed by protein pull down

using StrataClean resin.

High resolution single EV imaging flow cytometry

EVs were analysed at a single vesicle level by high resolution imaging flow cytometry

(IFCM; Amnis Cellstream, Cytek; equipped with 405, 488, 561 and 642 nm lasers) based on

previously optimized settings and protocols[61-63]. Briefly, fluorescently conjugated anti-

histone and anti-tetraspanin antibodies were used to probe for respective proteins on the EV

surface in pre-cleared conditioned media (CM) samples collected from HeLa cells for control

and H$_2$O$_2$ treatment. For each measurement, 25 µl of CM (and unused media as a control)

were incubated overnight at room temperature with the following antibodies: anti-CD9-APC

(Miltenyi Biotech, clone SN4, 130-128-037; 0.6 ng/µl), anti-CD63-APC (Miltenyi Biotec,

clone H5C6, 130-100-182; 0.6 ng/µL), anti-CD81-APC (Beckman Coulter, clone JS64,

A87789; 0.6 ng/µl), anti-H3-AF488 (Invitrogen, clone 17H2L9, MA7-02023-A488; 2

ng/µL), and anti-H4-FITC (Novus Biologicals; polyclonal, NB21-2044F; 2 ng/µL). For

detergent lysis controls, samples were treated with 0.5% (v/v) NP-40 for 30 min at RT post

staining. All samples were diluted 50-500-fold in PBS-HAT[64] before acquisition via the

autosampler of the Cellstream instrument with FSC turned off, SSC laser set to 40%, and all

other lasers set to 100% of the maximum power setting. Samples and controls were acquired

for 3-5 minutes at a flow rate of 3.66 µl/minute with CellStream software version 1.2.3 and

analysed with FlowJo Software version 10.8.1 (FlowJo LLC). Fluorescence calibration and

reporting of fluorescence data in molecules of equivalent soluble fluorophores (MESF) was

performed as described before[61, 62]. Dulbecco's PBS pH 7.4 (Gibco) was used as instrument

sheath fluid.

Histone analysis by mass spectrometry

Histones were prepared as described previously by Shechter *et al.*[56]. Briefly, histones were

acid-extracted with chilled 0.2 M sulfuric acid (5:1, sulfuric acid:pellet) and incubated with

constant rotation for 4 h at 4ºC, followed by precipitation with 33% TCA overnight at 4 ºC.

The pellet was dissolved in 50 mM ammonium bicarbonate (pH 8.0), and histones were

subjected to derivatization using 5 µl of propionic anhydride and 14 µl of ammonium

hydroxide to balance the pH at 8.0. The mixture was incubated for 15 min and the procedure

was repeated. Histones were then digested with 1 µg of sequencing grade trypsin (Promega)

diluted in 50 mM ammonium bicarbonate (1:20, enzyme:sample) overnight at room

temperature. The derivatization reaction was repeated to derivatize peptide N-termini. The

samples were dried in a vacuum centrifuge.

Samples were resuspended in 10 µl of 0.1% trifluoroacetic acid and loaded onto a Dionex

RSLC Ultimate 300 (Thermo Fisher Scientific), coupled online with an Orbitrap Fusion

Lumos (Thermo Fisher Scientific). Chromatographic separation was performed with a two-

column system, consisting of a C-18 trap cartridge (300 µm ID, 5 mm length) and a picofrit

analytical column (75 µm ID, 25 cm length) packed in-house with reversed-phase Repro-Sil

Pur C18-AQ 3 µm resin. Peptides were separated using a 30 min gradient from 1-30% buffer

B (buffer A: 0.1% formic acid, buffer B: 80% acetonitrile + 0.1% formic acid) at a flow rate

of 300 nl/min. The mass spectrometer was set to acquire spectra in a data-independent

acquisition (DIA) mode. Briefly, the full MS scan was set to 300-1100 m/z in the orbitrap

with a resolution of 120,000 (at 200 m/z) and an AGC target of $5x10^5$. MS/MS was

performed in the orbitrap with sequential isolation windows of 50 m/z with an AGC target of

$2x10^5$ and an HCD collision energy of 30.

Histone peptides raw files were imported into EpiProfile 2.0 software[65]. From the extracted

ion chromatogram, the area under the curve was obtained and used to estimate the abundance

of each peptide. To achieve the relative abundance of post-translational modifications

(PTMs), the sum of all different modified forms of a histone peptide was considered as 100%

and the area of the specific peptide was divided by the total area for that histone peptide in all

of its modified forms. The relative ratio of two isobaric forms was estimated by averaging the

ratio for each fragment ion with different mass between the two species.

Negative contrast staining and immuno-transmission electron microscopy

EVs were visualized by negative staining as described previously by Hedlund *et al.*[66]. A total

of 4 µL of sample was adsorbed to glow discharged, formvar/carbon coated copper grids.

Following two washes in ddH$_2$O, negative contrast staining was performed in 1.5% (w/v)

uranyl acetate for 30 s and dried at RT. Immuno-TEM on EVs was performed as previously

described[67] with slight modification. Samples were absorbed on to copper grids and blocked

by 0.1% (w/v) gelatin and 1% (w/v) BSA in PBS for 1 h at RT. After brief washing in PBS,

grids were incubated in primary antibodies (see the Antibodies section, above) diluted in PBS

containing 1% BSA and incubated overnight at 4ºC. Grids were washed in PBS and

incubated with 10 nm protein A-gold particles (PAG10, BBI Solutions) for 1 h at RT. After

washing using PBS, grids were washed 5 times in water and exposed to 1.5% uranyl acetate

for 30 s and dried at RT before imaging.

Immuno-TEM on cells was performed according to published methods[68]. OLN-93 cell

sections were processed according to the Tokuyashu method[69]. Cells were fixed in freshly

prepared EM fix solution containing 2% (w/v) paraformaldehyde and 0.2% (w/v)

glutaraldehyde. Optimal fixation was facilitated using a microwave incubator (PELCO

BioWave). Cells were then centrifuged for 5 min at 500 xg and the fixative solution was

replaced with PBS containing 10 mM glycine to quench free aldehyde groups. Thereafter, 3x

PBS washes were performed and cells were resuspended in 12% (w/v) gelatin and incubated

for 1 h at 37°C. Gelatin blocks were placed into 2.3 M sucrose overnight at 4°C on a rocking

platform. Blocks were mounted on the top of metal pins and snap frozen in liquid N_2. Cryo-

sectioning (70 nm sections) was performed in a Leica UC7 cryostat (Leica Microsystems),

and sections were mounted on copper 300x300 grids. Grids bearing several sections were

placed in PBS and incubated at 37°C for 1 h followed by incubation in 10 mM glycine in

PBS. After 3x washes in PBS the sections were blocked in 1% (w/v) BSA, 0.1% (w/v)

gelatin. Primary antibodies for histone H1 (most sensitive for this method) and CD63 were

added and incubated overnight at 4°C. After 5x washes in PBS, secondary antibody-gold or

protein-A gold conjugates were added. Finally, sections were washed extensively in PBS and

contrast enhanced with uranyl acetate/methylcellulose (1:4) solution. Control sections

processed without primary antibodies were prepared in parallel to estimate the specificity and

background of antibody binding. Specimens were examined with a JEM1230 (Jeol) or Talos

electron microscope (Thermo Fisher Scientific) at 80 kV and micrographs were collected

with a MSC 600CW CCD camera (Gatan) using DigitalMicrograph software (Gatan).

Statistics

Statistical analyses were performed using Graph Pad Prism 5.0c (GraphPad Software) or

SPSS 26 (IBM). Error bars represent the standard error of mean. Data was assessed for

normality using Shapiro-Wilk test. Results were analysed for statistical significance by T-

test, one-way or two-way analysis of variance (ANOVA) followed by the Bonferroni and

Sidak`s post hoc test. Treatment effects were assessed using paired sample T-test.

RESULTS

Histone association with extracellular and intraluminal vesicles

First, we characterized histone secretion into the conditioned media from the rat

oligodendrocyte progenitor cell line OLN-93. This line could be grown in 2D culture under

serum-free conditions for several days, facilitating purification of EVs by sequential

centrifugation [47]. In addition, it was possible to produced large numbers of EVs in 3D culture

in a bioreactor, enabling us to generate sufficient material for large scale biochemical

characterization.

Comparison of histones in the whole cell lysate (WCL) and EV pellet fraction ($EV^{100,000 \text{ xg}}$)

by western blotting (WB) showed significant enrichment of membrane localized tetraspanins

CD9 and CD63 and luminal markers tumor susceptibility gene 101 (Tsg101) and actin beta

(Actb) in EVs (Fig. 1A). Lack of contamination of with cellular organelles was confirmed by

absence of the cis-Golgi matrix protein GM130. As expected, robust signals were observed

for all nuclear core histones (H2A, H2B, H3 and H4) in the WCL, but also in EVs. We then

used immunogold staining together with transmission electron microscopy (immuno-TEM) to

confirm that all histones (core and linker) localized to EVs. Surface labelling in

unpermeabilized EVs suggested that like CD63 (Fig. 1B) and CD9 (Fig. S1A), histones also

localized to the EV surface (Fig. 1B and S1A)

To see whether histones colocalized with CD63 in intact cells, we first used

immunofluorescence staining in combination with confocal microscopy. Colocalization of

non-nuclear staining for all four core histones and CD63 was observed in cytoplasmic or

membrane puncta (Fig. 1C, Fig. S1B-D). Since the degree of colocalization observed was

greatest with H3/CD63 (Fig. 1C), we also tested this antibody combination in a proximity

ligation assay (Duolink PLA), which has a higher spatial resolution (within 40 nm) than

standard confocal fluorescence microscopy. PLA confirmed an association of anti-H3/CD63

antibodies in cytoplasmic puncta with the morphological appearance of MVBs (Fig. 1D). No

background signal was observed in the absence of primary antibodies (Fig. S1E) confirming

the specificity of the assay.

OptiPrep discontinuous density gradient centrifugation is a well-established method for the

purification of small EVs (sEVs) including smaller MVs and exosomes, from other

components such as larger microvesicles and apoptotic bodies[70, 71]. In EVs from mast cell and

erythroleukemic cell lines, histones localize to high denisty (HD; 1.14-1.19 g/ml) but not to

low density (LD; 1.10-1.13 g/ml) sEV fractions[30]. To investigate histone distribution in EVs

from OLN-93 cells, we fractionated $EV^{100,000\ xg}$ on 12-36% OptiPrep gradients. Fraction

densities were calculated based on a reference gradient (Fig. S2A). TEM was used to confirm

the purity and morphology of EVs in each fraction (Fig. S2B). The majority of EVs with an

intact morphology and ranging from 30-150 nm localized to fractions 3-7 (Fig. S2B). WB

confirmed enrichment of CD63 in fractions 3-8 (1.05-1.13 g/ml) and most abundantly in

fractions 6-7 (1.09-1.11 g/ml, Fig. 1E). CD9, Tsg101, and ALG-2-interacting protein X

(ALIX/PDCD6IP) showed shifted distributions in comparison to CD63, being enriched in

fractions 3-6 (Fig. 1E). Annexin A1, a marker of plasma membrane-derived microvesicles,

was most abundant in fraction 4 (1.06 g/ml). Flotillin-1 (membrane marker) spanned fractions

2-7, whereas extracellular acetylcholinesterase (AChE) and the NG2 chondroitin sulfate

proteoglycan, which is a membrane-anchored marker of oligodendrocyte progenitors,

localized to fractions 4-7. The Golgi marker GM130 was seen at low levels through fractions

3-9, but Golgin-97 was not detected (Fig.1E). Thus, we were able to resolve distinct but

overlapping EV populations ranging from Annexin A1 positive microvesicles (fraction 4),

through Tsg101/Alix positive sEVs in fractions 4-5 and CD9/CD63 tetraspanin positive sEVs

in fractions 5-7. Interestingly, in OLN-93 cells all four core histones and H1 localized to

fractions 4-7 and were most abundant in fraction 6, identifying them as components of sEVs

that co-purify with CD63, CD9, Tsg101, and Alix (Fig.1E).

To investigate the distribution of histones in EVs from other cell lines, we fractionated

$EV^{100,000 \ xg}$ pellets from a selection of mouse and human cells with different tissue origins

using OptiPrep gradients. H3 positive LD and HD fractions were identified for all cell lines

tested (HeLa, Caco-2, NSC-34, AC-16 and A549; Fig. S3) though with overlapping

distributions. H3 was found in fraction 8 (1.13 g/ml) in all lines tested and TEM analysis

confirmed the presence of EVs in histone positive fractions. Interestingly, HeLa EVs resolved

as two distinct H3 containing populations (Fig. S3; fraction 3-4 copurifying with Alix and

Tsg101 and fractions 6-8). Hence, H3 seems to localize to EVs that float over 3 different

density ranges – LD fractions 3-4 (mainly Tsg101+, Alix+; from HeLa and to a lesser extent

from Caco-2), HD fractions 5-8 (tetraspanin positive; all cell lines apart from A549) and

fractions 9+ (OLN-93, NSC-34, AC-16, A549).

$EV^{100,000 \ xg}$ and $EV^{OptiPrep}$ fractions may contain a mix of plasma membrane derived

microvesicles (indicated by the aforementioned presence of Annexin A1 in fractions 4-8; Fig.

1E) and exosomes that form as intraluminal vesicles (ILVs) in MVBs. To see if histones

could be identified in association with ILVs, we developed a protocol for immuno-TEM on

ultrathin sections of OLN-93 cells. All MVBs examined contained CD63 positive ILVs

(49%, ranging from 3-50 per MVB, Fig. 1F). The densest histone labelling was observed for

the histone H1 antibody where positive ILVs were identified in 32.9% of cases (Fig. 1G). We

also confirmed labelling for core histones (24-41% of ILVs; Fig. S4A-D).

We used immuno-TEM with secondary antibodies conjugated to differently sized gold

particles to verify the colocalization of histone and CD63 in MVBs. OLN-93 sections were

probed with anti-H1 (10 nm gold) and anti-CD63 (5 nm gold; Fig. 1H) in combination.

Quantification showed 36% of ILVs labelled positively for CD63 and 23% for H1.

Interestingly, 17% of ILVs co-labelled with both anti-CD63 and anti-H1 (Fig. 1I). We also

confirmed the colocalization of CD63 and H1 on EV^{OptiPrep} fractions 4-7 (Fig. 1J). The

presence of histone in ILVs and EV^{OptiPrep} from human HeLa (Fig. S5A-B) and Caco-2 cell

lines (Fig. S6A-B) was also confirmed. In summary, using a range of methodological

approaches, we have shown that histones are associated with ILVs inside MVBs and on

secreted EVs in several different cell lines. A proportion of these ILVs/EVs are positive for

both histones and CD63.

Histone secretion in response to cell stress

Cellular stress is a common denominator in pathological conditions where elevated

circulating histones have been observed[72] and we hypothesized that stress might enhance

histone secretion via EVs. Therefore, we developed a heat stress paradigm whereby we

harvested/changed the conditioned media on OLN-93 cells as a control condition ('Before')

and then cultured the cells at 42ºC for 3 h, followed by a 4 h recovery period at 37ºC ('Heat

stress'). Media was collected, changed, and harvested at a third time point after 17 h ('After';

Fig. 2A).

To quantify secreted histones, we first pulled down all proteins from 1 ml of conditioned

media using StrataClean resin and analysed by WB. We observed strong induction of the heat

shock protein Hsp70 in the After sample, confirming the efficacy of the treatment (Fig. 2B).

EV markers Tsg101, CD9 and Actb showed a slight reduction in the Heat stress sample,

likely due to the shorter period of media conditioning (7h versus 24 h for Before). However,

levels of H2B and H3 were equivalent in Before and Heat stress samples and increased in the

After sample (Fig. 2B), suggesting elevated secretion of histones in response to heat stress.

These findings were confirmed in EV$^{100,000 \ xg}$ fractions where a constant amount of protein

was analysed by WB to account for the different periods of media conditioning. Here, we

observed strong induction of Hsp70 in both Heat stress and After samples (Fig. 2C). The

levels of Tsg101, CD9, and Actb remained constant. However, heat stress resulted in

increased secretion of linker and core histones. Hence, EV associated histones accumulate in

the conditioned media and are increased in EVs in response to heat stress.

To see whether histone secretion is part of a common stress response, we exposed OLN-93

cells to several stress paradigms. EVs were harvested after 24 hours, though the period of

exposure to the stressor was different in each case (see Methods): Heat (42°C for 3 h), the

endotoxin (LPS; 24 h), low oxygen tension (hypoxia; 24 h), and oxidative stress (H_2O_2; 24 h).

Since histones released from dead/dying cells could complicate analyses, we first determined

the effect of the stresses on cell viability using live/dead (PI/Hoechst 33342) staining, and on

apoptosis using Annexin V/Cytox staining. A significant increase in cell death, indicated by

nuclear PI labelling, was observed after heat stress (Fig. S7A, B) but not after the other stress

paradigms. Similarly, a significant increase in Annexin V positive (early apoptotic), Cytox

positive (compromised plasma membrane, as for PI) and Annexin/Cytox double positive (late

apoptotic/dead) cells were observed following heat stress, but not for other stress conditions

(Fig, S7C, D). Interestingly, LPS treatment led to reduced Annexin V, Cytox and double

stained cells in comparisons to controls, indicating a potential protective effect of LPS

exposure in these cultures (Fig. S7D).

We then proceeded to quantify the relative levels of histones in cytoplasmic versus $EV^{100,000}$

^{xg} fractions under control and stress conditions. All stress interventions led to a decrease in

the cytoplasmic pool of H2B and H3 of between 10-64% (Fig. 2D, E). In contrast, large

increases in H2B and H3 were found in the $EV^{100,000\ xg}$ fractions under all stress conditions

ranging from 15-95-fold (Fig. 2D, E). To validate the results of WB in intact cells, we used

immunostaining followed by confocal microscopy. Enhanced colocalization of anti-H3 and

anti-CD63 staining was observed in cytoplasmic MVB-like compartments under all stress

conditions (Fig. S8A). In contrast, colocalization with the plasma membrane/microvesicle

marker Annexin A1 appeared unchanged (Fig. S8B). Duolink staining with anti-H3 and anti-

CD63 showed enhanced colocalization of these 2 markers under all stress conditions (Fig.

S8C). Further, immuno-TEM was used to quantify colocalization of histone and CD63 more

precisely. Increased numbers of anti-H1 labelled (10 nm gold) and anti-CD63 labelled (5 nm

gold) particles were found in the MVBs of cells exposed to all stress conditions (Fig. 2F,

S9A-E). The number of H1 or H1/CD63 double positive ILVs increased significantly under

all stress conditions apart from oxidative stress (Fig. 2G-H), which may reflect the short half-

life of the stressor (see Methods) and/or the longer recovery period available to cells

subjected to this stress paradigm.

To investigate if the size and quantity of EVs changed in response to stress, first we analysed

$EV^{100,000\ xg}$ from Before, Heat stress and After by NTA (Fig. S10A). There was an

approximately 3-fold increase in EV particle concentration in Heat stress and After samples

after normalization for the conditioning period (Before = 2.6×10^7, Heat stress = 7.1×10^7,

After = 8.6×10^7 particle/ml/h; Fig. S10B). We also observed an increase in 40-100 nm sized

particles (Fig. S10A). However, since the sensitivity of NTA declines for particles of less

than 50 nm[30], we used TEM to estimate the size of $EV^{OptiPrep}$. All stress conditions led to the

secretion of smaller diameter EVs (mean = 36-52 ±18-37 nm, mode = 30 nm; Fig. 2I-J) in

comparison to untreated samples (mean = 87 ±35 nm, mode = 100 nm; Fig. 2J). In summary,

our results show that cellular stress responses lead to an increase in the quantity and a

reduction in the size of EVs.

Co-dependence of histones and DNA for EV binding

Previous studies have focussed on the DNA/chromatin content of sEVs and concluded that

sEV histones are tightly associated with DNA as nucleosomes. However, the situation is

unclear since both double- and single-stranded (ds- or ss-, respectively) DNA have been

found inside and outside of EVs[40, 73]. As only dsDNA associates with histones in

nucleosomes, we sought to clarify both the possible co-dependence of histones and DNA for

EV binding, and their location in relation to EV membrane and lumen.

First, we quantified the amount of DNA (Qubit) and H3 (WB) in $EV^{OptiPrep}$ fractions from

OLN-93 cells cultured under control or stress conditions. H3 showed a dynamic distribution,

depending on the type of stress and period of exposure to the stressor (Fig. 3A). H3 increased

in fractions 6-7 under all stress conditions but DNA did not, with the majority of DNA/µg

total protein localizing to fractions 8-12. Notably, the distributions of DNA did not mirror the

distributions of H3 suggesting a limited co-dependency of histones and DNA for EV binding.

Next, we devised a protocol combining chemical extraction, ultracentrifugation, and

sonication to distinguish between histones or DNA loosely associated with the outer surface

of EVs from those tightly bound to, or integrated into, the membrane, or in the lumen (Fig.

3B). A high concentration of NaCl is commonly used to disrupt electrostatic interactions

whilst still being a non-denaturing salt[58]. The chaotropic salt lithium 3,5-diiodosalicylate can

solubilize membrane associated proteins at lower concentrations, while disrupting the

membrane completely at higher concentrations[59]. In addition, we used the detergent TX-100

(0.1%) to lyse EVs[74], or sonication to disrupt them. TEM was used to estimate EV

morphology and integrity. Following EV re-isolation via ultracentrifugation, we quantified

H3 and DNA in the supernatant or pellet fractions using WB or Qubit/Tape Station,

respectively.

EV^{OptiPrep} could be treated with up to 750 mM NaCl without adverse effect on morphology

(Fig. 3C, S11A). However, 1M NaCl led to visible damage to EV membranes (Fig. S11A).

Increasing concentrations of NaCl (250-750 mM) led to a displacement of approximately

40% of H3 into the supernatant, which became maximal at 500 mM NaCl (Fig. 3D-E). Up to

30% of EV associated DNA was displaced but did not reach a plateau, even at 750 mM NaCl

(Fig. 3E).

A significant proportion of H3 remained associated with the EV pellets following treatment

with 750 mM NaCl (Fig. 3F). To probe this in more detail, EV pellets recovered after 750

mM NaCl treatment, were sonicated to disrupt membranes, and expose luminal contents. We

observed that a minor proportion of H3 was released by this procedure, but the majority

remained associated with the pellet and could not be extracted with an additional 750 mM

NaCl treatment of sonicated EVs (Fig. 3G). Interestingly, following different stress

conditions a variable proportion of surface associated H3 (extracted by 750 mM NaCl) could

be removed. Heat stress EVs showed the least surface associated H3, while LPS and hypoxia

EVs showed the highest levels (Fig. S11B).

We used lithium 3,5-diiodosalicylate extraction to estimate the proportion of H3 that was

tightly associated or integral to the membrane. TEM analysis showed that EV membranes

remained largely intact in 25 mM lithium 3,5-diiodosalicylate (Fig. 3H) but were disrupted

by 50 mM (Fig. S11C). H3 was extracted progressively into the supernatant by 10-25 mM

and completely by 50 mM lithium 3,5-diiodosalicylate. TX-100 treatment led to extraction of

approximately 50% H3 into the supernatant (Fig. 3I). Taken together, our results suggests

that a significant proportion (50-60%) of EV histones are tightly associated with the EV

membrane. Some histones associate more loosely with the surface of EVs, likely via

electrostatic interaction, and a minor fraction may be luminal.

We treated EV^{OptiPrep} with a combination of nucleases (DNase I, S1 nuclease and

Exonuclease III) to remove DNA from EVs. In addition we sonicated EVs to expose luminal

contents. Digestion had no appreciable effect on H3 association in both intact and sonicated

EVs (Fig. 3J). However, based on Qubit estimation in the same samples, the amount of DNA

was significantly reduced and sonication had no additional effect (Fig. 3K). To determine

DNA size and amount more precisely, we purified EV DNA and analysed this by

TapeStation and Qubit in parallel. Nuclease digestion led to the degradation of EV DNA to

$\leq$100 bp (Fig. 3L), with $\approx$ 40% remaining (Fig. 3M), which was confirmed by Qubit (Fig.

3N).

We used a range of trypsin concentrations to probe exposure on the surface of EVs and

analysed histones and other proteins in the pellet by WB (Fig. 3O). Under conditions where

the membrane associated proteins Tsg101 and CD9 were protected from digestion, a

complete loss of the glycophosphatidylinositol (GPI)-linked outer membrane NG2 occurred

even at the lowest trypsin:protein ratio (1:1500). All histones were partially protected from

trypsin cleavage at 1:1500-1:500, although they were completely removed at 1:150.

Recombinant purified histones were used as control for trypsin digestion and, in contrast to

EV associated histones, did not show a similar pattern of partial cleavage products (Fig. S12).

Since all anti-histone antibodies tested were directed against C-terminal epitopes (see

Methods) we also examined whether N-terminal of C-terminal regions were equally

susceptible to cleavage in intact EVs. A HEK293 cell line stably expressing H3.1 with an N-

terminal HA- and a C-terminal 6xHis-tag (Fig. 3P) were generated. EVs were purified and

treated with trypsin as above. With an increasing trypsin:protein ratio, reactivity towards the

N-terminal HA-tag was lost without the appearance of smaller proteolytic fragments whereas

the C-terminal 6xHis antibody recognized one or more truncated species (Fig. 3Q). In

summary, our results show that histones are mainly EV membrane associated proteins.

Furthermore, although DNA is also present at the surface of EVs, the majority of histone

does not appear to be dependent on DNA for its membrane association.

Binding of histone octamers and nucleosomes to membrane lipids and EVs

To probe the potential interaction of histones with lipids, we prepared liposomes from OLN-

93 whole-cell lipids. After incubation with recombinant human histones or purified total

histones from calf thymus, liposome-histone mixtures were washed using 100,000 kDa

molecular weight cut-off ultrafiltration and analysed by WB. All histones bound to liposomes

and were retained by the filter even after thorough washing (Fig. 4A). Liposome-histone

mixtures were also resolved on 0-40% sucrose gradients[75]. Total histones from calf thymus,

or individual recombinant histones floated to lower density fractions and were found in the

least dense fractions 1-2 (Fig. 4B). These experiments confirm that recombinant histones

expressed in *E.coli*, which lack PTMs, bind directly to membrane lipids.

Next, to test if DNA enhances the binding of histones to liposomes, we used sucrose density

gradient floatation to compare the relative binding of recombinant histone octamers and

mono nucleosomes. His-tagged octamers/nucleosomes alone remained in fraction 6 (Fig. 4C,

D), whilst most of the added histone octamer bound to liposomes and floated in fractions 1-2

(Fig. 4C). In contrast, nucleosomes bound poorly to liposomes (Fig. 4D). EVs did not float

reliably on sucrose gradients when treated with nucleosomes (not shown), so we investigated

EV binding using a solid phase (dot blot) assay. First, we immobilized different amounts of

EVs (0.39-50 µg/ml) on a nitrocellulose membrane, then incubated with His-tagged

octamers/nucleosomes, followed by detection of binding with an anti-His tag antibody.

Results showed that octamers bound significantly higher to EVs in comparison to

nucleosomes, indicating the presence of DNA in the nucleosome interferes with

histone/octamer interaction to EV membranes. (Fig. 4E, F).

Origin of extracellular vesicle associated histones

Histones could localize to EVs via several different pathways: i) Sorting of soluble

cytoplasmic histones into MVBs via ESCRT-dependent or -independent pathways, thereby

partitioning to the luminal EV compartment. ii) Acquisition of histones to the outer surface of

EVs via the amphisome pathway, following fusion of autophagosomes to endosomes. iii)

Endocytosis of EV or non-EV derived histones and their subsequent sorting into ILVs, where

they would also localize to the outer surface of EVs. To investigate these 3 possibilities, we

used both siRNA-mediated knockdown and chemical modulators (drugs) targeting relevant

pathways. Drug concentrations were selected according to published observations (see Table

S1), which we then validated to make sure that the selected concentrations did not impact

significantly on cell morphology and viability (Fig. S13).

Knockdown of the ESCRT components *Alix* or *Tsg101* led to a reduction in H3 secretion

(Fig. 5A). Annexin A1 was expressed at a low level in OLN-93 cells (Fig. S8B) and EVs

(Fig. 1E) and did not change dramatically following siRNA treatments (Fig. 5A), indicating

that the microvesicle pathway was unable to compensate for H3 secretion via exosomes. On

the other hand, knockdown of the ceramide pathway using an siRNA directed against *neutral*

sphingomyelinase (*Smpd2/nSMNase2*) showed only a slight decrease in H3 secretion, which

was not significant. Due to the lack of specific antibody recognizing rat Smpd2 in WB, we

could not confirm the efficiency of Smpd2 knockdown. Hence, we used the nSMNase2

inhibitor GW4869 to block ceramide production[76, 77]. Treatment of OLN-93 cells with 5 μM

GW4869 completely blocked H3 secretion (Fig. 5B).

Based on the proteomic analysis of EVs and the non-vesicular secretome, it has been

proposed that EV-associated histones may originate from the nucleus and be secreted via an

autophagy-dependent amphisomal pathway[33]. To investigate the role of autophagy in histone

secretion, we treated OLN-93 cells with the autophagy inhibitor 3-methyadenine (3-MA),

which blocks phagophore formation. Treatment with 3-MA increased H3 secretion

significantly (Fig. 5B) and a significant increase in histone localization to ILVs (Fig. 5C). On

the other hand, inducers of mTOR-independent autophagy (trehalose and imipramine)

showed a significant decrease an opposite effect (Fig. 5B). The mTOR-dependent inhibitor

rapamycin showed a decrease in H3 secretion that was not significant. Further, we used

Bafilomycin A1 (Baf) to block the vacuolar H+-ATPase (V-ATPase) and thereby the fusion

of autophagosomes and late endosomes to lysosomes[78]. Like 3-MA, Baf treatment also

increased H3 secretion (Fig. 5B) and histone localization to ILVs (Fig. 5C). To investigate

the requirement for clathrin-dependent endocytosis, we used the dynamin inhibitor Hydroxy

Dynasore (Dyngo-4a)[79], which downregulated H3 secretion (Fig. 5B).

Histone translocation to, and function in, the nucleus is modulated by post-translational

modifications (PTMs), such as acetylation and methylation[80, 81]. Histone acetyl transferases

(HATs) modify histones on Lys (L) and Arg (K) residues, thereby reducing the net positive

charge of the protein. Histone acetylation is counteracted by the activity of deacetylases

(HDACs). To determine if acetylation status is important for histone secretion, we used drugs

to either inhibit or activate HDACs or HATs. The pan HDAC inhibitor Trichostatin A as well

as Romidepsin (HDAC1 and 2 selective) did not have a significant effect but both valproic

acid and sodium butyrate (HDAC1 selective) reduced H3 secretion (Fig. 5B). The p300/CBP

selective HAT inhibitors C646 and A-485 reduced, whereas the broad-spectrum HAT

activator YF-2 increased, H3 secretion. Hence, blocking ESCRT, ceramide and dynamin

pathways inhibited H3 secretion. In contrast, inhibition of autophagosome formation and

lysosomal degradation promoted accumulation of H3 in ILV/MVBs and secretion via EVs.

Interfering with histone acetylation status modulated H3 secretion though the relationship to

acetylation status was unclear.

Surface association of extracellular vesicle histones and their modifications

During the multiple centrifugation and washing steps used in purification, a significant

amount of EV surface proteins may be lost from the corona[82]. Alternatively, non-nuclear

histones might fuse with EV membranes under high centrifugal force[83, 84]. Therefore, to

confirm that histones are EV surface associated proteins, we used imaging flow cytometry

(IFCM) to characterize histone positive EVs in conditioned media without purification[61, 62, 85].

We identified fluorescently labelled anti-human H3 and H4 antibodies with good signal and

low noise (Fig. S14A) and combined these with previously characterized anti-human

tetraspanin antibodies (CD9, CD63 and CD81). We analysed EVs in the conditioned media

of HeLa cell cultures under either control conditions, or following acute exposure to H_2O_2

(Fig. 6A, S14).

Time course measurements over a 24 h period revealed that the mean fluorescence intensity

(MFI) of anti-H3 and -H4 labelling was already maximal by 2 h and remained constant out to

24 h (Fig. S14B). In contrast the MFI of anti-tetraspanin labelling increased from 2-24 h,

suggesting that H3/H4 localize to EVs but that the total amount of antibody accessible

histone remains constant over time. Interestingly, H3 and H4 showed different kinetics.

Under control conditions, H3 positive EVs showed a gradual increase, with the highest

concentration at 24 h. In H_2O_2 treated cultures, H3 positive EVs accumulated faster, with

highest concentrations measured at 4 h, followed by a gradual decrease (Fig. 6B). Thus, H_2O_2

treatment appeared to promote the release of H3 positive EVs. The different kinetics of stress

responsive compared to control EVs seen using IFCM may be due to a lower level of

antibody accessible H3 on the EV surface. In contrast, H4 positive EVs were detected in

control and H_2O_2 treated cells at similar levels up to the 8 h timepoint, and their concentration

increased markedly at 24 h in the H_2O_2 treated cultures (Fig. 6B). This indicates a difference

in the kinetics and/or accessibility of antibodies to H3 or H4 on EVs.

To further characterize H3 and H4 positive EVs, we performed double labelling to assess co-

occurrence of H3 or H4 together with CD9, CD63, or CD81, on the same EV. About 50-60%

of total fluorescence events were found to be double positive for both H3 and CD9 or CD63,

while only about one third were double positive for H3 and CD81 (Fig.6C and Fig. S14C).

Similarly, H4 was detected more on CD63 positive (~60%) and CD9 positive (~45%) EVs

than on CD81 positive EVs (~35%; Fig. 6D and S14C).

To address whether histone PTMs might influence their association with EVs, we

characterized the extent and pattern of PTMs on EV associated histones. We extracted

histones from OLN-93 EV^{OptiPrep} and the corresponding nuclei from control cells and those

exposed to either LPS or hypoxia (Fig. S15) and analysed PTMs using nano-liquid

chromatography coupled to high resolution mass spectrometry. Interestingly, common PTMs

including acetylation, methylation and phosphorylation were significantly reduced in EV

compared to nuclear histones (Fig. 6E). For example, in control EVs single H3K9ac and

H3K14ac marks were significantly lower in comparison to the nuclear fraction. More

complex modifications, such as trimethylation (me3) of lysine residues in H3 (Fig. 6E, F)

were also less abundant in EVs. Hence, histones lacking common modifications were more

enriched in EVs, as well as selected mono- or di-methylations (e.g. H3K79me2; Fig. 6F).

Phosphorylation on histone H3 (H3S10ph; Fig. 6F), was also more abundant in EVs,

particularly following LPS treatment, indicating that a proportion of EV-associated histones

may originate from mitotic chromatin. In support of our results with HDAC inhibition or

HAT activation, enrichment of unmodified histones in EVs is a novel finding.

DISCUSSION

Histones are identified consistently amongst the most enriched proteins in EV proteomes[32, 86,

87]. The prevailing view is that EV histones are present in nucleosomes directly associated[30,

31], or copurifying, with EVs[33, 34]. Alternatively, since histones are such abundant cellular

proteins, they may represent a component of the Contaminant Repository for Affinity

Purification ome (CRAPome)[49]. It is important to understand both the location and form of

EV histones since extracellular histones, cell-free DNA and nucleosomes have different

functions[88]. To our knowledge, this is the first study to investigate EV histones without

starting from the *a priori* point of view that they are nucleosomes.

Using a range of mammalian cell lines and experimental approaches, including density

gradient purification, analysis of ILVs within MVBs prior their release as exosomes,

biophysical/chemical fractionation, and IFCM to quantify histones on native EVs, we have

found that: i) Histones localize to ILVs before their secretion as exosomes and colocalize on

the surface of tetraspanin positive EVs (CD63>CD9>CD81). ii) Histone levels decrease in

the cytoplasm and accumulate in MVBs and on EVs in response to cellular stress. iii) The

majority of EV histones are bound to the EV membrane and cannot be removed by

neutralization of electrostatic interactions. Furthermore, unmodified histones and histone

octamers bind efficiently and directly to liposomes and EVs, whereas nucleosomes do not

and association of the majority of histone with EVs seems not to require DNA. iv) Histone

secretion can be blocked by interference with ESCRT-dependent or independent pathways of

EV biogenesis and depend on the autophagosome-lysosome pathway. v) EV-associated

histones are distinct from nuclear histones due to their relative lack of PTMs. The possible

pathways by which histones may associate with EVs are summarised in Fig. 7.

Histones are such well characterized chromatin proteins it is natural to assume that EV

histones are also DNA associated and exist as nucleosomes. A seminal study identifying

histones in dendritic cell derived exosomes[34] noted core histone association with apoptotic

material with a relatively high density (1.16-1.28 g/ml), which would equate to fractions 10-

12 on the OptiPrep gradients used in this study. Our findings indicate that whilst histones

localize to higher density fractions 10-12 in EVs from certain cell lines (e.g. AC-16), or

following cell stress in OLN-93 cells, this is the exception rather than the rule. The findings

of Théry *et al.* then contributed to a leading review that proposed histones as luminal cargo

within large microvesicles and/or apoptotic vesicles, rather than being on the surface of

exosomes[43]. More recently, it was proposed that histones are secreted as a complex with

dsDNA and can be separated from EVs in the non-vesicular fraction[33]. Our findings do not

support this hypothesis. The enrichment of chromosomal DNA in large microvesicles has

been confirmed[89]. However, such large EVs would have been removed by the purification

method used here. Other studies have addressed the localization of DNA at the surface of

EVs[30, 31], though neither of these have investigated the direct binding of free histones to EVs.

Importantly, one factor that has been overlooked consistently is the lack of stoichiometry

reported for core histones detected in EV proteomes[32]. As a specific example, the expected

chromatin/nucleosome H3:H4 ratio of 1:1 is detected in quantitative proteomics of cell lines

by Super-SILAC[90]. However, the same stoichiometry of core histones is not a consistent

feature of in EVs[86], which is confirmed in non-SILAC proteomics of highly purified EVs

intended for therapeutic application[91]. Hence, a parsimonious explanation of the available

data is that although nucleosomes may associate with EVs under certain circumstances and/or

cell types, most EV histones are not nucleosomes. Our results suggest that non-nucleosome

histones comprise a significant component of sEV histones and that they are not dependent

on DNA for EV association. This calls for a careful re-evaluation of the form of histones and

DNA present in EVs subpopulations from different cell types and we provide methodological

approaches to achieve this.

In agreement with proteomic analysis of EV membrane proteins[92], we found that only a small

fraction of EV-associated H3 is found in the luminal fraction of OLN-93 EVs. The majority

of H3 was membrane associated and we identified both a weak interaction with the EV

surface (disrupted with 750 mM NaCl), and a much stronger interaction with the membrane

fraction that could not be disrupted by ionic interference. Could histone EV binding be

mediated primarily by by be due to protein-protein or. protein-glycan interaction? In

Drosophila, H2A and H2B are associated intracellularly with lipid drops[93], which function as

a sink to supply histones for cell division and as antimicrobial effectors during bacterial

infection[94, 95]. Unilamellar lipid drop binding is mediated by the protein chaperone Jabba[95, 96]

and although histone-lipid drop association is conserved in mammals[94], a functional

homologue for Jabba has not been identified. We have not investigated specifically whether

histone-protein interaction contributes to EV binding, but our data shows that all histones can

bind independently to purified membrane derived liposomes. These findings agree with

previous reports. Histones bind directly to membrane phospholipids[97] and binding may occur

via the phosphodiester bonds of phosphate groups within phospholipids, mimicking the

binding of histones in nucleosomes to the phosphodiester bonds in DNA[12]. This is an

important point as it predicts direct competition between phospholipids and DNA for histone

binding, though this remains to be tested

Interestingly, there is also evidence for membrane penetration[97] and translocation[98, 99] of

histones. Penetration was enhanced by negative charge and inhibited by cholesterol,

indicating that phospholipid content and membrane asymmetry is important for histone

binding. In native cell membranes or EVs, negatively charged phospholipids are concentrated

at the inner leaflet, with positively charged phospholipids at the outer surface[100]. In

liposomes, this membrane asymmetry is lost, but in EVs it is unclear how histones might gain

access to negatively charged phospholipids on the inner leaflet. Since the negatively charged

phosphatidylinositol (PI) derivative $PI(3,5)P_2$ is enriched in endosomes-MVBs- lysosomes[101]

, it is tempting to speculate that PIP_2 plays a significant role in histone recruitment and

sequestration into ILVs/exosomes. Therefore, it is important to determine the affinity of

individual and multimerized histones, with and without DNA, for lipid membranes with

different phospholipid composition to see how the lipid environment dictates histone binding

capacity and location/conformation.

Upregulation of histone secretion was a common response to cellular stress. However,

interesting differences were observed in the kinetics, amount, and floating density of histone

positive EV fractions. Upregulation of EV histones was observed following heat stress,

together with a smaller EV size. In contrast, the levels of cytoplasmic histones were reduced

but within MVBs, histones increased, suggesting that stress induces the sequestration and

secretion of cytoplasmically located histones. LPS challenge has also been shown to lead to

an upregulation of EV-associated histones in mouse bone marrow-derived macrophages,

which was proposed to be due to migration of nuclear histones to the cytoplasm[102]. Whatever

the source of cytoplasmically located histones, accumulation can activate signalling pathways

that may lead to apoptosis[19, 103]. Hence, cells may sequester and secrete histones as part of a

stress-associated survival mechanism, whereby vesicles are released continually but histones

are loaded into them in a regulated manner.

In clinical contexts, circulating extracellular histones have been found in various

pathological[19, 104, 105] as well as non-pathological[106] states. Our findings support the idea that

at least some of these circulating histones could be EV associated. However, it is unknown

whether EV associated histones have a different function compared to other types of

extracellular histones identified in sepsis, trauma, or cell death[19, 23]. EVs might represent a

mechanism by which excess histones can be distributed to other cells[99, 107]. Alternatively,

histones may populate the secretory and endosomal pathways and function as innate immune

factors to rapidly respond to pathogenic stimuli, whilst buffering the detrimental effects of

their highly cationic sequences. Though our findings do not address these questions, they do

highlight the importance of accurate characterization of both the nature and the origin of

extracellular histone/DNA complexes since they may have different biological functions as

well as pathological effects.

EV associated histones may recycle from the extracellular space via endocytosis, as treatment

with Dyngo reduced histone secretion (Fig. 7A). However, we also found that both ESCRT-

dependent and ESCRT-independent pathways of EV biogenesis were important for histone

secretion (Fig. 7B, C) suggestive of an intracellular source. MVBs are a key node in the

regulation of secretory turnover, balancing degradation (via late endosome/lysosome fusion)

with recycling (via fusion to the plasma membrane)[108]. Modulation of autophagy and

lysosome maturation significantly affected the secretion of histones. Inhibition of phagophore

formation (3-MA treatment: Fig. 7D) led to a strong upregulation of H3 in EVs. Activation of

autophagy (trehalose, impiramine) had an opposing effect. Furthermore, inhibition of

lysosomal degradation (Baf) increased H3 secretion. Hence, this supports the conclusion that

a proportion of histones normally degraded via this pathway may be routed to MVBs for

secretion via exosomes. This is significant because histones lack a signal sequence and thus

sorting into MVBs/ILVs should occur via a non-canonical pathway. Recent findings have

begun to define pathways by which cytoplasmic histones are targeted to lysosomes for Lamp

2A-mediated degradation[109, 110]. An amphisome/lysosome origin for EV histones could also

explain why most histones associate with the outer surface of EVs, rather than the lumen, as

would be the case if they were sorted through canonical ESCRT-dependent-or ceramide

pathways from the cytoplasm. After being translated and before being incorporated into

chromatin, histones lack most modifications apart from the well-documented exception of

H3K9me and H4K5ac and K12ac[111-113]. Other modifications are deposited on histones after

their incorporation into chromatin. Since our data shows the presence of H3K27, H3K36, and

other H3 and H4 methyl PTMs on EV-associated histones, it indicates that these histones

likely originate from nuclear chromatin. Hence, although our data points towards a nuclear

source for histones and the importance of autophagy, it is unclear how cells may regulate the

routing/sorting of histones and/or DNA through MVBs and into ILVs/EVs, highlighting this

as an important area for further study.

1026

Our results do suggest that the removal of histone PTMs may be important for the

sorting/secretion process. PTMs are known to regulate the localization of histones to[114] and

within[115] the nucleus. The most studied PTMs are acetylation and methylation that are

involved in the regulation of gene expression[116] and are primarily regulated via HAT and

HDAC enzymes. HAT activation via YF-2 increased histone secretion, whereas HAT

inhibition showed the opposite effect. Inhibition of HDAC activity also led to significant

decreases in EV-associated H3 (Fig. 7E). Reduction of neutral PTMs on K/R residues would

increase histone net positive charge and may promote histone binding to lipids during the

sorting process. Since, we could not easily determine how modulation of PTM status related

to histone secretion, we determined the extent of histone PTMs to understand how they might

localize to EVs. The most striking finding was that EV associated histones are relatively

unmodified compared to their nuclear counterparts. Marks such as H3K4me3 and H3K9me3

were almost completely absent in EVs. These two PTMs are the most well characterized

occurrences of trimethylation and define active gene expression and constitutive

heterochromatin, respectively. These findings highlight that it is not just histones from open

or closed chromatin that translocate to EVs; we observed a general loss of modifications that

are known to have critical regulatory role in the nucleus.

1044

CONCLUSION

Our results show clearly that membrane associated histone proteins are secreted specifically

via the MVB/exosome pathway. Secretion is regulated by cellular stress responses, leading to

an upregulation of histones associated with a population of smaller sized EVs. Unexpectedly,

secreted histones do not depend on DNA for EV association and lack many of the of PTMs of

their nuclear counterparts. Understanding how histones are routed to and sorted in to MVB

for secretion, how they bind to membranes, as well as their functional significance are the

next steps.

1053

1054 **DATA AVAILABILITY**

1055 Mass spectrometry raw data availability. All raw mass spectrometry data files from this study

1056 have been submitted to the Chorus repository (https://chorusproject.org/pages/index.html)

1057 under project number 1847.

1058

1059 **AUTHOR CONTRIBUTIONS**

1060 Conception: SELA, MH, JG. Design: All authors. Execution: BS, MS, UM, BN, AG, StSt,

1061 LS, SiSi, JDG. Writing draft: BS, MS, UM, BN, MH, JDG. Reviewing, and editing: BS, MS,

1062 MH, StSt, AG, JZN, OPBW, SiSi, SELA, MH, JDG. All authors read and approved the final

1063 manuscript.

1064

ACKNOWLEDGEMENTS

We are most grateful to colleagues who have contributed time and materials and thank Prof.

C. Richter-Landsberg (Department of Biology, University of Oldenburg, Germany) for

providing OLN-93 cells and Dr. R. Lodge (Department of Medicine, Laval University,

Canada) for the pEGFP-C2 CD63-GFP. Hanna Levén May and Iwan Jones generated the

HA-H3-Flag construct, Sophie Rome, Giulia Corso and Mattias Hällbrink provided valuable

technical advice, and Fazal Oozeer and Adrian Pini contributed with early insights into

extracellular histones. Fransesca Aguilo supplied us with AChE and HDAC1 antibodies to

verify nuclear/cytoplasmic extracts and provided valuable input into the project. We

gratefully acknowledge Chaitali Chakraborty for assistance with TapeStation, the

Biochemical Imaging Center at Umeå University and the National Microscopy Infrastructure,

NMI (VR-RF 2019-00217) for microscopy support, and Umeå Centre for Electron

Microscopy (UCEM) in association with the NMI, VR-RFI2019-00217 and VR-2018-06478

for EM support and method development.

1079

1080   The study was supported by Umeå University, including the Foundation for Medical

1081   Research (Insamlingsstiftelsen), Stiftelsen Forskningsfonden För Klinisk Neurovetenskap

1082   Vid Norrlands Universitetsskjukhus, Västerbotten County Council/Region Västerbotten,

1083   Kempe Foundations, Stratneuro Initiative and Neurofonden.

1084

1085 **DECLARATION OF INTEREST STATEMENT**

1086   SELA is a founder of, and consultant for, Evox Therapeutics. SELA, JN and OW are

1087   shareholders in Evox Therapeutics. The remaining authors declare no competing interests.

1088   All authors declare no conflict of interest.

1089

1090 **REFERENCES**

1091

1.   Kossel, A. Ueber einen peptonartigen Bestandtheil des Zellkerns. (1884).

2.   Kornberg, R.D. Chromatin structure: a repeating unit of histones and DNA. *Science*
184, 868-871 (1974).

3.   Luger, K., Mäder, A.W., Richmond, R.K., Sargent, D.F. & Richmond, T.J. Crystal
structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251-260
(1997).

4.   Simpson, R.T. Structure of the chromatosome, a chromatin particle containing 160
base pairs of DNA and all the histones. *Biochemistry* **17**, 5524-5531 (1978).

5.   Parseghian, M.H. & Luhrs, K.A. Beyond the walls of the nucleus: the role of histones
in cellular signaling and innate immunity. *Biochem Cell Biol* **84**, 589-604 (2006).

6.   Singh, A., Verma, S., Modak, S.B., Chaturvedi, M.M. & Purohit, J.S. Extra-nuclear
histones: origin, significance and perspectives. *Mol Cell Biochem* **477**, 507-524
(2022).

7.   Brinkmann, V. *et al.* Neutrophil extracellular traps kill bacteria. *Science* **303**, 1532-
1535 (2004).

8.   Hiemstra, P.S. *et al.* Antimicrobial proteins of murine macrophages. *Infect Immun* **61**,
3038-3046 (1993).

9.   Hirsch, J.G. Bactericidal action of histone. *J Exp Med* **108**, 925-944 (1958).

10.   Miller, B.F., Abrams, R., Dorfman, A. & Klein, M. ANTIBACTERIAL
PROPERTIES OF PROTAMINE AND HISTONE. *Science* **96**, 428-430 (1942).

11.   Wiesner, J. & Vilcinskas, A. Antimicrobial peptides: the ancient arm of the human
immune system. *Virulence* **1**, 440-464 (2010).

12.   Abrams, S.T. *et al.* Circulating histones are mediators of trauma-associated lung
1115   injury. *Am J Respir Crit Care Med* **187**, 160-169 (2013).
1116   13.   Cheng, Z. *et al.* Circulating Histones Are Major Mediators of Multiple Organ
1117   Dysfunction Syndrome in Acute Critical Illnesses. *Crit Care Med* **47**, e677-e684
1118   (2019).
1119   14.   Huckriede, J. *et al.* Histone H3 Cleavage in Severe COVID-19 ICU Patients. *Front*
1120   *Cell Infect Microbiol* **11**, 694186 (2021).
1121   15.   Karki, P., Birukov, K.G. & Birukova, A.A. Extracellular histones in lung dysfunction:
1122   a new biomarker and therapeutic target? *Pulm Circ* **10**, 2045894020965357 (2020).
1123   16.   Kutcher, M.E. *et al.* Extracellular histone release in response to traumatic injury:
1124   implications for a compensatory role of activated protein C. *J Trauma Acute Care*
1125   *Surg* **73**, 1389-1394 (2012).
1126   17.   Lv, X. *et al.* Extracellular histones are clinically relevant mediators in the
1127   pathogenesis of acute respiratory distress syndrome. *Respir Res* **18**, 165 (2017).
1128   18.   Silk, E., Zhao, H., Weng, H. & Ma, D. The role of extracellular histone in organ
1129   injury. *Cell Death Dis* **8**, e2812 (2017).
1130   19.   Xu, J., Zhang, X., Monestier, M., Esmon, N.L. & Esmon, C.T. Extracellular histones
1131   are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. *J*
1132   *Immunol* **187**, 2626-2631 (2011).
1133   20.   Kawai, C. *et al.* Circulating Extracellular Histones Are Clinically Relevant Mediators
1134   of Multiple Organ Injury. *Am J Pathol* **186**, 829-843 (2016).
1135   21.   Ekaney, M.L. *et al.* Impact of plasma histones in human sepsis and their contribution
1136   to cellular injury and inflammation. *Crit Care* **18**, 543 (2014).
1137   22.   Yokoyama, Y. *et al.* Circulating histone H3 levels in septic patients are associated
1138   with coagulopathy, multiple organ failure, and death: a single-center observational
1139   study. *Thromb J* **17**, 1 (2019).
1140   23.   Xu, J. *et al.* Extracellular histones are major mediators of death in sepsis. *Nat Med* **15**,
1141   1318-1321 (2009).
1142   24.   Li, Y. *et al.* Circulating Histones in Sepsis: Potential Outcome Predictors and
1143   Therapeutic Targets. *Front Immunol* **12**, 650184 (2021).
1144   25.   Wygrecka, M. *et al.* Antihistone Properties of C1 Esterase Inhibitor Protect against
1145   Lung Injury. *Am J Respir Crit Care Med* **196**, 186-199 (2017).
1146   26.   Goldfinch, G.M. *et al.* The proteome of gastric lymph in normal and nematode
1147   infected sheep. *Proteomics* **8**, 1909-1918 (2008).
1148   27.   Mittal, A. *et al.* The proteome of rodent mesenteric lymph. *Am J Physiol Gastrointest*
1149   *Liver Physiol* **295**, G895-903 (2008).
1150   28.   Omenn, G.S. *et al.* Overview of the HUPO Plasma Proteome Project: results from the
1151   pilot phase with 35 collaborating laboratories and multiple analytical groups,
1152   generating a core dataset of 3020 proteins and a publicly-available database.
1153   *Proteomics* **5**, 3226-3245 (2005).
1154   29.   Waga, S., Tan, E.M. & Rubin, R.L. Identification and isolation of soluble histones
1155   from bovine milk and serum. *Biochem J* **244**, 675-682 (1987).
1156   30.   Lazaro-Ibanez, E. *et al.* DNA analysis of low- and high-density fractions defines
1157   heterogeneous subpopulations of small extracellular vesicles based on their DNA
1158   cargo and topology. *J Extracell Vesicles* **8**, 1656993 (2019).
1159   31.   Ghanam, J. *et al.* Extracellular vesicles transfer chromatin-like structures that induce
1160   non-mutational dysfunction of p53 in bone marrow stem cells. *Cell Discov* **9**, 12
1161   (2023).
1162   32.   Mathivanan, S. & Simpson, R.J. ExoCarta: A compendium of exosomal proteins and
1163   RNA. *Proteomics* **9**, 4997-5000 (2009).

33. Jeppesen, D.K. *et al.* Reassessment of Exosome Composition. *Cell* **177**, 428-445.e418
(2019).
34. Théry, C. *et al.* Proteomic analysis of dendritic cell-derived exosomes: a secreted
subcellular compartment distinct from apoptotic vesicles. *J Immunol* **166**, 7309-7318
(2001).
35. Turiák, L. *et al.* Proteomic characterization of thymocyte-derived microvesicles and
apoptotic bodies in BALB/c mice. *J Proteomics* **74**, 2025-2033 (2011).
36. Pan, B.T., Teng, K., Wu, C., Adam, M. & Johnstone, R.M. Electron microscopic
evidence for externalization of the transferrin receptor in vesicular form in sheep
reticulocytes. *J Cell Biol* **101**, 942-948 (1985).
37. Johnstone, R.M., Adam, M., Hammond, J.R., Orr, L. & Turbide, C. Vesicle formation
during reticulocyte maturation. Association of plasma membrane activities with
released vesicles (exosomes). *J Biol Chem* **262**, 9412-9420 (1987).
38. Laulagnier, K. *et al.* Mast cell- and dendritic cell-derived exosomes display a specific
lipid composition and an unusual membrane organization. *Biochem J* **380**, 161-171
(2004).
39. Valadi, H. *et al.* Exosome-mediated transfer of mRNAs and microRNAs is a novel
mechanism of genetic exchange between cells. *Nat Cell Biol* **9**, 654-659 (2007).
40. Thakur, B.K. *et al.* Double-stranded DNA in exosomes: a novel biomarker in cancer
detection. *Cell Res* **24**, 766-769 (2014).
41. Simons, M. & Raposo, G. Exosomes--vesicular carriers for intercellular
communication. *Curr Opin Cell Biol* **21**, 575-581 (2009).
42. Kim, Y.S. *et al.* The potential theragnostic (diagnostic+therapeutic) application of
exosomes in diverse biomedical fields. *Korean J Physiol Pharmacol* **22**, 113-125
(2018).
43. Colombo, M., Raposo, G. & Théry, C. Biogenesis, secretion, and intercellular
interactions of exosomes and other extracellular vesicles. *Annu Rev Cell Dev Biol* **30**,
255-289 (2014).
44. Yokoi, A. *et al.* Mechanisms of nuclear content loading to exosomes. *Sci Adv* **5**,
eaax8849 (2019).
45. Gilthorpe, J.D. *et al.* Extracellular histone H1 is neurotoxic and drives a pro-
inflammatory response in microglia. *F1000Res* **2**, 148 (2013).
46. Hsu, C. *et al.* Regulation of exosome secretion by Rab35 and its GTPase-activating
proteins TBC1D10A-C. *J Cell Biol* **189**, 223-232 (2010).
47. Krämer-Albers, E.M. *et al.* Oligodendrocytes secrete exosomes containing major
myelin and stress-protective proteins: Trophic support for axons? *Proteomics Clin*
Appl **1**, 1446-1461 (2007).
48. Trajkovic, K. *et al.* Ceramide triggers budding of exosome vesicles into
multivesicular endosomes. *Science* **319**, 1244-1247 (2008).
49. Mellacheruvu, D. *et al.* The CRAPome: a contaminant repository for affinity
purification-mass spectrometry data. *Nat Methods* **10**, 730-736 (2013).
50. Richter-Landsberg, C. & Heinrich, M. OLN-93: a new permanent oligodendroglia cell
line derived from primary rat brain glial cultures. *J Neurosci Res* **45**, 161-173 (1996).
51. Morissette, G., Lodge, R., Bouthillier, J. & Marceau, F. Receptor-independent,
vacuolar ATPase-mediated cellular uptake of histamine receptor-1 ligands: possible
origin of pharmacological distortions and side effects. *Toxicol Appl Pharmacol* **229**,
320-331 (2008).
52. Goldbaum, O. *et al.* Proteasome inhibition stabilizes tau inclusions in
oligodendroglial cells that occur after treatment with okadaic acid. *J Neurosci* **23**,
8872-8880 (2003).

53. Forterre, A. *et al.* Proteomic analysis of C2C12 myoblast and myotube exosome-like
vesicles: a new paradigm for myoblast-myotube cross talk? *PLoS One* **9**, e84153
(2014).
54. Al-Ani, A. *et al.* Oxygenation in cell culture: Critical parameters for reproducibility
are routinely not reported. *PLoS One* **13**, e0204269 (2018).
55. Loo, A.E., Ho, R. & Halliwell, B. Mechanism of hydrogen peroxide-induced
keratinocyte migration in a scratch-wound model. *Free Radic Biol Med* **51**, 884-892
(2011).
56. Shechter, D., Dormann, H.L., Allis, C.D. & Hake, S.B. Extraction, purification and
analysis of histones. *Nat Protoc* **2**, 1445-1457 (2007).
57. Li, J. *et al.* Serum-free culture alters the quantity and protein composition of
neuroblastoma-derived extracellular vesicles. *J Extracell Vesicles* **4**, 26883 (2015).
58. Perez-Jimenez, R., Godoy-Ruiz, R., Ibarra-Molero, B. & Sanchez-Ruiz, J.M. The
efficiency of different salts to screen charge interactions in proteins: a Hofmeister
effect? *Biophys J* **86**, 2414-2429 (2004).
59. Riendeau, D., Lemaire, J., Maestracci, D. & Lessard, L. Selective release of inner
core proteins from intestinal microvillus membrane by lithium diiodosalicylate. *Mol*
Cell Biochem **71**, 45-52 (1986).
60. Bittame, A. *et al.* Lipid Extraction from HeLa Cells, Quantification of Lipids,
Formation of Large Unilamellar Vesicles (LUVs) by Extrusion and in vitro Protein-
lipid Binding Assays, Analysis of the Incubation Product by Transmission Electron
Microscopy (TEM) and by Flotation across a Discontinuous Sucrose Gradient. *Bio-*
Protocol **6** (2016).
61. Tertel, T. *et al.* High-Resolution Imaging Flow Cytometry Reveals Impact of
Incubation Temperature on Labeling of Extracellular Vesicles with Antibodies.
Cytometry A **97**, 602-609 (2020).
62. Gorgens, A. *et al.* Optimisation of imaging flow cytometry for the analysis of single
extracellular vesicles by using fluorescence-tagged vesicles as biological reference
material. *J Extracell Vesicles* **8**, 1587567 (2019).
63. Dar, G.H. *et al.* GAPDH controls extracellular vesicle biogenesis and enhances the
therapeutic potential of EV mediated siRNA delivery to the brain. *Nat Commun* **12**,
6666 (2021).
64. Gorgens, A. *et al.* Identification of storage conditions stabilizing extracellular vesicles
preparations. *J Extracell Vesicles* **11**, e12238 (2022).
65. Yuan, Z.F. *et al.* EpiProfile 2.0: A Computational Platform for Processing Epi-
Proteomics Mass Spectrometry Data. *J Proteome Res* **17**, 2533-2541 (2018).
66. Hedlund, M. *et al.* Human placenta expresses and secretes NKG2D ligands via
exosomes that down-modulate the cognate receptor expression: evidence for
immunosuppressive function. *J Immunol* **183**, 340-351 (2009).
67. Stenqvist, A.C., Nagaeva, O., Baranov, V. & Mincheva-Nilsson, L. Exosomes
secreted by human placenta carry functional Fas ligand and TRAIL molecules and
convey apoptosis in activated immune cells, suggesting exosome-mediated immune
privilege of the fetus. *J Immunol* **191**, 5515-5523 (2013).
68. Tardoski, S. *et al.* Low-Intensity Ultrasound Promotes Clathrin-Dependent
Endocytosis for Drug Penetration into Tumor Cells. *Ultrasound Med Biol* **41**, 2740-
2754 (2015).
69. Möbius, W. & Posthuma, G. Sugar and ice: Immunoelectron microscopy using
cryosections according to the Tokuyasu method. *Tissue Cell* **57**, 90-102 (2019).

70.   Cantin, R., Diou, J., Bélanger, D., Tremblay, A.M. & Gilbert, C. Discrimination
between exosomes and HIV-1: purification of both vesicles from cell-free
supernatants. *J Immunol Methods* **338**, 21-30 (2008).
71.   Graner, M.W. *et al.* Proteomic and immunologic analyses of brain tumor exosomes.
Faseb j **23**, 1541-1557 (2009).
72.   Chen, R., Kang, R., Fan, X.G. & Tang, D. Release and activity of histone in diseases.
Cell Death Dis **5**, e1370 (2014).
73.   Balaj, L. *et al.* Tumour microvesicles contain retrotransposon elements and amplified
oncogene sequences. *Nat Commun* **2**, 180 (2011).
74.   Osteikoetxea, X. *et al.* Differential detergent sensitivity of extracellular vesicle
subpopulations. *Org Biomol Chem* **13**, 9775-9782 (2015).
75.   Bittame, A. *et al.* Lipid Extraction from HeLa Cells, Quantification of Lipids,
Formation of Large Unilamellar Vesicles (LUVs) by Extrusion and in vitro Protein-
lipid Binding Assays, Analysis of the Incubation Product by Transmission Electron
Microscopy (TEM) and by Flotation across a Discontinuous Sucrose Gradient. *Bio-
protocol* **6**, e1963 (2016).
76.   Essandoh, K. *et al.* Blockade of exosome generation with GW4869 dampens the
sepsis-induced inflammation and cardiac dysfunction. *Biochim Biophys Acta* **1852**,
2362-2371 (2015).
77.   Yao, Y. *et al.* Blockade of Exosome Release Suppresses Atrial Fibrillation by
Alleviating Atrial Fibrosis in Canines With Prolonged Atrial Pacing. *Front
Cardiovasc Med* **8**, 699175 (2021).
78.   Mauvezin, C. & Neufeld, T.P. Bafilomycin A1 disrupts autophagic flux by inhibiting
both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent
autophagosome-lysosome fusion. *Autophagy* **11**, 1437-1438 (2015).
79.   McCluskey, A. *et al.* Building a better dynasore: the dyngo compounds potently
inhibit dynamin and endocytosis. *Traffic* **14**, 1272-1289 (2013).
80.   Bártová, E., Krejcí, J., Harnicarová, A., Galiová, G. & Kozubek, S. Histone
modifications and nuclear architecture: a review. *J Histochem Cytochem* **56**, 711-721
(2008).
81.   Campos, E.I. *et al.* The program for processing newly synthesized histones H3.1 and
H4. *Nat Struct Mol Biol* **17**, 1343-1351 (2010).
82.   Wolf, M. *et al.* A functional corona around extracellular vesicles enhances
angiogenesis, skin regeneration and immunomodulation. *J Extracell Vesicles* **11**,
e12207 (2022).
83.   Nordin, J.Z. *et al.* Ultrafiltration with size-exclusion liquid chromatography for high
yield isolation of extracellular vesicles preserving intact biophysical and functional
properties. *Nanomedicine* **11**, 879-883 (2015).
84.   Heusermann, W. *et al.* Exosomes surf on filopodia to enter cells at endocytic hot
spots, traffic within endosomes, and are targeted to the ER. *J Cell Biol* **213**, 173-184
(2016).
85.   Ricklefs, F.L. *et al.* Imaging flow cytometry facilitates multiparametric
characterization of extracellular vesicles in malignant brain tumours. *J Extracell
Vesicles* **8**, 1588555 (2019).
86.   Kugeratski, F.G. *et al.* Quantitative proteomics identifies the core proteome of
exosomes with syntenin-1 as the highest abundant protein and a putative universal
biomarker. *Nat Cell Biol* **23**, 631-641 (2021).
87.   Das, R. & Plow, E.F. Phosphatidylserine as an anchor for plasminogen and its
plasminogen receptor, histone H2B, to the macrophage surface. *J Thromb Haemost* **9**,
339-349 (2011).

88. Marsman, G., Zeerleder, S. & Luken, B.M. Extracellular histones, cell-free DNA, or
nucleosomes: differences in immunostimulation. *Cell Death Dis* **7**, e2518 (2016).
89. Vagner, T. *et al.* Large extracellular vesicles carry most of the tumour DNA
circulating in prostate cancer patient plasma. *J Extracell Vesicles* **7**, 1505403 (2018).
90. Geiger, T., Cox, J., Ostasiewicz, P., Wisniewski, J.R. & Mann, M. Super-SILAC mix
for quantitative proteomics of human tumor tissue. *Nat Methods* **7**, 383-385 (2010).
91. Vo, N. *et al.* A novel multi-stage enrichment workflow and comprehensive
characterization for HEK293F-derived extracellular vesicles. *J Extracell Vesicles* **13**,
e12454 (2024).
92. Cvjetkovic, A. *et al.* Detailed Analysis of Protein Topology of Extracellular Vesicles-
Evidence of Unconventional Membrane Protein Orientation. *Sci Rep* **6**, 36338 (2016).
93. Cermelli, S., Guo, Y., Gross, S.P. & Welte, M.A. The lipid-droplet proteome reveals
that droplets are a protein-storage depot. *Curr Biol* **16**, 1783-1795 (2006).
94. Anand, P. *et al.* A novel role for lipid droplets in the organismal antibacterial
response. *Elife* **1**, e00003 (2012).
95. Li, Z. *et al.* Lipid droplets control the maternal histone supply of Drosophila embryos.
Curr Biol **22**, 2104-2113 (2012).
96. Stephenson, R.A. *et al.* Sequestration to lipid droplets promotes histone availability
by preventing turnover of excess histones. *Development* **148** (2021).
97. Kleine, T.J., Lewis, P.N. & Lewis, S.A. Histone-induced damage of a mammalian
epithelium: the role of protein and membrane structure. *Am J Physiol* **273**, C1925-
1936 (1997).
98. Rosenbluh, J. *et al.* Translocation of histone proteins across lipid bilayers and
Mycoplasma membranes. *J Mol Biol* **345**, 387-400 (2005).
99. Hariton-Gazal, E., Rosenbluh, J., Graessmann, A., Gilon, C. & Loyter, A. Direct
translocation of histone molecules across cell membranes. *J Cell Sci* **116**, 4577-4586
(2003).
100. Cockcroft, S. Mammalian lipids: structure, synthesis and function. *Essays Biochem*
65, 813-845 (2021).
101. Clifton, L.A. *et al.* Creation of distinctive Bax-lipid complexes at mitochondrial
membrane surfaces drives pore formation to initiate apoptosis. *Sci Adv* **9**, eadg7940
(2023).
102. Nair, R.R. *et al.* LPS-Challenged Macrophages Release Microvesicles Coated With
Histones. *Front Immunol* **9**, 1463 (2018).
103. Ibañez-Cabellos, J.S. *et al.* Extracellular histones activate autophagy and apoptosis via
mTOR signaling in human endothelial cells. *Biochim Biophys Acta Mol Basis Dis*
1864, 3234-3246 (2018).
104. Gould, T.J., Lysov, Z. & Liaw, P.C. Extracellular DNA and histones: double-edged
swords in immunothrombosis. *J Thromb Haemost* **13 Suppl 1**, S82-91 (2015).
105. Kim, J.E., Lee, N., Gu, J.Y., Yoo, H.J. & Kim, H.K. Circulating levels of DNA-
histone complex and dsDNA are independent prognostic factors of disseminated
intravascular coagulation. *Thromb Res* **135**, 1064-1069 (2015).
106. Holdenrieder, S. *et al.* Circulating nucleosomes in serum. *Ann N Y Acad Sci* **945**, 93-
102 (2001).
107. Sulkowski, P.L. *et al.* Horizontal Transfer of Histone H3 by Mammalian Cells.
bioRxiv, 2021.2006.2030.450524 (2021).
108. Cohen, M.J., Chirico, W.J. & Lipke, P.N. Through the back door: Unconventional
protein secretion. *Cell Surf* **6**, 100045 (2020).

109.   Hormazabal, J. *et al.* Chaperone mediated autophagy contributes to the newly
1361   synthesized histones H3 and H4 quality control. *Nucleic Acids Research* **50**, 1875-
1362   1887 (2022).

1363   110.   Ferreira, J.V. *et al.* LAMP2A regulates the loading of proteins into exosomes. *Sci Adv*
1364   **8**, eabm1140 (2022).

1365   111.   Sobel, R.E., Cook, R.G., Perry, C.A., Annunziato, A.T. & Allis, C.D. Conservation of
1366   deposition-related acetylation sites in newly synthesized histones H3 and H4. *Proc*
1367   *Natl Acad Sci U S A* **92**, 1237-1241 (1995).

1368   112.   Carraro, M. *et al.* DAXX adds a de novo H3.3K9me3 deposition pathway to the
1369   histone chaperone network. *Mol Cell* **83**, 1075-1092 e1079 (2023).

1370   113.   Loyola, A., Bonaldi, T., Roche, D., Imhof, A. & Almouzni, G. PTMs on H3 variants
1371   before chromatin assembly potentiate their final epigenetic state. *Mol Cell* **24**, 309-
1372   316 (2006).

1373   114.   Bernardes, N.E. & Chook, Y.M. Nuclear import of histones. *Biochem Soc Trans* **48**,
1374   2753-2767 (2020).

1375   115.   Bowman, G.D. & Poirier, M.G. Post-Translational Modifications of Histones That
1376   Influence Nucleosome Dynamics. *Chem Rev* **115**, 2274-2295 (2015).

1377   116.   Allfrey, V.G., Faulkner, R. & Mirsky, A.E. ACETYLATION AND METHYLATION
1378   OF HISTONES AND THEIR POSSIBLE ROLE IN THE REGULATION OF RNA
1379   SYNTHESIS. *Proc Natl Acad Sci U S A* **51**, 786-794 (1964).

1380

1381

FIGURE LEGENDS

Figure 1. Histones are associated with ILVs and EVs. (A) WB of whole cell lysate (WCL,

10 μg) from a stable OLN-93 cell line expressing human CD63-EGFP and EVs (10, 20 μg)

from 100,000 xg pellet. CD9, CD63, Tsg101, core histones (H2A, H2B, H3 and H4), Golgi

marker GM130 as well as Actb identified using specific antibodies. Approximate molecular

mass is shown in kDa. **(B)** Immuno-TEM micrographs of OLN-93 EV$^{100,000 \text{ xg}}$ using anti-

CD63 primary antibody and secondary antibody conjugated to with 10 nm gold. **(C)** Confocal

micrographs from a single z-plane of OLN-93 cells showing colocalization of anti-H3 (red)

with anti-CD63 (green). Nuclei were counterstained with DAPI (blue). Inset shows a cropped

and expanded region of the main image. **(D)** Confocal micrograph showing Duolink staining

(red) due to colocalization of anti-H3 (rabbit) and anti-CD63 (mouse) antibodies. **(E)** WB

showing localization of a range of membrane, EV and organelle markers together with linker

and core histones in OptiPrep purified EV$^{100,000 \text{ xg}}$ (300 μg) fractions from a 12%-36% (w/v)

equilibrium density gradient. One 20$^{\text{th}}$ of each fraction was loaded on the gel. Experiments

were repeated 3 times, and one representative image is shown. **(F)** Representative immuno-

TEM micrograph of OLN-93 cell sections showing anti-CD63. Inset panel shows a magnified

region containing an MVB. The column chart on the right shows quantification of the total

number of MVBs versus CD63^{+} MVBs (black), and total number of ILVs versus CD63^{+}

ILVs (red) as % of total. **(G)** as for (F) but showing anti-H1. Column chart on the right shows

the total number and H1-positive ILVs per MVB. Error bars represent the mean ±SD. **(H)**

Immuno-TEM showing differential labelling of anti-CD63 (5 nm gold, open arrowheads) and

anti-H1 (10 nm gold, solid arrowheads). Panels below (1 and 2) show higher magnification

images of the regions indicated in H. Solid arrowheads indicate anti-H1 labelling and open

arrowheads indicate anti-CD63 labelling. **(I)** Scatter plot showing quantification of CD63 and

H1 labelling and colocalization (on the same ILV). A total of 10 images from separate cells

and sections were quantified and the numbers of ILVs per MVB are represented by each

point. Error bars represent the mean ±SD. **(J)** Immuno-TEM micrographs of OLN-93

EV^{OptiPrep} showing anti-CD63 (10 nm gold, solid arrowheads) and anti-H1 (5 nm gold, open

arrowheads). Note gold particle sizes are opposite to those used in (F).

1411

Figure 2. **Cellular stress enhances the release of histones via ILVs/EVs.** **(A)** Schematic

diagram outlining the experimental procedure for the heat stress experiment, showing sample

collection times for Before, Heat stress and After conditions. **(B)** WB showing markers

(specific antibodies as shown) detected after pulldown with StrataClean resin of total protein

from 1 ml of conditioned media from Before, Heat stress and After samples, as well as fresh,

unconditioned media (Mock). **(C)** WB of EV$^{100,000\ xg}$ (11 µg per lane) isolated from the same

conditioned media samples shown in (B). **(D-E)** OLN-93 cells were stressed by heat, LPS,

hypoxia, and H_2O_2 followed by WB to quantify histone H2B and H3 in cytoplasm and

EV$^{100,000\ xg}$. Column charts show the ratio between band intensities for stress versus control

conditions. Three independent experiments were performed, and one representative WB

image is shown. **(F)** Immuno-TEM images of control and stressed OLN-93 cell sections

showing ILVs within an MVB. Anti-CD63 (5 nm gold; open arrowheads) and anti-H1 **(**10

nm gold**;** closed arrowheads). **(G)** Quantification of number of H1 positive ILVs per MVB

present under the various conditions. **(H)** Co-localisation of anti-CD63/H1 on the same ILV

under control and stress conditions. Error bars represent the SD. One-way ANOVA was used

to compare control and stress conditions. *, $p \leq 0.05$, ***, $p \leq 0.001$. **(I)** Representative TEM

micrographs of negatively stained OLN-93 EV^{OptiPrep} isolated from control and stress

conditions as indicated in each panel. **(J)** Scatter plots showing size distribution of EVs (n =

500 EVs measured from n=10 randomly selected images) measured in TEM micrographs

from each condition. Statistical significance between control and stress conditions (***, $p \leq$

0.001) using one-way ANOVA followed by Bonferroni's multiple comparison post-test.

Horizontal bars in each plot represent the mean diameter in nm, shown below with the ±SD,

as well as the mode size (in 10 nm bins).

1435

Figure 3. Histones associate with the EV membrane. (A) Line graphs (above) showing

quantification of protein (dashed lines) and DNA (solid lines) by Qubit in OptiPrep fractions

from different stress conditions. WBs (below) showing distribution of H3 in OptiPrep

fractions. **(B)** Schematic diagram showing the basic procedure used to extract histones from

EV^{OptiPrep}. **(C)** TEM images showing control (Ctrl) EVs or following 750 mM NaCl

treatment. **(D)** Extraction of histones from OptiPrep purified (fractions 4-7; EV^{OptiPrep}).

Approximately 10 µg EVs were treated with 150-750 mM NaCl for 24h at 4ºC, followed by

ultracentrifugation and estimation of H3 in the supernatants and pellets by dot blot. **(E)** Line

graphs showing the intensities of dot blots (H3 quantification, black) and DNA content

(Qubit assay). Statistical analysis between 150 mM NaCl as control and other NaCl

concentrations by one-way ANOVA, *, $p \leq 0.05$, **, $p \leq 0.01$. **(F)** Extraction of histones in

the supernatant (S) and remaining in the pellet (P) from EV^{OptiPrep} treated with 750 mM NaCl

and shown by WB. **(G)** S and P fractions of EV^{OptiPrep} after extraction with 750 mM NaCl,

then sonication and re-pelleting, followed by a further round of extraction with 750 mM NaCl

to remove luminal contents. The remaining pellet shows residual H3 that was not extracted

by 750 mM NaCl. **(H)** TEM images showing intact EV membranes after 25 mM lithium 3,5-

diiodosalicylate treatment. **(I)** Extraction of histones into the supernatant (S) or remaining in

the pellet of EV^{OptiPrep} following treatment with increasing concentrations of lithium 3,5-

diiodosalicylate (10-50 mM) or Triton X-100 (0.1% TX-100). **(J)** WB showing H3 present in

EV^{OptiPrep} (control or sonicated) with and without nuclease digestion (DNAseI, S1 nuclease

and Exosnuclease III). **(K)** Column chart showing DNA quantification (Qubit) in same

samples as (J). **(L)** TapeStation gel image of total EV DNA purified from control and

digested samples. **(M)** Column chart showing quantification of DNA by TapeStation and, **(N)**

Quantification of DNA in same samples using Qubit. **(O)** Analysis of OLN-93 EV^{OptiPrep} by

WB without (Ctrl) or following trypsin treatment (1:1500-1:150 µg trypsin:µg EV protein).

Presence of truncated bands for histone proteins following trypsin treatment are indicated on

right by open arrowheads. **(P)** Schematic diagram of epitope tagged H3 cassette used to

express the protein in a HEK293 stable cell line. HA-tag is located at the N-terminus and a

6xHis-tag at the C-terminus (not to scale). **(Q)** WB analysis of EV$^{100,000 \, xg}$ isolated from

stable HA-H3-6xHis HEK293 cell line after trypsin treatment. Location of full-length HA-

H3-6xHis, as well as partially degraded species are indicated by a black or open arrowheads,

respectively. Control recombinant histone digestion by trypsin is shown in supplementary

Fig. S12.

1469

Figure 4. Preferential membrane binding of histones and octamers versus nucleosomes

(A) Analysis of histone interaction with liposomes. Total histones or individual recombinant

histones (as indicated) were mixed with liposomes derived from OLN-93 total cellular lipids

and washed extensively in a 100,000 MWCO spin concentrator. The retentate was subjected

to WB with specific anti-histone antibodies. A histone protein-only control (without

liposomes) was used to control for lipid-independent retention of histones by the MWCO

filter. Histone proteins (200 ng as indicated) were loaded as references. **(B)** Liposome

flotation on a sucrose density gradient (0-40%). Histones as in (A) were mixed with

liposomes and then added to the bottom layer (40% sucrose) of the gradient and subjected to

ultracentrifugation at 100,000 xg. Fractions were collected and analysed by WB. **(C)**

Recombinant histone octamers (5 µg) proteins were mixed with liposomes (20 µg) and

separated as in (B). Octamers were detected by WB using anti-His tag antibody. **(D)** As for

(C) but using recombinant nucleosomes instead of octamers. **(E)** Dot blot showing binding of

octamers or nucleosomes to EV^{OptiPrep}. Serially diluted EVs were blotted onto a nitrocellulose

membrane and probed with 2 µg/ml octamers/nucleosomes. The bound fractions were

detected using an anti-His tag antibody. **(F)** Line graphs showing the intensity of dot blots

plotted from (E). Two-way ANOVA and Bonferroni multiple comparison test was

performed: **, $p \leq 0.01$, ***, $p \leq 0.001$,

1488

Figure 5. Histone secretion is influenced by multiple cellular pathways. **(A)** WBs

showing the effects of siRNA mediated knock-down of *Alix*, *Tsg101*, and *Smpd2* in OLN-93

cells. H3 and EV markers (Alix, Tsg101 as well as the plasma membrane/microvesicle

marker Annexin A1) were quantified in total protein recovered from the media by

StrataClean pulldown. The experiment was repeated 4 times, and a representative blot is

shown. Band intensities were quantified, and the column chart below shows the ratio of H3

(treated/control) from four independent experiments. **(B)** Pharmacological treatment of OLN-

93 cells, analysed as in (A). Control is without treatment. **(C)** Immuno-TEM micrographs

showing anti-H1 labelling associated with ILVs in OLN-93 cell sections, following treatment

with compounds that significantly increased (3-MA or bafilomycin) or decreased (sodium

butyrate) H3 secretion into the media. Right, scatter plot showing quantification of histone

positive ILVs under control and following treatment. One-way ANOVA: *, ≤ 0.05, **, $p \leq$

0.01, ***, $p \leq 0.001$.

1502

Figure 6: Histones are surface associated in intact EVs and carry less PTMs than their

nuclear counterparts (A) Plots showing representative IFCM data obtained from HeLa cell

conditioned media (CM) and media controls stained with either AF488 labelled anti-H3 or

FITC labelled anti-H4 antibodies, or a mixture of APC labelled anti-CD9/CD63/CD81

antibodies to detect total tetraspanin positive EVs. **(B)** Quantification of detected gateable

events positive for single stained H3, H4, or tetraspanins (n=3) in CM samples collected at

different time points from both control and H_2O_2 treated cell cultures. Samples were diluted

post staining as indicated on the *y*-axes. **(C)** H3 co-labelling and **(D)** H4 co-labelling with

anti-CD9, -CD63, or -CD81 antibodies in CM from 24 h H_2O_2 treated samples. See also Fig.

S15. Statistical comparisons by two-way ANOVA. **, ≤ 0.005, ***, $p \leq 0.001$, ****, $p \leq$

0.0001. **(E)** Radial plot showing relative abundance of H3 post-translational modifications

(PTMs) identified in $EV^{OptiPrep}$ versus nuclear histones. K = lysine, ac = acetylation, me =

mono- (me1), di- (me2) or tri- (me3) methylation, ph = phosphorylation. **(F)** Representative

examples of H3 PTMs that were selectively enriched in the nucleus, or in EVs. Lysine (K)

trimethylation (me3) results from the covalent attachment of three me groups to a K residue

and is therefore a relatively "slow" modification. H3 is poorly modified with me3 in EVs

relative to nuclei regardless of whether this PTM benchmarks actively transcribed chromatin

(H3K4me3) or silenced heterochromatin (H3K9me3). H3K9 is relatively unmodified in EVs,

and H3K79me2 is an example of a modification that is enriched in EV histones.

Phosphorylation of H3S10 is selectively enriched in EVs from LPS treated cells.

1523

Figure 7: Potential pathways for histone secretion via exosomes. Schematic diagram

summarizing the findings of this paper. **(A)** Extracellular histones (yellow diamonds)

internalized by endocytosis would undergo inward budding into MVBs and associate with the

outer surface of ILVs. Cytoplasmic histones (red diamonds) sorted during **(B)** ESCRT-

dependent or **(C)** –independent (ceramide pathway) ILV formation may localize as

internal/luminal cargo (red diamonds). **(D)** Histones originating from the nucleus (green

diamonds) and captured by autophagosomes may partition to ILVs after amphisomes or

lysosomes fuse with MVBs. **(E)** Posttranslational modification of histones in subcellular

compartments such as the cytoplasm may also be important for sorting to MVBs/ILVs. **(F)**

The possible route whereby histones may traffic from the nucleus to amphisomes is

unknown.

1535

1536   **Supplementary Table S1.** Summary of drugs used in Fig. 5B-C.

1537   **Supplementary figures S1-S15.** The legends text for supplementary figures is included in

1538   single PDF file.

OptiPrep fractions