Evidence that host-mediated epigenetic modifications regulate gene expression in *Wolbachia pipientis*

Stella Papaleo[1$], Simona Panelli[1], Ibrahim Bitar[2], Lodovico Sterzi[1], Riccardo Nodari[1,3], Francesco Comandatore[1*$]

1. Department of Biomedical and Clinical Sciences, Pediatric Clinical Research Center "Romeo and Enrica Invernizzi", University of Milan, 20157 Milan, Italy.
2. Biomedical Center, Faculty of Medicine, Charles University, Pilsen, Czechia.
3. Istituto Nazionale di Genetica Molecolare (INGM) "Romeo and Enrica Invernizzi", University of Milan, 20157 Milan, Italy.

* Corresponding author

\$ The authors contributed equally

Abstract

Wolbachia pipientis is an obligate intracellular bacterium, associated with several arthropods and filarial nematodes. *Wolbachia* establishes strict symbiotic relationships with its hosts, with the consequent loss of many genes and regulatory regions. Despite this, experimental studies show that *Wolbachia* gene expression is coordinated to host needs, but the mechanism is still unknown. The first published RNA-Seq study on *Wolbachia* evidenced a strong differential expression of a DNA methyltransferase (MTase). In bacteria, this enzyme methylates either adenines or cytosines on specific motifs, contributing to the regulation of gene expression. In this work, we tested the hypothesis that the activity of MTase modulates the expression of *Wolbachia* genes. We first determined the methylation motif of the *Wolbachia* MTase by expressing it in *Escherichia coli*. Surprisingly, the experiment revealed that the *Wolbachia* MTase methylates both adenine and cytosine, without recognising highly specific motifs. Then, re-analysing data from six RNA-Seq studies, we found that the nucleotide content of *Wolbachia* genes correlates with their expressions, with a pattern compatible to be a consequence of the DNA methylation. Lastly, we identified MTase as the *Wolbachia* gene with the most conserved binding site for the Ccka/CtrA signalling transduction system, a mechanism likely involved the host-bacterium communication. Overall, these findings suggest a cascade mechanism in which the host activates the *Wolbachia* Ccka/CtrA signalling system, thus inducing the expression of the MTase gene. Then, the subsequent DNA methylation will affect the expression of several *Wolbachia* genes on the basis of their cytosine and adenine content.

Introduction

Wolbachia pipientis (from here Wolbachia) is an intracellular bacterium belonging to the Alphaproteobacteria class. It is the most prevalent endosymbiotic microbe in the animal world, colonising many arthropods species and filarial nematodes(1–4). In arthropods, the bacterium can manipulate the reproduction of the host(5), protect it from viruses(6, 7) and provide nutrients(8). In filarial nematodes, genomic analyses and experimental evidence revealed that Wolbachia is essential for the host survival and development (9). The bacterium is strictly intracellular, with a tropism for microtubules and with the ability to colonise endoplasmic reticulum (ER), with important effects on host cell replication(2). It is mainly mother-to-offspring transmitted, being actively transported by specialised cells of the host from the soma to germ cells (2). Nevertheless, horizontal transfer among host individuals can occur (10, 11), ehnancing the bacterium spreading (11).

As for many other strictly host-associated bacteria, the Wolbachia genome underwent an important reduction during its evolution, reaching a size that ranges from ~0.8 Mb to ~1.8 Mb (BV-BRC representative genomes)(12). As observed for several endosymbiont bacterial species, genome reduction passes through the increasing of the number of mobile elements (e.g. Insertion Sequences) and the loss of many genes(13–15). During this process, the structure of the genome and the order of genes can change dramatically(14), leading to the disruption of several operons and loss of most gene promoters. Interestingly, RNA-seq studies have highlighted that Wolbachia gene expression can be influenced by host environmental/developmental conditions(16–22). This suggests that Wolbachia likely has a mechanism for regulating gene expression, despite the low number of promoters indicating it may not depend solely on transcription factors.

The first published Wolbachia RNA-seq study(16) compared gene expression of Wolbachia colonising somatic tissues and gonads in females and males of the filarial nematode

Onchocerca ochengi. One of the strongest differential expressions was reported for a gene

annotated as a S-adenosyl-methyonine-dependent (SAM)-dependent DNA

methyltransferase (hereafter "MTase"). In general, DNA methyltransferase enzymes are able

to methylate either adenine or cytosine on specific motifs of bacterial DNA. The most studied

MTase are the DNA adenine methyltransferase (Dam) and DNA cytosine methyltransferase

(Dcm) of *Escherichia coli*. The first one methylates the adenine residues in the sequence

GATC, while the other one targets the second cytosine within the motifs CCAGG and

CCTGG (23, 24). Furthermore, in *Alphaproteobacteria*, one of most characterised

methyltransferase is the CcrM, which methylates the adenine in the motif GAnTC, and it is

involved in the regulation of the cell cycle in *Caulobacter crescentus (25)*.

Despite DNA methylation has deeply been investigated in eukaryotes, as an epigenetic

mechanism of gene expression, its role in prokaryotes is still poorly investigated and no

studies have been published on *Wolbachia*.

Despite the effect of DNA methylation on gene expression has been deeply investigated in

eukaryotes (i.e. epigenetics), its role in prokaryotes physiology is still poorly understood and

no studies have been published on *Wolbachia.* At the state of the art, there is evidence that

DNA methyltransferase enzymes are involved in fundamental bacterial processes, including

proofreading during DNA replication, protection from foreign DNA and regulation of gene

expression. A specific type of DNA methyltransferase, called orphan DNA

methyltransferases, methylates gene promoters, changing their affinity for transcription

factors with important effects on gene expression (23, 24).

In this study, we investigated the hypothesis that DNA methylation is a key mechanism for

the regulation of the *Wolbachia* gene expression. First, we experimentally determined the

methylation motif of the MTase enzyme encoded in the genome of the *Wolbachia*

endosymbiont of *O. ochengi* (wOo). Then, we tested whether the frequency of this motif into

gene sequences correlates with their expression, and whether the intensity of the

phenomenon is related to the levels of expression of the MTase gene. Finally, we identified

regulatory regions upstream of the MTase gene in *Wolbachia* genomes. Our results suggest

that the host may specifically induce the expression of the *Wolbachia* MTase gene through a

signalling transduction system located on the bacterium's membrane. Subsequently, the

MTase methylates the *Wolbachia* genome, altering the expression of methylated genes

based on their nucleotide composition.

Material and methods

Experimental determination of the wOo SAM-dependent DNA methyltransferase

methylation pattern

The wOo SAM-dependent DNA methyltransferase (MTase) was artificially synthesised with

codon usage adaptation and cloned into the XhoI and SalI sites of the pCOLD III expression

vector (Takara Bio). The recombinant construct was transformed into *Escherichia coli*

StellarTM Competent Cells (*dam–/dcm–*) (Takara Bio) according to the supplier's protocol

(ClonTech; Protocol-at-a-Glance, PT5056-2). The *E. coli* strains were cultured in LB broth

medium supplemented with ampicillin. Before inducing expression, PCR with custom primers

targeting the gene of interest was performed on bacterial crude extracts to verify the

outcome of transformation. MTase expression was then induced at 15°C, according to the

protocol provided by the manufacturer for the Cold Shock Expression System of pColdTM

plasmids (Takara Bio). 24 hours after the induction, the expression of the wOo MTase gene

was verified by SDS-PAGE. DNA was then extracted using the NucleoSpin Microbial DNA

Mini kit (Macherey Nagel) and subsequently subjected to long reads sequencing using

Sequel I (Pacific biosciences). Assembly of reads was done using the Microbial Assembly

pipeline provided by SMRT Link v.10.1 with a minimum seed coverage of 30x. The

assembled genome was used as a reference in the downstream analysis. The methylation

status of each base of the obtained reads was then determined using the 'Base modification

Analysis' pipeline included in SMRT-Portal. The pipeline performs base modification and

modified base motifs detection. A control experiment was carried out transforming *E. coli*

Stellar TM Competent Cells with a pCOLD III vector without the wOo MTase gene inserted.

The 95th percentile of Mean QV and coverage values obtained from the control experiment

were set as minimum thresholds for the identification of methylated bases on the

transformed *E. coli* strain. These last analyses were performed using R.

Dataset reconstruction and data normalisation

The table S2 and gff map files of the supplementary information of Chung et al. 2020 were

retrieved (26). The table contains the gene expression quantifications computed by Chung

and colleagues (2020) re-analysing seven RNA-Seq studies on *Wolbachia* (16–22), for a

total of 128 samples (replicates) from 62 host conditions. More in details: i) four samples

from two conditions on *Wolbachia* endosymbiont of *Onchocerca ochengi* (wOo) from Darby

et al. 2012(16); ii) five samples from five conditions on *Wolbachia* endosymbiont of

Dirofilaria immitis (wDi) from Luck et al., 2014(17); iii) six samples from two conditions on

Wolbachia endosymbiont of *Drosophila melanogaster* (wMel), from Darby et al., 2014(18);

iv) 12 samples from seven conditions on wDi from Luck et al., 2015(19); v) 55 samples from

24 conditions on wMel, from Gutzwiller et al., 2015(20); vi) 14 samples from seven

conditions on *Wolbachia* endosymbiont of *Brugia malayi* (wBm) from Grote et al., 2017(21);

vii) 32 samples from 15 conditions on wBm from Chung et al., 2019(22). Preliminarily, we

excluded from the dataset all the samples from Luck et al., 2015(19) because only 8% of the

gene expression quantifications were greater than 0. Thus, gene expression analysis was

performed on a total of 116 samples from 55 host conditions, two for wOo, five for wDi, 22

for wBm and 26 for wMel.

Gene expression values were normalised to make them comparable among studies and

conditions. For each condition in each study the gene expression values were expressed as

$$\text{Log2expr} = \sum_s \left[\sum_i \log_2(\text{expr}_{s,i} / \sum_i \text{expr}_{s,i}) \right] / S_{num}$$

Where:

s = sample (or replicate); i = gene; $\text{expr}_{s,i}$ = the gene expression value of the gene i in the

sample (replicate) s; S_{num} = total number of samples. This normalisation formula derives from

the log2 fold change formula often used to compare two conditions in RNA-Seq studies:

$$\text{Log2FC} = \text{Log2expr}_{condition1} - \text{Log2expr}_{conditio21} =$$

$$\{ \sum_s [\sum_i \log_2(\text{expr}_{s,i} / \sum_i \text{expr}_{s,i})] / S \}_{condition1} - \{ \sum_s [\sum_i \log_2(\text{expr}_{s,i} / \sum_i \text{expr}_{s,i})] / S \}_{condition2}$$

Log2FC allows us to compare values from different conditions in the same experiment. To

make the normalised values more comparable among the studies, they were converted in

percentiles. Indeed, the percentile may be less biassed by sample specific features, such as

the number of genes.

Genes composition parameters determination

The genome assemblies of wBm (AE017321.1), wDi (www.nematodes.org: wDi 2.2), wOo

(HE660029.1) and wMel (AE017196.1) were retrieved and passed to Prodigal (27) for Open

Reading Frame (ORF) calling. For each ORF, Codon Bias Index (CBI) (which measure how

much a gene uses a subset of optimal codons, ranging from 0 for random using, to 1 for

maximum bias), the effective number of codons (Nc) and the ORF length were computed

using CodonW tool (Peden J.F. Analysis of codon usage. 1999. PhD Thesis, University of

Nottingham, UK), while %A, %T, %G, %C were computed using an in-house Perl script.

Conventionally, ORF sequences correspond to the transcribed messenger RNA (mRNA)

sequence, which is complementary to the DNA template strand. Thus, the nucleotide

composition of the genes is derived by complementing the ORF compositions: the %A of a

gene is determined as the %T of the corresponding ORF, and so on. Finally, the %C/%A

ratio is calculated. All subsequent analyses will focus on the nucleotide composition of the

genes rather than on ORF compositions.

Statistical analyses

The relationship between each gene composition parameter (CBI, Nc, length, %A, %T, %G,

%C and C/A ratio, see above) and gene expression was investigated. More in detail, for

each of the 55 samples included in the study, linear regression was used to assess the

correlation between the global gene expression reported for that sample and each of the

gene composition parameters. For each regression line, the slope and the p value were

obtained. Slope values indicate the strength of change in gene expression in relation to the

gene composition parameter: the higher the slope, the more the gene expression varies as a

consequence of that gene composition parameter. In a second analysis, to investigate if

MTase activity affects the expression of genes on the basis of their composition parameters,

the correlation between the linear regression slope and the MTase gene expression was

computed for each gene composition parameter.

The presence of other genes with expression patterns comparable to that of MTase gene

was tested as follows. Exploiting the gene orthologous information included in the Table S6

of Chung et al., 2020 (26), we performed a Spearman co-correlation among the expressions

of the 546 single copy core genes shared among wOo, wDi, wBm and wMel. The expression

patterns were also investigated by Principal Component Analysis (PCA), using R.

Comparison between species tree and MTase tree

A dataset including 112 high quality *Wolbachia* genome assemblies spanning the host

genetic diversity was reconstructed, including 111 assemblies retrieved from the Genome

Taxonomy DataBase (GTDB) and the wDi genome used for the previous analysis (indeed,

the wBm, wMel and wOo assemblies were already present in the GTDB database) (see

Table S1). The genome assemblies were subjected to Prodigal (27) for ORF calling, and the

amino acid sequences were passed to OrthoFinder (28) for orthologue analysis. The

nucleotide sequences of the single copy core genes were then retrieved, aligned using the

MUSCLE tool (29), tested for recombinations using the PhiPack tool (30), trimmed using the

trimal tool (-gt 0.5 setting) (31), and finally concatenated. The obtained concatenate was

passed to RAxML8 (32) for phylogenetic analysis using the GTR+I+G model, as previously

determined using ModelTest-NG tool (33). The orthologue group containing the MTase

genes was determined and subjected to phylogenetic analysis following the same flow

described above (using the TVM+I+G model and excluding the step of the concatenation of

the genes). The obtained trees were then compared using the Cophylo R library (34).

Analysis of the region upstream the methyltransferase gene

The positions and orientation of the MTase genes on the 112 genome assemblies were

determined by BLASTn searches. Then, for each genome assembly, the 100 nucleotides

upstream of the MTase gene were extracted and screened for the presence of some of the

most important regulatory regions: i) CtrA binding site (motif TTAA-N7-TTAA (35)); ii)

Pribnow box (motif TATAAT (36)); iii) CAAT box (motif HYYRRCCAWWSR(37)); iv) GC box

(motif WRDRGGHRKDKYYK(37)); v) TATA box (motif TATAWAWR(37)). Perfectly matching

motifs and one-mismatched motifs were considered for further analyses. The presence of

*cck*A and *ctr*A genes in the 112 *Wolbachia* genomes was then evaluated by BlastP search,

using as reference the *cck*A and *ctr*A gene sequences already reported in Lyndsey 2020

(38). Finally, the positions of the canonical regulatory regions upstream the MTase gene,

and the presence/absence of *cck*A/*ctr*A genes in the 112 *Wolbachia* genomes were

visualised using the gplots R library.

Analysis of the regions upstream all genes in the 112 *Wolbachia* genomes

On the basis of previous results on the MTase gene, TATA boxes were searched between

positions -20 and -70 upstream of all the genes of the 112 *Wolbachia* genomes, and CrtA

binding sites between the positions 0 and -30. The presence of short AT-rich sequences, like

TATA box and CtrA binding site, in AT-rich genomes (such as *Wolbachia*) could be due to

the chance. The possible functionality of these sequences has been assessed by

investigating whether TATA or CtrA binding sites are enriched upstream of specific genes in

the *Wolbachia* genomes. More in detail, for each of the ortholog genes previously identified

using OrthoFinder (see above), the frequency of *Wolbachia* strains having upstream TATA

or CtrA binding site was investigated using R. Lastly, both for TATA box and CtrA binding

site, the genes present in at least 100 out of 112 *Wolbachia* genomes and having the

regulatory box in >80% of the genomes were retrieved and annotated using the Clusters of

Orthologous Groups (COG) database.

Analysis of the genes with high and low G/T ratio

One of the gene composition parameters found to be more correlated to gene expression

and MTase action was C/A ratio. Thus, the expression of the genes with higher or lower C/A

ratio is expected to be more affected by the action of the MTase enzyme. To investigate the

effects of this mechanism on the *Wolbachia* metabolism and physiology, genes with low or

high C/A ratio values were studied as follows. All the amino acid sequences from the 112

Wolbachia strains included in the study were annotated on the basis of the Clusters of

Orthologous Genes (COG) database. For each strain, among the COG-annotated genes,

the 10 with lower C/A ratio ("low C/A ratio genes") and the 10 with higher C/A ratio ("high C/A

ratio genes") were retrieved. The pattern of presence/absence and C/A ratio value of all

these retrieved genes was graphically investigated by producing a heatmap using the gplots

R library.

Results

Experimental determination of the wOo SAM-dependent DNA methyltransferase

methylation pattern

The expression of the wOo MTase into the Stellar *E. coli dam*-/*dcm*- strain followed by the

SMRT PacBio sequencing allowed to identify 166 high quality methylated bases: 151 on

cytosine (classified as m4C - N4-methylcytosine) and 15 on adenine (classified as m6A - N6-

methyladenine) (see Table S2 and Figure S1). Investigation of the methylation pattern

revealed that m4c methylations occurred in a pattern that was not well conserved, whereas

m6A methylation involved a guanine-rich region at positions -4, -8 and -19 (Figure 1).

Dataset reconstruction

For each of the six datasets included in the study, the normalised gene expression values

were obtained as described in material and methods. The ORF sequences of the wOo, wDi,

wBm and wMel genome assemblies were analysed to determine the CBI, Nc, length, %A,

%T, %G, %C and %C/%A (from here C/A ratio). The obtained information about gene

composition parameters was merged to relative gene expression data (Supplementary

information available on repository...).

Statistical analyses

The slope and p value of the linear regression analyses performed between the eight gene

composition parameters (CBI, Nc, gene length, %A, %T, %G, %C and C/A ratio) and the

gene expression data obtained for each of the 55 samples are reported in Table S3. Length

and %C significantly correlate with gene expression in all the samples included in the study

(100%), while %T in 53/55 (96%), CBI 42/55 (78%), %A 40/55 (73%), C/A ratio 31/55 (56%),

%G 22/55 (40%) and Nc 7/55 (13%). The obtained slopes for the regression lines range as

follows: for gene length from 0.009 to 0.029, %C from 0.79 to 5.3, CBI from 30.9 to 71.7, %A

from -1.8 to 1.1, C/A ratio from 18.5 to 90.6, %G from -1.3 to 2.2 and Nc from 0.2 to 0.8.

Subsequently, for each of the gene composition parameters (and for each of the 55

samples) the correlation between the levels of expression of the MTase gene and the

regression slope obtained above (indicative of the strength of the correlation between the

gene expression data of the sample and the gene composition parameter) was tested by

linear regression. The aim was to investigate if MTase activity affects the expression of

genes on the basis of their composition parameters. The obtained correlations were

significant for all parameters except %G, which had a p-value of 0.08. For %G, %C, %T, C/A

ratio, Nc, and length, the slope increases with higher DNA MTase expression. Conversely,

%A and CBI exhibit a decreasing trend with increasing of the MTase expression (Figures 2-3

and Figures S2-S7).

Co-correlation analysis among the expression patterns of the single copy core genes shared

among wOo, wDi, wBm and wMel showed that no other gene has an expression pattern

overlapping that of the DNA MTase gene. Indeed, the highest correlation value was 0.84 and

the median value was 0.02 (Figure S8). In general, the absence of clusters of co-expressed

genes is also evident from PCA analysis (Figure S9).

Comparison between species tree and MTase tree

Orthology analysis identified a total of 3,561 orthologous groups, including 263 single copy

core genes. None of the single copy core genes were recombined. After trimming of gene

alignments, the obtained concatenate had a length of 223,833 bp. On the other side, the

MTase gene alignment was 1,342 bp long. The species tree obtained from the 263 single

copy genes and the MTgene tree are reported in Figure S10. The obtained topologies were

mainly congruent, with few signs of Horizontal Gene Transfer (HGT) events.

Analysis of the region upstream the methyltransferase gene

The MTase gene was found in all the 112 genome assemblies included in the study.

Unfortunately, for one genome assembly (accession GCF_018454445.1, *Wolbachia*

endosymbiont of *Rhagoletis cerasi* - wCer5) the MTase gene was located on the extreme of

a contig, and thus it was not possible to extract the 100 bp upstream of the gene

transcription initiation site. The presence of CtrA binding site, Pribnow, CAAT, GC and TATA

boxes upstream the MTase gene was assessed. Among these, at least one canonical

sequence (i.e. perfect match) was found only for CtrA binding site, TATA and Pribnow

boxes. In particular, 104 out of the 111 regions upstream of the MTase gene contained

canonical TATA boxes (Figure S11), 100 out of 111 canonical CtrA binding sites (Figure

S12) and five canonical Pribnow boxes (Figure S13).

Analysis of the regions upstream all genes in the 112 *Wolbachia* genomes

The assessment of the presence of CtrA binding site and TATA boxes upstream of all the genes of the 112 *Wolbachia* genomes included in the study (see above) led to the discovery of a total of 2,210 CtrA binding sites and 12,283 TATA boxes. The CtrA binding site was predominantly found upstream of the MTase gene, but it was also frequently present upstream of the Formate hydrogenlyase subunit 3/Multisubunit Na+/H+ antiporter, which is involved in energy production (see Table S4 and Figure S14). Instead, the TATA boxes were found with high frequency upstream of 12 genes (see Table S4 and Figure S15), including the gene coding for the CtrA DNA binding response regulator.

Analysis of the genes with high and low C/A ratio

A total of 57 low C/A ratio genes and 53 high C/A ratio genes were found. Among the low C/A ratio genes, the most frequent COG categories were "Translation, ribosomal structure and biogenesis" (J) with 13/57 (23%) genes, "Mobilome: prophages, transposons" (X) with 8/57 (14%), "Posttranslational modification, protein turnover, chaperones" (O) with 7/57 (12%) and "Transcription" (K) with 5/57 (9%) genes. Conversely, among the high C/A ratio genes, the most frequent COG categories were "Energy production and conversion" (C) with 15/53 (28%) genes, "Inorganic ion transport and metabolism" (P) with 11/53 (21%), "Lipid transport and metabolism" (I) with 5/53 (9%) genes. The presence/absence and C/A ratio patterns are shown in Figure 4.

Discussion

Wolbachia is one of the most widespread endosymbiotic bacteria (10), with an important impact on the survival and/or reproduction of several arthropod and filarial nematode host species (2, 5, 39)(2, 3, 39). The close symbiotic relationship with the host has had important consequences on the bacterial genome structure, including frequent rearrangements (14) and the loss of many genes and the loss of regulatory regions (40), with the disruption of

several operons and promoters. The loss of promoters could have limited evolutionary costs,

considering that the intracellular niche of the bacterium protects it from external environment

stresses. However, experimental studies showed the existence of *Wolbachia* genes

differentially expressed in host samples from different body compartments, developmental

stages and/or from hosts subjected to environmental conditions (26). Nevertheless, at the

state of the art, the gene regulation mechanisms in *Wolbachia* are poorly understood.

One of the most differentially expressed genes between *Wolbachia* colonising the soma and

gonads of the filarial nematode *Onchocerca ochengi* (wOo) was a SAM-dependent DNA

methyltransferase (MTase) (16). Generally, methylation of bacterial gene promoters is

known to influence the promoters affinity for transcription factors, leading to changes in gene

expression and epigenetic modifications (23, 41). In this work, we investigated the possible

role of the *Wolbachia* MTase as a mediator of gene expression regulation. In absence of

promoters, we hypothesise that intragenic DNA methylation could affect gene expression.

Indeed, experimental studies revealed that the efficiency of bacterial RNA polymerase is

influenced by epigenetic modifications (42–45).

To test our hypothesis, we i) experimentally determined the sequence motif recognised by

the wOo DNA MTase; ii) studied the expression data of 55 samples from six published RNA-

Seq studies on *Wolbachia*; iii) investigated the presence of regulatory regions upstream the

DNA MTase gene in *Wolbachia*.

In silico determination of the sequence motif recognized by MTase is challenging (46). This

is especially true in this case, considering that this is the first *Wolbachia* DNA MTase

enzyme ever characterised. An experimental approach to determining the methylation motif

of the enzyme is to express the MTase in an *Escherichia coli* strain depleted for any MTase

gene, and then identify the methylated bases and motifs using SMRT PacBio sequencing

(46). Surprisingly, wOo MTase was found to be able to methylate both adenine and cytosine,

without highly conserved patterns (Figure 1), especially for cytosine. To our knowledge this

is the first MTase enzyme described with these peculiar features. Our findings also suggest

that the enzyme has a 10-fold greater affinity for cytosine than adenine. This value has been

determined on an *E. coli* genome containing ~50% of AT. Instead, the genome of a

Wolbachia strain usually contains ~70% of AT and this partially reduces the MTase affinity

bias to 4-fold.

Regarding the analysis of gene expression, we started from the idea that if DNA methylation

within the gene affects its expression, we should observe a correlation between gene

nucleotide composition and gene expression. Furthermore, the direction (positive or

negative) and the strength (the slope of the regression line) of this correlation should change

with variation of DNA MTase gene expression. It is well-documented in literature that gene

length and codon usage could be correlated to gene expression (46, 47). In particular, highly

expressed genes tend to have lower Effective Number of Codons (Nc) and higher Codon

Bias Index (CBI). For each of the 55 samples, we computed the correlation between CBI,

Nc, gene length, %A, %T, %G, %C and C/A ratio versus the relative gene expression. The

result shows correlation between gene nucleotide composition and gene expression in many

samples. Interestingly, we found that the slope of the obtained regression lines often

correlates with the MTase expression in that sample. In more detail, the expression of genes

is positively correlated with their %C in all samples and the correlation slope increases

proportionally to the expression of DNA MTase (Figure S2). Regarding %A, the correlation

with gene expression was positive (the higher the %A the higher the gene expression) in

samples with poorly expressed MTase, while it became negative (the higher the %A the

lower the gene expression) in samples with high MTase gene expression (Figure S3).

The positive to negative slope trend observed for %A suggests that methylation of adenine

might play an important role in the regulation of gene expression, in particular considering

the AT-richness of the *Wolbachia* genome. Lastly, we found that the ratio of %C and %A

(C/A ratio) is a good parameter to study *Wolbachia* gene expression: it significantly positively

correlates with gene expression in all samples except for those with very low expression of

MTase (Figure 2). This strongly supports the hypothesis that the methylations of cytosine

and adenine could be the key to understanding the regulation of gene expression in

Wolbachia. Interestingly, as shown in Figure 4, the genes with low C/A ratio are mainly

involved in basal metabolism pathways (such as replication and translation) while those with

high C/A ratio are mostly part of pathways involved in stress response (such as the

production of energy and ion/molecule transport). Interestingly, among the low or high C/A

ratio genes there are, respectively, *fts*Z and *fts*W, both involved in the bacterium cell

replication. This suggests that the host could modulate the bacterium replication: a possible

key for a successful symbiotic relationship. In this picture, *Wolbachia* could have a stress-

related mitochondrion-like role for the host, as previously proposed by Darby and colleagues

(2012). Furthermore, it is interesting to note that highly expressed genes have high Codon

Bias Index (CBI) (as expected from literature) only for samples with low levels of MTase

expression; in those with highly expressed MTase this correlation became not significant

(Figure 3). This suggests that the MTase effect on gene expression might have evolved

independently from the codon usage adaptation.

The step forward in our analysis was to investigate how the expression of the MTase gene

can be regulated. To this aim we analysed a collection of representative *Wolbachia*

genomes, screening the 100-bp region upstream of the MTase gene for the presence of

regulatory sequences. This analysis highlighted the presence of CtrA binding site and TATA

box sequences highly conserved among *Wolbachia* strains. The CtrA binding site is a

sequence bounded by the CtrA response regulator protein, which is part of the two

component regulatory system Ccka/CtrA. This system is composed of Ccka, a membrane

sensor histidine kinase, and CtrA, an intracellular effector protein. When triggered by an

external stimulus, the Ccka protein activates CtrA, which in turn binds to the CtrA binding

site initiating the expression of the downstream genes. While the signalling mechanisms

behind the activation of Ccka are not fully understood, the downstream response regulator,

CtrA, is a well-described master regulator with transcription factor activity, conserved in

Alphaproteobacteria (48). The presence of a highly conserved CtrA binding site upstream

the MTase gene locus, strongly suggests that the CcKA/CtrA signal transduction pathway

could have a role in the regulation of the expression of this gene. Indeed, the host could

modulate the expression of the *Wolbachia* MTase gene by stimulating the Ccka receptor on

the membrane on the bacterial cell. Furthermore, the presence of a conserved CtrA binding

site upstream of only two genes suggests that this could be a highly regulated mechanism

subjected to a strong selective pressure.

Surprisingly, we also found highly conserved TATA boxes (i.e. eukaryotic regulatory

sequences) upstream of the MTase gene. In eukaryotes, TATA boxes tend to be placed

upstream of stress-responsive genes, whose expression must rapidly and variably be tuned

in response to specific environmental conditions and changing physiological needs (49, 50) .

The TATA box is bound by a specific protein, the TATA box binding protein (TBP), which

then recruits transcription factors. *Wolbachia* does not encode TBP proteins, so the

functionality of these TATA box sequences can only be hypothesised, e.g. assuming that

TBPs are supplied by the host. Interestingly, we found 12 *Wolbachia* genes having highly

conserved TATA boxes. These genes include *ctr*A response regulator (see above), genes

involved in energy production and a component of the Type IV secretion system.

To explain how the host could regulate gene expression in *Wolbachia*, we propose the

following model: i) specific molecules in the host cells activate the Ccka membrane histidine

kinase; ii) Ccka phosphorylates the intracellular CtrA response regulator; iii) the

phosphorylated CtrA binds to the CtrA binding site upstream of the MTase gene inducing its

expression in *Wolbachia*; iv) the MTase enzyme methylates adenine and cytosine of the

Wolbachia genome; v) the expression of several *Wolbachia* genes changes on the basis of

their nucleotide content, particularly enhancing the transcription of genes with a high ratio of

cytosine / adenine. The model is graphically summarised in Figure 5.

Moreover, the presence of a conserved TATA box upstream the *ctr*A gene could suggest a

possible role for the host in the regulation of *ctr*A gene expression.

While the hypothesised mechanism may not lead to highly precise regulation of gene

expression, it could be a rough yet functional mechanism, sufficient for the establishment of

a successful intracellular symbiosis. In the future, it would be interesting to investigate

whether similar mechanisms are present in other endosymbiont or intracellular organisms.

Despite the results presented in this work are far from being definitive, we hope to have

opened a new way in understanding the symbiotic relationship between *Wolbachia* and its

hosts. In our opinion, this is a magnificent example of how evolution works, recycling

metabolic pieces for similar aims, sometimes reaching surprisingly stable *equilibria*.

Captions

Figure 1. Methylation patterns of the DNA methyltransferase gene of *Wolbachia* endosymbiont of *Onchocerca ochengi* (wOo)

Sequence logo of the sequences flanking (from -20 to +20 positions) the methylated positions found in the genome of Stellar *Escherichia coli* dam-/dcm- strain expressing the DNA methyltransferase gene of *Wolbachia* endosymbiont of *Onchocerca ochengi* (wOo). The height of the letters depends on the conservation of the position among the sequences. a) the sequence logo of the m4C methylations on cytosines; b) the m6A methylation of adenine.

Figure 2. Correlation between the expression of genes and their %C / %A ratio

For each of the 55 samples included in the study, the correlation between the expression of the genes and the relative %C/%A ("C/A ratio") was investigated. The scatter plots of C/A ratio vs gene expression of three exemplificative samples selected from the 55 samples are reported: a) the correlation plot with highest slope (Hs), b) the correlation plot with the medium slope (Ms) and c) the correlation plot with the lowest slope (Ls). In each of these three plots, colour is green whether the correlation is significant (p value < 0.05) and grey if it is not, the regression line is coloured in dark grey and the statistics are reported on the chart. In figure d, the correlation plot between the slopes obtained for each of the 55 samples, and the relative expression of the DNA methyltransferase gene is reported. Points relative to significant correlations (p value < 0.05) are in green while non significant in grey. The regression line is in green and the statistics are reported on the chart.

Figure 3. Correlation between the expression of genes and their Codon Bias Index (CBI)

For each of the 55 samples included in the study, the correlation between the expression of the genes and the relative Codon Bias Index (CBI) was investigated. The scatter plots of CBI

vs gene expression of three exemplificative samples selected from the 55 samples are

reported: a) the correlation plot with highest slope (Hs), b) the correlation plot with the

medium slope (Ms) and c) the correlation plot with the lowest slope (Ls). In each of these

three plots, colour is green whether the correlation is significant (p value < 0.05) and grey if it

is not, the regression line is coloured in dark grey and the statistics are reported on the chart.

In figure d, the correlation plot between the slopes obtained for each of the 55 samples, and

the relative expression of the DNA methyltransferase gene is reported. Points relative to

significant correlations (p value < 0.05) are in green while non significant in grey. The

regression line is in green and the statistics are reported on the chart.

Figure 4. COG annotated genes with highest or lowest C/A ratio

The heatmap shows the pattern of the genes with high or low C/A ratio among the 112

Wolbachia strains included in the study. On the left, the Maximum Likelihood phylogenetic

tree obtained from concatenate of the single copy core genes. In the middle, heatmap

reports the pattern of the 10 lower C/A ratio COG annotated genes: rows refer to strains

(matching with the phylogenetic tree on the left) and columns to genes (COG annotation is

reported). Cell colours report the C/A ratio value following the legend on lower left. Asterisks

are reported for genes among those in the 10 lower C/A ratio for the relative strain. More on

the right, a heatmap reports the 10 higher C/A ratio COG annotated genes. Finally, on the

right, *Wolbachia* strains hosts and supergroups are shown.

Figure 5. Graphical representation of the mechanism proposed for the regulation of

the *Wolbachia* gene expression

The figure summarises the model here proposed for the *Wolbachia pipientis* gene

expression regulation. *Wolbachia* is an intracellular bacterium able to colonise the cytoplasm

of host cells. In the figure the bacterium, represented with violet membranes and white

cytoplasm, is surrounded by the host cytoplasm, coloured in green. At step 1 (on the left):

the host activates the Ccka sensor histidine kinase (in violet) placed on the *Wolbachia*

membrane (here we represented this activation as mediated by a molecular factor produced by the host, but other mechanisms are possible), then the Ccka phosphorilates the intracellular transcription regulator CtrA which, induces the transcription of the *Wolbachia* DNA *methyltransferase* gene (MTase) binding the CtrA binding site upstream it. At step 2 (in the middle): the MTase enzyme methylates the adenines and cytosines of the *Wolbachia* DNA and methyl groups are represented with dots coloured as the bases. At step 3 (on the right): RNA polymerases are represented by pink icons from which mRNA molecules originate. In highly expressed genes more RNA polymerases bind the gene simultaneously. The greater is the cytosines / adenines ratio in a gene the more that gene is expressed.

Acknowledgments

F.C. would like to thank his mentor, Claudio Bandi, for introducing him to the world of *Wolbachia* and for his helpful comments to this paper.

Funding

The study was supported by the National Institute of Virology and Bacteriology (Program EXCELES, ID project no. LX22NPO5103) funded by the European Union–Next Generation EU.

References

1.   R. Zug, P. Hammerstein, Still a host of hosts for Wolbachia: analysis of recent data
suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7**, e38544
(2012).

2.   J. Porter, W. Sullivan, The cellular lives of Wolbachia. *Nat. Rev. Microbiol.* **21**, 750–766
(2023).

3.   J. H. Werren, Biology of Wolbachia. *Annu. Rev. Entomol.* **42**, 587–609 (1997).

4.   C. Bandi, T. J. Anderson, C. Genchi, M. L. Blaxter, Phylogeny of Wolbachia in filarial
nematodes. *Proc. Biol. Sci.* **265**, 2407–2413 (1998).

5.   J. H. Werren, L. Baldo, M. E. Clark, Wolbachia: master manipulators of invertebrate
biology. *Nat. Rev. Microbiol.* **6**, 741–751 (2008).

6.   L. M. Hedges, J. C. Brownlie, S. L. O'Neill, K. N. Johnson, Wolbachia and virus
protection in insects. *Science* **322**, 702 (2008).

7.   L. Teixeira, A. Ferreira, M. Ashburner, The bacterial symbiont Wolbachia induces
resistance to RNA viral infections in Drosophila melanogaster. *PLoS Biol.* **6**, e2 (2008).

8.   I. L. G. Newton, D. W. Rice, The Jekyll and Hyde Symbiont: Could Be a Nutritional
Mutualist? *J. Bacteriol.* **202** (2020).

9.   C. Bandi, *et al.*, Effects of tetracycline on the filarial worms Brugia pahangi and
Dirofilaria immitis and their bacterial endosymbionts Wolbachia. *Int. J. Parasitol.* **29**,
357–364 (1999).

10.   M. Scholz, *et al.*, Large scale genome reconstructions illuminate Wolbachia evolution.
Nat. Commun. **11**, 5235 (2020).

11.   S. J. A. Tolley, P. Nonacs, P. Sapountzis, Horizontal Transmission Events in Ants: What
Do We Know and What Can We Learn? *Front. Microbiol.* **10**, 296 (2019).

12.   R. D. Olson, *et al.*, Introducing the Bacterial and Viral Bioinformatics Resource Center
(BV-BRC): a resource combining PATRIC, IRD and ViPR. *Nucleic Acids Res.* **51**,
D678–D689 (2023).

13.   A. Manzano-Marín, A. Latorre, Snapshots of a shrinking partner: Genome reduction in
Serratia symbiotica. *Sci. Rep.* **6**, 32590 (2016).

14.   F. Comandatore, *et al.*, Supergroup C Wolbachia, mutualist symbionts of filarial
nematodes, have a distinct genome structure. *Open Biol.* **5**, 150099 (2015).

15.   S. Mahmood, E. Nováková, J. Martinů, O. Sychra, V. Hypša, Supergroup F Wolbachia
with extremely reduced genome: transition to obligate insect symbionts. *Microbiome* **11**,
22 (2023).

16.   A. C. Darby, *et al.*, Analysis of gene expression from the Wolbachia genome of a filarial
nematode supports both metabolic and defensive roles within the symbiosis. *Genome*
Res. **22**, 2467–2477 (2012).

17. A. N. Luck, *et al.*, Concurrent transcriptional profiling of Dirofilaria immitis and its
Wolbachia endosymbiont throughout the nematode life cycle reveals coordinated gene
expression. *BMC Genomics* **15**, 1041 (2014).

18. A. C. Darby, *et al.*, Integrated transcriptomic and proteomic analysis of the global
response of Wolbachia to doxycycline-induced stress. *ISME J.* **8**, 925–937 (2014).

19. A. N. Luck, *et al.*, Tissue-specific transcriptomics and proteomics of a filarial nematode
and its Wolbachia endosymbiont. *BMC Genomics* **16**, 920 (2015).

20. F. Gutzwiller, *et al.*, Dynamics of Wolbachia pipientis Gene Expression Across the
Drosophila melanogaster Life Cycle. *G3* **5**, 2843–2856 (2015).

21. A. Grote, *et al.*, Defining Brugia malayi and Wolbachia symbiosis by stage-specific dual
RNA-seq. *PLoS Negl. Trop. Dis.* **11**, e0005357 (2017).

22. M. Chung, *et al.*, Drug Repurposing of Bromodomain Inhibitors as Potential Novel
Therapeutic Leads for Lymphatic Filariasis Guided by Multispecies Transcriptomics.
mSystems **4** (2019).

23. S. Papaleo, *et al.*, The red thread between methylation and mutation in bacterial
antibiotic resistance: How third-generation sequencing can help to unravel this
relationship. *Front. Microbiol.* **13**, 957901 (2022).

24. M. A. Sánchez-Romero, J. Casadesús, The bacterial epigenome. *Nat. Rev. Microbiol.*
18, 7–20 (2020).

25. S. S. Mohapatra, A. Fioravanti, E. G. Biondi, DNA methylation in Caulobacter and other
Alphaproteobacteria during cell cycle progression. *Trends Microbiol.* **22**, 528–535
(2014).

26. M. Chung, *et al.*, A Meta-Analysis of Transcriptomics Reveals a Stage-Specific
Transcriptional Response Shared Across Different Hosts. *G3* **10**, 3243–3260 (2020).

27. D. Hyatt, *et al.*, Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 119 (2010).

28. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

29. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

30. T. C. Bruen, H. Philippe, D. Bryant, A simple and robust statistical test for detecting the
presence of recombination. *Genetics* **172**, 2665–2681 (2006).

31. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).

32. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

33. D. Darriba, *et al.*, ModelTest-NG: A New and Scalable Tool for the Selection of DNA
and Protein Evolutionary Models. *Mol. Biol. Evol.* **37**, 291–294 (2020).

34. L. J. Revell, phytools 2.0: an updated R ecosystem for phylogenetic comparative
methods (and other things). *PeerJ* **12**, e16505 (2024).

35. M. C. Ouimet, G. T. Marczynski, Analysis of a cell-cycle promoter bound by a response
regulator. *J. Mol. Biol.* **302**, 761–775 (2000).

36. D. Pribnow, Nucleotide sequence of an RNA polymerase binding site at an early T7
promoter. *Proc. Natl. Acad. Sci. U. S. A.* **72**, 784–788 (1975).

37. P. Bucher, Weight matrix descriptions of four eukaryotic RNA polymerase II promoter
elements derived from 502 unrelated promoter sequences. *J. Mol. Biol.* **212**, 563–578
(1990).

38. A. R. I. Lindsey, Sensing, Signaling, and Secretion: A Review and Analysis of Systems
for Regulating Host Interaction in Wolbachia. *Genes* **11**, 813 (2020).

39. F. Comandatore, *et al.*, Phylogenomics and analysis of shared genes suggest a single
transition to mutualism in Wolbachia of nematodes. *Genome Biol. Evol.* **5**, 1668–1674
(2013).

40. C. Dale, N. A. Moran, Molecular Interactions between Bacterial Symbionts and Their
Hosts. *Cell* **126**, 453–465 (2006).

41. Q. Gao, *et al.*, Bacterial DNA methyltransferase: A key to the epigenetic world with
lessons learned from proteobacteria. *Front. Microbiol.* **14**, 1129437 (2023).

42. M. R. Rountree, E. U. Selker, DNA methylation inhibits elongation but not initiation of
transcription in Neurospora crassa. *Genes Dev.* **11**, 2383–2395 (1997).

43. F. Gracias, *et al.*, Homologues of epigenetic pyrimidines: 5-alkyl-, 5-hydroxyalkyl and 5-
acyluracil and -cytosine nucleotides: synthesis, enzymatic incorporation into DNA and
effect on transcription with bacterial RNA polymerase. *RSC Chem Biol* **3**, 1069–1075
(2022).

44. C. Rausch, *et al.*, Cytosine base modifications regulate DNA duplex stability and
metabolism. *Nucleic Acids Res.* **49**, 12870–12894 (2021).

45. M. Janoušková, *et al.*, 5-(Hydroxymethyl)uracil and -cytosine as potential epigenetic
marks enhancing or inhibiting transcription with bacterial RNA polymerase. *Chem.*
Commun. **53**, 13253–13255 (2017).

46. T. Ø. Jensen, *et al.,* Genome-wide systematic identification of methyltransferase
recognition and modification patterns. *Nat. Commun.* **10**, 3311 (2019).

47. T. E. F. Quax, N. J. Claassens, D. Söll, J. van der Oost, Codon Bias as a Means to
Fine-Tune Gene Expression. *Mol. Cell* **59**, 149–161 (2015).

48. S. Narayanan, L. Kumar, S. K. Radhakrishnan, Sensory domain of the cell cycle kinase
CckA regulates the differential DNA binding of the master regulator CtrA in Caulobacter
crescentus. *Biochim. Biophys. Acta Gene Regul. Mech.* **1861**, 952–961 (2018).

49. A. D. Basehoar, S. J. Zanton, B. F. Pugh, Identification and distinct regulation of yeast
TATA box-containing genes. *Cell* **116**, 699–709 (2004).

50. S.-H. Bae, H. W. Han, J. Moon, Functional analysis of the molecular interactions of
TATA box-containing genes and essential genes. *PLoS One* **10**, e0120848 (2015).

Lowest C/A

Highest C/A

G/T ratio

1. MTase EXPRESSION: The host induces the MTase expression

2. DNA METHYLATION: The MTase enzyme methylates adenine and cytosines

3. GENE REGULATION: Wolbachia genes with high C/A are more expressed

Host factor

CckA

P

CtrA

P

MTase

CtrA binding site

MTase enzyme

WOLBACHIA CELL

HOST CELL

MTase enzyme

A T G G A T C G C T A G A T C G
T A C C T A G C G A T C T A G C

● = Methylated Adenine

● = Methylated Cytosine

WOLBACHIA CELL

RNA polymerase

T A C C T C G C G C G C T A G C

T A A C T A G A G A G A T C G C

WOLBACHIA CELL