Distinct routes to thermotolerance in the fungal pathogen *Cryptococcus neoformans*

Mara J. W. Schwiesow[1,2†], Leah A. Farinella[3], Marina Ruzic[3], Jake T. Leinas[3], Nels C. Elde[1,2], Zoë A. Hilbert[1-3*]

[1]Department of Human Genetics, University of Utah, Salt Lake City, UT 84112, USA
[2] Howard Hughes Medical Institute, Salt Lake City, UT, USA
[3]Department of Biology, Boston College, Chestnut Hill, MA 02467, USA
[†]Current address: Department of Molecular and Cellular Biology, University of California at Merced, Merced, CA, USA

*Correspondence: hilbertz@bc.edu

Abstract

15
Increasing temperatures associated with climate change have the potential for far-
reaching impact on human health and disease vectors, including fungal pathogens. Pathogenic
fungi occupy a wide range of environments across the world, and their ranges have been slowly
expanding in recent decades due, in part, to climate change. Despite these links between
increasing temperature and higher prevalence of fungal disease, the direct effects of rising
environmental temperatures on the evolution of pathogenic fungi remains unclear. In this study,
we investigated how increasing temperatures drive adaptive evolution in the human fungal
pathogen *Cryptococcus neoformans*. First, we performed serial passages of a *C. neoformans*
environmental isolate with gradual changes in temperature over the course of 38 days. Through
this approach we identified several distinct thermally adapted isolates with competitive growth
advantages over the parental strain at high temperatures. We then characterized the phenotypic
and genetic changes acquired in these evolved isolates, which include alteration of cell size,
colony morphology, and, notably, antifungal resistance. Our genetic analyses further revealed
distinct genes that facilitate thermoadaptation in different populations—identifying new
molecular players in the regulation of this trait and revealing that there are multiple independent
routes to gaining thermotolerance. These results highlight the remarkable flexibility of fungi to
adapt rapidly to new environments and raise pressing questions about the impacts of rising
environmental temperatures on the future of infectious diseases and human health.
34

Introduction

The impacts of planetary warming on biological systems hold pivotal implications for human and animal health worldwide[1]. Climate change is likely to have particularly dramatic effects in the realm of infectious disease, where increased environmental temperatures could fundamentally alter the number, nature, and geographic distribution of pathogenic microbes, including fungi[2]. However, little work has been done to investigate how incrementally increasing temperatures will affect the evolution of these pathogens and what consequences this adaptation may have in the context of human infection.

For fungi, in particular, the inability of most species to grow at the high body temperatures characteristic of mammals has long been considered a primary limiting factor in the ability of these microbes to establish infection[3]. As global temperatures rise, however, adaptation to higher temperatures is expected to drive increases in the number of fungal species capable of surviving at mammalian body temperature, perhaps even driving the emergence of new species with pathogenic potential[4]. Increased environmental temperature will also result in range expansions for pathogenic fungi. Indeed, this has already been observed in several fungal species including *Coccidioides* spp., *Cryptococcus gattii*, and *Candida auris*, with *C. auris* emerging on three separate continents simultaneously[5–7]. Although changes in the geographic range for these particular species are in part due to expansion of their preferred environments, there is also evidence to suggest that these organisms are adapting to new, more stressful environmental conditions[8,9]. Given these trends, it can be expected that as global temperatures continue to rise, the ranges of these and other pathogenic fungi will continue to expand, the threat of thermal adaptation in these organisms will escalate, and human body temperature may no longer pose such an impediment to fungal infection.

In this study, we focus on the role of increasing environmental temperatures in shaping the evolution of the common human fungal pathogen, *Cryptococcus neoformans*. *C. neoformans* is a pathogenic basidiomycete fungus found ubiquitously in the environment, usually residing in bird guano, soil, and trees, and infection by this species is currently estimated to cause over 100,000 deaths annually[10,11]. Strains of *C. neoformans* occupy a wide range of temperature environments having been isolated from all continents, excluding Antarctica, both from environmental and clinical settings[12–16]. Despite this, our understanding of how thermotolerance evolved in this species and may vary across strains is relatively limited.

Some studies of laboratory reference isolates have revealed candidate genes and pathways in *C. neoformans* that are important for success at high temperature. For example, heat shock proteins, and certain other conserved genetic pathways, such as *CDC11*, calcineurin, and the HOG pathway, all play roles in adaptation to the host environment, including adaptation to host temperature[17–21]. However, these studies largely focus on gene knockouts of the laboratory reference strain and sidestep a more realistic view of how this organism might adapt by modulating expression or function of these or other genes to survive under changing conditions.

Experimental evolution is a powerful tool to investigate the processes and mechanisms of adaptation to new environments in wide-ranging species. Studies of thermal adaptation, in particular, include the pioneering work of Bennett and Lenski, who used long-term experimental evolution to study the adaptive changes that occur in the bacterial species *Escherichia coli* under high temperature stress[22]. In fungal species, similar serial passaging approaches at consistent, high temperatures have proven powerful in generating heat-tolerant strains for use in industrial settings[23]. Additionally, two studies in the yeasts *Saccharomyces cerevisiae* and *Saccharomyces uvarum* have shown that gradual and stepwise temperature ramps can increase the thermotolerance of fungi by 1.5°C or more, suggesting that these fungi have a remarkable plasticity for tolerating temperature fluctuations[24,25]. However, studies exploring the thermal adaptation capabilities of environmental pathogenic fungi and related impacts on pathogenicity have been largely missing.

In this study, we leverage experimental evolution approaches to study the process of
thermal adaptation in the clinically relevant *C. neoformans*. Through gradually increasing
temperatures over many generations, we found that not only can *C. neoformans* adapt to
produce genetically distinct strains with increased thermotolerance, but that this adaptation
occurs on relatively short timescales. In only 38 days of continuous culturing, we observed
increased thermotolerance of one environmental strain of *C. neoformans* characterized by
improved growth at high temperature with clear fitness advantages over the parental strain.
Thermal adaptation occurred across a range of media types and co-evolved with other
phenotypic changes in cell size, colony morphology, and antifungal resistance. We further
identify two genes, not previously known to regulate temperature sensitivity, which acquired
mutations in distinct evolved populations and contribute to improved growth at high temperature.
Our findings indicate that *C. neoformans* is capable of adapting to increasing global
temperatures and provide a foundation on which to build an understanding of how our changing
climate could impact human health and fungal disease.
100
Results
Variable thermotolerance across environmental isolates of *C. neoformans*
Environmental strains of *C. neoformans* inhabit diverse locations and are exposed to a
variety of temperature ranges with varying nutrient availability. Although environmental isolates
may generally experience more temperate climates, they are also likely subjected to wider
temperature swings in contrast to clinical isolates, which are better able to grow at the high
temperatures (37°C and above) of the human body. To test the thermotolerance properties of a
panel of environmental *C. neoformans* isolates, we performed spot dilution assays and
assessed the growth of these strains at two different temperatures: the standard laboratory
culture temperature of 30°C and at a high temperature of 37°C, representative of human body
temperature. To assess how thermotolerance might differ across different nutritional
environments, we additionally assayed strains under rich (Yeast extract Peptone Dextrose
[YPD]) and minimal (Yeast Nitrogen Base with glucose [YNB]) nutrient conditions. These assays
revealed that thermotolerance varies widely, even between closely related strains (Figure 1).
Some strains, such as Ftc217-1 and Jo278-1, grow well at high temperature, similar to the
laboratory reference strain, H99 (Figure 1b and c). Others, like Muc387-1, exhibited moderate
growth defects at high temperature. In general, growth differences at both temperatures were
more pronounced on the minimal YNB media than on YPD.
Among the set of strains tested here, one environmental isolate stood out: Ftc239-1.
This strain, which was isolated from a mopane tree in Botswana, is closely related to Ftc217-1,
but displays strikingly different growth across temperatures and nutrient conditions[14,15]. Unlike
Ftc217-1, Ftc239-1 has severely attenuated growth at 37°C on both YNB and YPD. This strain
also displays decreased growth at 30°C on YNB media (Figure 1b). The difference in growth on
these two media types, across all strains tested but particularly in Ftc239-1, suggests that
thermotolerance and nutrient conditions interface to control fungal growth.
Given the severe growth defects of Ftc239-1 at high temperatures, we reasoned that this
would be an ideal strain to use to explore the mechanisms by which thermotolerance evolves in
C. neoformans. In particular, the phenotypes of this strain gave us an opportunity to explore how
it might adapt to higher temperatures relevant for human infection, while also providing an
avenue to investigate the role of nutrient conditions in thermal adaptation.
131
132   **Experimental evolution can drive fungal adaptation to increasing temperatures**
133   We used laboratory-based experimental evolution to study the process of thermal
134   adaptation in the Ftc239-1 *C. neoformans* strain. Although temperature increases associated
135   with human-induced climate change are rapid for geologic time, they are much more gradual for
136   the organisms experiencing those changes. As such, we designed our experimental evolution

paradigm to mimic the gradual temperature increases environmental pathogens may experience in the coming decades, though over a greatly abbreviated timescale. Specifically, Ftc239-1 was serially passaged every 48 hours starting at 30°C with temperature increases of 0.5–1°C every 4–6 days until a temperature of 35°C was reached (Figure 2a). In total, the fungal cells were grown continuously in this paradigm for 19 passages (38 days), which we estimate represents <150 generations of growth—a relatively short course of evolution. From the originating Ftc239-1 strain, eight populations were evolved in parallel, with four passaged in a rich media environment (YPD) and four passaged under nutrient-limiting conditions (YNB). Given the differences we observed in growth across these two media types between strains (Figure 1), along with previous work highlighting the role of several metabolic pathways in thermal sensitivity, we reasoned that the differing nutrient conditions would reveal distinct evolutionary paths[26,27]. These independently evolving Thermal Adaptation populations were named TA1-4 for the populations evolved in YPD and TA5-8 for the YNB-passaged replicates (Figure 2a).

All of the populations were viable until the end of the evolution experiment (35°C), indicating that this *C. neoformans* strain has the ability to adapt to or at least withstand increased thermal stress. Following the end of passaging, we harvested and froze the evolved populations; we also recovered individual clonal isolates for further characterization (see Methods). Intriguingly, we observed variation in the appearances and morphology of the clonal samples at the end of the evolution. Some displayed a mucoid appearance, while others had increased colony size compared to the starting isolate, providing our first indication that our experimental evolution had yielded new and potentially adaptive phenotypes in these populations (Tables S1 and S2). Further, changes in these phenotypes were not always universal within populations, suggesting that there was both phenotypic and genetic heterogeneity at the end of our course of evolution.

Experimentally evolved isolates grow better on solid media at high temperatures

Following passaging, we first set out to determine whether any of the individual endpoint colonies displayed increased growth at high temperature compared to the parental strain. We, again, employed spot dilution assays to assess the growth phenotypes of our 36 recovered clonal isolates (Figure S1). This revealed that every independently evolved population contained fungi with enhanced growth at 35°C compared to the parental. These spot assays also confirmed our previous observations that there was phenotypic heterogeneity across isolates both between populations and within a single population. This was particularly clear among YPD-evolved isolates from populations TA1-TA4 where individual isolates varied both in ability to grow at high temperature (compare TA2-3 and 2-5 to other TA2 isolates) and in ability to grow on YNB minimal media (see isolates TA3-1, TA3-4, and all isolates from the TA4 population). These data indicate that there are multiple distinct paths to gaining thermotolerance in this *C. neoformans* strain.

From these experiments, we selected individual isolates from two independent populations for further characterization and follow-up (Figure 2b and c). One of these isolates was evolved in YPD (TA1-1) and one was evolved in YNB (TA8-2). At 30°C, both TA1-1 and TA8-2 had increased growth on YNB, while maintaining similar growth on YPD to the parental strain (Figure 2b and c, inner panels). At 35°C, both evolved isolates grew one or two dilutions further than the parental strain. This effect was consistent on both media types suggesting that thermotolerance in these particular isolates confers an advantage across nutrient conditions.

We next tested whether the increased thermotolerance in TA1-1 and TA8-2 conferred an enhanced ability to grow at high temperatures beyond the range of our evolution and whether there are tradeoffs for increased thermotolerance, perhaps affecting growth at temperatures below 30°C. Thermal adaptation studies with other microbial species, like *E. coli*, have shown that such paradigms can select for evolved isolates with highly specific temperature preferences i.e. optimized for growth at the temperatures using during the experiment[22]. To assess the

growth phenotypes of our strains outside the bounds of the experimental evolution, we looked at growth of our selected isolates in parallel at 26, 30, 35 and 37°C. Both thermally-adapted isolates showed increased growth at 37°C on both YPD and YNB medias (Figure 2b and c, outer panels). There was no substantial difference in growth from the parental strain for either isolate at 26°C on YPD, however there was a slight increase in growth of the evolved strains on YNB at this temperature. These results suggest that the gradual adaptation of this *C. neoformans* strain to a specific high temperature (35°C) confers a growth advantage in high temperatures generally, with no apparent tradeoffs for growth at lower temperatures.

Thermally adapted strains outcompete the parental strain at 35°C

To directly test whether the evolved strains gained a fitness advantage for growth at high temperature, we conducted co-culture competition experiments. In these competition experiments, evolved strains were grown together with a drug-resistant version of the parental strain in the same liquid media in which they were originally evolved (Figure 3a).

These experiments confirmed that each of the evolved strains has a substantial competitive advantage over the parental strain at 35°C. TA1-1 grew over five times better than the parental strain, and TA8-2 had a three to four-fold advantage in the media conditions used during the evolution experiment (Figure 3b and c). While this fitness effect is less robust at 30°C, the competitive advantages of the evolved strains were maintained at lower temperatures as well. These data indicate that the evolved isolates have successfully adapted to be able to grow at high temperatures (35°C) and further reveal, that these isolates have a fitness advantage over the parental strain not only at high temperature but across a range of temperatures.

Evolved isolates have altered colony morphologies and titan cell production

We next asked whether thermal adaptation in our evolved strains was accompanied by alteration of other physical or morphological properties of the fungi. High temperature is a known regulator of different aspects of cryptococcal biology, including pathogenicity-related phenotypes such as capsule production, cell size, and melanization, among others[28]. Investigation of colony morphology revealed differences in our harvested clonal samples. The parental strain, Ftc239-1, has smooth, small colonies that are cream-white in color. The evolved strains (TA1-1 and TA8-2) exhibited changes in one or more of these features. TA1-1 colonies were consistently larger than the parental colonies and took on a mucoid appearance (Figure 2b, Table S1). These mucoid colonies have a glossy appearance with smooth edges and often form a looser pellet when spun down in liquid culture[29]. TA8-2 maintained the smooth colony phenotype seen in the parental strain, but the colonies were markedly larger (Table S2). Interestingly, more than half of the remaining endpoint colonies appeared mucoid, which suggests that the mucoid phenotype confers some advantage for surviving thermal stress or that there may be overlap in the genetic pathways that control these two traits (Tables S1 and S2). We hypothesized that changes in mucoidy might indicate a perturbation of the cell surface of these evolved cells, including production of the polysaccharide capsule that serves as a protective coating on cryptococcal cells under host-like conditions. However, our preliminary analysis of capsule production in our selected clonal isolates showed no major differences in capsule size (Figures S2a and b). This suggests that the mucoidy we observe in isolates across our evolved populations is likely derived from other altered features of our thermoadapted strains.

Given that colony morphology varied across our evolved isolates, we were curious what other aspects of cell morphology might be altered in these strains. To this end, we looked at the ability of our evolved isolates to form titan cells: large (>10 μm), polyploid cells that are induced during *in vivo* infection and *in vitro* under host-like conditions, including elevated temperatures[30]. Larger cell sizes inhibit phagocytosis by immune cells and a polyploid genome enables

increased mutational flexibility, both of which allow the yeast to survive and adapt to host
conditions during infection. Using an *in vitro* titanization assay, we observed substantial changes
in titan cell formation in our selected thermoadapted isolates. Although TA1-1 showed robust
titan cell formation, the TA8-2 strain failed to titanize entirely (Figures S2a and c). A second
clonal isolate from the TA8 population, TA8-3, similarly failed to titanize under *in vitro* inducing
conditions, suggesting that thermal adaptation in the TA8 population was accompanied by a
cost for cell size variation (Figure S2C).
The opposing titanization phenotypes of TA1-1 and TA8-2 led us to ask whether titan cell
formation was preserved in our other thermally adapted populations and if there might be a
correlation between alterations in this phenotype and the media conditions each population
experienced during the evolution. To assess this, we repeated our titan cell assays with 12
additional endpoint colonies with representatives from every independent population (Figure
4a). The majority of the isolates we examined in these experiments showed increased growth at
high temperature compared to the parental starting strain (Figure S1). However, two isolates,
TA2-5 (from YPD-evolved population TA2) and TA6-2 (from YNB-evolved population TA6)
display parental-like temperature sensitive growth phenotypes and were included as important
reference points for correlating cell morphology phenotypes with thermoadaptation.
Although there was variation in the percentage of cells that exceeded the 10 μm
threshold, we observed that the vast majority of the isolates from our evolved populations
maintained the ability to titanize (Figure 4a). Besides TA8-2, only one other isolate failed to
produce cells larger than 10 μm: the YPD-evolved TA2-4. Interestingly, the TA2-5 isolate from
the same population, which shows impaired growth at high temperature, was still able to form
titan cells, suggesting that the lack of titanization in TA2-4 (and other strains) may be a direct
tradeoff caused by high temperature adaptation. Collectively, our investigations of cell
morphology across our evolved isolates showt that these features are variably altered across
and within independently evolved populations, regardless of media type, and include loss of key
virulence-related traits like titanization.
266
Antifungal drug susceptibility is altered in thermally adapted strains
In addition to thermal stress, fungal cells are subjected to a wide range of other stressors
both in the environment and during infection, prompting us to ask whether our evolved isolates
may also be more resistant to other types of cellular stress. We decided to explore this question
using exposure to the antifungal drug fluconazole as a clinically relevant test to investigate the
interface between thermal adaptation and other possible stressors.
In preliminary experiments assessing the minimum inhibitory concentrations (MICs) of
our strains, we observed that our selected clonal isolates showed opposite trends in their MICs
compared to the parental strain (Figure S2d). TA1-1 displayed enhanced fluconazole resistance
compared to the parental strain and TA8-2 showing decreased resistance (Figure S2d). To gain
a more comprehensive view of antifungal resistance in our thermally adapted strains, we
assessed the MICs across our wider panel of isolates (Figure 4b). Strikingly, we observed that
many strains displayed increased MICs compared to the parental Ftc239-1 strain, with a few
notable strains (e.g. TA3-5) showing robust resistance to fluconazole (Figure 4b and Table 1).
Only a single isolate from the evolution experiment, TA8-2, had a markedly decreased MIC in
these experiments.
To gain a more nuanced view of fluconazole's effects on growth in these isolates, we
used spot dilution assays to observe growth on fixed concentrations of fluconazole (Figure 4c
and Figure S3). We observed very few changes in growth for most strains on plates containing
a low (4 μg/mL) dose of fluconazole; in contrast, high concentrations (12 μg/mL) of fluconazole
triggered a much more variable growth inhibition. Although the parental strain and several
others display marked decreases in growth when exposed to this concentration of fluconazole,
many of the thermally adapted isolates continued to grow robustly showing that the increased

MIC values are indeed indicative of increased antifungal resistance. This includes the isolates:
TA1-1, TA3-5, TA5-1, TA6-3, and both isolates from the TA7 population.TA8-2, in contrast was
consistently among the worst performers in these fluconazole assays, aligning with its lower
MIC values. Interestingly, we observed that our isolates that remained temperature sensitive
after passaging (TA2-5 and TA6-2) showed similar sensitivity to fluconazole as the parental
strain, suggesting that drug resistance correlates with thermoadaptation in many of our
experimentally evolved populations.
297
Evolved isolates have distinct mutations that vary between populations and drive the
emergence of thermotolerance
Given the strong growth advantage of our passaged strains along with the phenotypic
differences across these isolates, we hypothesized that they likely carried adaptive genetic
lesions that underlie their phenotypes. To begin to assess the genetic basis of adaptation in our
thermally-adapted isolates, we returned our focus to TA1-1 and TA8-2, which differ in most traits
beyond thermoadaptation, suggesting that they may have taken different evolutionary and
genetic routes to adapt to high temperature. We performed whole-genome sequencing of our
parental isolate, TA1-1, and TA8-2, along with additional isolates from the TA8 population.
We observed unique mutations in colonies from the same population (TA8-2 vs. other
TA8 isolates) as well as across populations (YPD vs YNB, TA1-1 vs TA8-2). In the YPD-
passaged TA1-1, we identified four fixed mutations affecting a range of different genes and with
varying effects, from intronic mutations to frameshifting insertions (Table 2). In the TA8 YNB-
evolved population, we identified six mutations that were shared across all clonal isolates,
including TA8-2 (Table 3). These mutations were also at high frequency (>90%) in sequencing
data from the pooled population culture indicating they were fixed in the TA8 population. In
addition to the six mutations that were shared, we also identified several additional mutations
that were present in some—but not all—clonal isolates in the TA8 population (Table 4). Notably,
TA8-2 had no mutations beyond those that were fixed in the population. As with TA1-1 the
identified mutations in TA8-2 and other TA8 isolates were dispersed across a number of genes
and genomic locations and had a range of predicted effects on protein function.
Although there was no overlap in the mutated genes between TA1-1 and TA8-2, we did
note that both independent populations harbored mutations in core genes of the HOG pathway,
which regulates cellular stress responses in *C. neoformans* and other related species[31].In
particular, we identified a missense mutation in the *PBS2* gene of the TA1-1 isolate and a
frameshift mutation in the gene *SSK2* in the TA8 population strains[32]. The fact that we recovered
mutations in two key HOG-pathway genes in these independently evolved populations
highlights the importance of this pathway in responding to environmental stress and led us to
hypothesize that the HOG pathway may play an important role in regulating temperature
adaptation in this experiment.
To test this hypothesis and further probe the genetic basis of thermoadaptation in the
TA1-1 and TA8-2 isolates, we constructed a series of gene deletion mutants, focusing on genes
where we recovered coding sequence mutations. Spot assays of these mutants revealed that
deletion of the HOG pathway genes, *PBS2* and *SSK2*, had no impact on growth at high
temperature (Figure 5). Deletion of the genes *CNAG_07665* and *LMP1,* which carry
frameshifting mutations in TA1-1 and TA8-2, respectively, similarly had no impact on growth at
high temperature. Rather, we found that deletion of either the hexose transporter, *HXT1*
(CNAG_03438), or an unnamed, uncharacterized predicted serine-arginine protein kinase
(CNAG_02028) were sufficient for increased growth at both 35˚C and 37˚C when introduced to
the Ftc239-1 parental strain. The high temperature growth phenotypes of these deletion mutants
looked remarkably similar to those in our evolved clonal isolates, TA1-1 and TA8-2 which
acquired mutations in these genes via experimental evolution (Figure 5 and Tables 2 and 3).
These results strongly suggest that thermotolerance evolved through distinct mechanisms in

these two replicate populations: with mutation of *CNAG_02028* driving this trait in the TA1
population and an *HXT1* lesion promoting thermotolerance in TA8. Although it remains to be
seen what the impacts of the specific recovered mutations are on thermal adaptation along with
the molecular mechanisms by which these genes contribute to thermotolerance, these results
highlight the power of this experimental platform to uncover new understanding of the biology of
this critically important trait in fungi.
347
Discussion
Understanding how environmentally derived pathogenic microbes adapt to changes in
their environment, such as increasing temperatures, is of critical importance as climate change
advances. In this study, we employed experimental evolution to investigate the ability of
Cryptococcus neoformans, an important human fungal pathogen, to adapt to a high temperature
environment and how this alters consequential phenotypes.
We used an environmentally isolated *C. neoformans* strain, Ftc239-1, which exhibited
attenuated growth under normal laboratory culture conditions (30°C) and severe growth defects
at human body temperature (37°C). During serial passaging at increasing temperatures,
however, this strain was able to rapidly adapt, gaining the ability to grow better at high
temperature (35°C) than the parental strain after a very brief evolutionary timecourse (Figure 2).
The exact timing of when adaptive mutations arose in each population remains to be seen but
will further elucidate both the evolutionary trajectory of each independently evolved population
and how rapidly this new trait emerges in response to even gradual temperature shifts. Our
results also show that the growth advantage of our evolved strains was more pronounced at
37°C, which was outside the range of temperatures used during serial passaging. This suggests
that adaptation to modest temperature increases may affect growth at high temperatures more
broadly. But how far does this growth advantage extend? Probing whether there is an upper
temperature limit to the fitness advantage in these thermally adapted strains is an interesting
and important line of future inquiry.
Another critical question raised is whether adaptation to one form of stress (i.e.
temperature) confers resistance to other forms of stress and/or alters other infection-relevant
traits. Our data show that thermoadapted strains display a number of altered cellular features:
from widespread mucoidy and alterations in low-nutrient growth to changes in cell size variation
and antifungal drug resistance. Strikingly, our data characterizing antifungal drug resistance
across isolates from all populations of our evolution experiment suggest that rising
environmental temperatures may not only impact the prevalence of species capable of
establishing human infection but may also affect the treatment strategies for such infections
(Figure 4a and 4b). Many of our isolates displayed increases in resistance to the common
antifungal, fluconazole, as assessed through both MIC determinations and exposure to constant
high dose fluconazole. Although it remains to be seen whether this fluconazole resistance is
directly linked to thermoadaptation, this result was unexpected and highlights the ease with
which fungi can gain resistance to our limited therapeutic options—even in the absence of direct
exposure to these drugs. However, increased drug resistance was not a universal feature of our
evolved isolates; some, such as TA8-2 displayed increased sensitivity to antifungal drugs,
further underscoring the importance of studies examining the interplay between stress
resistance, generally, and thermal adaptation.
Through sequencing of two populations, we identified multiple, non-overlapping genetic
lesions which likely give rise to the diverse phenotypes we observe in our evolved isolates.
Among these, we identified a role for a predicted kinase, *CNAG_02028*, as well as the hexose
transporter *HXT1* in promoting thermotolerance in the TA1 and TA8 populations respectively
(Figure 5). In both cases, the mutations we recovered in these genes are missense mutations
and not clear loss of function alleles; however, the similarities in growth between the full deletion
mutants and our evolved clonal isolates hint that these missense mutations are likely to cause

loss of protein function. Neither of these genes have any reported role in nor regulation by high-
temperature growth in *C. neoformans.* Although sugar metabolism and, in particular, the
trehalose pathway in fungi, have been shown to be necessary for surviving high temperature
stress, it is particularly intriguing that mutation of just a single hexose transporter—among the
dozens encoded in the *C. neoformans* genome—can cause such a dramatic phenotype[26,33]. The
exact mechanisms through which *HXT1* participates in thermotolerance and whether there is a
link to nutrient conditions remain important outstanding questions.
Beyond the *HXT1* and *CNAG_02028* mutations, we hypothesize that other mutations in
our evolved isolates reflect one of several options. They could be passenger mutations that
were present when these high fitness impact mutations emerged or they could play more subtle
roles in adaptation to the stressful conditions of the experiment including both temperature and
other conditions, such as nutrient availability. Future work tracking the emergence of these
mutations over the course of the evolution experiment will help shed light on this question.
Collectively, our genetic data paired with the diversity of mutations and pathways identified by
sampling just two of our eight adapted populations supports the hypothesis that there may be
many routes by which this species can evolve thermotolerance.
Our work here begins a foray into the study of how our changing world will alter the
pathogens that infect us. As our findings suggest, host body temperature may not represent as
severe a barrier to *C. neoformans* and other fungal infections as environmental temperatures
increase. This work, therefore, provides a vital foundation to illuminate the consequences a
changing world will have for the health of future generations.
413
Materials and Methods
Strains and growth conditions
All strains were stored long term at –80°C in growth media—either yeast extract peptone
dextrose (YPD) or yeast nitrogen base (YNB)—supplemented with 20% glycerol. Strains were
inoculated by streaking out onto YPD plates and incubated for 2–3 days at 30°C before use in
experiments. A full list of strains used can be found in File S1.
420
Serial passaging of *C. neoformans* at increasing temperatures
For the first passage, an overnight culture of Ftc239-1 was inoculated from a single
colony and grown in YPD with shaking for 16 hours at 30°C. This culture was centrifuged at
3000 × g for 5 minutes and washed twice with autoclaved MilliQ water and resuspended in 5 mL
of water before the OD was measured. Cells were normalized to an OD of 0.05 (~4.04 x 10^6
cells/ml) in 5 mL of either YPD or YNB with 2% added glucose (4 replicate cultures of each).
The remaining starting inoculum culture was mixed 1:1 with 40% glycerol and aliquots were
frozen at –80°C.
The cultures were incubated at 30°C with shaking at 225 rpm for 48 hours. After 48
hours, the OD of each individual culture was measured and re-diluted to an OD of 0.05 in 5 mL
of the appropriate growth media. Passaging proceeded as above from this point. The
temperature was increased once the majority (>75%) of the cultures had reached three to four
doublings in the 48 hour culture period. Incubation temperature was increased by 1°C from
30°C to 33°C and 0.5°C from 33°C to 35°C. Aliquots from each individual culture were frozen
down as above at the start and end of each temperature, as well as every other passage.
Following the final passage at 35°C, multiple aliquots were frozen. Samples from each
population were also serially diluted and plated onto YPD plates for CFU determination. For all
of the populations except for TA1, serial dilution plates had >50 colonies. Five individual
colonies from each of these plates were selected randomly, picked into 5 mL of YPD, grown
overnight, and frozen for use in phenotyping experiments (Figure S1, Tables S1 and S2). For
population TA1, we only recovered 2 colonies after serial dilution and one was a bacterial
contaminant (see discussion below), so the only recovered *C. neoformans* isolate was saved as

above. Collectively, this gave us 36 clonal isolates from our eight evolved populations for further
phenotyping.
Though the experimental passaging was carried out based on OD_{600} measurements, the
actual amount of yeast passaged may not have been adequately reflected by the OD. In the
course of our characterization of experimentally evolved strains, we determined that both of the
populations discussed in depth here and likely all of the populations were contaminated with
Staphylococcus hominis, which absorbs at the same OD. This means that passage bottlenecks
were more severe than intended. However, this contamination did not affect our ability to isolate
thermally adapted fungal clones. Further, our data suggest that the bacterial contaminant has
not significantly impacted the thermal adaptation of the *Cryptococcus*.
453
Yeast spot assays
C. neoformans strains of interest were grown overnight in YPD at 30°C, subcultured to
an OD of 0.2 and grown to mid-log phase (OD_{600}=0.8-1). The cultures were then centrifuged at
3000 × g for 5 minutes, washed twice with autoclaved MilliQ water, and resuspended in 5 mL of
MilliQ water. The OD_{600} of each culture was measured, and the OD was adjusted to OD 1. Ten-
fold serial dilutions of each culture were then made down to OD 0.0001. 2.5 microliters of each
dilution was spotted onto desired media: YPD, YNB, or YPD plates with the indicated
concentrations of fluconazole. Spot plates were incubated at the desired temperature for a total
of 72 hours. Plates were imaged on Gel Doc XR+ (BioRad) or using a homemade imaging
setup after 48 and 72 hours. Unless otherwise indicated, representative images from spot
assays show growth after 48 hours.
465
Strain construction and molecular biology
All genetic manipulations were done using CRISPR-mediated gene editing and the
TRACE system following previously published protocols[34,35]. Briefly, gRNAs were generated
using guide-specific primer and amplification from pBHM2329 plasmid. *CAS9* was amplified for
transformation from the pBHM2403 plasmid.
For insertion of the hygromycin B cassette as the Safe Haven 2 locus (SH2), we used a
VNB lineage-specific guide targeting this locus. A repair template plasmid carrying the ~1 kb of
sequence immediately 5' and 3' to this gRNA sequence from the H99 genome was constructed
using HiFi DNA Master Mix (NEB Cat#E2621) with a standard HYG^R cassette inserted between
the two flanks. Repair templates were then PCR amplified from this plasmid for CRISPR
transformations. Gene deletion repair templates were generated following previously published
methods. Specifically, the 1000bp of sequence immediately 5' to the start codon and 3' to the
stop codon were cloned into the pUC19 vector backbone flanking the standard HYG^R cassette.
These plasmids were sequence verified and then used as template for CRISPR repair template
amplification. DNA concentrations in all CRISPR transformations were as follows: 250 ng Cas9,
100 ng gRNA, and 2-2.5 µg repair template.
482
Transformant selection and genotyping
Following electroporation of CRISPR reagents, strains were plated onto selective media
(YPD+HYG) and allowed to grow for 2-3 days at 30C. Individual colonies were then patched
onto non-selective YPD media for 2-3 passages before being patched back onto selective
media to identify stable genomic integrants.
Colonies were genotyped by colony PCR following previously published protocols[36].
Genomic sequences were amplified from gDNA extracted from candidate colonies and
integration of the drug cassette at the SH2 locus or at each individual gene locus was confirmed
by Sanger sequencing. PCRs were also performed to ensure that no strains had integrated the
gRNA or *CAS9* constructs. All primer sequences for generating CRISPR reagents and
genotyping mutants are listed in File S2.

Competition assays

For competition experiments, *C. neoformans* strains of interest were grown at 30°C
overnight in YPD. For experiments done in YPD, the OD was measured for each test strain and
for a parental Ftc239-1 strain carrying a HYG-resistance cassette at the SH2 locus. Cell
suspensions at a concentration of ~4.04 x 10^6 cells/mL were made for each individual strain.
Cell suspensions were mixed 1:1 (test strain:labeled parental) in a total of 40 mL YPD media
before adding 5 mL to individual culture tubes. Each competition was performed in biological
triplicate for each temperature (30°C or 35°C). CFUs for the initial mixtures were determined by
serially diluting the suspensions and plating onto both YPD and YPD+HYG plates. Competition
cultures were grown for 24 hours at 225 rpm at their respective temperatures. After 24 hours,
serial dilutions were plated onto nonselective and selective media plates as above. Competitive

Index here is calculated as: $Competitive\ Index = \dfrac{\frac{24\ hr\ CFU\ Test\ Strain}{24\ hr\ CFU\ HYG\ Control}}{\frac{0\ hr\ CFU\ Test\ Strain}{0\ hr\ CFU\ HYG\ Control}}$

For experiments done in YNB, overnight YPD cultures were washed twice and
resuspended in 5 mL of MilliQ water before setting up the experiment as above just in the
alternative YNB media.
510

Fluconazole MIC determinations

Overnight cultures of *C. neoformans* strains were adjusted to a density of ~5 x 10^6
cells/mL and were spread onto either YPD-agar or RPMI-agar medium with L-glutamine and
without sodium bicarbonate. Fluconazole Etest strips (bioMérieux) were placed on the agar
surface of one of the replicate plates once the cultures had dried. Plates were incubated at 35°C
and imaged after 48 and 72 hours. Only strains with robust growth on control YPD plates were
imaged and used to determine MIC values.
518

Titan cell induction and capsule assays

Titan cell induction was performed following previously published protocols[37]. Briefly,
strains were cultured overnight in YPD at 30°C. Cultures were centrifuged at 3000 × g for 5
minutes, washed three times in PBS, and resuspended in 3 mL of PBS. The cells were diluted
to ~5 x 10^3 cells/ml in 4 mL of serum-free RPMI 1640 media. One milliliter of each strain dilution
was plated into three replicate wells of a 24-well plate. The plate was incubated at 37C with 5%
CO_2 without shaking for 7 days. Cells were then fixed with formaldehyde, stained 1:1 with India
Ink, and imaged on a Zeiss Axio Observer 7 microscope at 100x.
Capsule was induced following previously published protocols[38]. *C. neoformans* strains
of interest were grown overnight (16-18 hours) in YPD at 30°C. Cultures were centrifuged,
washed twice with PBS, and resuspended in 5 mL of PBS. OD_{600} was measured for each strain,
and the strains were subcultured to an OD of 0.3 in 4 mL of DMEM media containing 10% FBS,
non-essential amino acids, and penicillin/streptomycin. Two milliliter aliquots of each dilution
were incubated in a 6-well plate at 37°C with 5% CO_2. After 48 hours of incubation, the cells
were fixed and imaged as above.
For both capsule and titan inductions, cell size measurements for each strain were made
manually for at least 50 cells under each condition. Capsule thickness here was calculated as:
capsule thickness = (total diameter – cell body diameter)/2.
537

Genomic DNA extraction and sequencing

Single colonies or pooled culture stocks were inoculated into 50 mL of YPD and grown
overnight, or until the culture was fully saturated, with shaking at 30C. Cells were collected by
centrifugation and lyophilized overnight. High-molecular-weight genomic DNA was isolated
following the CTAB protocol. Following CTAB extraction, samples were further purified using the

ProNex Size-Selective Purification System (Promega). Concentrations were measured by Qubit
dsDNA broad range assay. For Nanopore sequencing of Ftc239-1, one microgram of HMW
gDNA was prepared for sequencing using the SQK_LSK110 Kit and run on an R9 (FLO-
MIN106) flow cell. For Illumina sequencing, samples were sheared with a Covaris S2 Focused-
ultrasonicator and libraries were prepared using the New England Biolabs NEBNext Ultra II DNA
Library Prep kit (cat#E7634L) with an average insert size of 450 bp. Ten strains were barcoded
and pooled, and 150x150 bp paired-end reads were collected on a NovaSeq S4 Flow Cell with
the v1.5 Reagent Kit. Illumina library preparation and sequencing was performed by the
University of Utah's High Throughput Genomics core facility at the Huntsman Cancer Institute.
Raw sequencing datasets are available through the SRA under the BioProject Accession
PRJNA1096649.
554
Sequencing analysis
A chromosome scale assembly of the parental Ftc239-1 strain was drafted largely as
previously published[36]. Briefly, fast5 files from Nanopore sequencing were re-basecalled using
guppy-gpu (v6.4.2) and the resulting fastq files were assembled using Canu (v 2.1.1) with an
estimated genome size of 20 megabases and default settings[39]. The Canu assembly had 14
chromosome-scale contigs. Shorter contigs mapping to the mitochondrial genome or repetitive
genomic regions were manually trimmed. This assembly was polished once with Nanopore
reads using Medaka (1.7.2; https://github.com/nanoporetech/medaka) and then polished and
additional five times with Illumina reads from the parental strain using pilon (v1.24)[40]. The
mitochondrial genome was assembled from Illumina reads using GetOrganelle (v1.7.5.3) with
the fungus_mt setting and confirmed by comparison to the trimmed contigs assembled by
Canu[41]. This mitochondrial genome was manually added to the assembly. Chromosomes were
named based on comparison to other reference assemblies. The complete draft assembly was
annotated *de novo* using funannotate (v1.8.15) and publicly available RNA-seq datasets of *C.*
neoformans strains grown under a variety of conditions (SRA Accession Numbers: SRR646263,
SRR639772, SRR646268, SRR646265)[42]. *De novo* annotations were functionally annotated by
BLAST to the reference (H99) proteome available from Uniprot and annotation files were
updated using AGAT (v1.0.0)[43].
For Illumina sequencing data of isolates from the end of the evolution, adapters were
trimmed using Trimmomatic (v0.39) and reads were aligned to our *de novo* assembly using
bwa-mem2 (v 2.2.1)[44,45]. Duplicates were marked and read groups added with Picard and the
resulting files were sorted and indexed with SAMtools[46]. Variants were called using sorted BAM
files with FreeBayes (v0.9.21.7) with the following options: -p 1 --min-alternate-fraction 0.8 --
standard-filters[47]. The predicted impacts of called variants were assessed using SnpEff using
our *de novo* assembly as reference[48]. Called variants were also cross-checked by analysis with
breseq (v0.37.1) with settings altered to reject indels and variants predicted in regions of long
homopolymer runs (options: --consensus-reject-indel-homopolymer-length 3 --consensus-reject-
surrounding-homopolymer-length 3)[49]. All sequencing analysis was performed using resources
through the University of Utah Center for High Performance Computing.
584
585 **Data Availability Statement**
586 Strains generated by this study are available upon request. File S1 contains a list of all strains
587 used in this work. Sequencing data are available under the NCBI BioProject Accession
588 PRJNA1096649. Draft genome assembly and annotations for the Ftc239-1 parental isolate have
589 been uploaded to the GSA Figshare portal.
590
Acknowledgements and Funding

We thank John Perfect for providing the environmental isolates used in this study. This
work was funded by a Helen Hay Whitney Foundation postdoctoral fellowship and Boston
College startup funds to ZAH, a Burroughs Wellcome Fund Investigator in the Pathogenesis of
Infectious Disease award to NCE, and an NIH R35 (R35GM1 34936) awarded to NCE.

Author Contributions
Conceptualization: M.J.W.S., Z.A.H.
Methodology: M.J.W.S., L.A.F., M.R., Z.A.H.
Software: Z.A.H.
Validation: M.J.W.S., Z.A.H.
Formal Analysis: M.J.W.S., L.A.F., M.R., J.T.L., Z.A.H.
Investigation: M.J.W.S., L.A.F., M.R., J.T.L., Z.A.H.
Data Curation: M.J.W.S., L.A.F., M.R., J.T.L., Z.A.H.
Writing-Original Draft: M.J.W.S., Z.A.H.
Writing-Review and Editing: M.J.W.S., Z.A.H., N.C.E.
Visualization: M.J.W.S., Z.A.H.
Supervision: Z.A.H., N.C.E.
Funding acquisition: N.C.E., Z.A.H.

Competing Interests
The authors declare no competing financial interests.

References

1. *Climate Change 2022: Impacts, Adaptation, and Vulnerability - Summary for Policymakers*.
https://www.ipcc.ch/report/ar6/wg2/chapter/summary-for-policymakers/ (2022).
2. Garcia-Solache, M. A. & Casadevall, A. Global Warming Will Bring New Fungal Diseases
for Mammals. *MBio* **1**, e00061-10 (2010).
3. Robert, V. A. & Casadevall, A. Vertebrate Endothermy Restricts Most Fungi as Potential
Pathogens. *J. Infect. Dis.* **200**, 1623–1626 (2009).
4. Huang, J. *et al.* Pan-drug resistance and hypervirulence in a human fungal pathogen are
enabled by mutagenesis induced by mammalian body temperature. *Nat. Microbiol.* **9**,
1686–1699 (2024).
5. Gorris, M. E., Treseder, K. K., Zender, C. S. & Randerson, J. T. Expansion of
Coccidioidomycosis Endemic Regions in the United States in Response to Climate
Change. *GeoHealth* **3**, 308–327 (2019).
6. Cogliati, M. Global warming impact on the expansion of fundamental niche of Cryptococcus
gattii VGI in Europe. *Environ. Microbiol. Rep.* **13**, 375–383 (2021).
7. Casadevall, A., Kontoyiannis, D. P. & Robert, V. Environmental Candida auris and the
Global Warming Emergence Hypothesis. *MBio* **12**, e00360-21 (2021).
8. Cavicchioli, R. *et al.* Scientists' warning to humanity: microorganisms and climate change.
Nat. Rev. Microbiol. **17**, 569–586 (2019).
9. Nnadi, N. E. & Carter, D. A. Climate change and the emergence of fungal pathogens. *PLoS*
Pathog. **17**, e1009503 (2021).
10. Idnurm, A. *et al.* Deciphering the Model Pathogenic Fungus Cryptococcus Neoformans.
Nat. Rev. Microbiol. **3**, 753–764 (2005).
11. Rajasingham, R. *et al.* The global burden of HIV-associated cryptococcal infection in adults
in 2020: a modelling analysis. *Lancet Infect. Dis.* **22**, 1748–1755 (2022).
12. Rhodes, J. *et al.* Tracing Genetic Exchange and Biogeography of Cryptococcus
neoformans var. grubii at the Global Population Level. *Genetics* **207**, 327–346 (2017).

13. Mitchell, T. G., Castañeda, E., Nielsen, K., Wanke, B. & Lazéra, M. S. Environmental Niches for Cryptococcus neoformans and Cryptococcus gattii. in *Cryptococcus* 235–259 (John Wiley & Sons, Ltd, 2010).

14. Desjardins, C. A. *et al.* Population genomics and the evolution of virulence in the fungal pathogen Cryptococcus neoformans. *Genome Res.* **27**, 1207–1219 (2017).

15. Chen, Y. *et al.* Comparative analyses of clinical and environmental populations of Cryptococcus neoformans in Botswana. *Mol. Ecol.* **24**, 3559–3571 (2015).

16. Litvintseva, A. P. *et al.* Evidence that the human pathogenic fungus Cryptococcus neoformans var. grubii may have evolved in Africa. *PLoS One* **6**, e19688 (2011).

17. Brown, S. M., Campbell, L. T. & Lodge, J. K. Cryptococcus neoformans, a fungus under stress. *Curr. Opin. Microbiol.* **10**, 320–325 (2007).

18. Glazier, V. E. & Panepinto, J. C. The ER stress response and host temperature adaptation in the human fungal pathogen Cryptococcus neoformans. *Virulence* **5**, 351–356 (2014).

19. Leach, M. D. & Cowen, L. E. Surviving the Heat of the Moment: A Fungal Pathogens Perspective. *PLoS Pathog.* **9**, e1003163 (2013).

20. Stempinski, P. R. *et al.* Genetic contribution to high temperature tolerance in Cryptococcus neoformans. *Genetics* **217**, iyaa009 (2021).

21. Odom, A. *et al.* Calcineurin is required for virulence of Cryptococcus neoformans. *EMBO J.* **16**, 2576–2589 (1997).

22. Bennett, A. F. & Lenski, R. E. Evolutionary Adaptation to Temperature II. Thermal Niches of Experimental Lines of Escherichia coli. *Evolution* **47**, 1–12 (1993).

23. Garcia-Rios, E. *et al.* Thermo-adaptive Evolution to Generate Improved Saccharomyces cerevisiae Strains for Cocoa Pulp Fermentations. *Int. J. Food Microbiol.* **342**, 109077 (2021).

24. Huang, C.-J., Lu, M.-Y., Chang, Y.-W. & Li, W.-H. Experimental Evolution of Yeast for High-Temperature Tolerance. *Mol. Biol. Evol.* **35**, 1823–1839 (2018).

25. Longan, E. R. & Fay, J. C. Experimental evolution of Saccharomyces uvarum at high temperature yields elevation of maximal growth temperature and loss of the mitochondrial genome. *microPublication Biology* (2023) doi:10.17912/micropub.biology.000831.

26. Petzold, E. W. *et al.* Characterization and regulation of the trehalose synthesis pathway and its importance in the pathogenicity of Cryptococcus neoformans. *Infect. Immun.* **74**, 5877–5887 (2006).

27. Lee, I. R., Chow, E. W. L., Morrow, C. A., Djordjevic, J. T. & Fraser, J. A. Nitrogen metabolite repression of metabolism and virulence in the human fungal pathogen Cryptococcus neoformans. *Genetics* **188**, 309–323 (2011).

28. Montoya, M. C., Magwene, P. M. & Perfect, J. R. Associations between Cryptococcus Genotypes, Phenotypes, and Clinical Parameters of Human Disease: A Review. *Journal of Fungi* **7**, 260 (2021).

29. Guerrero, A., Jain, N., Goldman, D. L. & Fries, B. C. Phenotypic switching in Cryptococcus neoformans. *Microbiology* **152**, 3–9 (2006).

30. Dambuza, I. M. *et al.* The Cryptococcus neoformans Titan cell is an inducible and regulated morphotype underlying pathogenesis. *PLoS Pathog.* **14**, e1006978 (2018).

31. Bahn, Y.-S. & Jung, K.-W. Stress Signaling Pathways for the Pathogenicity of Cryptococcus. *Eukaryot. Cell* **12**, (2013).

32. Bahn, Y.-S., Geunes-Boyer, S. & Heitman, J. Ssk2 Mitogen-Activated Protein Kinase Kinase Kinase Governs Divergent Patterns of the Stress-Activated Hog1 Signaling Pathway in Cryptococcus neoformans. *Eukaryot. Cell* **6**, 2278–2289 (2007).

33. Liu, T.-B. *et al.* The glucose sensor-like protein Hxs1 is a high-affinity glucose transporter and required for virulence in Cryptococcus neoformans. *PLoS One* **8**, e64239 (2013).

34. Fan, Y. & Lin, X. Multiple Applications of a Transient CRISPR-Cas9 Coupled with
Electroporation (TRACE) System in the Cryptococcus neoformans Species Complex.
Genetics **208**, 1357–1372 (2018).
35. Huang, M. Y. *et al.* Short homology-directed repair using optimized Cas9 in the pathogen
Cryptococcus neoformans enables rapid gene deletion and tagging. *Genetics* **220**, iyab180
(2021).
36. Hilbert, Z. A. *et al.* Distinct pathways of adaptive evolution in Cryptococcus neoformans
reveal a mutation in adenylyl cyclase with trade-offs for pathogenicity. *Curr. Biol.* **33**, 4136-
4149.e9 (2023).
37. Saidykhan, L. *et al.* An in vitro method for inducing titan cells reveals novel features of
yeast-to-titan switching in the human fungal pathogen Cryptococcus gattii. *PLoS Pathog.*
18, e1010321 (2022).
38. Zaragoza, O. & Casadevall, A. Experimental modulation of capsule size in Cryptococcus
neoformans. *Biol. Proced. Online* **6**, 10–15 (2004).
39. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).
40. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection
and genome assembly improvement. *PLoS One* **9**, e112963 (2014).
41. Jin, J.-J. *et al.* GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of
organelle genomes. *Genome Biol.* **21**, 241 (2020).
42. Funannotate documentation — Funannotate 1.8.16 documentation.
https://funannotate.readthedocs.io/en/latest/.
43. Welcome to AGAT's documentation! — AGAT documentation.
https://agat.readthedocs.io/en/latest/.
44. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).
45. Vasimuddin, M., Misra, S., Li, H. & Aluru, S. Efficient Architecture-Aware Acceleration of
BWA-MEM for Multicore Systems. in *2019 IEEE International Parallel and Distributed*
Processing Symposium (IPDPS) 314–324 (IEEE, 2019). doi:10.1109/IPDPS.2019.00041.
46. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).
47. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.
arXiv [q-bio.GN] (2012).
48. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-
2; iso-3. *Fly* **6**, 80–92 (2012).
49. Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. *Methods Mol. Biol.* **1151**, 165–188
(2014).
729
730

Figure Legends

Figure 1. Phenotypic variation of *C. neoformans* environmental isolate growth at high temperatures. (a) Phylogenetic tree of *C. neoformans* strains used in this study adapted from Desjardins *et. al*. The primary strain used in subsequent figures, Ftc239-1, is indicated in bold and boxed. (b and c) Growth of *C. neoformans* strains on YNB (b) or YPD (c) agar plates at 30°C (left) or 37°C (right).

Figure 2. Serial passaging of *C. neoformans* selects for strains with increased thermotolerance. (a) Schematic of the serial passaging set-up used in this study. See also Materials and Methods. Populations TA1-TA4 were passaged in YPD; TA5-TA8 were passaged in YNB. (b and c) Growth of serially passaged isolates on YPD (b) or YNB (c) agar plates. Strains were tested for growth at temperatures they experienced during the experimental evolution passaging (30 and 35°C, middle panels). Growth was also assessed at temperatures outside of the range used during passaging (26°C and 37°C, outer panels).

Figure 3. Thermally-adapted isolates outcompete the parental strain at high temperature. (a) Schematic of the competition experiment set-up between the HYG^R-labeled parental strain and serially-passaged isolates at 30°C or 35°C. See also Materials and Methods. (b and c) Competition experiments between HYG^R-labeled parental and unlabeled parental (blue), TA1-1 (dark red), TA8-2 (orange) strains in either YPD (b) or YNB (c) at both 30°C and 35°C. Competitive indices (CIs) are calculated as indicated in Methods. A CI=1 (dotted line) indicates equal growth of the two strains. Plotted values indicate the average value ± SEM of three replicate experiments carried out on three separate days; dots indicate the average value of biological replicates from a single experiment.

Figure 4. Thermal adaptation alters titan cell formation and antifungal drug susceptibility in serially-passaged isolates. (a) Measurements of cell body diameter from cells grown under titan inducing conditions. Percentages above each strain indicate the percent of cells with a diameter >10 µm (dotted line). Error bars show mean ± SD for one representative experiment. 100 cells per strain were measured with the exception of TA2-4 which displayed a growth defect under these conditions and only 60 cells could be quantified. (b) MIC values for expanded panel of clonal isolates as determined using fluconazole E-tests. Bars show the mean and ± SD of two replicate experiments with individual values plotted as dots. Mean MIC numerical values are also provided in Table 1. (c) Growth of a subset of isolates on control YPD plates, or YPD plates containing 4 or 12 µg/mL of the antifungal fluconazole. Images show 72 hours of growth at 35˚C. The parental strain along with the two main isolates examined are boxed in color.

Figure 5. Deletion of the genes *HXT1* or *CNAG_02028* are sufficient for growth at 37˚C in the Ftc239-1 background. (a) Comparison of growth across temperatures for the TA1-1 evolved isolate and gene deletion mutants for genes that carry genetic lesions in TA1-1. (b) Comparison of growth of the TA8-2 isolate compared to deletion mutants for the genes that carry coding sequence mutations in TA8-2. All images show serial dilutions plated on YPD and incubated at the indicated temperatures.

777 **Tables**
778 **Table 1. Average fluconazole MIC values for experimentally evolved isolates.** Numerical
779 values of the data plotted in Figure 4b. Averages are for two MIC readings taken from different
780 experimental replicates (on different days).
781

Strain	Evolution Media	Growth at 37˚C	Average FLC MIC (µg/mL)
Ftc239-1	--	-	6
TA1-1	YPD	+	12
TA2-4	YPD	+	7
TA2-5	YPD	-	9
TA3-1	YPD	+	10
TA3-5	YPD	+	36
TA4-1	YPD	+	6
TA4-5	YPD	+	11
TA5-1	YNB	+	10
TA5-5	YNB	+	7
TA6-2	YNB	-	12
TA6-3	YNB	+	10
TA7-3	YNB	+	14
TA7-5	YNB	+	10
TA8-2	YNB	+	3.5

782
783 **Table 2. Fixed mutations in the TA1-1 YPD-passaged strain.**
784

Chromosome	Position	Gene	Gene Name and Info	Mutation Effect	REF	ALT
1	2,000,229	*CNAG_00769*	*PBS2*, HOG pathway MAPKK	Y372*	C	A
2	973,304	*CNAG_03735*	*CAP4*, predicted glycosyl transferase	Intronic	GAAAAAAAAG	GAAAAAAAAG
6	1,422,809	*CNAG_02028*	predicted serine-arginine protein kinase	G153E	C	T
7	930,986	*CNAG_07665*	hypothetical protein	4 nt Insertion, Frameshift	C	GGGCG

Table 3. Fixed mutations in the TA8 YNB-passaged population. These mutations are present in all of the sequenced TA8 clonal isolates.

Chromosome	Position	Gene	Gene Name and Info	Mutation Effect	REF	ALT
2	1,592,272	CNAG_06765	LMP1, low mating performance protein	1 nt deletion, Frameshift	TT	T
4	328,289	CNAG_05063	SSK2, HOG pathway MAPKKK	1 nt deletion, Frameshift	TT	T
5	723,192	Intergenic			G	C
8	435,015	CNAG_03438	HXT1, hexose transporter	G63R	G	C
10	1,058,846	Intergenic (CNAG_04922 3' UTR)	CNAG_04922, hypothetical protein		C	A
14	928,714	Intergenic	Intergenic		C	T

Table 4. Additional fixed mutations in TA8-derived strains. These mutations are only present in the indicated colony from the TA8 population.

Strain	Chromosome	Position	Gene	Gene Name and Info	Mutation Effect	REF	ALT
TA8-1	7	289,398	Intergenic (CNAG_05884 5' UTR)	KAP123, predicted importin beta family		CTTTTTTTTTC	CTTTTTTTTC
TA8-3	8	858,513	Intergenic			G	C
TA8-3	10	843,898	CNAG_04855	hypothetical protein	A248S	G	T
TA8-4	3_11	308,737	CNAG_02599	predicted aldolase	N24N	A	G

Figure S1. Phenotypic variation of *C. neoformans* parental and evolved isolates. (a and b)
Growth of *C. neoformans* parental strain and YPD (A) and YNB (B) evolved isolates on YPD
(left) or YNB (right) agar plates at 30 and 35°C.

**Figure S2. Evolved isolates from two populations display changes in titan cell formation
and antifungal resistance.** (a) Representative images from cells in titan-inducing (left) and
capsule-inducing (right) conditions for parental and evolved strains. Images from different
conditions are at the same scale with all scale bars representing 10 µm. (b) Measurements of
capsule thickness from cells grown under capsule inducing conditions. Capsule thicknesses
were calculated as indicated in Materials and Methods. Median and quartiles are indicated with
dashed lines for cells from one representative experiment (c) Measurements of cell body
diameter from cells grown under titan inducing conditions. Percentages above each strain
indicate the percent of cells with a diameter >10 µm (dotted line). Error bars show mean ± SD
for one representative experiment. (d) Fluconazole minimum inhibitory concentrations (MICs) for
the parental and evolved isolates on RPMI agar plates with E-test strips. For (b) and (c), 50 cells
were measured per strain. Significance was assessed in comparison to the parental Ftc239-1
strain by ordinary one-way ANOVA with Dunnett's multiple comparisons test.

**Figure S3. Evolved isolates from many independent populations show enhanced
resistance to fluconazole.** Growth phenotypes of 14 experimentally evolved isolates on control
YPD plates (left) or plates containing the indicated concentrations of fluconazole (right). All
plates were incubated at 35°C and images show 72h of growth. Images for strains shown in
Figure 4c are identical in this panel and included to allow for comparisons across all of the
tested isolates.

**Table S1. Colony phenotypes of individual colony isolates of the YPD-passaged
populations (TA1-TA4).** Isolates discussed in this study are indicated by darkened background.

Population	Colony #	Colony Appearance	Colony Size
Ftc239-1	--	Smooth	Small
TA1	1	*Mucoid*	*Normal*
	1	*Mucoid*	*Normal*
	2	*Mucoid*	*Normal*
	3	Smooth	*Normal*
	4	*Mucoid*	*Normal*
TA2	5	Smooth	*Normal*
	1	*Mucoid*	*Normal*
	2	*Mucoid*	*Normal*
	3	*Mucoid*	*Normal*
	4	*Mucoid*	*Normal*
TA3	5	Smooth	*Normal*
	1	*Mucoid*	*Normal*
	2	*Mucoid*	*Normal*
TA4	3	*Mucoid*	*Normal*

| | 4 | Mucoid | Normal |
| | 5 | Mucoid | Normal |

824
825
Table S2. Colony phenotypes of individual colony isolates of the YNB-passaged populations (TA5-TA8). Isolates discussed in this study are indicated by darkened backgrounds.

Population	Colony #	Colony Appearance	Colony Size
Ftc239-1	--	Smooth	Small
	1	Smooth	Small
	2	Smooth	*Normal*
	3	Smooth	Small
	4	Smooth	Small
TA5	5	Smooth	Small
	1	Smooth	*Normal*
	2	Smooth	*Normal*
	3	Smooth	Small
	4	Smooth	*Normal*
TA6	5	Smooth	Small
	1	*Mucoid*	*Normal*
	2	*Mucoid*	*Normal*
	3	*Mucoid*	*Normal*
	4	*Mucoid*	*Normal*
TA7	5	*Mucoid*	*Normal*
	1	Smooth	*Normal*
	2	Smooth	*Normal*
	3	Smooth	Small
	4	Smooth	*Normal*
TA8	5	Smooth	Small

830
831
Supplemental Files
File S1. List of strains used in this study.
File S2. List of primers used in this study.

	Control	4 µg/mL FLC	12 µg/mL FLC
Ftc239-1			
TA1-1			
TA2-4			
TA2-5			
TA3-1			
TA3-5			
TA4-1			
TA4-5			
TA5-1			
TA5-5			
TA6-2			
TA6-3			
TA7-3			
TA7-5			
TA8-2			