Plasmodium falciparum protein phosphatase PP7 is required for early ring-stage development

Avnish Patel[1], Aline Fréville[1], Joshua A. Rey[1], Helen R. Flynn[3], Konstantinos Koussis[2], Mark J. Skehel[3], Michael J. Blackman[2,1] and David A. Baker[1]*

1. Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, United Kingdom.

2. Malaria Biochemistry Laboratory, The Francis Crick Institute, London, United Kingdom

3. Proteomics laboratory, The Francis Crick Institute, London, United Kingdom and Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, United Kingdom.

*Corresponding author

david.baker@lshtm.ac.uk

Abstract

We previously reported that the *Plasmodium falciparum* putative serine/threonine protein phosphatase 7 (PP7) is a high confidence substrate of the cAMP-dependent protein kinase (PKA). Here we explore the function of PP7 in asexual *P. falciparum* blood stage parasites. We show that conditional disruption of PP7 leads to a severe growth arrest. We show that PP7 is a calcium-dependent phosphatase which interacts with calmodulin and calcium-dependent protein kinase 1 (CDPK1), consistent with a role in calcium signalling. Notably, PP7 was found to be dispensable for erythrocyte invasion, but was crucial for ring-stage development, with PP7-null parasites arresting shortly following invasion and showing no

21   transition to ameboid forms. Phosphoproteomic analysis revealed that PP7 may regulate certain PKAc

22   substrates. Its interaction with calmodulin and CDPK1 further emphasise a role in calcium signalling,

23   while its impact on early ring development and PKAc substrate phosphorylation underscores its

24   importance in parasite development.

Introduction

Protein phosphorylation is an essential post-translational modification event that is dynamically

controlled by the action of protein kinases and protein phosphatases in all eukaryotic cells. The major

human malaria pathogen *Plasmodium falciparum* possesses 90 putative protein kinases (1, 2) and 27

putative protein phosphatases (3). Genome-wide screens in *P. berghei*, a rodent malaria species, has

identified the stage-specific essentiality of kinases and phosphatases (4, 5). With the widespread

adoption of conditional gene disruption technologies in *Plasmodium*, several of the essential kinases

have been further studied for their cell-specific roles (6–11). By contrast, essential protein phosphatases

have remained relatively under-explored (12), yet an improved understanding of their function is a

requisite to discerning the global role of protein phosphorylation dynamics in the complex malaria

parasite life cycle, and can also inform target-based drug discovery efforts for rational drug design.

Previous studies have assigned functions for essential phosphatases to processes late in the asexual

blood stage cycle, revealing roles for calcineurin for merozoite host cell attachment (13, 14), PP1 in

schizont DNA replication and egress (15), and NIF4 in merozoite invasion (16). However, to our

knowledge no malarial protein phosphatase has been shown to play a role in the early developmental

phases of asexual blood stage parasites.

We previously identified 39 potential substrates of the cAMP dependent protein kinase catalytic subunit

(PKAc), an essential kinase required for merozoite invasion of host erythrocytes. Phosphosites within

these protein substrates were found to be upregulated following disruption of adenylyl cyclase β or

PKAc, and correspondingly downregulated following deletion of phosphodiesterase β (17) (7). Amongst

the putative PKAc substrates identified, PP7 was considered of particular interest in our efforts to

elucidate the cell signalling events underlying erythrocyte invasion due to its predicted essentiality and

its peak transcriptional expression in mature schizonts.

Here we investigate the function of *P. falciparum* PP7. We find it to be indispensable for asexual blood

stage parasite growth, and specifically for the development of early ring-stage parasites. We show that

PP7 is expressed in late schizont stages, that its activity is Ca^{2+}-dependent, and that it interacts with both

calmodulin and the Ca^{2+}-dependent protein kinase CDPK1. Finally, we use global phospho-proteomics to

identify sites that are differentially regulated by PP7.

Results

The architecture of PP7 includes calcium-binding domains

PP7 is annotated as a member of the Phospho-Protein Phosphatases (PPP) family (PlasmoDB (18)

PF3D7_1423300). Its overall structure comprises a central predicted serine/threonine protein

phosphatase domain (19) flanked by putative Ca^{2+}-binding-related domains: an N-terminal calmodulin

binding IQ domain (20) and two C-terminal Ca^{2+}-binding EF hands (21) (Fig 1A). This architecture is

characteristic of members of one of the five sub-families of the phosphoprotein phosphatase (PPP)

family, PPEF/PP7, with EF referring to the presence of an EF-hand domain downstream of the

phosphatase catalytic domain. Figure S1 shows the domain structure of the *P. falciparum* PP7 and

alignments with orthologues from other eukaryotes (22).

P. falciparum PP7 is expressed in mature schizonts and is essential for asexual blood stage parasite

growth

To investigate the biological function of PP7 in *P. falciparum* blood stages, we generated a transgenic

parasite line (called PP7-HA:loxP) in a manner designed to allow determination of both the subcellular

location of PP7 and the consequences of conditional disruption of PP7. The mutant line was generated

on the genetic background of a 3D7 *P. falciparum* line that stably expresses dimerisable Cre (DiCre), the

Cre-recombinase activity of which is induced in the presence of rapamycin (RAP) (23). The PP7 gene was

'floxed' such that treatment with RAP would lead to excision of DNA sequences encoding the

phosphatase catalytic domain and all downstream features (Fig 1B). At the same time the gene was also

modified by fusion to sequence encoding a C-terminal triple hemagglutinin (HA) epitope tag (Fig 1B).

Successful modification of the PP7 locus in PP7-HA:loxP parasites was verified by PCR (Fig 1B and C) and

western blot, which also showed loss of expression following RAP-induced truncation of the tagged PP7-

HA (Fig 1D). Immunofluorescence analysis (IFA) of the transgenic line revealed a partly peripheral signal

in individual merozoites within mature segmented schizonts (Fig 1E).

To investigate the essentiality of PP7, highly synchronised young ring-stage cultures of PP7-HA:loxP were

treated with RAP to induce excision of the sequence encoding the phosphatase catalytic domain at 4

hours (h) post invasion (Fig 1B). RAP-treated PP7-HA:loxP parasites seem to mature normally through

the cycle of RAP-treatment   (cycle 0). The RAP-treated schizonts produce the same number of

merozoites (Fig S2A), have the same DNA content as schizonts treated with Vehicle (DMSO) only (Fig

S2B), egress and re-invade normally (Fig S2C and D, movie S1). However, early in the next cycle (cycle 1),

the parasites underwent severe growth arrest (Fig 2). Examination of the cycle 1 parasites by Giemsa-

staining showed that the development of early ring stages was adversely affected in RAP-treated PP7-

null parasites, with an accumulation of pycnotic and abnormal parasites that appeared immediately

after invasion. In contrast, new rings of control PP7-HA:loxP parasites treated with DMSO, developed

normally and went on to form schizonts (Fig 2 inset, Fig 5A and 5B). It was concluded that PP7 is

essential for asexual blood stage survival and that it plays a role in formation of ring-stage parasites soon

after erythrocyte invasion.

PP7 is a calcium-dependent phosphatase

Since the domain structure of *P. falciparum* PP7 includes putative domains involved in calcium sensing,

we sought to investigate whether PP7 is a calcium-dependent phosphatase. For this, we isolated PP7

from extracts of PP7-HA:loxP parasites by immuno-precipitation with magnetic anti-HA conjugated

beads (Fig 3A). The immobilized PP7 was then examined for phosphatase activity in the absence or

presence of Ca^{2+} or chelating agents. As shown in in Fig 3B, PP7 displayed phosphatase activity which

was strongly enhanced by the presence of Ca^{2+} and inhibited by calcium chelators (Fig 3B). These results

indicate that parasite-derived PP7 is a functional phosphatase and that its activity is Ca^{2+} dependent.

PP7 interacts with calmodulin and CDPK1

The observed calcium-dependence of PP7 activity raised the possibility that it may play a role in calcium-

mediated signalling. To evaluate this and to gain further mechanistic insight into the role of PP7, we

sought to identify potential PP7-interacting partner proteins. For this, immuno-precipitated material

from PP7-HA:loxP and 3D7 DiCre (negative control) schizonts were analysed by mass spectrometry to

identify co-purifying protein species with and without the presence of additional Ca^{2+}. As shown in Fig

4A and Supplementary Table 1, this analysis revealed that PP7 interacts with calmodulin and calcium-

dependent protein kinase 1 (CDPK1), an enzyme involved in parasite invasion and development (9). This

finding was confirmed by western blot analysis of PP7 immuno-precipitates using a CDPK1-specific

antibody (Fig 4B). The mass spectrometric analysis also identified a conserved protein of unknown

function (PF3D7_0423300) that was only enriched in the absence of added Ca^{2+}, suggesting a calcium-

dependent interaction of PP7 and PF3D7_0423300. Collectively, these results were interpreted as

indicating a role for PP7 in calcium-mediated signalling in *P. falciparum*.

Loss of PP7 results in phosphoproteomic changes

To gain further insight into the mechanisms through which PP7 controls early ring-stage development,

we profiled the PP7-dependent phosphoproteome from schizonts of PP7-null and control parasites

cultures, reasoning that these are most likely to represent the developmental lifecycle stages in which

Ca^{2+}-dependent signalling could plausibly exert control over PP7 activity. Phosphopeptides enriched

from trypsin-digested protein extracts of control and RAP-treated PP7-HA:loxP parasites were examined

by tandem mass spectrometry using isobaric labelling for quantification. This showed that whilst

phosphorylation sites of PP7 were significantly down-regulated in the PP7-null parasites, as expected,

phosphosites of only four other proteins showed significant down- or up-regulation: the up-regulated

sites were on the peptidyl-prolyl cis-trans isomerase FKBP35 (PF3D7_1247400) and on PfMGET

(Pf3D7_1469900), whilst the down-regulated sites were on a putative EF hand domain-containing

protein (PF3D7_1221300) which was found to be expressed in late schizonts in a transcriptomic study

(24), on a putative peroxiredoxin (PfnPRx-Pf3D7_1027300) and on Pf3D7_1021700; a VPS13 domain-

containing protein (Fig 4C, Table S2). Gene ontology (GO) enrichment analysis on hyper- and hypo-

phosphorylated proteins suggested, primarily, deregulation in protein dephosphorylation and NADH

dehydrogenase activity (Fig S3). These results confirmed that loss of PP7 results in a selective set of

phosphoproteomic changes in the parasite that could underlie the ring development phenotype.

PP7 is not required for invasion but is essential for early ring-stage development

To further investigate how loss of PP7 affects asexual parasite development, we quantified and

characterized schizont and resultant ring formation over time in control and RAP-treated (PP7-null) PP7-

HA:loxP parasites. This showed no effects of PP7 deletion on the apparent numbers of schizonts and

subsequent ring stage formation, as determined by flow cytometry (Fig 5C). However, microscopic

analysis of Giemsa-stained parasites 2 h post-invasion showed that newly intraerythrocytic PP7-null

parasites did not develop post-invasion, instead displaying a predominantly dot-like morphology (Fig 5A,

5B and Fig 6A). To analyse this ring-stage development defect in more detail we modified the PP7-

HA:loxP line to express a cytosolic mNeon-Green marker (7), resulting in the PP7-HA-cyto-mNeon:loxP

line in which we could observe the morphology of the cell cytosol by green fluorescence in live cells.

Imaging of control or RAP-treated PP7-HA-cyto-mNeon:loxP parasites that had additionally been stained

with a membrane marker BODIPY Tr ceramide and DNA stain Hoechst 33342 showed that RAP-treated

(PP7-null) parasites formed pycnotic structures that failed to expand, in contrast with the clearly-defined

ameboid structures in control parasites (Fig 6B). Together these data indicate a crucial role for PP7 in

early ring-stage ameboid development.

Discussion

The findings presented in this study shed light on the crucial role of PP7, a biochemically active

serine/threonine protein phosphatase, in *P. falciparum* blood stages, expanding our understanding of

the signalling machinery of this deadly parasite.

Our data unequivocally demonstrate that PP7 is indispensable for asexual blood stage survival in *P.*

falciparum. Conditional disruption of PP7 led to a severe growth arrest, emphasizing its role in early

asexual blood stage parasite replication. This aligns with previous findings from a global *P. berghei*

knockout screen (4), reinforcing the notion that PP7 plays a conserved and vital role in *Plasmodium*

species. While PP7 was found to be dispensable for erythrocyte invasion, its requirement for early ring-

stage development is clear. The striking accumulation of pycnotic parasites in early ring stages upon PP7

disruption led us to investigate and confirm a role in amoeboid formation. This amoeboid form has

recently been described (25) and remains an understudied aspect of early invaded parasite

development. The developmental arrest observed in PP7-null rings may suggest a crucial role in the

organization or maturation of structures critical for the development of newly invaded parasites that

contribute to the amoeboid morphology. Interestingly, the *Toxoplasma* orthologue of PP7, *Tg*PP7, is also

essential for parasite replication and virulence *in vivo* (26). In contrast to our findings with *P. falciparum*

PP7, however, *Tg*PP7 was found to have a role in tachyzoite invasion. This implies the evolutionary co-

option of PP7 to suit species-specific roles in apicomplexan invasion and early invaded parasite

development. Further work is warranted to elucidate the precise mechanisms underlying these

differences.

The discovery of PP7 as a calcium-dependent phosphatase is a significant revelation. The pronounced

increase in PP7 activity in the presence of Ca^{2+}, which could be reversed by calcium chelation, implies a

role in calcium signalling, which is pivotal during various transitions during the asexual blood stage

lifecycle, including erythrocyte invasion and egress (6, 9, 27, 28). Our results suggest that the calcium-

dependent phosphatase activity of PP7 plays a crucial role in regulating key calcium-dependent events in

the parasite life cycle.

These conclusions are supported by our identification of calmodulin and calcium-dependent protein

kinase 1 (CDPK1) as interacting partners of PP7. Calmodulin is a known calcium sensor, and CDPK1 plays

a crucial role in parasite invasion and development. The apparent interaction of PP7 with these proteins

strongly suggests a role for PP7 in orchestrating crucial calcium-dependent signalling cascades. Given

that PP7 is a high confidence PKAc substrate (7), it is plausible to suggest that PP7 may act as molecular

player in the interaction between the PKAc and calcium-mediated cell signalling pathways in blood stage

schizonts. Indeed, cross-talk between these two pathways has been previously observed (29, 30) but the

precise molecular processes governing this remain to be elucidated. We suggest that this role could be

fulfilled in part by PP7.

(26)

The phosphoproteomic profiling of PP7-dependent processes in mature schizonts represents a rich

dataset for further exploration. We detected five proteins other than PP7, of which the phosphorylation

status is altered significantly upon loss of PP7. Among them, three (PF3D7_1021700, PF3D7_1027300,

PF3D7_1221300) were also found to be deregulated in the PKAc null parasite phosphoproteome (7). The

putative EF hand domain-containing protein PF3D7_1221300 is predicted to be non-essential (31), but

nonetheless underscores the potential role of PP7 in calcium signaling. The peptidyl-prolyl cis-trans

isomerase FKBP35 (PF3D7_1247400) is predicted to be essential (31) and is most highly expressed in the

early schizont stage (24). FK506-binding proteins have been shown to catalyze the cis-trans

isomerization of proline imidic peptide bonds (32) and have a role in protein folding (33) but little is

known of the specific cellular roles in model organisms. The potential role of PP7 in regulation of protein

folding processes presents an interesting avenue for future research. The only other protein predicted

to be essential is Pf3D7_1021700 (31)and harbors a VPS13 domain. VPS13 proteins are known lipid

transporters, targeted to distinct membranes and particularly membrane contact sites. In yeast, VPS13

is involved in membrane expansion(34). Although plausible that it has a similar role in *Plasmodium*,

contributing to the ablation of ring formation, no mechanistic evidence is currently available. It will be

interesting to investigate further whether PP7 disruption affects phosphorylation of PKAc substrates,

especially in the presence of calcium signalling, which will imply a role in regulating PKAc activity. Given

the importance of cAMP and calcium signalling in parasite biology, the modulation of PKAc or PKAr

phosphorylation by PP7 is an exciting avenue for future research.

In conclusion, our findings underscore the multifaceted role of PP7 in *P. falciparum* asexual blood stage

development. The essentiality of PP7, its calcium-dependent phosphatase activity, and its interactions

with key signalling molecules, together with its role in some PKAc substrate phosphorylation in

schizonts, emphasize the importance of PP7 in the cell biology of the parasite. Our work opens new

avenues for research into the mechanisms underlying PP7 functions, its contribution to parasite survival,

and its potential as a target for anti-malarial drug development.

Acknowledgements

This work was also supported by funding from the Wellcome Trust to DAB (220318/Z/20/Z) and to MJB

from the Wellcome Trust (220318/A/20/Z) and the Francis Crick Institute (https://www.crick.ac.uk/)

which receives its core funding from Cancer Research UK (CC2129), the UK Medical Research Council

(CC2129), and the Wellcome Trust (CC2129).

 Data availability

221 The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

222 the PRIDE (35) partner repository (http://proteomecentral.proteomexchange.org) with the dataset

223 identifier PXD051277.

Materials and Methods

P. falciparum **culture and synchronisation**

P. falciparum erythrocytic stages were cultured in human erythrocytes (National Blood Transfusion

Service, UK) and RPMI 1640 medium (Life Technologies) supplemented with 0.5% Albumax type II

(Gibco), 50 μM hypoxanthine, and 2 mM L-glutamine at 37^0C and supplied with (gas composition).

Synchronous parasite cultures were obtained as described previously (36). Briefly, late segmented

schizonts were enriched by centrifugation on a 63% Percoll (GE Healthcare) cushion, followed by the

addition of fresh erythrocytes to allow invasion for 1–2 h with continuous shaking. Remaining schizonts

were then lysed by sorbitol treatment to yield highly synchronous ring-stage cultures. In all cases,

induction of DiCre activity when required was by treatment for 2–4 h with 100 nM rapamycin (RAP;

Sigma) as described previously (37, 38). Control parasites were treated with vehicle only (0.1% v/v

DMSO).

The PP7-HA:loxP line was generated from the DiCre-expressing 3D7 *P. falciparum* clone (23) using SLI

(39) with a plasmid containing a SERA2loxPint followed by a triple-HA tag and an in frame *Thosea asigna*

virus 2A (T2A) ribosomal skip peptide and NeoR cassette with a downstream loxP and PbDT 3⬛UTR

sequences as described previously (7). To ensure appropriate recombination to drive gene excision and

C-terminal tagging, a re-codonised version of the C-terminal portion of PP7, containing the phosphatase

catalytic domain and downstream EF hands, was synthesised commercially (IDT) and inserted

downstream of the SERA2loxPint and upstream of the 3×HA tag (Figure S4). The sequence of the ~800

bp upstream homology region is shown in Figure S4A.

Oligonucleotide primer sequences used in diagnostic PCR to detect integration and excision of

transgenes, and the sequences of re-codonised regions, are provided below in Tables 1 and S4.

PP7-HA:loxP mNeon was generated by transfection of PP7-HA:loxP with a construct targeting the p230p

locus. A linearised donor DNA fragment which inserted mNeon in-frame with a T2A peptide and BSD

selection marker when integrated, and a pDC2-based p230p targeting Cas9 gRNA plasmid was co-

transfected. Parasites were left to grow for two days post transfection followed by treatment with 5

µg/ml blasticidin (BSD) to select for integrants. After the emergence of BSD-resistant parasites gDNA

was screened for correct integration.

Parasite sample preparation and western blot

Parasite extracts were prepared from Percoll-purified schizonts treated with 0.15% w/v saponin to

remove erythrocyte material. To solubilise parasite proteins, PBS-washed saponin-treated parasite

pellets were resuspended in three volumes of NP-40 extraction buffer (10 mM Tris, 150 mM NaCl, 0.5

mM EDTA, 1% NP40, pH 7.5, with 1× protease inhibitors (Roche). Samples were gently vortexed and

incubated on ice for 10 min followed by centrifugation at 12,000g for 10 min at 4°C. For western blot,

SDS-solubilised proteins were electrophoresed on 4%-15% Mini-PROTEAN TGX Stain-Free Protein Gels

(Bio-Rad) under reducing conditions and proteins transferred to nitrocellulose membranes using a

semidry Trans-Blot Turbo Transfer System (Bio-Rad). Antibody reactions were carried out in 1% skimmed

milk in PBS with 0.1% Tween-20 and washed in PBS with 0.1% Tween-20. Appropriate horseradish

peroxide-conjugated secondary antibodies were used, and antibody-bound washed membranes were

incubated with Clarity Western ECL substrate (Bio-Rad) and visualised using a ChemiDoc (Bio-Rad).

Antibodies used for western blots presented in this work were as follows: anti-HA monoclonal antibody

(mAb) 3F10 (diluted 1:2,000) (Roche); mouse anti-GAPDH mAb (1:20,000).

Immunofluorescence assays

Thin blood films were fixed with 4% formaldehyde in PBS and permeabilised with PBS containing 0.1%

(v/v) Triton X-100. Blocking and antibody binding was performed in PBS 3% BSA w/v at room

temperature. Slides were mounted with ProLong Gold Antifade Mountant containing DAPI (Thermo

Fisher Scientific). Images were acquired with a NIKON Eclipse Ti fluorescence microscope fitted with a

Hamamatsu C11440 digital camera and overlaid in ICY bioimage analysis software. Antibodies used for

IFA were as follows: anti-HA monoclonal antibody (mAb) 3F10 (diluted 1:100) (Roche); rabbit anti-GAP45

mAb (1:200).

Flow cytometry

For growth assays, synchronous ring-stage parasites were adjusted to a 0.1% parasitaemia 1%

haematocrit suspension and dispensed in triplicate into six-well plates. Triplicate samples of 100 µL were

harvested at days 0, 2, 4 and 6 for each well and fixed with 4% formaldehyde 0.2% glutaraldehyde in

PBS. Fixed samples were stained with SYBR green and analysed by flow cytometry.

For the measurement of egress and ring formation of highly synchronized DMSO- and RAP-treated PP7-

HA:loxP parasites. A culture of PP7-HA:loxP with a 1 h invasion window was seeded in duplicate at 1%

hematocrit at 2% parasitaemia and treated with DMSO or RAP at 4 h post invasion for 2 h. Triplicate 100

μL samples were taken and fixed with 4% formaldehyde 0.2% glutaraldehyde in PBS at hourly intervals

from 45-53 h post invasion and at 69 h post invasion the subsequent day. Fixed samples were stained

with SYBR green and analysed by flow cytometry. Schizont parasitaemia was determine by gating high

signal SYBR positive cells. Ring parasitaemia was determined similarly but by gating low signal SYBR

positive cells.

Analysis of parasite development using Giemsa-stained samples

To analyse and measure the development of PP7-HA:loxP parasites, tightly synchronized schizonts were

allowed to re-invade fresh erythrocytes. At 4 hours post invasion, the rings were treated with DMSO or

rapamycin for 2 hours (cycle 0). At the indicated time point after invasion (cycle 0 and cycle 1), the

parasites were pelleted, smeared on a microscope slides, fixed with methanol and stained with Giemsa.

The parasites were then imaged and counted using an Olympus BX51 microscope equipped with an

Olympus SC30 camera and a 100x oil objective, controlled by CellSens software.

Merozoite number

The quantification of merozoites per schizont was conducted using Giemsa-stained smears of Percoll-

enriched mature parasites. The cells were imaged on an Olympus BX51 microscope equipped with an

Olympus SC30 camera and a 100x oil objective, controlled by cellSens software. Two independent

experiments were conducted with a minimum of 15 schizonts counted per experiment. Statistical

analysis was performed using GraphPad Prism v10.

Parasite invasion rate

Parasite invasion assays were performed as described previously in (40). Briefly, parasites treated with

either DMSO or rapamycin were diluted to a parasitaemia of 1% and a haematocrit of 2%. A 50 µl

starting sample (designated as H0) was collected and fixed (PBS solution containing 8%

paraformaldehyde, 0.01% glutaraldehyde). After a 4-hour incubation period, a second sample was

collected and fixed (designated as H4). The parasites were labelled with SYBR Green I (1:5000, Life

Technologies). The parasitaemia was quantified using an Attune cytometer (Thermofisher). Parasite

invasion rate was determined as the ratio of parasitaemia at H4 to H0. Each experiment was conducted

in triplicate, with a minimum of three biological replicates. Statistical analysis was performed using

GraphPad Prism v10.

Schizont DNA content analysis

Highly synchronous schizonts were fixed and labelled with SYBR Green I (1:5000, Life Technologies). The

fluorescence of SYBR Green was analysed using an Attune cytometer (Thermofisher) with the following

laser settings: forward scatter at 125 V, side scatter at 350V, and blue laser (BL1) 530:30 at 280V. For

each sample, at least 100,000 cells were analysed with three biological replicates performed in

triplicate. Data analysis was carried out using FlowJo software.

Parasite egress assay and time-lapse and live fluorescence microscopy.

Highly synchronous schizonts were isolated using Percoll centrifugation and incubated for 4 hours in

medium containing the PKG inhibitor C2 to arrest egress (1 µM, (38). DMSO-treated parasites were

treated with Hoechst to stain the nuclei and mixed with the unstained RAP-treated parasites. Prior to

analysis, the schizonts were resuspended without C2 in fresh serum-free RPMI at 37°C to allow egress

and immediately loaded onto a poly-L-lysine coated µ-slide VI 0.4 (Ibidi). The slides were transferred to a

Nikon Ti-E inverted microscope housed in a pre-warmed 37 °C chamber with an atmosphere of 5% CO_2.

Egress was imaged using a 63× oil immersion objective and an ORCA-Flash 4.0 CMOS camera

(Hamamatsu). DIC (differential inference contrast) images were taken at a rate of 1 frame/10 sec for 30

minutes, fluorescence images were taken every 2 min to avoid bleaching. Videos were acquired and

processed using Nikon NIS-Elements and Image J software.

Analysis of PP7 rings stage morphology

A culture of PP7-HA:loxP-mNeonGreen parasites with 1 h synchronicity was split and treated with DMSO

or RAP. Cultures were held pre-egress with 1 µM PKG inhibitor (C2) until 50 h post invasion at which

point C2 was removed by washing with pre-warmed complete medium. Cultures were left to invade

new host cells for 1 h with continuous shaking and staining with BODIPY-Tr ceramide (Thermo).

Subsequently a sample of each culture was taken. For Giemsa-stained blood film analysis samples were

smeared and fixed with methanol followed by Giemsa staining. For fluorescence microscopy, samples

were washed and diluted to 0.1% haematocrit, and placed into a chamber of a poly-L lysine coated Ibidi

chamber slide to adhere for 10 minutes at 37 degrees. Adherent cells were then imaged on a NIKON

Eclipse Ti fluorescence microscope fitted with a Hamamatsu C11440 digital camera and overlaid in ICY

bioimage analysis software.

Immuno-precipitation

Tightly synchronised schizonts (~45 h post invasion) of PP7-HA:loxP and 3D7DiCre parental parasites

were enriched on a 70% Percoll cushion. The schizonts were treated for 3 h with 1 µM C2 (to arrest

egress) after which the cultures were treated with 0.15% saponin in PBS containing cOmplete Mini

EDTA-free Protease and PhosSTOP Phosphatase inhibitor cocktails (both Roche) for 10 min at 4°C to lyse

the host erythrocytes. Samples were washed twice in PBS containing protease and phosphatase

inhibitors, snap-frozen and pellets stored at −80°C. Parasite pellets (70-100 µl packed volume) were

resuspended in three volumes of NP-40 extraction buffer (10 mM Tris, 150 mM NaCl, 0.5 mM EDTA, 1%

NP40, pH 7.5, with 1× protease inhibitors (Roche). Samples were gently vortexed and incubated on ice

for 10 min followed by centrifugation at 12,000g for 10 min at 4°C. Clarified lysates were then added to

anti-HA antibody-conjugated magnetic beads (Thermo Scientific) which had been equilibrated in NP-40

extraction buffer. Samples were incubated at room temperature for 2 h on a rotating wheel after which

beads were precipitated using a magnetic sample rack. The supernatant was removed, and beads

washed three times with NP-40 extraction buffer followed by three washes with extraction buffer

lacking detergent.

Mass spectrometry of immuno-precipitated material

Washed beads were resuspended in trypsinisation buffer (50 mM ammonium bicarbonate, 40 mM 2-

chloroacetamide and 10 mM Tris-(2-carboxyethyl) phosphine hydrochloride) and samples reduced and

alkylated by heated to 70°C for 5 minutes. 250 ng of trypsin was added to the samples and heated at

37°C overnight with gentle agitation followed by filtration using a 0.22 µm Costar® Spin-X® centrifuge

tube filter (Sigma). Samples were then run on a LTQ-Orbitrap-Velos mass spectrometer (Thermo

Scientific). Search engines, Mascot (http://www.matrixscience.com/) and MaxQuant

(https://www.maxquant.org/) were used for mass spectrometry data analysis. The PlasmoDB database

was used for protein annotation. Peptide and proteins having minimum threshold of 95% were used for

further proteomic analysis and peptide traces analysed using Scaffold5. Enrichment was determined by

comparing results from tagged lines with those of immuno-precipitated material from 3D7DiCre

parental parasites.

Phosphatase assay

Assays were carried out using the EnzChek Phosphatase Assay Kit (Therno) according to the manufacturer's procedures. Briefly, percoll purified schizont pellets of 50 µl were prepared for PP7-HA:loxP or 3D7DiCre parental parasites. Samples were processed by the methodology for immuno-precipitation. PP7-HA conjugated beads were then resuspended in a total volume of 650 µl of assay buffer (20 mM Tris, 150 mM NaCl, pH 7.5,). 50 µl of suspended bead slurry was dispensed in triplicate into wells of a 96-well plate for each test condition. 50 µl of reaction buffer containing additives for the conditions tested and difluoro-4-methylumbelliferyl phosphate (DiFMUP) substrate were added, to give a final concentration of 100 µM DiFMUP. A set of control wells were made with the addition of reaction buffer lacking DiFMUP. Plates were incubated at 37°C for 1 h and then read with Ex/Em of 358/455 nm on a spectramax i5 plate reader (Molecular Devices). Averaged arbitrary fluorescence for each condition was calculated by normalising to the negative control wells.

Phosphoproteomics

The phosphoproteomics data presented are from an isobaric labelling experiment. Tightly synchronised, ring-stage PP7-HA:loxP were treated with 100 nM RAP or vehicle only (DMSO) and schizonts (about 45 h old) enriched on an approximately 70% Percoll cushion. The schizonts were treated for 2 h with 1 µM C2 (to arrest egress) and then washed to allow egress for 25 min, after which the cultures were treated with 0.15% saponin in PBS containing cOmplete Mini EDTA-free Protease and PhosSTOP Phosphatase inhibitor cocktails (both Roche) for 10 min at 4°C to lyse the host erythrocytes. Samples were washed twice in PBS containing protease and phosphatase inhibitors, snap-frozen in liquid nitrogen, and pellets stored at –80°C. Parasite pellets were resuspended in 1 mL 8 M urea in 50 mM HEPES, pH 8.5, containing protease and phosphatase inhibitors and 100 U/mL benzonase (Sigma). Proteins were extracted from

the pellets using three 15 s bursts with a probe sonicator followed by a 10 min incubation on ice and a

30 min centrifugation at 14,000 rpm at 4°C. Protein content was estimated by a BCA protein assay and a

200 μg aliquot of each sample was taken for further processing. Samples were reduced with 10 mM

dithiothreitol for 25 min at 56°C and then alkylated with 20 mM iodoacetamide for 30 min at room

temperature. The alkylation reaction was quenched with an additional 10 mM dithiothreitol, and then

each sample was diluted with 50 mM HEPES to reduce the urea concentration to <2 M prior to

digestion. Proteolytic digestion was carried out by the addition of 4 μg LysC (WAKO) and incubated at

37°C for 2.5 h followed by the addition of 10 μg trypsin (Pierce) and overnight incubation at 37°C. After

acidification, C18 MacroSpin columns (Nest Group) were used to clean up the digested peptide solutions

and the eluted peptides dried by vacuum centrifugation. Samples were resuspended in 50 mM HEPES

and labelled using the 0.8 mg Tandem Mass Tag 10plex isobaric reagent kit (Thermo Scientific)

resuspended in acetonitrile. Labelling reactions were quenched with hydroxylamine, and a pool was

made of each set of samples. Acetonitrile content was removed from the pooled TMT solution by

vacuum centrifugation and then acidified before using a Sep-Pak C18 (Waters) to clean up the labelled

peptide pool prior to phosphopeptide enrichment. The eluted TMT-labelled peptides were dried by

vacuum centrifugation and phosphopeptide enrichment was subsequently carried out using the

sequential metal oxide affinity chromatography (SMOAC) strategy with High Select TiO_2 and Fe-NTA

enrichment kits (Thermo Scientific). Eluates were combined prior to fractionation with the Pierce High

pH Reversed-Phase Peptide Fractionation kit (Thermo Scientific). The dried TMT-labelled

phosphopeptide fractions generated were resuspended in 0.1% TFA for LC-MS/MS analysis using a

U3000 RSLCnano system (Thermo Scientific) interfaced with an Orbitrap Fusion Lumos (Thermo

Scientific). Each peptide fraction was pre-concentrated on an Acclaim PepMap 100 trapping column

before separation on a 50-cm, 75-μm I.D. EASY-Spray Pepmap column over a 3-h gradient run at 40°C,

eluted directly into the mass spectrometer. The instrument was run in data-dependent acquisition mode

with the most abundant peptides selected for MS/MS fragmentation. Two replicate injections were

made for each fraction with different fragmentation methods based on the MS2 HCD and MSA SPS MS3

strategies described. The acquired raw mass spectrometric data were processed in MaxQuant (version

1.6.2.10) for peptide and protein identification; the database search was performed using the

Andromeda search engine against the Homo sapiens canonical sequences from UniProtKB (release

2018_05) and *P. falciparum* 3D7 sequences from PlasmoDB (18). Fixed modifications were set as

Carbamidomethyl (C) and variable modifications set as Oxidation (M) and Phospho (STY). The estimated

false discovery rate was set to 1% at the peptide, protein, and site levels. A maximum of two missed

cleavages were allowed. Reporter ion MS2 or Reporter ion MS3 was appropriately selected for each raw

file. Other parameters were used as preset in the software. The MaxQuant output file PhosphoSTY

Sites.txt, an FDR-controlled site-based table compiled by MaxQuant from the relevant information about

the identified peptides, was imported into Perseus (v1.4.0.2) for data evaluation.

For a phosphorylation site to be considered regulated, the following cut-offs were applied: P-value

<⬚0.05, Welch difference >⬚0.7 or < -0.7 and localisation probability >0.7. Significantly hyper- and hypo-

phosphorylated proteins were used for GO enrichment analysis. GO IDs were extracted from the *P.*

falciparum annotation file (PlasmoDB.org). Enriched GO terms were identified as described before

(PMID: 25011111). Plots were made using R package ggplot2 (https://ggplot2.tidyverse.org).

Figure legends

Figure 1

(1A) Cartoon of the predicted *P. falciparum* PP7 protein showing the positions of the individual domains.

Amino acid numbers are shown in black. The arrow indicates the point at which the protein product is

truncated when the modified locus is excised in the transgenic parasites.

(1B) Schematic representation of the SLI strategy (39) used to produce the PP7-HA:loxP DiCre line and

resultant RAP-induced disruption of the modified gene. Double-headed arrows represent the regions

amplified by PCR in (1C). Red arrowheads, *loxP* sites, yellow lollipops, translational stop codons, white

box, phosphatase domain, light blue box, regions of re-codonised sequence (R.R.).

(1C) Diagnostic PCR analysis of genomic DNA (gDNA) from a transgenic PP7 parasite line verifying

successful modification of target loci by SLI to produce PP7-HA:loxP. Efficient excision of 'floxed'

sequences is observed upon treatment with RAP. Track C represents amplification of a control locus

(PKAr) to check gDNA integrity. PCRs 1-4 are represented in the schematic locus in panel 1B. PCR 1

screens for the WT locus, PCR 2 for 5' integration, PCR 3 for the excision of the 'floxed' sequence and

PCR 4 for 3' integration. See Table 1 for sequences of all primers used for PCR. Sizes for expected

amplification products are as follows: C, control locus (primers 5 and 6) 836 b.p. PCR 1 (primers 7 and 8)

1296 b.p, PCR2 (primers 7 and 9) 3559 b.p, PCR3 (primers 7 and 10) 1362 b.p. (RAP) and PCR 4 (primers

11 and 8) 1013 b.p.

(1D) Western blot analysis of expression (DMSO, control) and ablation (RAP) of PP7-HA from highly

synchronous mature schizonts. Expression of GAPDH (PF3D7_1462800) is shown as a loading control.

(1E) IFA analysis showing the diffuse peripheral localisation of PP7-HA and the loss of expression upon

RAP treatment (16 h post invasion). Over 99% of all RAP-treated PP7-HA:loxP schizonts examined by IFA

showed diminished HA expression in three independent experiments. Scale bar, 2 μm.

Figure 2

Growth curves showing parasitaemia as measured by flow cytometry of PP7-HA:loxP parasites treated

with DMSO (vehicle only control, blue) or RAP (red). Means from three independent biological replicates

are plotted. Error bars, SD. Inset, Giemsa-stained thin blood films showing ring-stage parasites following

egress of synchronous DMSO- and RAP-treated PP7-HA:loxP schizonts. Ring formation occurs in DMSO-

treated PP7-HA parasites, parasites did not develop beyond the early ring stage in RAP-treated

parasites. Student's t test for the comparison of conditions between DMSO and RAP treatments, *

:P<0.05,*** :P<0.001.

Figure 3

(3A) Western blot showing immunoprecipitation (IP) of PP7-HA from schizont extracts. Black arrow

indicates the predicted mass of PP7-HA. Minor lower molecular weight degradation products are seen.

Images are representative of two independent biological repeats.

(3B) Phosphatase assay of immuno-precipitated PP7-HA conjugated beads. PP7 conjugated bead

conditions are shown in green and control beads from 3D7 DiCre parental schizont lysate immuno-

precipitation are displayed in blue. Results are averages of three independent biological repeats. Error

bars, SD, * :P<0.05, ** :P<0.01, *** :P<0.001, Student's t test for the comparison between conditions.

Figure 4

(4A) Mass spectrometric identification of interacting partners of PP7. The right panel shows immuno-

precipitation with no added Ca^{2+} ions, left panel shows immuno-precipitation in the presence of

additional 2 mM Ca^{2+} (present during detergent extraction, immuno-precipitation and washes). Volcano

plot of P values versus the corresponding log2 fold change in abundance compared to 3D7DiCre control

samples (Fischer's exact test). Plotted by analysing proteins enriched through IP (panel 3A) by mass

spectrometry. Green line indicates p=-2log10 and green dots represent peptides where p<-2log10.

Peptides for PP7 were enriched to p<-19log10.

(4B) Western blot analysis of immuno-precipitated PP7:HA-loxP schizont lysates with anti-HA beads,

using the HA antibody (upper), the CDPK1 antibody (middle) and the GAPDH antibody (lower) (used as a

negative control cytosolic marker to detect potential non-specific binding).

(4C) Volcano plot showing the changes in detection of phospho-sites between DMSO- and RAP-treated

PP7-HA:loxP. The negative log10 transform of the p-value–derived Welch-corrected t test comparing

five DMSO- and five RAP-treated replicates is plotted against the log2-transformed fold change in

reporter ion intensity (DMSO/RAP). Significance of $p < 0.05$ is denoted by the horizontal line. Vertical

lines indicate −0.7 and 0.7 log2 fold change. Light blue circles correspond to significantly

hypophosphorylated peptides and red circles to significantly hyperphosphorylated peptides.

Figure 5

(5A) Giemsa-stained thin blood films showing development of PP7-HA:loxP parasites treated with DMSO

or rapamycin during cycle 0 at 4 h post invasion for 2 h. Scale bars, 5 µm.

(5B), Microscopic quantification of parasite developmental stages at each time point. Rapamycin-treated

parasites displayed an accumulation of pycnotic and abnormal looking parasites at early life cycle stage.

Counts are means of results of two independent experiments. Rapamycin treated parasites develop

normally during the first cycle and are able to re-invade. Early on during the next cycle, an accumulation

of abnormal rings and pycnotic forms appear, the parasites did not develop beyond this cycle.

(5C) Flow cytometry analysis of ring formation in DMSO- and RAP-treated PP7-HA:loxP parasites treated

in the previous cycle. Samples from highly-synchronized cultures treated at the ring stage the cycle prior

to invasion and analysis, were taken in triplicate at the stated time intervals post invasion and stained

with the DNA stain SYBR green. Samples were analyzed by flow cytometry, and the schizont and ring

parasitemia values determined by gating high-signal and low-signal SYBR-positive cells, respectively.

Mean parasitemia values (starting schizontemia adjusted to 2%) from two independent experiments are

plotted. Error bars, SD; Student's t test for the comparison of conditions between DMSO and RAP

treatments (ns: nonsignificant). Insets are smears taken from cultures at 24h post invasion. Scale bars,

5 µm.

Figure 6

PP7 ablation results in arrest of ring-stage parasite development. (A) Tightly synchronized DMSO- or RAP-treated PP7-HA:loxP parasites were arrested with C2 block and subsequently washed and left to invade in fresh pre-warmed medium. After a further 2 h, thin blood smears were taken, and Giemsa stained. Ring-stage parasite morphology was classified by manual counting. A minimum of 100 cells were counted per replicate and scored for morphology, scale bar 2 μm. (B) Tightly synchronized DMSO- or RAP-treated PP7-HA-Cyto-mNeon:loxP parasites were arrested with C2 and subsequently washed and left to invade in fresh pre-warmed medium. Parasites were then stained with BODIPY TR ceramide and Hoechst 3342 to visualise membranes and nuclei respectively. Parasite samples were placed in an Ibid chamber slide and imaged by fluorescence microscopy. A minimum of 40 ring-stage cells were counted per condition and scored for ameboid or pycnotic morphology of parasites, scale bar 2 μm. Mean data are plotted for three independent experiments. Error bars, SD, ** :P<0.01, **** :P<0.0001, Student's t test for the comparison of conditions between DMSO and RAP treatments.

Table 1-primers used in this study

Number	Primer name	Sequence 5'-3'
1	PP7 5HR F	GCGGCCGCAGATCTCTCGAGGGAAGAAAGCGCCTGTACTATG
2	PP7 5HR R	CGAAGTTATTGTATATTATTTTTTTTTATTTACAATTAATTTGGTATCTTTGGACTTTTTG
3	PP7 Reco F	GTATATATATATATATTTATATATTTTATATTCTTTTAGATATTAGGAGACGTTCATGGACAG
4	PP7 Reco R	CAGATCCGCCTGAACCGGATCCATTATTGCTGTATATATATGTAGGTTTG
5	PKAr F	GCGGCCGCAGATCTCTCGAGCCTAAAGTTTATAAGAAAGATGAAAAAG
6	PKAr R	CAGATCCGCCtGAACCGGATCCATTTTCATCAATACAAGTTGTATCCAAAC
7	5' screen F PP7	CAACATTGAAGATGTTGTTATGATATATG
8	3' screen R PP7	GGTGAAATATATTCAGTCGATGC
9	HA R	GCATAGTCAGGAACATCGTAAGG
10	screen L7 myc R	CCGTTCAAATCTTCTTCAGAAATCAAC

11	pL7 3' screen F	CAGCTATGACCATGATTACGCC
12	p230p Int screen F	CTATATGGTATCCAAAACCTTTAAATTATATAGC
13	p230p WT screen R	GAGGAATTTTTAAATATGATATACCTTTATCATTAG
14	p230p 5 Int screen R	CCAATGGCCCCTTTCCGC
15	p230p 3 Int screen F	CTAGCGGCTTGTCGACG

References

1. Miranda-Saavedra D, Gabaldón T, Barton GJ, Langsley G, Doerig C. 2012. The kinomes of
apicomplexan parasites. Microbes Infect 14:796–810.

2. Ward P, Equinet L, Packer J, Doerig C. 2004. Protein kinases of the human malaria parasite
Plasmodium falciparum: The kinome of a divergent eukaryote. BMC Genomics 5.

3. Wilkes JM, Doerig C. 2008. The protein-phosphatome of the human malaria parasite Plasmodium
falciparum. BMC Genomics 9.

4. Guttery DS, Poulin B, Ramaprasad A, Wall RJ, Ferguson DJP, Brady D, Patzewitz EM, Whipple S,
Straschil U, Wright MH, Mohamed AMAH, Radhakrishnan A, Arold ST, Tate EW, Holder AA,
Wickstead B, Pain A, Tewari R. 2014. Genome-wide functional analysis of plasmodium protein
phosphatases reveals key regulators of parasite development and differentiation. Cell Host
Microbe 16:128–140.

5. Tewari R, Straschil U, Bateman A, Böhme U, Cherevach I, Gong P, Pain A, Billker O. 2010. The
systematic functional analysis of Plasmodium protein kinases identifies essential regulators of
mosquito transmission. Cell Host Microbe 8:377–387.

6. Dvorin JD, Martyn a C, Patel SD, Grimley JS, Collins CR, Hopp CS, Bright AT, Westenberger S,
Winzeler E, Blackman MJ, Baker DA, Wandless TJ, Duraisingh MT. 2010. A plant-like kinase in
Plasmodium falciparum regulates parasite egress from erythrocytes. Science 328:910–2.

7. Patel A, Perrin AJ, Flynn HR, Bisson C, Withers-Martinez C, Treeck M, Flueck C, Nicastro G, Martin
SR, Ramos A, Gilberger TW, Snijders AP, Blackman MJ, Baker DA. 2019. Cyclic AMP signalling
controls key components of malaria parasite host cell invasion machinery. PLoS Biol 17.

8.   Ganter M, Goldberg JM, Dvorin JD, Paulo JA, King JG, Tripathi AK, Paul AS, Yang J, Coppens I, Jiang
567   RHY, David A. 2017. Plasmodium falciparum CRK4 directs continuous rounds of DNA replication
568   during schizogony 1–48.

570   9.   Kumar S, Kumar M, Ekka R, Dvorin JD, Paul AS, Madugundu AK, Gilberger T, Gowda H, Duraisingh
571   MT, Keshava Prasad TS, Sharma P. 2017. PfCDPK1 mediated signaling in erythrocytic stages of
572   Plasmodium falciparum. Nat Commun 8.

574   10.   Koussis K, Withers-Martinez C, Baker DA, Blackman MJ. 2020. Simultaneous multiple allelic
575   replacement in the malaria parasite enables dissection of PKG function. Life Sci Alliance 3.

577   11.   Davies H, Belda H, Broncel M, Ye X, Bisson C, Introini V, Dorin-Semblat D, Semblat JP, Tibúrcio M,
578   Gamain B, Kaforou M, Treeck M. 2020. An exported kinase family mediates species-specific
579   erythrocyte remodelling and virulence in human malaria. Nat Microbiol 5:848–863.

581   12.   Fréville A, Gnangnon B, Khelifa AS, Gissot M, Khalife J, Pierrot C. 2022. Deciphering the Role of
582   Protein Phosphatases in Apicomplexa: The Future of Innovative Therapeutics? Microorganisms
583   10.

585   13.   Paul AS, Saha S, Engelberg K, Jiang RHY, Coleman BI, Kosber AL, Chen C-T, Ganter M, Espy N,
586   Gilberger TW, Gubbels M-J, Duraisingh MT. 2015. Parasite Calcineurin Regulates Host Cell
587   Recognition and Attachment by Apicomplexans. Cell Host Microbe 18:49–60.

589   14.   Philip N, Waters AP. 2015. Conditional Degradation of Plasmodium Calcineurin Reveals Functions
590   in Parasite Colonization of both Host and Vector. Cell Host Microbe 18:122–131.

592   15.   Paul AS, Miliu A, Paulo JA, Goldberg JM, Bonilla AM, Berry L, Seveno M, Braun-Breton C, Kosber
593   AL, Elsworth B, Arriola JSN, Lebrun M, Gygi SP, Lamarque MH, Duraisingh MT. 2020. Co-option of
594   Plasmodium falciparum PP1 for egress from host erythrocytes. Nat Commun 11.

596   16.   Zhu X, Li S, Wang C, Yu Y, Wang J, He L, Siddiqui FA, Chen L, Zhu L, Zhou D, Qin J, Miao J, Cui L,
597   Cao Y. 2022. The Plasmodium falciparum Nuclear Protein Phosphatase NIF4 Is Required for
598   Efficient Merozoite Invasion and Regulates Artemisinin Sensitivity. mBio 13.

17. Flueck C, Drought LG, Jones A, Patel A, Perrin AJ, Walker EM, Nofal SD, Snijders AP, Blackman MJ,
Baker DA. 2019. Phosphodiesterase beta is the master regulator of cAMP signalling during
malaria parasite invasion. PLoS Biol 17:e3000154.

18. Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, Gao X, Gingle A, Grant G,
Harb OS, Heiges M, Innamorato F, Iodice J, Kissinger JC, Kraemer E, Li W, Miller JA, Nayak V,
Pennington C, Pinney DF, Roos DS, Ross C, Stoeckert CJ, Treatman C, Wang H. 2009. PlasmoDB: a
functional genomic database for malaria parasites. Nucleic Acids Res 37.

19. Kamada R, Kudoh F, Ito S, Tani I, Janairo JIB, Omichinski JG, Sakaguchi K. 2020. Metal-dependent
Ser/Thr protein phosphatase PPM family: Evolution, structures, diseases and inhibitors.
Pharmacol Ther 215.

20. Jang DJ, Ban B, Lee JA. 2011. Characterization of Novel Calmodulin Binding Domains within IQ
Motifs of IQGAP1. Mol Cells 32:511.

21. Heizmann CW. 2019. Ca2+-Binding Proteins of the EF-Hand Superfamily: Diagnostic and
Prognostic Biomarkers and Novel Therapeutic Targets. Methods Mol Biol 1929:157–186.

22. Steele FR, Washburn T, Rieger R, O'Tousa JE. 1992. Drosophila retinal degeneration C (rdgC)
encodes a novel serine/threonine protein phosphatase. Cell 69:669–676.

23. Knuepfer E, Napiorkowska M, Van Ooij C, Holder AA. 2017. Generating conditional gene
knockouts in Plasmodium - a toolkit to produce stable DiCre recombinase-expressing parasite
lines using CRISPR/Cas9. Sci Rep 7.

24. Toenhake CG, Fraschka SAK, Vijayabaskar MS, Westhead DR, van Heeringen SJ, Bártfai R. 2018.
Chromatin Accessibility-Based Characterization of the Gene Regulatory Network Underlying
Plasmodium falciparum Blood-Stage Development. Cell Host Microbe 23:557-569.e9.

25. Grüring C, Heiber A, Kruse F, Ungefehr J, Gilberger TW, Spielmann T. 2011. Development and
host cell modifications of Plasmodium falciparum blood stages in four dimensions. Nat Commun
2.

634 26. Liang QL, Nie LB, Li TT, Elsheikha HM, Sun LX, Zhang ZW, Zhao DY, Zhu XQ, Wang JL. 2022.
635   Functional Characterization of 17 Protein Serine/Threonine Phosphatases in Toxoplasma gondii
636   Using CRISPR-Cas9 System. Front Cell Dev Biol 9:738794.

638 27. Brochet M, Billker O. 2016. Calcium signalling in malaria parasites. Mol Microbiol 00:1–12.

640 28. Brochet M, Collins MO, Smith TK, Thompson E, Sebastian S, Volkmann K, Schwach F, Chappell L,
641   Gomes AR, Berriman M, Rayner JC, Baker DA, Choudhary J, Billker O. 2014. Phosphoinositide
642   Metabolism Links cGMP-Dependent Protein Kinase G to Essential Ca2+ Signals at Key Decision
643   Points in the Life Cycle of Malaria Parasites. PLoS Biol 12.

645 29. Furuyama W, Enomoto M, Mossaad E, Kawai S, Mikoshiba K, Kawazu SI. 2014. An interplay
646   between 2 signaling pathways: Melatonin-cAMP and IP3–Ca2+ signaling pathways control
647   intraerythrocytic development of the malaria parasite Plasmodium falciparum. Biochem Biophys
648   Res Commun 446:125–131.

650 30. Beraldo FH, Almeida FM, Da Silva AM, Garcia CRS. 2005. Cyclic AMP and calcium interplay as
651   second messengers in melatonin-dependent regulation of Plasmodium falciparum cell cycle. J
652   Cell Biol 170:551.

654 31. Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner IF, Casandra D,
655   Mayho M, Brown J, Li S, Swanson J, Rayner JC, Jiang RHY, Adams JH. 2018. Uncovering the
656   essential genes of the human malaria parasite *Plasmodium falciparum* by saturation
657   mutagenesis. Science (1979) 360.

659 32. Neye H. 2007. Peptidyl - Prolyl Isomerases. xPharm: The Comprehensive Pharmacology Reference
660   1.

662 33. Jo GA, Lee JM, No G, Kang DS, Kim SH, Ahn SH, Kong IS. 2015. Isolation and characterization of a
663   17-kDa FKBP-type peptidyl-prolyl cis/trans isomerase from Vibrio anguillarum. Protein Expr Purif
664   110:130–137.

666 34. Dziurdzik SK, Conibear E. 2021. The Vps13 Family of Lipid Transporters and Its Role at Membrane
667   Contact Sites. Int J Mol Sci 22:2905.

35.   Deutsch EW, Bandeira N, Perez-Riverol Y, Sharma V, Carver JJ, Mendoza L, Kundu DJ, Wang S,
Bandla C, Kamatchinathan S, Hewapathirana S, Pullman BS, Wertz J, Sun Z, Kawano S, Okuda S,
Watanabe Y, MacLean B, MacCoss MJ, Zhu Y, Ishihama Y, Vizcaíno JA. 2023. The
ProteomeXchange consortium at 10 years: 2023 update. Nucleic Acids Res 51:D1539–D1548.

36.   Harris PK, Yeoh S, Dluzewski AR, O'Donnell RA, Withers-Martinez C, Hackett F, Bannister LH,
Mitchell GH, Blackman MJ. 2005. Molecular identification of a malaria merozoite surface
sheddase. PLoS Pathog 1:0241–0251.

37.   Jones ML, Das S, Belda H, Collins CR, Blackman MJ, Treeck M. 2016. A versatile strategy for rapid
conditional genome engineering using loxP sites in a small synthetic intron in Plasmodium
falciparum. Sci Rep 6:21800.

38.   Collins CR, Das S, Wong EH, Andenmatten N, Stallmach R, Hackett F, Herman JP, Müller S,
Meissner M, Blackman MJ. 2013. Robust inducible Cre recombinase activity in the human malaria
parasite Plasmodium falciparum enables efficient gene deletion within a single asexual
erythrocytic growth cycle. Mol Microbiol 88:687–701.

39.   Birnbaum J, Flemming S, Reichard N, Soares AB, Mesén-Ramírez P, Jonscher E, Bergmann B,
Spielmann T. 2017. A genetic system to study Plasmodium falciparum protein function. Nat
Methods 14:450–456.

40.   Mohring F, Rawlinson T, Draper S, Moon R. 2020. Multiplication and Growth Inhibition Activity
Assays for the Zoonotic Malaria Parasite, Plasmodium knowlesi. Bio Protoc 10.

(A) PP7-HA IP

	Input	Unbound	Wash	Beads
250 kDa				
150 kDa				
100 kDa				← ~112KDa
75 kDa				
50 kDa				
37 kDa				
25 kDa				

(B) Phosphatase activity of immuno-precipitated schizont extracts

(A)

No added Ca²⁺

2mM Ca²⁺

PF3D7_1423300 (PP7)

PF3D7_1434200 (Calmodulin)
PF3D7_0423300 (Conserved Protein)

PF3D7_0217500 (CDPK1)

(B)

αHA

αCDPK1

αGAPDH

(C)

P. falciparum schizonts
PP7 RAP/PP7 DMSO

(A)

Cycle 0 | Cycle 1

5hpi | 24hpi | 30hpi | 48hpi | 6hpi

DMSO

RAP

(B)

Parasite stage (%)

5hpi D | 5hpi R | 24 hpi D | 24 hpi R | 30 hpi D | 30 hpi R | 48 hpi D | 48 hpi R | 6hpi D | 6hpi R

Cycle 0 | Cycle 1

- Ring
- Trophozoites
- Late Trophozoites
- Schizonts
- Pycnotic
- Abnormal Rings

(C)

PP7 invasion time course

DMSO | RAP

% Parasitaemia

Hours: 0, 2, 4, 24

ns

- DMSO Schizonts
- DMSO Rings/Pycnotic/Abnormal rings
- RAPA Schizonts
- RAPA Rings /Pycnotic/Abnormal rings

(A)

Ring like | Dot like | Blob like | Wisp like

Percentage of parasites

ring | dot | blob | wisp

■ DMSO 2 h □ RAP 2 h

(B)

Percentage of parasites

DMSO 2 h | RAP 2 h

■ Ameboid □ Pycnotic