Monoclonal antibodies derived from B cells in subjects with cystic fibrosis reduce

***Pseudomonas aeruginosa* burden in mice**

Malika Hale[1*], Kennidy K. Takehara[2*], Christopher D. Thouvenel[1], Dina A. Moustafa[3], Andrea Repele[1], Mary F. Fontana[2], Jason Netland[2], Sharon McNamara[4], Ronald L. Gibson[4,5], Joanna B. Goldberg[3], David J. Rawlings[1,2,5,6] and Marion Pepper[2, 6]

*These authors contributed equally.

Affiliations

[1]Center for Immunity and Immunotherapies, Seattle Children's Research Institute, Seattle, WA

[2]Department of Immunology, University of Washington School of Medicine, Seattle, WA

[3]Division of Pulmonary, Asthma, Cystic Fibrosis, and Sleep, Department of Pediatrics, Emory University School of Medicine, Atlanta, GA

[4]Cystic Fibrosis Center, University of Washington/Seattle Children's Hospital, Seattle, WA

[5]Department of Pediatrics, University of Washington School of Medicine, Seattle, WA

[6]Address correspondence to: mpepper@uw.edu and drawling@uw.edu

Abstract:

Pseudomonas aeruginosa (PA) is an opportunistic, frequently multidrug-resistant pathogen that can cause severe infections in hospitalized patients. Antibodies against the PA virulence factor, PcrV, protect from death and disease in a variety of animal models. However, clinical trials of PcrV-binding antibody-based products have thus far failed to demonstrate benefit. Prior candidates were derivations of antibodies identified using protein-immunized animal systems and required extensive engineering to optimize binding and/or reduce immunogenicity. Of note, PA infections are common in people with cystic fibrosis (pwCF), who are generally believed to mount normal adaptive immune responses. Here we utilized a tetramer reagent to detect and isolate PcrV-specific B cells in pwCF and, via single-cell sorting and paired-chain sequencing, identified the B cell receptor (BCR) variable region sequences that confer PcrV-specificity. We derived multiple high affinity anti-PcrV monoclonal antibodies (mAbs) from PcrV-specific B cells across 3 donors, including mAbs that exhibit potent anti-PA activity in a murine pneumonia model. This robust strategy for mAb discovery expands what is known about PA-specific B cells in pwCF and yields novel mAbs with potential for future clinical use.

Introduction

Pseudomonas aeruginosa (PA) is a gram-negative bacteria responsible for significant morbidity and mortality in vulnerable individuals. Treatment is challenging because of intrinsic and acquired antibiotic resistance to most antibiotic drug classes. PA is one of the most common pathogens in severe healthcare associated infections (Mancuso et al., 2021; Weiner-Lastinger et al., 2020). Due to the significant mortality caused by multi-drug resistant strains and the lack of alternative therapies, new anti-pseudomonal medicines are urgently needed.

Monoclonal antibodies (mAbs) that bind to key PA virulence factors, such as PcrV, have shown promise in animal models. PcrV is a 28kDa surface protein that forms the distal tip of the type III secretion system required for toxin injection into host cells (Sato & Frank, 2011). A humanized anti-PcrV antibody fragment (Fab) (Baer et al., 2009; François et al., 2012; Frank et al., 2002) and a bi-specific antibody containing an anti-PcrV binding moiety, each originally derived from antibody sequences elicited by immunizing mice with purified recombinant PcrV (Chastre et al., 2022; Warrener et al., 2014) have demonstrated safety in ventilated patients at high risk for PA pneumonia, but failed to achieve the targeted efficacy outcomes in Phase 2 trials. In prior studies of Covid-19 and malaria, our groups have generated potent anti-pathogen mAbs by sequencing the variable regions of the B cell receptor (BCR) derived from antigen-specific memory B cells (MBCs) that arise following natural infection in humans (Hale et al., 2022; Rodda et al., 2021; Thouvenel et al., 2021). Therefore, we hypothesized that protective anti-PcrV monoclonal antibodies could be derived from B cells in individuals previously infected with PA.

People with cystic fibrosis (pwCF) experience intermittent or chronic airway PA infections (Cramer et al., 2023; Lund-Palau et al., 2016). Intermittent infections early in

childhood eventually progress to a state of persistent, chronic airway colonization. CF is a multi-organ disease caused by mutations in the gene encoding cystic fibrosis transmembrane conductance regulator (CFTR), an apical epthelial ion channel involved in chloride and bicarbonate transport (Shteinberg et al., 2021). Several mechanisms related to the biology of the CF airway have been proposed to explain susceptibility to PA (Rossi et al., 2020). Importantly, pwCF do not have intrinsic deficits in adaptive immunity (Alicandro et al., 2023; Ong & Ramsey, 2023). Class-switched anti-PcrV antibodies are present in CF serum and sputum, at higher concentrations than the general population (Mauch & Levy, 2014; Moss et al., 2001). However, the PcrV-specific B cells that give rise to antibody-secreting cells had not been previously studied. Here, we show that pwCF have more PcrV-specific circulating B cells compared to non-CF controls. Building on this finding, we utilized single-cell sequencing to identify PcrV-specific paired-chain BCR sequences in pwCF. From BCR sequences derived from PcrV-specific B cells and MBCs, we generated novel anti-PcrV mAbs, including several that confer robust protection against a virulent strain of PA in an *in vivo* pneumonia challenge model.

Results

PcrV-specific B cells are enriched in CF individuals

We previously described techniques to generate a fluorophore-linked tetramers to isolate antigen-specific B cells (Krishnamurty et al., 2016). To identify PcrV-specific B cells, we generated a tetramer using recombinant PcrV protein. In parallel, we also generated a decoy tetramer consisting of irrelevant proteins linked to a tandem fluorophore (Figure 1A). To test the tetramer reagent, we immunized C57BL/6 mice with 250 μg purified, recombinant PcrV protein

resuspended in Sigma Adjuvant System (SAS, Sigma). Immunization induced an expanded population of class-switched PcrV[+] B cells on day 7, demonstrating the specificity of the reagent (Taylor et al., 2012) (Figure 1–figure supplement 1).

We next obtained peripheral blood samples from a cohort of 14 young adults with CF who were followed in CF clinic at Seattle Children's Hospital. As B cells specific for any single antigen are rare, tetramer-bound cells were enriched on an anti-fluorophore magnetic column prior to analysis. After enrichment, PcrV-specific cells made up $\geq$ 5% of B cells in 10 out of 14 (71%) of CF donors and demonstrated significant expansion in cell number over non-CF individuals (Figure 1B). Similar differences between CF *vs.* non-CF donors were found when the number of PcrV-specific B cells in each sample was normalized to lymphocyte count (Figure 1C).

Generation of mAbs from IgM[+] B cells derived from a CF donor with PA

We next sought to isolate and sequence B cell receptors from PcrV-specific B cells in order to generate monoclonal antibodies. PBMCs were isolated from a blood sample from a CF donor (Donor 1), who was chronically infected with PA based on at least 2 PA+ oropharyngeal cultures in 3 consecutive years collected as part of routine care (Green et al., 2012; Mogayzel et al., 2014; Rosenfeld et al., 2022) (see Table 1 for clinical characteristics). Following single cell sorting of PcrV-specific cells (Figure 2–figure supplement 2A), we performed paired chain sequencing of the heavy and light chain variable regions. Our FACS sorting method allows single-cell surface phenotyping data to be linked to the BCR sequence for individual cells (Krishnamurty et al., 2016). Interestingly, only two PcrV-specific B cells sequenced in Donor 1 expressed the canonical MBC surface markers (CD21[+]CD27[+]) (gating strategy shown in Figure 2–figure supplement 2B). Consistent with these observations, BCR sequencing of PcrV-specific

B cells revealed predominantly germline sequences with little or no evidence for somatic hypermutation (Figure 2A).

We next generated mAbs that employed the BCR variable regions from 10 tetramer-bound B cells isolated from Donor 1. As the surface-expressed isotype of all these PcrV-specific B cells was IgM (Figure 2–figure supplement 2C), we first expressed the BCR sequences as pentameric IgM antibodies. When tested for binding to recombinant PcrV in a plate-bound ELISA assay, supernatants collected from cells co-transfected with heavy, light, and J chain plasmids (to make pentameric IgM) exhibited strong binding to PcrV for 3 of 10 CF-derived BCRs (denoted BCRs 408, 411, and 421) (Figure 2B). The light chain for BCR 421 contains several mutations in CDR1. Despite having germline sequences, mAbs from clones 408 and 411 had high affinity as measured by ELISA. As clinical application would likely require large-scale production in a similar system, we chose to move forward with the best binders that also were most efficiently expressed in the 293T cell line: 408 and 411 (data not shown).

We have previously found that for mAbs that target divergent pathogens, including SARS-CoV-2 and *Plasmodium falciparum*, multimerized antibodies (eg. pentameric IgM) have enhanced binding and protection properties *in vitro* (Hale et al., 2022; Thouvenel et al., 2021). However, nearly all current monoclonal antibody therapeutics employ the IgG1 isotype (McConnell, 2019). Therefore, we next sought to compare the activity of these mAbs in both the IgM and IgG1 formats, and generated expression plasmids containing the heavy chain variable regions upstream of the gamma1 constant region to enable expression as IgG1 mAbs. After co-transfection with the paired light chain (and co-expressed J chain for IgM production) and antibody purification, we were able to compare binding to recombinant PcrV for BCR sequences 408 and 411 when expressed as IgG *vs*. IgM mAbs (Figure 2C). CF BCRs 408 and 411 exhibited

binding to PcrV as both IgM and IgG mAbs; with higher binding as IgM, reflecting the avidity of IgM multimers (Hale et al., 2022; Thouvenel et al., 2021).

One challenge in studying candidate human PcrV mAbs is the lack of readily available comparators. V2L2MD is a heavily engineered anti-PcrV mAb that was generated by phage display binding optimization from a precursor candidate identified by hybridoma screening from PcrV-immunized, human-variable region transgenic mice (Warrener et al., 2014). The clinical candidate bi-specific, gremubamab, consists of the paired heavy-and light chain variable regions of V2L2MD fused to an anti-polysaccharide (Psl) single chain variable fragment (scFv) and the human IgG1 constant region (Chastre et al., 2022; DiGiandomenico et al., 2014). To prepare for *in vivo* protection assays, we introduced gremubamab's anti-PcrV variable regions (V2L2MD) into expression plasmids using the same strategies we have employed for expressing CF BCRs. This approach enabled us to produce and purify positive control human V2L2MD IgG and IgM mAbs in parallel with our candidate CF BCR-derived mAbs. To confirm specificity, we measured binding to recombinant PcrV for purified IgG and IgM mAbs produced in parallel from the newly cloned V2L2MD and the best candidate (411), together with commercially sourced gremubamab (Figure 2D). As IgM mAbs, 411 and V2L2MD exhibited similar binding to PcrV, on par with the commercially-sourced gremubamab.

Anti-PcrV mAbs derived from CF B cells protect mice from PA pneumonia

Next, we used a challenge model of pneumonia to test the anti-PA activity of candidate mAbs (**Figure 3A**). Mice treated with a single, 20 µg intranasal dose of anti-PcrV IgG (408 and 411) exhibited an ~2 log reduction in burden of bacteria in the lungs at 24 h in comparison with an off-target control mAb or vehicle (PBS) only (Figure 3B). The CF-derived mAbs performed equivalently to the positive control V2L2MD mAb.

Anti-PcrV mAbs derived from MBCs from pwCF exhibit in vivo protection.

The above findings demonstrate that isolation of PcrV-specific B cells from a chronically-infected CF donor enabled the rapid discovery of 2 protective mAbs. In our earlier studies of other pathogens, antigen-specific MBCs had been excellent sources for high-affinity, protective BCR sequences (Rodda et al., 2021; Thouvenel et al., 2021). Therefore, we hypothesized that higher affinity, anti-PcrV mAb might be isolated from antigen-experienced, somatically hypermutated, PcrV-specific MBCs. We obtained peripheral blood samples from 2 additional pwCF: Donor 2, who had tested culture positive for PA for the first time 6 months prior to sample collection and was negative on most recent testing, and Donor 3, who like Donor 1, was considered to be chronically infected at the time of blood sample (Table 1) (Green et al., 2012; Rosenfeld et al., 2022). Using improved methods that enhanced the efficiency and depth of sequencing (manuscript in preparation as Thouvenel et al.), we performed single cell BCR sequencing of PcrV-specific B cells from each donor. Sequences derived from PcrV-specific cells identified by surface phenotype as likely MBCs (CD21$^+$CD27$^+$) carried an average somatic hypermutation load of 2.5% in Donor 2 (Figure 4A) and <1% in Donor 3 (Figure 4–figure supplement 3A).

There was significant breadth in V gene usage within and between donors (Figure 4–figure supplement 3B). The small number of cells analyzed, compared to B cell repertoire studies that employ bulk sorted populations, limits the interpretation of our data. However, a strength of singly sorted B cell sequencing is the ability to obtain paired chain information. The heavy chain IGHV3-23 is used, with mutations, in the V2L2MD antibody, and is a very common VH gene segment in most human B cell repertoire studies (Rubelt et al., 2016). Accordingly, IGHV3-23 was also well represented in this dataset. Notably, none of the PcrV-specific B cells sequenced

for our study paired VH3-23 heavy chains with IGKV1-6, the V2L2MD light chain variable

gene (Figure 4–figure supplement 3C) and IGKV1-6 was infrequent in our dataset. Heavy- and

light-chain pairings for CD21$^+$CD27$^+$ cells are shown in Figure 4B.

To pursue our goal of identifying potential protective mAbs among MBCs, 12 of these

BCRs were randomly selected for cloning into IgG1 expression plasmids. Supernatants from

transfected cells were screened for binding to PcrV (Figure 4–figure supplement 4). BCRs 435

and 442 were the most striking binders in the transfection screen. Because our initial screen did

not adequately control for antibody concentration in the supernatant, antibodies that are less

efficiently expressed by 293T cells might appear to be poor binders. Based on performance in the

screen or for reasons of special interest (for example, the raw heavy chain sequence that became

BCR clone 439 included the IGHA constant region that defines the IgA antibody isotype, the

predominant isotype in respiratory secretions), 5 MBC-derived BCRs were chosen for

production as purified IgG mAbs. When this panel of purified mAbs was evaluated at matched

concentrations (Figure 4C), 2 of 5 of MBC-derived mAbs matched or exceeded PcrV binding

exhibited by the protective, donor 1 mAb, 411. Further, 4 of 5 matched or exceeded the other

protective donor-1 derived protective mAb, 408 IgG.

Next, we tested a moderate-affinity MBC-derived mAb (439) in addition to the two best

performing MBC-derived mAbs (437 and 442) for analysis of *in vivo* protection. Strikingly, all

three CF MBC-derived mAbs including 437 (originally derived from an IgG MBC), 439 (derived

from an IgA MBC), and 442 (derived from an IgM MBC) achieved significant reductions in

bacterial burden at 24 h when compared to an off-target control mAb or saline alone (Figure 5).

In summary, we obtained 175 unique paired heavy-light chain BCR sequences of PcrV-

binding B cells from 3 donors with CF. After expressing 20 BCRs as human mAbs, we

confirmed at least moderate *in vitro* PcrV binding for the majority screened (12 of 20, 60%), and

high-affinity binding for 7 (35%). Of the 5 mAbs tested in an established *in vivo* model of PA

pneumonia, all produced significant reductions in lung bacterial burden. In comparison to

standard methods for antibody discovery using immunized mice and screening hundreds to

thousands of hybridomas, our approach was more efficient. In addition, our results reveal a

diverse PcrV-specific repertoire for PA-specific B cells in pwCF.

Discussion

In the face of rising antibiotic resistance by pathogenic bacteria, new antimicrobial treatment

modalities are needed. Unfortunately, relative to the projected need, few novel antimicrobials are

under active pharmaceutical development (Theuretzbacher et al., 2020). Antibacterial mAbs

could fill the widening gap as alternatives, or adjuncts, to traditional antibiotics. Because

mechanisms of intrinsic resistance already limit therapeutic options, multidrug-resistant PA is

especially difficult to treat (Yaeger et al., 2021). Efforts to develop mAbs targeting PcrV, a

critical component of the PA toxin injection apparatus, are bolstered by strong evidence for

antibody-mediated protection in animal models (Goure et al., 2005; Moriyama et al., 2009; Sawa

et al., 1999; Sécher et al., 2019). Unfortunately, two engineered antibody-like drug candidates (a

Fab fragment and a bi-specific mAb) derived from mice failed to achieve efficacy in human

trials (Chastre et al., 2022; Jain et al., 2018; Yaeger et al., 2021). New approaches are needed to

identify fully human mAbs with robust anti-PA protective activity, reduced immunogenicity and

improved pharmacokinetics.

Traditional methods of mAb development begin with immunization of mice with a

recombinant protein. To address the immunogenicity of the murine constant region, mouse-

derived antibodies must then be iteratively re-humanized *in vitro* and/or modified into antibody fragments (Fab) which lack effector functions and are subject to rapid renal clearance. Although fully *in vitro* strategies like phage display can enable the use of human variable domains (De Tavernier et al., 2016; DiGiandomenico et al., 2012), the high-throughput screening processes are vulnerable to bottleneck effects and drift, and lack key features of *in vivo* B cell development, such as selection against autoreactivity.

Here, we generated multiple anti-PcrV mAbs directly from the BCR variable regions of antigen-specific B cells derived from CF donors with intermittent (past) or persistent Pa infection. For practical reasons, we tested only a subset of paired antibody sequences obtained. However, our antibody discovery strategy was extremely efficient, requiring expression of very few (<12) BCRs to yield protective anti-pseudomonal mAbs. Strikingly, in *in vivo* challenge studies, 4 of 5 mAbs directly cloned from human B cells achieved control of PA infection that matched V2L2MD IgG, the highly engineered PcrV-specific binding component of gremubamab. While assessment of lung bacterial burden requires sacrifice of the treated mice, doses achieving similar reductions in lung bacterial burdens have resulted in 100% survival in cohorts of mice in prior studies of V2L2MD (Warrener et al., 2014). Our findings employing BCR sequences from multiple isotypes are consistent with contemporaneous work by Simonis et al showing protection in neutropenic mice by 3 anti-PcrV mAbs discovered by single-cell BCR sequencing of IgG-expressing B cells from two CF donors (Simonis et al., 2023). The history of prior anti-PA therapeutics shows that promising results in animal models do not guarantee clinical success. However, mAbs derived from human B cells have important theoretical advantages over existing, animal-derived anti-pseudomonal candidates like KB001 and

gremubamab, including reduced immunogenicity (Strand et al., 2017) and fully human effector functions.

While the goal of this study was to generate novel protective mAbs for therapeutic use, our findings also offer new insight into the cellular sources of the humoral response to PA in pwCF, expanding on studies which have thus far been limited to the measurement of secreted antibodies (Mauch & Levy, 2014). As we hypothesized, given the uniquely high frequency of PA infections in the CF population, circulating PcrV-specific B cells in CF individuals were expanded compared to non-CF individuals. While the ability to generate protective mAbs *ex vivo* from CF B cells demonstrates that B cells with the technical capacity to produce functional anti-PA antibodies are present in the peripheral blood, and recent work has shown that anti-PcrV antibodies capable of neutralizing PA *in vitro* are present in some pwCF (Simonis et al., 2023), multiple *in vivo* challenges may limit PA clearance. Significant future work will be required to fully characterize the nature and dynamics of PA-specific B cells in CF individuals. Better understanding of the endogenous anti-PA response might also reveal insight that could enhance future therapeutic antibody products, enabling optimization of isotype, glycosylation, delivery site or other features.

We have previously shown that protective mAbs may be built from antigen-specific MBCs that target the parasite *Plasmodium falciparum* and the virus SARS-CoV-2 (Hale et al., 2022; Rodda et al., 2021; Thouvenel et al., 2021). Besides eliminating the humanization requirement and concern for residual immunogenicity, human MBC-derived mAbs are the product of evolved B cell development and maturation processes that encompass tremendous diversity and include selection against auto-reactivity and, for cells that have exited GCs, pre-optimized binding to antigen via affinity maturation. When combined with the results of our

present study, human antigen-specific B cells may be broadly considered as an underutilized high-yield resource in the critical endeavor to discover new treatments for infectious disease.

Materials and Methods

PBMC and serum collection

Under protocols approved by the Seattle Children's Institutional Review Board (SCH#10325), CF donors were recruited from patients receiving care that day at Seattle Children's Cystic Fibrosis Clinic in March-June 2019 and April-July 2021. Individuals who declined to participate or were unable to safely donate at least 30 mL of blood were excluded. Blood was transported at room temperature to the University of Washington Department of Immunology and processed within 4 hours of collection. Serum was collected by centrifugation at 1500 g for 10 min, and then frozen at -80 C. Peripheral mononuclear blood cells (PBMCs) were collected using SepMate-50 PBMC Isolation Tubes (STEMCELL Technologies) and frozen slowly at -80 C before transfer to liquid nitrogen for long-term storage. Non-CF donor samples were provided by BloodWorks Northwest from regular blood donors or from non-mobilized healthy donors through the Fred Hutch Hematopoietic Cell Procurement and Processing core.

Tetramer production

Recombinant his-tagged PcrV was first expressed and isolated from DH5α pET16b pcrV (kind gift from Dr. Timothy Yahr, University of Iowa; (Sawa et al., 1999)) and purified from filtered bacterial supernatant on a His affinity column. To enable tetramerization, PcrV was biotinylated using EZ-Link Sulfo-NHS-LC Biotinylation Kit (ThermoFisher). The production of antigen-specific B cell tetramer reagents has been previously described (Krishnamurty et al., 2016). Briefly, biotinylated PcrV was co-incubated with streptavidin-fluorophore (SA-PE or SA-APC;

Agilent). Decoy tetramers are produced by tetramerization of an irrelevant protein with a matched conjugated fluorophore (*eg.* PE-Cy7 for use in experiments requiring the PE-conjugated PcrV tetramer).

Identification of PcrV-specific cells

Our methods for isolating antigen-specific B cells using tetramer reagents are described in detail elsewhere (Taylor et al., 2012). Flow cytometry was performed on a BD LSR II (Becton Dickinson). For BCR sequencing experiments, single cells were sorted into a 96 well-microplate using a BDFACS Aria II that was contained within a biosafety cabinet.

BCR sequencing

cDNA was amplified from singly sorted B cells using SMART-Seq v4 (Takara Bio) at half reaction volumes. Initial BCR sequencing for donor 1 followed protocols we have previously described in detail (Hale et al., 2022; Rodda et al., 2021; Thouvenel et al., 2021). Briefly, a single, multiplex PCR was performed for each B cell using a universal primer for the template switch region and pooled constant region primers for the μ, γ, α, κ, and λ constant regions. Amplicons were then purified and sequenced by Sanger sequencing. For donors 2 and 3, a custom protocol that enables high-quality, single cell BCR sequencing by MiSeq was employed (manuscript in preparation as Thouvenel et al). Alignment of all trimmed sequences was performed using both TRUST4 (Song et al., 2021) and IGMT/HighV-QUEST (Alamyar et al., 2012). Rare conflicts (*eg.* differences in reported %SHM) were resolved by review of the raw sequence data and individual analysis in IgBlast (Ye et al., 2013).

BCR cloning

Our methods for cloning BCR variable region sequences into antibody expression plasmids were described previously (Hale et al., 2022; Rodda et al., 2021; Thouvenel et al., 2021). Briefly, each

light chain was cloned into vectors of its isotype, κ or λ, following the manufacturer's protocol for in-fusion cloning (Takara Bio). All heavy chains were similarly cloned into IgG1 and IgM plasmids in parallel. Concordance with the parental cDNA was confirmed by Sanger sequencing of the cloned plasmids.

For the V2L2MD IgG and IgM mAbs, heavy and light chain sequences were synthesized as a gBlock (IDT) by introducing mutations to match the amino acid sequence for the anti-PcrV VH/VL in gremubamab (INN 10909; 17) in the closest human VH and VL nucleotide sequences. The resultant V(D)J sequences were then synthesized as a gBlock (IDT) and then cloned into expression plasmids using the same methods described for generation of our human B cell BCR sequences.

Whole-plasmid sequences were obtained from a subset of plasmids as a quality control measure (Primordium Labs).

Production of mAbs

IgG mAbs were produced in HEK-293T cells (ATCC) by co-transfection of heavy- and light-chains in polyethylenimine as previously described (Thouvenel et al., 2021). Production of IgM mAbs was carried out as described in Hale et al., using the same human J chain plasmid. For initial screens, supernatant was harvested at 4 days post-transfection and concentrated and buffer-exchanged into PBS using 50,000 MWCO Millipore Amicon Ultra-15 Centrifugal Filter units (Thermo Fisher Scientific). Purification was carried out following manufacturer's instructions on HiTrap Protein G HP purification column for IgG mAbs (GE Healthcare), and a POROS CaptureSelect IgM-XL Affinity Matrix Column (Thermo Fisher Scientific) for IgM. Antibodies were concentrated and buffer-exchanged into phosphate buffered saline to at least 1 mg/mL, and then stored at -80C in 120-200 µL aliquots.

ELISAs

To prepare for antigen-specific antibody ELISAs, recombinant PcrV was prepared as described above and diluted to 2 µg/ml in PBS and incubated on high-binding 96 well plates (Corning) overnight at 4° C. Plates were then washed thoroughly with PBS and 0.05% Tween 20 (PBS-T). Next, non-specific interactions were blocked using 200 µl/well of PBS-T with 3% bovine serum albumin (3%) at room temperature for 1 – 3 hours. Samples of interest were serially diluted in PBS-T immediately prior to use. Dilution series for each sample were incubated on washed plates for 2 hours at room temperature. After thorough washing with PBS-T, bound antibodies were detected by incubation for 1 hour at room temperature with HRP-conjugated goat anti-human IgG (diluted 1:3000 in PBS-T; SouthernBiotech), IgM (1:1500; SouthernBiotech), or IgA (1:1500; ThermoFisher), washed again, and developed using $1\times$ 3,39,5,50-tetramethylbenzidine (Invitrogen) and 1 M H_2SO_4. OD was measured on a SpectroMax i3X (Molecular Devices, San Jose, CA) at 450 and 570 nm, and $OD_{450-750}$ was analyzed in Prism (v9.5; GraphPad). When un-purified transfectant supernatants were used, ELISAs for total IgM or IgG were also performed in parallel using human uncoated IgM or IgG ELISA kits (Invitrogen), following manufacturer's instructions.

Murine pneumonia challenge

All animal procedures were conducted according to the guidelines of the Emory University Institutional Animal Care and Use Committee (IACUC), under approved protocol number PROTO 201700441. The study was carried out in strict accordance with established guidelines and policies at Emory University School of Medicine, and recommendations in the Guide for Care and Use of Laboratory Animals, as well as local, state, and federal laws. *P. aeruginosa* strain PA103 (ATCC 29260, sputum isolate) was grown on Difco™ Pseudomonas isolation agar for 16 to 18 h at 37°C and

suspended in PBS. Inocula were adjusted spectrophotometrically to obtain the desired challenge dose in a volume of 10-20 µl. Eight-to ten-week-old female BALB/c mice (Jackson Laboratories, Bar Harbor, ME) were anesthetized by intraperitoneal injection of 0.2 ml of a cocktail of ketamine (100 mg/ml) and xylazine (5 mg/ml) and intranasally instilled with approximately 10^5 CFU *P. aeruginosa* PA103 (in 10-20 µL of PBS) as described (Moustafa et al., 2023). At 15 minutes post-infection, monoclonal antibodies or PBS were delivered via the same route in a 20 µl volume. Mice were euthanized at 24 hours post-infection, and whole lungs were collected aseptically, weighed, and homogenized for 20 seconds in 1 ml of PBS, followed by serial dilution onto Pseudomonas isolation agar, and plated for CFU enumeration.

Acknowledgements

This work was supported by a Seattle Children's/University of Washington Cystic Fibrosis Research & Translation Center pilot grant (M. Pepper). Clinical core and pilot project funding are supported by National Institutes of Health (NIH P30 DK089507). Sample collections at Seattle Children's Hospital were supported by the Clinical Core within the UW Cystic Fibrosis Foundation RDP grant SINGH19R0. M. Hale is supported by the University of Washngton Medical Scientist Training Program (NIH 5T32GM007266). D.J. Rawlings holds the Tom Hansen Investigator for Pediatric Innovation Endowment at Seattle Children's. Jennifer Haddock provided indispensable administrative support. We thank the people with cystic fibrosis for their participation in this study.

Author Contributions

Conceptualization: K.K. Takehara, M.F. Fontana, J. Netland, J.B. Goldberg, C.D. Thouvenel, D.A. Moustafa, M. Hale, M. Pepper and D.J. Rawlings. Supervision/direction: S. McNamara, R.L. Gibson, J.B. Goldberg, M. Pepper and D.J. Rawlings. Funding acquisition: J.B. Goldberg, M. Pepper and D.J. Rawlings. Investigation: K.K. Takehara, C.D. Thouvenel, M. Hale, D.A. Moustafa, A. Repele, M.F. Fontana and J. Netland. Data analysis: K.K. Takehara, C.D. Thouvenel, M. Hale, D.A. Moustafa, A. Repele, M.F. Fontana, J. Netland, M. Pepper and D.J. Rawlings. Writing and editing: M. Hale, M. Pepper and D.J. Rawlings.

References

Alamyar, E., Duroux, P., Lefranc, M.-P., & Giudicelli, V. (2012). IMGT(®) tools for the nucleotide analysis of immunoglobulin (IG) and T cell receptor (TR) V-(D)-J repertoires, polymorphisms, and IG mutations: IMGT/V-QUEST and IMGT/HighV-QUEST for NGS. *Methods in Molecular Biology*, *882*, 569–604. https://doi.org/10.1007/978-1-61779-842-9_32

Alicandro, G., Orena, B. S., Rosazza, C., Cariani, L., Russo, M., Zatelli, M., Badolato, R., Gramegna, A., Blasi, F., & Daccò, V. (2023). Humoral and cell-mediated immune responses to BNT162b2 vaccine against SARS-CoV-2 in people with cystic fibrosis. *Vaccine*, *41*(28), 4114. https://doi.org/10.1016/j.vaccine.2023.05.041

Baer, M., Sawa, T., Flynn, P., Luehrsen, K., Martinez, D., Wiener-Kronish, J. P., Yarranton, G., & Bebbington, C. (2009). An Engineered Human Antibody Fab Fragment Specific for Pseudomonas aeruginosa PcrV Antigen Has Potent Antibacterial Activity. *Infection and Immunity*, *77*(3), 1083–1090. https://doi.org/10.1128/IAI.00815-08

Chastre, J., François, B., Bourgeois, M., Komnos, A., Ferrer, R., Rahav, G., De Schryver, N., Lepape, A., Koksal, I., Luyt, C.-E., Sánchez-García, M., Torres, A., Eggimann, P., Koulenti, D., Holland, T. L., Ali, O., Shoemaker, K., Ren, P., Sauser, J., … The COMBACTE-MAGNET EVADE Study Group. (2022). Safety, efficacy, and pharmacokinetics of gremubamab (MEDI3902), an anti-Pseudomonas aeruginosa bispecific human monoclonal antibody, in P. aeruginosa-colonised, mechanically ventilated intensive care unit patients: A randomised controlled trial. *Critical Care*, *26*(1), 355. https://doi.org/10.1186/s13054-022-04204-9

Cramer, N., Klockgether, J., & Tümmler, B. (2023). Microevolution of Pseudomonas aeruginosa in the airways of people with cystic fibrosis. *Current Opinion in Immunology*, *83*, 102328. https://doi.org/10.1016/j.coi.2023.102328

De Tavernier, E., Detalle, L., Morizzo, E., Roobrouck, A., De Taeye, S., Rieger, M., Verhaeghe, T., Correia, A., Van Hegelsom, R., Figueirido, R., Noens, J., Steffensen, S., Stöhr, T., Van de Velde, W., Depla, E., & Dombrecht, B. (2016). High Throughput Combinatorial Formatting of PcrV Nanobodies for Efficient Potency Improvement. *Journal of Biological Chemistry*, *291*(29), 15243–15255. https://doi.org/10.1074/jbc.M115.684241

DiGiandomenico, A., Keller, A. E., Gao, C., Rainey, G. J., Warrener, P., Camara, M. M., Bonnell, J., Fleming, R., Bezabeh, B., Dimasi, N., Sellman, B. R., Hilliard, J., Guenther, C. M., Datta, V., Zhao, W., Gao, C., Yu, X.-Q., Suzich, J. A., & Stover, C. K. (2014). A multifunctional bispecific antibody protects against Pseudomonas aeruginosa. *Science Translational Medicine*, *6*(262), 262ra155-262ra155. https://doi.org/10.1126/scitranslmed.3009655

DiGiandomenico, A., Warrener, P., Hamilton, M., Guillard, S., Ravn, P., Minter, R., Camara, M. M., Venkatraman, V., MacGill, R. S., Lin, J., Wang, Q., Keller, A. E., Bonnell, J. C., Tomich, M., Jermutus, L., McCarthy, M. P., Melnick, D. A., Suzich, J. A., & Stover, C. K. (2012). Identification of broadly protective human antibodies to Pseudomonas aeruginosa exopolysaccharide Psl by phenotypic screening. *Journal of Experimental Medicine*, *209*(7), 1273–1287. https://doi.org/10.1084/jem.20120033

Ehrenmann, F., & Lefranc, M.-P. (2011). IMGT/3Dstructure-DB: Querying the IMGT database for 3D structures in immunology and immunoinformatics (IG or antibodies, TR, MH, RPI, and FPIA). *Cold Spring Harbor Protocols*, *2011*(6), 750–761. https://doi.org/10.1101/pdb.prot5637

François, B., Luyt, C.-E., Dugard, A., Wolff, M., Diehl, J.-L., Jaber, S., Forel, J.-M., Garot, D., Kipnis, E., Mebazaa, A., Misset, B., Andremont, A., Ploy, M.-C., Jacobs, A., Yarranton, G., Pearce, T., Fagon, J.-Y., & Chastre, J. (2012). Safety and pharmacokinetics of an anti-PcrV PEGylated monoclonal antibody fragment in mechanically ventilated patients colonized with Pseudomonas aeruginosa: A randomized,double-blind, placebo-controlled trial. *Critical Care Medicine*, *40*(8), 2320–2326. https://doi.org/10.1097/CCM.0b013e31825334f6

Frank, D. W., Vallis, A., Wiener-Kronish, J. P., Roy-Burman, A., Spack, E. G., Mullaney, B. P., Megdoud, M., Marks, J. D., Fritz, R., & Sawa, T. (2002). Generation and characterization of a protective monoclonal antibody to Pseudomonas aeruginosa PcrV. *The Journal of Infectious Diseases*, *186*(1), 64–73.

Goure, J., Broz, P., Attree, O., Cornelis, G. R., & Attree, I. (2005). Protective Anti-V Antibodies Inhibit Pseudomonas and Yersinia Translocon Assembly within Host Membranes. *The Journal of Infectious Diseases*, *192*(2), 218–225. https://doi.org/10.1086/430932

Green, D. M., Collaco, J. M., McDougal, K. E., Naughton, K. M., Blackman, S. M., & Cutting, G. R. (2012). Heritability of Respiratory Infection with *Pseudomonas aeruginosa* in Cystic Fibrosis. *The Journal of Pediatrics*, *161*(2), 290-295.e1. https://doi.org/10.1016/j.jpeds.2012.01.042

Hale, M., Netland, J., Chen, Y., Thouvenel, C. D., Smith, K. N., Rich, L. M., Vanderwall, E. R., Miranda, M. C., Eggenberger, J., Hao, L., Watson, M. J., Mundorff, C. C., Rodda, L. B., King, N. P., Guttman, M., Gale, M., Abraham, J., Debley, J. S., Pepper, M., & Rawlings, D. J. (2022). IgM antibodies derived from memory B cells are potent cross-variant neutralizers of SARS-CoV-2. *Journal of Experimental Medicine*, *219*(9), e20220849. https://doi.org/10.1084/jem.20220849

Jain, R., Beckett, V. V., Konstan, M. W., Accurso, F. J., Burns, J. L., Mayer-Hamblett, N., Milla, C., VanDevanter, D. R., Chmiel, J. F., & KB001-A Study Group. (2018). KB001-A, a novel anti-inflammatory, found to be safe and well-tolerated in cystic fibrosis patients infected with Pseudomonas aeruginosa. *Journal of Cystic Fibrosis*, *17*(4), 484–491. https://doi.org/10.1016/j.jcf.2017.12.006

Krishnamurty, A. T., Thouvenel, C. D., Portugal, S., Keitany, G. J., Kim, K. S., Holder, A., Crompton, P. D., Rawlings, D. J., & Pepper, M. (2016). Somatically Hypermutated Plasmodium-Specific IgM+ Memory B Cells Are Rapid, Plastic, Early Responders upon Malaria Rechallenge. *Immunity*, *45*(2), 402–414. https://doi.org/10.1016/j.immuni.2016.06.014

Lund-Palau, H., Turnbull, A. R., Bush, A., Bardin, E., Cameron, L., Soren, O., Wierre-Gore, N., Alton, E. W. F. W., Bundy, J. G., Connett, G., Faust, S. N., Filloux, A., Freemont, P., Jones, A., Khoo, V., Morales, S., Murphy, R., Pabary, R., Simbo, A., … Davies, J. C. (2016). Pseudomonas aeruginosa infection in cystic fibrosis: Pathophysiological mechanisms and therapeutic approaches. *Expert Review of Respiratory Medicine*, *10*(6), 685–697. https://doi.org/10.1080/17476348.2016.1177460

Mancuso, G., Midiri, A., Gerace, E., & Biondo, C. (2021). Bacterial Antibiotic Resistance: The Most Critical Pathogens. *Pathogens*, *10*(10), 1310. https://doi.org/10.3390/pathogens10101310

Mauch, R. M., & Levy, C. E. (2014). Serum antibodies to Pseudomonas aeruginosa in cystic fibrosis as a diagnostic tool: A systematic review. *Journal of Cystic Fibrosis*, *13*(5), 499–507. https://doi.org/10.1016/j.jcf.2014.01.005

McConnell, M. J. (2019). Where are we with monoclonal antibodies for multidrug-resistant infections? *Drug Discovery Today*, *24*(5), 1132–1138. https://doi.org/10.1016/j.drudis.2019.03.002

Mogayzel, P. J., Naureckas, E. T., Robinson, K. A., Brady, C., Guill, M., Lahiri, T., Lubsch, L., Matsui, J., Oermann, C. M., Ratjen, F., Rosenfeld, M., Simon, R. H., Hazle, L., Sabadosa, K., & Marshall, B. C. (2014). Cystic Fibrosis Foundation Pulmonary Guideline. Pharmacologic Approaches to Prevention and Eradication of Initial Pseudomonas aeruginosa Infection. *Annals of the American Thoracic Society*, *11*(10), 1640–1650. https://doi.org/10.1513/AnnalsATS.201404-166OC

Moriyama, K., Wiener-Kronish, J. P., & Sawa, T. (2009). Protective effects of affinity-purified antibody and truncated vaccines against Pseudomonas aeruginosa V-antigen in neutropenic mice. *Microbiology and Immunology*, *53*(11), 587–594. https://doi.org/10.1111/j.1348-0421.2009.00165.x

Moss, J., Ehrmantraut, M. E., Banwart, B. D., Frank, D. W., & Barbieri, J. T. (2001). Sera from Adult Patients with Cystic Fibrosis Contain Antibodies to Pseudomonas aeruginosa Type III Apparatus. *Infection and Immunity*, *69*(2), 1185–1188. https://doi.org/10.1128/IAI.69.2.1185-1188.2001

Moustafa, D. A., DiGiandomenico, A., Raghuram, V., Schulman, M., Scarff, J. M., Davis, M. R., Varga, J. J., Dean, C. R., & Goldberg, J. B. (2023). Efficacy of a Pseudomonas aeruginosa serogroup O9 vaccine. *Infection and Immunity*, *91*(12), e0024723. https://doi.org/10.1128/iai.00247-23

Ong, T., & Ramsey, B. W. (2023). Cystic Fibrosis: A Review. *JAMA*, *329*(21), 1859–1871. https://doi.org/10.1001/jama.2023.8120

Rodda, L. B., Netland, J., Shehata, L., Pruner, K. B., Morawski, P. A., Thouvenel, C. D., Takehara, K. K., Eggenberger, J., Hemann, E. A., Waterman, H. R., Fahning, M. L., Chen, Y., Hale, M., Rathe, J., Stokes, C., Wrenn, S., Fiala, B., Carter, L., Hamerman, J. A., … Pepper, M. (2021). Functional SARS-CoV-2-Specific Immune Memory Persists after Mild COVID-19. *Cell*, *184*(1), 169-183.e17. https://doi.org/10.1016/j.cell.2020.11.029

Rosenfeld, M., Faino, A. V., Onchiri, F., Aksit, M. A., Blackman, S. M., Blue, E. E., Collaco, J. M., Gordon, W. W., Pace, R. G., Raraigh, K. S., Zhou, Y.-H., Cutting, G. R., Knowles, M. R., Bamshad, M. J., & Gibson, R. L. (2022). Comparing encounter-based and annualized chronic pseudomonas infection definitions in cystic fibrosis. *Journal of Cystic Fibrosis*, *21*(1), 40–44. https://doi.org/10.1016/j.jcf.2021.07.020

Rossi, E., La Rosa, R., Bartell, J. A., Marvig, R. L., Haagensen, J. A. J., Sommer, L. M., Molin, S., & Johansen, H. K. (2020). Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. *Nature Reviews Microbiology*, 1–12. https://doi.org/10.1038/s41579-020-00477-5

Rubelt, F., Bolen, C. R., McGuire, H. M., Vander Heiden, J. A., Gadala-Maria, D., Levin, M., Euskirchen, G. M., Mamedov, M. R., Swan, G. E., Dekker, C. L., Cowell, L. G., Kleinstein, S. H., & Davis, M. M. (2016). Individual heritable differences result in unique cell lymphocyte receptor repertoires of naïve and antigen-experienced cells. *Nature Communications*, *7*, 11112. https://doi.org/10.1038/ncomms11112

Sato, H., & Frank, D. (2011). Multi-Functional Characteristics of the Pseudomonas aeruginosa Type III Needle-Tip Protein, PcrV; Comparison to Orthologs in other Gram-negative Bacteria. *Frontiers in Microbiology*, *2*(142). https://doi.org/10.3389/fmicb.2011.00142

Sawa, T., Yahr, T. L., Ohara, M., Kurahashi, K., Gropper, M. A., Wiener-Kronish, J. P., & Frank, D. W. (1999). Active and passive immunization with the Pseudomonas V antigen protects against type III intoxication and lung injury. *Nature Medicine*, *5*(4), 392–398. https://doi.org/10.1038/7391

Sécher, T., Dalonneau, E., Ferreira, M., Parent, C., Azzopardi, N., Paintaud, G., Si-Tahar, M., & Heuzé-Vourc'h, N. (2019). In a murine model of acute lung infection, airway administration of a therapeutic antibody confers greater protection than parenteral administration. *Journal of Controlled Release*, *303*, 24–33. https://doi.org/10.1016/j.jconrel.2019.04.005

Shteinberg, M., Haq, I. J., Polineni, D., & Davies, J. C. (2021). Cystic fibrosis. *The Lancet*, *397*(10290), 2195–2211. https://doi.org/10.1016/S0140-6736(20)32542-3

Simonis, A., Kreer, C., Albus, A., Rox, K., Yuan, B., Holzmann, D., Wilms, J. A., Zuber, S., Kottege, L., Winter, S., Meyer, M., Schmitt, K., Gruell, H., Theobald, S. J., Hellmann, A.-M., Meyer, C., Ercanoglu, M. S., Cramer, N., Munder, A., … Rybniker, J. (2023). Discovery of highly neutralizing human antibodies targeting Pseudomonas aeruginosa. *Cell*, *186*(23), 5098-5113.e19. https://doi.org/10.1016/j.cell.2023.10.002

Song, L., Cohen, D., Ouyang, Z., Cao, Y., Hu, X., & Liu, X. S. (2021). TRUST4: Immune repertoire reconstruction from bulk and single-cell RNA-seq data. *Nature Methods*, *18*(6), 627–630. https://doi.org/10.1038/s41592-021-01142-2

Strand, V., Balsa, A., Al-Saleh, J., Barile-Fabris, L., Horiuchi, T., Takeuchi, T., Lula, S., Hawes, C., Kola, B., & Marshall, L. (2017). Immunogenicity of Biologics in Chronic Inflammatory Diseases: A Systematic Review. *BioDrugs*, *31*(4), 299–316. https://doi.org/10.1007/s40259-017-0231-8

Taylor, J. J., Pape, K. A., & Jenkins, M. K. (2012). A germinal center–independent pathway generates unswitched memory B cells early in the primary response. *Journal of Experimental Medicine*, *209*(3), 597–606. https://doi.org/10.1084/jem.20111696

Theuretzbacher, U., Outterson, K., Engel, A., & Karlén, A. (2020). The global preclinical antibacterial pipeline. *Nature Reviews Microbiology*, *18*(5), 275–285. https://doi.org/10.1038/s41579-019-0288-0

Thouvenel, C. D., Fontana, M. F., Netland, J., Krishnamurty, A. T., Takehara, K. K., Chen, Y., Singh, S., Miura, K., Keitany, G. J., Lynch, E. M., Portugal, S., Miranda, M. C., King, N. P., Kollman, J. M., Crompton, P. D., Long, C. A., Pancera, M., Rawlings, D. J., & Pepper, M. (2021). Multimeric antibodies from antigen-specific human IgM+ memory B cells restrict Plasmodium parasites. *Journal of Experimental Medicine*, *218*(4), e20200942. https://doi.org/10.1084/jem.20200942

Warrener, P., Varkey, R., Bonnell, J. C., DiGiandomenico, A., Camara, M., Cook, K., Peng, L., Zha, J., Chowdury, P., Sellman, B., & Stover, C. K. (2014). A Novel Anti-PcrV Antibody Providing Enhanced Protection against Pseudomonas aeruginosa in Multiple Animal Infection Models. *Antimicrobial Agents and Chemotherapy*, *58*(8), 4384–4391. https://doi.org/10.1128/AAC.02643-14

Weiner-Lastinger, L. M., Abner, S., Edwards, J. R., Kallen, A. J., Karlsson, M., Magill, S. S., Pollock, D., See, I., Soe, M. M., Walters, M. S., & Dudeck, M. A. (2020). Antimicrobial-resistant pathogens associated with adult healthcare-associated infections: Summary of data reported to the National Healthcare Safety Network, 2015–2017. *Infection Control and Hospital Epidemiology*, *41*(1), 1–18. https://doi.org/10.1017/ice.2019.296

Yaeger, L. N., Coles, V. E., Chan, D. C. K., & Burrows, L. L. (2021). How to kill Pseudomonas—Emerging therapies for a challenging pathogen. *Annals of the New York Academy of Sciences*, *1496*(1), 59–81. https://doi.org/10.1111/nyas.14596

Ye, J., Ma, N., Madden, T. L., & Ostell, J. M. (2013). IgBLAST: An immunoglobulin variable domain sequence analysis tool. *Nucleic Acids Research*, *41*(Web Server issue), W34-40. https://doi.org/10.1093/nar/gkt382

Figures

Figure 1. PcrV-specific B cells in people with cystic fibrosis (pwCF). A) Schematic of primary human B cells binding to PcrV tetramer reagent and decoy (left). Representative flow plot for B cells after enrichment. Cells binding only to the PcrV tetramer are indicated with the red box. B-C) Percentage (B) and number (C) of PcrV-specific B cells in pwCF (CF; n = 14) vs. control, blood bank donors (non-CF; n = 14). Statistics determined by 2-tailed Mann-Whitney tests. p = ***: <0.001, ****: <0.0001. Error bars represent mean and SD.

Figure 2. **Generation of anti-PcrV mAbs derived from B cells isolated from a PA-infected CF donor.** A) Percentage of somatic hypermutation (SHM) detected in B cell receptor (BCR) sequences from cells of the indicated phenotype for CF donor 1. B) ELISAs showing PcrV binding using supernatants derived from 293T cells transfected with IgM expression plasmids containing the indicated BCR sequences. C) Area under the curve (AUC) for representative ELISAs of purified antibodies 408 and 411 expressed alternatively as IgM vs. IgG in a single experiment. D) AUC for representative ELISA of purified mAbs V2L2 and 411 expressed,

alternatively as, IgG vs. IgM; and for the commercially sourced, clinical, bi-specific mAb, gremubamab.

Figure 3. **pwCF-derived, germline, anti-PcrV-specific mAbs exhibit robust anti-PA activity in an *in vivo* mouse pneumonia model.** A) Schematic illustrating the experimental PA infection and mAb delivery protocol. B) Bacterial load in mouse lungs at 24 h post-infection for mice that received 20 µg intranasal dose of off-target, control IgG1 mAb (Ctl), indicated anti-PcrV mAbs, or diluent alone (PBS). Asterisks show significance in Dunn's test versus animals treated with diluent only (PBS); p = *: <0.05.

Figure 4. **High affinity anti-PcrV mAbs derived from memory B cells isolated from CF Donor 2.** A) Somatic hypermutation (%SHM) rates in BCR sequences from individual B cells

with the indicated surface phenotype isolated from CF donor 2. Each circle represents a heavy or light chain sequence from a singly-sorted cell. B) Heatmap showing paired heavy (x-axis) and light (y-axis) V genes for 79 MBCs in donor 3. The color gradient depicts SHM for the heavy chains in each pairing. Clone numbers for BCRs subsequently expressed as mAbs are included above their corresponding box. C) PcrV binding for purified mAbs generated from five MBC BCRs (colored lines) screened in a single ELISA. The two Donor 1-derived mAbs with *in vivo* protective activity (408 and 411) are used as benchmarks for relative binding activity (dashed lines). The off-target control (anti-SARS-CoV-2 RBD) line appears hidden because it overlaps with the blue line (432).

Figure 5. mAbs derived from CF MBC BCR sequences control PA infection in a murine pneumonia model. Lung bacterial load for mice treated intranasally with 20 µg of the indicated anti-PcrV mAb, or vehicle only (PBS). Combined data from two independent experiments is shown (n = 10 mice per condition). Asterisks show significance in Dunn's test versus animals treated with the off-target control antibody (Ctl); p = *: <0.05, **: <0.01, ***: <0.001, n.s.: not significant.

Tables

Donor	Age	On ETI?	Lung disease severity	PA infectious history
1	33	Yes	Moderate	Chronic
2	19	Yes	Minimal	Culture-negative at time of blood draw. Culture-positive 6 mo. prior. Previously, 5 y. of negative cultures.
3	18	Yes	Minimal	Chronic

Table 1. Clinical characteristics of subjects from whom PcrV-specific BCR sequences were obtained. ETI: elexacaftor/tezacaftor/ivacaftor triple therapy. Chronic infection is defined here

as positive PA cultures in at least 2 of 3 consecutive years (Green et al., 2012; Rosenfeld et al., 2022).