Natural variation in salt-induced changes in root:shoot ratio

reveals SR3G as a negative regulator of root suberization

and salt resilience in Arabidopsis

Maryam Rahmati Ishka[1], - ORCID - 0000-0002-5447-643X - mri23@cornell.edu

Hayley Sussman[1], - ORCID - 0000-0002-1331-3319 - hs995@cornell.edu

Yunfei Hu[2], - ORCID - 0009-0004 -9799-7798 - huyf19@lzu.edu.cn

Mashael Daghash Alqahtani[3], - ORCID - 0000-0003-0043-0847 - mdalqahtani@pnu.edu.sa

Eric Craft[4], - ORCID - 0000-0003-0966-8296 - ejc35@cornell.edu

Ronell Sicat[5], - ORCID - 0000-0001-7037-1614 - ronell.sicat@kaust.edu.sa

Minmin Wang[6], - ORCID - 0000-0001-9545-7895 - minwang@ucdavis.edu

Li'ang Yu[1], - ORCID - 0000-0002-9556-011X - ly76@cornell.edu

Rashid Ait-Haddou[5, 6], - ORCID - 0000-0002-2606-127X - rachid.aithaddou@kfupm.edu.sa
Bo Li[2] - ORCID - 0000-0002-3603-3352 - li_bo@lzu.edu.cn

Georgia Drakakaki[7], - ORCID – 0000-0002-3949-8657 - gdrakakaki@ucdavis.edu

Andrew Nelson[1], - ORCID - 0000-0001-9896-1739 - an425@cornell.edu

Miguel Pineros[4], ORCID - 0000-0002-7166-1848 - map25@cornell.edu

Arthur Korte[8], - ORCID - 0000-0003-0831-1463 - arthur.korte@uni-wuerzburg.de

Lukasz Jaremko[9], lukasz.jaremko@kaust.edu.sa

Christa Testerink[10], - ORCID - 0000-0001-6738-115X - christa.testerink@wur.nl

Mark Tester[3], - ORCID - 0000-0002-5085-8801 mark.tester@kaust.edu.sa

Magdalena M. Julkowska[1,3]* - ORCID - 0000-0002-4259-8296

*- Corresponding author mmj55@cornell.edu

Affiliations:

- Boyce Thompson Institute, Ithaca, NY, USA

- Lanzhou University, Lanzhou, China

- Center for Desert Agriculture, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

- USDA-ARS, Ithaca, NY, USA

- Visualization Core Lab, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

- King Fahd University of Petroleum and Minerals, Department of Mathematics

- UC Davis, Davis, CA, USA

- Julius-von-Sachs-Institute & Center for Computational and Theoretical Biology, JMU, Wuerzburg, Germany

- King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

- Wageningen Research and University, Wageningen, The Netherlands

Abstract

Soil salinity is one of the major threats to agricultural productivity worldwide. Salt stress exposure alters root and shoot growth rates, thereby affecting overall plant performance. While past studies have extensively documented the effect of salt stress on root elongation and shoot development separately, here we take an innovative approach by examining the coordination of root and shoot growth under salt stress conditions. Utilizing a newly developed tool for quantifying the root:shoot ratio in agar-grown Arabidopsis seedlings, we found that salt stress results in a loss of coordination between root and shoot growth rates. We identify a specific gene cluster encoding domain-of-unknown-function 247 (DUF247), and characterize one of these genes as Salt Root:shoot Ratio Regulator Gene (SR3G). Further analysis elucidates the role of SR3G as a negative regulator of salt stress tolerance, revealing its function in regulating shoot growth, root suberization, and sodium accumulation. We further characterize that *SR3G* expression is modulated by *WRKY75* transcription factor, known as a positive regulator of salt stress tolerance. Finally, we show that the salt stress sensitivity of *wrky75* mutant is completely diminished when it is combined with *sr3g* mutation. Together, our results demonstrate that utilizing root:shoot ratio as an architectural feature leads to the discovery of new stress resilience gene. The study's innovative approach and findings not only contribute to our understanding of plant stress tolerance mechanisms but also open new avenues for genetic and agronomic strategies to enhance crop environmental resilience.

Introduction

Salt stress is predominant in the dryland areas where evaporation rate exceeds water input. As all water contains dissolved ions, the prolonged exposure to drought stress results in increased accumulation of salts in the upper soil layers (Abrol et al., 1988; Fischer et al., 2021; Richards, 1969). Dryland areas of South America, southern and western Australia, Mexico, southwest United States, and South Africa were previously identified as primary hotspots for soil salinization (Hassani et al., 2021). As fresh water becomes scarcer, understanding the mechanisms of salt stress resilience becomes of paramount importance in the context of global agricultural productivity and sustainability.

Salt stress detrimentally affects plant growth and development, decreasing the activity of the meristem (West et al., 2004). Studies utilizing main root elongation as an indicator of salt tolerance have led to identification of many crucial components of salt stress signaling and tolerance, including genes involved in salt-overly sensitive signaling (SOS) pathway, jasmonate , ethylene, abscisic acid (ABA), and reactive oxygen species (ROS) in glycophytes and halophytes alike (Fu et al., 2019; Geng et al., 2013; Hu et al., 2023; Şekerci et al., 2023; Valenzuela et al., 2016; Wu et al., 1996). Additionally, negative halotropism response, that describes a directional growth of main root away from the salt gradient, was used to identify a wide range of molecular components regulating the auxin redistribution and cell-wall remodeling, to actively avoid high salt (Deolu-Ajayi et al., 2019; Galvan-Ampudia et al., 2013; Korver et al., 2020; Yu et al., 2022; Y. Zou et al., 2022). The development of lateral roots is affected by salinity, and the differential effect on main and lateral roots within specific genotype can result in substantial reprogramming of root system architecture (Julkowska et al., 2017, 2014). The shoot development is impacted by salt stress, as the salt impacts the plant transpiration and photosynthetic efficiency instantaneously (Awlia et al., 2021, 2016; Julkowska et al., 2016). Many forward genetic studies were highly successful in associating natural variation in root and shoot growth with allelic variation in gene coding and promoter regions, thereby identifying potential new target traits for improved stress resilience (Awlia et al., 2021; Julkowska et al., 2017, 2016). However, these studies typically focus on either above or below ground tissue.

The coordination of root and shoot growth rate is an important aspect of seedling establishment, and becomes even more critical under abiotic stress (Bacher et al., 2022; Chen et al., 2022; Karcher et al., 2008; Wang et al., 2018; Xu et al., 2015). The importance of increased root:shoot ratio, maximizing water scavenging while limiting transpiration, is well studied under drought stress (Asch et al., 2005; Chen et al., 2022; Karcher et al., 2008; Uga et al., 2013). However, the effect of salt stress on root:shoot ratio is yet to be described, and its contribution to salt stress remains unknown. Here, we developed a tool to quantify root:shoot ratio from agar-grown Arabidopsis seedlings, and re-analyzed the natural diversity panel of Arabidopsis exposed to salt stress (Julkowska et al., 2017). We have observed that while salt stress does not elicit a clear response in root:shoot ratio, it results in loss of coordination of root and shoot growth rates. The locus associated with salt-induced changes in root:shoot ratio consisted of a gene cluster encoding domain-of-unknown-function 247 (DUF247), with majority of SNPs clustering into one specific gene (At3g50160), that we named as Salt Root:shoot Ratio Regulator Gene (SR3G). Further characterization of SR3G revealed its function as a negative regulator of salt stress resilience, through regulation of root suberization, shoot growth, and Na^+ accumulation. We found WRKY75 transcription factor, a positive regulator of salt stress tolerance, acts upstream of SR3G, binding directly to the SR3G promoter in yeast. While *sr3g* and *wrky75* mutants showed contrasting salt stress response, the double mutants of *sr3g/wrky75* displayed reduced sensitivity to salt stress.

Materials & Methods

Screening HapMap population for the increase in root and shoot size

The images collected within (Julkowska et al., 2017) were re-analyzed in this manuscript to examine salt-induced changes in the root:shoot ratio for the data presented here. 360 accessions of the Arabidopsis HapMap population (Weigel and Mott, 2009) were screened over 7 experimental batches, using 4 biological replicates per genotype per condition. The seeds were surface sterilized, incubated in sterile MQ water at 4°C for 48 h, and germinated on media containing ½ Murashige-Skoog, 0.5% sucrose, 0.1% MES buffer, pH 5.8 (KOH) and 1% Dashin agar. The 4-day-old seedlings were subsequently transferred to media additionally supplemented with 0, 75 or 125 mM NaCl. The plates were scanned every 2nd day using a flat-bed scanner for 8 consecutive days.

The collected images were analyzed for green and white pixels, corresponding to the projected surface of the shoot and root, respectively (https://github.com/ronellsicat/PlantSeg). A Matlab-based tool was developed to simplify and speed up the segmentation and analysis pipeline. For automatic segmentation, the tool uses a combination of image operations (histogram equalization), thresholding on different color spaces (e.g., RGB, YCbCr, Lab, HSV), and binary image processing (boundary and islands removal). As the tool is digitalizing various color scales and classifies pixels into either white (root), green (shoot) or blue (background) categories, the adjustment for white balance is obsolete. For analysis, the tool uses connected components analysis to extract the plants (combined roots and shoots) and can guide the user to find possible mistakes which can be manually corrected via the user interface. After segmentation, the number of pixels for roots and shoots are counted and recorded automatically. More details are provided in the publicly available repository for the tool (https://github.com/ronellsicat/PlantSeg). The collected images and corresponding data acquired from them using the tool can be accessed at Zenodo Repository (experimental batch 1: https://doi.org/10.5281/zenodo.7272155; experimental batch 2: https://doi.org/10.5281/zenodo.7272344; experimental batch 3: https://doi.org/10.5281/zenodo.7272647; experimental batch 4: https://doi.org/10.5281/zenodo.7271841; experimental batch 5: https://doi.org/10.5281/zenodo.7272139; experimental batch 6: https://doi.org/10.5281/zenodo.7273936; experimental batch 7: https://doi.org/10.5281/zenodo.7273937; quantified data: https://doi.org/10.5281/zenodo.7268673). The tool's precision was evaluated by growing the Col-0 seedlings on agar plates, analyzing their root and shoot projected area on the final day using the tool, and correlating it with the recorded fresh and dry weight of the seedlings (**Fig. S1**). The collected data from the HapMap panel was analyzed using R for potential outliers (samples identified to be more than 3x standard deviation from the mean of each genotype within each condition), and the genotype-specific mean was calculated using the stats package (Team et al., 2018). The overall increase in root and shoot area was investigated for differences between control and salt stress conditions (**Fig. S2**). The root and shoot growth were estimated by fitting a linear function for a log-transformed projected root or shoot area over the non-transformed timeline of the experiment (https://rpubs.com/mjulkowska/HapMap-root-shoot-data). The data were subsequently used to explore differences between the experimental batches and the standard deviations within individual accessions. The genotypes above 3 standard deviations from the population's mean were excluded from further analysis. The salt tolerance index (STI) was calculated for root and shoot growth by dividing the average trait value at salt stress by the genotype-specific average at control conditions (**Fig. S3**). The correlations between individual traits were examined using the genotypic mean (**Fig. S4**). All quantitative data for root and shoot growth was visualized using ggplot2, corrplot and ggpubr packages

143 (Kassambara, n.d.; Wei and Simko, 2017; Wilke, 2016). For all measured traits, we estimated broad sense
144 heritability (H^2) using the MVApp (Julkowska et al., 2018) (**Table S1**). The data used as an input for GWAS
145 can be found in **Table S2**.

Genome-Wide Association Study

The genotype-specific mean was used as an input to GWAS. The GWAS was conducted as described in
(Awlia et al., 2021), initially using 250 k SNPs (Horton et al., 2012) and subsequently using 4 M SNPs
(1001genomes.org). The GWAS associations were evaluated for minor allele count (MAC) and association
strength above the Bonferroni threshold with -log10(p-value/#SNPs), calculated for each sub-population
of SNPs above threshold MAC (**Table S3**, Bonf.threshold.MAC.specific). The GWAS data was processed for
associations with all SNPs used to generate Manhattan and QQ-plots (R notebook for processing GWAS
data: https://rpubs.com/mjulkowska/GWAS-root-shoot-Arabidopsis-salt-250kSNPs).   Additionally,   we
prioritized the genetic loci where multiple SNPs were associated with individual traits, or multiple traits
were associated with one genetic locus. Our final selection yielded 14 associations with traits measured
under control conditions, while 9 and 23 associations were specific to traits measured under 75 and 125
mM NaCl, respectively (**Table S3**). The GWAS output files can be accessed at Zenodo Repository (GWAS
using   250   k   SNPs: https://doi.org/10.5281/zenodo.7271751;   GWAS   using   4   M   SNPs:
https://doi.org/10.5281/zenodo.7271741). For the most promising loci (**Table S4**), the sequence
information of 147 accessions belonging to the HapMap population was downloaded from the 1001
genomes project website (1001genomes.org) and aligned with ClustalO as described in (Julkowska et al.,
2016). More details on plotting the genome divergence in the loci of interest can be found at
https://www.protocols.io/view/plotdivergence-for-plotting-the-natural-variation-x54v98y84l3e/v1.   The
input   and   output   files   of   Plot   Divergence   analysis   can   be   found   in   Zenodo   repository
(https://doi.org/10.5281/zenodo.7278021).

Haplotype analysis

For the most promising candidate loci (**Table S4**), we have identified the gene open reading frames that
were located within the genome-wide linkage-disequilibrium (LD) of the associated SNPs. The LD was
expanded if multiple SNPs were identified within the region, and the region of interest was expanded
based on the number of coding genes within the LD window. The protein-coding genes were subsequently
used for haplotype analysis. All the SNPs located within the gene coding region were used to perform a
haplotype   analysis.   The   detailed   protocol   used   for   haplotype   analysis   can   be   accessed   at
https://www.protocols.io/view/haplotype-analysis-of-the-cds-in-arabidopsis-thali-5jyl86n8v2wp/v1. The
accessions   were   grouped   into   haplotype   groups   depending   on   the   combination   of   SNPs   within   the
protein-coding region of the gene of interest. The haplotypes represented by less than 3 accessions were
excluded from further analysis. The phenotypes of individual haplotypes were explored for significant
differences using ANOVA. The results for all investigated haplotypes can be accessed at Zenodo Repository
(https://doi.org/10.5281/zenodo.7277703).

Identification of SR3G orthologs and paralogs

DUF247 domain-containing orthologs from *Arabidopsis thaliana* and eight other species were identified
by   reciprocal   best   BLAST   within   CoGe   ((Nelson   et   al.,   2018);
https://genomevolution.org/coge/CoGeBlast.pl) using *SR3G* (At3g50160) nucleotide sequence as query
with an E-value cutoff of 1.00E-10). The gene ID, species, and gene names been used in this study are
listed in **Table S5**. Sequence homologs were further confirmed by multiple alignments of those hits within
each species using Geneious Prime. Sequences with missing or incomplete DUF247 domains were

removed, with the exception of the mis-annotated DUF427 homologs in *Camelina sativa* which contained
an extraneous intron that, after removal, resulted in a set of intact DUF247 open reading frame containing
genes.

Phylogenetic and positive selection analysis of SR3G orthologs

Identified orthologs were translated into amino acid, then aligned by MUSCLE. Alignments were carefully
validated by manual correction. Briefly, diverged insertions were trimmed, shifted frames were adjusted,
and conserved regions after translation were retained for phylogeny reconstruction. Phylogenetic
reconstruction was performed using the RAxML (Stamatakis, 2014) plug-in within Geneious (Geneious
Prime 2022.1.1) with the GAMMA-GTR nucleotide model, rapid bootstrapping with a parsimony search,
and 100 bootstraps. The distribution of *SR3G* and other DUF247 orthologs, root lengths for each branch,
were visualized by software Geneious Prime.

To characterize any potential positive selection that might exist along certain branches of the
inferred phylogeny, including SR3G and its orthologs, we performed a branch-site model A test on
branches leading to SR3G, and branches leading to the clade containing SR3G and its closet orthologs
using PAML4.0 (Stamatakis, 2014; Yang, 2007) as performed previously (Beilstein et al., 2015; Nelson et
al., 2014). Specific branches are denoted as $\omega0$, $\omega1$, $\omega2$, $\omega3$, and ω-rest (Table S???). In the null hypothesis
(H_0): we assigned a substitution rate ($\omega = dN/dS$) along with $0 < \omega < 1$ or $\omega = 1$ to all branches (represented
by H_0: $\omega0 = \omega1 = \omega2 = \omega3 = \omega$-rest). The alternative hypotheses (HAs) here allow for differential selection
to be applied to respective branch(es). Specifically, the five alternative hypotheses are listed here: H_1: ω
-rest branches $= \omega0 = \omega3 = \omega1 \neq \omega2$, H_2: ω -rest branches $= \omega0 = \omega1 = \omega2 \neq \omega3$, H_3: ω -rest branches $= \omega0$
$\neq \omega1 \neq \omega2 \neq \omega3$, H_4: ω -rest branches $\neq \omega0 \neq \omega1 \neq \omega2 \neq \omega3$, H_{sep}: each branch has its unique ω value. To
test these hypotheses, we further validate or reject alternative hypotheses by conducting a Chi-square
test based on likelihood scores (Ln-L score) derived from the respective branch-site model A tests (Chi-
square test cutoff: $P < 0.05$).

Screening candidate genes identified through GWAS

The genes located within the selected loci (**Table S4**) were further inspected for their contributions to root
development and root:shoot ratio under control and salt stress conditions. The T-DNA insertion lines
(**Table S4**) were genotyped using primers listed in **Table S4**, to confirm the homozygous insertion of the
T-DNA within the phenotype lines. All the mutant lines were used for seed propagation in the controlled
growth chamber with 16/8h of light/dark period, at 22°C, 60% humidity and 200 µmole m^{-1} s^{-1} under long-
day conditions. The seeds of plants identified as homozygous were subsequently used for the experiment
on agar plates, as described before for the Genome-Wide Association Study. A detailed protocol for plate
assays can be accessed at https://www.protocols.io/view/plate-assay-for-identification-of-salt-stress-
indu-eq2lyd1mwlx9/v1. The plate images were analyzed for root system architecture using Smart Root
(Lobet et al., 2011) and the root:shoot ratio using the tool described before for Genome-Wide Association
Study. The analysis of the mutant lines was performed in R. The data was first curated for selecting
possible outliers within a genotype and subsequently used to fit growth functions for root and shoot
tissues using exponential functions.

Screening SR3G mutants within root:shoot locus

The following T-DNA lines (listed in **Table S4**) were used for SR3G (AT3G50160) as the main GWAS
candidate: *sr3g-4* (SAIL_690_E12, glufosinate resistant) and *sr3g-5* (SAIL_608_C06, glufosinate resistant).
All T-DNA insertion alleles were in the background of Col-0 accession. The position of T-DNA insertions

was verified using PCR (primers listed in **Table S4**). RNAi lines targeting the neighboring of the *SR3G* or a
combination of *SR3G* and its neighboring *DUF247* genes included CSHL#3 targeting DUF130, CSHL#4
targeting SR3G, CSHL#5 targeting DUF130 and DUF170, CSHL#6 targeting DUF130 and DUF140, CSHL#7
targeting DUF10 and DUF150, CSHL#8 targeting DUF140 and DUF190, CSHL#10 targeting DUF180,
CSHL#11 targeting DUF150 and DUF170, CSHL#12 targeting DUF130, DUF140, DUF170, DUF180, and
DUF190, CSHL#13 targeting DUF140 and DUF170, CSHL#15 targeting DUF140, DUF170, DUF180, and
DUF190, CSHL#16 targeting DUF130, SR3G, and DUF170, CSHL#18 targeting DUF150 and DUF190,
CSHL#19 targeting DUF130, DUF140, DUF150, SR3G, DUF170, DUF180, and DUF190, CSHL#20 targeting
DUF130 and SR3G, CSHL#21 targeting DUF130 and DUF150, and CSHL#22 targeting DUF150 and SR3G.
The RNAi lines were verified using RT-qPCR, with the primers listed in **Table S4**. The following T-DNA
insertion alleles (also listed in **Table S4**) were used to characterize the neighboring genes of the *SR3G*
(AT3G50160): 120.2 (SAIL_382_A09), 140.1 (SALK_044685), 140.2 (SALK_005466), 140.3 (SALK_122700),
150.1 (SALK_003824), 150.3 (SALK_071080), 170.1 (SALK_009186), 170.2 (SALK_112602), 170.3
(SALK_008710), 170.4 (SALK_145999), 170.5 (SALK_072937), 190.2 (SALK_137791), 200.1 (SALK_113759),
200.2 (SALK_093701), 200.3 (SALK_129634). The primers used for genotyping PCR are shown in **Table S4**.

Screening WRKY75 mutant lines and generation of *wrky75/sr3g* double mutant

The following T-DNA insertion alleles were used for *WRKY75* (AT5G13080): *wrky75-1* (SALK_004954,
kanamycin resistant) and *wrky75-3* (SALKseq_046273, kanamycin resistant). All T-DNA insertion alleles
were in the background of Col-0 accession. The primers used for genotyping PCR are shown in **Table S4**.
Two sets of double mutant combinations were generated for *wrky75-3/sr3g-5*, with *wrky75-3* serving as
the maternal parent in one (referred to as line d) and *sr3g-5* serving as the maternal parent in the other
(referred to as line e).
250

251 **Plasmid construction and plant transformation**
252

For plant expression, all clones were constructed using the GreenGate cloning system (Lampropoulos et
al., 2013). We have generated full-length SR3G constructs, as well as three fragments - fragment A (from
amino acid number 1 to 216), fragment B (from amino acid number 217 to 468) and fragment C (from
amino acid number 469 to 503). Fragments A, B, AB, and ABC (full length) were cloned into entry vector
pGGC (GreenGate) with mVenus as N-terminal tag. SR3G fragments were PCR amplified using high-fidelity
DNA polymerase, and cloned as full-length cDNA or truncated version. The entry clone was evaluated
using Sanger sequencing to verify the absence of mutations. Promoters used included Ubiquitin 10
promoter (Norris et al., 1993). The list of constructs generated within this project can be found in **Table
S7.** The SR3G constructs generated in this paper and their corresponding vector maps can be accessed in
Zenodo   repository
(https://zenodo.org/records/10788996?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6IjI2ZjkzZGRlLTQ3M2YtND
BmZS04MmM5LTNhMjQ5MGFmOWY0NiIsImRhdGEiOnt9LCJyYW5kb20iOiI3OGNmY2UwM2YyMGFkNW
YwYjM2ZjAwOWYyM2VjNDNiNCJ9.NRWsSWHBe4srG720PncdfFjeDXQ0dATuI2BjSnca5sD1P_GAW5vhD_
bxOSBIH2f8txZObh5YRWH4Hkva0pzqXQ). The final constructs were validated via vector digestion, using
BaeGI for the constructs with fragments A, B, and AB and PvuI for the full length SR3G construct.
Subsequently, we transformed them into *Agrobacterium tumerfaciens* strain *GV3101*. Plant
transformations were conducted using the floral dip method (Clough and Bent, 1998). Transgenic plants
were selected for BASTA resistance (10 mg/L) for three generations, to ensure the homozygous and non-
segregating population of seedlings for each construct.

Agar-based plate experiments for mutant lines

Arabidopsis seeds were sterilized for 10 min with 50% bleach and rinsed five times using milli-Q water, and germinated on ½ strength Murashige and Skoog (MS) medium containing 0.5% (w/v) sucrose, 0.1% (w/v) 4-morpholineethanesulfonic acid (MES), and 1% (w/v) agar, with PH adjusted to 5.8 with KOH. After 24 h of vernalization at 4°C in the dark, the plates were placed in the Conviron growth chamber with the light intensity of 130–150 µmol x m^{-2} x s^{-1} in a 16 h light / 8 h dark cycle at 21°C and 60% humidity. For plate assays, 4 days after germination, the seedlings were transferred to ½ MS media supplemented with 0, 75, or 125 mM NaCl as indicated. The plates were scanned using EPSON scanner every other day, starting from 4 days after germination until the plants were 14 days old. To analyze root system architectural traits from the scanned plate images, we used SmartRoot plugin (Lobet et al., 2011) in ImageJ to trace the root, and extracted root-related features in the CSV format followed by data analysis in R (https://rpubs.com/mri23/913282)

Evaluation of plant performance under salt stress in soil

The seeds were germinated in ½ MS media for one week, as described for the agar-based plate experiments. One week after germination, the seedlings were transplanted to the pot (12 x 4 cm insert) containing the Cornell Mix soil (per batch combine: 0.16 m^3 of peat moss, 20.84 kg of vermiculite, 0.59 kg of Uni-Mix fertilizer, and 2.27 kg of lime) watered to 100% water holding capacity and placed in the walk-in growth chamber with the 16 h light / 8 h dark period, 22°C and 60% relative humidity throughout the growth period. When all of the pots dried down to the weight corresponding to 50% of their water holding capacity, they were soaked for 1 h in tap water or a 200 mM NaCl solution, resulting in an effective concentration of 100 mM NaCl based on the 50% soil water holding capacity, which corresponded to a moderate level of salt stress (Awlia et al., 2016). The control pots were soaked for the same length of time in 0 mM NaCl solution, to account for the soil saturation effect. We then allowed the pots to be drained for 2-3 h to eliminate excess moisture. The pots were placed under phenotyping rigs equipped with an automated imaging system (Yu et al., 2023) and the pot weight was measured daily to maintain the reference weight corresponding to 50% of the soil water holding capacity throughout the experiment. We would like to note that this gravimetric based method for application of salt stress has been developed for soils typically used for pot-grown plants, with relatively high water holding capacity (Awlia et al., 2016). Within these specific conditions, no drought stress symptoms were observed.

To evaluate the growth of soil-grown plants, we used an in-house developed automated imaging system using Raspberry Pi cameras to capture the growth dynamics over the period of two weeks (Yu et al., 2023). The collected images were processed using the PlantCV pipeline (Gehan et al., 2017), followed by the data analysis in R, as described here (https://rpubs.com/mjulkowska/sr3g_Maryam2021OctNov).

Transient expression of SR3G constructs in tobacco epidermal leaf

Agrobacterium tumefaciens strain *GV3101* containing cellular marker free GFP (*35S::GFP* with rifampicin and spectinomycin resistant)*,* generously provided by Professor Maria Harrison Lab at Boyce Thompson Institute, or mVenus-SR3G constructs (including *UBQ10::mVenus-SR3G-full length*, *UBQ10::mVenus-SR3G-fragment A*, *UBQ10::mVenus-SR3G-fragment B*, and *UBQ10::mVenus-SR3G-fragment AB*, all with spectinomycin and tetracycline resistant) were grown in liquid YEP medium with appropriate antibiotics overnight at 28°C while shaking. The following concentrations were used for antibiotics when used for transient transformation: spectinomycin (50 ug/ml), tetracycline (10 ug/ml), and rifampicin (12.5 ug/ml). Rifampicin and kanamycin (50 ug/ml) were used for P19 plasmid (generously provided by Maria Harrison Lab at BTI) as a silencing inhibitor. The following day, OD$_{600}$ was measured to be 0.3-0.5. The cultures were then centrifuged at 8000x g for 2 min. The Agrobacterium pellet was resuspended in an induction buffer

containing 10 mM MgCl₂, 10 mM MES, pH 5.6, and supplemented with 200 µM acetosyringone and was incubated for 2–4 h in the dark at room temperature. Leaves of 4-weeks-old *Nicotiana benthamiana* plants were infiltrated using a syringe. After 48 hours, the fluorescence was visualized in epidermal cells of leaf discs using confocal microscopy.

Confocal microscopy images were taken using the Leica TCS SP5 Laser Scanning Confocal Microscope with the following setting: Excitation at wavelengths of 488 nm (GFP), 514 nm (mVenus), and 561 nm (chlorophyll autofluorescence) were provided using an argon or DPSS (Diode-Pumped Solid State) laser. A spectral emission range of 505–525 nm for GFP, 526–560 nm for mVenus, and 680–700 nm for chlorophyll autofluorescence was used.

Elemental analysis

The elemental analysis was performed using Inductively Coupled Plasma Atomic Emission Spectroscopy (ICP-AES). For analysis of Na^+ and K^+ ions in roots and shoots of plate-grown seedlings, the plants were grown on ½ x MS plates as described above. After two weeks of exposure to 0, 75 or 125 mM NaCl, root and shoot tissues were harvested and measured for fresh weight, rinsed in milli-Q water and collected into separate paper bags that were dried at 60 °C for 2-3 days. Subsequently, the dry weight was recorded. Samples were digested in double distilled HNO3, followed by the addition of 60/40 nitric/perchloric acid continuing incubation at 150 °C. Samples were processed for ICP-AES analysis using a Thermo iCap 7000 ICP-AES after diluted to 10 ml with deionized water. For soil-grown plants, the entire rosette leaves were harvested at the end of week four and underwent a similar procedure as described above. Ion content was calculated per dry weight for each sample, followed by data analysis in R (sr3g:https://rpubs.com/mri23/913285, wrky75/sr3g double mutants: https://rpubs.com/mri23/1157092 for plate-grown and sr3g:https://rpubs.com/mri23/913299 for soil-grown seedlings).

Electrolyte leakage analysis

The electrolyte leakage measurement was done as previously described (Hu et al., 2015) with slight modifications. In brief, for each plant, three leaf discs were incubated in 2 ml of distilled water for 24 h at the light at room temperature with gentle shaking, followed by measuring the initial electrolyte leakage using a conductivity meter (Horiba LAQUAtwin EC-11 Compact Conductivity Meter, # 3999960125). The samples were then subjected to 80 °C for 2 h to release the total electrolyte, and cooled at room temperature for 24 h. Final electrolyte leakage was measured the following day. The electrolyte leakage percentage was calculated by dividing initial conductivity by final conductivity * 100 and followed by data analysis in R (https://rpubs.com/mri23/913288 and https://rpubs.com/mri23/1102053).

RT-qPCR analysis

Total RNA was isolated from Arabidopsis seedlings root and shoot treated with or without 75 or 125 mM NaCl for one week using the RNeasy Plant Mini Kit (Qiagen). RNA samples were cleaned at least twice using RNase-free DNase (Qiagen) to remove genomic DNA contamination. One microgram of total RNA was used for a reverse transcriptase reaction using iScript cDNA Synthesis Kit (Catalog #170-8891; Bio-Rad). RT-qPCR analysis was conducted using iQ SYBRGreen Supermix (Bio-Rad) according to manufacturer's instructions in the CFX96 real-time PCR system (Bio-Rad). *AT4G04120* (*transposable_element_gene*) was used as a housekeeping gene for data normalization. RT-qPCR experiments were conducted using three independent experiments, each with at least three biological replicates. The list of primers used is shown in **Table S4**.

Suberin staining

Suberin content was examined in 6-days-old Arabidopsis roots treated with or without 75 mM NaCl for two days using Fluorol Yellow 088 (FY; Santa Cruz Biotechnology Incorporation) staining. Seedlings grown on agar plates (as described above for mutant screens) were incubated in a freshly prepared solution of FY (0.01% [w/v] in lactic acid 85% [W/W]) at 70°C for 30 min, rinsed in milli-Q water for three times with 5 min each, and mounted on slides with 50% glycerol before confocal microscopy examination using excitation at wavelength of 488 nm and a spectral emission range of 500–550 nm. Suberin quantification was performed using the ImageJ software followed by data analysis in R (https://rpubs.com/mri23/935019).

Suberin monomer quantification using Gas Chromatography–Mass Spectrometry (GC-MS)

100 - 150 Arabidopsis seedlings were grown on the plates for four complete days, transferred to 75 mM salt plate for two days, and roots were harvested in distilled water to remove any remaining agar. Each plate was treated as one biological replicate, and five plates were used for each genotype/condition. Each plate of roots was collected into a microcentrifuge tube and dried in a speed vacuum concentrator (Thermo Electron, Waltham, MA), and weighed. Approximately 1 mg of dried root were collected from each plate. Suberin monomer quantification was performed according to (Delude et al., 2017; Kreszies et al., 2019) with modifications. Dried root tissue was incubated in 1% cellulase plus 1% pectinase in 10 mM citrate buffer (pH 3.0) for two weeks, with the incubation solution changed every three days. After the cellulase/pectinase solution was removed, chloroform/methanol solution (2:1 and 1:2, v/v) were mixed with the root tissue for three days respectively, to remove lipids. The remaining root tissue was mixed with 1 ml freshly prepared 5% H_2SO_4 in methanol (v/v) and 200 µl of toluene containing 20 µg of heptadecanoic acid as internal standard, sealed in an 8 ml capped tube and heated at 85°C for 90 min. After samples were cooled to room temperature, 200 µl hexane and 1.5 ml 0.9% sodium chloride in water are added to the capped tube with shaking. After centrifuging at 1000 g for 1 min, the upper organic phase was transferred to a glass insert of GC vial and dried. 40 µl pyridine was added to the glass insert, 40 µl N,O-bis(trimethylsilyl)trifluoroacetamide (Restek, Bellefonte, PA) was added subsequently, incubated at 65°C for 2 h. Standards of 1 µl derivatized sample was injected into GCMS (Agilent 6890-5975, Santa Clara, CA), with the inlet temperature of 250°C, column helium flow of 1 ml/min, and a split ratio of 5:1. The oven temperature was held at 50°C for 1 min, raised to 200°C at the rate of 25°C/min, held at 200°C for 1 min, raised to 320°C at the rate of 10°C/min, and finally held at 320°C for 5 min. 1-Docosanol (22:0-OH), 1-eicosanol (20:0-OH), hexadecanedioic acid (16:0-DCA), 1-hexadecanol (16:0-OH), 16-hydroxy hexadecanoic acid (16:0-ωOH), C4 - C24 Even Carbon Saturated FAMEs (Sigma 49453-U), 20-hydroxy arachidic acid (20:0-ωOH) (Cayman Chemical Company, Ann Arbor, Michigan) standards were input at 2.5 µg, 5 µg, 10 µg, and 20 µg into a mixture and processed with the same procedure as the tested samples to build the standard curve. The MS spectrum of derivatized products with ±2n carbon number of these standards was calculated and identified, referring tom the retention order in (Delude et al., 2017).

Yeast One-hybrid Assay

The respective combinations of *pB42AD*-fusion effectors and *pLacZ* reporters were co-transformed to yeast strain EGY48 (Shanghai Maokang, China). Transformants were selected and grown on SD/-Trp-Ura medium. The putative interactions were observed on SD/-Trp/-Ura + X-Gal after a 3d-incubation at 30°C. Yeast transformation and growth assay were performed as described in the Yeast Protocols Handbook (BD Clontech, USA). In brief, 631 bp of the *DUF247-150* (AT3G50150) and 953 bp of the *SR3G* (At3g50160) promoter region were separately ligated into *pLacZ* vector as the reporter, while the CDS of *WRKY75* (AT5G13080) was cloned into *pB42AD* vector as the effector. The *SR3G::pLacZ* and *WRKY75-pB42AD* were

transformed into the yeast strain EGY48(Shanghai Maokang, China). The respective combinations of
pB42AD-fusion effectors and *pLacZ* reporters were co-transformed into yeast strain EGY48.
Transformants were selected and grown on SD/-Trp-Ura medium. The putative interactions were
observed on SD/-Trp/-Ura + X-Gal after a 3d-incubation at 30°C. Yeast transformation and liquid assay
were conducted as described in the Yeast Protocols Handbook (BD Clontech, USA). The primers used for
the Y1H assay are shown in **Table S4.**

Results

Natural variation in salt stress-induced changes in root:shoot ratio

To investigate salt stress-induced changes in root:shoot ratio, we developed a custom tool that quantifies green and white pixels, corresponding to shoot and root of Arabidopsis seedlings grown on agar plates (**Fig. 1**). The projected area of root and shoot showed high correlation with root and shoot fresh weight (**Fig. S1**), indicating that projected root and shoot area can be used as an accurate proxy to non-destructively measure the increase in root and shoot biomass from the images of seedlings grown on agar plates.

Figure 1. Salt stress is causing discoordination of root and shoot growth. Arabidopsis HapMap population was screened for salt stress-induced changes in root:shoot ratio. The increase in the projected area of shoot and root (Fig. S2) was used to estimate **(A)** shoot and **(B)** root exponential growth rate. **(C)** The root:shoot growth rate ratio was calculated per genotype. The histograms represent the number of accessions across three studied conditions (0, 75 and 125 mM NaCl), whereas the population average is indicated using the dashed line. Additionally, the distribution of the genotypes within each treatment was visualized using the error plots (lower panel), where population average and standard error is indicated using a colored point and a line, respectively. Individual grey points represent individual genotypes.

Using our custom tool, we re-analyzed the data consisting of 360 Arabidopsis accessions grown on plates supplemented with 0, 75, and 125 mM NaCl (Julkowska et al., 2017), for salt stress-induced changes in the root:shoot ratio (**Fig. 1, Fig. S2**). The growth of root and shoot for each plant was estimated using the exponential function. Salt stress-induced reduction in growth rate was observed to be dose-dependent for both root and shoot (**Fig. 1A-B**). In general, the root growth was more sensitive to salt stress compared to shoot growth, with the population-wide relative growth rates reduced to a fraction of 0.67 and 0.41 of the growth rates observed at control for 75 and 125 mM NaCl, respectively. In comparison, the growth rates of the shoot were significantly reduced to 0.71 and 0.43 of the control in 75

and 125 mM NaCl treatments, respectively (**Fig. S3**). While the mean value of root:shoot growth rate did
not change upon salt stress treatment, the variance in the root:shoot ratio significantly expanded with
the increasing concentrations of salt (**Fig. 1C**). These results suggest that while root and shoot growth are
well coordinated under non-stress conditions, salt stress exposure results in loss of coordination of organ
growth across Arabidopsis accessions. To test whether there is a clear directional correlation between the
change in root:shoot ratio and overall salt stress tolerance, we have used the overall seedling size as a
proxy for plant salt tolerance (**Fig. S4, S5**). No significant correlation was found between the root:shoot
growth ratio and total seedling size (**Fig. S4, S5**), indicating that the relationship between coordination of
root and shoot growth and salt tolerance during the early seedling establishment is complex.

Identification of genetic components underlying salt-induced changes in root:shoot ratio
To identify genetic components underlying salt-induced changes in root:shoot ratio, we used the collected
data as an input for GWAS. The associations were evaluated based on the p-value, the number of SNPs
within the locus, and the number of traits associated with individual loci. As Bonferroni threshold differs
depending on the minor allele count (MAC) considered, we identified significant associations based on a
Bonferroni threshold for each subpopulation of SNPs based on MAC (**Table S3**). While we conducted a
GWAS on directly measured traits, as well as their Salt Tolerance Index (STI) values, however the amount
of associations with STI was much lower compared to directly measured traits (**Table S3**). This observation
aligns with the understanding that plastic responses to environmental conditions tend to be genetically
more complex. This complexity likely stems from the involvement of more genetic regulators compared
to low-plasticity phenotypes.
The initial GWAS was performed using 250 k SNPs (Horton et al., 2012), and led to the
identification of 14 significantly associated loci, of which 3 and 7 loci were associated with traits measured
at 75 or 125 mM NaCl treatments, respectively. We identified one salt specific association on chromosome
2 (**Fig. S6**), where two SNPs were identified within AT2G27300 encoding NAC-domain containing
transcription factor (NTL8). Additionally, salt-induced changes in root:shoot ratio and shoot:total seedling
area were associated with SNPs on chromosome 3, where 2 significantly associated SNPs were flanking
AT3G22430, encoding an unknown gene encoding a domain of unknown function (**Fig. S7**).
To refine our analysis and further prioritize candidate genes for validation, we repeated the GWAS
using a higher-density 4M SNP panel (1001 Genomes Consortium, 2016). This increased SNP density
provided more associations and highlighted loci with greater confidence due to the higher resolution. The
higher-density panel also allowed us to capture different linkage disequilibrium (LD) blocks across the
genome. One locus on chromosome 2, detected by both the 250K and 4M SNP panels, appeared to be in
a region of strong LD or possibly under selection, consistently identified by both panels. However, most
loci were not detected by the lower-density 250K panel, likely due to differences in the panels' resolution
and statistical power.
The new GWAS with the 4M SNP panel revealed 32 additional loci, with only one locus (locus 30,
Table S3). being detected by both the 250K and 4M SNP panels. We prioritized the identified loci based
on several factors, including the number of significantly associated SNPs, MAC, and the number of traits
associated with each locus. This led to the identification of three loci of major interest (**Table S4**). We
further examined the regions in LD with the significantly associated SNPs for sequence divergence and
open reading frames (**Fig. 2, Fig. S8, Fig. S9**).
Out of identified associations, locus 30 stood out as it consisted of nine SNPs associated with p-
value above the Bonferroni threshold, and was associated with both root:shoot and shoot:total seedling
area at 6 and 8 days after transfer of the seedling to 125 mM NaCl (**Fig. 2A-B, Fig. S8, Table S4**). The
significantly associated SNPs were located in and around the gene coding region of AT3G50160, which
encodes a domain of unknown function 247 (DUF247). We investigated the region for sequence
divergence, and found that the region contained not only a high degree of sequence diversity but also

potential structural variants (**Fig. 2C**). To identify allelic variation underlying the initial associations, we
performed haplotype analysis, where accessions with shared SNP pattern within the protein-coding part
of AT3G50160 were grouped into individual haplotype-groups (**Fig. 2D**). Each unique haplotype,
represented by at least three accessions, was subsequently studied for differences in observed
phenotypes. We observed that Haplotype 1, represented by 46 accessions including Col-0, was
significantly different from Haplotype 11, represented by 11 accessions including Blh-1, in shoot-per-total
seedling area at 6 days after transfer to 125 mM NaCl (**Fig. 2D**), as well as shoot growth rate and relative
shoot growth rate at 125 mM NaCl (**Fig. S9**). Upon further investigation and sequencing of AT3G50160 in
both Col-0 and Blh-1 accessions, we discovered that the Blh-1 allele carries a large insertion at 644
nucleotides downstream from the ATG of AT3G50160, followed by multiple other sequence variations
(**Fig. S10**). The first insertion leads to a premature STOP-codon within the DUF247 domain after Gly-215
(**Fig. 2E, Fig. S11**), resulting in a truncated protein that still carries a nuclear localization signal and coiled-
coil domain, but is missing a major part of the calcium-binding domain and a predicted transmembrane
domain (**Fig. S12**). As there are other DUF247s within the locus associated with salt-induced changes in
root:shoot ratio, we decided to name the gene containing majority of the associated SNPs (At3g50160) as

Figure 2. SR3G is associated with salt stress-induced changes in shoot-per-total seedling area ratio. The salt induced changes in root-to-shoot and shoot-per-total seedling area ratios were used as an input for GWAS. Manhattan plots representing the associations of shoot-per-total seedling area recorded 6 days after transfer to 125 mM NaCl and **(A)** 250 k SNPs and **(B)** 4 M SNPs. **(C)** Significant associations were found with the SNPs forming a locus on chromosome 3 in and around SR3G (AT3G50160), encoding Domain of Unknown Function 247 (DUF247). The natural variation in the LD region was studied in 162 accessions sequenced by 1001 Genomes Project. The upper panel represents portion of the missing data, upper middle panel represents deletions relative to Col-0, while lower middle panel represents the sequence similarity compared to Col-0. The bottom panel represents the open reading frames (ORFs) within the LD, and the location of associated SNPs is indicated with the dashed lines. Red dashed lines represent associations above Bonferroni threshold in 250k SNP mapping, while grey lines represent associations with -log10(p-value) > 5 in the 4M SNP mapping. **(D)** The haplotype analysis performed using SNPs located within the coding region of the SR3G revealed significant differences in shoot-per-total seedling area recorded 6 days after transfer to 125 mM NaCl between Haplotype groups 1 (represented by 46 accessions, including Col-0) and 11 (represented by 11 accessions, including Blh-1). The significant differences between individual haplotype groups were tested using ANOVA with Tukey HSD to identify significantly different groups. **(E)** Upon further sequencing of SR3G in accessions from Haplotype 1 and 11, revealed two 200 bp insertions within SR3G exons in 4 tested accessions belonging to haplotype 11 group. These 200 bp insertions resulted in a premature STOP-codon within the DUF247 domain (after Gly-215).

487 Salt Root:shoot Ratio Regulator Gene (SR3G). Together, these results suggest that the allelic variation in
488 SR3G protein sequence might underlie the observed differences in root:shoot ratio under salt stress
489 conditions.

Evolutionary context of SR3G locus reveals its origin in local genome duplication

SR3G sits in what appears to be a tandemly duplicated array of DUF247s along Arabidopsis chromosome
3. The genome-wide identification of SR3G included eight DUF247 paralogs, AT3G50120, AT3G50130,
AT3G50140, AT3G50150, AT3G50170, AT3G50180, AT3G50190, and AT3G50200. Reciprocal best BLAST
within CoGe's BLAST tool of these nine DUF247 paralogs against the genomes of seven other Brassicaceae
(see **Table S5**) revealed a total of 50 potential orthologs used for downstream analysis. While the copy
number of DUF247 loci are similar across the examined species, the noted exceptions are those with
recent whole genome duplication events (*Brassica rapa* and *Camelina sativa*). Given these data, it appears
that DUF247 loci are commonly tandemly duplicated, although the eight paralogs observed in Arabidopsis
represents an extreme case for this gene family.
To examine conservation within the predicted DUF247 functional domains, we generated an
amino-acid based multiple sequence alignment using MUSCLE. All of the identified DUF247 proteins
contained the DUF247 domain near the center of the translated sequence. While partial deletions were
identified outside of the DUF247 domain at the C-terminus of a few DUF247s, such as *CARHR159180*,
Csa09g052970, and *Bra01902*, in general the DUF247 domains were highly conserved across Brassicaceae.
The amino acid sequences spanning the "coiled-coil domain" (Amino acid position: Met153-Lys173),
putative Ca^{2+} binding domain (Amino acid position: Gly210-Phe372), and super polar fragment (Amino
acid position: Ser301-Ile320) were further selected for phylogenetic reconstruction (**Fig. 3B**). The resulting
phylogeny demonstrated that SR3G is sister to a clade containing AT3G50150 and representatives from
C. sativa, *Capsella rubella,* and *A. lyrata*, with representative DUF247s from *B. rapa* and
Eutrema salsugineum serving as a monophyletic outgroup. These data support a hypothesis where SR3G
is a recent duplicate (likely A/C Clade specific; Forsythe et al., 2020) of AT3G50150 and that representative
SR3G sequences in *C. sativa*, *C. rubella*, and *A. lyrata* have been lost.
Given the specific retention of AT3G50150 in *A. thaliana*, we next tested whether this locus is
experiencing directional selection. To test this assumption, we grouped the branches leading to SR3G into
four groups (**Figure 3B**: ω0, ω1, ω2, and ω-rest) and then performed a branch-site model A hypothesis
test**.** In our null hypothesis (H_0), we hypothesized that all branches have the same selection pressure (H_0:
ω -rest branches = ω0 = ω3 = ω1 = ω2; see **Table S6** for omega values for each hypothesis). For our
alternative hypothesis, we created five different ω groups to cover scenarios where specific branches may
be experiencing directional selection (See Methods). Based on our analysis, the branch leading to SR3G is
experiencing directional selection (likelihood score = 20625.5, Chi-square derived P value; P = 0.00418).
Interestingly, no other branches appeared to be experiencing the same selection. To pinpoint specific
amino acids within SR3G under selection, we performed the Bayes Empirical Bayes (BEB) test, identifying
three sites with a posterior probability > 0.90, including 124C (P = 0.992), 206A (P = 0.951), and 209A (P =
0.988). Two out of three sites are located at the putative Ca^{2+} binding site and all three sites possess a
unique amino acid compared with the other 49 amino acid sequences (**Fig. 3C**).

Figure 3. Phylogeny and positive selection analysis of SR3G and its orthologs from other species. The orthologous genes of SR3G (AT3G50160) from other seven species were identified using Go-GeBlast (E-value < 1.00E-10). **(A)** The protein sequences of the 50 homologous genes were aligned by MUSCLE and then an unrooted phylogenetic tree was reconstructed by a Maximum -likelihood (ML)-derived style by RAxML (Bootstrap number: 100). **(B)** A positive selection analysis was conducted by branch-site model A test: Specifically, we hypothesized the differential or constant dN/dS (ω) substitution rate among the closely distant branches leading to DUF247 (AT3G50160) (See Method & Materials for details). These branches were marked as ω1 (light brown branch), ω2 (DUF247-specific branch), ω3 (light blue branch), and ω-resvt (light purple branch). **(C)** The gene structure of DUF247 (AT3G50160) was illustrated by wide arrows (grey arrow: 5 and 3' UTR, yellow arrow: coding-region). The regions used for cloning (Fragment A, B, and C) were marked by narrow arrows along with "Coil-Coil Domain" and "Super-polar Fragment" marked on Fragment A and B, respectively. Lastly, the Bayes Empirical Bayes (BEB) was performed to test the probability of sites along with ω > 1 over the DUF247 (AT3G50160). Three sites were generated with a posterior probability > 0.90: 124C (P = 0.992), 206A (P = 0.951), and 209A (P = 0.988). These three sites were marked with an orange asterisk and assumed to have been undergoing positive selection.

SR3G expression is enhanced in root by salt stress

To identify the developmental window and specific plant tissues where the *SR3G* gene is active, we
examined its expression pattern. Based on Arabidopsis eFP Browser (http://bar.utoronto.ca/), *SR3G*
expression is relatively low, and restricted to the early stages of seedling development (**Fig. S13A**). Upon
examining the expression profiles specific to different tissues, we observed that *SR3G* is predominantly
expressed in the root stele (**Fig. S13B**). Moreover, *SR3G* expression was induced by salt stress (**Fig. S13C**).
To validate these results, we performed RT-qPCR on 11 days old seedlings of Col-0 exposed to 0 or 75 mM
NaCl for 7 days. In Col-0 seedlings, the transcript levels of *SR3G* were significantly higher in shoot

compared to the root tissue (**Fig. 4A**). Interestingly, the *SR3G* transcripts increased only in root upon salt
stress exposure but not in shoot (**Fig. 4A**). These
results suggest that *SR3G* is the most prominent
DUF247 studied within our study to affect root
development under salt stress.

We further examined expression profiles of other
DUF247 genes that directly neighbor *SR3G* in
response to salt stress using the eFP Browser. We
observed that the expression of DUF247-120
(At3g50120) remained undetectable under control
conditions, except for a slight increase in rosette
leaves after 6 hours of salt stress exposure (**Fig.
S14**). For DUF247-130 (At3g50130), -140
(At3g50140), -150 (At3g50150), and -170
(At3g50170), high expression was detected in roots
under non-stress conditions, but expression levels
decreased upon salt stress (**Fig. S14**). The
expression of DUF247-180 (At3g50180) remained
unchanged with or without salt stress (**Fig. S14**).
DUF247-190 (At3g50190) exhibited high expression
in roots during early salt stress exposure, which then
declined at later stages (**Fig. S14**). For DUF247-200
(At3g50200), its already low detectable expression

Figure 4. Transcript abundances for the two closely related DUF247s. Expression of **(A)** SR3G (AT3G50160) and **(B)** its closest homologue DUF247-150 (AT3G50150) were measured in Col-0 seedlings grown with and without salt stress. RT-qPCR analyses were conducted using seedlings grown on 1/2x MS for 4 d and then followed by transferring to the treatment plates with or without 75 mM NaCl for one week. Mean values are shown ± SE, with number of replicates (n) shown in each graph. AT4G04120 was used as a reference gene for normalization. Significance was determined by the Tukey–Kramer HSD test in JMP.

disappeared entirely during prolonged salt stress exposure (**Fig. S14**). Because *DUF247-150*
(At3g50150) is closely located to the *SR3G* in chromosome 3, we investigated its gene expression
in young seedlings. We observed that its expression was significantly higher in shoot compared to
the root tissue. However, its expression remained unchanged in both root and shoot under salt
stress (**Fig. 4B**). Together, these results suggest that *SR3G* is the most prominent DUF247 studied
herein involve in regulation of root development under salt stress.

SR3G relocates from plasma membrane to nucleus/cytosol after removal of its transmembrane

domain

SR3G protein is predicted to contain a nuclear localization signal (Lys15-Ala19), a coiled-coil domain
(Met153-Lys173), a Ca^{2+} binding pocket (Gly210-Phe372), and a trans-membrane domain (Pro469-
Thr503) (**Fig. S15**). We generated constitutive expression constructs of full-length *SR3G* gene CDS, and a
combination of individual domains (refer to as A, B, and AB domains) to identify their subcellular
localization (**Fig. S15**). Fragment A comprised a nuclear localization signal, a coiled-coil domain, and the
initial segment of the Ca^{2+} binding pocket, encompassing seven amino acids. Fragment B encompassed
the remaining segment of the Ca^{2+} binding pocket. Fragment C only included the transmembrane domain.
All constructs contained N-terminal fusion of mVenus. The individual constructs were used in transient
expression assay in tobacco leaves. Our result showed that full-length SR3G localizes to the plasma
membrane (**Fig. 5A**) as its signal overlapped with FM4-64, the plasma membrane marker (**Fig. 5B**). The
SR3G fragments from which the transmembrane domain was excluded—namely fragments A, B, and AB—
exhibited localization in both the nucleus and cytosol, resembling the pattern observed with free-GFP
localization (**Fig. 5C-F**). It is noteworthy that while the signal from fragment A was directed to both the

cytosol and nucleus, its intensity was notably stronger in the nucleus (**Fig. 5C**), aligning with the prediction
that it contains the nuclear localization signal. Together, these findings indicate that the positioning of
SR3G within the plasma membrane is due to the transmembrane domain encoded within the fragment C,
and its removal could potentially release the protein into both the nucleus and cytosol.

(C-E)

Figure 5. SR3G full-length protein resides in the plasma membrane while removal of its transmembrane domain results in protein relocation to nucleus and cytosol. (A-B) SR3G full-length protein or (C-E) truncated versions fused to mVenus at the N-terminus were agro-infiltrated into the Nicotiana benthamiana leaves for transient expression. Shown are a bright-field image of transfected leaf cells and mVenus-SR3G-mediated fluorescence as well as GFP-based subcellular marker. **(A)** Localization of full-length SR3G alone or **(B)** in combination with the known plasma membrane dye, FM4-64. Localization of truncated versions of SR3G are shown for **(C)** SR3G-Fragment A, **(D)** SR3G-Fragment B, and **(E)** SR3G-Fragment AB. **(F)** Free GFP was used as a nuclear and cytosolic marker. Scale bar = 25 μm.

Lower expression of *SR3G* results in reduced lateral root development under salt stress

To investigate role of *SR3G* in salt-induced changes in root growth and development, we studied the root system architecture of two T-DNA insertion lines: *sr3g-4* (SAIL_690_E12) and *sr3g-5* (SAIL_608_C06) localized in second exon and first intron region of the *SR3G* gene, respectively (**Fig. S16A**). Among these two mutant lines, *SR3G* expression was reduced in the *sr3g-5* mutant compared to Col-0 (**Fig. S16B**), indicating the *sr3g-5* is a knockdown mutant. We did not observe any differences in *SR3G* transcript abundance between *sr3g-4* and Col-0 (**Fig. S16B**). Moreover, we examined the transcript abundance of the closely related gene (i.e., *DUF247-150, At3g50150*) in the *sr3g-5* mutant, and found no significant difference. In the case that *DUF247-150* would act redundantly to *SR3G*, we would expect that the expression of *DUF247-150* would be increased in *SR3G* knockdown mutant. As this is not the case, these results further contribute to the evidence that *SR3G* and *DUF247-150* are not acting redundantly (**Fig. S16C**).

Two T-DNA insertion lines were subsequently studied for salt-induced alterations in root architecture (**Fig. 6 and Fig. S17 and S18**). Among the various features of the root system architecture that we investigated (**Fig. S17**), *sr3g-5* exhibited a significantly reduced growth rate of lateral roots under 75 mM NaCl, but not under 125 mM NaCl, when compared to Col-0 (**Fig. 6A**). Likewise, the *sr3g-5* mutant displayed a significant reduction in the Stress Tolerance Index (STI), calculated as the ratio of growth rate under salt to growth rate under control conditions, for lateral root length under 75 mM NaCl (**Fig. 6B**). No significant difference was observed between *sr3g-4* and Col-0 (**Fig. S18**), except for Stress Tolerance Index (STI) calculated using growth rates of lateral root length under 125 mM salt stress (Fig. S18-E). In this case, the sr3g-4 mutant exhibited a non-significant decrease compared to Col-0, consistent with the observations made for the *sr3g-5* mutant line (**Fig. 6B).** Based on the RT-qPCR results (**Fig. S16**), we proceeded to further investigate the role of *SR3G* in salt stress, focusing on the sr3g-5 mutant in our subsequent investigations.

As Na^+ exclusion from the shoot is one of the main mechanisms of salt tolerance (Julkowska et al., 2017; Møller et al., 2009; Munns and Tester, 2008), we further investigated the role of *SR3G* in ion exclusion using plate-grown seedlings. We used root and shoot samples collected for Col-0 and *sr3g-5* mutant to examine Na^+ and K^+ accumulation using ICP-AES at 14 days after exposure to salt stress. We observed no significant differences between Col-0 and *sr3g-5* mutant for Na^+ and K^+ accumulations and their corresponding ratio upon salt stress exposure (**Fig. S17**, **Fig. 6C**). These results suggest that *SR3G* plays a role in root development under the salt, but does not affect ion accumulation at early developmental stages.

To evaluate the effects of other closely related DUF247 genes on root development, we have investigated the T-DNA insertion lines targeting the neighboring DUF247 genes (AT3G50120, AT3G50130, AT3G50140, AT3G50150, AT3G50170, AT3G50180, AT3G50190, AT3G50200), and RNAi lines targeting one or more DUF247s within the interval (**Table S4, Fig. S19-S23**). For the screened T-DNA insertion lines (**Fig. S19-20**), we have not observed consistent changes in main or lateral root development in either control or salt stress conditions across alleles targeting the same gene. The only exception was *DUF247-200* encoded by AT3G50200, where two out of three studied T-DNA insertion alleles exhibited increased salt tolerance index for main root growth rate (**Fig. S20**).

For the screened RNAi lines (**Table S4, Fig. S21-S23**), we observed the most interesting root architecture phenotypes in lines targeting the neighboring genes of the *SR3G* (RNAi line #04, targeting AT3G50150, RNAi line #16 targeting AT3G50130, *SR3G*, and AT3G50170, and RNAi line #19 targeting AT3G50130, AT3G50140, AT3G50150, *SR3G*, AT3G50170, AT3G50180, and AT3G50190). Three out of five screened independent transformation lines from RNAi line #04 showed a decreased relative main root length (Salt / Control, CSHL #4 B2a, D4a, and G1d), whereas two lines showed a decrease in relative lateral root number (CSHL #4 D4a and F1b), and one line showed decreased lateral root length at 75 mM NaCl

(CSHL #4 D4a **Fig. S21-23**). Two out of two independent transformants for RNAi line #16 showed a
decreased relative main root length and increased lateral root number at 75 mM NaCl (CSHL #16 A2c and
B1a, **Fig. S21-23**). In three of the four screened independent transformants of line #19 we observed an
increased lateral root length at 75 mM NaCl (CSHL #19 F1a, I3c, and L1c **Fig. S21-23**). We did not observe
strong reduction in lateral root development across RNAi lines targeting two closely related DUF247 genes
(i.e., RNAi line #22, targeting AT3G50150 and *SR3G*), as in the single mutant (**Fig. 6, Fig. S21**). The above
results further support that these two genes do not act redundantly in the context of lateral root
development.

Figure 6. *sr3g* mutant displays reduced growth rate for the lateral root length. Root system
architecture analysis of Col-0 and *sr3g-5* plants under various concentrations of NaCl are shown
here. **(A)** The representative images of 13-day-old Col-0 and sr3g-5 genotypes that experienced
9 days of salt treatment at indicated concentrations as well as growth rate for lateral root. **(B)** Salt
Tolerance Index (STI) for the main root length, average lateral root length, and lateral root number
at 75 mM NaCl. The STI was calculated by dividing the growth rate measured under salt stress by
the growth rate measured under control condition for each genotype. **(C)** Na$^+$/K$^+$ ratio in root and
shoot of Col-0 and *sr3g-5* after two weeks on 75 mM salt are shown. Each dot in (A) and (B)
represents individual replicate per genotype. Lines in (A) and (B) graphs represent median and
average, respectively. In (A) and (B), the asterisks above the graphs indicate significant differenc-
es between Col-0 and the *sr3g-5* mutant, as determined by the Student 's t-test: *P < 0.05. Statisti-
cal analysis in (C) was done by comparison of the means for all pairs using Tukey–Kramer HSD
test. Levels not connected by the same letter are significantly different (P < 0.05). Root system
architecture and ICP-AES analyses of Col-0 and *sr3g-5* mutant are shown in details in Figure S17.

Loss of *SR3G* results in increased salt tolerance in soil

To investigate *SR3G's* contribution to salt stress tolerance beyond seedling stage and agar plate
conditions, we evaluated the performance of Col-0 and *sr3g-5* mutant in soil. We exposed the soil-grown
plants to salt stress and imaged the rosette size every 30 minutes for two weeks using the automated
imaging platform (Yu et al., 2023). As salt-induced changes depend on the developmental stage at which
the stress is applied (Julkowska et al., 2017), we examined the role of *SR3G* exposed to salt stress at 2 or
3 weeks after germination, corresponding to early and late vegetative stages. We calculated a daily growth
rate for each plant, using the projected rosette area. No significant difference was observed between Col-
647

0 and *sr3g-5* under control conditions (**Fig. 7A-B**), indicating that *SR3G* does not play a role in rosette
growth and development under non-stress conditions. Significant difference was observed between Col-
0 and *sr3g-5* for plants exposed to early salt treatment (**Fig. 7A-B**), where *sr3g-5* mutant displayed an
increased rosette growth rate as compared to Col-0 (**Fig. 7A**). Late stress exposure resulted in significantly
higher *sr3g-5* growth rate at two and three days after exposure to salt stress (**Fig. 7A**). Similarly, the *sr3g-*
5 mutant showed a significantly higher rosette area compared to Col-0 exclusively when plants were
exposed to salt stress at an earlier stage of their development (**Fig. 7B**). No significant difference was
observed between Col-0 and *sr3g-5* mutant in terms of water content under either of conditions tested,
providing no substantial evidence for SR3G to ~~to~~ regulate water content (**Fig. S24**).

To investigate the role of *SR3G* in the maintenance of cellular integrity during salt stress, we
evaluated the salt-induced changes in cell membrane integrity by measuring ion leakage in the leaf discs
at the end of the experiment (**Fig. 7C**). While ion leakage increased in Col-0 under early salt stress
conditions, no noticeable salt-induced change was observed for *sr3g-5* compared to control condition
(**Fig. 7C**). Although the *sr3g-5* mutant exhibited significant increase in ion leakage in response to late salt
stress, this phenomenon may be more closely associated with the onset of senescence rather than direct
exposure to salt stress (**Fig. 7C**).

We further explored the role of *SR3G* in ion exclusion by quantifying the accumulation of Na^+ and
K^+ ions in the rosette at the end of the experiment (4-weeks-old plants). We observed a reduced
accumulation of Na^+ in *sr3g-5* mutants compared to Col-0 for plants exposed to salt stress at an earlier
timepoint (**Fig. 7D**). No significant differences in Na^+ accumulation was observed between the Col-0 and
sr3g-5 at the late salt stress regime (**Fig. 7D**). A subtle, non-significant, reduction in K^+ content was
observed in *sr3g-5* mutant compared to Col-0 across all three conditions studied (**Fig. 7E**). In summary,
these results suggest that *SR3G* plays a negative role in maintenance of plant growth, cellular integrity
and Na^+ accumulation during the salt stress, especially when the salt stress is applied during early
vegetative development.

Figure 7. sr3g mutant has higher growth rate and larger rosette area while displaying less ion leakage and Na⁺ accumulation in shoot under salt stress. The salt stress responses of 2-week-old (referred to as "early salt stress") or 3-week-old (referred to as "late salt stress") soil-grown Col-0 and *sr3g-5* plants that were exposed to a final concentration of 100 mM NaCl were examined here. **(A)** Rosette growth rate and **(B)** area were monitored over a period of two weeks, during which the "early salt stress" and "late salt stress" groups were exposed to two or one week of salt stress respectively. Each line represents the trajectory of individual plant throughout time of experiments, where red and blue lines indicate Col-0 and *sr3g-5* plants respectively. The dashed lines represent the mean values of the genotype per condition, whereas the grey band represents the confidence interval. The significant differences between genotypes, determined by one-way analysis of variance, are illustrated in a plot below each graph, with dashed red line representing a threshold corresponding to *p-value* of 0.05. Representative images of 4-week-old Col-0 and *sr3g-5* plants were shown in (A). **(C)** Ion leakage percentage, **(D)** Na⁺ and **(E)** K⁺ contents in shoot were measured in 4-week-old plants. Statistically different groups were determined using Tukey–Kramer HSD test. Groups that are not assigned the same letter are significantly different (P < 0.05).

sr3g mutant displays enhanced root suberization

As *SR3G* limits shoot growth, promotes cell damage and sodium translocation into shoot (**Fig. 7**), and is expressed in the most inner layers of the root (i.e., stele, **Fig. S13**), we hypothesized that *SR3G* plays a role in the regulation of suberization within the Casparian strips located in the endodermis, the layer adjacent to the stele. Salt stress exposure increases root suberization, thereby limiting Na$^+$ loading into the transpiration stream and accumulation in the shoot (Baxter et al., 2009). We examined root suberization patterns in Col-0 and *sr3g-5* under non-stress and salt stress conditions. Root suberization was visualized using lipophilic stain Fluorol Yellow (FY), and staining intensity was examined under a confocal microscope. Additionally, we examined the developmental transition in root suberization through imaging four regions along the main root (**Fig. 8**). Visually, we observed that roots of *sr3g-5* were more suberized compared to Col-0 under both control and salt stress conditions (**Fig. 8A**). The quantification of FY signal intensity revealed non-significant, but substantially higher levels of root suberization in *sr3g-5* compared to Col-0 in sections one to three of the root under control condition (**Fig. 8B**). Only in root section four, the level of suberization significantly decreased in *sr3g-5* compared to Col-0 (**Fig. 8B**). In plants exposed to salt stress, we observed a non-significant difference between *sr3g-5* and Col-0 throughout the four sections, although, in section one which is the root tip proximal, *sr3g-5* showed more suberization compared to Col-0 (section 1, **Fig. 8A**). These results suggest that *SR3G* may play a negative role in root suberization.

To validate the FY histochemical analysis, we quantified root suberin monomers in the Arabidopsis seedlings using GC-MS analysis. Suberin monomers that were detected in the Arabidopsis roots included fatty acid (FA), fatty alcohol (OH-FA), α,ω-dicarboxylic acids (DCA), and ω-hydroxy fatty acid (ωOH-FA). Roots of *sr3g-5* contained significantly more total FA and DCA monomers compared to Col-0 under control condition, whereas the total OH-FA and ωOH-FA remained unchanged (**Fig. 8C**). In plants exposed to salt stress, the *sr3g-5* mutant showed an increased abundance of three suberin monomers, with DCA and ωOH-FA showing the most significant changes (**Fig. 8C**). The total FA decreased significantly in the mutant compared to Col-0 (**Fig. 8C**). Together, these results further corroborate with FY staining, suggesting that SR3G plays a negative role in root suberization.

Figure 8. sr3g mutant root is more suberized than Col-0. (A) Representative images of fluorol yellow (FY) staining of four root sections (# 1 to 4, from root tip to root base as shown by white rectangles across the seedling) were shown for Col-0 and sr3g-5 mutant with or without 75 mM NaCl. Seeds were germinated on the 1/2 MS plates for 4 d and then seedlings were transferred to the treatment plates with or without 75 mM NaCl for 2 more days. **(B)** Quantification of FY signal intensity for each root section. The FY quantification was done using ImageJ. Scale bar is equal to 500 μm in all. **(C)** Suberin monomers detected using Gas Chromatography–Mass Spectrometry (GC-MS) in the Col-0 and sr3g-5 mutant roots with or without 75 mM NaCl included fatty acid (FA), fatty alcohol (OH-FA), α,ω-dicarboxylic acids (DCA), and ω-hydroxy fatty acid (ωOH-FA). The significant differences between Col-0 and sr3g-5 mutant were evaluated using Student 's t-test.

SR3G expression is regulated by *WRKY75* transcription factor

To identify potential transcriptional regulators of *SR3G*, we examined the *SR3G* promoter for cis-regulatory elements using the CisCross tool (https://plamorph.sysbio.ru/ciscross/), and made a list of transcription factors that putatively interact with *SR3G* promoter (**Fig. 9A**). As the root stele showed the highest enrichment in the *SR3G* transcript (**Fig. S13**), we examined the transcription factors that were previously identified to be expressed in root stele (Brady et al., 2011). Cross-referencing of these two datasets revealed one transcription factor, *WKRY75* (AT5G13080) that was both enriched in root stele, and potentially binding to *SR3G* promotor.

      WRKY75 was previously described to act as a positive regulator of salt stress response by enhancing *SOS1* expression (Lu et al., 2023). To investigate whether the *SR3G* expression depends on *WRKY75*, we examined the *SR3G* expression in the *wkry75* T-DNA insertion lines. *SR3G* expression was undetected in *wrky75* knock-down mutants (**Fig. 9B**), suggesting that *WKRY75* regulates *SR3G* expression. In comparison, expression of *WRKY75* remained unchanged in the *sr3g-5* mutant when compared to Col-0 (**Fig. 9C**). To investigate whether *WRKY75* directly binds to the *SR3G* promoter, we performed Y1H assay. Similar to the two positive controls, the strains transformed with WRKY75-AD and SR3G:pLacZ grew on the SD/-Trp/-Ura + X-Gal medium, thereby confirming the interaction between *WRKY75* and *SR3G* promoter region (**Fig. 9D**). In contrast, *WRKY75* did not show binding affinity to the promoter region of the adjacent gene to the *SR3G* (i.e., DUF247-150, AT3g50150), thereby reinforcing the uniqueness of the interaction between the *SR3G* and *WRKY75* (**Fig. 9E**).

Figure 9. WRKY75 binds to the promoter region of the SR3G but not to its neighboring DUF247. (A) *In silico* searches in CisCross (https://plamorph.sysbio.ru/ciscross/CCL_index.html) shows potential transcription factors with binding sites on the main SR3G (At3g50160) promoter region. A black-arrow points to the WRKY75 (AT5G13080). RT-qPCR showing expression of SR3G (AT3G50160) **(B)** and WRKY75 (AT5G13080) **(C)** genes, respectively, in Col-0, sr3g-5 mutant, and two different mutant alleles of the wrky75, i.e., wrky75-1 and wrky75-3. RT-qPCR analyses were conducted using seedlings grown on 1/2x MS for 4 d and then followed by transferring to the treatment plates with or without 75 mM NaCl for one week. Mean values are shown ± SE, with three biological replicates used in each condition and genotype. AT4G04120 (transposable_element_gene) was used as a reference gene for normalization. Significance was determined by the Tukey–Kramer HSD test in JMP. Levels not connected by same letter are significantly different. Yeast one-hybrid (Y1H) assay showing WRKY75 (438 bp) binds to the promoter region of the main SR3G (AT3G50160, 953 bp) **(D)** but not to its neighboring promoter, i.e. DUF247-150 (At3g50150, 631 bp) **(E)**. pB42AD (AD) and pLacZ were used as effector and reporter construct, respectively. Effector and reporter constructs were co-transformed into yeast strain EGY48. Transformants were selected and grown on SD/-Trp-Ura medium. The interactions were observed on SD/-Trp/-Ura + X-Gal medium. Empty vector expressing AD domain and pLacZ were used as negative control. The two positive controls are NIGT1.4-GolS2 and NIGT1.4-GAE1. The oligo sequences used for Y1H and luciferase assays were provided in Table S4.

722
To further examine the functional consequences of the relationship between SR3G and WRKY75,
we generated *wrky75-3/sr3g-5* double mutants, and examined their root system architecture alongside
the single mutants and Col-0. Under control conditions, both *sr3g-5* and *wrky75-3* mutants showed
significant reduction in main root length compared to Col-0 and the two double mutants (**Fig. 10A**). Under

75 mM salt, only *wrky75-3* showed significant reduction in main root length compared to all other
genotypes tested, confirming the previously described salt sensitivity for *wrky75* mutant under salt stress
(Lu et al., 2023).   We have not observed any differences in main root length under 125 mM salt across all
genotypes tested (**Fig. 10A**). Regarding lateral root number (**Fig. 10B**) and length (**Fig. 10C**), we observed
significant reduction in lateral root number and length for the two single mutants, *sr3g-5* and *wrky75-3*,
compared to other genotypes only under control condition (**Fig. 10B-C**). We further investigated Na$^+$ and
K$^+$ accumulations as well as Na$^+$/K$^+$ ratio in plate grown seedlings (**Fig. 10D, Fig. S25**). At 75 mM salt
concentration, a subtle non-significant reduction in root Na$^+$/K$^+$ ratio was observed in the *wrky75* single
mutant compared to the other genotypes tested. Conversely, shoot Na$^+$/K$^+$ ratio exhibited a slight non-
significant decrease in the two double mutants compared to the other three genotypes at 75 mM salt
concentration. At 125 mM salt concentration, root Na$^+$/K$^+$ ratio non-significantly decreased in the two
double mutants compared to Col-0 and the two single mutants. Similarly, the two double mutants
maintained lower Na$^+$/K$^+$ ratios compared to Col-0 and the two single mutants in shoot at 125 mM salt
concentration (**Fig. 10D**), indicating that the *SR3G* mutation indeed compensated for the increased Na$^+$
accumulation observed in the *wrky75* mutant under salt stress.

Figure 10. wrky75/sr3g double mutants roots exhibit reduced sensitivity to salt stress and accumulate lower levels of Na+ in their shoots.
(A-C) Root system architecture analysis for the Col-0, sr3g-5, wrky75-3, and two double mutants under various concentrations of NaCl are shown here. Main root growth rate **(A)**, Lateral root number increase **(B)**, and Lateral root length increase **(C)** are shown for the indicated genotypes at various salt concentrations. **(D)** Na+/K+ ratio in root and shoot of indicated genotypes after two weeks on treatment plates. Each dot represents individual replicate per genotype. The asterisks above the graphs (A-C) indicate significant differences between Col-0 and the genotype, as determined by the Student 's t-test: *P < 0.05 and **P < 0.01. ns denotes no statistically significant. Statistical analysis in (D) was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not connected by the same letter are significantly different (P < 0.05).

Discussion

Exposure to abiotic stress alters plant architecture by affecting biomass distribution between individual organs (Guo et al., 2020; Julkowska and Testerink, 2015). Exposure to drought alters the distribution of the biomass between root and shoot, to optimize water uptake, while limiting transpiration (Asch et al., 2005; Chen et al., 2022; Karcher et al., 2008; Uga et al., 2013). While the processes underlying salt-induced changes in shoot and root growth dynamics and architecture have been systematically described (Ariel et al., 2010; Awlia et al., 2021; Deolu-Ajayi et al., 2019; Duan et al., 2013; Geng et al., 2013; Julkowska et al., 2017; Yutao Zou et al., 2022), the methodological exploration of biomass distribution between root and shoot for plants exposed to salt stress is underexplored in the current body of research. Here, we developed a tool that allows a non-destructive quantification of root and shoot projected area of the seedlings grown on agar plates (**Fig. 1A**). This new tool allows us to efficiently describe changes in biomass accumulation and stress-induced alterations in root and shoot growth. Although the method is restricted to small plants and early developmental stages, it provides valuable insight into stress responses. The application of this tool to an Arabidopsis natural diversity panel revealed that salt stress exposure results in loss of coordination between root and shoot growth (**Fig. 1D**), rather than a clear directional change, as in the case of drought. To our knowledge, this is the first systematic study of changes in root:shoot ratio. The developed tool for quantifying root and shoot size in plate-grown Arabidopsis has the potential to significantly advance research by enabling developmental and stress studies, focusing on coordination of root and shoot development during seedling establishment. Facilitating detailed comparisons of root:shoot ratios, this tool can aid in formulating new hypotheses about the mechanisms contributing to seedling vigor and stress tolerance.

The natural variation in salt-induced changes in root:shoot ratio revealed a clear association with SNPs on chromosome 3 within *SR3G*, that is situated within a region containing eight other *DUF247* genes (**Fig. 2**). SR3G was previously associated with self-incompatibility in grasses (Crain et al., 2020; Lian et al., 2021), as well as with lesions and dwarfism in Arabidopsis (Kondou et al., 2013) through constitutive activation of defense responses. However, the *SR3G* associated with constitutive defense response was AT3G60470 (Kondou et al., 2013), which is located outside the interval associated with salt stress-induced changes in root:shoot ratio (**Fig. 2C**). Because we did not observe any consistent phenotypes in the T-DNA insertion lines of *DUF247s* neighboring the *SR3G* (**Fig. S19-20**), we examined the RNAi lines targeting multiple *DUF247s* including *SR3G* (**Fig. S21-23**). Our root system architecture analyses revealed no redundancy between *SR3G* and other DUF247s, and the reduced lateral root development in double mutants of RNAi lines simply mimics the phenotype of *sr3g-5* single mutant (**Fig. 6, Fig. S21**). Further, we demonstrated that only *SR3G* is regulated by *WRKY75* (**Fig. 9B-E**). As these genes are result of tandem duplication, they are expected to accumulate a lot of variation in the cis-regulatory regions. Therefore, differential regulation of two closely related genes can be expected by the accumulated variation within the promoter region.

The results acquired in this study provide multiple layers of evidence that support the role of *SR3G* as a negative regulator of salt stress response in Arabidopsis. Loss-of-function *SR3G* maintained higher shoot growth rate and larger rosette area when exposed to salt stress in the soil (**Fig. 7A-B**), probably due to reduced Na^+ accumulation (**Fig. 7D**), which in turn resulted in lower rates of cellular damage (**Fig. 7C**). This increased salt tolerance feature can be explained by the increased root suberization (**Fig. 7**) as suberin-deficient mutants show susceptibility to salt stress (Barberon et al., 2016; Shukla et al., 2021; Ursache et al., 2021). For example, Arabidopsis *CYP86A1/HORST* (At5g58860) encodes a cytochrome P450-dependent enzyme that is responsible for catalyzing fatty acid ω-hydroxylation. It plays a pivotal role in the biosynthesis of aliphatic suberin, particularly expressed in the root endodermis. Knock-out mutations of *CYP86A1/HORST* result in a substantial decrease of approximately 60% in total root suberin

(Höfer et al., 2008; Li et al., 2007). Consequently, the diminished aliphatic suberin content in the roots of
cyp86a1 mutants leads to increased permeability to water and solutes (Ranathunge and Schreiber, 2011)*.*
Interestingly, *SR3G* expression is enhanced by salt stress (**Figs. 4, Fig. S13**), specifically in the root stele
(**Fig. S13B**). The root stele encompasses vascular bundles, procambium and pericycle, which gives rise to
the secondary growth and lateral root primordia respectively (Birnbaum et al., 2003). The stele is central
to the systemic transport of water, minerals, and nutrients within the plant (Li et al., 2017; Ramachandran
et al., 2020; Zarebanadkouki et al., 2019), and to the exclusion of salt ions from the transpiration stream
(Møller et al., 2009).

The deposition of hydrophobic residues of lignin and suberin in the Casparian strips poses a
significant barrier to apoplastic transport (Foster and Miklavcic, 2017). Previously, silencing genes involved
in Casparian strips formation resulted in increased lateral root development (Wang et al., 2020). As the
suberization of the Casparian strips increases upon salt stress, it limits the accumulation of Na^+ and loss
of K^+ ions from the root stele (Barberon et al., 2016). However, the effects of increased root suberization
under salt stress have thus far not been described to affect lateral root development. Here, we have
identified *SR3G* as a negative regulator of root suberization under salt stress (**Fig. 8**), which provides an
additional layer of protection against sodium accumulation (**Fig. 8**), but negatively impacts lateral root
development (**Fig. 6A-B**).

As SR3G was previously not described in the context of salt stress, we attempted to identify its
molecular context through its transcriptional regulation. Through our *in-silico* analysis, we have identified
WRKY75 as a putative regulator of *SR3G* expression, and confirmed it using RT-qPCR and Y1H assay (**Fig.**
9). WRKY75 was previously described to act as a positive regulator of salt stress tolerance through
activation of *SOS1* expression (Lu et al., 2023). Similar to previous study, we also observed *wrky75* mutants
to be more sensitive to salt stress within our experiments (**Fig. 10A-C**). Nevertheless, when the *wrky75*
mutation was coupled with *sr3g-5*, the roots displayed a phenotype identical to Col-0 (**Fig. 10A**), implying
that the salt sensitivity of the *wrky75* mutant was counteracted by the *SR3G* mutation. Considering
WRKY75's role in positively regulating *SOS1* expression during salt stress, it is plausible that its binding to
the *SR3G* promoter inhibits its expression, thereby enhancing salt tolerance on a larger scale.

One of the main players in Na^+ transport is High-affinity K^+ Transporter (HKT1) (Laurie et al., 2002).
HKT1 is expressed in root stele (Mäser et al., 2002), where it retrieves Na^+ from the transpiration stream
and limits ion accumulation in the shoot, resulting in higher salt stress tolerance at later developmental
stages (Møller et al., 2009). At early developmental stages, *HKT1* overexpression reduces the
development of lateral root primordia (Julkowska et al., 2017), thus creating less Na^+ entry gate into the
main root, resulting in less Na^+ accumulation in shoot. Similarly, *sr3g-5* mutants developed fewer lateral
roots that are consistent with the lower Na^+ levels in the root upon salt stress (**Fig. 6**). The effects of HKT1
and SR3G differ in the magnitude of repression of lateral root development, with HKT1 mutants resulting
in almost complete cessation of lateral root development (Julkowska et al., 2017), whereas repression of
SR3G function has much more subtle phenotype. Whereas the effect of cell-type specific overexpression
of HKT1 on plant performance was negative during the early developmental stage, due to impairing lateral
root development, the *sr3g* mutants showed increased salt tolerance when exposed to salt stress
treatment during their early vegetative growth (**Fig. 7**). These results suggest that reduced lateral root
development under salt stress can have a gradient of responses leading to increased and decreased salt
tolerance, and the effect is highly dependent on the wider physiological mechanisms, and their
contributions to ion exclusion and maintained plant development throughout salt stress exposure.

In conclusion, the study of salt-induced changes in the root:shoot ratio, alongside the
development of new tools to quantify these changes during the seedling establishment stage, represents
an important new perspective into mechanisms underlying salt stress tolerance. Identification of SR3G as
a novel regulator of root suberization, shoot growth, and Na^+ accumulation provides better understanding

of molecular mechanisms underlying plant responses to salt stress, but also offers a new avenue to

explore more nuanced and tissue-specific strategies for enhancing salt stress tolerance.

Acknowledgments

The authors would like to thank BTI and KAUST Greenhouse Teams for their care of the plants. The authors would like to acknowledge support from the NSF-IOS #2023310 (ADLN) and NSF-IOS #2102120 (ADLN). The majority of funding for this work was generously provided from KAUST baseline funding awarded to Mark Tester, and BTI's startup funds awarded to Magdalena Julkowska.

Author contributions

M.M.J. and M.R.I. conceived the project and wrote the manuscript with inputs from all coauthors, C.T. and M.A.T. helped with conceiving the overall idea for the project, infrastructure and funding, R.A.H and R.S. developed the tool to quantify root and shoot size from Arabidopsis grown on agar plates, A.K. assisted with GWAS analysis, M.R.I. conducted all experiments for functional characterization of *SR3G*, H.S., and M.D.A. performed molecular cloning of DR3G, H.S. screened other DUF247s T-DNA and RNAi mutant lines, L.J. performed the analysis of DUF247 domains that informed the cloning of individual SR3G fragments, G.D. and M.W. conducted GC-MS on suberin monomers, M.P. and E.C. performed ICP-AES, A.D.L.N. and L.Y. performed phylogenetic analysis, B.L. and Y.H. conducted Y1H.

Competing interests

The authors declare no competing interests

Correspondence and requests for materials

should be addressed to Dr. Magdalena M. Julkowska.

 Figure legends

Figure 1. Salt stress is causing discoordination of root and shoot growth. Arabidopsis HapMap population
was screened for salt stress-induced changes in root:shoot ratio. The increase in the projected area of
shoot and root (Fig. S2) was used to estimate **(A)** shoot and **(B)** root exponential growth rate. **(C)** The
root:shoot growth rate ratio was calculated per genotype. The histograms represent the number of
accessions across three studied conditions (0, 75 and 125 mM NaCl), whereas the population average is
indicated using the dashed line. Additionally, the distribution of the genotypes within each treatment was
visualized using the error plots (lower panel), where population average and standard error is indicated
using a colored point and a line, respectively. Individual grey points represent individual genotypes.

Figure 2. SR3G is associated with salt stress-induced changes in shoot-per-total seedling area ratio. The
salt induced changes in root-to-shoot and shoot-per-total seedling area ratios were used as an input for
GWAS. Manhattan plots representing the associations of shoot-per-total seedling area recorded 6 days
after transfer to 125 mM NaCl and **(A)** 250 k SNPs and **(B)** 4 M SNPs. **(C)** Significant associations were
found with the SNPs forming a locus on chromosome 3 in and around *SR3G* (AT3G50160), encoding
Domain of Unknown Function 247 (DUF247). The natural variation in the LD region was studied in 162
accessions sequenced by 1001 Genomes Project. The upper panel represents portion of the missing data,
upper middle panel represents deletions relative to Col-0, while lower middle panel represents the
sequence similarity compared to Col-0. The bottom panel represents the open reading frames (ORFs)
within the LD, and the location of associated SNPs is indicated with the dashed lines. Red dashed lines
represent associations above Bonferroni threshold in 250k SNP mapping, while grey lines represent
associations with -log10(p-value) > 5 in the 4M SNP mapping. **(D)** The haplotype analysis performed using
SNPs located within the coding region of the SR3G revealed significant differences in shoot-per-total
seedling area recorded 6 days after transfer to 125 mM NaCl between Haplotype groups 1 (represented
by 46 accessions, including Col-0) and 11 (represented by 11 accessions, including Blh-1). The significant
differences between individual haplotype groups were tested using ANOVA with Tukey HSD to identify
significantly different groups. **(E)** Upon further sequencing of SR3G in accessions from Haplotype 1 and
11, revealed two 200 bp insertions within SR3G exons in 4 tested accessions belonging to haplotype 11
group. These 200 bp insertions resulted in a premature STOP-codon within the DUF247 domain (after Gly-
215).

Figure 3. Phylogeny and positive selection analysis of *SR3G* and its orthologs from other species. The
orthologous genes of *SR3G* (AT3G50160) from other seven species were identified using GoGeBlast (E-
value < 1.00E-10). **(A)** The protein sequences of the 50 homologous genes were aligned by MUSCLE and
then an unrooted phylogenetic tree was reconstructed by a Maximum -likelihood (ML)-derived style by
RAxML (Bootstrap number: 100). **(B)** A positive selection analysis was conducted by branch-site model A
test: Specifically, we hypothesized the differential or constant dN/dS (ω) substitution rate among the
closely distant branches leading to DUF247 (AT3G50160) (See Method & Materials for details). These
branches were marked as $\omega1$ (light brown branch), $\omega2$ (DUF247-specific branch), $\omega3$ (light blue branch),
and ω-rest (light purple branch). **(C)** The gene structure of DUF247 (AT3G50160) was illustrated by wide
arrows (grey arrow: 5 and 3' UTR, yellow arrow: coding-region). The regions used for cloning (Fragment
A, B, and C) were marked by narrow arrows along with "Coil-Coil Domain" and "Super-polar Fragment"
marked on Fragment A and B, respectively. Lastly, the Bayes Empirical Bayes (BEB) was performed to test
the probability of sites along with $\omega > 1$ over the DUF247 (AT3G50160). Three sites were generated with
a posterior probability > 0.90: 124C (P = 0.992), 206A (P = 0.951), and 209A (P = 0.988). These three sites
were marked with an orange asterisk and assumed to have been undergoing positive selection.

Figure 4. Transcript abundances for the two closely related *DUF247s.* Expression of **(A)** *SR3G*
(AT3G50160) **(B)** and its closest homologue *DUF247-150* (AT3G50150) were measured in Col-0 seedlings
grown with and without salt stress. RT-qPCR analyses were conducted using seedlings grown on 1/2x MS
for 4 d and then followed by transferring to the treatment plates with or without 75 mM NaCl for one
week. Mean values are shown ± SE, with number of replicates (n) shown in each graph. *AT4G04120* was
used as a reference gene for normalization. Significance was determined by the Tukey–Kramer HSD test
in JMP.

Figure 5. SR3G full-length protein resides in the plasma membrane while removal of its transmembrane
domain results in protein relocation to nucleus and cytosol. (A-B) SR3G full-length protein or (C-E)
truncated versions fused to mVenus at the N-terminus were agro-infiltrated into the *Nicotiana*
benthamiana leaves for transient expression. Shown are a bright-field image of transfected leaf cells and
mVenus-SR3G-mediated fluorescence as well as GFP-based subcellular marker. **(A)** Localization of full-
length SR3G alone or **(B)** in combination with the known plasma membrane dye, FM4-64. Localization of
truncated versions of SR3G are shown for **(C)** SR3G-Fragment A, **(D)** SR3G-Fragment B, and **(E)** SR3G-
Fragment AB. **(F)** Free GFP was used as a nuclear and cytosolic marker. Scale bar = 25 µm.

Figure 6. *sr3g* mutant displays reduced growth rate for the lateral root length. Root system architecture analysis of Col-0 and *sr3g-5* plants under various concentrations of NaCl are shown here. **(A)** The representative images of 13-day-old Col-0 and *sr3g-5* genotypes that experienced 9 days of salt treatment at indicated concentrations as well as growth rate for lateral root. **(B)** Salt Tolerance Index (STI) for the main root length, average lateral root length, and lateral root number at 75 mM NaCl. The STI was calculated by dividing the growth rate measured under salt stress by the growth rate measured under control condition for each genotype. **(C)** Na^+/K^+ ratio in root and shoot of Col-0 and *sr3g-5* after two weeks on 75 mM salt are shown. Each dot in (A) and (B) represents individual replicate per genotype. Lines in (A) and (B) graphs represent median and average, respectively. In (A) and (B), the asterisks above the graphs indicate significant differences between Col-0 and the *sr3g-5* mutant, as determined by the Student 's t-test: *$P < 0.05$. Statistical analysis in (C) was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not connected by the same letter are significantly different ($P < 0.05$). Root system architecture and ICP-AES analyses of Col-0 and *sr3g-5* mutant are shown in details in Figure S17.

Figure 7. *sr3g* mutant has higher growth rate and larger rosette area while displaying less ion leakage and Na^+ accumulation in shoot under salt stress. The salt stress responses of 2-week-old (referred to as "early salt stress") or 3-week-old (referred to as "late salt stress") soil-grown Col-0 and *sr3g-5* plants that were exposed to a final concentration of 100 mM NaCl were examined here. **(A)** Rosette growth rate and **(B)** area were monitored over a period of two weeks, during which the "early salt stress" and "late salt stress" groups were exposed to two or one week of salt stress respectively. Each line represents the trajectory of individual plant throughout time of experiments, where red and blue lines indicate Col-0 and *sr3g-5* plants respectively. The dashed lines represent the mean values of the genotype per condition, whereas the grey band represents the confidence interval. The significant differences between genotypes, determined by one-way analysis of variance, are illustrated in a plot below each graph, with dashed red line representing a threshold corresponding to p-value of 0.05. Representative images of 4-week-old Col-0 and *sr3g-5* plannts were shown in (A). **(C)** Ion leakage percentage, **(D)** Na^+ and **(E)** K^+ contents in shoot were measured in 4-week-old plants. The asterisks above the graphs indicate significant differences between Col-0 and *sr3g-5* plants, as determined by one-way analysis of variance: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$. Statistically different groups were determined using Tukey–Kramer HSD test. Groups that are not assigned the same letter are significantly different ($P < 0.05$).

Figure 8. *sr3g* mutant root is more suberized than Col-0. (A) Representative images of fluorol yellow (FY) staining of four root sections (# 1 to 4, from root tip to root base as shown by white rectangles across the

seedling) were shown for Col-0 and *sr3g-5* mutant with or without 75 mM NaCl. Seeds were germinated

on the 1/2 MS plates for 4 d and then seedlings were transferred to the treatment plates with or without

75 mM NaCl for 2 more days. **(B)** Quantification of FY signal intensity for each root section. The FY

quantification was done using ImageJ. Scale bar is equal to 500 μm in all. Three to five biological replicates

were used for FY signal quantification, with n=3 for Col control, n=4 for sr3g control, n=5 for Col under salt

stress, and n=5 for sr3g under salt stress. **(C)** Suberin monomers detected using Gas Chromatography–

Mass Spectrometry (GC-MS) in the Col-0 and *sr3g-5* mutant roots with or without 75 mM NaCl included

fatty acid (FA), fatty alcohol (OH-FA), α,ω-dicarboxylic acids (DCA), and ω-hydroxy fatty acid (ωOH-FA).

The significant differences between Col-0 and *sr3g-5* mutant were evaluated using Student 's t-test. Five

biological replicates per genotype per condition used for this experiment.

Figure 9. WRKY75 binds to the promoter region of the *SR3G* but not to its neighboring DUF247. (A) In

silico searches in CisCross (https://plamorph.sysbio.ru/ciscross/CCL_index.html) shows potential

transcription factors with binding sites on the main *SR3G* (At3g50160) promoter region. A black-arrow

points to the *WRKY75* (AT5G13080). **(B)** RT-qPCR showing expression of *SR3G* (AT3G50160) and **(C)**

WRKY75 (AT5G13080) genes, respectively, in Col-0, *sr3g-5* mutant, and two different mutant alleles of the

wrky75, i.e., wrky75-1 and *wrky75-3*. RT-qPCR analyses were conducted using seedlings grown on 1/2x

MS for 4 d and then followed by transferring to the treatment plates with or without 75 mM NaCl for one

week. Mean values are shown ± SE, with three biological replicates used in each condition and genotype.

AT4G04120 (transposable_element_gene) was used as a reference gene for normalization. Significance

was determined by the Tukey–Kramer HSD test in JMP. Levels not connected by same letter are

significantly different. **(D)** Yeast one-hybrid (Y1H) assay showing *WRKY75* (438 bp) binds to the promoter

region of the main *SR3G* (AT3G50160, 953 bp) **(E)** but not to its neighboring promoter, i.e. DUF247-150

(At3g50150, 631 bp). *pB42AD* (AD) and *pLacZ* were used as effector and reporter construct, respectively.

Effector and reporter constructs were co-transformed into yeast strain EGY48. Transformants were

selected and grown on SD/-Trp-Ura medium. The interactions were observed on SD/-Trp/-Ura + X-Gal

medium. Empty vector expressing AD domain and *pLacZ* were used as negative control. The two positive

controls are NIGT1.4-GolS2 and NIGT1.4-GAE1. The oligo sequences used for Y1H and luciferase assay

were provided in **Table S4.**

Figure 10. *wrky75/sr3g* double mutants roots exhibit reduced sensitivity to salt stress and accumulate

lower levels of Na⁺ in their shoots. (A-C) Root system architecture analysis for the Col-0, *sr3g-5, wrky75-*

3, and two double mutants under various concentrations of NaCl are shown here. **(A)** Main root growth

rate, **(B)** Lateral root number increase, and **(C)** Lateral root length increase are shown for the indicated

genotypes at various salt concentrations. **(D)** Na$^+$/K$^+$ ratio in root and shoot of indicated genotypes after

two weeks on treatment plates. Each dot represents individual replicate per genotype. The asterisks

above the graphs (A-C) indicate significant differences between Col-0 and the genotype, as determined

by the Student 's t-test: *$P < 0.05$ and **$P < 0.01$. ns denotes no statistically significant. Statistical analysis

in (D) was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not

connected by the same letter are significantly different ($P < 0.05$).

Supplemental Figure and Table Legends

Figure S1. Evaluation of the tool's precision for estimating seedling size. The tool used to quantify root and shoot size from the agar plates was tested on the Arabidopsis Col-0 seedlings exposed to various concentrations of salt stress (0, 75 and 125 mM NaCl). The fresh weight of root and shoot of individual plants were recorded at the last day of experiment (8 days after transfer to treatment plates, 12 days after germination), and the correlation analysis between individual organ's fresh weight and area was performed in R. The Correlation co-efficient and the p-value for individual correlations were calculated in R using stats package.

Figure S2. Salt stress reduced the increase in root and shoot area over time across HapMap accession. Four-days-old seedlings exposed to 0, 75 and 125 mM NaCl grown on agar plates were scanned every 2nd day and analyzed for root and shoot area using a custom tool. The change in genotype-specific **(A)** shoot and **(B)** root area was calculated in R using stats package. The thick dashed lines represent population-wide average in root and shoot area over the time of the experiment, while individual transparent lines indicate change in root and shoot area for individual accession. The differences between control and individual salt stress treatments were evaluated using ANOVA. The *, **, *** and **** represent p-values below 0.05, 0.01, 0.001 and 0.0001 respectively.

Figure S3. Salt stress induced changes in relative growth of shoot, root and root:shoot ratio. The stress tolerance index was calculated for shoot, root and root:shoot ratio by dividing genotype specific value recorded at salt stress over the genotypic mean of the trait recorded under non-stress conditions. Individual points represent individual genotypes. The population mean for 75 and 125 mM NaCl treated plants is represented by the colored point, whereas the population-wide standard error by whiskers.

Figure S4. The correlation between all measured traits and calculated stress tolerance indices (STI). The correlations between all measured and calculated traits was performed using the Pearson's correlation coefficient. All non-significant correlations were replaced by blank circles. The size and color of individual circles indicates the strength and significance of individual correlation between two traits. Blue and red shades indicate positive and negative correlations respectively.

Figure S5. The correlation between total seedling area and salt-induced changes in the root-to-shoot ratio. The correlation between the total seedling area quantified using the custom tool at the last day of experiment (8 days after transfer and 12 days after germination), and the ratio of root-and-shoot growth

rate was examined across Arabidopsis HapMap population and various concentrations of salt stress (0, 75

and 125 mM NaCl). The correlation analysis was performed in R, and the Correlation co-efficient (r) and

the p-values for individual correlations were calculated in R using stats package.

Figure S6. Region around NTL8 is associated with salt stress-induced changes in root-per-shoot growth.

The salt induced changes in root-to-shoot and shoot-per-total seedling area ratios were used as an input

for GWAS. **(A)** Manhattan plot represents the associations of root-per-shoot growth factors recorded at

75 mM NaCl with 250 k SNPs while **(B)** represents the QQ plot for this association study. **(C)** Significant

associations were found with the SNPs forming a locus on chromosome 2 in and around AT2G27300,

encoding a NAC-domain containing transcription factor (NTL8). The natural variation in the LD region was

studied in 162 accessions sequenced by 1001 Genomes Project. The upper panel represents portion of

the missing data, upper middle panel represents deletions relative to Col-0, while lower middle panel

represents the sequence similarity compared to Col-0. The bottom panel represents the open reading

frames (ORFs) within the LD, and the location of associated SNPs is indicated with the dashed lines. Red

dashed lines represent associations above Bonferroni threshold in 250k SNP mapping, while grey lines

represent associations with -log10(p-value) > 5 in the 4M SNP mapping.

Figure S7. Region around unknown gene is associated with salt stress-induced changes in relative

changes in shoot-per-total seedling area. The salt induced changes in root-to-shoot and shoot-per-total

seedling area ratios were used as an input for GWAS. **(A)** Manhattan plot represents the associations of

relative change (125 mM / 0 mM) in shoot-per-total seedling area at 4 days after transfer to 125 mM NaCl,

while **(B)** represents the QQ plot for this association study. **(C)** Significant associations were found with

the SNPs forming a locus on chromosome 3 in and around AT3G22430, encoding an unknown gene

encoding a domain of unknown function. The natural variation in the LD region was studied in 162

accessions sequenced by 1001 Genomes Project. The upper panel represents portion of the missing data,

upper middle panel represents deletions relative to Col-0, while lower middle panel represents the

sequence similarity compared to Col-0. The bottom panel represents the open reading frames (ORFs)

within the LD, and the location of associated SNPs is indicated with the dashed lines. Red dashed lines

represent associations above Bonferroni threshold in 250k SNP mapping, while grey lines represent

associations with -log10(p-value) > 5 in the 4M SNP mapping.

Figure S8. The QQ-plots for GWAS models with shoot-per-total seedling area. The QQ-plots for

association between shoot-per-total seedling area recorded 6 days after transfer to 125 mM NaCl and **(A)**

250 k SNPs and **(B)** 4 M SNPs. The red line indicates the expected vs. observed ratio of association, while

black points indicate the observed associations.

Figure S9. Haplotype analysis of AT3G50160 (SR3G). The haplotype analysis performed using SNPs

located within the coding region of the SR3G (AT3G50160) revealed significant differences in **(A)** Relative

shoot growth (as fraction of shoot growth at 125 mM NaCl and 0 mM NaCl) and **(B)** Shoot growth at 125

mM NaCl. The significant differences between individual haplotype groups (each haplotype represented

by at least 3 accessions), were tested using ANOVA with Tukey HSD to identify significantly different

groups. Significant differences were identified between Haplotype groups 1 (represented by 46

accessions, including Col-0) and 11 (represented by 11 accessions, including Blh-1).

Figure S10. Nucleotide sequence alignment of SR3G CDS between Col-0 and Blh-1 alleles. The nucleotide

sequences from SR3G CDS were obtained from Col-0 and Blh-1 cDNA, and compared to Col-0 cDNA

sequence on TAIR. The cDNA sequences were subsequently aligned in JalView using ProbconsWS

alignment. The intensity of purple hue indicates the nucleotide identity, while dash marks indicate

insertion in Blh-1 compared to Col-0 allele.

Figure S11. Amino acid sequence alignment of SR3G CDS between Col-0 and Blh-1 alleles. The amino

acid sequences from SR3G CDS were obtained by translating Col-0 and Blh-1 cDNA into amino acid

sequence in JalView using Standard settings. The amino acid sequences were subsequently aligned in

JalView using ProbconsWS alignment. The sequence length markers for are indicated above the aligned

sequence. The intensity of purple hue is indicating the nucleotide identity, while dash marks indicate

insertion in one of the alleles. * indicate STOP codons.

Figure S12. Sequence alignment between Col-0 and Blh-1 alleles for SR3G promotor. The nucleotide

sequences from SR3G promotor were obtained from Col-0 and Blh-1 genomic DNA, and compared to Col-

0 sequence on TAIR. The gDNA sequences were subsequently aligned in JalView using ProbconsWS

alignment. The intensity of purple hue is indicating the nucleotide identity, while dash marks indicate

insertions.

Figure S13. Gene expression profile of *SR3G* across various developmental stages and salt stress. The

absolute gene expression of *SR3G* is depicted across **(A)** various developmental stages, **(B)** root cell types,

and **(C)** in response to salt stress. The images were acquired through the eFP Browser using data sources

specified for the developmental map, abiotic stress, and tissue specific.

Figure S14. Gene expression profile of other *DUF247s* under salt stress. Absolute gene expression of

other *DUF247s* is illustrated in response to salt-induced changes. The images were acquired through the

eFP Browser using data sources specified for the abiotic stress. The *SR3G* is shown in a green box for

comparison.

Figure S15. SR3G predicted protein domains. The SR3G protein sequence (Col-0 allele) was used to

identify predicted protein domains. The identified domains include: nuclear localization signal (NLS): K15-

F19; coiled-coil domain: M153-K173; potential Ca2+ binding pocket: G210-F372; and transmembrane

domain P469-T503. Informed by these predicted protein domains, we designed SR3G Fragment A, B and

C for subsequent localization studies and functional characterization. The pink mark at Y216 indicates a

position of a premature STOP codon in Blh-1 allele.

Figure S16. The expression of *DUF247-150* remains unaltered in the *sr3g* mutant. (A) Schematic diagram

of main *SR3G* gene model and the location of T-DNA insertions. The location of T-DNA insertions is shown

for *sr3g-4*, and *-5* mutants using triangles. RT-qPCR showing expression of *SR3G* gene (AT3G50160) **(B)**

and its neighboring gene, *DUF247-150* (At3g50150) **(C)** in Col-0 and two *sr3g* mutants. RT-qPCR analyses

were conducted using seedlings grown on 1/2x MS for 4 d and then followed by transferring to the

treatment plates with or without 75 mM NaCl for one week. Mean values are shown ± SE, with the number

of replicates (n) shown in each graph. *AT4G04120* (transposable_element_gene) was used as reference

gene for normalization. Significance was determined by the Tukey–Kramer HSD test in JMP. Levels not

connected by same letter are significantly different.

Figure S17. *sr3g-5* mutant displays no difference in main root length and lateral root numbers compared

to Col-0. Root system architecture analysis of Col-0 and *sr3g-5* plants under various concentrations of

NaCl are shown here for **(A)** main root length and **(B)** increase in lateral root number. 13-day-old Col-0

and *sr3g-5* genotypes experienced 9 days of salt treatment at indicated concentrations were used here.

Each dot in (A) and (B) represents individual replicate per genotype. Statistical analysis was performed

between Col-0 and the *sr3g-5* mutant using the Student 's t-test: *$P < 0.05$. ns stands for not significant.

(C) Na$^+$ and **(D)** K$^+$ contents in root and shoot of Col-0 and *sr3g-5* after two weeks on 75 mM salt are shown.

Statistical analysis in (C-D) was done by comparison of the means for all pairs using Tukey–Kramer HSD

test. Levels not connected by the same letter are significantly different ($P < 0.05$).

Figure S18. *sr3g-4* mutant has no alteration in root system architectures compared to Col-0. Root system

architecture analysis of Col-0 and *sr3g-4* plants under various concentrations of NaCl are shown here for

(A) main root length, **(B)** lateral root length, and **(C)** lateral root number. Salt Tolerance Index (STI) for the

main root length, average lateral root length, and lateral root number are shown at **(D)** 75 and **(E)** 125

mM NaCl, respectively. The STI was calculated by dividing the growth rate measured under salt stress by

the growth rate measured under control condition for each genotype. Statistical analysis was performed

between Col-0 and the *sr3g-4* mutant using the Student 's t-test: *P < 0.05.

Figure S19. Root system architecture analysis of neighboring DUF247 mutants under the salt. (A) Main

root length, **(B)** lateral root number, and **(C)** lateral root length are shown for neighboring *DUF247s* of the

SR3G, including At3g50120 (shown as 120, with one T-DNA representative, 120-2(SAIL_382_A09)),

At3g50140 (shown as 140, with three T-DNA representatives, 140-1 (SALK_044685), 140-2

(SALK_005466), and 140-3 (SALK_122700)), At3g50150 (shown as 150, with two T-DNA representatives,

150-1 (SALK_003824) and 150-3 (SALK_071080)), and At3g50170 (shown as 170, with two T-DNA

representatives, 170-1 (SALK_009186) and 170-2 (SALK_112602)) compared to Col-0 under various

concentrations of NaCl as indicated. The corresponding Salt Tolerance Index (STI) was calculated for **(D)**

main root growth, **(E)** increase in lateral root number, and **(F)** lateral root growth rate. Statistical analysis

was performed between Col-0 and the other mutants using the Student 's t-test: *P < 0.05. The root

system architecture analysis was performed similar to what described for Figure 6.

Figure S20. Root system architecture analysis of neighboring DUF247 mutants under the salt, continues.

(A) Main root length, **(B)** lateral root number, and **(C)** lateral root length are shown for neighboring of the

SR3G, including At3g50170 (shown as 170, with three T-DNA representatives, 170-3F (SALK_008710), 170-

4F (SALK_145999), 170-5A (SALK_072937)), At3g50190 (shown as 190, with one T-DNA representative,

190-2B (SALK_137791)), and At3g50200 (shown as 200, with three T-DNA representatives, 200-1A

(SALK_113759), 200-2B (SALK_093701), and 200-3B (SALK_129634)) compared to Col-0 under various

concentrations of NaCl as indicated. The corresponding Salt Tolerance Index (STI) was calculated for **(D)**

main root growth, **(E)** increase in lateral root number, and **(F)** lateral root growth rate. Statistical analysis

was performed between Col-0 and the other mutants using the Student 's t-test: *P < 0.05. The root

system architecture analysis was performed similar to what described for Figure 6.

Figure S21. Root system architecture analysis of RNAi lines targeting neighboring DUF247s. (A) Main

root length, **(B)** lateral root number, and **(C)** lateral root length; **(D)** lateral root length tolerance index, **(E)**

lateral root number, and **(F)** lateral root number tolerance index are shown for RNAi lines targeting

AT3G50140 (CSHL3-F1a), targeting AT3G50130 and AT3G50140 (CSHL6-C2d), targeting AT3G50150 and

AT3G50170 (CSHL11-F1b), targeting AT3G50140 and AT3G50170 (CSHL13-F2a), targetingAT3G50150 and

AT3G50190 (CSHL18-K4c), targeting AT3G50130, AT3G50140, AT3G50150, AT3G50160, AT3G50170,

AT3G50180, and AT3G50190 (CSHL19-E2c), targeting AT3G50130 and AT3G50150 (CSHL21-E2a), and

targeting AT3G50150 and AT3G50160 (CSHL22-F1c). Statistical analysis was performed between Col-0 and

the RNAi lines using a Student's t-test: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Figure S22. Root system architecture analysis of RNAi lines targeting neighboring DUF247s continued.

(A) Main root length, **(B)** lateral root number, and **(C)** lateral root length; **(D)** lateral root length tolerance

index, **(E)** lateral root number, and **(F)** lateral root number tolerance index are shown for RNAi lines

targeting AT3G50160 (CSHL4-B2a and CSHL4G1d), targeting AT3G50140 and AT3G50150 (CSHL7-A3a),

targeting AT3G50140 and AT3G50190 (CSHL8-G2b), targeting AT3G50180 (CSHL10-I1c), targeting

AT3G50130, AT3G50140, AT3G50170, AT3G50180, and AT3G50190 (CSHL12-A1a), targeting AT3G50140,

AT3G50170, AT3G50180, and AT3G50190 (CSHL15-E1b), targeting AT3G50130, AT3G50160, and

AT3G50170 (CSHL-A2c and CSHL16-B1a), targeting AT3G50130 and AT3G50160 (CSHL20-G3b), and

targeting AT3G50130 and AT3G50150 (CSHL21-L2c and CSHL21-M1d). Statistical analysis was performed

between Col-0 and the RNAi lines using a Student's t-test: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P <$

0.0001.

Figure S23. Root system architecture analysis of RNAi lines targeting neighboring DUF247s continued.

(A) Main root length, **(B)** lateral root number, and **(C)** lateral root length; **(D)** lateral root length tolerance

index, **(E)** lateral root number, and **(F)** lateral root number tolerance index are shown for RNAi lines

targeting AT3G50140 (CSHL3-F3a), targeting AT3G50160 (CSHL4-D1c3, CSHL4-D4a, and CSHL4-F1b),

targeting AT3G50130 and AT3G50170 (CSHL5-D3b and CSHL5-E2b), targeting AT3G50130 and AT3G50140

(CSHL6-B1c), targeting AT3G50140 and AT3G50150 (CSHL7-C1b2), targeting AT3G50140 and AT3G50190

(CSHL8-D3b and CSHL8-H1d), targeting AT3G50180 (CSHL10-I1b), targeting AT3G50150 and AT3G50170

(CSHL11-E5b), targeting AT3G50130, AT3G50140, AT3G50170, AT3G50180, and AT3G50190 (CSHL12-B1a

and CSHL12-D1c), targeting AT3G50140 and AT3G50170 (CSHL13-F1b and CSHL13-F4d), targeting

AT3G50140, AT3G50170, AT3G50180, and AT3G50190 (CSHL15-G1a1 and CSHL15-H2b), targeting

AT3G50150 and AT3G50190 (CSHL18-E4c and CSHL18-I2d), targeting AT3G50130, AT3G50140,

AT3G50150, AT3G50160, AT3G50170, AT3G50180, and AT3G50190 (CSHL19-F1a, CSHL19-I3c, and

CSHL19-L1c), and targeting AT3G50130 and AT3G50160 (CSHL20-F2b). Statistical analysis was performed

between Col-0 and the RNAi lines using a Student's t-test: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P <$

0.0001.

Figure S24. Additional phenotypes of soil grown sr3g mutant and Col-0 wild-type plants. Salt Tolerance Index (STI) was calculated by divididng the value of each trait for each genotype by its corresponding mean value for that trait and timepoint. The STI was calculated for **(A)** Rosette growth rate and **(B)** area. Each line represents the relative trajectory of individual plant throughout time of experiments, where red and blue lines indicate Col-0 and *sr3g-5* plants respectively. The dashed lines represent the mean values of the genotype per condition, whereas the grey band represents the confidence interval. The significant differences between genotypes, determined by one-way analysis of variance, are illustrated in a plot below each graph, with dashed red line representing a threshold corresponding to p-value of 0.05. **(C)** The percentage of plant water content was calculated using the formula: (Fresh Weight–Dry Weight)/Fresh Weight×100 for the 2-week-old (referred to as "early salt stress") or 3-week-old (referred to as "late salt stress") soil-grown Col-0 and sr3g-5 plants that were exposed to a final concentration of 100 mM NaCl. "early salt stress" group experienced two weeks of salt stress while the "late salt stress" group experienced one week. Statistical analysis was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not connected by the same letter are significantly different (P < 0.05).

Figure S25. *wrky75/sr3g* double mutants exhibit higher shoot K$^+$ contents compared to Col-0 and each individual single mutant in the presence of 125 mM salt. (A) Salt Tolerance Index (STI) for the main root length (MR.STI), average lateral root length (aLRL.STI), and lateral root number (LRno.STI) at 75 mM and **(B)** 125 mM NaCl are shown here. The STI was calculated by dividing the growth rate measured under salt stress by the growth rate measured under control condition for each genotype. The STI reported here were correspond to root system architecture analysis in Figure 10. Each dot in (A) and (B) represents individual replicate per genotype. In (A) and (B), significant differences between Col-0 and the *other genotypes was* determined by the Student 's t-test: *P < 0.05. ns stands for not significant. **(C)** Na$^+$ and **(D)** K$^+$ contents in root and shoot of Col-0, *sr3g-5, wrky75-3,* and two double mutants under various concentrations of NaCl are shown here. Statistical analysis in (A-B) was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not connected by the same letter are significantly different (P < 0.05).

Figure S26. *wrky75/sr3g* double mutants exhibit no noticeable differences in rosette area compared to their individual single mutants when grown in saline soil. The salt stress responses of 2-week-old soil-grown plants that were exposed to a final concentration of 100 mM NaCl were examined here. Rosette area **(A)** and growth rate for each individual day were shown **(B-M)**. Growth rate is shown for 1 **(B)**, 2 **(C)**, and 3 **(D)** days before the stress, respectively. Growth rate is shown for 1 **(E)**, 2 **(F)**, 3 **(G)**, 4 **(H)**, 5 **(I)**, 6 **(J)**, 7 **(K)**, 8 **(L)**, and 9 **(M)** days of stress application, respectively. Na+/K+ **(N)** ratio in shoot were measured in 4-week-old plants. Each line in (A) represents the trajectory per genotype through time. The effect of the genotype within individual treatment in (B-M) was tested using ANOVA, and the significantly different

genotypes were additionally determined using Tukey HSD test (pvalue < 0.05). Statistical analysis in (N)

was done by comparison of the means for all pairs using Tukey–Kramer HSD test. Levels not connected by

the same letter are significantly different ($P < 0.05$).

Table S1. The individual traits recorded and calculated from the HapMap accessions exposed to the

three different levels of salt stress. The root and shoot projected area were quantified using a custom

developed tool. The change in the root and shoot area over time was used to calculate Growth Factor (GF)

by fitting an exponential function. The broad sense heritability (H2) was calculated for each trait over

individual days and conditions using MVApp.

Table S2. Genotypic mean data used as an input for GWAS. The genotypic mean data was calculated

from at least 4 biological replicates per genotype per condition. The root and shoot area were extracted

using a custom developed tool. The Growth Factors (GF) for root and shoot were calculated using

exponential growth function. Total seedling slze (Tot) was calculated by adding root and shoot area for

individual day. Shoot per Total (SpT) ratio was calculated for individual days by dividing genotype specific

shoot area by the total seedling area for that day and condition. Root per Shoot ratio (RpS) was calculated

by dividing genotype specific root area by the shoot area for that day and condition.

Table S3. Significant associations identified with GWAS of all used traits. The location of individual SNPs

is listed according to its location on Chromosome (Chr) and position (Pos). The individual SNP Minor Allele

Count (MAC) and Minor Allele Frequency (MAF) is calculated based on specific SNP set used (4 Million or

250 Thousand – listed in "Mapping" column). The traits associated with individual SNPs are abbreviated

as RpS for Root-per-Shoot, SpT for Shoot-per-Total seedling area, Tot for Total seedling area, GR for

growth rate and SHIIT for Shoot Ion Independent Tolerance index (Salt / Control).

Table S4. Most important GWAS associations. The most important associations identified through GWAS

(Table S3) were inspected further for their functions using either T-DNA or RNAi mutant lines. List of

oligoes used for genotyping, RT-qPCR, cloning, and Y1H assay are provided.

Table S5. The gene ID, species, and gene names used for identification of SR3G orthologs and

paralogs.

Table S6. Background and foreground values of omega.

Table S7. Links to cloning construct maps generated in this study.

1227

References

1001 Genomes Consortium. 1001 Genomes Consortium. 2016. 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell* **166**:481–491. doi:10.1016/j.cell.2016.05.063

Abrol IP, Yadav JSP, Massoud FI. 1988. Salt-affected Soils and Their Management. Food & Agriculture Org.

Ariel FD, Diet A, Crespi M, Chan RL. 2010. The LOB-like transcription factor MtLBD1 controls Medicago truncatula root architecture under salt stress. *Plant Signal Behav* **5**:1666–1668. doi:10.4161/psb.5.12.14020

Asch F, Dingkuhn M, Sow A, Audebert A. 2005. Drought-induced changes in rooting patterns and assimilate partitioning between root and shoot in upland rice. *Field Crops Res* **93**:223–236. doi:10.1016/j.fcr.2004.10.002

Awlia M, Alshareef N, Saber N, Korte A, Oakey H, Panzarová K, Trtílek M, Negrão S, Tester M, Julkowska MM. 2021. Genetic mapping of the early responses to salt stress in Arabidopsis thaliana. *Plant J* **107**:544–563. doi:10.1111/tpj.15310

Awlia M, Nigro A, Fajkus J, Schmoeckel SM, Negrão S, Santelia D, Trtílek M, Tester M, Julkowska MM, Panzarová K. 2016. High-Throughput Non-destructive Phenotyping of Traits that Contribute to Salinity Tolerance in Arabidopsis thaliana. *Front Plant Sci* **7**:1414. doi:10.3389/fpls.2016.01414

Bacher H, Sharaby Y, Walia H, Peleg Z. 2022. Modifying root-to-shoot ratio improves root water influxes in wheat under drought stress. *J Exp Bot* **73**:1643–1654. doi:10.1093/jxb/erab500

Barberon M, Vermeer JEM, De Bellis D, Wang P, Naseer S, Andersen TG, Humbel BM, Nawrath C, Takano J, Salt DE, Geldner N. 2016. Adaptation of Root Function by Nutrient-Induced Plasticity of Endodermal Differentiation. *Cell* **164**:447–459. doi:10.1016/j.cell.2015.12.021

Baxter I, Hosmani PS, Rus A, Lahner B, Borevitz JO, Muthukumar B, Mickelbart MV, Schreiber L, Franke RB, Salt DE. 2009. Root suberin forms an extracellular barrier that affects water relations and mineral nutrition in Arabidopsis. *PLoS Genet* **5**:e1000492. doi:10.1371/journal.pgen.1000492

Beilstein MA, Renfrew KB, Song X, Shakirov EV, Zanis MJ, Shippen DE. 2015. Evolution of the Telomere-Associated Protein POT1a in Arabidopsis thaliana Is Characterized by Positive Selection to Reinforce Protein-Protein Interaction. *Mol Biol Evol* **32**:1329–1341. doi:10.1093/molbev/msv025

Birnbaum K, Shasha DE, Wang JY, Jung JW, Lambert GM, Galbraith DW, Benfey PN. 2003. A gene expression map of the Arabidopsis root. *Science* **302**:1956–1960. doi:10.1126/science.1090022

Brady SM, Zhang L, Megraw M, Martinez NJ, Jiang E, Yi CS, Liu W, Zeng A, Taylor-Teeples M, Kim D, Ahnert S, Ohler U, Ware D, Walhout AJM, Benfey PN. 2011. A stele-enriched gene regulatory network in the Arabidopsis root. *Mol Syst Biol* **7**:459. doi:10.1038/msb.2010.114

Chen Q, Hu T, Li X, Song C-P, Zhu J-K, Chen L, Zhao Y. 2022. Phosphorylation of SWEET sucrose transporters regulates plant root:shoot ratio under drought. *Nat Plants* **8**:68–77. doi:10.1038/s41477-021-01040-7

Clough SJ, Bent AF. 1998. Floral dip: a simplified method forAgrobacterium-mediated transformation ofArabidopsis thaliana. *Plant J* **16**:735–743. doi:10.1046/j.1365-313x.1998.00343.x

Crain J, Larson S, Dorn K, Hagedorn T, DeHaan L, Poland J. 2020. Sequenced-based paternity analysis to improve breeding and identify self-incompatibility loci in intermediate wheatgrass (Thinopyrum intermedium). *Theor Appl Genet* **133**:3217–3233. doi:10.1007/s00122-020-03666-1

Delude C, Vishwanath SJ, Rowland O, Domergue F. 2017. Root Aliphatic Suberin Analysis Using Non-extraction or Solvent-extraction Methods. *Bio-protocol* **7**. doi:10.21769/BioProtoc.2331

Deolu-Ajayi AO, Meyer AJ, Haring MA, Julkowska MM, Testerink C. 2019. Genetic Loci Associated with Early Salt Stress Responses of Roots. *iScience* **21**:458–473. doi:10.1016/j.isci.2019.10.043

Duan L, Dietrich D, Ng CH, Chan PMY, Bhalerao R, Bennett MJ, Dinneny JR. 2013. Endodermal ABA

signaling promotes lateral root quiescence during salt stress in Arabidopsis seedlings. *Plant Cell*
25:324–341. doi:10.1105/tpc.112.107227
Fischer, G., Nachtergaele, O. F, Velthuizen V, T. H, Chiozza, F., Franceschini, G., Henry, M., Muchoney, D.,
Tramberend, S. 2021. Global Agro-Ecological Zones v4 – Model documentation. Food & Agriculture
Org.
Foster KJ, Miklavcic SJ. 2017. A Comprehensive Biophysical Model of Ion and Water Transport in Plant
Roots. I. Clarifying the Roles of Endodermal Barriers in the Salt Stress Response. *Front Plant Sci*
8:1326. doi:10.3389/fpls.2017.01326
Fu Y, Yang Y, Chen S, Ning N, Hu H. 2019. Arabidopsis IAR4 Modulates Primary Root Growth Under Salt
Stress Through ROS-Mediated Modulation of Auxin Distribution. *Front Plant Sci* **10**:522.
doi:10.3389/fpls.2019.00522
Galvan-Ampudia CS, Julkowska MM, Darwish E, Gandullo J, Korver RA, Brunoud G, Haring MA, Munnik T,
Vernoux T, Testerink C. 2013. Halotropism is a response of plant roots to avoid a saline
environment. *Curr Biol* **23**:2044–2050. doi:10.1016/j.cub.2013.08.042
Gehan MA, Fahlgren N, Abbasi A, Berry JC, Callen ST, Chavez L, Doust AN, Feldman MJ, Gilbert KB, Hodge
JG, Hoyer JS, Lin A, Liu S, Lizárraga C, Lorence A, Miller M, Platon E, Tessman M, Sax T. 2017.
PlantCV v2: Image analysis software for high-throughput plant phenotyping. *PeerJ* **5**:e4088.
doi:10.7717/peerj.4088
Geng Y, Wu R, Wee CW, Xie F, Wei X, Chan PMY, Tham C, Duan L, Dinneny JR. 2013. A spatio-temporal
understanding of growth regulation during the salt stress response in Arabidopsis. *Plant Cell*
25:2132–2154. doi:10.1105/tpc.113.112896
Guo W, Chen L, Herrera-Estrella L, Cao D, Tran L-SP. 2020. Altering Plant Architecture to Improve
Performance and Resistance. *Trends Plant Sci* **25**:1154–1170. doi:10.1016/j.tplants.2020.05.009
Hassani A, Azapagic A, Shokri N. 2021. Global predictions of primary soil salinization under changing
climate in the 21st century. *Nat Commun* **12**:6663. doi:10.1038/s41467-021-26907-3
Höfer R, Briesen I, Beck M, Pinot F, Schreiber L, Franke R. 2008. The Arabidopsis cytochrome P450
CYP86A1 encodes a fatty acid omega-hydroxylase involved in suberin monomer biosynthesis. *J Exp*
Bot **59**:2347–2360. doi:10.1093/jxb/ern101
Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, Muliyati NW, Platt A, Sperone FG,
Vilhjálmsson BJ, Nordborg M, Borevitz JO, Bergelson J. 2012. Genome-wide patterns of genetic
variation in worldwide Arabidopsis thaliana accessions from the RegMap panel. *Nat Genet* **44**:212–
216. doi:10.1038/ng.1042
Hu Y, Wu Q, Sprague SA, Park J, Oh M, Rajashekar CB, Koiwa H, Nakata PA, Cheng N, Hirschi KD, White
FF, Park S. 2015. Tomato expressing Arabidopsis glutaredoxin gene AtGRXS17 confers tolerance to
chilling stress via modulating cold responsive components. *Hortic Res* **2**:15051.
doi:10.1038/hortres.2015.51
Hu Y, Zeng L, Lv X, Guo J, Li X, Zhang X, Wang D, Wang J, Bi J, Julkowska MM, Li B. 2023. NIGT1.4
maintains primary root elongation in response to salt stress through induction of ERF1 in
Arabidopsis. *Plant J* **116**:173–186. doi:10.1111/tpj.16369
Julkowska MM, Hoefsloot HCJ, Mol S, Feron R, de Boer G-J, Haring MA, Testerink C. 2014. Capturing
Arabidopsis root architecture dynamics with ROOT-FIT reveals diversity in responses to salinity.
Plant Physiol **166**:1387–1402. doi:10.1104/pp.114.248963
Julkowska MM, Klei K, Fokkens L, Haring MA, Schranz ME, Testerink C. 2016. Natural variation in rosette
size under salt stress conditions corresponds to developmental differences between Arabidopsis
accessions and allelic variation in the LRR-KISS gene. *J Exp Bot* **67**:2127–2138.
doi:10.1093/jxb/erw015
Julkowska MM, Koevoets IT, Mol S, Hoefsloot H, Feron R, Tester MA, Keurentjes JJB, Korte A, Haring MA,
de Boer G-J, Testerink C. 2017. Genetic Components of Root Architecture Remodeling in Response

to Salt Stress. *Plant Cell* **29**:3198–3213. doi:10.1105/tpc.16.00680

Julkowska MM, Saade S, Agarwal G, Gao G, Pailles Y, Awlia M, Tester M. 2018. MVAPP—multivariate
analysis application for streamlined data analysis and curation. *Plant Physiol* **23**:44.

Julkowska MM, Testerink C. 2015. Tuning plant signaling and growth to survive salt. *Trends Plant Sci*
20:586–594. doi:10.1016/j.tplants.2015.06.008

Karcher DE, Richardson MD, Hignight K, Rush D. 2008. Drought tolerance of tall fescue populations
selected for high root/shoot ratios and summer survival. *Crop Sci* **48**:771–777.
doi:10.2135/cropsci2007.05.0272

Kassambara A. n.d. ggpubr:'ggplot2'Based Publication Ready Plots. R package version 0.4. 0. *]*
https://cran-r-project org/web/packages/ggpubr/indes ….

Kondou Y, Noguchi K, Kutsuna S, Kawashima M, Yoneda A, Ishibashi M, Muto S, Ichikawa T, Nakazawa
M, Matsui M, Manabe K. 2013. Overexpression of DWARF AND LESION FORMATION 1 (DLE1)
causes altered activation of plant defense system in Arabidopsis thaliana. *Plant Biotechnol* **30**:385–
392. doi:10.5511/plantbiotechnology.13.0605a

Korver RA, van den Berg T, Meyer AJ, Galvan-Ampudia CS, Ten Tusscher KHWJ, Testerink C. 2020.
Halotropism requires phospholipase Dζ1-mediated modulation of cellular polarity of auxin
transport carriers. *Plant Cell Environ* **43**:143–158. doi:10.1111/pce.13646

Kreszies T, Shellakkutti N, Osthoff A, Yu P, Baldauf JA, Zeisler-Diehl VV, Ranathunge K, Hochholdinger F,
Schreiber L. 2019. Osmotic stress enhances suberization of apoplastic barriers in barley seminal
roots: analysis of chemical, transcriptomic and physiological responses. *New Phytol* **221**:180–194.
doi:10.1111/nph.15351

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J. 2013. GreenGate---a novel,
versatile, and efficient cloning system for plant transgenesis. *PLoS One* **8**:e83043.
doi:10.1371/journal.pone.0083043

Laurie S, Feeney KA, Maathuis FJM, Heard PJ, Brown SJ, Leigh RA. 2002. A role for HKT1 in sodium
uptake by wheat roots. *Plant J* **32**:139–149. doi:10.1046/j.1365-313x.2002.01410.x

Lian X, Zhang S, Huang G, Huang L, Zhang J, Hu F. 2021. Confirmation of a Gametophytic Self-
Incompatibility in Oryza longistaminata. *Front Plant Sci* **12**:576340. doi:10.3389/fpls.2021.576340

Li B, Kamiya T, Kalmbach L, Yamagami M, Yamaguchi K, Shigenobu S, Sawa S, Danku JMC, Salt DE,
Geldner N, Fujiwara T. 2017. Role of LOTR1 in Nutrient Transport through Organization of Spatial
Distribution of Root Endodermal Barriers. *Curr Biol* **27**:758–765. doi:10.1016/j.cub.2017.01.030

Li Y, Beisson F, Koo AJK, Molina I, Pollard M, Ohlrogge J. 2007. Identification of acyltransferases required
for cutin biosynthesis and production of cutin with suberin-like monomers. *Proc Natl Acad Sci U S A*
104:18339–18344. doi:10.1073/pnas.0706984104

Lobet G, Pagès L, Draye X. 2011. A novel image-analysis toolbox enabling quantitative analysis of root
system architecture. *Plant Physiol* **157**:29–39. doi:10.1104/pp.111.179895

Lu K-K, Song R-F, Guo J-X, Zhang Y, Zuo J-X, Chen H-H, Liao C-Y, Hu X-Y, Ren F, Lu Y-T, Liu W-C. 2023.
CycC1;1-WRKY75 complex-mediated transcriptional regulation of SOS1 controls salt stress
tolerance in Arabidopsis. *Plant Cell* **35**:2570–2591. doi:10.1093/plcell/koad105

Mäser P, Eckelman B, Vaidyanathan R, Horie T, Fairbairn DJ, Kubo M, Yamagami M, Yamaguchi K,
Nishimura M, Uozumi N, Robertson W, Sussman MR, Schroeder JI. 2002. Altered shoot/root Na+
distribution and bifurcating salt sensitivity in Arabidopsis by genetic disruption of the Na+
transporter AtHKT1. *FEBS Lett* **531**:157–161. doi:10.1016/s0014-5793(02)03488-9

Møller IS, Gilliham M, Jha D, Mayo GM, Roy SJ, Coates JC, Haseloff J, Tester M. 2009. Shoot Na+
exclusion and increased salinity tolerance engineered by cell type-specific alteration of Na+
transport in Arabidopsis. *Plant Cell* **21**:2163–2178. doi:10.1105/tpc.108.064568

Munns R, Tester M. 2008. Mechanisms of Salinity Tolerance. *Annual Review of Plant Biology*.
doi:10.1146/annurev.arplant.59.032607.092911

Nelson ADL, Forsythe ES, Gan X, Tsiantis M, Beilstein MA. 2014. Extending the model of Arabidopsis telomere length and composition across Brassicaceae. *Chromosome Res* **22**:153–166. doi:10.1007/s10577-014-9423-y

Nelson ADL, Haug-Baltzell AK, Davey S, Gregory BD, Lyons E. 2018. EPIC-CoGe: managing and analyzing genomic data. *Bioinformatics* **34**:2651–2653. doi:10.1093/bioinformatics/bty106

Norris SR, Meyer SE, Callis J. 1993. The intron of Arabidopsis thaliana polyubiquitin genes is conserved in location and is a quantitative determinant of chimeric gene expression. *Plant Mol Biol* **21**:895–906. doi:10.1007/BF00027120

Ramachandran P, Augstein F, Nguyen V, Carlsbecker A. 2020. Coping With Water Limitation: Hormones That Modify Plant Root Xylem Development. *Front Plant Sci* **11**:570. doi:10.3389/fpls.2020.00570

Ranathunge K, Schreiber L. 2011. Water and solute permeabilities of Arabidopsis roots in relation to the amount and composition of aliphatic suberin. *J Exp Bot* **62**:1961–1974. doi:10.1093/jxb/erq389

Richards LA. 1969. Diagnosis and Improvement of Saline and Alkali Soils. United States Department of Agriculture.

Şekerci K, Higashitani N, Ozgur R, Uzilday B, Higashitani A, Turkan I. 2023. A stratagem for primary root elongation under moderate salt stress in the halophyte Schrenkiella parvula. *Physiol Plant* **175**:e13937. doi:10.1111/ppl.13937

Shukla V, Han J-P, Cléard F, Lefebvre-Legendre L, Gully K, Flis P, Berhin A, Andersen TG, Salt DE, Nawrath C, Barberon M. 2021. Suberin plasticity to developmental and exogenous cues is regulated by a set of MYB transcription factors. *Proc Natl Acad Sci U S A* **118**. doi:10.1073/pnas.2101730118

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**:1312–1313. doi:10.1093/bioinformatics/btu033

Team RC, Team MRC, Suggests M, Matrix S. 2018. Package stats. *The R Stats Package*.

Uga Y, Sugimoto K, Ogawa S, Rane J, Ishitani M, Hara N, Kitomi Y, Inukai Y, Ono K, Kanno N, Inoue H, Takehisa H, Motoyama R, Nagamura Y, Wu J, Matsumoto T, Takai T, Okuno K, Yano M. 2013. Control of root system architecture by DEEPER ROOTING 1 increases rice yield under drought conditions. *Nat Genet* **45**:1097–1102. doi:10.1038/ng.2725

Ursache R, De Jesus Vieira Teixeira C, Dénervaud Tendon V, Gully K, De Bellis D, Schmid-Siegert E, Grube Andersen T, Shekhar V, Calderon S, Pradervand S, Nawrath C, Geldner N, Vermeer JEM. 2021. GDSL-domain proteins have key roles in suberin polymerization and degradation. *Nat Plants* **7**:353–364. doi:10.1038/s41477-021-00862-9

Valenzuela CE, Acevedo-Acevedo O, Miranda GS, Vergara-Barros P, Holuigue L, Figueroa CR, Figueroa PM. 2016. Salt stress response triggers activation of the jasmonate signaling pathway leading to inhibition of cell elongation in Arabidopsis primary root. *J Exp Bot* **67**:4209–4220. doi:10.1093/jxb/erw202

Wang X, Zhang Y, Wang L, Pan Z, He S, Gao Q, Chen B, Gong W, Du X. 2020. Casparian strip membrane domain proteins in Gossypium arboreum: genome-wide identification and negative regulation of lateral root growth. *BMC Genomics* **21**:340. doi:10.1186/s12864-020-6723-9

Wang Y, Meng B, Zhong S, Wang D, Ma J, Sun W. 2018. Aboveground biomass and root/shoot ratio regulated drought susceptibility of ecosystem carbon exchange in a meadow steppe. *Plant Soil* **432**:259–272. doi:10.1007/s11104-018-3790-7

Weigel D, Mott R. 2009. The 1001 genomes project for Arabidopsis thaliana. *Genome Biol* **10**:107. doi:10.1186/gb-2009-10-5-107

Wei T, Simko V. 2017. R package" corrplot": Visualization of a Correlation Matrix (Version 0.84). *Retrieved from, https://github com/taiyun/corrplot*.

West G, Inzé D, Beemster GTS. 2004. Cell cycle modulation in the response of the primary root of Arabidopsis to salt stress. *Plant Physiol* **135**:1050–1058. doi:10.1104/pp.104.040022

Wilke CO. 2016. cowplot: streamlined plot theme and plot annotations for "ggplot2." *CRAN Repos*.

Wu SJ, Ding L, Zhu JK. 1996. SOS1, a Genetic Locus Essential for Salt Tolerance and Potassium
Acquisition. *Plant Cell* **8**:617–627. doi:10.1105/tpc.8.4.617
Xu W, Cui K, Xu A, Nie L, Huang J, Peng S. 2015. Drought stress condition increases root to shoot ratio via
alteration of carbohydrate partitioning and enzymatic activity in rice seedlings. *Acta Physiol Plant*
37. doi:10.1007/s11738-014-1760-0
Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**:1586–1591.
doi:10.1093/molbev/msm088
Yu L'ang, Sussman H, Khmelnitsky O, Ishka MR, Srinivasan A, Nelson ADL, Julkowska MM. 2023.
Development of a mobile, high-throughput, and low-cost image-based plant growth phenotyping
system. *bioRxiv*. doi:10.1101/2023.07.18.549560
Yu B, Zheng W, Xing L, Zhu J-K, Persson S, Zhao Y. 2022. Root twisting drives halotropism via stress-
induced microtubule reorientation. *Dev Cell* **57**:2412–2425.e6. doi:10.1016/j.devcel.2022.09.012
Zarebanadkouki M, Trtik P, Hayat F, Carminati A, Kaestner A. 2019. Root water uptake and its pathways
across the root: quantification at the cellular scale. *Sci Rep* **9**:12979. doi:10.1038/s41598-019-
49528-9
Zou Y, Gigli-Bisceglia N, van Zelm E, Julkowska MM, Zeng Y, Cheng Y, Koevoets IT, Jørgensen B, Giesbers
M, Vroom J, Ketelaar T, Petersen BL, Zhang Y, Testerink C. 2022. Cell wall extensin arabinosylation is
required for root directional response to salinity. *bioRxiv*. doi:10.1101/2022.06.22.497042
Zou Y, Gigli-Bisceglia N, Zelm E, Kokkinopoulou P. 2022. Cell wall extensin arabinosylation is required for
root directional response to salinity. *bioRxiv*.