Title: *CDC42 Inhibitors Alter Patterns of Vessel Arborization in Skin and Tumors in vivo.*

Authors: *Linh M Vuong[1], Stephanie Hachey[2], Jessica Shiu[1], Danny F Xie[3], Noel Salvador[4], Nicoletta Brindani[5], Sine Mandrup Bertozzi[6], Maria Summa[7], Rosalia Bertorelli[7], Andrea Armirotti[8], Rachel Pham[1], Vance SH Ku[1], Terry Nguyen[1], Benard Choi[3], Christopher CWW Hughes[2], Marco De Vivo[5], Anand K Ganesan[1]*

[1]*Department of Dermatology, University of California, Irvine 92697*

[2]*Department of Biochemistry and Molecular Biology, University of California, Irvine 92697*

[3]*Beckman Laser Institute, University of California, Irvine 92697*

[4]*Department of Biological Chemistry, University of California, Irvine 92697*

[5]*Molecular Modeling and Drug Discovery Lab, Istituto Italiano di Tecnologia, Genova, Italy 16163*

[6]*Translational Pharmacology Facility, Istituto Italiano di Tecnologia, Genova, Italy 16163*

[7]*Translational Pharmacology Facility, Istituto Italiano di Tecnologia, Genova, Italy 16163*

[8]*Analytical Chemistry Facility, Istituto Italiano di Tecnologia, Genova, Italy 16163*

Abstract: Tumors that arise in the epidermis must develop a vascular supply to grow beyond a millimeter in depth. Despite this dependence on angiogenesis for tumor growth, melanoma tumors are minimally responsive to anti-angiogenesis agents. Existing methods to measure changes in angiogenesis rely on two dimensional approaches that have limited ability to capture changes in the three-dimensional architecture of the tumor vascular network. Here we integrate antibody infusion, optical tissue clearing, multiphoton imaging, and three-dimensional semi-automated tracing to quantify changes in the vascular architecture of skin and tumors. Initial studies used this new vessel visualization approach to demonstrate that RhoJ knockout mice have marked differences in vessel arborization in the skin despite having similar numbers of endothelial cells. Small molecules that inhibit CDC42 GTPases, which include RhoJ, RhoQ, and CDC42, inhibit tumor growth and vessel branching within tumors to a similar degree as Braf inhibitors. In contrast to Braf inhibitors, which only affect vessel arborization in tumors, CDC42 inhibitors alter vessel arborization in normal skin, colon, and human skin organoids. Taken together, these studies identify a new class of pharmacologic agents that inhibit vessel branching in both normal skin and tumors which could have utility in treating skin cancer and skin disorders characterized by pathologic angiogenesis.

Introduction:

Vascular morphogenesis in tissues is a dynamic process where endothelial cells modulate their cytoskeleton and cell matrix adhesions to properly respond to environmental cues (*1*). The dynamic rearrangement of endothelial cells' cytoskeleton allows them to migrate both forward and backward within the growing blood vessel (*2-4*). Generation of the capillary plexus depends on the regression of existing vessels during the maturation process, and similar types of vessel regression are observed in conditions where the adult vasculature needs to be remodeled (*5*). All tumors, particularly those that arise in the epidermis that lack their own vasculature, are dependent on the ability to induce the growth of vessels around them (*6-8*). Tumors rapidly accelerate angiogenesis by producing a cornucopia of angiogenesis-promoting factors, inducing a haphazard, tortuous vessel tree that is unable to deliver nutrients evenly to the tumor (*9, 10*).

Combining tissue clearing with semi-automated tracing has provided great insight into how angiogenesis occurs in the skin and other tissues (*11-13*). This approach has been recently applied to tumors (*14, 15*). The tumor vasculature is characterized by large tortuous vessels with leaky basement membranes and unperfused voids (*16, 17*). While longitudinal imaging approaches have provided structural insight into how vessels develop or repair after injury (*5, 18*), this information alone has not identified better therapies for conditions characterized by pathologic angiogenesis, such as inherited skin vascular anomalies (*19, 20*) or melanoma tumors (*6*), where existing anti-angiogenic therapies are many times toxic or only minimally effective (*21*).

Many intracellular signaling pathways are known to activate angiogenesis (*22*). Vascular endothelial growth factors, which activate intracellular signaling cascades that induce actin remodeling and focal adhesion assembly (*23*), are major chemo-attractants for migrating endothelial cells. For example, spatial activation of VEGFR2 signaling is critical in the maintenance of the adult vasculature (*5*). Semaphorins have also been identified as regulators of vasculogenesis- Semaphorin 3E binds to the PlexinD1 receptor to induce receptor internalization, focal adhesion disassembly, and, subsequently, vessel regression (*24, 25*). CDC42 GTPases are critical mediators of both attractive and repulsive cues generated by activation of VEGF and Plexin receptors, respectively (*26*). RhoJ, a protein with 55% homology to CDC42, is highly expressed in endothelial cells, induces actin depolymerization, focal adhesion disassembly, and endothelial cell contraction (*27, 28*). CDC42, in contrast, activates focal adhesion assembly and stimulates the migration of endothelial cells (*29-32*). CDC42 GTPases mediate crosstalk between these receptors, as RhoJ is suppressed by VEGFR2 and activated by Plexin while CDC42 is activated

by VEGFR2 and suppressed by Plexin (*28*). RhoJ deficiency results in impaired VEGF-induced endothelial cell migration (*33*), impaired retinal vascular angiogenesis early during development (*34, 35*), and impaired tumor angiogenesis (*36*). Targeting CDC42 GTPases with small molecules has been problematic secondary to their globular structure and lack of readily apparent druggable pockets (*37, 38*). We recently used a structure-based drug design approach to discover a class of CDC42 GTPase inhibitors that block CDC42 downstream effector interactions (*39, 40*). These agents blocked S6 and ERK activation, inhibited angiogenesis in vascularized tumor organoids, and inhibited the growth of mouse and patient-derived tumors *in vivo* (*39, 40*).

Here we sought to determine how RhoJ and drugs that inhibit CDC42 signaling modulate angiogenesis *in vivo*. We developed a platform combining tissue clearing and vascular labeling with fluorescent antibodies and three-dimensional semi-automated vessel tracing to demonstrate that RhoJ deficient mice had significantly altered vessel arborization in their skin despite having only modest changes in the number of skin endothelial cells. We applied our new imaging approach to image patient-derived xenografts treated with CDC42 inhibitors and other standard-of-care melanoma care therapies (BRAF inhibitors), demonstrating that both agents had similar tumor suppressive effects and inhibited blood vessel branching and the accumulation of vessel endpoints. In contrast to vemurafenib, however, CDC42 inhibitors inhibited vessel branching in the skin and colon of mice without apparent tissue toxicity, identifying these agents as novel therapies for disorders characterized by disordered angiogenesis in skin or colon.

Results:

RhoJ deletion disrupts the vascular network in the skin of RhoJ knockout (KO) mice.

RhoJ knockout mice have decreased branching of the retinal vasculature during development, a phenotype that improves with age (*26*). RhoJ is also known to play a role in adult vascular homeostasis, as RhoJ knockout mice had impaired tumor angiogenesis (*36*) and impaired skin wound healing (*41*). Initial work sought to examine if RhoJ knockout mice had altered vascular architecture in the skin as compared to RhoJ wild-type mice. We previously combined tissue clearing with intravascular infusion of DyLight 649 labeled tomato-lectin to visualize normal vasculature in the brain (*42, 43*). Here we modified an existing iDISCO (**i**mmunolabeling three-**D**imensional **I**maging of **S**olvent-**C**leared **O**rgans) (*44*) protocol to clear skin tissue harvested from RhoJ knockout and wild-type mice infused with Tomato-lectin prior to sacrifice (Fig 1A). Cleared skin was imaged with a Leica SP8 multiphoton microscope to generate Z-stack images of skin and measure differences in vessel architecture between wild type and RhoJ knockout mice using

two different analytic pipelines.   A gross comparison of flattened images generated from RhoJ knockout and wild-type mouse skin revealed that RhoJ knockout mice had a decrease in the number of skin vessels (Fig 1B). Differences in vascular architecture in the skin were quantified by two different methods. AngioTool is a software package that takes 2D images, quantifies pixels in grayscale, and measures junctions and endpoints. This package has been used to measure differences in the vascular architecture of the murine embryonic hindbrain, post-natal retina, and allantois explant(45). NeuTube, in contrast, takes 3D z-stacks to trace neurons and can quantify terminal nodes, branch nodes, and the total number of neurons (46).   We compared the ability of these two methods to detect differences in vessel architecture in RhoJ knockout (RhoJ-KO) and wild-type (RhoJ-WT) skin.   AngioTool analysis revealed that the skin of RhoJ-KO mice had a decreased number of vessel junctions and endpoints as compared to RhoJ-WT skin (Fig 1C). Analysis of the same images with neuTube revealed that RhoJ-KO skin had fewer terminal nodes, branch nodes, and vessels as compared to RhoJ-WT skin (Fig S1A-C).   Despite the observed decrease in vessel density, RhoJ-KO skin showed no obvious pathological changes in the epidermis and dermis as visualized by haematoxylin and eosin staining (Fig 1D).   We did, however, observe a slight decrease in thickness of the superficial dermal layer of RhoJ-KO skin, as highlighted by a Verhoeff-Van Gieson stain (Fig 1D).   Flow cytometry analysis revealed that RhoJ-KO skin and RhoJ-WT skin had similar number of endothelial cells (CD31+ cells (47)) and fibroblasts (PDGRα+ cells (48)) in their skin (Figure 1E) and spleen (Figure1F).   Taken together, these results indicate that while RhoJ deletion affects vascular arborization in the skin, it does not grossly affect skin structure or the viability of fibroblasts or endothelial cells. Of note, three-dimensional quantitative approaches were able to better capture differences in the arborization of the vasculature in mice, and this approach was used to measure changes in vessel arborization in tumors, which are thicker structures than the skin.

CDC42 inhibitors alter vascular arborization patterns in patient-derived xenografts implanted into mice.

Previous work identified a class of CDC42 interaction inhibitors that could inhibit tumor growth *in vivo (39, 40)* and block angiogenesis in vascularized microtumors *in vitro* (*40*).   To examine how CDC42 inhibitors modulate tumor angiogenesis *in vivo*, we implanted patient-derived xenografts into the flanks of NSG mice. When the tumors reached a volume of 150-200 mm^3, we treated the mice with the indicated doses of the CDC42 inhibitor ARN22089, vemurafenib, or vehicle twice daily for two weeks (Figure 2A). We observed that ARN22089 inhibited the growth of tumors as effectively as vemurafenib despite the fact that the reported Kd for vemurafenib binding to its

target Braf is >10-fold lower (*49*) than the IC50 of ARN22089 for CDC42 (*40*). To compare how ARN22089 and vemurafenib affected arborization of vessels in tumors, we: 1) sacrificed the mice after the two week treatment; 2) infused the mice with tomato lectin intravascularly; 3) cleared the tumors using our iDISCO protocol; 4) imaged these tumors with multiphoton microscopy to obtain three-dimensional image stacks; 5) converted 3D images to grayscale images; and 6) quantified vascular branching using neuTube (Figure 2B). Gross examination of stacked images from drug-treated mice revealed a decrease in vasculature in treated mice as compared to controls (Fig 2C, upper images).   Upon neuTube tracing of the vessel architecture (Fig 2C, lower images), ARN22089 affected vessel number and branching to a greater extent than vemurafenib. NeuTube analysis of images revealed that ARN22089 had a dose-responsive effect on the number of branches, the number of terminal nodes, vessel tortuosity, and vessel length (Fig 2D). We also performed daily tail vein injection of a second CDC42 lead inhibitor (ARN25062) and observed that this additional lead candidate: 1) stunted tumor growth (Fig S2A); and 2) reduced vessel arborization in tumors grossly (Fig S2B). Additionally, vessel arborization changes induced by these analogues could be detected using both AngioTool and neuTube (Fig S2C,D). Finally, we stained ARN22089 treated tumors with CD31 and revealed that treated tumors had fewer CD31 stained vessels histologically (Fig 2E).

CDC42 inhibitors block angiogenesis in human-derived vascularized microtumors *in vitro*. Previous work had demonstrated that ARN22089 could inhibit vessel growth in human derived vascularized microtumors *in vitro* using single-chamber VMT models (*50*). To more easily determine whether CDC42 inhibitors had direct effects on tumor vasculature, we used microtumor VMT devices where two vascular chambers had an intervening tumor chamber so that one could measure the effects of drugs after infusion on one side of the chamber versus the infusion of the other with vehicle (Fig 3A). ARN22089 disrupted the vasculature of the drug-treated but not the control side of the vasculature with a concomitant effect on tumor growth (Fig 3B), similar to the effects that were observed previously (*40*), with quantifiable effects on vessel length (Fig 3C). Next, we sought to examine whether ARN25062 and vemurafenib could also inhibit vessel growth in similar devices where tumor cells and blood vessels are spatially separated (Fig 3D). We observed that both ARN25062 and vemurafenib visually and quantitatively inhibited tumor growth and blood vessel length in vascularized microtumors that were comprised of A375 cells (Fig 3E,F) and WM3248 melanoma cells (3G). Taken together, these results confirm that CDC42 inhibitors block the arborization of either mouse or human derived vessels around human melanoma tumors.

CDC42 inhibitors disrupt skin vasculature in mice in vivo and in human-derived vascularized microorgans in vitro

RhoJ knockout mice had notable changes in the vascular arborization of skin compared to wild-type mice (Fig 1). As CDC42 inhibitors should also block RhoJ function, we next compared the effects of ARN22089 or vemurafenib treatment on the arborization of vessels in the skin. For these experiments: 1) C57B6 mice were treated with either ARN22089 (4, 12, or 40 mg/kg), vehicle, or 25 mg/kg vemurafenib twice daily for one week; 2) mice were infused with Tomato-lectin intravascularly prior to sacrifice; 3) skin tissues were cleared and imaged with three-dimensional multiphoton microscopy in 3D; 4) changes in vessel arborization were quantified using AngioTool. Gross examination of image stacks revealed that CDC42 inhibitors altered skin vascular arborization patterns, which were even more obvious when examining flattened images (Fig 4A, S3B). AngioTool was used to quantify changes in branching in the 2D images, revealing that ARN22089 had a dose-responsive effect on the number of observed junctions and endpoints (Fig 4B). In contrast, vemurafenib had no effect on the number of observed junctions and endpoints (Fig 4B). ARN22089 did not have any gross effects on skin appearance (Fig S3A) but did have modest effects on the thickness of the superficial dermal layer as detected by VVG staining and standard haematoxylin and eosin staining (Fig 4C), similar to the effects observed in RhoJ knockout mice.

To further verify that ARN22089 affects skin vascularization, we harvested skin from the NSG mice bearing tumors that were treated with CDC42 inhibitors, vehicle, or vemurafenib at regions that were at a minimum 5 mm away from the implanted tumor. We observed that ARN22089 treatment at either 20 mg/kg or 40 mg/kg BID inhibited the accumulation of vessel junctions and endpoints as measured by AngioTool (Fig S4A). Notably, vemurafenib did not significantly alter either the accumulation of vessel junctions or endpoints (Fig S4B), similar to what was observed in C57B6 skin.

Next, we examined how ARN22089, ARN25062, and vemurafenib affected the vascularization of entirely human-derived vascularized micro-organs that are comprised of human-derived endothelial cells, pericytes, and fibroblasts but not tumor cells (Fig 4D). We observed gross inhibition of vessel structure formation in these vascularized micro-organs that were treated with ARN22089 or ARN25062 but not with vemurafenib or vehicle (Fig 4E). AngioTool quantitation of

VMO images revealed that ARN22089 and ARN25062 affected vessel length and the observable number of vessel branchpoints (Fig 4F).

To examine tissue toxicity, we next examined how ARN22089 affected the vascularization of other organs. ARN22089 had no effect on blood vessel accumulation in the brain (Fig S5A), although pK data did reveal that ARN22089 could cross the blood brain barrier to a small degree (Fig S5B). Haematoxylin and eosin staining of cardiac tissue harvested from treated animals revealed no gross effects on the heart (Fig S5C) and drug treatment had no effect on mouse weight (Fig S5D). Intriguingly, we did observe an effect on the accumulation of vasculature in the colon (Fig S5E) without any observable effects on the histology of the colon (data not shown).

CDC42 inhibitors affect the vascularization of the skin by similar mechanisms observed in RhoJ knockout mice.

CDC42 inhibitors block both the interaction of RhoJ and CDC42 with their downstream effectors (*40*), and both RhoJ knockout mice and CDC42 inhibitor-treated mice had similar skin vessel arborization patterns. To examine whether CDC42 inhibitors modulate skin angiogenesis by inhibiting RhoJ, we treated wild-type and RhoJ knockout mice with the CDC42 inhibitor ARN22089 and used the same approach as described in Figure 4 to measure changes in vascularization. Analysis of stacked and flattened images from the treated mice revealed no apparent differences in the vessel arborization patterns of RhoJ knockout mice after ARN22089 treatment (Fig 5A). Similarly, we observed no differences in junctions or endpoints as measured by AngioTool (Fig 5B). Taken together, these results are consistent with a model in which both CDC42 and RhoJ are critical regulators of blood vessel progression and regression in skin and tumors, such that inhibiting one or both of these pathways can alter the dynamics of blood vessels such that they affect arborization in skin and tumors (Fig 5C).

Discussion

While many anti-angiogenic agents have been identified to treat cancer and other conditions, all of the existing agents have significant and frequent side effects in nearly half of treated patients, such as skin rash, diarrhea, and hypertension (*51, 52*). The cumulative side effects of these therapies preclude their use in conditions that arise in infancy, such as hereditary hemorrhagic telangiectasia, or in chronic diseases that require continuous treatment, such as inflammatory skin diseases. Moreover, the effects of VEGF inhibitors on vasculature are transient, as the normal vessel structures can return even one or two days after treatment cessation (*53, 54*).

Existing methods to measure changes in vasculature rely on 2D quantitative approaches (*55, 56*), although several different 3D approaches have recently been used that integrate tissue clearing to measure tumor vascularization (*14, 16, 17*). We observe that a 3D quantitative approach (neuTube) provides more insight into vascular structure than approaches where 3D images are flattened into 2D. This simple and straightforward approach can be applied to measure changes in vessel arborization in tumors.

Here, we sought to use our approach to examine how targeting angiogenesis pathways other than the conventional VEGF pathway, namely CDC42 GTPase-regulated pathways, affects angiogenesis. In contrast to the transient effects of RhoJ deletion on retinal vascular angiogenesis (*34, 35*), RhoJ knockout mice had a persistent decrease in vessel number in adult skin, suggesting that RhoJ has an important role in skin vascular homeostasis (Fig 1C). RhoJ deletion did not affect the number of endothelial cells and fibroblasts in skin (Fig 1E), in contrast to the significant differences in vessel branching observed. This suggests that RhoJ deletion affects endothelial cell behavior rather than survival. It is important to note that while the observed effects can be seen in normal skin, deletion does not grossly affect skin structure as the epidermis was intact and hair follicles were observed in RhoJ-KO mouse skin.

CDC42 inhibitors largely recapitulated the effects observed with RhoJ deletion, and their activity required RhoJ as they did not inhibit angiogenesis in RhoJ-KO mouse skin (Fig 5). CDC42 inhibitors did not affect the vasculature of the heart or the brain but did affect vasculature in the colon (Fig S5E). It is known that vessel regression, which is regulated by RhoJ (*29, 57*), is important in the maintenance of the vascular plexus in the skin (*5*). Whether the observed tissue-specific differences are a consequence of different roles/expression patterns of RhoJ in distinct endothelial populations will be a topic for further study. Of note, other studies have shown that inhibiting receptors that activate RhoJ (Glutamate synthetase) can block inflammation in the skin induced by imiquimod (*41*), which further supports the role of RhoJ in skin vascular homeostasis. We observed that the effects in mice were preserved in human vascular organoids (Fig 4D-F), indicating that human blood vessels were also responsive to these drugs. The observation that CDC42 inhibitors can block angiogenesis in both mice and vascular organoid models suggested that CDC42 inhibitors could be used to treat conditions characterized by increased skin vascularity, such as rosacea, psoriasis, and photoaging. These agents may also be appropriate in treating conditions characterized by pathologic angiogenesis (port-wine birthmarks, hereditary hemorrhagic telangiectasias) where VEGF inhibitors are not indicated secondary to systemic toxicity.

Melanoma tumors are minimally responsive to anti-angiogenesis agents(*6*). Tumor vessels can rapidly regrow after the cessation of VEGF-targeted therapies (*53, 54, 58*), and VEGF inhibitors seem to affect the blood vessels in the periphery but not the central part of the tumor(*17*). Here we used a combination of antibody administration, tissue clearing, and multiphoton microscopy to compare the effects of established melanoma therapies (BRAF inhibitors) and CDC42 inhibitors in tumors. CDC42 inhibitors normalized vessel tortuosity and decreased vessel number, as did vemurafenib, a finding that has been reported for VEGF inhibitors (*17*). A closer examination of the histology of tumors revealed that vemurafenib-treated tumors were characterized by large areas of tissue necrosis, in contrast to CDC42 inhibitors where such broad necrosis was not observed (data not shown). Vessel tracing results suggest that CDC42 inhibitors also affect large vessels in the tumor, in contrast to vemurafenib where large vessels were mostly preserved (Fig 2C). Moreover, vemurafenib did not affect skin vasculature, in contrast to CDC42 inhibitors. These observations suggest that vemurafenib likely affects vessel number by inducing the death and subsequent necrosis of melanoma cells with secondary effects on vasculogenesis, in contrast to CDC42 inhibitors which inhibit vasculogenesis directly in both skin and tumors.

In summary, the work presented here provides evidence that CDC42 GTPases play differing roles in vascular homeostasis in adult tissues. While it is currently not clear whether the effects of CDC42 inhibition will be more permanent than VEGF inhibitors, the relative lack of systemic toxicity, particularly as CDC42 inhibitors do not affect cerebral vasculature whereas bevacizumab crosses the blood-brain barrier, suggests that these agents could be used, at minimum, as topical therapies for skin cancer and vascular/ inflammatory skin disease.

Materials and Methods:

In vivo skin and tumor experiments. All animal experiments were approved by the UC Irvine Institutional Animal Care and Use Committee (IACUC) (AUP-20-161). C57BL6 mice and RhoJ knockout mice in the C57B6 background were used in studies examining skin, colon, and brain vascularization while NOD.Cg-*Prkdcscid IL2rgtm1Wj*/SzJ (NSG) mice were used for the patient-derived xenograft experiments. An equal number of male and female mice were used in treatment with inhibitors. For skin treatment, wild-type or RhoJ knockout C57BL6 mice were treated with inhibitors by oral gavage twice daily for one week. For PDX tumor experiments, tumors were maintained by passaging into NSG mice before treatment. Inoculation of tumors in NSG mice was performed as previously described (*50*). When tumors reached ~150-200 mm^3, mice were treated

by oral gavage twice daily with 20 or 40 mg/kg ARN22089 or with vemurafenib at 10 or 25 mg/kg. Tail-vein injection was done once a day at 10 mg/kg ARN22089. All tumor experiments involved treating animals for two weeks while skin experiments involved treatment for 1 week.

Cardiac perfusion and tissue collection. At time of harvest, mice were infused with lectin-DyLight-649 (200 uL, 25% lectin-DyLight and 75% PSB) through the tail vein injection to label endothelial cells. One hour later, mice were euthanized by cardiac perfusion with 50 mL saline followed by 4% methanol-free paraformaldehyde (PFA) as described previously(*43*). For skin, the fur was shaved and depilated and dorsal skin was removed from the back. Tumors were removed from the flanks of the dorsal region. Brain, kidney, intestines, livers, hearts, and tissues were placed in 4% PFA for 24-48 h and then placed in 1xPBS at 4°C for subsequent tissue clearing.

Tissue clearing and imaging analysis. A modified iDISCO protocol was used to clear tissues. Tissues were dehydrated in graded series (20, 40, 60, 80, 100, 100%) of methanol in water for 48 h at RT. Samples were incubated with 66% dichloromethane (DCM) and 33% methanol for 48 h at RT. Next, samples were washed twice for 15 min each with DCM and placed in dibenzyl ether (DBE) for storage and used as a medium for imaging. The Leica TCS SP8 X instrument was used to image tissues and processed with a LAS X Navigator suite. The sample was placed on a makeshift holder attached to a microscope slide and filled with DBE; a coverslip was placed over the sample, making sure the sample and liquid touches the coverslip. Images were captured using an HC PL FLOUTAR 10x/0.30 objective. Image dimension: 1024 x 1024 pixel dimension and 1107.14 μm by 1107.14 μm, 5 μm per z-stack. Vascular fluorescence was detected by scanning for signals in the DyLight649 spectrum (670-708 nm). For image analysis, Z-stack images were opened with Fiji ImageJ, and color images were converted to black background and white pixels using color-split and saved as a tiff file either in three-dimenasional (3D) or compressed into 2D image using Z stack projection. 3D tiff files were analyzed using neuTube software(*46*), while 2D images were analyzed using AngioTool software(*45*). Both automated and manual tracing were performed on Tiff images in neuTube. Automated tracing was applied initially, with manual curating as needed due to images with low contrast between vessels and background and where large vessels cannot be correctly auto-traced. A custom MATLAB script was used to extract the number of branch and end points, vessel length, vessel diameter, and tortuosity in the neuTube SWC format files (individual nodes with x, y, z coordinates, radius and node connectivity). In the SWC format, vessel structures are simplified into spherical nodes with the radius of the vessel at a given location. Branch points (green nodes in images) appear where branches occur, and end points (yellow nodes in images) appear at the end of a vessel segment (all other nodes are red

in images). A vessel segment is defined as a series of nodes that are bounded by two branch points or one branch point and one end point. The length of a vessel segment is defined as the distance between its pair of boundary points. The diameter of the vessel is calculated using the radius of the nodes. Parameters for AngioTool analysis were set the same for all images analyzed. For NSG skin stack images, the first and last few images in the stacks were removed to reduce background noise; all stacks from C57BL6 skin were analyzed. For image analysis >3 tumors or skin tissues were cleared, imaged and analyzed.

Statistics. GraphPad Prism software was used to generate graphs and perform statistical significance, using one- and two-way ANOVA test and unpaired T-test. One-way ANOVA and unpaired T-test were applied on vessel analysis and two-way ANOVA test was used to determine significance for tumor growth curves. Custom R script was used to convert pixel value to micrometer for diameter and distance; it was also used to calculate the frequency of branch nodes, end nodes, number of vessels, and tortuosity.

Microfluidic device fabrication Microfluidic device fabrication and loading have been previously described (5-6). In summary, a custom polyurethane master mold was created using a two-part polyurethane liquid plastic (Smooth Cast 310, Smooth-On Inc.). Subsequently, a PDMS layer was replicated from this master mold, and holes were punched to create inlets and outlets. The platform was assembled in two stages: first, the PDMS layer was chemically glued and subjected to 2 minutes of oxygen plasma treatment to affix it to the bottom of a 96-well plate. Following this, a 150 μm thin transparent membrane was bonded to the bottom of the PDMS device layer through an additional 2-minute treatment with oxygen plasma. The fully assembled platform was then placed in a 60°C oven overnight, covered with a standard 96-well plate polystyrene lid, and sterilized using UV light for 30 minutes before cell loading.

Cell culture and microfluidic device loading To establish the vascular chamber, normal human lung fibroblasts and ECFC-ECs (endothelial colony forming cell endothelial cells) or HUVECs (human umbilical vein endothelial cells) were harvested and resuspended in fibrinogen solution at a concentration of 3×10^6 cells/mL and 7×10^6 cells/mL, respectively. For VMT, A375 and WM3248 melanoma cells were introduced into the tumor chamber at a concentration of 1×10^5 to 2×10^5 cells/mL fibrinogen solution. Fibrinogen solution was prepared by dissolving 70% clottable bovine fibrinogen (Sigma- Aldrich) in EBM2 basal media (Lonza) to a final concentration of 5 mg/mL. The cell-matrix suspension was mixed with thrombin (50 U/mL, Sigma-Aldrich) at a concentration of 3 U/mL, quickly seeded into the microtissue chambers, and allowed to polymerize in a 37 ◦C incubator for 15 minutes. Laminin (1 mg/mL, LifeTechnologies) was then

introduced into the microfluidic channels through medium inlets and incubated at 37 ∘C for an additional 15 minutes. After incubation, culture medium (EGM-2, Lonza) was introduced into the microfluidic channels and medium wells. The medium was changed every other day, and the hydrostatic pressure head re-established daily to maintain interstitial flow.

Drug treatment in the VMO and VMT Following a culture period of 4–5 days to facilitate the development of a perfused vasculature within each VMO or VMT, the culture medium was replaced with a medium containing the specified drug concentrations. Drugs were administered to the microtissues through the newly formed vascular bed via gravity-driven flow. Specifically, ARN25062, ARN22089, and vemurafenib were used at a 2 µM dose. A375 VMT, WM3248 VMT, and VMO were randomly assigned to one of four conditions: control (vehicle only), 2 µM ARN25062, 2 µM ARN22089, or 2 µM vemurafenib. For experiments with two vascular side chambers, the left chamber received 2 µM ARN22089, while the right chamber served as control (vehicle only). Both VMO and VMT underwent a 48-hour treatment period, with complete medium replacement every 48 hours. Fluorescent micrographs of VMT were captured every 48 hours for 6 days post-treatment, and the quantification of tumor and vasculature growth was performed.

Fluorescence imaging and analyses Fluorescence images were acquired with a Biotek Li-onheart fluorescent inverted microscope using automated acquisition and standard 10x air ob-jective. To test vessel perfusion, 25 µg/mL FITC- or rhodamine- conjugated 70 kDa dextran was added to the medium inlet prior to treatment. For quantifying vessel length in VMOs and VMTs, REAVER software (MATLAB (7)) was employed. ImageJ software (National Institutes of Health) was utilized to determine the total fluorescence intensity (mean grey value) for each tumor image, providing a measure of tumor growth. Normalization to baseline was performed for each chamber. In VMTs, tumor growth was quantified by measuring the total fluorescence intensity in the color channel representing the tumor cells. This measurement accounted for both the area and depth of individual tumors, considering that thicker areas appear brighter. Any image adjustments made were applied uniformly to ensure consistency across all images in the experimental group.

Tissue Histology and Immunohistochemistry Tissues were fixed in 4% formaldehyde and washed in 1xPBS. For embedding, tissues were incubated in 65% ethanol for 30 min at RT and kept in 70% ethanol before sending the samples to Experimental Tissue Resource (ETR) at UCI for embedding, VVG (*59*) and H&E staining (*60*).

Flow Cytometry Back of mouse was first shaved and depilated. A large surface area (6 cm x 3 cm) of the skin on the back was cut and fat removed. The skin was minced, and digest buffer

added (RPMI without Ca/Mg (no FBS/EDTA), 23.2 mM HEPES and 2.32 mM sodium pyruvate, 0.25 mg/mL Liberase, 46.4 unit DNase) and incubated for 1.5 h at 37°C shaking. Spleen samples were removed from the animals, minced and RPMI added to minced tissue. All samples were filtered with 70 micrometer cell strainer, and washed with FAC buffer (5% FBS, 2 mM EDTA, 1% non-essential amino acids, 3.9x beta-mercapthoethanol (from 1000x)) twice and spun at 18.0 xg for 10 min 4°C. Skin samples were blocked with TruStain FcX PLUS and spleen samples with Fc block, both, at 1:100 for 15 min at 4°C. All samples washed twice with FAC buffer and incubated in primary Cd31 (1:50, brilliant violet 421 Invitrogen 404031182), PDGFRa (1:100, APA5 Life Technology 12140181), viability (1:200, eBioscience dyefluor 780 65086514) in FAC buffer for 30 min at 4°C nutating. Samples were then washed 2x and fixed with 1% formaldehyde for 15 min and washed twice again. Samples were resuspended with 400 uL FAC buffer and ran on the BD Fortessa X20 Flow cytometer. Compensation and FMO for both skin and spleen were run before running the experimental samples. Samples were analyzed using FlowJo.

Quantification of brain levels of arn22089 following oral administration Animals were treated with ARN22089 at 10 mg/Kg by oral administration, as described (39). Mouse brains were collected at 1, 2, 4 and 8 hours after administration. The brains were homogenized and, following protein precipitation with acetonitrile, the amount of ARN22089 was quantified by LC-MS/MS as described (39).

Acknowledgement: Funding for this work was largely derived from (CA-244571) from the National Cancer Institute and S10OD028698. This study was made possible in part through access to the Optical Biology Core Facility, a shared resource supported by the Cancer Center Support Grant (CA-62203) and UCI Skin (AR-075047) at the University of California, Irvine.

Author contributions:
Conceptualization: AKG, CCWWH, MDV, BC, LMV
Funding acquisition and project administration: AKG
Data curation: LMV, SJH, JS, SMB, NB, MS, RB, AA
Formal analysis: LMV, SJH, JS, SMB, NB, MS, RB, AA, VSHK
Investigation: LMV, SJH, JS, NS, DFX, TN
Writing – original draft: LMV, AKG
Writing – review and editing: LMV, AKG, CCWWH, MDV, BC, DFX, SJH, JS
Methodology: NB, SMB, MS, RB, AA, TN, NS, DFX, RP, VSHK
Conflicts of interest: No competing interests were reported by the authors.

References:

1. H. Gerhardt *et al.*, VEGF guides angiogenic sprouting utilizing endothelial tip cell filopodia. *J Cell Biol* **161**, 1163-1177 (2003).
2. L. Jakobsson *et al.*, Endothelial cells dynamically compete for the tip cell position during angiogenic sprouting. *Nat Cell Biol* **12**, 943-953 (2010).
3. H. W. Lee, J. H. Shin, M. Simons, Flow goes forward and cells step backward: endothelial migration. *Exp Mol Med* **54**, 711-719 (2022).
4. S. Arima *et al.*, Angiogenic morphogenesis driven by dynamic and heterogeneous collective endothelial cell movement. *Development* **138**, 4763-4776 (2011).
5. C. Y. Kam *et al.*, Mechanisms of skin vascular maturation and maintenance captured by longitudinal imaging of live mice. *Cell* **186**, 2345-2360 e2316 (2023).
6. Z. Wu *et al.*, The role of angiogenesis in melanoma: Clinical treatments and future expectations. *Front Pharmacol* **13**, 1028647 (2022).
7. W. C. Cho, G. Jour, P. P. Aung, Role of angiogenesis in melanoma progression: Update on key angiogenic mechanisms and other associated components. *Semin Cancer Biol* **59**, 175-186 (2019).
8. D. R. Bielenberg, B. R. Zetter, The Contribution of Angiogenesis to the Process of Metastasis. *Cancer J* **21**, 267-273 (2015).
9. R. Lugano, M. Ramachandran, A. Dimberg, Tumor angiogenesis: causes, consequences, challenges and opportunities. *Cell Mol Life Sci* **77**, 1745-1770 (2020).
10. D. Hanahan, J. Folkman, Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* **86**, 353-364 (1996).
11. E. J. Lee *et al.*, Three-dimensional visualization of cerebral blood vessels and neural changes in thick ischemic rat brain slices using tissue clearing. *Sci Rep* **12**, 15897 (2022).
12. K. Takahashi *et al.*, An analysis modality for vascular structures combining tissue-clearing technology and topological data analysis. *Nat Commun* **13**, 5239 (2022).
13. H. Kolesova, V. Olejnickova, A. Kvasilova, M. Gregorovicova, D. Sedmera, Tissue clearing and imaging methods for cardiovascular development. *iScience* **24**, 102387 (2021).
14. N. Frenkel *et al.*, Tissue clearing and immunostaining to visualize the spatial organization of vasculature and tumor cells in mouse liver. *Front Oncol* **13**, 1062926 (2023).
15. S. Kostrikov *et al.*, Optical tissue clearing and machine learning can precisely characterize extravasation and blood vessel architecture in brain tumors. *Commun Biol* **4**, 815 (2021).
16. B. J. Stolz *et al.*, Multiscale topology characterizes dynamic tumor vascular networks. *Sci Adv* **8**, eabm2456 (2022).
17. M. Dobosz, V. Ntziachristos, W. Scheuer, S. Strobel, Multispectral fluorescence ultramicroscopy: three-dimensional visualization and automatic quantification of tumor morphology, drug penetration, and antiangiogenic treatment response. *Neoplasia* **16**, 1-13 (2014).
18. N. MacRitchie, P. Maffia, Light sheet fluorescence microscopy for quantitative three-dimensional imaging of vascular remodelling. *Cardiovasc Res* **117**, 348-350 (2021).
19. K. Ricci, Medical Therapeutics for the Treatment of Vascular Anomalies: Part 3. *Oral Maxillofac Surg Clin North Am* **36**, 125-136 (2024).
20. R. Gupta, Propranolol for Vascular Anomalies: Efficacy and Complications in Pediatric Patients. *J Indian Assoc Pediatr Surg* **28**, 194-205 (2023).
21. R. N. Gacche, Changing landscape of anti-angiogenic therapy: Novel approaches and clinical perspectives. *Biochim Biophys Acta Rev Cancer* **1878**, 189020 (2023).
22. Z. L. Liu, H. H. Chen, L. L. Zheng, L. P. Sun, L. Shi, Angiogenic signaling pathways and anti-angiogenic therapy for cancer. *Signal Transduct Target Ther* **8**, 198 (2023).

23. M. Shibuya, Vascular Endothelial Growth Factor (VEGF) and Its Receptor (VEGFR) Signaling in Angiogenesis: A Crucial Target for Anti- and Pro-Angiogenic Therapies. *Genes Cancer* **2**, 1097-1105 (2011).

24. W. J. Oh, C. Gu, The role and mechanism-of-action of Sema3E and Plexin-D1 in vascular and neural development. *Semin Cell Dev Biol* **24**, 156-162 (2013).

25. J. Kim, W. J. Oh, N. Gaiano, Y. Yoshida, C. Gu, Semaphorin 3E-Plexin-D1 signaling regulates VEGF function in developmental angiogenesis via a feedback mechanism. *Genes Dev* **25**, 1399-1411 (2011).

26. A. Uemura, Y. Fukushima, Rho GTPases in Retinal Vascular Diseases. *Int J Mol Sci* **22**, (2021).

27. T. T. Shi, G. Li, H. T. Xiao, The Role of RhoJ in Endothelial Cell Biology and Tumor Pathology. *Biomed Res Int* **2016**, 6386412 (2016).

28. Y. Fukushima *et al.*, RhoJ integrates attractive and repulsive cues in directional migration of endothelial cells. *EMBO J* **39**, e102930 (2020).

29. A. Sundararaman, H. Mellor, A functional antagonism between RhoJ and Cdc42 regulates fibronectin remodelling during angiogenesis. *Small GTPases* **12**, 241-245 (2021).

30. Y. Wakayama, S. Fukuhara, K. Ando, M. Matsuda, N. Mochizuki, Cdc42 mediates Bmp-induced sprouting angiogenesis through Fmnl3-driven assembly of endothelial filopodia in zebrafish. *Dev Cell* **32**, 109-122 (2015).

31. J. Ma *et al.*, Role of activated Rac1/Cdc42 in mediating endothelial cell proliferation and tumor angiogenesis in breast cancer. *PLoS One* **8**, e66275 (2013).

32. G. D. Hu *et al.*, The generation of the endothelial specific cdc42-deficient mice and the effect of cdc42 deletion on the angiogenesis and embryonic development. *Chin Med J (Engl)* **124**, 4155-4159 (2011).

33. S. Kaur *et al.*, RhoJ/TCL regulates endothelial motility and tube formation and modulates actomyosin contractility and focal adhesion numbers. *Arterioscler Thromb Vasc Biol* **31**, 657-664 (2011).

34. H. Takase *et al.*, Genome-wide identification of endothelial cell-enriched genes in the mouse embryo. *Blood* **120**, 914-923 (2012).

35. S. Kusuhara *et al.*, Arhgef15 promotes retinal angiogenesis by mediating VEGF-induced Cdc42 activation and potentiating RhoJ inactivation in endothelial cells. *PLoS One* **7**, e45858 (2012).

36. C. Kim *et al.*, Vascular RhoJ is an effective and selective target for tumor angiogenesis and vascular disruption. *Cancer Cell* **25**, 102-117 (2014).

37. M. D. M. Maldonado, J. I. Medina, L. Velazquez, S. Dharmawardhane, Targeting Rac and Cdc42 GEFs in Metastatic Cancer. *Front Cell Dev Biol* **8**, 201 (2020).

38. M. D. M. Maldonado, S. Dharmawardhane, Targeting Rac and Cdc42 GTPases in Cancer. *Cancer Res* **78**, 3101-3111 (2018).

39. N. Brindani *et al.*, Design, Synthesis, In Vitro and In Vivo Characterization of CDC42 GTPase Interaction Inhibitors for the Treatment of Cancer. *J Med Chem* **66**, 5981-6001 (2023).

40. S. Jahid *et al.*, Structure-based design of CDC42 effector interaction inhibitors for the treatment of cancer. *Cell Rep* **39**, 110760 (2022).

41. G. Eelen *et al.*, Role of glutamine synthetase in angiogenesis beyond glutamine synthesis. *Nature* **561**, 63-69 (2018).

42. M. Loren, C. Crouzet, A. Bahani, V. Vasilevko, B. Choi, Optical clearing potential of immersion-based agents applied to thick mouse brain sections. *PLoS One* **14**, e0216064 (2019).

43. K. Khouri *et al.*, Simple methodology to visualize whole-brain microvasculature in three dimensions. *Neurophotonics* **8**, 025004 (2021).

44. N. Renier *et al.*, iDISCO: a simple, rapid method to immunolabel large tissue samples for volume imaging. *Cell* **159**, 896-910 (2014).

45. E. Zudaire, L. Gambardella, C. Kurcz, S. Vermeren, A computational tool for quantitative analysis of vascular networks. *PLoS One* **6**, e27385 (2011).
46. L. Feng, T. Zhao, J. Kim, neuTube 1.0: A New Design for Efficient Neuron Reconstruction Software Based on the SWC Format. *eNeuro* **2**, (2015).
47. N. V. Goncharov *et al.*, Markers of Endothelial Cells in Normal and Pathological Conditions. *Biochem (Mosc) Suppl Ser A Membr Cell Biol* **14**, 167-183 (2020).
48. L. Yao *et al.*, Temporal control of PDGFRalpha regulates the fibroblast-to-myofibroblast transition in wound healing. *Cell Rep* **40**, 111192 (2022).
49. G. Bollag *et al.*, Vemurafenib: the first drug approved for BRAF-mutant cancer. *Nat Rev Drug Discov* **11**, 873-886 (2012).
50. S. Jahid *et al.*, Structure-based design of CDC42 effector interaction inhibitors for the treatment of cancer. *Cell Rep* **39**, 110641 (2022).
51. R. J. Bodnar, Anti-Angiogenic Drugs: Involvement in Cutaneous Side Effects and Wound-Healing Complication. *Adv Wound Care (New Rochelle)* **3**, 635-646 (2014).
52. F. Elice, F. Rodeghiero, Side effects of anti-angiogenic drugs. *Thromb Res* **129 Suppl 1**, S50-53 (2012).
53. M. R. Mancuso *et al.*, Rapid vascular regrowth in tumors after reversal of VEGF inhibition. *J Clin Invest* **116**, 2610-2621 (2006).
54. T. Inai *et al.*, Inhibition of vascular endothelial growth factor (VEGF) signaling in cancer causes loss of endothelial fenestrations, regression of tumor vessels, and appearance of basement membrane ghosts. *Am J Pathol* **165**, 35-52 (2004).
55. Q. Zhao *et al.*, Quantitative analysis on in vivo tumor-microvascular images from optical-resolution photoacoustic microscopy. *J Biophotonics* **12**, e201800421 (2019).
56. F. Grizzi *et al.*, Quantitative evaluation and modeling of two-dimensional neovascular network complexity: the surface fractal dimension. *BMC Cancer* **5**, 14 (2005).
57. A. Sundararaman, Y. Fukushima, J. C. Norman, A. Uemura, H. Mellor, RhoJ Regulates alpha5beta1 Integrin Trafficking to Control Fibronectin Remodeling during Angiogenesis. *Curr Biol* **30**, 2146-2155 e2145 (2020).
58. C. G. Lee *et al.*, Vascular endothelial growth factor (VEGF) induces remodeling and enhances TH2-mediated sensitization and inflammation in the lung. *Nat Med* **10**, 1095-1103 (2004).
59. M. A. Piccinin, J. Schwartz, in *StatPearls*. (Treasure Island (FL) ineligible companies. Disclosure: Janice Schwartz declares no relevant financial relationships with ineligible companies., 2024).
60. A. H. Fischer, K. A. Jacobson, J. Rose, R. Zeller, Hematoxylin and eosin staining of tissue and cell sections. *CSH Protoc* **2008**, pdb prot4986 (2008).

Figure 1: RhoJ regulates skin vessel arborization in skin.

Fig. 1: RhoJ regulates skin vessel arborization in skin. (A) Diagram of optical tissue clearing and vessel analysis. Each image (1.107 x 1.107 mm; 1024 x 1024 pixels) has about >50 z-stacks (5 µm/stack). Z-stacks are compressed into 2D Tiff file, converted to grayscale, and analyzed with AngioTool. Images were generated from three mice per group. (B) Representative images of skin vasculature from RhoJ wild type (WT) and knockout (KO) mice. Fluorescence images of lectin labeled structures were obtained, converted to grayscale, and traced with AngioTool. Scale bar is 50 µm. (C) Scatter bar plot of the frequency that vessel branches and termini were observed in (B); $^{*,***}p < 0.013, 0.0009$; n= 3 mice per genotype. (D) VVG elastin and H and E staining of skin from RhoJ-WT and RhoJ-KO mice. Note the presence of an intact epidermis and hair follicles indicating that RhoJ deletion did not induce skin necrosis. (E) Scatterplot showing the percentage of $Cd31^+$ or $Pdgfra^+$ live cells in the skin and spleen from >4 mice per group (RhoJ-WT and RhoJ-KO).

Figure 2: Cdc42 inhibitors alter vascular arborization patterns in tumors.

Fig. 2: CDC42 inhibitors alter vascular arborization patterns in tumors. (A) Plot of tumor volume as a function of time in mice treated with vehicle (blue line), vemurafenib (orange line), or ARN22089 (purple [20 mg/kg] and red [40 mg/kg] lines) at the indicated doses. Graph represents mean+SEM. *p-value < 0.03 **p-value < 0.0071; ***p-value < 0.0001 (n=5 mice per group). (B) Schematic diagram of tissue clearing, visualization, and vessel analysis of the tumors. Each image (1.107 x 1.107 mm; 1024 x 1024 pixels) has >200 z-stacks (5 µm/stack). (C) Representative z-stacks of fluorescent images from tumors in mice treated with vehicle, vemurafenib (25 mg/kg twice daily) or ARN22089 (4, 12 or 40 mg/kg twice daily) orally are shown. Bottom represents corresponding grayscale vessel images; scale is 100 µm. (D) Scatter plot with bar graphs show the statistical quantification of vessel parameters (number of branch/end points and vessels, average diameter, length, and tortuosity) after 3D vessel tracing for >3 tumors per group. . *,**,***,****p < 0.02, 0.01, 0.002, 0.0001. (E) Representative H and E and CD31 staining of tumors showing a decrease in vessel-stained structures in tumors that were ARN22089 treated.

Figure 3: Cdc42 inhibitors disrupt vasculature in tumor organoids.

Figure 3. CDC42 inhibitors disrupt vasculature in tumor organoids. A) Diagram showing a double chamber microfluidic design where two independently fed vascular chambers (green and red) flank a central 200 μm wide tumor chamber (blue) separated by 3 PDMS posts on each side. The left vascular chamber (green) was treated with ARN22089 (2 μM) on day 6, while the right side received vehicle only (control, red). B) Representative fluorescent micrographs of double chamber VMT with A375 (green) in the center and vasculature (red) on either side. C) Quantification of A375 tumor-associated vasculature showing fold change in vessel length from baseline for left side (treated) vs. right side (control). D) Schematic showing a single dual-chamber microfluidic device. The vascular chamber connects to the tumor chamber (800 μm wide), separated by 6 PDMS posts spaced 50 μm apart that serve as burst valves to prevent the gel from traversing the chamber. EC and LF are introduced into loading port L1, and cancer cells are introduced separately into loading port L2. Loading is facilitated by a pressure regulator (PR). Tissues are maintained via hydrostatic pressure generated across microfluidic channels connecting media reservoirs M1-M2. Physiological flow rates are established by microfluidic resistors. E) Representative fluorescent micrographs of dual-chamber VMT containing A375 tumor (green) and vasculature (red), treated with either control (vehicle only), ARN25062 (2 μM), or vemurafenib (2 μM) for 48 hours starting on day 4. Media was refreshed on day 6, day 8, and day 10. Scale bar = 500 μm. F) Quantification of A375 tumor growth and tumor associated vasculature in the VMT showing fold change from baseline $^*p < 0.05$, $^{**}p < 0.01$. G) Quantification of WM3248 tumor growth and tumor-associated vasculature showing fold change in tumor vessel length from baseline. $^*p < 0.05$, $^{**}p < 0.01$.

Figure 4: Cdc42 inhibitors alter vessel arborization in skin.

Fig. 4: CDC42 inhibitors alter vessel arborization in skin. (A) Skin of mice treated with vehicle, vemurafenib, and ARN22089 at the indicated doses were cleared and imaged to generate Z-stacks as described, with representative stack shown. Bottom of each stack includes the 2D AngioTool tracing of pixels in grayscale. Scale bar is 100 μm. Each image (1.107 x 1.107 mm; 1024 x 1024 pixels) has >50 z-stacks (5 μm/stack). (B) Scatter plot with bar graphs show the frequency of branching and termini quantified from flattened images using the 2D AngioTool software. ***p-value < 0.0001, >3 mice per condition. (C) VVG elastin and H and E staining of skin from WT and RhoJ KO mice. Similar to the RhoJ KO, the presence of an intact epidermis and hair follicles are observed in all samples, with a slight decrease in VVG highlighted areas in treated animals. (D) Schematic showing a single dual-chamber microfluidic device. The vascular chamber is 800 μm wide, separated by 6 PDMS posts spaced 50 μm apart that serve as burst valves to prevent the gel from traversing the chamber. EC and LF are introduced into loading port L1. Loading is facilitated by a pressure regulator (PR). Tissues are maintained via hydrostatic pressure generated across microfluidic channels connecting media reservoirs M1-M2. Physiological flow rates are established by microfluidic resistors. (E) Representative fluorescent micrographs of dual chamber VMO showing vasculature (greyscale), treated with either control (vehicle only), ARN25062 (2 μM), ARN22089 (2 μM) or vemurafenib (2 μM) for 48 hours starting on day 4. Media was refreshed on day 6, day 8, and day 10. (F) Quantification of VMO-associated vasculature showing fold change in vessel length and number of branchpoints compared to baseline. *p < 0.05, **p < 0.01.

Figure 5: Cdc42 inhibitors block skin angiogenesis in a RhoJ dependent manner.

Fig. 5: CDC42 inhibitors block skin angiogenesis in a RhoJ dependent manner. (A) Representative grayscale and AngioTool tracing of RhoJ KO skin vessels (treated with vehicle or 40 mg/kg ARN22089). (B) Scatter plot showing the frequency of branches and termini as measured by the 2D AngioTool software; n.s. no significant difference. At least 3 mice per condition were tested. (C) Schematic of RhoJ/Cdc42 affecting angiogenesis. RhoJ is known to affect vessel regression while CDC42 affects vessel elongation. Inhibitors can block both functions resulting in altered vessel arborization patterns in skin and tumors.