MICROGLIA AGING IN THE HIPPOCAMPUS ADVANCES THROUGH INTERMEDIATE STATES THAT DRIVE INFLAMMATORY ACTIVATION AND COGNITIVE DECLINE

Jeremy M. Shea[1]# and Saul A. Villeda[1-3],#

1. Department of Anatomy, University of California San Francisco, San Francisco, California 94143, USA

2. Department of Physical Therapy and Rehabilitation Science, University of California San Francisco, San Francisco, California 94143, USA

3. Bakar Aging Research Institute, University of California San Francisco, San Francisco, California, 94143, USA

#Corresponding Authors:

Dr. Saul Villeda, PhD	Dr. Jeremy Shea, PhD
University of California San Francisco	University of California San Francisco
Department of Anatomy	Department of Anatomy
513 Parnassus Ave, Box 0452	513 Parnassus Ave, Box 0452
San Francisco, CA 94143	San Francisco, CA 94143
Email: saul.villeda@ucsf.edu	Email: jeremy.shea@ucsf.edu
Tel: 415-502-1929	Tel: 617-460-4122

SUMMARY

During aging, microglia – the resident macrophages of the brain – exhibit dystrophic phenotypes and contribute to age-related neuroinflammation. While numerous hallmarks of age-related microglia dystrophy have been elucidated, the progression from homeostasis to dysfunction during the aging process remains unresolved. To bridge this gap in knowledge, we undertook complementary cellular and molecular analyses of microglia in the mouse hippocampus across the adult lifespan and in the experimental aging model of heterochronic parabiosis. Single-cell RNA-Seq and pseudotime analysis revealed age-related transcriptional heterogeneity in hippocampal microglia and identified intermediate states of microglial aging that also emerge following heterochronic parabiosis. We tested the functionality of intermediate stress response states via TGFβ1 and translational states using pharmacological approaches *in vitro* to reveal their modulation of the progression to an inflammatory state. Furthermore, we utilized single-cell RNA-Seq in conjunction with an *in vivo* adult microglia-specific *Tgfb1* conditional genetic knockout mouse model, to demonstrate that microglia advancement through intermediate aging states drives inflammatory activation and associated hippocampal-dependent cognitive decline.

INTRODUCTION

Microglia are one of few immune cells with long term residence in the brain parenchyma[1]. Microglia support neural development by synaptic pruning and neurogenic function in the developing brain, followed by homeostatic maintenance in the adult brain[2]. Alternatively, in the aged brain microglia exhibit dystrophic phenotypes, aggravating perceived insults rather than responding productively to environmental signals[3]. While microglia have been identified as principal mediators of multiple age-related neurodegenerative pathologies[4], mechanisms leading to microglia dysfunction during aging remain obscure.

Numerous studies have indicated that aged microglia are inflamed[5], have reduced phagocytic capacity[6,7], and have decreased motility[8]. Microglia exhibit several hallmarks of aging that potentially contribute to their age-related dysfunction, such as shortened telomeres[9], altered intercellular communication[10], molecular alterations[11], and a loss of proteostasis[12]. Furthermore, many recent studies have started to reveal the molecular changes that define microglial aging. Single cell RNA-Seq (scRNA-Seq) analyses indicate that microglia isolated from the entire brain lose homeostasis and activate inflammatory transcriptional profiles with age[13,14]. Data rich studies have also revealed partial overlaps between aging microglia and those from disease models, including Alzheimer's disease[15-19]. Interestingly, the microglial response to aging shows regional variation, as white matter-rich regions induce an interferon signature in aged microglia[20]. Studies using aged plasma administration and heterochronic blood exchange demonstrate that microglia aging is in part driven by the aged systemic environment [21,22]. However, the genesis of age-related dystrophic microglial phenotypes has not been extensively investigated. So, we set out to characterize the progression of age-related hippocampal microglial dysfunction, aiming to uncover intermediate states that could be intrinsic to the aging process. To do so, we undertook complementary cellular and molecular analyses of microglia across the adult lifespan and in heterochronic parabiosis - an experimental model of aging in which the circulatory systems of young adult and aged animals are joined[23].

In this study, we report that microglia in the adult mouse hippocampus, a brain region responsible for learning and memory and susceptible to age-related cognitive decline, advance through intermediate states that drive inflammatory activation during aging. We utilize scRNA-Seq across the adult lifespan to identify intermediate transcriptional states of microglial aging that emerge following exposure to an aged systemic environment. Functionally, we tested the role of these intermediate states using *in vitro* microglia approaches and an *in vivo* temporally controlled adult microglia-specific *Tgfb1* conditional genetic knockout mouse model to demonstrate that

118 intermediates represent checkpoints in the progression of microglia from homeostasis to
119 inflammatory activation, with functional implications for hippocampal-dependent cognitive decline.
120

RESULTS
Single-cell transcriptional analysis reveals the dynamics of hippocampal microglia aging
Previously, single-cell transcriptional profiling of microglia uncovered a diversity of responses
during development and in response to pathology aggregated over multiple brain
regions[13,14,16,19,24,25]. Microglia have region-specific transcriptional states, so we decided to focus
our analyses on the hippocampus[11], a region with well-reported functional differences during
aging[26]. Correspondingly, we investigated age-related transcriptional heterogeneity by performing
scRNA-Seq on hippocampal CD11b-positive cells isolated from mice at 6, 12, 18, and 24 months
of age using a commonly employed pooling strategy[18,27,28] (n = 1 pool of 5 biological replicates for
each age) to reveal the dynamics of microglia aging. 82% of cells isolated with CD11b were
identified as microglia using canonical markers with a small subset being proliferative (Figure
S1A-C). Dimensionality reduction revealed that non-proliferative microglia were grouped into
interconnected clusters that contained homeostatic, transition, and activated states, along with a
semi-distinct cluster with interferon activation (Figure 1a, S1D-G). Overlaying microglia ages onto
the clusters revealed progression from a homeostatic state in younger microglia to an activated
state in old microglia, suggesting that gradual transcriptional changes occur during microglial
aging (Figure 1B,C, S1H). Furthermore, the heterogeneity of microglial transcriptional programs
increased with age (Figure S1I). Expression of *Cx3cr1* and *Itgam* exemplified the homeostatic
state, while *B2m*, *Apoe*, *Cd48*, and *Lyz2* depict the activated state (Figure 1D). Interestingly,
several immediate early genes (*Jun*, *Klf2*, *Fos*), as well as the purinergic receptor gene *P2ry12*,
exhibited transient increases in expression at middle-age (Figure 1D). Alternatively, the interferon
cluster represents a small proportion of the overall microglia population with little age-related
changes in the number of these cells, but augmented expression of interferon genes during aging
(Figure 1A,C,D, S1G). Thus, gene expression changes exhibited unique patterns during aging,
with certain genes showing age-related decreases (e.g. *Tgfbr1*) or increases (e.g. *Apoe*), while a
subset of genes (e.g. *Tgfb1*) have peaks of expression at middle age (Figure 1D,E,G; Table S1).
Several complement genes implicated in age-related synaptic loss[10,29](e.g. *C1q* and *C3*)
demonstrated progressive age-related transcriptional increases, and the increased complement
activity in microglia was confirmed by immunohistochemistry (Figure 1E,F).
The effects of aging on microglia could be progressive or sudden, so we investigated if we
could find evidence of early aging of hippocampal microglia. We observe relatively few

differentially genes between 6- and 12-month (Figure 1G,H, Table S1). The number of
differentially expressed genes gradually increased through the 24-month time point (Figure 1G-J,
Table S1), suggesting that hippocampal microglia progressively accumulate aged-related
expression changes. Furthermore, when investigating the outcome for genes with significant
changes for each age, we find that genes that are differentially expressed at 12 months lose their
expression change, while genes that change at later ages show progressive expression
differences (Figure 1H). GO analysis of the microglia from 6- and 12-month-old mice revealed
that activation and translational programs are modestly enriched in genes with increased
expression during early aging (Figure 1I). The latter stages of aging display more prominent
enrichment of immune activation and translation processes in genes with increased expression,
while revealing that genes that regulate transcriptional processes exhibit decreased expression
(Figure 1J,K). We corroborated the progressive increase in microglial activation by
immunohistochemical analysis for CD68 and p65 (Figure S1J,K). Thus, microglia progressively
age with subsets of cells at each age group being forerunners towards inflammatory activation or
refractory to aging by remaining homeostatic. This analysis further suggests that microglia could
pass through transitory states during aging.
168

Trajectory analysis indicates that aging microglia pass through intermediate states
To determine the transcriptional progression of microglia aging towards an inflammatory state,
we performed pseudotime analysis of the scRNA-Seq data and observed several trajectories
(Figure 2A). We find that the pseudotime trajectories progressed along the biological ages of the
mice, indicating the suitability of this analysis for modeling aging progression in microglia (Figure
2B). We focus our analysis on one trajectory with two semi-distinct subbranches that terminated
in inflammatory activation, since the alternate branch had minimal transcriptional differences
between the beginning and end of the trajectory (Figure 2B,C). To gain insight into the
transcriptional states of microglial aging along this inflammatory trajectory, we used spatial
autocorrelation analysis and identified five expression modules of coregulated genes (Figure
2B,C, Table S2). We named these modules according to the most enriched GO pathways and
known roles of prominent genes for microglia. This trajectory proceeds from high expression of
homeostatic genes and mitochondrial processes (module 1) to transient expression of stress
response genes and *Tgfb1* (module 2), of which TGFβ signaling is critical for microglia
development[30]. A concerted increase in expression of ribosomal genes (module 3) precedes one
subbranch of inflammatory activation (module 4), while the other myeloid activation subbranch
(module 5), characterized by *B2m* and *C1qc* expression, proceeds independently of increased

ribosomal gene expression. GO analysis of genes specifically upregulated at 12-months showed

an enrichment of stress response pathways (Figure S2A,B), corroborating the findings of the

pseudotime analysis that intermediate stages of microglial aging pass through stress response

pathways. Thus, pseudotime trajectory analysis identifies a progression of intermediate states of

microglial aging with potential involvement in age-related maladaptive inflammation.

The gene level dynamics of microglia aging are further revealed when interrogating the

modules at each timepoint. Expression of several genes in the homeostatic cluster, including

Cx3cr1 and *Tgfbr1*, progressively decline throughout the aging timeline (Figure 2D). Stress

response gene expression peaks at early timepoints during aging, while those genes in the

translation module have coherent upregulation at 24-months of age (Figure 2D). The top ranked

genes in the inflammation module gradually increase their expression throughout aging in a

consistent manner (Figure 2D). Alternatively, several genes in the myeloid activation module have

expression peaks at 12- (*Cst3*, *Selplg*) *or* 18-months (*Ctss*, *Cd9*) of age (Figure 2D). When the

modules are collapsed into metagenes, we observe that homeostatic module is reduced early on

in aging, while the stress response module follows later (Figure 2E-G). The ribosomal and

inflammatory modules display augmented expression at latter ages (Figure 2E-G). Alternatively,

the myeloid activation consistently has slightly increased expression throughout aging. These

results suggest that with advancing age microglia lose homeostatic and stress response

functions, while increasing their translational and inflammatory capacity.

Next, we utilized a publicly available scRNA-Seq datasets of pro-aging systemic

interventions and age-related neurodegenerative disease models to determine their impact on

hippocampal microglia aging trajectories. We used the heterochronic parabiosis model[31] to

investigate whether exposure to an aged systemic environment could promote progression of

young adult microglia along the identified aging trajectory (Figure 2H). First, using

immunohistochemistry for IBA1 and CD68, we reveal that an aged systemic environment

activates microglia (Figure S2C). Next, we find that hippocampal microglia from young adult

heterochronic parabionts have decreased expression of homeostatic genes (module 1), while

having increased expression of ribosomal genes (modules 3), inflammatory activation genes

(module 4) and myeloid activation genes (module 5) (Figure 2I, S2D). We used the *App*$^{NL-G-F}$

transgenic mouse model at 12 months of age to determine the effects of Alzheimer's disease

pathology on microglial aging[17] (Figure 2J). We observe exaggerated advancement along the

aging trajectory in *App*$^{NL-G-F}$ mice compared to controls, as expression is shifted towards the age-

associated modules (Figure 2K, S2E,F). These shift in gene expression posit that the aged

systemic and diseased environments drive adult microglia advancement along an aging-
associated inflammatory trajectory.

Intermediate states of microglia aging act as checkpoints on inflammatory progression
Next, we sought to corroborate microglial age-related molecular changes associated with
individual intermediate states in the aging hippocampus by RNAscope and immunohistochemical
analysis. Specifically, we assessed changes in the intermediate stress response (module 2) and
translation (module 3) modules by examining expression of *Tgfb1* (Figure 3A,B) and the ribosomal
protein S6 (Figure 3G,H), respectively. The vast majority (>95%) of *Tgfb1* signal localized to
microglia at every age (Figure S3A), indicating microglia are the predominant source for
hippocampal TGFB1. Consistent with scRNA-Seq data of aging microglia (Figure 1E), we
observed highest *Tgfb1* expression in microglia by middle-age (Figure 3B). TGFBR1 activation
was reduced during aging in hippocampal microglia (Figure S3B), and genes involved in the TGFβ
signaling pathway exhibited age-related expression changes with several having peak expression
at 12 months of age (Figure S3C). Additionally, we observed an increase in translational
components found during the scRNA-Seq analysis by immunohistochemistry for the ribosomal
protein S6 (Figure 3H)[32].

To complement hippocampal aging analysis, we assessed the impact of the aging
systemic milieu on both the TGFβ pathway and translational components in the young adult
hippocampus following heterochronic parabiosis. We observed a decrease in microglia *Tgfb1* and
TGFβ pathway expression in young adult heterochronic parabionts compared to age-matched
young adult isochronic controls[31] (Figure 3C, S3D). Additionally, expression of the translation
module in microglia increases following exposure to an aged systemic environment in the
heterochronic parabiosis model (Figure 3I).

To investigate the role of the intermediate stress response (module 2) and translation
(module 3) modules in mediating advancement through microglial activation states, we used an
in vitro approach. Primary microglia were treated with LPS, which induced gene expression
changes with a significant overlap to aging, as well as expression changes in aging modules
genes (Figure S3E,F; Table S3). We administered TGFB1 to activate TGFβ signaling (module 2)
(Figure 3A) and CX-5461 to inhibit RNA Pol I synthesis and interfere with translation (module 3)
(Figure 3G) in the context of microglial activation. TGFB1 treatment modified the transcriptional
states of LPS-treated microglia (Figure S3G) and caused decreased expression of inflammatory
immune processes after LPS stimulation (Figure 3D). TGFB1 treatment reduced expression of
later aging modules (modules 2, 3 and 4) and restored expression of the homeostatic gene *Cx3cr1*

(Figure 3E,F), suggesting that TGFBβ signaling acts as an aging checkpoint during microglial
stress response. Targeting translation (module 3) with CX-5461 attenuated expression of immune
processes after LPS stimulation (Figure 3J; Table S3), as well as decreased expression of genes
in modules 2, 3, and 4, while restoring expression of homeostatic genes in module 1 (Figure
3K,L). Interestingly, CX-5461 treatment had no effect on genes in module 5, which is located on
an independent myeloid activation trajectory from altered translational gene expression (Figure
3K,L). These *in vitro* perturbation data identify active roles for intermediate states in mediating
advancement along aging-associated inflammatory trajectories.

Mimicking age-related changes in microglia-derived TGFB1 promotes microglial
advancement along an aging-associated inflammatory trajectory *in vivo*
Having uncovered roles for intermediate states in advancing microglia along aging-associated
inflammatory trajectories *in vitro*, we next investigated the role of individual aging modules in adult
microglia *in vivo*. We elected to disrupt the transition from homeostasis to inflammatory activation
by targeting microglia-derived TGFB1, a key marker of the stress response intermediate state
(module 2). While TGFB1 is critical for microglia development[33], the role of microglia-derived
TGFB1 in regulating microglia homeostasis during aging remains to be defined.

270   We generated *Tgfb1^{flox/flox}*, *Tgfb1^{flox/wt}*, and *Tgfb1^{wt/wt}* mice carrying an inducible *Cx3cr1-*
Cre-ER gene, in which *Tgfb1* is excised specifically in adult microglia upon tamoxifen
administration (*Tgfb1* cKO, *Tgfb1* Het, and WT, respectively) (Figure 4A). To disrupt the age-
related increase in microglia *Tgfb1* observed between mature and middle-age (Figure 1E and
Figure 3B), mature adult mice were administered tamoxifen and molecular changes were
assessed two months later (Figure 4A). We find that microglia activation, as measured by
CD68/IBA1 immunohistochemistry, exhibited genotype-dependent effects (Figure S4A). We
performed scRNA-Seq on microglia isolated from WT, *Tgfb1* Het and *Tgfb1* cKO mature mice (n
= 2 pools of 3 animals per genotype), and detected genotype-dependent changes in
*Tgfb1*expression levels, as well as effectors of TGFB1 signaling (Figure S4B). Dimensionality
reduction and clustering of cells from scRNA-Seq revealed readily identifiable clusters based on
marker expression (Figure 4B). Composition of these clusters were dependent on the microglia
genotype, with WT microglia representing the largest fraction of the homeostatic cluster and *Tgfb1*
cKO microglia representing the largest fraction of the activated cluster (Figure 4C, S4C). Next,
we assessed whether deletion of *Tgfb1* in mature adult microglia impacted advancement along
aging-associated inflammatory trajectories. Interestingly, when we analyze the effects of *Tgfb1*
dosage on the aging modules, we find that the loss of a single *Tgfb1* allele reduces expression of

homeostatic genes (module 1), while increasing expression of translation-related genes (module
3) (Figure 4D,E). In addition to a decrease in expression of homeostatic genes (module 1), loss
of both *Tgfb1* alleles caused a greater increase in expression of translation-related genes (module
3), and led to a profound increase in inflammatory activation genes (module 4) (Figure 4D,F).
Results were corroborated in an independent cohort of mature *Tgfb1* cKO mice, in which we
detected altered levels of immune- and age-related cell surface markers (Figure S4D,E),
transcriptional alterations that significantly overlapped with aging (Figure S4F-H; Table S4), and
progression along the aging-associated inflammatory trajectories (Figure S4I,J). These data
indicate that microglia-derived TGFB1 is both necessary for expression of youth-associated
homeostatic genes and prevents aberrant microglia inflammatory activation, further suggesting
that TGFBβ signaling acts as a checkpoint in mediating advancement along aging-associated
inflammatory trajectories.

300 Mimicking age-related changes in microglia-derived TGFB1 impairs hippocampal-
301 dependent cognitive function
It is becoming increasingly apparent that microglia interact with other cell types in the brain to
influence cognitive function[2,34]. Given loss of microglia-derived TGFB1 promoted microglia
progression along aging-associated inflammatory trajectories, we next investigated its functional
consequence on cognition. We assessed hippocampal-dependent learning and memory in WT,
Tgfb1 Het and *Tgfb1* cKO mature mice using novel object recognition and contextual fear
conditioning - behavioral paradigms that are sensitive to age-related impairments[35,36]. During
novel object recognition testing, WT mice were biased toward a novel object relative to a familiar
object while neither *Tgfb1* Het or *Tgfb1* cKO mice showed any preference (Figure 4G). During
contextual fear conditioning testing, *Tgfb1* Het and *Tgfb1* cKO mice exhibited less freezing
compared to WT controls (Figure 4H). Alternatively, we assessed amygdala-dependent fear
memory by cued fear conditioning (Figure S4K)[37] and short-term spatial reference memory by Y
maze (Figure S4L)[38], and observed no differences across genotypes. As a control, we also
profiled general health using an open field paradigm and observed no differences in total distance
traveled or time spent in the center of the open field, indicative of normal motor and anxiety
functions (Figure S4M). These behavioral data indicate that loss of microglia-derived TGFB1 not
only impacts aging-associated inflammatory trajectories but also promote hippocampal-
dependent cognitive impairments characteristic of age-related functional decline.

320 DISCUSSION

Cumulatively, we demonstrate that microglia aging in the hippocampus advances through
intermediate states that precede inflammatory activation, with functional implications for age-
related cognitive decline. These transient states, such as activation of the stress response and
TGFβ signaling, act as checkpoints on further age-related activation. Furthermore, microglia
asynchronously progress through this inflammatory aging trajectory that is in turn influenced by
the systemic environment and neurodegenerative disease states.

Analysis of our scRNA-Seq data elucidate cryptic, but necessary, intermediate states of
microglia progression to age-related inflammatory activation. These states are present in middle-
aged mice and constitute stress response pathways likely responsive to internal and external
cellular insults[39,40]. Pseudotime analysis suggests a highly plausible sequence of molecular
alterations that lead to age-related inflammatory activation of hippocampal microglia. First,
mitochondrial dysfunction triggers a stress response, and products of this stress response (e.g.
TGFB1) attempt to return microglia to homeostasis[41]. Here, chronic activation of the stress
response leads to subsequent increased translational capacity that promotes inflammatory
activation[42]. Alternatively, another branch of microglial activation advances independently of
increased translational capacity, evidenced by translational inhibition having little effect on its
activation following LPS stimulation. Thus, we find that increases in translational capacity
predicate age-related inflammatory phenotypes, identifying potential targets to counter microglial
aging. Analysis of scRNA-Seq data further identify intermediate states of microglial aging in young
adult heterochronic parabionts, suggesting a pivotal role for the aged systemic environment in
driving advancement towards age-related inflammatory activation.

While we find evidence of the beginnings of inflammatory activation in the hippocampus
by middle age (12 months), consistent with the timing recently reported in the whole brain[14], it
should be noted that scRNA-Seq analysis of the whole brain did not observe intermediate stages
in microglial aging[14]. These disparate observations could likely be the result of analyzing multiple
regions with highly differentiated transcriptional programs[11] that could obfuscate regional aging
trajectories. This further underscores the need for both temporal and regional specificity in aging
studies, given the level of cellular heterogeneity observed across ages and brain regions.
Moreover, studying microglia - or other cell types - at early stages of aging in a region-specific
manner could also represent useful models to investigate the ontogeny of aging without the
confounding effects of accumulated functional deficits.

Of note, we find that an inflammatory trajectory of microglia aging can be mitigated *in vitro*
by environmental TGFβ signaling, while loss of microglia-derived TGFB1 promoted progression
along aging-associated inflammatory trajectories *in vivo*. Together, these data indicate that TGFβ

signaling is necessary for maintaining youth-associated homeostasis and represents a checkpoint

in mediating advancement along aging-associated inflammatory trajectories. Functionally, loss of

microglia-derived TGFB1 resulted in hippocampal-dependent cognitive impairments, suggesting

that advancement through aging-associated inflammatory trajectories may facilitate age-related

cognitive decline. TGFβ likely acts in an autocrine fashion to return microglia to a homeostatic

state after encountering stressful stimuli. Given that a transient increase in *Tgfb1* expression is

observed in middle-aged microglia, it is possible that maintaining stress response-mediated

signaling into older ages may also provide a means to delay age-related microglial dysfunction

and maintain cognitive function. Alternatively, TGFβ signaling impairs microglial responses to

Alzheimer's disease pathology[43], suggesting that the dynamic regulation of TGFβ signaling in

response to aging-associated pathologies is necessary for tuning microglial responses.

While microglia have been demonstrated to mediate the establishment of cognitive

function during development[2] and play prominent roles in neurodegeneration[44], little is known

about how age-related changes in microglia affect cognitive function. The specificity of the

cognitive deficits that we observe in *Tgfb1* Het and *Tgfb1* cKO mice suggests that age-related

changes in microglia-derived TGFB1 predominantly impacts hippocampal-dependent cognitive

decline. Furthermore, the deficits observed in *Tgfb1* heterozygous mice point to the importance

of maintaining youthful levels of microglia-derived TGFB1 to counteract stressors that accrue

during aging.

Altogether, our results highlight the need to investigate aging across the lifespan in a

region-specific manner to reveal necessary cellular intermediates in the aging process. While the

present study focuses on age-related changes in microglia, our findings have broad implications

for other studies in determining the extent to which aging sequelae are conserved across different

cell types in the hippocampus. Furthermore, therapeutic interventions targeting intermediate

states in the aging process hold far reaching potential to counteract the development of age-

related pathologies, including those promoting cognitive dysfunction.

389 **ACKNOWLEDGMENTS**

390 We thank Dr. Erik Ullian for critically reading manuscript. **Funding**: This work was funded by

391 National Institute on Aging (AG055292 (J.M.S.), AG055797 (S.A.V.), AG077816 (S.A.V.)),

392 Simons Foundation (S.A.V). The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

J.M.S and S.A.V. developed concept and designed experiments. J.M.S conducted experiments, collected data, and performed analysis. J.M.S and S.A.V wrote manuscript. S.A.V supervised all aspects of the project. All authors had the opportunity to discuss results and comment on manuscript.

457 **COMPETING INTEREST STATEMENT**

458 The authors declare no competing interests.

Fig. 1 | Hippocampal microglia exhibit progressive age-related transcriptional inflammatory activation. A, UMAP plot of microglia separated into transcriptional clusters. (n = 1 pool of 5 animals for each age) **B,** Superimposition of ages onto the UMAP plot with dashed lines signifying relative cluster demarcation. **C,** Percent composition of each cluster by age. **D,** Dot plot of expression of cluster markers sorted by age. Percent of cells expressing the gene and average normalized expression are represented. **E,** Violin plots of genes with dynamic age-related expression patterns. **F,** Representative images and quantification of C1q (yellow), C3 (red), and Iba1 (cyan) staining in the hilus and molecular layer (ML) of 6- and 24-month-old mice. (n=4-5 mice per group; mixed effects analysis followed by Dunnett's multiple comparisons; *P<0.05, **P<0.01, ****P<0.0001). **G,** Number of differentially expressed genes from the 6-month timepoint at each age. Bars above the intersect represent increased expression and those below represent decreased expression. **H,** The average expression change at all ages for genes differentially regulated genes at individual ages represented by the color scheme in **B**. **I, J, K,** Volcano plots of differentially expressed genes in microglia between 6- and 12-months (**I**), 6- and 18-months (**J**), and 6- and 24-months (**K**) and corresponding gene ontology analysis of genes with significantly increased (brown) or decreased (green) expression for each comparison. Data are shown as mean±s.e.m.

Fig. 2 | Hippocampal microglia aging advances through intermediate states that respond
to systemic interventions or disease states. A, Pseudotime trajectories of microglia from an
anchor point located in 6-month microglia presented in a UMAP plot. (n = 1 pool of 5 animals for
each age) **B,** Microglia ages superimposed over pseudotime trajectories. **C,** Gene expression
modules representing sections of the inflammatory aging trajectory over the right half of the UMAP
plot (left). Modules were discovered using Moran's I autocorrelation test. Top Gene Ontology
terms and representative genes in each module (right). **D,** Dotplot of pseudotime modules sorted
by age. Percent of cells expressing the gene and average normalized expression are represented.
E, F, G, Average gene expression changes for each aging module represented as log2 fold
change of 12-months (**E**), 18-months (**F**), or 24-months (**G**) over 6-months. **H,** Diagram of the
heterochronic parabiosis model with the comparisons made in scRNA-Seq. **I,** Average gene
expression changes for each aging module represented as log2 fold change of heterochronic
young (HY) over isochronic young (IY) adult parabionts. Data from [31]. **J,** Diagram of microglia
surrounding an Aβ plaque. **K,** Average gene expression changes for each aging module
represented as log2 fold change of the App^{NL-G-F} genotype (AD) over wildtype (WT). Data from[17].
(one sample T-test with the expected value of 0 (no change); *P<0.05, **P<0.01, ***P<0.001,
****P<0.0001).

Fig. 3 | Intermediate states of microglia aging act as checkpoints on inflammatory progression. A, Representation of the microglia aging trajectory over the UMAP plot highlighting the region of peak *Tgfb1* expression. **B,** Representative RNAscope images and quantification of *Tgfb1* (red) expression in IBA1 (cyan) cells across ages. (n=5 per group; one-way ANOVA with Dunnett's post-hoc test; *P<0.05, **P<0.01)**. C,** Schematic of the heterochronic parabiosis model and quantification of hippocampal microglia expression of *Tgfb1* from isochronic young (IY) and heterochronic young (HY) adult parabionts. Data derived from Pálovics *et al*[31]. **D,** Top gene ontology terms of genes with significantly decreased expression in bulk microglia RNA-Seq following TGFB1 treatment compared to control (DMSO) in LPS treated microglia. (n=5 per group) **E,** Heatmap of top 10 genes in each aging module following TGFB1 compared to DMSO in LPS treated microglia. **F,** Average gene expression changes for each aging module represented as log2 fold change of TGFB1 treatment over DMSO. (one sample T-test with the expected value of 0 (no change); *P<0.05, ***P<0.001, ****P<0.0001). **G,** Representation of the microglia aging trajectory over the UMAP plot highlighting the stage where CX-5461 modulates the trajectory. **H,** Representative images of S6 (magenta) and Iba1 (cyan) staining in the hippocampus of 6- and 24-month-old mice and quantification across aging. (n=3 mice per group; one-way ANOVA with Tukey's post hoc test; *P<0.05, **P<0.005, ****P<0.0001). **I,** Schematic of the heterochronic parabiosis model and quantification of hippocampal microglia expression of translation module from isochronic young (IY) and heterochronic young (HY) adult parabionts. Data derived from Pálovics *et al*[31]. (one sample T-test with the expected value of 0 (no change); **P<0.01). **J,** Top gene ontology terms of genes with significantly decreased expression in bulk microglia RNA-Seq following CX-5461 treatment compared to control (DMSO) in LPS treated microglia. (n=3 per group) **K,** Heatmap of top 10 genes in each aging module following CX-5461 compared to DMSO in LPS treated microglia. **L,** Average gene expression changes for each aging module represented as log2 fold change of CX-5461 treatment over DMSO. (one sample T-test with the expected value of 0 (no change); *P<0.05, **P<0.01, ****P<0.0001).

Fig. 4 | Targeting age-related changes in microglia-derived TGFB1 promotes microglia advancement along inflammatory trajectories in the hippocampus and impairs cognition. **A,** Schematic of *in vivo* manipulation schema of Module 2. Adult (7-8 months) littermate *Cx3cr1-Cre-ER$^{+/wt}$; Tgfb1$^{flox/flox}$* (cKO), *Cx3cr1-Cre-ER$^{+/wt}$; Tgfb1$^{flow/wt}$* (Het), and *Cx3cr1-Cre-ER$^{+/wt}$; Tgfb1$^{wt/wt}$* (WT) mice were administered tamoxifen and subject to hippocampal microglia scRNA-Seq and behavior analysis two months later. **B,** UMAP plot of microglia separated into transcriptional clusters. (n = 2 pools of 3 animals per genotype) **C,** Stacked bar plot of the normalized relative percentage of cells of each genotype in the identified clusters. **D,** Dot plot of expression of aging module markers sorted by genotype. Percent of cells expressing the gene and average normalized expression are represented. **E,F,** Average gene expression changes for each aging module represented as log2 fold change of either Het over WT (**E**) or KO over WT (**F**). (one sample T-test with the expected value of 0 (no change); *P<0.05, ****P<0.0001). **G,** Novel object recognition task. (Top) Diagram of the training and testing phases of the novel object recognition paradigm. (Bottom) Quantification of the NOR testing phase represented as percentage of time spent with the novel object (over the total time spent interacting with the objects). (n=9-13 per genotype; one sample T-test with the expected value of 50 (equal time spent with each object); ***P<0.001)(differences between groups determined by one-way ANOVA; **P<0.01). **H,** Contextual fear conditioning. (Top) Diagram of the fear conditioning paradigm. (Bottom) Quantification of the percentage of time mice froze in the contextual fear conditioning testing phase. (n=14-16 per genotype; one-way ANOVA; *P<0.05). Data are shown as mean±s.e.m.

METHODS

Mice

All animal procedures were performed in accordance with protocols approved by the UCSF

IACUC. Animals were housed in SPF barrier facilities and provided continuous food and water

along with environmental enrichment. Aging characterizations were performed on an inhouse

C57BL/6J mouse colony where 2-month-old mice were purchased from Jackson Laboratories

(Stock 000664) and aged in the UCSF Parnassus barrier facility. Timed pregnant females were

obtained for isolation of primary microglia from newborn pups. Male mice were used in all

experiments. B6.129P2(Cg)-*Cx3cr1*[tm2.1(cre/ERT2)Litt]/WganJ (*Cx3cr1*CreER) (Stock 021160) and

C57BL/6J-Tgfb1[em2Lutzy]/Mmjax (*Tgfb1*-flox) (Stock 65809-JAX) mice were obtained from Jackson

Laboratories. Mice were bred to obtain *Cx3cr1*CreER +/wt with either *Tgfb1* wt/wt, *Tgfb1* fl/fl. As

*Cx3cr1*CreER is a knock-in/knock-out allele for the microglia homeostatic gene *Cx3cr1*, all mice

tested were *Cx3cr1*CreER +/wt. To induce recombination and deletion of *Tgfb1* by CreER, we

injected tamoxifen for 5 days at 90mg/kg daily, and mice were analyzed after 60 days.

Parabiosis

Parabiosis surgery followed previously described procedures[45]. Mirror-image skin incisions at the

left and right flanks were made through the skin, and shorter incisions were made through the

peritoneum. The peritoneal openings of the adjacent parabionts were sutured together with

chromic gut suture (MYCO Medical, GC635-BRC). Apposing elbow and knee joints from each

parabiont were sutured together (Coated VICRYL Suture, Ethicon, J386) and the skin of each

mouse was stapled (9-mm Autoclip, Clay Adams, 427631) to the skin of the adjacent parabiont.

Each mouse was injected subcutaneously with Carpofen, Enrofloxacin, and Buprenex as directed

for pain and monitored during recovery. For overall health and maintenance behavior, several

recovery characteristics were analyzed at various times after surgery, including pair weights and

grooming behavior.

Bulk Microglia RNA-Seq

Microglia were isolated from C57BL6/J, *Tgfb1* cHet, and *Tgfb1* cKO using FACS. Briefly, mice

were sedated with ketamine followed by perfusion with 30mL of ice-cold PBS. The entire brain

was removed, then the hippocampus was sub-dissected. Single-cell suspensions were generated

by enzyme-mediated (papain) and mechanical dissociation using Miltenyi Neural Dissociation Kits

(P) (Miltenyi, 130-092-628) according to the manufacturer's instructions. Myelin was depleted

from the suspensions using Myelin Removal Beads (Miltenyi, 130-096-731). Cells were labeled

with CD11b-APC (eBioscience, 17-0112-82, RRID:AB_469343) and CD45-PE (eBioscience, 12-

0451-82, RRID:AB_465668), then sorted at 4°C into Tri Reagent with a FACS AriaII (BD). The

gating strategy for microglia isolation is presented in Figure S1D, showing the collection of

Cd11b$^+$Cd45^{Intermediate} cells. During the collection of control and *Tgfb1* microglia, CD48 was

detected in Cd11b$^+$Cd45^{Intermediate} cells using CD48-Pacific Blue (BioLegend, 103417,

RRID:AB_756139). FlowJo was used to analyze flow cytometry data. RNA was isolated from

microglia using Tri Reagent (Sigma-Aldrich, T9424). RNA pellets were resuspended in 5uL TE

buffer.

RNA was transformed into RNA-Seq libraries using an adapted version of the Smart-Seq2

protocol[46]. Briefly, 5ng of total RNA was reverse transcribed with SuperScript II (Thermo Fisher,

18064014) supplemented with betaine and MgCl$_2$ using an oligo-dT RT primer

(AAGCAGTGGTATCAACGCAGAGTACT(30)VN) with a PCR binding site and a template

switching oligonucleotide (5′ AAGCAGTGGTATCAACGCAGAGTACATrGrG+G) with a

homotypic PCR binding site. Betaine and MgCl2 were added to enhance the reaction. After

reverse transcription, whole transcriptome amplification by PCR was performed for 12 cycles

using KAPA HiFi HotStart ReadyMix (Roche, 7958935001) with a PCR primer

(AAGCAGTGGTATCAACGCAGAGT). The PCR reaction was cleaned up with Ampure XP beads

(Beckman Coulter, A63881). In addition, after the whole transcriptome PCR amplification step,

qPCR was performed to determine the presence of microglia-specific transcripts (*Cx3cr1*) and

housekeeping genes (*Gapdh*). The amplified DNA was diluted to a concentration of 0.5ng/ul, and

subjected to tagmentation with the Illumina Nextera XT kit (Illumina, FC-131-1096). Each sample

was PCR amplified with a unique set of Nextera indices. Bulk microglia RNA-Seq libraries were

sequenced on an Illumina HiSeq, while *in vitro* microglia RNA-Seq libraries were sequenced on

an Illumina Nova-Seq.

RNA-Seq Analysis

FASTQ reads were pseudo aligned to the mouse transcriptome (GRCm39 cDNA from Ensembl)

using kallisto[47] with default parameters. Next, transcript abundance estimates were imported into

DeSeq2[48]. Differential expression analysis was performed using DeSeq2 using the Wald

significance test. Gene ontology analysis was performed with Panther[49-51]. Principal component

analysis plots were generated with the plotPCA function included with DeSeq2 and graphed with

ggplot2. Heatmaps were generated with the pheatmap package in R.

The overlap between gene expression differences in aging and LPS treatment were

determined using genes significantly changed between the 6- and 24-month-old timepoints in the

scRNA-Seq data and the control and LPS treated primary microglia. χ^2 test was used to determine

the significance of the overlap between these two datasets. The Venn diagram was generated

using the VennDiagram package in R.

10x Genomics Single Cell RNA-Sequencing

10x Single-Cell RNA-Seq libraries were generated from Cd11b+ cells isolated from the

hippocampi of an aging cohort of mice consisting of 6-, 12-, 18-, and 24-month-old C57BL/6J mice

that were all collected and processed on the same day in an interspersed order. Mice were

sedated with ketamine followed by perfusion with ice-cold PBS. Subsequently, the entire brain

was removed from the mouse followed by sub-dissection of the hippocampus. For each age,

hippocampi from five mice were pooled during the dissection step in HBSS at 4°C. Single-cell

suspensions were generated by enzyme-mediated (papain) and mechanical dissociation using

Miltenyi Neural Dissociation Kits (P). The papain dissociation was done at 37°C for 10 minutes

with 3 trituration steps. All other processing steps were performed at 4°C. Myelin was depleted

from the suspensions using Myelin Removal Beads (Miltenyi). Microglia were enriched using

Cd11b magnetic beads (Miltenyi, 130-126-725). Cell viability was determined to be over 95%.

Single-cell RNA-Seq 3' libraries were generated from the cell suspension using 10x Genomics

Chromium Single-Cell 3' Solution. Libraries were sequenced on the Illumina Nova-Seq. Cell

Ranger demultiplexed and mapped (using bcl2fastq) reads, followed by alignment (with STAR),

and generation of single-cell expression matrices[52].

The isolation protocol for the *Tgfb1* genetic mouse model was modified. We added

transcriptional and translational inhibitors (Actinomycin D (Sigma-Aldrich, A1410), Anisomycin

(Sigma-Aldrich, A9789), and Triptolide (Sigma-Aldrich, T3652) to prevent transcriptional changes

due to *ex vivo* activation [53]. We utilized the concentrations presented in Marsh *et al*. 2022[53].

Single-cell RNA-Seq Analysis

Seurat: The single-cell gene expression matrices generated by Cell Ranger were loaded into

Seurat[54].

Aging microglia dataset. Cells were filtered according to the following parameters: Genes > 1000

& RNA count > 10000 & mitochondrial percentage < 5. The count data was log normalized. The

2,500 most variable genes were identified using the "vst" method. The expression data was scaled

and centered. Principal component analysis was performed, and the first 25 PCAs were used to

identify nearest neighbors, and interconnected clusters were identified with a resolution of 0.4.

UMAP was used to visualize the results. Preliminary dimensionality reduction and visualization

with UMAP using 25 PCAs and 25 nearest neighbors in Seurat identified a large microglia cluster,

plus sparse clusters consisting of astrocytes, vascular cells, neutrophils, and macrophages. *Hexb*,

P2ry12, *Tmem119*, and *Slc2a5* are used as markers to identify microglia in the preliminary

clustering, while *Nrxn1* (neurons), *Tm4sf1* (vasculature), *Dclk*1 (astrocytes), *Sdpr* (vasculature

smooth muscle), *Abcb1a* (pericyte/vasculature), and *Foxq1* (macrophages) identified other cell

types. The data was then subsetted on the non-proliferating microglial clusters (to avoid

confounds associated with the cell cycle). The top 1,500 most variable genes in the subsetted

microglia were identified using the "vst" method. The data was scaled and centered. Principal

component analysis was performed, and the first 8 PCAs were used to identify nearest neighbors,

and interconnected clusters were identified with a resolution of 0.25. UMAP was used to visualize

the results using 8 PCAs and 100 nearest neighbors. Gene enrichment and differential expression

analysis was performed on the filtered dataset to determine differences between ages.

Tgfb1 cKO microglia dataset. Cells were prefiltered according to the following parameters: Hexb

> 4 & Tmem119 > 1 & Cx3cr1 > 4 & Top2a < 1 & Mki67 < 1 nFeature_RNA > 1000 &

nFeature_RNA < 6000 & nCount_RNA > 2000 & nCount_RNA < 40000 & percent.mt <10. Data

was integrated using IntegrateData and scaled. Principal component analysis was performed,

and the first 50 PCAs were used to identify nearest neighbors, and interconnected clusters were

identified with a resolution of 0.35. UMAP was used to visualize the results. Preliminary

dimensionality reduction and visualization with UMAP using 50 PCAs identified a large microglia

cluster, plus clusters of other cell types. The non-proliferative microglia clusters were subsetted

for further analysis (the macrophage cluster was differentiated from the microglia cluster based

on increased expression of *Mcp1, Pf4,* and *Ms4a7*). The data was scaled and centered. Principal

component analysis was performed, and the first 10 PCAs were used for UMAP dimensionality

reduction and to identify nearest neighbors, and interconnected clusters were identified with a

resolution of 0.20.

Monocle 3: Monocle 3 was used to generate the pseudotime analysis[55-58]. Data was imported

from Seurat analysis and a Monocle dataset was generated. The dataset was preprocessed with

normalization of the data and PCA generation. Next, the preprocessed data underwent further

non-linear dimensionality reduction using UMAP. Cells were clustered, and trajectories were

determined for cells within a cluster. For determining "pseudotime" aging of microglia, the root

node of the pseudotime trajectory was manually placed in the cluster of 6-month cells. The graph

segments of the pseudotime trajectories resulting in inflammatory activation were chosen for

further analysis. Graph autocorrelation analysis was performed on the inflammatory trajectory

using Moran's I, which determines if genes are expressed in focal regions of graph space. A cutoff

q-value of 0.005 was set as significant for focal expression. Subsequently, coregulated modules

were determined using Louvain community analysis at an optimized resolution of 0.0078. The

modules were then overlaid onto the UMAP plot to determine the region of the trajectory with focal

expression of the module. The ten most significant genes of each module, as determined by q-

value were used to construct the heatmaps in Figures 3F and 3I. The significant genes of each

module were used to determine the effects of manipulations in the presence of LPS in Figure 3J.

Immunohistochemistry

Mice were sedated with ketamine followed by perfusion with 30mL of ice-cold PBS. Brains were

collected from PBS perfused mice and fixed overnight at 4°C in 4% paraformaldehyde in PBS.

The brains were washed with PBS and immersed in 30% sucrose (in PBS), and then stored at

4°C until the brains sank. 40μM coronal sections were sliced using a microtome at -20°C. Sections

were stored in cryopreservation buffer (40% PB Buffer (0.02 M Sodium Phosphate Monobasic,

0.08 M Sodium Phosphate Dibasic, pH 7.4), 30% Glycerol, 30% Ethylene Glycol) at -20°C until

staining. Hippocampal sections were washed three times in TBST. Sections were permeabilized

in pretreatment buffer (0.1% Triton-X in TBST) with rocking for 1 hour at room temperature. After

three washes in TBST, sections were blocked in 5% goat serum for 2 hours (for IBA1/CD68

double stain) or 5% donkey serum (NFKB/Iba1 and C1q/C3/Iba1 stains). Blocking buffer was

replaced with primary antibody mixture and incubated overnight at 4°C with rocking. The following

primary antibodies were used: rabbit anti-IBA1 (1:1000, Wako, Cat# 019-19741,

RRID:AB_839504), rat anti-CD68 (1:250, Bio-Rad, Cat# MCA1957, RRID:AB_322219), rabbit

anti-NFKB p65 (1:500, Santa Cruz, Cat# sc-372, RRID:AB_632037), guinea pig anti-Iba1 (1:1000,

Synaptic Systems, Cat# 234-004, RRID:AB_2493179), rabbit anti-C1q (1:1000, Abcam, Cat#

ab182451, RRID:AB_2732849), rat anti-C3 (1:1000, Abcam, Cat# ab11862, RRID:AB_2066623),

and rabbit anti-S6 (1:500, Cell Signaling, Cat# 2217, RRID:AB_331355). The sections were

washed three times with TBST. Subsequently, secondary antibody solution was added to the

sections, and incubated at room temperature with rocking for two hours. The following secondary

antibodies were used: goat anti-rabbit AlexaFluor 488 (Life Technologies, Cat# A-11008,

RRID:AB_143165), goat anti-rat AlexaFluor 555 (Life Technologies, Cat# A-21434,

RRID:AB_2535855), donkey anti-guinea pig AlexaFluor 488 (Jackson Immuno, Cat# 706-545-
148, RRID:AB_2340472), donkey anti-rabbit AlexaFluor 555 (Life Technologies, Cat# A-31572,
RRID:AB_162543), and donkey anti-rat AlexaFluor 647 (Jackson Immuno, Cat# 712-606-150,
RRID:AB_2340695). The sections were washed three times with TBST, with the first wash
containing Hoescht 33342 (Invitrogen, H3570) at a concentration of 1μg/mL. The sections were
transferred to phosphate buffer and mounted on frosted slides. Coverslips were mounted with
Prolong Gold Antifade (Thermo Fisher, P10144) reagent after the sections were dried. Slides
were imaged at 20x magnification on a Zeiss LSM 800 or LSM900 for a final resolution of
0.312μM/pixel. 8-9 z-planes were imaged at 3μM intervals. The hippocampus was imaged, and
the tiled images were stitched together with Zeiss ZenPro software. 2-3 hippocampal sections
were imaged from each sample.

Images were analyzed with FIJI. Maximum intensity projections of Z-stacked images were
constructed. Images were converted to 8-bit images and consistent adjustments were made for
each channel within an experiment. The Iba1 channel was used to quantify microglia using the
particle counter function in FIJI and the counts were validated by manual visual counting. The
presence of CD68, NFKB(p65), C3, or C1q in microglia was determined by creating ROIs within
the microglia in a field using the create selection function on the Iba1 images. The ROIs of the
hippocampal subregions and the microglia were combined, and CD68 puncta meeting a minimum
particle size threshold (100 pixels, equating to 31.2μM) within microglia determined if the microglia
were activated (CD68+IBA1+), while the measure function determined the intensity of NFKB(p65),
C3, or C1q. To determine percent activation of microglia with IBA1/CD68 staining, we divided the
number of microglia that were activated (CD68+IBA1+) by the total number of microglia (IBA1+)
in the analyzed regions and multiplied by 100.

***In vitro* treatment of microglia**
Mixed glial cultures were generated from p1 male pups. Cortices and hippocampi were isolated
from the pups, and meninges were removed. The tissue was broken up by repeatedly pipetting
up and down, followed by dissociation with Trypsin. Single cell suspensions were plated on poly-
L-lysine coated flasks in DMEM (Thermo Fisher, 11965-118) supplemented with 10% FBS
(Gemini Bio-Products, 900-208) and antibiotic mix (Penicillin/Streptomycin)(Genesee Scientific,
25-512) and incubated at 37°C in 5% CO_2. Media was changed after 4 days to remove debris.
Microglia were isolated after 14 days by agitating the plates to dislodge microglia, then transferring
the supernatant of each sample evenly amongst 4 wells of a 24 well plate. Microglia were

incubated for 16 hours followed by replacement with serum free macrophage SFM media (Thermo

Fisher, 12065074) supplemented with M-CSF (10ng/m) (Peprotech, 315-02) and antibiotic mix.

After 24 hours microglia were treated with RNA polymerase I Inhibitor 2, CX-5461 (100nM)(EMD

Millipore, 509265) or human TGFβ1 recombinant protein (10ng/mL)(ThermoFisher Scientific,

PHG9204) for 24 hours followed by addition of lipopolysaccharides (LPS, 200ng/uL) from *E. coli*

(Sigma-Aldrich, L2018) for 8 hours and ATP (10nM)(InvivoGen, tlrl-atpl) for 30 minutes. Tri

Reagent was added to the culture plate, and the plates were shaken for 5 minutes. RNA was

isolated according to manufacturer's instructions. RNA-Seq libraries were generated as above for

the bulk RNA-Seq.

RNAscope

RNAscope was performed with the RNAscope® Multiplex Fluorescent Reagent Kit v2 (ACD,

323110). 40μM coronal sections were washed with TBST, then incubated with hydrogen peroxide

for 45 minutes at 25°C. After washing with TBST, the sections were incubated in Target Retrieval

Buffer at 95°C for 10 minutes. The sections were mounted on slides and dried overnight. The

sections were treated with RNAscope Protease III for 10 minutes at 40°C. Sections were washed

in water four times. Sections were incubated with *Tgfb1* probe (ACD, 443571-C2) for 2 hours at

40°C. The sections were washed twice with RNAscope wash buffer. The sections were incubated

with AMP1 for 30 minutes, AMP2 for 30 minutes, AMP3 for 15 minutes, HRP-C2 for 15 minutes,

TSA Plus Cy3 (1:1000, Perkin Elmer, SKU NEL744001KT) for 30 minutes, and HRP-Blocker for

15 minutes; all at 40°C with washes with RNAscope wash buffer between every incubation. RNA-

Protein Co-Detection Ancillary Kit (ACD, 323180) buffer was used to block the sections before

incubating with rabbit anti-IBA1 (1:500, Wako, Cat# 019-19741, RRID:AB_839504) overnight at

4°C. Sections were washed three times in TBST. Sections were incubated with secondary

antibody solution with goat anti-rabbit AlexaFluor 488 (1:1000, Life Technologies, Cat# A-11008,

RRID:AB_143165) for two hours at room temperature. The sections were washed three times

with TBST, with the first wash containing Hoescht 33342 (Invitrogen, H3570) at a concentration

of 1μg/mL Coverslips were mounted with Prolong Gold Antifade (Thermo Fisher, P10144) reagent

after the sections were dried. Slides were imaged at 40x magnification on a Zeiss LSM 900. 10

z-planes were imaged at 1μM intervals.

Images were analyzed with FIJI. Maximum intensity projections of Z-stacked images were

constructed. Images were converted to 8-bit images and the *Tgfb1* was thresholded. For each

section, the intensity of 5-7 individual *Tgfb1* puncta were averaged to get the average puncta

intensity of that section. The intensity of all *Tgfb1* signal in clearly identifiable individual IBA1 cells

was measured for 20-25 cells in the molecular layer of the hippocampus. The mean *Tgfb1* count

per microglia for each sample was calculated by dividing the average *Tgfb1* intensity in IBA1 cells

by the average puncta intensity.

Contextual fear conditioning

In this task, mice learned to associate the environmental context (fear conditioning chamber) with

an aversive stimulus (mild foot shock; unconditioned stimulus, US) during the training phase

enabling testing for hippocampal- dependent contextual fear conditioning. To also assess

amygdala-dependent cued fear conditioning, the mild foot shock was paired with a light and tone

cue (conditioned stimulus, CS) during the training phase. Conditioned fear was displayed as

freezing behavior. Specific training parameters are as follows: tone duration is 30 seconds; level

is 70 dB, 2 kHz; shock duration is 2 seconds; intensity is 0.6 mA. This intensity is not painful and

can easily be tolerated but will generate an unpleasant feeling. More specifically, on day 1 each

mouse was placed in a fear-conditioning chamber and allowed to explore for 2 min before delivery

of a 30-second tone (70 dB) and light ending with a 2-second foot shock (0.6 mA). Two minutes

later, a second CS-US pair was delivered. On day 2, each mouse was first placed in the fear-

conditioning chamber containing the same exact context, but with no CS or foot shock. Freezing

was analyzed for 2 minutes and represented as the percentage of time that the mouse froze over

the 2 minutes. One hour later, the mice were placed in a new context containing a different odor,

floor texture, chamber walls and shape. Animals were allowed to explore for 2 minutes before

being re-exposed to the CS. Freezing was analyzed for 30 seconds following the CS and

represented as the percentage of time that the mouse froze over those 30 seconds. Determination

of freezing behavior was performed using FreezeScan video tracking system and software

(Cleversys, Inc). Single outliers were removed using the extreme studentized deviate method

(Grubbs' test) with an alpha of 0.05.

Novel object recognition

The novel object recognition task was adapted from a previously described protocol[59].

Specifically, during the habituation phase (day 1), mice could freely explore an empty open field

arena for 10 minutes (motor activity and anxiety (time in center) were measured during this

phase). During the training phase (day 2), two identical objects were placed in the habituated

arena, and mice could explore the objects for 5 minutes. For the testing phase (day 3), one object

was replaced with a novel object, and mice could explore the objects for 5 minutes. Time spent
exploring each object was quantified using the Smart Video Tracking Software (Panlab; Harvard
Apparatus). Two different sets of objects are used. To control for any inherent object preference,
half of the mice are exposed to object A as their novel object and half to object B. To control for
any potential object-independent location preference, the location of the novel object relative to
the trained object is also varied. The objects were chosen based on their ability to capture the
animal's interest, independent of genetic background or age. To determine percent time with
novel object, we calculate (Time with novel object)/(Time with Trained Object + Time with Novel
Object) * 100. In this preference index, 100% indicates full preference for the novel object, and
0% indicates full preference for the trained object. A mouse with a value of 50% would have spent
equal time exploring both objects. Mice that did not explore both objects for 5 seconds during the
training phase or testing phase were excluded from analysis. Single outliers were removed using
the extreme studentized deviate method (Grubbs' test) with an alpha of 0.05.

Y maze

The Y Maze task was conducted using an established forced alternation protocol[60]. During the
training phase, mice were placed in the start arm facing the wall and allowed to explore the start
and trained arm for 5 minutes, while entry to the 3rd arm (novel arm) was blocked. The maze was
cleaned between each mouse to remove odor cues, and the trained arm was alternated between
mice. The mouse was then removed to its home cage. After 30 minutes, the block was removed,
and the mouse was returned to the start arm and allowed to explore all 3 arms for 5 minutes. Time
spent in each arm was quantified using the Smart Video Tracking Software (Panlab; Harvard
Apparatus). The percent time in the novel arm was defined as time in the novel arm divided by
time spent in the novel and trained arms during the task.

Datasets

All RNA-Seq and scRNA-Seq data have been deposited in the Gene Expression Omnibus and
are publicly available as of the date of publication. The following datasets were generated for this
manuscript and deposited in Gene Expression Omnibus: aging single-cell RNA-Seq
(GSE179358), *in vitro* treated primary microglia (GSE179611), and *Tgfb1* cKO microglia RNA-
Seq (GSE190007).

Statistics

Researchers were blinded throughout histological assessments with groups being un-blinded at
the end of each experiment upon statistical analysis. Data are expressed as mean ± s.e.m with
individual sample values being shown. Statistical analysis was performed with Prism 8.0
(GraphPad), R, DESeq2, Seurat, or Monocle 3. Comparisons of means in histology experiments
were analyzed with multiple T-tests followed by Holm-Sidak correction or mixed effects analysis
followed by Dunnett's multiple comparisons (Prism). Changes in expression for aggregated gene
sets were determined with one sample T-tests with the expected value of 0 in log2 space (null
hypothesis of no change). Differential expression analysis in DESeq2 was based on Wald's
significance test of the negative binomial distribution. Differential expression analysis of single-
cell RNA-Seq data in Seurat was performed using non-parametric Wilcoxon rank sum tests.
Autocorrelation analysis to determine focal expression Monocle 3 was done using Moran's I. The
significance of gene set overlap was determined using the χ^2 test with R. All data generated or
analyzed in this study are included in this article.

SUPPLEMENTAL MATERIAL

Fig. S1 | Hippocampal microglia exhibit age-related heterogeneity during aging. A, UMAP plot of all Cd11b+ cells. Several adjacent clusters of microglia were identified, as well as a cluster of proliferating microglia. Smaller populations of peripheral immune cell types – macrophages and neutrophils – were identified. Clusters of astrocytes and vascular cells were also found. Overall, greater than 82% of cells were microglia. (n = 1 pool of 5 animals for each age) **B,** Dot plot showing expression (average expression and percent of cells expressing) of top two markers for each cluster. **C,** UMAP plots with expression levels of microglia markers superimposed onto cells. Notice that peripheral immune cells express microglia markers; however, they are distinguished from microglia based on marker expression from (b). **D-G,** Volcano plots of differential gene expression for the clusters identified in Figure 1A compared to every other cluster for Homeostatic (**D**), Transition (**E**), Activation (**F**), and Interferon (**G**) microglia clusters. **H,** Cluster composition of non-proliferating microglia by age. **I,** Standardized variation of non-proliferating microglia for each age. **J,** Representative images of NFκB (p65) (yellow) and IBA1 (cyan) staining in the hippocampus of 3- and 24-month-old mice and quantification of staining in 3-, 6-, 12-, 18-, and 24-month-old mice. (n=5 mice per group; one-way ANOVA with Dunnett's post hoc test; *$P<0.05$, ***$P<0.001$)**. K,** Representative images of CD68 (red) and IBA1 (cyan) staining in the hippocampus of 3- and 24-month-old mice and quantification of staining in 3-, 6-, 12-, 18-, and 24-month-old mice. (n=4-5 mice per group; one-way ANOVA with Dunnett's post hoc test; *$P<0.05$, **$P<0.01$)**. Data are shown as mean±s.e.m.

Figure S2 | Hippocampal microglia aging advances through intermediate states that
respond to systemic interventions or disease states. **A**, Volcano plot of differential gene
expression of 12-month microglia versus all other ages with significant genes in teal. **B**, Gene
ontology analysis of biological processes enriched in those genes with increased expression at
12-months of age. **C**, Representative images and quantification of CD68 (red) and IBA1 (cyan)
staining in the hippocampus of isochronic young (IY) and heterochronic young (HY) along with a
diagram of the comparisons. (n=5 mice per group; unpaired Student's T-test; ***$P<0.001$)**.** Data
are shown as mean±s.e.m. **D,** Dotplot of pseudotime modules in young (Y), isochronic young (IY),
and heterochronic young (HY) parabiont microglia. Data is from Palovics *et al*[61]. Percent of cells
expressing the gene and average normalized expression are represented. **E,** Volcano plots of
differential gene expression of App^{NL-G-F} genotype (AD) over wildtype (WT) microglia at 6- (left)
and 12-months of age. Significant genes are in teal. Data is from Frigerio[62]. **F,** Dotplot of
pseudotime modules across ages (3-, 6-, 12-, and 21-months old) and genotypes (App^{NL-G-F}
genotype (AD) over C57Bl/6 (WT)). Data is from Frigerio[17]. Percent of cells expressing the gene
and average normalized expression are represented.

Fig. S3 | Intermediate states of microglia aging act as checkpoints on inflammatory progression. **A,** Quantification of percentage of *Tgfb1* RNAscope signal within IBA1 cells. **B**, Quantification of IHC of pTGFBR1 signal in IBA1 cells across ages. (n=4-5 mice per group; one-way ANOVA; *P<0.05) **C,** Dotplot of the expression values of TGFB1 signaling components from scRNA-Seq of aging hippocampal microglia (6-, 12-, 18-, and 24-month-old). Percent of cells expressing the gene and average normalized expression are represented. **D**, Dotplot of the expression values of TGFB1 signaling components in young (Y), aged (A), isochronic young (IY), and heterochronic young (HY) parabiont microglia. Data is from Palovics *et al*[61]. Percent of cells expressing the gene and average normalized expression are represented. UMAP plot of *Tgfb1* scRNA-Seq colored by genotype. **E**, Heatmap of gene expression changes in the top 10 genes in each aging module induced by LPS. **F**, Overlap between microglia gene expression changes induced by LPS and aging. (χ^2<2.2e-16). **G**, PCA plot of pharmacological manipulations with or without LPS treatment.

Fig. S4 | Targeting age-related changes in microglia-derived TGFB1 promotes microglia advancement along inflammatory trajectories in the hippocampus and impairs cognition.
A, Representative images and quantification of IBA1 (cyan)/CD68 (red)-positive microglia in wildtype, *Tgfb1* cHet, and *Tgfb1* cKO hippocampi. (n=3-5; one way ANOVA with Tukey post hoc test; *P<0.05) **B,** Dotplot of the expression values of TGFB1 signaling components from scRNA-Seq of *Tgfb1* WT, cHet, and cKO microglia. (n = 2 pools of 3 animals per genotype) **C,** UMAP plot of *Tgfb1* scRNA-Seq colored by genotype. **D,** FACS plots of gating strategy for microglia isolation for RNA-Seq in cHet and cKO hippocampi. Quantification of the flow cytometry analysis of CD11b and CD45. (n=3-5; mixed effects analysis; *P<0.05, ****P<0.0001). **E**, Example histogram and quantification of CD48 in control and *Tgfb1* cKO FACS-sorted microglia. (n=3-5 per group; T-test; ****P<0.0001). **F,** Top gene ontology terms associated with genes significantly increased in microglia bulk RNA-Seq in *Tgfb1* cKO samples (n=3-5 samples per genotype). **G**, Top gene ontology terms associated with genes significantly decreased in microglia bulk RNA-Seq in *Tgfb1* cKO samples. **H**, Overlap between concordant microglia gene expression changes induced by *Tgfb1* knockout and aging. (χ^2<2.2e-16). **I**, Heatmap of gene expression of the top 10 genes in each aging module in control and *Tgfb1* cKO microglia. **J**, Average gene expression changes for each aging module represented as log2 fold change of *Tgfb1* cKO over control. (one sample T-test with the expected value of 0 (no change); ****P<0.0001). **K**, Freezing during the cued fear conditioning phase of the fear conditioning test. The percent freezing was calculated for the last 30 seconds of the test following the conditioned stimulus. (n=13-16 per genotype) **L**, Diagram and results of Y-maze test. Results are presented as preference for the novel arm versus the trained armed (time in novel arm/ (time in novel arm + trained arm) x 100). (n=14-16 per genotype) **M**, Percent time in the center of the open field and distance traveled during the open field test. (n=14-16 per genotype) Data are shown as mean±s.e.m.

Table S1 | Gene markers for each age group and differential expression between the 6- and
12-, 18-, or 24-month-old microglia in single-cell RNA-Seq of aging hippocampal microglia.
The tables contain genes selectively increased or decreased in each age group when compared
to every other age group combined and genes differentially expressed between 6- and 12-, 18-,
or 24-month microglia. The tables contain the average log fold change, the percentage of cells in
each group with expression, and the adjusted p-value.

1133 **Table S2 | Pseudotime trajectories of aging microglia.** The pseudotime analysis contains the
1134 spatial autocorrelation analysis. Moran's I was used to detect focal expression of genes that were
1135 consequently constructed into co-regulated modules using Louvain community analysis. The
1136 table contains the adjusted p-value (q-value) and module information.

Table S3 | RNA-Seq analysis of primary microglia treated with LPS and TGFB1 or CX-5461.
The tables contain the RNA-Seq differential expression analysis of control and LPS activated
primary microglia, DMSO and CX-5461 treated primary microglia activated by LPS, and DMSO
and TGFB1 treated primary microglia activated by LPS, respectively. The tables contain
expression values of genes for the samples that were compared along with mean expression
values for all samples combined, log fold change between sample groups, and adjusted p-values.
in the RNA-Seq differential expression analysis of control and LPS activated primary microglia,
DMSO and CX-5461 treated primary microglia activated by LPS, and DMSO and TGFB1 treated
primary microglia activated by LPS, respectively.

 Table S4 | Differential expression analysis of control and *Tgfb1* cKO hippocampal
microglia. The table contains expression values of genes for the samples that were compared
along with mean expression values for all samples combined, log fold change between sample
groups, and adjusted p-values.

REFERENCES

Waisman, A., Ginhoux, F., Greter, M. & Bruttger, J. Homeostasis of Microglia in the Adult Brain: Review of Novel Microglia Depletion Systems. *Trends in immunology* **36**, 625-636, doi:10.1016/j.it.2015.08.005 (2015).

Salter, M. W. & Beggs, S. Sublime microglia: expanding roles for the guardians of the CNS. *Cell* **158**, 15-24, doi:10.1016/j.cell.2014.06.008 (2014).

Mosher, K. I. & Wyss-Coray, T. Microglial dysfunction in brain aging and Alzheimer's disease. *Biochem Pharmacol* **88**, 594-604, doi:10.1016/j.bcp.2014.01.008 (2014).

Bachiller, S. *et al.* Microglia in Neurological Diseases: A Road Map to Brain-Disease Dependent-Inflammatory Response. *Front Cell Neurosci* **12**, 488, doi:10.3389/fncel.2018.00488 (2018).

Sierra, A., Gottfried-Blackmore, A. C., McEwen, B. S. & Bulloch, K. Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia* **55**, 412-424, doi:10.1002/glia.20468 (2007).

Marschallinger, J. *et al.* Lipid-droplet-accumulating microglia represent a dysfunctional and proinflammatory state in the aging brain. *Nature neuroscience* **23**, 194-208, doi:10.1038/s41593-019-0566-1 (2020).

Pluvinage, J. V. *et al.* CD22 blockade restores homeostatic microglial phagocytosis in ageing brains. *Nature*, doi:10.1038/s41586-019-1088-4 (2019).

Damani, M. R. *et al.* Age-related alterations in the dynamic behavior of microglia. *Aging Cell* **10**, 263-276, doi:10.1111/j.1474-9726.2010.00660.x (2011).

Flanary, B. E., Sammons, N. W., Nguyen, C., Walker, D. & Streit, W. J. Evidence that aging and amyloid promote microglial cell senescence. *Rejuvenation Res* **10**, 61-74, doi:10.1089/rej.2006.9096 (2007).

Udeochu, J. C., Shea, J. M. & Villeda, S. A. Microglia communication: Parallels between aging and Alzheimer's disease. *Clin Exp Neuroimmunol* **7**, 114-125, doi:10.1111/cen3.12307 (2016).

Grabert, K. *et al.* Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nature neuroscience* **19**, 504-516, doi:10.1038/nn.4222 (2016).

Burns, J. C. *et al.* Differential accumulation of storage bodies with aging defines discrete subsets of microglia in the healthy brain. *Elife* **9**, doi:10.7554/eLife.57495 (2020).

Hammond, T. R. *et al.* Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity* **50**, 253-271 e256, doi:10.1016/j.immuni.2018.11.004 (2019).

Li, X. *et al.* Transcriptional and epigenetic decoding of the microglial aging process. *Nat Aging* **3**, 1288-1311, doi:10.1038/s43587-023-00479-x (2023).

15 Silvin, A. *et al.* Dual ontogeny of disease-associated microglia and disease inflammatory
macrophages in aging and neurodegeneration. *Immunity* **55**, 1448-1465 e1446,
doi:10.1016/j.immuni.2022.07.004 (2022).

16 Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting Development
of Alzheimer's Disease. *Cell* **169**, 1276-1290 e1217, doi:10.1016/j.cell.2017.05.018
(2017).

17 Sala Frigerio, C. *et al.* The Major Risk Factors for Alzheimer's Disease: Age, Sex, and
Genes Modulate the Microglia Response to Abeta Plaques. *Cell Rep* **27**, 1293-1306
e1296, doi:10.1016/j.celrep.2019.03.099 (2019).

18 Lee, S. *et al.* APOE modulates microglial immunometabolism in response to age, amyloid
pathology, and inflammatory challenge. *Cell Rep* **42**, 112196,
doi:10.1016/j.celrep.2023.112196 (2023).

19 Masuda, T. *et al.* Spatial and temporal heterogeneity of mouse and human microglia at
single-cell resolution. *Nature* **566**, 388-392, doi:10.1038/s41586-019-0924-x (2019).

20 Kaya, T. *et al.* CD8(+) T cells induce interferon-responsive oligodendrocytes and microglia
in white matter aging. *Nature neuroscience* **25**, 1446-1457, doi:10.1038/s41593-022-
01183-6 (2022).

21 Yousef, H. *et al.* Aged blood impairs hippocampal neural precursor activity and activates
microglia via brain endothelial cell VCAM1. *Nat Med* **25**, 988-1000, doi:10.1038/s41591-
019-0440-4 (2019).

22 Rebo, J. *et al.* A single heterochronic blood exchange reveals rapid inhibition of multiple
tissues by old blood. *Nat Commun* **7**, 13363, doi:10.1038/ncomms13363 (2016).

23 Villeda, S. A. *et al.* The ageing systemic milieu negatively regulates neurogenesis and
cognitive function. *Nature* **477**, 90-94, doi:10.1038/nature10357 (2011).

24 Li, Q. *et al.* Developmental Heterogeneity of Microglia and Brain Myeloid Cells Revealed
by Deep Single-Cell RNA Sequencing. *Neuron* **101**, 207-223 e210,
doi:10.1016/j.neuron.2018.12.006 (2019).

25 Kracht, L. *et al.* Human fetal microglia acquire homeostatic immune-sensing properties
early in development. *Science (New York, N.Y.)* **369**, 530-537,
doi:10.1126/science.aba5906 (2020).

26 Fan, X., Wheatley, E. G. & Villeda, S. A. Mechanisms of Hippocampal Aging and the
Potential for Rejuvenation. *Annu Rev Neurosci* **40**, 251-272, doi:10.1146/annurev-neuro-
072116-031357 (2017).

27 Millet, A., Ledo, J. H. & Tavazoie, S. F. An exhausted-like microglial population
accumulates in aged and APOE4 genotype Alzheimer's brains. *Immunity* **57**, 153-170
e156, doi:10.1016/j.immuni.2023.12.001 (2024).

28   Harrington, E. P. *et al.* MHC class I and MHC class II reporter mice enable analysis of
immune oligodendroglia in mouse models of multiple sclerosis. *Elife* **12**,
doi:10.7554/eLife.82938 (2023).

29   Shi, Q. *et al.* Complement C3-Deficient Mice Fail to Display Age-Related Hippocampal
Decline. *J Neurosci* **35**, 13029-13042, doi:10.1523/jneurosci.1698-15.2015 (2015).

30   Butovsky, O. *et al.* Identification of a unique TGF-beta-dependent molecular and functional
signature in microglia. *Nature neuroscience* **17**, 131-143, doi:10.1038/nn.3599 (2014).

31   Pálovics, R. *et al.* Molecular hallmarks of heterochronic parabiosis at single-cell resolution.
Nature **603**, 309-314, doi:10.1038/s41586-022-04461-2 (2022).

32   Yi, Y. W., You, K. S., Park, J. S., Lee, S. G. & Seong, Y. S. Ribosomal Protein S6: A
Potential Therapeutic Target against Cancer? *Int J Mol Sci* **23**, doi:10.3390/ijms23010048
(2021).

33   Spittau, B., Dokalis, N. & Prinz, M. The Role of TGFbeta Signaling in Microglia Maturation
and Activation. *Trends in immunology* **41**, 836-848, doi:10.1016/j.it.2020.07.003 (2020).

34   Cserép, C., Pósfai, B. & Dénes, Á. Shaping Neuronal Fate: Functional Heterogeneity of
Direct Microglia-Neuron Interactions. *Neuron* **109**, 222-240,
doi:10.1016/j.neuron.2020.11.007 (2021).

35   Horowitz, A. M. *et al.* Blood factors transfer beneficial effects of exercise on neurogenesis
and cognition to the aged brain. *Science (New York, N.Y.)* **369**, 167-173,
doi:10.1126/science.aaw2622 (2020).

36   Traschütz, A., Kummer, M. P., Schwartz, S. & Heneka, M. T. Variability and temporal
dynamics of novel object recognition in aging male C57BL/6 mice. *Behav Processes* **157**,
711-716, doi:10.1016/j.beproc.2017.11.009 (2018).

37   Phillips, R. G. & LeDoux, J. E. Differential contribution of amygdala and hippocampus to
cued and contextual fear conditioning. *Behav Neurosci* **106**, 274-285, doi:10.1037//0735-
7044.106.2.274 (1992).

38   Kraeuter, A. K., Guest, P. C. & Sarnyai, Z. The Y-Maze for Assessment of Spatial Working
and Reference Memory in Mice. *Methods Mol Biol* **1916**, 105-111, doi:10.1007/978-1-
4939-8994-2_10 (2019).

39   Wek, R. C. Role of eIF2alpha Kinases in Translational Control and Adaptation to Cellular
Stress. *Cold Spring Harb Perspect Biol* **10**, doi:10.1101/cshperspect.a032870 (2018).

40   Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of
aging. *Cell* **153**, 1194-1217, doi:10.1016/j.cell.2013.05.039 (2013).

41   Chovatiya, R. & Medzhitov, R. Stress, inflammation, and defense of homeostasis. *Mol Cell*
54, 281-288, doi:10.1016/j.molcel.2014.03.030 (2014).

42   Xu, Z. X. *et al.* Elevated protein synthesis in microglia causes autism-like synaptic and
behavioral aberrations. *Nat Commun* **11**, 1797, doi:10.1038/s41467-020-15530-3 (2020).

43   Yin, Z. *et al.* APOE4 impairs the microglial response in Alzheimer's disease by inducing
TGFbeta-mediated checkpoints. *Nat Immunol* **24**, 1839-1853, doi:10.1038/s41590-023-
01627-6 (2023).

44   Sarlus, H. & Heneka, M. T. Microglia in Alzheimer's disease. *J Clin Invest* **127**, 3240-3249,
doi:10.1172/JCI90606 (2017).

45   Smith, L. K. *et al.* beta2-microglobulin is a systemic pro-aging factor that impairs cognitive
function and neurogenesis. *Nat Med* **21**, 932-937, doi:10.1038/nm.3898 (2015).

46   Trombetta, J. J. *et al.* Preparation of Single-Cell RNA-Seq Libraries for Next Generation
Sequencing. *Curr Protoc Mol Biol* **107**, 4 22 21-17, doi:10.1002/0471142727.mb0422s107
(2014).

47   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525-527, doi:10.1038/nbt.3519 (2016).

48   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).

49   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. *Nat Genet* **25**, 25-29, doi:10.1038/75556 (2000).

50   Gene Ontology, C. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids
Res* **49**, D325-D334, doi:10.1093/nar/gkaa1113 (2021).

51   Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P. D. PANTHER version 14:
more genomes, a new PANTHER GO-slim and improvements in enrichment analysis
tools. *Nucleic Acids Res* **47**, D419-D426, doi:10.1093/nar/gky1038 (2019).

52   Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat
Commun* **8**, 14049, doi:10.1038/ncomms14049 (2017).

53   Marsh, S. E. *et al.* Dissection of artifactual and confounding glial signatures by single-cell
sequencing of mouse and human brain. *Nat Neurosci* **25**, 306-316, doi:10.1038/s41593-
022-01022-8 (2022).

54   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*
36, 411-420, doi:10.1038/nbt.4096 (2018).

55   Qiu, X. *et al.* Single-cell mRNA quantification and differential analysis with Census. *Nat
Methods* **14**, 309-315, doi:10.1038/nmeth.4150 (2017).

56   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat
Methods* **14**, 979-982, doi:10.1038/nmeth.4402 (2017).

57   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386,
doi:10.1038/nbt.2859 (2014).

58   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.
Nature **566**, 496-502, doi:10.1038/s41586-019-0969-x (2019).

59   Leger, M. *et al.* Object recognition test in mice. *Nat Protoc* **8**, 2531-2537,
doi:10.1038/nprot.2013.155 (2013).

60   Belarbi, K. *et al.* Beneficial effects of exercise in a transgenic mouse model of Alzheimer's
disease-like Tau pathology. *Neurobiol Dis* **43**, 486-494, doi:10.1016/j.nbd.2011.04.022
(2011).

61   Palovics, R. *et al.* Molecular hallmarks of heterochronic parabiosis at single-cell resolution.
Nature **603**, 309-314, doi:10.1038/s41586-022-04461-2 (2022).

62   Pasciuto, E. *et al.* Microglia Require CD4 T Cells to Complete the Fetal-to-Adult Transition.
Cell **182**, 625-640 e624, doi:10.1016/j.cell.2020.06.026 (2020).

Figure 1

A Pseudotime trajectory

B Microglia age overlayed onto pseudotime trajectory

C Gene expression changes associated with specific regions of pseudotime trajectory

Module | Gene Ontologies

1 — System Development
Tgfbr1, Atf4, mtNd1, Mef2c, Cx3cr1, Mafb

2 — Gene Expression, Response to Stress
Atf3, Klf4, Jun, Fosb, Tgfb1, Ddit3, Nfkbiz

3 — Translation
Rpl's (75 genes), Eif3f, Eef2, Bglap3

4 — Cell Activation, Immune Response, Exocytosis
Il1b, Axl, Ftl1, Lyz2, Lpl, Lgals3, C3

5 — Immune Response, Myeloid Activation
B2m, Rxrg, Cst3, C1qc, Trem2

D

E 12Month/6Month Log2 Fold Change

F 18Month/6Month Log2 Fold Change

G 24Month/6Month Log2 Fold Change

H Heterochronic / Isochronic

I Aging module gene expression changes after heterochronic parabiosis — HY/IY Log2 Fold Change

J Alzheimer's Disease

K Aging module gene expression in response to Aβ — Log2 FC (AD/WT)

Figure 2

Figure 3

Figure 4