Fibroblast Stromal Support Model for Predicting Human Papillomavirus-Associated

Cancer Drug Responses.

Claire D. James[a], Rachel L. Lewis[a], Alexis L. Fakunmoju[a], Austin J. Witt[a], Aya H.

Youssef[a], Xu Wang[a], Nabiha M. Rais[a], Apurva Tadimari Prabhakar[a], J. Mathew

Machado[a], Raymonde Otoa[a], Molly L. Bristol [a,b,#]

[a] Philips Institute for Oral Health Research, School of Dentistry, Virginia Commonwealth

University (VCU), Richmond, Virginia, USA

[b] VCU Massey Comprehensive Cancer Center, Richmond, Virginia, USA

Running Title: Stroma Translational Model for HPV+ Cancer Responses

#Address correspondence to Molly L. Bristol, mlbristol@vcu.edu.

Abstract

Currently, there are no specific antiviral therapeutic approaches targeting Human

papillomaviruses (HPVs), which cause around 5% of all human cancers. Specific

antiviral reagents are particularly needed for HPV-related oropharyngeal cancers

(HPV[+]OPCs) whose incidence is increasing and for which there are no early diagnostic

tools available. We and others have demonstrated that the estrogen receptor alpha (ERα)

is overexpressed in HPV[+]OPCs, compared to HPV-negative cancers in this region, and

that these elevated levels are associated with an improved disease outcome. Utilizing this

HPV[+] specific overexpression profile, we previously demonstrated that estrogen

attenuates the growth and cell viability of HPV$^+$ keratinocytes and HPV$^+$ cancer cells *in*

vitro. Expansion of this work *in vivo* failed to replicate this sensitization. The role of

stromal support from the tumor microenvironment (TME) has previously been tied to

both the HPV lifecycle and *in vivo* therapeutic responses. Our investigations revealed that

in vitro co-culture with fibroblasts attenuated HPV$^+$ specific estrogen growth responses.

Continuing to monopolize on the HPV$^+$ specific overexpression of ERα, our co-culture

models then assessed the suitability of the selective estrogen receptor modulators

(SERMs), raloxifene and tamoxifen, and showed growth attenuation in a variety of our

models to one or both of these drugs *in vitro.* Utilization of these SERMs *in vivo* closely

resembled the sensitization predicted by our co-culture models. Therefore, the *in vitro*

fibroblast co-culture model better predicts *in vivo* responses. We propose that utilization

of our co-culture *in vitro* model can accelerate cancer therapeutic drug discovery.

Importance

Human papillomavirus-related cancers (HPV$^+$ cancers) remain a significant public health

concern, and specific clinical approaches are desperately needed. In translating drug

response data from *in vitro* to *in vivo*, the fibroblasts of the adjacent stromal support

network play a key role. Our study presents the utilization of a fibroblast 2D co-culture

system to better predict translational drug assessments for HPV$^+$ cancers. We also

suggest that this co-culture system should be considered for other translational

approaches. Predicting even a portion of treatment paradigms that may fail *in vivo* with a

co-culture model will yield significant time, effort, resource, and cost efficiencies.

Keywords

Estrogen, ERα, stroma, HPV, human papillomavirus, oropharyngeal cancer, Raloxifene,

Tamoxifen, therapeutics

Introduction

49   Human papillomaviruses (HPVs) are small, double-stranded DNA viruses, and

high-risk HPVs are known carcinogens[1–7]. HPV is the most common sexually transmitted

infection in the United States (U.S.), and estimated to infect more than 80% of the

population at least once in their lifetime[1–3,8–12]. HPV16 is the most prevalent genotype,

accounting for at least 50% of cervical cancers and approximately 90% of HPV[+]

oropharyngeal cancers (HPV[+]OPCs)[9,13,14]. While prophylactic HPV vaccines have

already begun to show remarkable efficacy in preventing infection and related diseases,

HPV continues to account for ~5% of worldwide cancer, and disproportionately affects

marginalized populations both in the U.S. and around the world[1–3,8–12,15–17]. As such, the

lack of specific antiviral therapeutics available for combatting HPV-related cancers is of

significant concern.

60   While many cancers are on the decline, the last two decades have shown a sharp

increase in HPV[+]OPCs, for which there are no early diagnostic tools available[4–6,18,19].

HPV[+]OPCs are found at 4-fold higher levels in men than in women, suggesting there are

sex-related differences in the development of these cancers[4,18]. Using data from The

Cancer Genome Atlas (TCGA), we and others have shown that the estrogen receptor

alpha (ERα) is overexpressed in HPV[+]HNC (head and neck cancers including OPCs) and

that these elevated levels are associated with an improved disease outcome[18,20–26].

67   We have also previously demonstrated that 17-β estradiol (estrogen) attenuates

the growth and cell viability of HPV[+] keratinocytes and HPV[+] cancer cells *in vitro*, but

not HPV negative (HPV⁻) keratinocytes or HPV⁻ cancer cells[26]. Sensitization occurs via

numerous mechanisms: 1) at the level of viral transcription, 2) via interactions with E6

and E7, 3) through manipulation of cell survival and cell death pathways[26]. Here, we

report that the expansion of estrogen treatment into *in vivo NOD-scid IL2Rg^null* (NSG)

mice revealed a lack of response to estrogen alone or in combination with radiotherapy

(IR).

75   Previously, estrogen has been shown to promote HPV-induced cervical disease in

immunocompetent mice, yet this enhancement was lost in NSGs[27–32]. HPV oncogenes, in

conjunction with estrogen, were shown to fundamentally reprogram the tumor

microenvironment (TME)[27,28,33]. We therefore sought to determine if the TME,

specifically the stromal support provided by fibroblasts, alters the estrogenic effects in

our model systems[33–42]. *In vitro* co-culture studies revealed that stromal interactions

markedly change cell growth and viability in response to estrogen in our HPV⁺ models.

82   While our estrogen studies did not prove effective *in vivo*, ERα remains

overexpressed in HPV⁺OPC and HPV⁺ cervical cancers[24–27,29,31–33,43]. Selective estrogen

receptor modulators (SERMs) have proven to provide a multitude of therapeutic

applications[44–50]. Analysis of K14E6/E7 transgenic mice models have previously shown

the possible utility of raloxifene on reduction of recurrence of cervical neoplastic disease,

and earlier literature suggested the utility of tamoxifen to lengthen the latent period for

cervical dysplasia and carcinoma in carcinogen-induced models[45,51]. We utilized our co-

culture system to assess the efficacy of raloxifene and tamoxifen and showed significant

growth repression to one or both SERMs in a number of cancer cell lines. Our

preliminary assessment of these SERMs in an *in vivo* HPV⁺HNC cell line correlated with

our *in vitro* observations and suggests that these drugs may be useful adjuvant approaches

for further investigation. To our knowledge, our analysis is the first to provide evidence

on the utility of SERMs for HPV[+]OPCs.

95   Overall, this report further expands upon the analysis of utilizing estrogen related

signaling in the quest for HPV-specific antiviral approaches[25]. While estrogen presented

compelling evidence *in vitro*, this report demonstrates that estrogen treatment did not

translate to *in vivo* models[26]. The development of co-culture models utilizing fibroblast

"feeder" cells demonstrated that these supporting fibroblasts altered the response to

estrogen *in vitro*, modeling what was observed *in vivo*. Analysis of SERMs with this co-

culture model demonstrated the utility of altering estrogen-related signaling both *in vitro*

and *in vivo*. Therefore, co-culture with fibroblasts offers a simple and more

physiologically relevant environment by stimulating more of the cellular interactions

present in solid tumors. This co-culture model may serve to better predict drug responses

in other translational paradigms, not limited to HPV[+] cancers. Co-culture allows for an

examination of the complex cellular responses to drug effects *in vivo*, thereby enhancing

the accuracy of *in vitro* therapeutic evaluations for more successful translational

approaches.

Results

Estrogen sensitization is not observed *in vivo*

We have previously reported that estrogen attenuates the growth of epithelial cells

in an HPV[+] dependent manner *in vitro*[26]. We found that this occurred via both a

repression of transcription from the HPV16 long control region and through interactions

with the viral oncogenes, E6 and E7[26]. Furthermore, estrogen treatment enhanced

irradiation-induced cell death in an HPV[+] dependent manner[26]. A logical progression was

to assess the response of HPV[+] cancers to estrogen *in vivo*. Previously, our laboratory as

well as others have demonstrated that HeLa cells are highly responsive to estrogen

treatment[26,52,53]. Consequently, experiments were designed to assess the combination of

estrogen and radiation treatment on HeLa xenografts in female NSG mice. Contrary to

our *in vitro* data, Figure 1A shows that estrogen alone, or in combination with radiation,

had no impact on tumor response in mice. Of note, animal weights remained consistent

throughout the study with all treatments (Figure 1B).

123   Previously, we had observed HPV[+] dependent cell death following estrogen

treatment regardless of sex, tissue of origin, or viral genome integration status[26]. To

determine if any of these factors played a role *in vivo*, we decided to expand our analysis

to include the male, episomal, HPV16[+] oropharyngeal cancer cell line UMSCC104[54]. In

addition, we added "early" estrogen supplementation that began the same day as

xenografts were injected, to determine if a temporal relationship was essential to the

estrogen treatment response. As our previous radiation treatment using 10 Gy had some

off-target effects in our NSGs, radiation was reduced to 5 Gy in these studies. As

observed in Figure 1C (animal weight Figure 1D), estrogen again had no impact on tumor

response in any of these conditions. These results complement the previous observations

by the Lambert laboratory utilizing MmuPV1[27]. The Lambert observations demonstrated

that estrogenic effects *in vivo* are reliant, at least in part, on estrogen's suppression of the

host immune system in immunocompetent mice; whereas estrogenic enhancement of

disease progression were not observed in immunodeficient NSG mice[27]. Our results

indicate that the lack of estrogenic alterations in disease progression may not be

papillomavirus species specific, but further studies are needed to confirm this

observation.

Stroma alters HPV-specific estrogen growth response in keratinocyte models

There are numerous differences when moving from *in vitro* to *in vivo* models. An

increasingly recognized component of *in vivo* responses is the adjacent stromal support

network, or "tumor microenvironment" (TME)[33,55,56]. In regard to HPV, evidence also

supports a significant role for stroma during the viral lifecycle and HPV-induced disease

[33,39,41,42]. Fibroblasts are a key component of this stromal support and can significantly

alter cancer resistance and therapeutic responses[33–42]. We already utilize feeder layers of

mitomycin C (MMC) growth-arrested murine 3T3-J2 fibroblasts (referred to as J2s

moving forward) in the immortalization and maintenance of our primary keratinocyte

derived HPV[+] epithelial cultures. Thus, we applied the same approach to investigate

whether this co-culture system affects the response to estrogen. MMC inactivation is a

supported approach to arrest the proliferation of fibroblasts, allowing for the

establishment of a supportive feeder layer that maintains its ability to synthesize RNA

and protein and provide stromal regulation of neighboring cells of interest[57]. This

approach is widely accepted as necessary for maintenance of the HPV genome, as well as

a necessary component of 3D models for HPV lifecycle analysis[58–64]. N/Tert-1 cells

(telomerase immortalized foreskin keratinocytes, HPV negative), as well as HFK+E6E7

(foreskin keratinocytes immortalized by the viral oncogenes only), and HFK[+]HPV16

(foreskin keratinocytes immortalized by the entire HPV16 genome, replicating as an

episome), were cultured in the presence or absence of J2 cells, and treated with 15µM

estrogen, or vehicle control[26]. Compared to untreated, non-cocultured N/Tert-1 cells, no

significant alterations in growth rate were observed following estrogen treatment or co-

culture with J2s (Figure 2A). Conversely, estrogen significantly repressed the growth of

both the E6E7 and HPV16 immortalized cell lines in the absence of stromal support

(Figure 2B and 2C, respectively). The presence of stromal support fibroblasts

significantly rescued this growth suppression (Figure 2B and 2C); coculture with

fibroblasts mitigates the growth suppressive effects of estrogen, *in vitro*. This implies that

the presence of stromal support is responsible for the differences in response to estrogen

treatment that we observed between our *in vitro* and *in vivo* models (Figure 1)[26].

Stroma alters HPV-specific estrogen growth response in cancer models

We sought to expand our co-culture investigations to include cancer lines. A

hallmark of transformation in many cancer cell lines is anchorage independent growth

and loss of adherence in cell culture[65]. To improve our cancer cell count analysis, we

developed a novel, quantifiable co-culture system. Using the Nuclight lentivirus system

(Sartorius), we transduced our cells of interest with nuclear mKate2-red and developed

stable cell lines; this system enabled automated cell counting and could distinguish

between our cells of interest and the non-labeled J2s. An additional advantage of this

system was the ability to monitor cellular morphology and observe any co-culture

influences upon colony formation and cellular distribution. We sought to determine

whether J2s alter cancer cell line responses to estrogen, as observed in our keratinocyte

models (Figure 2). We utilized four cancer cell lines in our co-culture experiments: HN30

– an HPV$^-$ head and neck cancer (p53wt), HeLa – an HPV18$^+$ integrated cervical cancer

(p53wt), UMSCC47 – an HPV16$^+$ integrated head and neck cancer (p53wt), and

UMSCC104 – an HPV16$^+$ episomal head and neck cancer (p53wt). As HeLa cells are

highly sensitive to estrogen, 1.5µM estrogen was utilized in experiments with these cells;

all other cell lines were treated with 15µM as previously described[26,52]. As previously

shown, estrogen treatment significantly repressed cell growth in an HPV$^+$ dependent

manner (Figures 3A,C,E,F)[26]. Conversely, estrogenic sensitivity was no longer observed

in HPV$^+$ cancer cells grown with feeder cells (Figures 3B,D,F,H). This was similar to the

loss of estrogenic sensitivity in our HPV$^+$ keratinocytes (Figure 2), and in our mouse

studies (Figure 1A, and 1C). Again, demonstrating that stromal support alters estrogenic

sensitivity of HPV$^+$ cancer models.

To determine if this repression of estrogenic sensitization was specific to mouse

fibroblast support in co-culture, HeLa cells were grown in the presence or absence of

conditioned media collected from replicating 3T3-J2s cultures, or grown in co-culture

with MMC-inactivated human dermal mesenchymal fibroblasts (HDFMs). Again Figure

4A demonstrates that HeLa cells in monoculture undergo significant growth repression in

the presence of estrogen. Similarly, estrogenic growth repression was observed in the

presence of J2 conditioned media (Figure 4B). Co-culture with HDFMs rescued

estrogenic growth repression in HeLa cells (Figure 4C), similar to what was observed

with J2s (Figure 3B). HDFMs did not repress the growth of HeLa alone (Figure 4C),

contrary to the growth repression observed in the presence with J2 (Figure 3B). While we

do not currently understand the mechanism behind this altered growth potential with

alternative species' fibroblasts, we find it noteworthy that estrogenic sensitivity is lost

with both co-culture methods, but not by incubation with media, indicating that cell-cell

contact is required. Moreover, this loss of sensitivity is not dependent on fibroblast

growth alteration of cancer cells.

Stroma does not alter the response to cisplatin in cancer models.

To assess the predictiveness of our co-culture model, we found it important to investigate whether stroma would alter the response to other chemotherapeutic approaches. HeLa, UMSCC47, UMSCC104, and HN30 cells were therefore evaluated for their responsiveness to cisplatin (Figure 5A,C,E,G). Fibroblasts did not rescue the growth arrest observed in all cell lines (Figures 5B,D,F,H). Cisplatin is a well-established clinical treatment paradigm for HPV[+] and HPV[-]OPCs, cervical cancer, and many other cancers[66–74]. We predicted that fibroblasts would not be able to rescue cancer cells from this conventional chemotherapeutic agent. We postulate that fibroblasts are unlikely to change the response to most currently accepted cancer treatment modalities. Instead, we suggest that this model may better predict which novel therapeutics may fail translational approaches. This is something we are currently investigating, and we encourage others to consider this approach as well when translating treatments from *in vitro* to *in vivo*.

Stroma alters HPV long control region (LCR) transcriptional regulation

We previously published that estrogen represses transcription of the HPV16 long control region (LCR) both in our N/Tert-1 and C33a transcription models[26]. This transcriptional repression of the LCR, in turn, downregulated the expression of early viral genes in our numerous HPV[+] keratinocyte and HPV[+] cancer models[26]. Expanding upon these previous observations, we further wanted to determine the impact of stromal support on estrogenic HPV16-LCR transcriptional regulation. N/Tert-1 cells grown in the presence of stromal support had significantly enhanced levels of HPV16-LCR transcription and this transcriptional regulation was no longer significantly repressed by estrogen (Figure 6A); this may be one of the many contributing factors involved in the

loss of estrogen sensitivity. ERα was also assessed, and stroma did not appear to alter

protein expression in N/Tert-1 cells (Figure 6B). These results suggest that stroma is

highly supportive of HPV16-LCR transcriptional regulation.

Fibroblasts are routinely utilized to support HPV genome maintenance and the

viral life cycle in keratinocyte models[33,41,57,75–82]. While this model is accepted, the full

mechanism of how fibroblasts aid maintenance of the viral genome as an episome has yet

to be elucidated. Figure 6A demonstrates that stromal support enhances HPV16-LCR

transcription. It was therefore important to confirm that this transcriptional regulation had

downstream effects. Previously, elements of the HPV upstream regulatory region,

another term for the viral LCR, have been implicated as a requirement for long term viral

persistence in keratinocytes[83]. The McBride laboratory demonstrated that the chromatin

architecture of this region is important for genome partitioning and may influence

integration[83]. Figure 6C confirms that HFK[+]HPV16 begin to integrate the viral genome

when grown in the absence of J2 for one week[84,85]. The accepted practice of fibroblast co-

culture, may in part maintain viral episomes via influencing the transcription of LCR.

Thus, stroma supports viral protein expression and alters host signaling pathways in

keratinocytes

We next investigated viral protein expression and host protein signaling observed

in the presence or absence of fibroblast support. HFK+E6/E7, HFK[+]HPV16, and N/Tert-

1 cells (Figure 6D) were grown in the presence or absence of J2s. J2s are washed off

before harvesting samples, and are murine-derived so there should be limited detection

via human-specific antibodies. To confirm that altered protein levels observed were not

due to any residual J2s, lane 1 in Figure 6D demonstrates that significant specific bands

were not observable with the majority of our chosen antibodies with 100μg of J2 protein

input. SIRT1 does have a notable lower band which is not unexpected due to the gene

homology between mouse and human SIRT1 and that the immunogen was developed

from the amino acids 1-131 of mouse Sir2α. γH2AX has a low-level detectible band due

to cross reactivity of the antibody; it should be noted that the significant observations

discussed have taken this into account. Figure 6D demonstrates that HFK+E6E7, which

do not rely on the LCR for early gene expression, do not have altered E7 protein levels in

the presence or absence of fibroblasts (lanes 2,3). As fibroblasts were shown to enhance

LCR transcription (Figure 6A), HFK[+]HPV16 cells, which rely on the LCR for early gene

expression, have enhanced E7 levels when grown in the presence of J2s (lanes 4,5).

HFK+E6E7 had limited levels of p53 expression, and HFK[+]HPV16 had low expression

of p53, due to E6-targeted degradation; however significant enhancement of p53 levels in

the presence of J2 in both cell lines were striking[63,64,79,86–91]. To our knowledge, this is the

first report to suggest why p53 is not always fully degraded in HPV[+] cell lines; E6

degradation of p53 may depend on whether or not keratinocytes are maintained on feeder

cells[79,88,90,92–97]. Upregulation of p53 in the presence of J2 was also observed in N/Tert-1

cells (lanes 6,7) so this observation is independent from E6 or full genome expression. It

is worth noting that HFK[+]HPV16 cells are consistently maintained in co-culture with J2s,

and that J2s were not supplemented for the "control" conditions. This also demonstrates

that the stromal induced alterations of p53 protein levels are reversible (Figure 6D). Total

levels of pRb were not altered N/Tert-1 cells nor in HFK+E6E7; this corresponds to

unaltered E7 levels in HFK+E6E7 cells. Alternatively, pRb was reduced in HFK[+]HPV16

grown in the presence of J2s. Again, suggesting fibroblasts are important for viral

regulation of keratinocyte signaling, possibly through the LCR. A consistent observation

was the upregulation of γH2AX in the presence of J2s in all of the cell lines (Figure 6D).

Numerous reports have demonstrated HPV viral integrity and genome stability is highly

reliant on DNA repair machinery, including γH2AX; J2 enhancement of γH2AX may be

another key mechanism of fibroblast regulation of viral genome stability in keratinocyte

models[7,76,98–106]. Upstream of p53 and γH2AX, SIRT1 is known to target histones and

non-histone substrates such as p53, and has been shown to decrease in response to DNA

damage[107]. N/Tert-1 and HFK⁺HPV16 grown in the presence of J2s demonstrated a

significant decrease in SIRT1. The lower band observed can be associated with SIRT1

post-translational modifications; while this was altered by fibroblasts in both HFK+E6E7

and HFK⁺HPV16, conclusions were not made due to the band observed with the J2 input

sample [40,107–110]. While we are still investigating additional signaling mechanisms

involved, these observations highlight the importance of properly controlling for whether

or not cell lines are grown in the presence or absence of feeder cells when considering

viral and host signaling events.

Selective estrogen receptor modulators reduce growth rates *in vitro* and *in vivo*

While estrogen had no impact on the tumor response in mice, alone or in

combination with radiation (Figure 1), we expanded our analysis to assess if the selective

estrogen receptor modulators (SERMs) raloxifene or tamoxifen would be a useful

approach to continue to take advantage of the HPV⁺ specific overexpression of ERα. It

should be noted that SERMs can be agonists or antagonists of the ERα, and these

responses are dependent on region, cell type, and the localization and availability of

estrogen response elements (EREs)[44,46,47,111]. Preclinical data supports the utility of

SERMs for cervical cancer, particularly the utility of raloxifene on the reduction of

recurrent neoplastic disease;   however, SERMs have yet to be evaluated in

HPV^{+}OPC[47,111].

HeLa, UMSCC47, and UMSCC104 HPV^{+} cancer cells grown without fibroblast

support exhibited significant growth repression to both SERMs (Figures 7A,C,E). While

fibroblast support did not alter the response to SERMs in all the cell lines, rescued growth

was observed in HeLa cells treated with tamoxifen while remaining responsive to

raloxifene (Figures 7B,D,F). It is worth noting that previous *in vivo* and clinical analysis

have predicted the utility of raloxifene in HPV^{+} cervical cancer response, whereas

tamoxifen is not recommended[45–47,51,111]. We therefore sought to analyze SERM

responsiveness in an HPV^{+}HNC *in vivo* model. Expansion of SERM treatment *in vivo*

was therefore conducted in UMSCC104 cells alone or in combination with IR to assess

their utility as well as determine if our co-culture model could be useful in future

translational approaches. For this study, 4 Gy IR was chosen to further reduce the off-

target issues relating to radiation use in NSGs[112]. As observed in Figure 8A, radiation

alone significantly reduced tumor volume starting on Day 14, while raloxifene (Figure

8A) and tamoxifen alone (Figure 8B) were able to significantly reduce tumor volume

starting on Day 32. Furthermore, in comparison to radiation (IR) alone, tamoxifen+IR

significantly reduced tumor volume starting on Day 42 (Figure 8B). Kaplan-Meier

survival analysis demonstrated 50% survival for tamoxifen+IR on day 70 in comparison

to 10% survival at the same time point for IR alone or raloxifene+IR at our chosen

endpoint (Figure 8C).   With all treatments, animal weights remained consistent

throughout the study (Figure 8D). Our *in vitro* co-culture system modeled our *in vivo*

responses, as well as those previously observed, to both estrogen and SERMs in HPV$^+$

cervical *in vivo* models (Figure 1,3,7) [27,32,43,45–47,51,111]. Co-culture also predicted the

responsiveness of SERMs in an HPV$^+$HNC *in vivo* model (Figure 7,8) and suggests that

the utility of SERMs for the treatment of HPV$^+$OPC are worth further investigation.

Altogether, we suggest that the predictiveness of this co-culture system should be

considered in more translational approaches. Current studies are investigating the

molecular mechanisms behind the alterations observed when cells are grown in fibroblast

co-culture. These mechanistic approaches may further expand the predictive utility of our

co-culture model.

Discussion

We previously established that estrogen attenuates the growth of HPV$^+$

keratinocyte and cancer cell lines in both an LCR and E6E7 dependent manner[25,26]. Of

note, when these studies were conducted, we did not supplement fibroblast support

during drug treatment[26]. Conversely, when estrogen was utilized to treat HPV$^+$ xenografts

in NSG mice, responsiveness was no longer observed (Figure 1). This data is supportive

of previous observations by the Lambert laboratory[27]. We acknowledge the importance of

the immune system in the context of HPV, the responsiveness to estrogen, and its

potential impact on translational approaches as well[27,28,45,113]. As NSG mice possess a

significantly compromised immune system, we then sought to investigate the role that

stromal support may play in the altered responsiveness to estrogen when moving from *in*

vitro to *in vivo* models. In doing so, we found that there is an HPV-specific change in

response to estrogen when comparing co-culture to non-fibroblast conditions (Figure

2,3,4). We find that this response is, at least in part, HPV16 LCR dependent; however,

there are likely other mechanisms at play (Figure 4). Additional mechanisms behind the

observed stromal growth alterations, as well as the response to therapeutics, are currently

under investigation in our laboratory and will be the subject of future reports.

Of note, the analysis of alternative species fibroblasts and mesenchymal cell types

on basement membrane arrangement and growth characteristics of organotypic raft

cultures were conducted many years ago[114]. Tissue-specific, species-specific, and spatial-

specific alterations were found to impact numerous epithelial phenotypes[114]. Extracellular

matrix components were found to have the greatest impact in this publication. We

propose that the differences in extracellular matrix components between mouse and

human might have moderate plating efficiency or growth efficiency alterations, and

contribute to the altered growth potential observed in HeLa cells (Figure 3,4). As seen in

Figure 3, UMSCC47 cell growth was not impacted by J2s; alternatively, UMSCC104 and

HN30 cells grew better in the presence of J2. Regardless of the altered growth potential

of cell lines in the presence or absence of fibroblasts, the observation that estrogenic

sensitivity was lost in co-culture remains the same.

Anecdotally, is it known in the HPV field that J2s are crucial to the culture of

primary keratinocytes for the maintenance of viral genome stability[75,76,104]. Expanding the

accepted fibroblast co-culture system for HPV[+] primary keratinocytes, our data

demonstrates that this model promotes translational utility. When keratinocytes or cancer

cells were grown in this co-culture system, *in vivo* results were more predictable. These

models allowed for us to translate our estrogen results into estrogen pathway targeting

drugs. Previous studies have highlighted the potential utility of SERMs in HPV[+] cervical

cancers, most specifically the use of raloxifene to reduce neoplastic recurrance[45,46,48,111].

We have expanded SERM analysis from cervical cancer to oropharyngeal cancer. We

now demonstrate that SERMs may present clinical applicability as adjuvant approaches

in HPV[+]OPC and future investigations are warranted. While our observations apply both

to primary keratinocytes and cancer models of HPV, we are currently in collaboration to

expand analysis to other cancer treatment models. Nevertheless, stroma is recognized to

contribute to the response to therapeutics and the development of resistance, and we

conclude that it should be more often considered in future translational

approaches[33,35,39,40,92].

Materials and Methods

Cell Culture

HN30 (generous gift from Hisashi Harada, VCU Philips Institute), UMSCC47 (Millipore;

Burlington, MA, USA), and HeLa (generous gift from Alison McBride, NIAID) cells

were grown in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen; Carlsbad,

CA, USA) supplemented with 10% charcoal/dextran stripped fetal bovine serum (Gemini

Bio-products; West Sacramento, CA, USA). UMSCC104 (Millipore) cells were grown in

Eagle's minimum essential medium (EMEM) (Invitrogen) supplemented with

nonessential amino acids (NEAA) (Gibco) and 20% charcoal/dextran stripped fetal

bovine serum. N/Tert-1 cells and all derived cell lines have been described previously

and were maintained in keratinocyte-serum free medium (K-SFM; Invitrogen),

supplemented with a 1% (vol/vol) PenStrep (Gibco) and previously described

antibiotics[20–22,115–120]. HFK[+]HPV16 have been previously described and were grown in

Dermalife-K complete media (Lifeline Technology), and maintained on inactivated

fibroblast feeder cells (described below)[121]. HFK+E6/E7 were grown in Dermalife-K

complete media (Lifeline Technology), and maintained on inactivated fibroblast feeder

cells (described below); the immortalization process is described below. Of note, we have

no issues with fibroblast plating efficiency, or keratinocyte viral episome maintenance

utilizing these keratinocyte complete media kits over the traditional use of E-media[121,122].

For all cells not directly purchased from companies, the cell type was confirmed by Johns

Hopkins or MD Anderson cell line authentication services, and the cells were maintained

at 37°C in a 5% CO_2–95% air atmosphere, passaged every 3 or 4 days, and routinely

monitored for mycoplasma (Sigma, MP0035).

Flank xenografts for *in vivo* drug trials

HeLa and UMSCC104 cells were stably transduced with a lentiviral vector for pLX304

Luciferase-V5 blast (generous gift from Renfeng Li, originally obtained from Kevin

Janes, Addgene plasmid # 98580)[123]. Cell lines were selected with 10μg/ml blasticidin.

Expression was verified with bioluminescent imaging, further outlined and defined in

transcriptional activity analysis method detailed below.

Xenografting was performed using previously described methodology, in collaboration

with the Virginia Commonwealth University (VCU) Cancer Mouse Models Core

Laboratory (CMMC)[124,125]. All experiments were conducted in accordance with animal

protocol AD10002330 approved by VCU Institutional Animal Care and Use Committee.

NOD-SCID-IL2γ receptor null (NSG) mice (6-8 weeks old) were injected with 1×10^6

cells suspended in PBS and Cultrex™ basement membrane extracts (BME) (Bio-

techne/R&D Systems) into the right flank. HeLa studies were conducted in female mice;

UMSCC104 studies were conducted in male mice (chosen to mimic sex from human

donors). Days 1-3 post-xenograft and at varying times throughout the studies,

bioluminescence imaging was performed using a Xenogen IVIS-100 system (Calipers

Life Sciences, Hopkinton, MA) to verify xenograft establishment, growth, and possible

metastasis based on previously established protocols[126]. Tumor volume was also

measured on noted dates and calculated as $V = AB2 (\pi/6)$, where A is the longest

dimension of the tumor, and B is the dimension of the tumor perpendicular to A. Data

points presented on the graph are representative for each condition while more than 70%

of the animals remained in each group. HeLa cells were palpable on day 10, UMSCC104

cells were palpable on day 7. It should be noted that in our hands, UMSCC104 xenografts

are prone to ulceration. Animals were humanely euthanized when ulcerations were

observed.

NSG estrogen delivery

A combinational approach for estrogen delivery was based on modified protocols

designed in collaboration with the source of our obtained control and estrogen beeswax

pellets (0.4mg estrogen, Huntsman Cancer Institute, University of Utah)[127,128]. Estrogen

was also delivered in drinking water using a protocol kindly shared online by the Wicha

lab[129]. Briefly, a 2.7 mg/mL stock of 17-estradiol (Sigma # E2758) in ethanol was diluted

to a final concentration of 8 µg/mL in sterile drinking water. Pellets (control or estrogen)

were implanted after tumors were palpable. Drinking water supplementation also began

at this time point. "Early estrogen" in UMSCC104 denotes that water estrogen

supplementation began the same day as xenograft injections (pellets were still implanted

when tumors became palpable on day 7).

NSG SERM delivery

Treatment of mice with Raloxifene was performed as previously described by the

Lambert Laboratory[45,111]. The human formulation of raloxifene hydrochloride (60mg

tablets; EVISTA; Eli Lilly) were purchased from Virginia Commonwealth University

Health System Pharmacy. Tablets were resuspended in PBS for a final concentration of

10 mg/ml. Mice were administered a 150 μl drug suspension (equivalent to 1.5 mg) via

i.p. injection. Treatment of Tamoxifen was performed as previously described[130].

Tamoxifen (Sigma, T-5648) was resuspended in corn oil (Sigma C-8267) at 37°C for a

final concentration of 10 mg/ml. Mice were administered 100 μl drug suspension

(equivalent to 1.0 mg) via i.p. injection. SERM treatments began after tumors were

palpable. Mice received treatment 5 days a week for 4 weeks, for a total of 20 injections.

Small Animal Radiation Research Platform (SARRP) ionizing radiation delivery

1 day following pellet implantation to allow for mouse recovery, targeted ionizing

radiation (IR) was delivered utilizing the Xstrahal SARRP. HeLa studies utilized 10 Gy.

Due to the radiation sensitivity of NSG mice, UMSCC104 studies reduced IR dose to 5

Gy, and finally to 4 Gy.

Culture, conditioned media collection, and mitomycin C (MMC) inactivation of

3T3-J2 mouse feeder cells

3T3-J2 immortalized mouse fibroblasts (J2) were grown in DMEM and supplemented

with 10% FBS. Fresh media was exchanged twice a week; conditioned media was spun

down as 500 rcf to remove any residual cells[131,132]. 80-90% confluent plates were

supplemented with 4μg/ml of MMC in DMSO (Cell Signaling Technology) for 4-6 hours

at 37°C. MMC-supplemented medium was removed and cells were washed with 1xPBS.

Cells were trypsinized, spun down at 500 rcf for 5 mins, washed once with 1xPBS, spun

again, and resuspended. Quality control of inactivation (lack of proliferation) was

monitored for each new batch of mitomycin-C. 100-mm plate conditions were

supplemented with $1\text{x}10^6$ J2 and 6-well plate conditions were supplemented with $1\text{x}10^5$

every 2-3 days; for longer term cultures, any remaining J2s were washed off with 1x PBS

and new J2 were continually supplemented every 2-3 days.

Culture and mitomycin C (MMC) inactivation of human dermal mesenchymal

fibroblast feeder cells

Human dermal mesenchymal fibroblasts (HDFM) were grown, treated, and quality

controlled as described for the above J2 protocol. 6-well plate conditions were

supplemented with $1\text{x}10^5$ every 2-3 days.

Generation of E6E7-immortalized keratinocytes

Primary keratinocytes from single donors were obtained from LifeLine Cell

Technologies[121]. Cells were cultured on collagen-coated plates for lentiviral delivery of

HPV16 E6E7, using the pLXSN16E6E7 plasmid (Addgene plasmid # 52394, a gift from

Denise Galloway)[133]. Following selection with G418 (72mM), cells were cultured on

mitomycin-C inactivated 3T3-J2 fibroblasts. HFKs were cultured in DermaLife-K

Complete media (LifeLine Cell Technologies) and E6E7 expression was confirmed by

qRTPCR. Of note, these cells were generated at the same time and utilizing the same

donors as the previously described HFK-HPV16[121].

Co-culture of keratinocytes in the presence or absence of inactivated fibroblasts

N/Tert-1, HFK E6E7, or HFK HPV16 cells were seeded at $5\text{x}10^5$ in 100-mm plates in the

presence or absence of previously seeded J2s (at least 6 hours prior). Twenty-four hours

later, noted cells were supplemented with 15 μM 17β-estradiol (estrogen). Forty-eight

hours after estrogen supplementation, plates were washed to remove residual J2 and cells

were trypsinized and counted. For analysis of 1 week time point, HFK E6E7 or HFK

HPV16 cells were seeded at $1x10^5$ in 100-mm plates in the presence or absence of

previously seeded J2s. J2s were re-supplemented every 2-3 days as previously described.

Pellets from these experiments were utilized for subsequent immunoblotting or DNA

analysis, detailed below.

Generation of stable nuclear labeled cells with Incucyte® Nuclight Lentivirus

mKate2 Incucyte® Nuclight Lentivirus (puro) cells were generated according to the

Sartorius product guide protocol (Sartorius cat# 4476), using a MOI of 3 or 6, depending

on cell type. Cells generated were maintained in 1µg/ml puromycin supplemented media.

Fluorescence was routinely monitored by BZ-X TexasRed filter via the Keyence BZ-

X800 inverted fluorescence microscope.

Co-culture of nuclear-labeled cancer cell lines in the presence or absence of

inactivated fibroblasts

Stable mKate2-puro HeLa, UMSCC47, UMSCC104, or HN30 cells were seeded in

triplicate at $1x10^4$ per well in 6-well plates in the presence or absence of previously

seeded J2 cells, or HDFM cells (at least 6 hours prior – fibroblast type is noted for each

experiment). Twenty-four hours after seeding, day 0 images were captured in brightfield

and TexasRed with the Keyence BZ-X800 Image Cytometer. Noted drugs were

supplemented immediately after this initial imaging: 1.5µM (HeLa) or 15 µM 17β-

estradiol (Sigma), 10µM Tamoxifen (MP Biomedicals), 10µM Raloxifene (Cayman

Chemical Company), or 10-20µM Cisplatin (APExBIO). Cytometry images were again

captured on day 1 and day 3. UMSCC104 were washed on day 3 after imaging, new J2

and drug were supplemented, and additional images were captured on day 5 and 7. Ten

fields of view were randomized per well for all conditions. Cell count image cytometry

batch analysis was performed using the Keyence BZ-X800 Image Analyzer software. All

conditions utilized set analysis conditions from a single randomized control image and

applied to all data points automatically to reduce variability and bias. Data is presented as

fold of control from day 0. Representative images are presented in Figure 9.

Immunoblotting

Specified cells were washed with 1X PBS and trypsinized. Pellets were washed with 1X

PBS and resuspended in 5X packed cell volume of NP40 buffer (50mM Tris-HCl Ph 7.5,

150mM NaCl, 1%NP-40, 5mM EDTA) supplemented with Roche cOmplete protease

inhibitor and Roche PhosSTOP phosphatase inhibitor. Cell-lysis buffer suspension was

incubated on ice for 30 min with occasional agitation, then centrifuged for 15 min at

14,000 rcf at 4 °C. Supernatant protein concentration was measured via the Bio-Rad

protein estimation assay according to manufacturer's instructions. 100 μg protein samples

were heated at 95 °C in 4x Laemmli sample buffer (Bio-Rad) for 5 min. Noted samples

were run down a Novex 4–12% Tris-glycine gel (Invitrogen) and transferred onto

a nitrocellulose membrane (Bio-Rad) at 30V overnight, or 100V for 1 hour using the wet-

blot transfer method. Membranes were blocked with Odyssey (PBS) blocking buffer

(diluted 1:1 with 1X PBS) at room temperature for 1 hour and probed with indicated

primary antibody diluted in Odyssey blocking buffer. Membranes were washed with PBS

supplemented with 0.1% Tween (PBS-Tween) and probed with the indicated Odyssey

secondary antibody 1:10,000 (goat anti-mouse IRdye 800CW or goat anti-rabbit IRdye

680CW) diluted in Odyssey blocking buffer. Membranes were washed three times with

PBS-Tween and an additional wash with 1X PBS. Infrared imaging of the blot was

performed using the Odyssey CLx Li-Cor imaging system. Immunoblots were quantified

using ImageJ utilizing GAPDH as the internal loading control. The following primary

antibodies were used for immunoblotting in this study at 1:1000, unless otherwise noted:

ERα (Abcam, ab32063), GAPDH 1:10,000 (Santa Cruz, sc-47724), pRb (Santa Cruz, sc-

102), p53 (Cell Signaling Technology, CST-2527 and CST-1C12), γH2AX 1:500 (Cell

Signaling Technology, CST-80312 and CST-20E3), SIRT1 (EMD Millipore 07-131), β-

actin (Santa Cruz, sc-47778), E7 1:500 (Santa Cruz, sc-6981).

Transfection and transcriptional activity analysis.

N/Tert-1 cells were plated at a density of 5×10^5 in 100-mm dishes. The following day,

the previously described plasmids for pGL3 basic, pGL3 control, or pHPV16-LCR-Luc

were transfected Lipofectamine 2000 (according to the manufacturer's instructions,

ThermoFisher Scientific). Twenty-four hours after transfection, cells were washed, and

noted cells were supplemented with 15 μM 17β-estradiol; J2s were also supplemented at

this time point for noted conditions to reduce likelihood of altered transfection efficiency.

Forty-eight hours after transfection, cells were harvested utilizing Promega reporter lysis

buffer and analyzed for luciferase using the Promega luciferase assay system.

Concentrations were normalized to protein levels, as measured by the Bio-Rad protein

assay dye. Relative fluorescence units (RFU) were measured using the BioTek Synergy

H1 hybrid reader.

Exonuclease V assay

PCR based analysis of viral genome status was performed using methods described by

Myers *et al.* [84]. Briefly, 20 ng genomic DNA was either treated with exonuclease V

(RecBCD, NEB), in a total volume of 30 ul, or left untreated for 1 hour at 37°C followed

by heat inactivation at 95°C for 10 minutes. 2 ng of digested/undigested DNA was then

quantified by real time PCR using a 7500 FAST Applied Biosystems thermocycler with

SYBR Green PCR Master Mix (Applied Biosystems) and 100 nM of primer in a 20 μl

reaction. Nuclease free water was used in place of the template for a negative control.

The following cycling conditions were used: 50°C for 2 minutes, 95°C for 10 minutes, 40

cycles at 95°C for 15 seconds, and a dissociation stage of 95°C for 15 seconds, 60°C for

1 minute, 95°C for 15 seconds, and 60°C for 15 seconds. Separate PCR reactions were

performed to amplify HPV16 E6 F: 5'- TTGCTTTTCGGGATTTATGC-3' R: 5'-

CAGGACACAGTGGCTTTTGA-3',   HPV16   E2   F: 5'-

TGGAAGTGCAGTTTGATGGA-3' R: 5'- CCGCATGAACTTCCCATACT-3', human

mitochondrial   DNA   F: 5'-CAGGAGTAGGAGAGAGGGAGGTAAG-3'   R: 5'-

TACCCATCATAATCGGAGGCTTTGG -3', and human GAPDH DNA F: 5'-

GGAGCGAGATCCCTCCAAAAT-3' R: 5'- GGCTGTTGTCATACTTCTCATGG-3'

Reproducibility, research integrity, and statistical analysis

All *in vitro* experiments were carried out at least in triplicate in all of the cell lines

indicated. All cell lines were bought directly from sources indicated, or typed via cell line

authentication services. All images shown are representatives from triplicate experiments.

In vivo experiments were designed in collaboration with the VCU Massey Cancer Center

animal core and biostats core for sample size justification and statistical power analysis.

Quantification is presented as mean +/- standard error (SE). Student's t-test or analysis of

variance (ANOVA) were used to determine significance as appropriate: * $p<0.05$,

**$p<0.01$, ***$p<0.001$

Data availability

Following the 2023 NIH data management and sharing policy, all data resulting from the development of projects will be available in scientific communications presented at conferences and in manuscripts that will be published in peer-reviewed scientific journals. Data will be deposited in the Open Science Framework (OSF) platform. OSF can be accessed at https://osf.io. VCU is an OSF institutional member, and OSF is an approved generalist repository for the 2023 NIH data management and sharing policy.

Acknowledgements

This work was supported by the VCU Philips Institute for Oral Health Research, the National Institute of Dental and Craniofacial Research/NIH/DHHS R03 DE029548, and the National Cancer Institute-designated Massey Cancer Center grant P30 CA016059. Mouse services and products in support of the research project were generated by the Virginia Commonwealth University Cancer Mouse Models Core Laboratory, supported, in part, with funding from NIH-NCI Cancer Center Support Grant P30 CA016059.

References

1. Brianti, P., De Flammineis, E. & Mercuri, S. R. Review of HPV-related diseases and cancers. *New Microbiol.* **40**, 80–85 (2017).

2. de Martel, C., Plummer, M., Vignat, J. & Franceschi, S. Worldwide burden of cancer attributable to HPV by site, country and HPV type. *Int. J. Cancer* **141**, 664–670 (2017).

3. McBride, A. A. & Münger, K. Expert Views on HPV Infection. *Viruses* **10**, 94 (2018).

4.   Marur, S., D'Souza, G., Westra, W. H. & Forastiere, A. A. HPV-associated head

and neck cancer: a virus-related cancer epidemic. *Lancet Oncol.* **11**, 781–789

(2010).

5.   Lowy, D. R. & Munger, K. Prognostic implications of HPV in oropharyngeal

cancer. *N. Engl. J. Med.* **363**, 82–84 (2010).

6.   Sheedy, T. & Heaton, C. HPV-associated oropharyngeal cancer. *Jaapa* **32**, 26–31

(2019).

7.   Porter, V. L. & Marra, M. A. The Drivers, Mechanisms, and Consequences of

Genome Instability in HPV-Driven Cancers. *Cancers* **14**, 4623 (2022).

8.   Chesson, H. W., Dunne, E. F., Hariri, S. & Markowitz, L. E. The estimated lifetime

probability of acquiring human papillomavirus in the United States. *Sex. Transm.*

Dis. **41**, 660–664 (2014).

9.   Kang, S. D. *et al.* Effect of Productive Human Papillomavirus 16 Infection on

Global Gene Expression in Cervical Epithelium. *J. Virol.* **92**, e01261-18 (2018).

10.   McLaughlin-Drubin, M. E. & Meyers, C. Evidence for the coexistence of two

genital HPV types within the same host cell in vitro. *Virology* **321**, 173–180 (2004).

11.   McLaughlin-Drubin, M. E. & Münger, K. Oncogenic activities of human

papillomaviruses. *Virus Res.* **143**, 195–208 (2009).

12.   McLaughlin-Drubin, M. E., Meyers, J. & Munger, K. Cancer associated human

papillomaviruses. *Curr. Opin. Virol.* **2**, 459–466 (2012).

13.   zur Hausen, H. Papillomaviruses in the causation of human cancers - a brief

historical account. *Virology* **384**, 260–265 (2009).

14. Psyrri, A. & DiMaio, D. Human papillomavirus in cervical and head-and-neck

cancer. *Nat. Clin. Pract. Oncol.* **5**, 24–31 (2008).

15. Huh, W. K. *et al.* Final efficacy, immunogenicity, and safety analyses of a nine-

valent human papillomavirus vaccine in women aged 16–26 years: a randomised,

double-blind trial. *The Lancet* **390**, 2143–2159 (2017).

16. Kavanagh, K. *et al.* Changes in the prevalence of human papillomavirus following a

national bivalent human papillomavirus vaccination programme in Scotland: a 7-

year cross-sectional study. *Lancet Infect. Dis.* **17**, 1293–1302 (2017).

17. Meeting, I. W. G. on the E. of C. R. to H. & Cancer, I. A. for R. on. *Human*

Papillomaviruses. (World Health Organization, 2007).

18. Sung, H. *et al.* Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence

and Mortality Worldwide for 36 Cancers in 185 Countries. *CA. Cancer J. Clin.* **71**,

209–249 (2021).

19. Roman, B. R. & Aragones, A. Epidemiology and incidence of HPV-related cancers

of the head and neck. *J. Surg. Oncol.* **124**, 920–922 (2021).

20. Nulton, T. J., Olex, A. L., Dozmorov, M., Morgan, I. M. & Windle, B. Analysis of

The Cancer Genome Atlas sequencing data reveals novel properties of the human

papillomavirus 16 genome in head and neck squamous cell carcinoma. *Oncotarget*

8, 17684–17699 (2017).

21. Evans, M. R. *et al.* An oral keratinocyte life cycle model identifies novel host

genome regulation by human papillomavirus 16 relevant to HPV positive head and

neck cancer. *Oncotarget* **8**, 81892–81909 (2017).

22. Evans, M. R. *et al.* Human Papillomavirus 16 E2 Regulates Keratinocyte Gene

641   Expression Relevant to Cancer and the Viral Life Cycle. *J. Virol.* **93**, e01941-18

642   (2019).

643   23. Nulton, T. J., Kim, N.-K., DiNardo, L. J., Morgan, I. M. & Windle, B. Patients with

644   integrated HPV16 in head and neck cancer show poor survival. *Oral Oncol.* **80**, 52–

645   55 (2018).

646   24. Kano, M. *et al.* Expression of estrogen receptor alpha is associated with

647   pathogenesis and prognosis of human papillomavirus-positive oropharyngeal

648   cancer. *Int. J. Cancer* **145**, 1547–1557 (2019).

649   25. James, C. D., Morgan, I. M. & Bristol, M. L. The Relationship between Estrogen-

650   Related Signaling and Human Papillomavirus Positive Cancers. *Pathog. Basel*

651   *Switz.* **9**, 403 (2020).

652   26. Bristol, M. L., James, C. D., Wang, X., Fontan, C. T. & Morgan, I. M. Estrogen

653   Attenuates the Growth of Human Papillomavirus-Positive Epithelial Cells. *mSphere*

654   **5**, e00049-20 (2020).

655   27. Wang, W. *et al.* Stress keratin 17 and estrogen support viral persistence and

656   modulate the immune environment during cervicovaginal murine papillomavirus

657   infection. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2214225120 (2023).

658   28. Spurgeon, M. E. *et al.* Human papillomavirus oncogenes reprogram the cervical

659   cancer microenvironment independently of and synergistically with estrogen. *Proc.*

660   *Natl. Acad. Sci. U. S. A.* **114**, E9076–E9085 (2017).

29. Chung, S.-H., Wiedmeyer, K., Shai, A., Korach, K. S. & Lambert, P. F.

Requirement for estrogen receptor alpha in a mouse model for human

papillomavirus-associated cervical cancer. *Cancer Res.* **68**, 9928–9934 (2008).

30. Jabbar, S. F., Abrams, L., Glick, A. & Lambert, P. F. Persistence of high-grade

cervical dysplasia and cervical cancer requires the continuous expression of the

human papillomavirus type 16 E7 oncogene. *Cancer Res.* **69**, 4407–4414 (2009).

31. Son, J., Park, J. W., Lambert, P. F. & Chung, S.-H. Requirement of estrogen

receptor alpha DNA-binding domain for HPV oncogene-induced cervical

carcinogenesis in mice. *Carcinogenesis* **35**, 489–496 (2014).

32. Chung, S.-H., Franceschi, S. & Lambert, P. F. Estrogen and ERalpha: culprits in

cervical cancer? *Trends Endocrinol. Metab. TEM* **21**, 504–511 (2010).

33. Spurgeon, M. E. & Lambert, P. F. Human Papillomavirus and the Stroma:

Bidirectional Crosstalk during the Virus Life Cycle and Carcinogenesis. *Viruses* **9**,

219 (2017).

34. Jensen, C. & Teng, Y. Is It Time to Start Transitioning From 2D to 3D Cell Culture?

Front. Mol. Biosci. **7**, (2020).

35. Alkasalias, T., Moyano-Galceran, L., Arsenian-Henriksson, M. & Lehti, K.

Fibroblasts in the Tumor Microenvironment: Shield or Spear? *Int. J. Mol. Sci.* **19**,

1532 (2018).

36. Truffi, M., Sorrentino, L. & Corsi, F. Fibroblasts in the Tumor Microenvironment.

Adv. Exp. Med. Biol. **1234**, 15–29 (2020).

37. Monteran, L. & Erez, N. The Dark Side of Fibroblasts: Cancer-Associated

Fibroblasts as Mediators of Immunosuppression in the Tumor Microenvironment.

Front. Immunol. **10**, (2019).

38. Mao, X. *et al.* Crosstalk between cancer-associated fibroblasts and immune cells in

the tumor microenvironment: new findings and future perspectives. *Mol. Cancer* **20**,

131 (2021).

39. De Nola, R., Loizzi, V., Cicinelli, E. & Cormio, G. Dynamic crosstalk within the

tumor microenvironment of uterine cervical carcinoma: baseline network, iatrogenic

alterations, and translational implications. *Crit. Rev. Oncol. Hematol.* **162**, 103343

(2021).

40. Han, L. *et al.* Senescent Stromal Cells Promote Cancer Resistance through SIRT1

Loss-Potentiated Overproduction of Small Extracellular Vesicles. *Cancer Res.* **80**,

3383–3398 (2020).

41. Barros, M. R. *et al.* Activities of stromal and immune cells in HPV-related cancers.

J. Exp. Clin. Cancer Res. **37**, 137 (2018).

42. Raikhy, G. *et al.* Suppression of Stromal Interferon Signaling by Human

Papillomavirus 16. *J. Virol.* **93**, 10.1128/jvi.00458-19 (2019).

43. Chung, S.-H., Shin, M. K., Korach, K. S. & Lambert, P. F. Requirement for Stromal

Estrogen Receptor Alpha in Cervical Neoplasia. *Horm. Cancer* **4**, 50–59 (2013).

44. Martinkovich, S., Shah, D., Planey, S. L. & Arnott, J. A. Selective estrogen receptor

modulators: tissue specificity and clinical utility. *Clin. Interv. Aging* **9**, 1437–1452

(2014).

45.   Spurgeon, M. E., Chung, S.-H. & Lambert, P. F. Recurrence of cervical cancer in

mice after selective estrogen receptor modulator therapy. *Am. J. Pathol.* **184**, 530–

540 (2014).

46.   Munger, K. Are selective estrogen receptor modulators (SERMs) a therapeutic

option for HPV-associated cervical lesions and cancers? *Am. J. Pathol.* **184**, 358–

361 (2014).

47.   Castle, P. E. Do Selective Estrogen Receptor Modulators Treat Cervical Precancer

and Cancer? Time to Pool Data from Relevant Trials. *Int. J. Cancer J. Int. Cancer*

128, 997–998 (2011).

48.   Patel, H. K. & Bihani, T. Selective estrogen receptor modulators (SERMs) and

selective estrogen receptor degraders (SERDs) in cancer treatment. *Pharmacol.*

Ther. **186**, 1–24 (2018).

49.   Johnston, S. R. Endocrine manipulation in advanced breast cancer: recent advances

with SERM therapies. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **7**, 4376s–

4387s; discussion 4411s-4412s (2001).

50.   Moshi, M. R. *et al.* The Clinical Effectiveness of Denosumab (Prolia®) for the

Treatment of Osteoporosis in Postmenopausal Women, Compared to

Bisphosphonates, Selective Estrogen Receptor Modulators (SERM), and Placebo: A

Systematic Review and Network Meta-Analysis. *Calcif. Tissue Int.* **112**, 631–646

(2023).

51.   Sengupta, A., Dutta, S. & Mallick, R. Modulation of Cervical Carcinogenesis by

Tamoxifen in a Mouse Model System. *Oncology* **48**, 258–261 (2009).

52. Li, D. *et al.* Estrogen-Related Hormones Induce Apoptosis by Stabilizing Schlafen-

12 Protein Turnover. *Mol. Cell* **75**, 1103-1116.e9 (2019).

53. Liu, Y., Tian, L.-B., Yang, H.-Y. & Zhang, H.-P. Effects of estradiol and

progesterone on the growth of HeLa cervical cancer cells. *Eur. Rev. Med.*

Pharmacol. Sci. **21**, 3959–3965 (2017).

54. Tang, A. L. *et al.* UM-SCC-104: a new human papillomavirus-16 containing head

and neck squamous cell carcinoma cell line. *Head Neck* **34**, 1480–1491 (2012).

55. Bożyk, A., Wojas-Krawczyk, K., Krawczyk, P. & Milanowski, J. Tumor

Microenvironment—A Short Review of Cellular and Interaction Diversity. *Biology*

11, 929 (2022).

56. Jin, M.-Z. & Jin, W.-L. The updated landscape of tumor microenvironment and drug

repurposing. *Signal Transduct. Target. Ther.* **5**, 1–16 (2020).

57. Doorbar, J. *et al.* The biology and life-cycle of human papillomaviruses. *Vaccine* **30**

Suppl 5, F55-70 (2012).

58. Psyrri, A. & DiMaio, D. Human papillomavirus in cervical and head-and-neck

cancer. *Nat. Clin. Pract. Oncol.* **5**, 24–31 (2008).

59. Huh, W. K. *et al.* Final efficacy, immunogenicity, and safety analyses of a nine-

valent human papillomavirus vaccine in women aged 16–26 years: a randomised,

double-blind trial. *The Lancet* **390**, 2143–2159 (2017).

60. Kavanagh, K. *et al.* Changes in the prevalence of human papillomavirus following a

national bivalent human papillomavirus vaccination programme in Scotland: a 7-

year cross-sectional study. *Lancet Infect. Dis.* **17**, 1293–1302 (2017).

61. Meeting, I. W. G. on the E. of C. R. to H. & Cancer, I. A. for R. on. *Human*

Papillomaviruses. (World Health Organization, 2007).

62. Vats, A., Trejo-Cerro, O., Thomas, M. & Banks, L. Human papillomavirus E6 and

E7: What remains? *Tumour Virus Res.* **11**, 200213 (2021).

63. The Expression of HPV E6/E7 mRNA In Situ Hybridization in HP... : International

Journal of Gynecological Pathology.

https://journals.lww.com/intjgynpathology/Fulltext/2023/01000/The_Expression_of

_HPV_E6_E7_mRNA_In_Situ.2.aspx.

64. Basukala, O. & Banks, L. The Not-So-Good, the Bad and the Ugly: HPV E5, E6

and E7 Oncoproteins in the Orchestration of Carcinogenesis. *Viruses* **13**, 1892

(2021).

65. Comprehensive understanding of anchorage-independent survival and its

implication in cancer metastasis | Cell Death & Disease.

https://www.nature.com/articles/s41419-021-03890-7.

66. Kiyota, N. *et al.* Weekly Cisplatin Plus Radiation for Postoperative Head and Neck

Cancer (JCOG1008): A Multicenter, Noninferiority, Phase II/III Randomized

Controlled Trial. *J. Clin. Oncol.* **40**, 1980 (2022).

67. Helfenstein, S. *et al.* 3-weekly or weekly cisplatin concurrently with radiotherapy

for patients with squamous cell carcinoma of the head and neck – a multicentre,

retrospective analysis. *Radiat. Oncol.* **14**, 32 (2019).

68. Cheng, Y., Li, S., Gao, L., Zhi, K. & Ren, W. The Molecular Basis and Therapeutic

Aspects of Cisplatin Resistance in Oral Squamous Cell Carcinoma. *Front. Oncol.*

11, 761379 (2021).

69.   Rades, D. *et al.* Chemoradiation with Cisplatin vs. Carboplatin for Squamous Cell

772   Carcinoma of the Head and Neck (SCCHN). *Cancers* **15**, 3278 (2023).

773   70.   Gold, J. M. & Raja, A. Cisplatin. in *StatPearls* (StatPearls Publishing, Treasure

774   Island (FL), 2024).

775   71.   Ozols, R. F. *et al.* Phase III trial of carboplatin and paclitaxel compared with

776   cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a

777   Gynecologic Oncology Group study. *J. Clin. Oncol. Off. J. Am. Soc. Clin. Oncol.*

778   **21**, 3194–3200 (2003).

779   72.   Kondagunta, G. V. *et al.* Combination of paclitaxel, ifosfamide, and cisplatin is an

780   effective second-line therapy for patients with relapsed testicular germ cell tumors.

781   *J. Clin. Oncol. Off. J. Am. Soc. Clin. Oncol.* **23**, 6549–6555 (2005).

782   73.   Wang, H. *et al.* Cisplatin prevents breast cancer metastasis through blocking early

783   EMT and retards cancer growth together with paclitaxel. *Theranostics* **11**, 2442–

784   2459 (2021).

785   74.   Nguyen, V. T. *et al.* Dose-Intense Cisplatin-Based Neoadjuvant Chemotherapy

786   Increases Survival in Advanced Cervical Cancer: An Up-to-Date Meta-Analysis.

787   *Cancers* **14**, 842 (2022).

788   75.   Chapman, S., Liu, X., Meyers, C., Schlegel, R. & McBride, A. A. Human

789   keratinocytes are efficiently immortalized by a Rho kinase inhibitor. *J. Clin. Invest.*

790   **120**, 2619–2626 (2010).

791   76.   Warburton, A., Markowitz, T. E., Katz, J. P., Pipas, J. M. & McBride, A. A.

792   Recurrent integration of human papillomavirus genomes at transcriptional

793   regulatory hubs. *NPJ Genomic Med.* **6**, 101 (2021).

77.   Meyers, C. Organotypic (raft) epithelial tissue culture system for the differentiation-
dependent replication of papillomavirus. *Methods Cell Sci.* **18**, 201–210 (1996).

78.   Lambert, P. F. *et al.* Using an immortalized cell line to study the HPV life cycle in
organotypic 'raft' cultures. *Methods Mol. Med.* **119**, 141–155 (2005).

79.   Bienkowska-Haba, M. *et al.* A new cell culture model to genetically dissect the
complete human papillomavirus life cycle. *PLOS Pathog.* **14**, e1006846 (2018).

80.   Roberts, S., Evans, D., Mehanna, H. & Parish, J. L. Modelling human
papillomavirus biology in oropharyngeal keratinocytes. *Philos. Trans. R. Soc. Lond.*
B. Biol. Sci. **374**, 20180289 (2019).

81.   Stanley, M. A., Pett, M. R. & Coleman, N. HPV: from infection to cancer. *Biochem.*
Soc. Trans. **35**, 1456–1460 (2007).

82.   Warburton, A., Della Fera, A. N. & McBride, A. A. Dangerous Liaisons: Long-
Term Replication with an Extrachromosomal HPV Genome. *Viruses* **13**, 1846
(2021).

83.   Coursey, T. L., Van Doorslaer, K. & McBride, A. A. Regulation of Human
Papillomavirus 18 Genome Replication, Establishment, and Persistence by
Sequences in the Viral Upstream Regulatory Region. *J. Virol.* **95**, e0068621 (2021).

84.   Myers, J. E., Zwolinska, K., Sapp, M. J. & Scott, R. S. An Exonuclease V–qPCR
Assay to Analyze the State of the Human Papillomavirus 16 Genome in Cell Lines
and Tissues. *Curr. Protoc. Microbiol.* **59**, e119 (2020).

85.   James, C. D. *et al.* HPV16 genome structure analysis in oropharyngeal cancer PDXs
identifies tumors with integrated and episomal genomes. *Press* (2024).

86.   Thomas, M., Pim, D. & Banks, L. The role of the E6-p53 interaction in the

molecular pathogenesis of HPV. *Oncogene* **18**, 7690–7700 (1999).

87.   Li, S. *et al.* Ubiquitination of the HPV Oncoprotein E6 Is Critical for E6/E6AP-

Mediated p53 Degradation. *Front. Microbiol.* **10**, (2019).

88.   Scheffner, M., Huibregtse, J. M., Vierstra, R. D. & Howley, P. M. The HPV-16 E6

and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of

p53. *Cell* **75**, 495–505 (1993).

89.   Graham, S. V. Keratinocyte Differentiation-Dependent Human Papillomavirus Gene

Regulation. *Viruses* **9**, 245 (2017).

90.   Huibregtse, J. M., Scheffner, M. & Howley, P. M. A cellular protein mediates

association of p53 with the E6 oncoprotein of human papillomavirus types 16 or 18.

EMBO J. **10**, 4129–4135 (1991).

91.   Hoppe-Seyler, K., Bossler, F., Braun, J. A., Herrmann, A. L. & Hoppe-Seyler, F.

The HPV E6/E7 Oncogenes: Key Factors for Viral Carcinogenesis and Therapeutic

Targets. *Trends Microbiol.* **26**, 158–168 (2018).

92.   Sahebali, S. *et al.* Stromal issues in cervical cancer: a review of the role and

function of basement membrane, stroma, immune response and angiogenesis in

cervical cancer development. *Eur. J. Cancer Prev.* **19**, 204–215 (2010).

93.   Deng, H., Hillpot, E., Mondal, S., Khurana, K. K. & Woodworth, C. D. HPV16-

Immortalized Cells from Human Transformation Zone and Endocervix are More

Dysplastic than Ectocervical Cells in Organotypic Culture. *Sci. Rep.* **8**, 15402

(2018).

94. Dakic, A. *et al.* ROCK inhibitor reduces Myc-induced apoptosis and mediates
immortalization of human keratinocytes. *Oncotarget* **7**, 66740–66753 (2016).

95. Epithelial–Stromal Interactions Modulating Penetration of Matrigel Membranes by
HPV 16-Immortalized Keratinocytes. *J. Invest. Dermatol.* **109**, 619–625 (1997).

96. Waggoner, S. E. *et al.* Human Cervical Cells Immortalized in Vitro with Oncogenic
Human Papillomavirus DNA Differentiate Dysplastically in Viva.

97. Deng, H., Hillpot, E., Yeboah, P., Mondal, S. & Woodworth, C. D. Susceptibility of
epithelial cells cultured from different regions of human cervix to HPV16-induced
immortalization. *PLOS ONE* **13**, e0199761 (2018).

98. Moody, C. A. & Laimins, L. A. Human Papillomaviruses Activate the ATM DNA
Damage Pathway for Viral Genome Amplification upon Differentiation. *PLOS*
Pathog. **5**, e1000605 (2009).

99. Gautam, D. & Moody, C. A. Impact of the DNA Damage Response on Human
Papillomavirus Chromatin. *PLOS Pathog.* **12**, e1005613 (2016).

100. Anacker, D. C. & Moody, C. A. Modulation of the DNA damage response during
the life cycle of human papillomaviruses. *Virus Res.* **231**, 41–49 (2017).

101. Moody, C. A. Mechanisms by which HPV Induces a Replication Competent
Environment in Differentiating Keratinocytes. *Viruses* **9**, 261 (2017).

102. Mac, M. & Moody, C. A. Epigenetic Regulation of the Human Papillomavirus Life
Cycle. *Pathogens* **9**, 483 (2020).

103. Prati, B., Marangoni, B. & Boccardo, E. Human papillomavirus and genome
instability: from productive infection to cancer. *Clinics* **73**, e539s (2018).

104. McBride, A. A. & Warburton, A. The role of integration in oncogenic progression

of HPV-associated cancers. *PLoS Pathog.* **13**, e1006211 (2017).

105. Knipe, D. M. *et al.* Snapshots: chromatin control of viral infection. *Virology* **435**,

141–56 (2013).

106. Sakakibara, N., Mitra, R. & McBride, A. a. The papillomavirus E1 helicase

activates a cellular DNA damage response in viral replication foci. *J. Virol.* **85**,

8981–95 (2011).

107. Ouyang, C., Lu, G., He, W., Bay, B.-H. & Shen, H.-M. Post-translational

Modification in Control of SIRT1 Stability during DNA Damage Response. *Int. J.*

Biol. Sci. **18**, 2655–2669 (2022).

108. Flick, F. & Lüscher, B. Regulation of Sirtuin Function by Posttranslational

Modifications. *Front. Pharmacol.* **3**, 29 (2012).

109. Yi, J. & Luo, J. SIRT1 and p53, effect on cancer, senescence and beyond. *Biochim.*

Biophys. Acta **1804**, 1684–1689 (2010).

110. Lee, Y.-H., Kim, S.-J. & Surh, Y.-J. Role of Post-translational Modification of

Silent Mating Type Information Regulator 2 Homolog 1 in Cancer and Other

Disorders. *J. Cancer Prev.* **27**, 157–169 (2022).

111. Chung, S.-H. & Lambert, P. F. Prevention and treatment of cervical cancer in mice

using estrogen receptor antagonists. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 19467–

19472 (2009).

112. Shultz, L. D. *et al.* Human Lymphoid and Myeloid Cell Development in NOD/LtSz-

scid IL2Rγnull Mice Engrafted with Mobilized Human Hemopoietic Stem Cells 12.

J. Immunol. **174**, 6477–6489 (2005).

113. Saulters, E. L. *et al.* Differential Regulation of the STING Pathway in Human

Papillomavirus-Positive and -Negative Head and Neck Cancers. *Cancer Res.*

Commun. **4**, 118–133 (2024).

114. Smola, H. *et al.* Dynamics of Basement Membrane Formation by Keratinocyte–

Fibroblast Interactions in Organotypic Skin Culture. *Exp. Cell Res.* **239**, 399–410

(1998).

115. Bristol, M. L. *et al.* DNA Damage Reduces the Quality, but Not the Quantity of

Human Papillomavirus 16 E1 and E2 DNA Replication. *Viruses* **8**, 175 (2016).

116. James, C. D. *et al.* Human Papillomavirus 16 E6 and E7 Synergistically Repress

Innate Immune Gene Transcription. *mSphere* **5**, e00828-19 (2020).

117. James, C. D. *et al.* Restoring the DREAM Complex Inhibits the Proliferation of

High-Risk HPV Positive Human Cells. *Cancers* **13**, 489 (2021).

118. James, C. D. *et al.* SAMHD1 Regulates Human Papillomavirus 16-Induced Cell

Proliferation and Viral Replication during Differentiation of Keratinocytes. *mSphere*

4, e00448-19 (2019).

119. Gauson, E. J. *et al.* Failure to interact with Brd4 alters the ability of HPV16 E2 to

regulate host genome expression and cellular movement. *Virus Res.* **211**, 1–8

(2016).

120. Gauson, E. J. *et al.* Regulation of human genome expression and RNA splicing by

human papillomavirus 16 E2 protein. *Virology* **468–470**, 10–18 (2014).

121. Fontan, C. T. *et al.* A Critical Role for p53 during the HPV16 Life Cycle.

Microbiol. Spectr. **10**, e0068122 (2022).

122. Prabhakar, A. T. *et al.* Human Papillomavirus 16 E2 Interaction with TopBP1 Is

Required for E2 and Viral Genome Stability during the Viral Life Cycle. *J. Virol.*

97, e00063-23 (2023).

123. Bajikar, S. S. *et al.* Tumor-Suppressor Inactivation of GDF11 Occurs by Precursor

Sequestration in Triple-Negative Breast Cancer. *Dev. Cell* **43**, 418-435.e13 (2017).

124. Nagatani, M. *et al.* Comparison of biological features between severely immuno-

deficient NOD/Shi-*scid Il2rg^{null}* and NOD/LtSz-*scid Il2rg^{null}* mice. *Exp. Anim.* **68**,

471–482 (2019).

125. Sannigrahi, M. K. *et al.* HPV E6 regulates therapy responses in oropharyngeal

cancer by repressing the PGC-1α/ERRα axis. *JCI Insight* **7**, e159600.

126. Turk, A. N., Byer, S. J., Zinn, K. R. & Carroll, S. L. Orthotopic xenografting of

human luciferase-tagged malignant peripheral nerve sheath tumor cells for in vivo

testing of candidate therapeutic agents. *J. Vis. Exp. JoVE* 2558 (2011)

doi:10.3791/2558.

127. DeRose, Y. S. *et al.* Patient-derived models of human breast cancer: protocols for in

vitro and in vivo applications in tumor biology and translational medicine. *Curr.*

Protoc. Pharmacol. **Chapter 14**, Unit14.23 (2013).

128. Guillen, K. P. *et al.* A breast cancer patient-derived xenograft and organoid platform

for drug discovery and precision oncology. 2021.02.28.433268 Preprint at

https://doi.org/10.1101/2021.02.28.433268 (2021).

129. Animal Procedures and Information | Lab Manual and Procedures | Wicha Lab.

https://www.med.umich.edu/wicha-lab/labmanual6.html.

130. Whitfield, J., Littlewood, T. & Soucek, L. Tamoxifen Administration to Mice. *Cold*

Spring Harb. Protoc. **2015**, 269–271 (2015).

131. Caneparo, C., Baratange, C., Chabaud, S. & Bolduc, S. Conditioned medium

produced by fibroblasts cultured in low oxygen pressure allows the formation of

highly structured capillary-like networks in fibrin gels. *Sci. Rep.* **10**, 9291 (2020).

132. Ghani, N. 'Izzah A. *et al.* Effect of Different Collection Times of Dermal Fibroblast

Conditioned Medium (DFCM) on In Vitro Re-Epithelialisation Process.

Biomedicines **10**, (2022).

133. Halbert, C. L., Demers, G. W. & Galloway, D. A. The E7 gene of human

papillomavirus type 16 is sufficient for immortalization of human epithelial cells. *J.*

Virol. **65**, 473–478 (1991).

939
940
Figure Legends Figure

Figure 1: Estrogen fails to sensitize *in vivo*.

1A. HeLa cells are an integrated HPV18+ female cervical cancer cell line, we therefore

chose to utilize female NOD-scid IL2Rgnull (NSG) mice for this treatment paradigm; we

did not choose to ovariectomize these mice. Day 0 marks the date at which cells were

injected for xenografts. Tumors were palpable on day 10, mice were randomized.

Estrogen alone (E=0.4mg pellet + 8 µg/ml water supplementation *ad libitum*), radiation

alone (IR=10 Gy), as well as the combinational approach (E+IR) were monitored for

effects on tumor volume by calipers. 1B. Mice were monitored for weight throughout the

study. 1C. UMSCC104 cells are an episomal HPV16+ male oropharyngeal line, we

therefore chose to utilize male NSG mice for this treatment paradigm. Day 0 marks the

date at which cells were injected for xenografts; early estrogen water supplementation

began on this day, pellets were later added when tumors were palpable (8 µg/ml water

supplementation *ad libitum*). Tumors were palpable on day 7. Estrogen alone (0.4mg

pellet + 8 µg/ml water supplementation ad libitum), radiation alone (5 Gy), as well as the

combinational approach were monitored for effects on tumor volume by calipers. 1D.

Mice were monitored for weight throughout the study.

Figure 2: Fibroblasts significantly reduce HPV-specific estrogenic sensitization in

keratinocytes. (A) N/Tert-1, (B) HFK+E6E7, and (C) HFK+HPV16 cells were seeded

on day 0 and grown in the presence or absence of J2s that had been seeded at least 6

hours prior. Cells were washed to remove J2s in noted conditions, trypsinized, and

counted on day 3. *, $p < 0.05$; ***, $p < 0.001$.

Figure 3: Fibroblasts significantly reduce HPV-specific cancer estrogenic

sensitization. J2s were seeded in the morning and noted nuclear-labeled cancer cells

were seeded at least 6 hours after: HeLa (A,B), UMSCC47 (C,D), UMSCC104 (E,F),

HN30 (G,H). Co-culture images for quantitation were taken the following morning and

are set at day 0, estrogen (E) was added immediately after initial imaging on day 0. Cells

were again imaged at day 1 and day 3. UMSCC104 cells were grown for an additional

time point; these were replenished with new J2s and estrogen on day 3 (post-imaging)

and day 5 and imaged again on day 7. Within same graphs *$p<0.05$, ***$p<0.001$. J2s

altered growth rates for some of the cell lines and graphs are presented as separate for

visual simplicity, but experiments were run concurrently; comparing top and bottom

graphs \$p<0.05 J2 increased growth, &p<0.05 J2 decreased growth.

Figure 4: Human fibroblasts (HDFM) significantly reduce HPV-specific cancer

estrogenic sensitization. HeLa were seeded into standard media (A), J2 conditioned

media (B), or HDFMs were seeded in the morning and nuclear-labeled HeLa cells were

seeded at least 6 hours after (C). Co-culture images for quantitation were taken the

following morning and are set at day 0, estrogen (E) was added immediately after initial

imaging on day 0. Cells were again imaged at day 1 and day 3. *$p<0.05$

Figure 5: Fibroblasts do not alter cisplatin sensitivity. J2s were seeded in the morning

and noted nuclear-labeled cancer cells were seeded at least 6 hours after: HeLa (A,B),

UMSCC47 (C,D), UMSCC104 (E,F), HN30 (G,H). Co-culture images for quantitation

were taken the following morning and are set at day 0, noted concentrations of cisplatin

were added immediately after initial imaging on day 0. Cells were again imaged at day 1

and day 2. New J2s or media were replenished on day 3, and imaged again on day

5. Within same graphs ****$p<0.001$. J2s altered growth rates for some of the cell lines

and graphs are presented as separate for visual simplicity, but experiments were run

concurrently; comparing top and bottom graphs \$$p<0.05$ J2 increased growth, &$p<0.05$

J2 decreased growth.

Figure 6: Stroma supports transcriptional regulation, viral protein expression, and

episomal maintenance; estrogenic transcriptional regulation is lost with stroma. (A)

N/Tert-1 cells were transfected with 1 µg of pgl3 basic backbone, 1 µg of pgl3 control

(positive control), or 1 µg LCR and grown in the presence or absence of 15 µM estrogen

and/or J2s that had previously been seeded. Forty-eight hours after transfection, a

luciferase-based assay was utilized to monitor levels of LCR transcription. Data are

presented as relative fluorescence units (RFU), normalized to total protein concentration

as monitored by a standard bovine serum albumin (BSA) assay. ANOVA **, $P < 0.01$;

***, $P < 0.001$ (B) N/Tert-1 cells were grown in the presence or absence of J2s. Cells

were washed to remove J2, then lysed and analyzed via western blotting for ERα.

GAPDH was used as a loading control. (C) HFK$^+$HPV16 cells were seeded on day 0 and

grown in the presence or absence of J2s for 1 week. Cells were washed to removed J2,

then lysed and analyzed for DNA expression of E2 and E6 via the exonuclease V assay,

in comparison to GAPDH and mitochondrial DNA controls. Results are presented as

percent integration as calculated from the cut ratio of matched GAPDH. *$P < 0.05$. (D)

HFK+E6E7 (lanes 2,3) and HFK$^+$HPV16 (lanes 4,5) and N/Tert-1 (lanes 6,7) cells were

seeded on day 0 and grown in the presence or absence of J2s for 1 week. Lane 1 is an

input lysate from J2 alone to control for any background level of expression in fibroblasts

that were not removed via washing. Cells were washed to remove J2s in noted conditions,

trypsinized, lysed, and analyzed via western blotting for SIRT1, pRb, p53, γH2aX, and

E7. GAPDH and β-actin were utilized as loading controls.

1016

Figure 7: Fibroblast co-culture demonstrates SERMs are worth assessing in an

HPV+ *in vivo* model. J2s were seeded in the morning and noted nuclear-labeled cancer

cells were seeded at least 6 hours after: HeLa (A,B), UMSCC47 (C,D), UMSCC104

(E,F). Co-culture images for quantitation were taken the following morning and are set at

day 0, 10 μg raloxifene (R) or 10 μg tamoxifen (T) were added immediately after initial

imaging on day 0. Cells were again imaged at day 1 and day 3. UMSCC104 cells were

grown for an additional time point; these were replenished with new J2s and drugs on day

3 (post-imaging) and day 5 and imaged again on day 7. Within same graphs *p<0.05,

***p<0.001. J2s altered growth rates for some of the cell lines and graphs are presented

as separate for visual simplicity, but experiments were run concurrently.

1027

Figure 8: Fibroblast *in vitro* co-culture SERM assessment in UMSCC104 cells

predicted the utility *in vivo*. As described in Figure 1, UMSCC104 cells were injected

for xenografts in male NOD-scid IL2Rgnull (NSG) mice. Day 0 marks the date at which

cells were injected for xenografts. (A,B) Tumors were palpable on day 7. 1-Control, 2-

Raloxifene alone (1.5 mg), 3-Tamoxifen alone (1.0 mg), 4-radiation alone (4 Gy) (IR), as

well as the combinational approaches (5-Raloxifene+IR, 6-Tamoxifen+IR) were

monitored for effects on tumor volume by calipers. These experiments were run

concurrently but presented on separate graphs for visual clarity (groups have been

numbered for this clarity). Data points shown for Mean Tumor Volume are representative

of conditions with at least 70% of animals remaining. (C) Kaplan-Meier Survival plots

the survival curve of the animals treated. This data includes any mice that needed to be

sacrificed due to tumor ulceration. (D) Mice were monitored for weight throughout the

study. *p<0.05, **p<0.01 ***p<0.001 from control, $p<0.05 from IR alone, colors of *

or $ are matched to colors of corresponding conditions that are significant from control or

radiation alone, respectively.

1043

1044 **Figure 9. Representative cell culture images quantified via Keyence.** Images

1045 presented are day 5 of UMSCC104 cell cultures in the presence or absence of J2s. Images

1046 were captured at 10x in brightfield and TexasRed. All cell lines at all time points were

1047 counted utilizing the same control-established automated parameters to ensure

1048 reproducibility.

A.

B.

C.

D.

A.

B.

C.

A. HeLa

B. HeLa +J2

C. UMSCC47

D. UMSCC47 +J2

E. UMSCC104

F. UMSCC104 +J2

G. HN30

H. HN30 +J2

A.

B.

C.

D.

E.

F.

A.

UMSCC104
Mean Tumor Volume ±SEM

B.

UMSCC104
Mean Tumor Volume ±SEM

C.

Kaplan-Meier Survival Plot

D.

UMSCC104
Mean Body Weight ±SEM

Traditional Cell Culture
UMSCC104

Fibroblast Co-Culture
UMSCC104 + J2

mKate$_2$-red

brightfield

merged