RSL4 regulatory network acts as an early driver of root hair growth at low temperature in *Arabidopsis thaliana*

Tomás Urzúa Lehuedé[1,2,3,*], Victoria Berdion Gabarain[4,*], Tomas Moyano[1,3], Lucia Ferrero[5], Miguel Angel Ibeas[1,2,3], Hernan Salinas-Grenet[1,3], Gerardo Núñez-Lillo[6], Romina Acha[1,2], Jorge Perez[1,2], Florencia Perotti[5], Virginia Natali Miguel[5], Fiorella Paola Spies[5], Miguel A. Rosas[7], Michitaro Shibata[8], Diana R. Rodríguez-García[4], Adrian A. Moreno[1], Keiko Sugimoto[8], Karen Sanguinet[7], Claudio Meneses[2,9], Raquel L. Chan[5], Federico Ariel[5], Jose M. Alvarez[1,3], José M. Estevez[1,2,3,4,†]

[1]Centro de Biotecnología Vegetal, Facultad de Ciencias de la Vida, Universidad Andres Bello, Santiago, Chile.

[2]ANID - Millennium Science Initiative Program - Millennium Nucleus for the Development of Super Adaptable Plants (MN-SAP), Santiago, Chile.

[3]ANID - Millennium Science Initiative Program - Millennium Institute for Integrative Biology (iBio) Santiago, Chile.

[4]Fundación Instituto Leloir and IIBBA-CONICET. Av. Patricias Argentinas 435, Buenos Aires C1405BWE, Argentina.

[5]Instituto de Agrobiotecnología del Litoral, Universidad Nacional del Litoral-CONICET, Facultad de Bioquímica y Ciencias Biológicas, CCT Santa Fe, Colectora Ruta Nacional 168 km 0, Santa Fe 3000, Argentina.

[6]Escuela de Agronomía, Facultad de Ciencias Agronómicas y de los Alimentos, Pontificia Universidad Católica de Valparaíso, Calle San Francisco s/n, La Palma, Quillota 2260000, Chile.

[7]Department of Crop and Soil Sciences and Molecular Plant Sciences Graduate Program, Washington State University, Pullman, WA, 99164, USA.

[8]RIKEN Center for Sustainable Resource Science, Yokohama, 230-0045 and Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.

[9]Departamento de Fruticultura y Enología, Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Santiago, Chile; Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile; Fondo de Desarrollo de Áreas Prioritarias, Center for Genome Regulation, Santiago, Chile.

*Co-first authors.

†Correspondence should be addressed. Email: jose.estevez@unab.cl (J.M.E).

Abstract

The root hair (RH) cells can elongate to several hundred times their initial size, and are

an ideal model system for investigating cell size control. Their development is

influenced by both endogenous and external signals, which are combined to form a

integrative response. Surprisingly, a low temperature condition of 10°C causes an

increased RH growth in *Arabidopsis* and in several monocots, even when the

development of the rest of the root and aerial parts of the plant are halted. Previously,

we demonstrated a strong correlation between the growth response and a significant

decrease in nutrient availability in the medium under low temperature conditions.

However, the molecular basis responsible for receiving and transmitting signals related

to the availability of nutrients in the soil, and their relation to plant development,

remain largely unknown. We decided to further investigate the intricate molecular

processes behind the particular responsiveness of this root cell type at low temperature.

In this study, we have discovered a gene regulatory network (GRN) controlling early

transcriptome responses to low temperature. This GNR is commanded by specific

transcription factors (FTs), namely *ROOT HAIR DEFECTIVE 6-LIKE 4* (RSL4), a member of

the homeodomain leucine zipper (HD-Zip I) group I 13 (AtHB13), the trihelix TF *GT2-*

LIKE1 (GTL1), and a previously unidentified MYB-like TF (AT2G01060). Furthermore, we

have identified four downstream RSL4 targets *AtHB16*, *AtHB23*, *EARLY-RESPONSIVE TO*

DEHYDRATION 7 (ERD7) and ERD10 suggesting their participation in the regulation of

RH development under these conditions. Functional analysis shows that such

components of the RSL4-dependent transcriptional cascade influence the subsequent

RH growth response to low temperature. These discoveries enhance our

comprehension of how plants synchronize the RH growth in response to variations in

temperature and nutrient availability at the cellular level.

Abstract Word counts 285

Text Word counts 5,944

Figures 1-5

Figure S1-S17

References 70

Introduction

Root hairs (RHs) are single cells that develop as a tubular protrusion from the root epidermis in a polar manner up to several hundred times their original size. RH differentiation is a complex process and consists of 3 stages: differentiation, initiation and elongation. The first is controlled by a developmental program involving transcription factor complexes (TFs) allowing or repressing the expression of the GL2 (GLABRA2) protein. This protein blocks RH development (Lin *et al.,* 2015) by inhibiting the transcription of the RHD6 (ROOT HAIR DEFECTIVE 6) regulator, which together with RSL1 can induce the expression of other factors such as RSL2 (Root Hair Defective 6 like 2) and RSL4 (ROOT HAIR DEFECTIVE 6 LIKE 4), thus allowing RH growth (Menand *et al.,* 2007; Yi *et al.,* 2010; Datta *et al.,* 2015; Proust *et al.,* 2016; Vijayakumar *et al.,* 2016; Lopez *et al.,* 2023). RHs are able to grow within hours in order to reach water-soluble nutrients in the soil, to promote interactions with the local microbiome, and to support the anchoring of the root. The overall fitness of plants is reduced in various loss-of-function RH mutants that lack RHs when grown in challenging soil conditions (Wu *et al.,* 2007, Haling *et al.,* 2013, Miguel *et al*., 2015). On the other hand, genotypes with longer RHs, achieved through the overexpression of ROOT HAIR DEFECTIVE 6-LIKE transcription factors (RSLs), have shown beneficial effects in species such as wheat (Han *et al.*, 2016, 2017), rice (Moon *et al*., 2019), and Brachypodium (Kim & Dolan, 2016). This highlights the deep importance of RHs in relation to plant development and physiology, despite not always receiving full recognition. It is known that intrinsic factors regulate RH elongation at the cellular level and external signals, among which we find morphogenetic programming, nutrient availability, and the action of different hormones (Casal & Estevez, 2021; Lopez *et al.,* 2023). Several works previously showed that RH growth is highly responsive to increasing nutrient concentrations in the media (Yi *et al.,* 2010; Bhosale *et al.,* 2018; Moison *et al.,* 2021; Jia *et al.,* 2023). While high nutrient concentrations impair RH development and growth, low concentrations of nutrients such as inorganic phosphate (Pi) and/or nitrates trigger a strong growth response mediated by the phytohormone auxin (Bhosale *et al.,* 2018; Mangano *et al.,* 2017; Jia *et al.,* 2023; Lopez *et al.,* 2023).

Recently it was shown that low temperature (10°C) is able to enhance RH growth possibly as a result of the restriction of nutrient mobility and accessibility by the root (Moison *et al.,* 2021; Martínez-Pacheco *et al.,* 2021). Conversely, an increase in agar concentration in the medium that restrains nutrient mobility resulted in the reversion of low temperature-induced RH elongation (Moison *et al.,* 2021). Altogether, these observations suggested that low temperatures restrict nutrient mobility and availability in the culture medium, leading to the promotion of polar RH growth. Despite these previous findings, the signals that trigger the RH cell elongation under these conditions are still unknown, but it is presumed that RHs could be highly susceptible to

environmental stresses such as temperature changes, triggering differences at the transcriptomic and proteomic levels. Specifically, a novel ribonucleoprotein complex composed by the lncRNA AUXIN-REGULATED PROMOTER LOOP (APOLO) and the TF WRKY42 forms a regulatory hub to activate RHD6 by shaping its epigenetic environment and integrating low temperature signals governing RH growth (Moison *et al.,* 2021; Martínez-Pacheco *et al.,* 2021). Recently, new molecular components related to cell wall-apoplastic related peroxidases, PRX62 and PRX69, were identified as important factors for low temperature triggered RH growth by inducing changes in ROS homeostasis and cell wall glycoprotein EXTENSIN insolubilization (Martinez-Pacheco *et al.,* 2022). An autocrine regulation pathway between Rapid Alkalinization Factor 1 (RALF1) and the *Catharanthus roseus* RLK1-like (CrRLK1L) FERONIA (FER) receptor phosphorylates the early translation factor eIF4E1 and produces RH-related proteins including RSL4 to promote RH formation (Zhu *et al.,* 2020a,b). Finally, at low-temperature, RHs showed a physical link between FER and TARGET OF RAPAMYCIN COMPLEX (TORC). Mutants for *fer-4* and TORC components responded differently to low nitrate than Col-0, and RHs remained stunted throughout the bulge stage (Martinez-Pacheco *et al.,* 2023a). This FER-TORC pathway connects nutrients perception at the cell surface with downstream responses in RH growth at low temperatures (Martinez-Pacheco *et al.,* 2022). Although advances were achieved in our understanding on how RH growth occurs at low temperature, the dynamic transcriptional cascade controlling RH growth remains largely unknown, providing us with the opportunity to find new key regulators in the process of RH elongation. Here, we provide a deep characterization of how the TF RSL4 and main transcriptional network nodes AtHB13, GTL1/DF1 and MYB-like as well as the downstream selected components AtHB16/AtHB23 and ERD10/ERD7 are able to direct at early time points the gene expression program that allows RH growth to respond to the low temperature. We functionally validated these transcriptional nodes by gene expression assay, phenotypic characterization of mutant/overexpressor and evaluation of genetic interactions. These findings improve our understanding of how plants coordinate the development of their RHs in response to changes in temperature and nutrient availability at the cellular level.

Results

Low temperature effect on RH growth depends on nutrient availability. Root hair (RH) growth is a well conserved adaptive response to external low nutrients in the media-soil. Previously, it was shown that at low temperature growth conditions (10°C), nutrients in the media are reduced, which triggers a strong response in RH growth (Moison *et al.,* 2021; Martinez-Pacheco *et al.,* 2021). Then, we assessed the effect of the low temperature on RH growth, through a detailed temporal characterization of root development (**Figure 1A**). Starting at 2 days of 10ºC, a significant effect on RH

elongation compared to control conditions was detected. This induction of growth reached its peak at 3 days of low temperature treatment. Then, we decided to test if small temperature changes around 10°C also trigger a similar RH growth effect. We performed a temperature gradient from 22°C to 4°C in 2°C steps and maximum growth was detected at 8-10°C (**Figure 1B**). From 8°C down to 4°C, the low temperature had a negative effect on RH growth suggesting that the low temperature itself is more deleterious than the linked low-nutrient effect. Based on this, we tested if a medium with no nutrient content is able to repress the low temperature induced growth effect on RHs. To this end, we used agarose that lacks any detectable nutrients and salts and tested the low temperature growth effect (**Figure 1C**). In agreement with previous evidence, when no nutrient is in the medium no low temperature effect is observed on the RH growth, confirming that the signal is nutritional and not related to the temperature itself as suggested before (Moison *et al.*, 2021; Martinez-Pacheco *et al.*, 2021; Martinez-Pacheco *et al*., 2023). Our data allowed us to use the low temperature effect on RHs as a proxy of complex nutritional signals changes coming from the medium. When we evaluated the effect of different pretreatment growth times, it was observed that the final RH lengths in plants exposed to low temperatures were equivalent whether plants were pre-grown for 7 or 14 days at 22°C, which was also the case for their basal RH lengths (**Figure S1**). Since root light exposure generates a stress that affects growth, hormonal signaling, abiotic responses, or nutrient starvation adaptation compared with roots grown in the dark (Silva-Navas *et al.,* 2016), we decided to test if direct light exposure might affect the RH growth intensity recorded before. For this purpose, we used the system called D-Roots originally developed to cultivate roots in darkness, including dark agar plugs and black-colored vertical plates (Silva-Navas *et al.,* 2015). Using this D-root system, we have confirmed the previously measured RH phenotypes at low temperature although with slightly higher magnitude (**Figure 1C**). These results together showed that 8-10°C were the main temperature trigger of RH growth affecting nutrient accessibility to the root surfaces and light exposure does not significantly affect this growth process in RHs. Finally, we tested if this enhancement on RH growth at low temperature is also present in monocots (**Figure 1D**) including *Brachypodium distachyon* (Bd21), rice (Kitaake), and wheat (Chinese Spring). In these three cases we observed the same effect on RH growth at low temperature as in Arabidopsis, although in these cases we tested them using 16 h:8 h day:dark cycle (see methods). This last result highlighted that low temperature effect on RH growth might be a broader response in plant roots, also applicable to these monocots also.

Genome wide transcriptional changes in roots exposed to low temperature. RSL4 and RSL2 have been identified as important regulators of RH growth in response both at 22°C and at low temperature, although RSL4 has been shown to be of higher relevance than RSL2 in this role (Yi *et al.*, 2010; Moison *et al.,* 2021). Therefore, the loss of

function of RSL4, and RSL2 to a lower extent, would lead to changes in the transcript abundance of direct and indirect RH gene targets. To identify all genes regulated by the transcription factor (FT) RSL4 in the context of low temperature response, and to find molecular factors involved in RH development that are dependent on RSL4, RSL2, or both, we analyzed the transcriptome of *rsl2*, *rsl4*, and *rsl2 rsl4* genotypes in response to low temperature treatments (2h and 6h) and compared it with the transcriptome of Wt ecotype Columbia-0 (Col-0) plants (**Figure 2A**). Differential expression analysis was performed and the number of genes was filtered considering an FC>|0.5| in at least one condition. A total of 6,756 differentially expressed genes (DEG) were identified between Col-0 and any of the mutant genotypes (**Figure S2** and **Figure S3**). The heatmap in **Figure S2** shows the total number of DEGs identified, which were clustered into 5 groups. Cluster 5 composed of 1,157 genes, which are positively regulated by RSL2 and RSL4, is enriched with several GO terms related to RH development. To identify genes whose expression depends on the interaction between low temperature and the genotypes, we performed a two-way analysis of variance (ANOVA) analysis as described previously (Alvarez *et al.,* 2014), considering temperature (T) and each genotype (G). This analysis allows us to uncover genes for the interaction of TxG factors whose response to temperature is affected in the mutants analyzed. Our analysis identified 2,365 genes with a significant TxG factor for the *rsl4* mutant (**Figure 2B**). These results were consistent with RSL4 being more influential in the RH response to low temperature as compared to RSL2 (Moison *et al.*, 2021) in agreement with previous evidence that show that RSL4 levels define the final RH length (Datta *et al.,* 2016). By analyzing by heat map these 2,365 DEG, we identified three main clusters regulated by RSL4 and also affected by low temperature (**Figure 2B**). GO analysis revealed several processes related to RNA, virus replication and cold. In addition, the expression in trichoblast and RH cells of RSL4 (and the related RHD6) were increased at early time points of low temperature treatment while RSL2 was decreased (**Figure S4**). All this evidence together indicates that the TF RSL4 is a major regulator of RH growth at low temperature.

RSL4 controls low temperature response through AtHB13-GTL1-MYB like transcription factors. To get insight into the regulatory cascade triggered by RSL4, we focused on the 2,365 genes that showed an altered response to low temperature in the *rsl4* mutant (significant TxG factor). To investigate which TF might mediate the RSL4 effect of gene expression in response to low temperature, we exploited the TF-target data allocated in the ConnecTF database (Brooks *et al.,* 2021). The ConnecTF platform provides access to a library of experimentally confirmed TF–target gene interaction datasets. To this end, we used the list of differentially expressed TFs that depend on RSL4 and checked which TF has available validated *in vitro* TF-binding data obtained by DNA Affinity Purification sequencing (DAP-seq) (O'Malley *et al.*, 2016). We then reduced the list of possible target genes to those genes highly or uniquely expressed in RH cells previously

identified by single cell RNA-seq or GFP-enriched protoplast studies (Brady *et al*., 2007; Denyer *et al*., 2019; Jean-Baptiste *et al.*, 2019; Ryu *et al.*, 2019; Shulse *et al.*, 2019; Zhang *et al.*, 2019) (**Figure 2C**). Then, we selected the top three TFs of the best five-ranked list: (i) AtHB13, a class I HD-Zip TF previously described as involved in low temperature response but not related to RH growth (Cabello *et al.,* 2012), (ii) a MYB-like (AT2G01060) and (iii) GTL1, a previously characterized negative regulator of RH growth under high nutrient condition, and its homologous DF1 (Shibata *et al.*, 2018, 2022). Based on this analysis, a RH gene regulatory network composed of 110 genes controlled by RSL4 at low temperature was proposed (**Figure 2D**). Downstream of these main regulators, we selected several members of the Arabidopsis HD-Zip I subfamily, including AtHB16/AtHB23 and two TFs *EARLY-RESPONSIVE TO DEHYDRATION* ERD7/ERD10 for further characterization. Notably, none of these TFs (unless RSL4) were previously linked to RH growth at low temperature. This emphasizes the prognostic capability of these integrated methodologies.

The RSL4-dependent regulatory network regulates RH growth at low temperature. To validate the function of these RSL4-regulated gene regulatory networks in RH growth at low temperature, we performed detailed RH phenotype screenings at 22°C and at 10°C in loss of function mutants and overexpression lines for each component of the primary and secondary regulatory network nodes as well as the upstream RSL4-regulator RHD6 (**Figures S5-S13** and **Figure 3A**). These include the TFs RSL4, AtHB13, GTL1/DF1 and MYB-like as well as the downstream selected components AtHB16/AtHB23 and ERD7/ERD10 (**Table S1**) (**Figure 3A-B**). Not surprisingly, several of the TFs analyzed exhibited a changed RH growth at 10°C (**Figures S5-S13** and **Figure 3A**) providing strong evidence of the predictive power of this combined bioinformatics-experimental approach to identify key roles of specific TFs in certain environmental conditions. According to our functional screening, the main nodes RSL4 (and RSL2) and the downstream gene AtHB16 are positive regulators of RH growth at low temperature while, on the other hand, GTL1-DF1 and AtHB3,13,20,23 emerged as negative RH growth regulators at low temperature. A higher order of mutants for multiple HBs as the quadruple mutant *athb3 (hat7) athb13 athb20 athb23* showed an enhanced RH long phenotype at low temperature while single mutants *athb13* and *athb23* alone did not, suggesting a high level of functional redundancy of these four HBs in RH growth at low temperature. Surprisingly, *athb16* has a very short RH phenotype at low temperature. This may indicate at least two contrasting roles for these HBs (AtHB16 vs AtHB3,13,20,23) in RH growth at low temperature. To support the role of each TF on the primary network on RH growth, we have quantified the expression levels of the network's main nodes (*RSL4, AtHB13, GTL1/DF1* and *MYB-like*) at early temporal points (0h (Basal, B), 2h, and 6h at 10°C) in selected mutant lines previously isolated and characterized (**Figure S14-17**) (**Figure 3A-B**). In this manner, we are able to measure the impact and influence of each of the main regulatory nodes identified in our network

over the expression of the selected gene from the proposed GRN. We found that RSL4-GTL1 and GTL1-AtHB13 TFs form positive transcriptional feedback loops while RSL4-AtHB13 forms a negative feedback loop (**Figure 3C-D**). This highlights a complex transcriptional landscape behind a coherent phenotypic response, in this case at low temperature even for single plant cells. To determine whether RSL4 controls the expression of these main gene regulatory network nodes by direct binding to their promoter regions, we searched for putative RSL4-binding sites (Hwang *et al.,* 2017) in open chromatin regions, according to publicly available ATAC-seq datasets (Maher *et al.,* 2018). According to ChIP-qPCR using RSL4:RSL4-GFP plants and anti-GFP antibody, GTL1, AtHB13 and MYB-like are direct RSL4 targets in the primary gene network and AtHB16/AtHB23-ERD10 are their targets in the secondary network (**Figure 4A**), as revealed in comparison to the previously identified direct target EXPANSIN 7 (EXP7) (Hwang *et al.,* 2017). Overall, these results confirm that RSL4 positively controls the expression of GTL1 and represses the expression of AtHB13 thus impacting RH growth. It was also tested that presence of the GTL1 binding sites in the selected gene network by ChIP-seq and positive peaks indicate direct regulation close to the starting transcriptional sites for AtHB16 and ERD7/ERD10 only at room temperature suggestion that the low temperature repressed its activity in these genes (**Figure 4B**). Finally, we measured the expression levels on protein translational reporters for RSL4:RSL4-GFP (and the related RHD6:RHD6-GFP), as well as the node GTL1:GTL1-GFP (and the related DF1:DF1-GFP) (**Figure 4C-D** and **Figure S18**) both at room and at low temperature. All of them except RSL4 were induced at low temperature in epidermal cells close to the RH development and also in the differentiation zone. In addition, GUS staining of AtHB13, AtHB16 and AtHB23 driven lines were also characterized (**Figure 4E**) and a strong signal was detected for AtHB16, and to a lower extent, in AtHB23 in both low temperature and room temperature. In sum, our systems biology approach identified new components and relevant TF–target gene interactions that control RH growth in responses to moderate cold. Our results indicate that RSL4 initiates an interconnected transcriptional cascade by regulating GTL1, AtHB13, MYB-like and AtHB16/AtHB23 and ERD10 as downstream gene targets. Consequently, GTL1 directly controls the expression of RSL4 in a feedback loop and downstream moderate cold-responsive genes (**Figure 5**). This last result in HBs mutants and reporters lines highlights the major role of AtHB16 in controlling RH growth at low temperature. Overall, we were able to experimentally confirm the function of these new regulatory components in low temperature-controlled RH growth in Arabidopsis, establishing a complex regulatory circuitry that integrates an environmental signal with RH development.

Discussion

In this work, we have extended the RH growth responses at low temperature to three
monocots in addition to *Arabidopsis thaliana* suggesting a more general growth
response than anticipated. We uncovered a regulatory map of RH growth under low
temperature in Arabidopsis by focusing on early time points (hours). We have
uncovered a new gene regulatory network controlled by RSL4 under low temperature
conditions that mediates RH growth using an RNA-seq filtered by DAP-seq and validated
expression profiling using mutant lines, coupled to ChIP approaches. Under the control
of RSL4 there are several TFs in the main network nodes such as AtHB13, GTL1/DF1 and
MYB-like as well as the downstream selected components AtHB16/AtHB23 and
ERD7/ERD10 (**Figure 5**). None of them were previously associated with low
temperature (unless HB13 and ERD7) and related to RH growth highlighting the
predictive power of this approach. Previously, based on single-cell RNA sequencing, it
was identified several TFs similar to our gene regulatory network (GRN) such as HB3
(HAT7) and GTL1 as responsive to brassinosteroids in the cortex during cell elongation
(Nolan et al. 2023). This suggests that similar GRN might be sub-functionalized in
different cellular contexts and under different environmental and hormonal cues. One
of the negative regulator of RH growth identified here is the trihelix TF GT2-LIKE1 (GTL1)
and its closest homolog, DF1, which stops RH cell growth by directly repressing RSL4 but
also bind to RHD6 preventing the activation of RSL4 (Shibata *et al.*, 2022) under excess
nutrient conditions (Shibata *et al.*, 2018; Shibata *et al.* 2022) but it was not linked to low
temperature conditions. In our work, we found that GTL1 is positively controlled by
RSL4 and negatively regulates RH growth at low temperature with a similar function
described before linked to the nutrient availability and status present in the media.

Several members of the Arabidopsis HD-Zip I subfamily, AtHB13, AtHB16 and AtHB23,
were identified in this study as relevant TFs in the regulation of RH growth at moderate
low temperature (**Figure 5**). They belong to clade V (Arce *et al.,* 2011). They have a
leucine zipper (Zip) downstream the homeodomain (HD), located in the middle of the
protein (Viola *et al.,* 2016). Arabidopsis HD-Zip I members were classified based on their
exon-intron structure (Henriksson *et al.,* 2005) and subsequently based on
uncharacterized conserved motifs outside the HD-Zip (Arce *et al.,* 2011). HD-Zip I
proteins are only found in plants and have been linked to environmental stresses
(Perotti *et al.,* 2017) but also to many developmental processes (Ribone *et al.,* 2015;
Capella *et al.,* 2015; Perotti *et al.,* 2021; Perotti et al. 2022). AtHB13 provides enhanced
tolerance to both biotic and abiotic stresses, plays a role in pollen hydration and seed
germination (Hanson *et al.,* 2002; Cabello *et al.*, 2012; Gao et al. 2014; Silva *et al.,* 2016)
while AtHB23 participates in the elongation of the hypocotyl and the expansion of
cotyledons (Choi *et al.,* 2014), with functions in lateral root development (Perotti *et al.,*
2019; Perotti *et al.,* 2020; Perotti et al. 2021), together with PHL1 promotes
carbohydrate transport from pedicel-silique nodes to seeds (Spies *et al.,* 2022), and is a
developmental modulator of root growth in normal and salinity conditions,   by

interacting with PHL1 and MYB68 (Spies et al., 2023). Here, we found that AtHB13 is negatively regulated by RSL4-GTL1 and positively regulated by MYB-like at low temperature and participates in RH growth. It is plausible that AtHB13 do not act alone since in the RSL4-regulated RH network at low temperature were also identified several more HD-Zip I proteins. It remains to be tested if the contrasting growth roles of AtHB16 and AtHB3,13,20,23 in the RH response to low temperature are conducted as protein complexes or as individual TFs.

In addition, in this work, we uncovered one uncharacterized MYB-like TF as part of the GRN that controls RH growth at low temperature. The MYB family, one of the biggest TF groups in seed plants, consists of 197 members and this group is characterized by the presence of a DNA-binding domain called the MYB DNA-binding domain (Riechmann *et al.,* 2000) with a minimum of four repetitions of around 52 amino acids, which assemble into a helix-turn-helix shape. The MYB-CC family consists of 15 members, which are distinguished by the presence of both the MYB domain and a coiled-coil (CC) domain (Rubio *et al.,* 2001). Here we identified a MYB-like TF (AT2G01060), currently not well characterized, that regulates RH growth at low temperature. Finally, downstream RSL4-GTL1 regulators, we have identified ERD7 and ERD10 proteins that belong to a small family of highly conserved proteins that contain a PAM2 motif that may interact with poly(A)-binding protein (Aalto et al., 2012). ERD7 is strongly upregulated by both biotic and abiotic stresses such as cold, salt, excess light and Pi starvation (Cheng et al., 2013; Rasmussen et al., 2013) and, it was found to be associated with lipids droplets via its C-terminal senescence domain (SD) (Doner et al. 2021). Specifically, it binds to negatively charged phospholipids, such as phosphatidylinositol (PI) and phosphatidic acid (PA) that could induce structural and mechanical changes in the membrane that affect membrane fluidity (Barajas-Lopez et al. 2021). In particular and in agreement with our study, ERD7 is associated with low temperature stress. Overall, although our low temperature condition is moderate, ERD7 and, possibly ERD10, might play a protective role in the membrane stability during the RH cell expansion process.

Through our investigation of gene-expression patterns, we were able to uncover potential TFs that potentially control RH growth responses to low temperature. The incorporation of RH specific genes coupled to an early temporal data has a significant favorable effect on the accuracy of our network predictions. Furthermore, our findings served to identify novel TFs participating in the low temperature response triggering RH growth. This highlights the significance of using spatiotemporal regulatory networks when investigating specific biological inquiries. Our study enabled us to ascertain that genes sensitive to low temperature and their associated biological processes are regulated in a coordinated manner in both spatial and temporal dimensions. Together with RSL4, the TFs AtHB3,13,20,23, AtHB16 and GTL1/DF1, have significant functions in controlling the low temperature response in RH growth and the downstream changes in gene expression eventually result in the enhancement of RH growth. These discoveries

enhance our comprehension of how plants synchronize the RH growth in response to
variations in temperature and nutrient availability at the cellular level. Future
challenges are related to the definition of the fine tune mechanisms of all these TFs
acting in real time in the control of RH genes. As a long term goal, these new RH
regulators identified here could be used to generate crops with enhanced RH growth to
tailored nutrient uptake to specific soil and temperature conditions.

Materials and methods

Plant Material and Growth Conditions. All the *Arabidopsis thaliana* lines used were in
the Columbia-0 (Col-0) background. All the plant lines sussed in this study are detailed
(**Table S1**). Seeds were surface sterilized and stratified in darkness at 4°C for 3 days
before being germinated on ½ strength 0,5X MS-MES (Duchefa, Netherlands) on 0.8%
Plant Agar™ (Duchefa, Netherlands) on 120 x 120 mm square petri dishes (Deltalab,
Spain) in a plant growth chamber in continuous light (120 µmol s^{-1} m^{-2}). For the RH
phenotype characterization, seeds were surface sterilized and stratified in darkness for
3 days at 4°C. Then grown on ½ strength MS agar plates, in a plant growth chamber at
22 °C in continuous light (120 µmol s^{-1} m^{-2}) for 7-14 days depending on the experiment
at 22°C as a pretreatment and then at 10°C (moderate-low temperature treatment) for
12, 24, 36, 48, 60 and 72 hours. Measurements were made between 7-17 days. For
quantitative analysis of RH phenotype, 10 fully elongated RHs were measured (using
the ImageJ software) from each root (n=20) grown by three biological replicates on
vertical agar plates. After treatment only new RH was measured. Images were captured
with a Leica EZ4 HD Stereo microscope (Leica, Germany) equipped with the LAZ ez
software.Results were expressed as the mean ± SD using the GraphPad Prism 8.0.1 (USA)
statistical analysis software.

RH phenotype in monocots. Plant species and varieties used include *Brachypodium*
distachyon (Bd21), rice (Kitaake), and wheat (Chinese Spring). All seeds were imbibed in
water for 2 hours and then surface sterilized with 0.6% hypochlorite and 1% Triton X-
100 with gentle shaking for 8 minutes. Seeds were then rinsed five times with sterile
water and then placed uniformly across ½ Murashige and Skoog (MS; PhytoTechnology
laboratories, USA), pH 5.7, 3% Gelzan (PhytoTechnology laboratories, USA) plates and
stored at 4°C for 3 d in the dark. For Arabidopsis, seeds rinsed with 70% ethanol and
then surface sterilized with 0.6% hypochlorite and 0.05% Triton X-100 with gentle
shaking for 5 minutes. Seeds were then rinsed five times with sterile water and placed
uniformly across ½ Murashige and Skoog (MS; PhytoTechnology laboratories, USA), 1%
sucrose, pH 5.7, 3% Gelzan (PhytoTechnology laboratories, USA) plates and stored at
4°C for 3 days in the dark. Plants were grown in a Conviron Adaptis growth chamber
model CMP6010 with a 16 h : 8 h, day : night cycle at a light intensity of 100 µmol m^{-2}s^{-1}

at 22°C. 3 fully elongated root hairs from the maturation zone were measured per root after 8 d at 22°C and after 5 d at 22°C + 3d at 10°C. Images were captured using a M205 FA Leica Stereo microscope. The root length data is represented as a box blot with the solid color dot representing the mean. The root hair density data are the mean±SE (n=10–15 roots). A t-test was used to determine significant differences between the means.

RNA-seq analysis. For the RNA-seq analysis, seedlings were grown on ½ strength MS agar plates, in a plant growth chamber at 22 °C in continuous light (120 µmol s^{-1} m^{-2}) for 14 days at 22°C as a pre-treatment and then at 10°C (moderate-low temperature treatment) for 2 and 6 hours. We analyzed a dataset with 12 factor groups (tree time points: B, 2h, 6h) and four genotypes (Col-0, *rsl2*, *rsl4* and *rsl2 rsl4*) each with three biological replicates giving 36 samples in total. Total RNA was extracted from 20–30 mg of frozen root tissue. Frozen root samples were ground in liquid nitrogen and total RNAs were extracted using E.Z.N.A Total RNA Kit I (Omega Bio-tek, Georgia, USA). RNA quantity and purity were evaluated with a Qubit®2.0 fluorometer (Invitrogen™, Carlsbad, CA, USA) using a Qubit™ RNA BR assay kit. RNA integrity and concentration were assessed by capillary electrophoresis using an automated CE Fragment Analyzer™ system (Agilent Technologies, Santa Clara, CA, USA) with the RNA kit DNF-471-0500 (15nt). Total RNA-seq libraries were prepared according to the TruSeq Stranded Total RNA Kit (Illumina, San Diego, CA, USA) following the manufacturer's instructions. Finally, the constructed libraries were sequenced using Macrogen sequencing services (Seoul, Korea) in paired end mode on a HiSeq4000 sequencer. For total RNA differential expression analysis, a quality check was performed with FASTQC software (Andrews, 2010). Then, the adapter sequences were removed, reads with a quality score less than 30 and length less than 60 nucleotides were eliminated using Flexbar (Dodt *et al.,* 2012). Resulting filtered reads were aligned against *Arabidopsis thaliana* Araport 11 genome with the STAR aligner software. A total of 36 RNA libraries were sequenced, obtaining an average of 84,741,219 reads for each one, with a minimum and maximum value of 72,658,864 and 102,182,242 reads, respectively. After filtering them by quality and removing adapters, an average of 98.6% of the reads remained and after aligning them against the *Arabidopsis thaliana* reference genome, between 88.1% (69,820,848) and 96.5% (93,740,042) of total reads were correctly aligned. For each library, the feature Counts software from the Rsubread package (Liao *et al.,* 2019) was applied to assign expression values to each uniquely aligned fragment. Differential gene expression analysis was performed using the Bioconductor R edgeR package (Robinson *et al.,* 2010). Differentially expressed genes (DEGs) were selected with an FDR < 0.05 and a Log2 FC > |0.5|. To search for genetic functions and pathways overrepresented in the DEG lists, genetic enrichment analysis was performed using the Genetic Ontology (GO) database with the R package ClusterProfiler v4.0.5 (Yu *et al.,* 2012), using the compareCluster function. The parameters used for this analysis were: lists of differentially expressed

genes for each comparison in ENTREZID, enrichGO sub-function, the universe from the total of differentially expressed genes that present annotation as genetic background, Benjamini-Hochberg statistical test and a filter of FDR less than 0.05. Subsequently, the semantics filter of GO terms was performed using the simplify function of the same package using a p-value and q-value cutoff less than 0.05.

RT-qPCR analyses. Total root RNA was extracted from these samples using the E.Z.N.A. Total RNA Kit (Omega Bio-tek, Georgia, USA) according to the product's protocol. One microgram of total RNA was reverse transcribed using an oligo(dT)$_{20}$ primer (Macrogen, South Korea) and M-MLV Reverse Transcriptase™ (Invitrogen, USA) according to the manufacturer's instructions. cDNA was then diluted 3-fold before qPCR analysis. qPCR was performed on an AriaMx™ Real-time PCR System (Agilent, USA) using 10 uL of Brilliant III Ultra-Fast SYBR Green Master-Mix (Stratagene, USA), 1 μL of cDNA, and 0.25 μM of each primer to a total volume of 20 μL per reaction. The *TAFII15* (AT4g31720) gene was used as reference for normalization of gene expression levels. Statistical analysis was performed using either parametric or non-parametric T-tests, after determining the normality of the populations via the Shapiro-Wilk test. Results were plotted showing mean and standard deviation. Primer used is detailed in **Table S2**.

Network construction. Direct and indirect targets of RSL4 were identified using the RSL4 cis-motif reported in Hwang *et al.,* 2017 and the ConnecTF database. ConnecTF was used to connect RSL4 to its indirect targets via TF2s that are themselves regulated by RSL4 according to our RNA-seq analysis. We queried the ConnecTF database for all the DAP-seq *in vitro* binding datasets. We restricted the results returned using Target Genes filter with the list of differentially expressed genes according to our ANOVA results for *rsl4* mutants (GxT genes). For this query, the TF2s were restricted using the Filter TFs option with the list of TFs regulated by RSL4. Finally, the resulting network was exported in the Network tab, combined with the direct TF-target information based on RSL4 cis-motif, uploaded, and visualized using Cytoscape.

TF-target list enrichment. Using the Target Enrichment tab in ConnecTF, all the TF2s that are enriched for RSL4 indirect targets were analyzed and visualized. The significance of the overlap between TF–targets captured by DAP-seq in the list of RSL4 regulated targets was calculated. The P-values are calculated using Fisher's exact test adjusted with the Bonferroni correction. The background set of genes used for the calculation, which is by default all protein-coding genes for the Arabidopsis instances of ConnecTF, were manually set by using the Background genes option in the query page.

Chromatin immunoprecipitation (ChIP) PCR assay. Chromatin immunoprecipitation (ChIP) assays were performed on RSL4:RSL4-GFP plants (Moison *et al.* 2021) mainly as described in (Ariel *et al.,* 2020). Plants were grown for 10 d in plates containing MS 0.5×

medium (pH 5.7; 0.8% agar) placed vertically in a culture chamber at 22 °C and continuous light (120 μmol s^{-1} m^{-2}). After 10 d, the plates were placed horizontally and treated with β-estradiol 10 μM solution for 3 h. The expression of RSL4 was checked by qPCR. Chromatin was crosslinked with formaldehyde 1% (v/v) for 10 min at room temperature. Crosslinking was stopped by adding glycine (125 mM final concentration) and incubating for 10 min at room temperature. Crosslinked chromatin was extracted by cell resuspension, centrifugation, cell membrane lysis, and sucrose gradient as previously described (Ariel *et al.,* 2020). Nuclei were resuspended in Nuclei Lysis Buffer and chromatin was sonicated using a water bath Bioruptor Pico (Diagenode; 30 s on/30 s off pulses, at high intensity for 10 cycles). Chromatin samples were incubated for 12 h at 4 °C with Protein G Dynabeads (Invitrogen) precoated with the antibodies anti-FLAG (Abcam ab18230) or anti-IgG (Abcam ab6702) as a negative control. Immunoprecipitated DNA was recovered using Phenol:Chloroform: Iso-amilic Acid (25:24:1; Sigma) and analyzed by qPCR using the primers listed in **Table S2.** One upstream region of the EXP7 gene was used as a positive control (Hwang *et al.,* 2017). Untreated sonicated chromatin was processed in parallel and considered the input sample. The GraphPad Prism 6 software was used to analyze the data and produce the graphs.

GTL1 ChIP-seq analysis. Illumina sequencing reads derived from ChIP and Input DNA of GTL1 were mapped to the TAIR10 reference genome employing Bowtie2 for alignment. Subsequently, the elimination of duplicate reads was performed. For the identification of GTL1 ChIP peaks in relation to its corresponding Input DNA control, the MACS2 software was utilized with a significance threshold set at a q-value of 0.05. Post peak-calling, gene annotation of the identified peaks was conducted utilizing BEDtools, which incorporated a 2 kb region upstream from the transcription start site (TSS) for consideration. The preparation of the sequencing data for visualization involved filtering, ordering, and scaling of bam files, followed by conversion to bigwig format. This conversion leveraged the "bamCoverage" function within deepTools 2.0, configured with a bin size of 10 bp, RPKM normalization, and a smooth length of 60. Visual representations of the genomic data displayed in Figure 4B were generated using the Integrative Genomics Viewer (IGV)(reference: 10.1093/bib/bbs017).

Confocal Microscopy. Confocal laser scanning microscopy for the lines RSL4:RSL4-GFP, RSL2:RSL2-GFP, RHD6:RHD6-GFP, GTL1:GTL1-GFP and DF1:DF1-GFP was performed using Zeiss LSM5 Pascal (Zeiss, Germany) (Excitation: 488 nm argon laser; Emission: 490–525 nm, Zeiss Plain Apochromat 10X/0.30 or 40X/1.2 WI objectives according to experiment purpose). Z stacks were done with an optical slice of 1 μm, and fluorescence intensity was measured at the RH tip. The scanning was performed using confocal laser scanning microscopy Zeiss LSM 710 (Carl Zeiss, Germany). For image acquisition, 20x/1.0 NA Plan-Apochromat objective for root tips were used. The GFP signal was

excited with a 488 nm argon laser at 4% laser power intensity and emission band of 493-549 nm. Propidium Iodide signal was excited with a 488 nm argon laser at 4% laser power intensity and emission band 519-583 nm. GFP signal at the RH tip were quantified using the ImageJ software. Fluorescence AU was expressed as the mean ± SD using the GraphPad Prism 8.0.1 (USA) statistical analysis software. Results are representative of two independent experiments, each involving 10–15 roots.

Acknowledgements

We would like to thank xxxx for the seed lines. We thank NASC (Ohio State University) for providing T-DNA lines seed lines, Trevor Nolan (Phillip Benfey group) for the quadruple HB mutant, J.M.E., F.A. and R.L.C. are investigators of the National Research Council (CONICET) from Argentina. M.I. is supported by ANID FONDECYT POSTDOCTORADO [grant 3220138]. This work was supported by grants from ANPCyT (PICT2019-0015 and PICT2021-0514), by ANID – Programa Iniciativa Científica Milenio ICN17_022, NCN2021_010 and Fondo Nacional de Desarrollo Científico y Tecnológico [1200010] to J.M.E. J.M.A lab is supported by ANID FONDECYT 1210389, Programa Iniciativa Científica Milenio ICN17_022, and NSF Plant Genome Grant NSF-PGRP: IOS-1840761.

Author Contribution

T.U.L. and V.B.G performed most of the experiments, analyzed the data and helped in the writing process of the manuscript. M.A.I. and H.S-G. performed some experiments and helped in the writing process. G.N-L. and C.M. performed the RNA-seq analysis, T.M. and J.M.A. performed the gene network analysis and helped on the writing process. L.F. and F.A. carried out the ChIP experiment of RSL4 and helped on the writing process. R.M. and K.S. carried out the root hair characterization in monocots. A.A.M. helped in the writing process. M.S. and K.S. provided seeds and helped in the writing process. R.L.C. provide AtHBs lines and help in the writing process. J.M.E. designed research, analyzed the data, supervised the project, and wrote the paper. All authors commented on the results and the manuscript. This manuscript has not been published and is not under consideration for publication elsewhere. All the authors have read the manuscript and have approved this submission.

Competing financial interest

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to J.M.E. (Email: jose.estevez@unab.cl). HD-Zip mutants and overexpressor lines must be requested to R.L.C.

Figure 1. Low temperature effect on RH growth in Arabidopsis and monocots. (**A**) Low-temperature RH growth up to 3 days of treatment. (**B**) Temperature fluctuations impact on RH growth and growth maxima was detected at 8-10°C. (**C**) A nutrient-free medium can suppress the low-temperature growth impact on RHs. The previously recorded RH traits at low temperature were verified with somewhat larger magnitude using this D-root method. (A-C) Each point is the mean of the length of the 5 longest RHs identified in the maturation zone of a single root. Data are the mean ± SD (N= 30 roots), two-way ANOVA followed by a Tukey–Kramer test; (****) $p < 0.001$, NS= non-significant. Results

are representative of three independent experiments. Asterisks indicate significant
differences between Col-0 and the corresponding treatment at the same temperature
or between the same treatment at different temperatures. Representative images of
each genotype are shown on the right. Scale bars= 500 µm. (**D**) Low-temperature
triggers an enhancement on RH growth in the monocots *Brachypodium distachyon*
(Bd21), rice (Kitaake), and wheat (Chinese Spring). Representative pictures are shown in
the bottom. Scale bars= 500 µm. Data are the mean ± SD (N= 30 roots), two-way
ANOVA followed by a Tukey–Kramer test; (**) $p < 0.01$, (***) $p < 0.001$, NS= non-
significant.

614
615
Figure 2. Identification of the RH specific RSL4-mediated gene regulatory network
acting at low temperature on an early temporal window. (A) Dual bioinformatic-
experimental strategies using RNA-seq plus DAP-seq prediction followed to identify the

RH specific RSL4-transcriptional network at early times at low temperature. (**B**)
Differential expression analysis by a heat map of the transcriptome of Wt Col-0 and *rsl4*
genotypes in response to low temperature treatments (2h and 6h). A two-way analysis
of variance (ANOVA) analysis was carried out, considering temperature (T) and each
genotype (G) as factors and controlling type I error using the false discovery rate. A total
of 2,365 DEG showed an altered response to low temperature in the *rsl4* mutant
(significant TxG factor) were grouped into 3 clusters. (**C**) Best 5-ranked list of TFs based
on the overall overlap of 2,365 genes controlled by RSL4 with DAP-seq common targets
(on the left) and RSL4-specifically expressed in RHs (on the right) using the ConnecTF
database. (**D**) Gene regulatory RSL4-network acting on RH growth at low temperature
after 6h of exposure. Genes are drawn as triangles (TF, transcription factors) and
squares (target genes). Lines indicate predicted transcriptional activation or predicted
transcriptional repression. Yellow lines indicate RSL4, red line AtHB13 and green lines
MYB-like putative TFs binding site by DAP-seq in the upstream region of the
corresponding genes.

Figure 3. Major nodes of the RSL4-transcriptional primary and secondary network
impact on RH phenotypes linked to gene expression changes at low temperature. (A)

Heatmap of the RH phenotype at 22°C and 10°C of the different mutants and overexpression lines of the main network nodes RSL4 (and RSL2), AtHB13, GTL1/DF1 and MYB-like as well as the downstream selected components AtHB16/AtHB23 and ERD7/ERD10. Blue color scale represents the length of RH in μm (0-1500). Pair comparisons are made between Col-0 and each line (see **Table S1** and **Figures S5-S12**). Data are the mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; p-values are shown. (**B**) Representative images of RH phenotype of Col-0, GTL1, MYB-like (At2G01060) and AtHB13 as well as RSL2-RSL4 quantified in (A). Scale bars = 1000 μm. (**C**) Heatmap of the gene expression analysis of the network main nodes TFs RSL4, AtHB13 and GTL1/DF1 and as well as the downstream selected components ERD7/ERD10 in each mutant/overexpression line previously isolated and characterized (**Figures S12-S16**). Data are the mean of three biological replicates, two-way ANOVA followed by a Tukey–Kramer test; p-values are shown. (**D**) Main transcriptional nodes of the RSL4 network with the detailed effect of each component at transcriptional and RH phenotypic levels. Arrows indicate positive gene expression regulation and blunt arrows indicate transcriptional repression. Green curved line means positive RH growth regulation and red curved line means negative RH growth regulation.

Figure 4. Regulation of RSL4 on the main transcriptional nodes and selected downstream target genes and low temperature effect on main TFs reporters expression.

(**A**) ChIP-qPCR analysis of RSL4 binding to primary and secondary RSL4 nodes gene targets promoter regions at low temperature. Schemes of the loci showing the location of the fragments analyzed by ChIP-qPCR are shown in the upper part (A = Primer A; B = Primer B; F = Primer forward; R = Primer reverse). Primers were designed analyzing ATAC-seq experiments in regions where the chromatin is accessible (Maher *et al.,* 2018). *EXP7* oligos were used as positive controls (xxxx). The data are mean ± SD of 2 technical replicates. The experiment was performed twice with similar results. The enrichment was measured relative to the negative control ACTIN. (**B**) ChIP-seq analysis of GTL1 binding on HB16, ERD7/ERD10 as secondary RSL4 nodes gene targets promoter regions only at room temperature. Schemes of the loci. (**C-D**). Effect of the low temperature on the expression of GTL1 and RSL4 in the elongation and differentiation root zones. (**C**) Confocal images of GTL1:GTL1-GFP in *gtl-1* translational reporter line of at 22°C and after transfer from ambient to low temperature (22°C→10°C). Quantitative evaluation of the GFP fluorescence intensity across the root elongation zones at 22°C and after transfer from ambient to low temperature. Fluorescence intensity is expressed in arbitrary units (A.U.), N=6-8 roots. Results are representative of three independent experiments. Scale bars=50 µm. (**D**) Confocal images of RSL4:RSL4-GFP translational reporter line of at 22°C and after transfer from ambient to low temperature (22°C→10°C). Quantitative evaluation of the GFP fluorescence intensity across the root elongation zones at 22°C and after transfer from ambient to low temperature. Fluorescence intensity is expressed in arbitrary units (A.U.), N=6-8 roots. Results are representative of three independent experiments. Scale bars=50 µm. (**D**) Confocal images of RSL4:RSL4-GFP translational reporter line of at 22°C and after transfer from ambient to low temperature (22°C→10°C). Quantitative evaluation of the GFP fluorescence intensity across the root elongation zones at 22°C and after transfer from ambient to low temperature. Fluorescence intensity is expressed in arbitrary units (A.U.), N=8 roots. Results are representative of three independent experiments. Scale bars=50 µm. (**E**) Expression in roots of GUS reporter lines for AtHB13, AtHB16 and AtHB23 at 22°C and after transfer from ambient to low temperature (22°C→10°C). Images are representative of three independent experiments. Scale bars=200 µm.

Figure 5. Proposed model of RSL4-regulation of RH growth at low temperature at early times (2-6 hs). Main transcriptional nodes of the RSL4-regulated network with the detailed effect of each component at transcriptional and RH phenotypic levels. Red circles indicate that repress RH growth at low temperature and Green circles are enhancers of RH growth. Direct regulation at 10°C/22°C (D10/D22) was assessed by ChIP results. Arrows indicate positive gene expression regulation and blunt arrows indicate transcriptional repression. (?) Indicates that needs to be experimentally validated. RHS= Root Hair Specific genes.

Figure S1. Influence of room temperature pretreatment time length on low

temperature root hair growth in Arabidopsis thaliana. (**A**) quantification of RH length

at different treatment times at 22°C prior to cold exposure (7 and 14 days) and different

cold exposure times (0, 12, 24, 36, 48, 60 and 72 hours) in both pretreatment conditions.

(**B**) visualization by stereo microscopy of the phenotypic data represented. Data are the

mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) $p < 0.05$,

(**) $p < 0.01$, (***) $p < 0.001$, NS=non-significant. Results are representative of three

independent experiments. Asterisks indicate significant differences between Col-0 and

the corresponding genotype at the same temperature. Representative images of each

genotype are shown below. Scale bars= 500 µm.

708
709
Figure S2. **DEG analysis between Col-0 and *rsl2, rsl4* and *rsl2rsl4* mutants at low**
temperature at 2h and 6h. (A) Root samples of *Arabidopsis thaliana* cv. Col-0 were

compared with the mutant samples *rsl2*, *rsl4* and the double mutant *rsl2 rsl4* in a basal
state at 22ºC (B) and two cold treatments for 2 and 6 hours at 0ºC (2h and 6h). 6,756
differentially expressed genes (DEG) in at least one condition were represented in a
blue-red scale heatmap. Each column represents the expression of each replicate.
Expression data was scaled considering the mean centered divided to standard
deviation. (**B**) Clustering analysis of differential expressed genes was made using the
Ward method. Continuous line represents each cluster tendency. (**C**) Gene ontology
term enrichment analysis of genes belonging to each one of the identified 1-5 clusters.
GO in cluster 5 showed several RH development processes. The blue-red scale color
represents adjusted p-value and the point size represents gene ratio.
722

724
725

Figure S3. Differential expression analysis between Col-0 and rsl2 *rsl4* mutant samples
in low temperature treatment. (**A**) Volcano plot for differentially expressed genes
comparing Col-0 and *rsl2* single mutant. Candidate genes were filtered with a *p-value* <
0.01. Down (light blue points) and up (dark blue points) regulated *rsl2* genes were
selected with an FC < -0.5 and FC > 0.5, respectively. (**B**) Volcano plot for differentially
expressed genes comparing Col-0 and *rsl4* single mutant. Candidate genes were filtered
with a *p-value* < 0.01. Down (green points) and up (dark green points) regulated *rsl4*
genes were selected with an FC < -0.5 and FC > 0.5, respectively. (**C**) Volcano plot for
differentially expressed genes comparing Col-0 and *rsl2rsl4* double mutant. Candidate
genes were filtered with a *p-value* < 0.01. Down (red points) and up (dark red points)
regulated *rsl2rsl4* genes were selected with an FC < -0.5 and FC > 0.5, respectively. (**D**)
Venn diagram comparing the down (left side) and up (right side) regulated transcripts
between *rsl2, rsl4* and *rsl2rsl4* candidate genes.

739
740
741

Figure S4. **Expression of *RHD6*, *RSL4* and *RSL2* under low temperature.** *RHD6, RSL2* and *RLS4* transcript levels were measured by RT-qPCR in roots of Wt Col-0 roots in parallel after 10 days at 22°C, followed by 6h, 24h and 36h at 10°C. Statistical differences between populations were assessed by one-way ANOVAs followed by Tukey post-tests. Significant differences were represented by letters. The average Cts for the technical duplicates of each transcript were normalized using the constant housekeeping transcript of *TAFII15*, which was evaluated parallelly in the same fashion. (a) p <0.05, (ab) p <0.005, (b) p <0.001.

750
751

Figure S5. (**A**) AtHB13 gene structure and the insertion of the T-DNA mutants. Arrow
indicates the position of the primers. (**B**) Scatter-plot of RH length of Col-0, *athb13-2*
mutant and overexpressor 35S:AtHB13 grown at 22°C or at 10°C. Each point is the mean
of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD
(N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; ns=non-significant.
Results are representative of three independent experiments. Asterisks indicate
significant differences between Col-0 and the corresponding genotype at the same
temperature. (**C**) Representative images of each genotype are shown below. Scale bars=
500 μm. (**D**) AtHB13 expression levels in Col-0, *athb13-2* and 35S:HB13 lines.

Figure S6. (**A**) MYB-like gene structure and the insertion of the T-DNA mutant. (**B**) MYB like expression levels in Col-0, *myb-like-1* line. (**C**) Scatter-plot of RH length of Col-0, mutants grown at 22°C or at 10°C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; ns=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature. (**D**) Representative images of each genotype. Scale bars= 500 μm.

Figure S7. (**A**) GTL1 gene structure and the insertion of the T-DNA mutant *gtl1-1*. (**B**) GTL1 expression levels in Col-0, *gtl1-1* and the overexpressor EXP7:GTL1-GFP lines. (**C**) Scatter-plot of RH length of Col-0 and *gtl1-1* mutant, overexpressos EXP7:GTL1-GFP grown at 22°C or at 10°C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Data are the mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; (***) $p < 0.001$, NS=non-significant. Results are representative of three independent experiments. Asterisks indicate significant differences between Col-0 and the corresponding genotype at the same temperature. (**D**) Representative images of each genotype are shown below. Scale bars= 500 µm.

782
783
Figure S8. (**A**) RSL4 expression levels in Col-0, *rsl2/rsl4* and the overexpressor
EXP7:RSL4-GFP lines (**B**) Scatter-plot of RH length of Col-0, *rsl2-1* and *rsl4-1* mutants,
overexpressor EXP7:RSL4-GFP, *rhd6* grown at 22°C or at 10°C. Each point is the mean of
the length of the 10 longest RHs identified in a single root. Data are the mean ± SD
(N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; (*) p <0.05, (***) p
<0.001, NS=non-significant. Results are representative of three independent
experiments. Asterisks indicate significant differences between Col-0 and the
corresponding genotype at the same temperature. (**C**) Representative images of each
genotype are shown below. Scale bars= 500 μm.

Figure S9. (**A**) AtHB16 gene structure and the insertion of the T-DNA mutants. (**B**)

Scatter-plot of RH length of Col-0, *athb16-1* and *athb16-2* mutants and 35S:AtHB16

lines (2,14,30) grown at 22°C or at 10°C. Each point is the mean of the length of the 10

longest RHs identified in a single root. Data are the mean ± SD (N=20 roots), two-way

ANOVA followed by a Tukey–Kramer test; (***) p <0.001, NS=non-significant. Results

are representative of three independent experiments. Asterisks indicate significant

differences between Col-0 and the corresponding genotype at the same temperature.

(**C**) Representative images of each genotype are shown below. Scale bars= 500 µm. (**D**)

AtHB13 expression levels in Col-0, *athb16-1* and *athb16-2* mutants and 35S:AtHB16

overexpression lines (2,14,30).

804
Figure S10. (**A**) AtHB23 gene structure and target of the amiRNA *athb23* silenced lines.
(**B**) Scatter-plot of RH length of Col-0, amiRNA *athb23* (IB, 9C, 20C) lines and 35S:AtHB23
lines (5H, 6A,7F) grown at 22°C or at 10°C. Each point is the mean of the length of the
10 longest RHs identified in a single root. Data are the mean ± SD (N=20 roots), two-way
ANOVA followed by a Tukey–Kramer test; (**) p <0.01, (***) p <0.001, NS=non-
significant. Results are representative of three independent experiments. Asterisks
indicate significant differences between Col-0 and the corresponding genotype at the
same temperature. (**C**) Representative images of each genotype are shown below. Scale
bars= 500 μm. (**D**) AtHB13 expression levels in Col-0, amiRNA *athb23-1* and amiRNA
athb23 (IB, 9C, 20C) lines and 35S:AtHB23 line.

815
Figure S11. (**A**) Scatter-plot of RH length of Col-0, and quadruple *athb3 athb13 athb20*
athb23 grown at 22°C or at 10°C. Each point is the mean of the length of the 10 longest
RHs identified in a single root. Data are the mean ± SD (N=20 roots), two-way ANOVA
followed by a Tukey–Kramer test; (*) $p < 0.05$. Results are representative of three
independent experiments. Asterisks indicate significant differences between Col-0 and
the corresponding genotype at the same temperature. (**B**) Representative images of
each genotype are shown below. Scale bars= 500 µm.

Figure S12. (**A**) ERD7 gene structure and the insertion of the T-DNA mutants. (**B**)
Scatter-plot of RH length of Col-0, *erd7-1* mutant grown at 22°C or at 10°C. Each point is
the mean of the length of the 10 longest RHs identified in a single root. Data are the
mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test; ns=non-
significant. Results are representative of three independent experiments. Asterisks
indicate significant differences between Col-0 and the corresponding genotype at the
same temperature. (**C**) Representative images of each genotype are shown below. Scale
bars= 500 μm.

832
Figure S13. (**A**) ERD10 gene structure and the insertion of the T-DNA mutants. (**B**)
Scatter-plot of RH length of Col-0 and *erd10-1* mutant grown at 22°C or at 10°C. Each
point is the mean of the length of the 10 longest RHs identified in a single root. Data are
the mean ± SD (N=20 roots), two-way ANOVA followed by a Tukey–Kramer test;
ms=non-significant. Results are representative of three independent experiments.
Asterisks indicate significant differences between Col-0 and the corresponding
genotype at the same temperature. (**C**) Representative images of each genotype are
shown below. Scale bars= 500 μm.

841
842 **Figure S14**. Expression levels of the RH main gene regulatory network nodes RSL4,
843 AtHB13 and GTL1, and MYB-like (AT2G01060) in Wt and in the mutants *rsl2/rsl4*,
844 *athb13-2*, *gtl1-1* and *myb-like*. The mean and standard deviation is plotted, followed by
845 parametric and non-parametric T-tests on populations paired treatment-wise.

846
Figure S15. Expression levels of the RSL4-downstream genes ERD7 and ERD10 in Wt and
in the mutants *rsl2/rsl4*, *athb13-2*, *gtl1-1* and *myb-like 1*. The mean and standard
deviation is plotted, followed by parametric and non-parametric T-tests on populations
paired treatment-wise.

Figure S16. Expression levels of the RH main gene regulatory network nodes RSL4,
AtHB13 and GTL1, and MYB-like (AT2G01060) in Wt and in the overexpression lines
EXP7:RSL4-GFP, EXP7:GTL1-GFP and 35S:AtHB13. The mean and standard deviation is
plotted, followed by parametric and non-parametric T-tests on populations paired
treatment-wise.

857
Figure S17. Expression levels of the RSL4-downstream genes ERD7 and ERD10 in Wt and
in the overexpression lines EXP7:RSL4-GFP, EXP7:GTL1-GFP and 35S:AtHB13. The mean
and standard deviation is plotted, followed by parametric and non-parametric T-tests
on populations paired treatment-wise.

Figure S18. **Effect of the low temperature on the expression of DF1 and RHD6 in the**
elongation and differentiation root zones. (**A**) Confocal images of DF1:DF1-GFP in *df1-1*
translational reporter line of at 22°C and after transfer from ambient to low
temperature (22°C→10°C). Quantitative evaluation of the GFP fluorescence intensity
across the root elongation zones at 22°C and after transfer from ambient to low
temperature. Fluorescence intensity is expressed in arbitrary units (A.U.), N=6-8 roots.
Results are representative of three independent experiments. Scale bars=50 μm. (**B**)
Confocal images of RHD6:RHD6-GFP translational reporter line of at 22°C and after
transfer from ambient to low temperature (22°C→10°C). Quantitative evaluation of the
GFP fluorescence intensity across the root elongation zones at 22°C and after transfer
from ambient to low temperature. Fluorescence intensity is expressed in arbitrary units
(A.U.), N=8 roots. Results are representative of three independent experiments. Scale
bars=50 μm.

Table S1. Mutants and transgenic lines used and generated in this study.

			Mutants and transgenic lines		
Locus	**Gene name**	**Gene expression effect**	**Line name**	**ABRC name/stock number**	**Reference**
AT1G27740	*RSL4*	mutant	*rsl2 rsl4*		Yi et al., 2010
		O.E.	EXP7:RSL4		Yi et al., 2010
AT1G69780	*AtHB13*	mutant	*athb13-1*	SAIL_893_G05	Ribone et al., 2015
		mutant	*athb13-2*	GABI_859H06	Ribone et al., 2015
		O.E.	35S:AtHB13		Cabello et al., 2012
		reporter GUS	HB13:GUS		Ribone et al., 2015
AT2G01060	*MYB-like*	mutant	*myb-like 1*	SALKseq_8995	this study
AT1G33240	*GTL1*	complemented	GTL1:GTL1-GFP/*gtl1-1*		Shibata et al., 2018
		O.E.	EXP7:GTL1-GFP/Col-0		Shibata et al., 2018
		mutant	*gtl1-1*	WiscDsLox413-416C9	Shibata et al., 2018
		mutant	*gtl1-1 df1-1*		Shibata et al., 2018
AT1G76880	*DF1*	complemented	DF1:DF1-GFP/*df1-1*		Shibata et al., 2018
		O.E.	EXP7:DF1-GFP/Col-0		Shibata et al., 2018
		mutant	*df1-1*	SALK_106258	Shibata et al., 2018
AT4G40060	*AtHB16*	mutant	*athb16-1*		This study
		mutant	*athb16-2*		This study
		O.E.	35S:AtHB16-2		This study
		O.E.	35S:AtHB16-14		This study
		O.E.	35S:AtHB16 30		This study
		reporter GUS	HB16:GUS		This study
AT5G39760	*AtHB23*	O.E.	35S:AtHB23-7F		Perotti et al. 2021, 2022
		O.E.	35S:AtHB23-5H		Perotti et al. 2021, 2022
		O.E.	35S:AtHB23-6A		Perotti et al. 2021, 2022
		silenced	amiR-AtHB23-1B		Perotti et al. 2021, 2022
		silenced	amiR-AtHB23-20C		Perotti et al. 2021, 2022
		silenced	amiR-AtHB23-9C		Perotti et al. 2021, 2022
		reporter GUS	HB23:GUS		Perotti et al. 2021
AT5G15150 AT1G69780	*AtHB3* *AtHB13*	mutant mutant	*Athb3 athb13 athb20*		Nolan et al. 2023

AT3G01220 AT5G39760	*AtHB20* *AtHB23*	mutant mutant	*athb23*		
AT2G17840	*ERD7*	mutant	*erd7-1*	WISCDSLOX452E10	this work
AT1G20450	*ERD10*	mutant	*erd10-1*	SALK_087789C	this work

880 **Table S2**. Oligonucleotide list for RT-qPCR analysis. Primer sequences along with their
881 respective Tms are shown next to their corresponding targeted genes.
882

RT-qPCR primer list		
Gene	**Primer probe**	**Sequence (5' - 3')**
AT2G01060	Forward	TCGATTGGTGAGAGGTGCAG
	Reverse	AAAGGCGAGGACATGGTCTG
ERD7	Forward	AAGGGCTCGATGCAGCAGGATA
	Reverse	GGTTTTTGCAAGCGTCGAAGGC
ERD10	Forward	ACCGATTGCTGACATCCCGGAG
	Reverse	TCTCCAGTGGTCTTGGCGTGAT
GTL1	Forward	ATGGAATTGTTTGAAGGTTTGG
	Reverse	GACATGACCTCGTGTTCTCG
AtHB13	Forward	TCTCAACTGCGCCGCCATCAAA
	Reverse	GCGACGGTGGGAAGAAGTGTCG
RHD6	Forward	TGATTTGGTGACAATGCTTGA
	Reverse	GGAGAGAATGGCATCAATGG
RSL2	Forward	CCCCAATGGAACAAAGGTC
	Reverse	TCTCGGTGAGCTGAGACCAA
RSL4	Forward	GTGCCAAACGGGACAAAAGT
	Reverse	TTGTGATGGAACCCCATGTC
TAFII15	Forward	GAATCACGGCCAACAATC
	Reverse	ACTCTTAGCCAAGTAGTGCTCC

Genotyping primer list		
Gene	**Primer probe**	**Sequence (5' - 3')**
myb-1 (SALKseq_8995.3)	Left primer	CCATGAGAGGGAAAGCCTATC
	Right primer	AGGACATGGTCTGACTGATGG
myb-2 (SALKseq_034222)	Left primer	AGGAGCAATTTGAACTCCCTC
	Right primer	TTGGAAACCTGGATTGTTGAC
erd7-1 (WISCDSLOX452E10)	Left primer	TTTAATGCAATTTCCGAGACG
	Right primer	CAATCTCTTTCCTTTCCCCAG
erd10-1 (SALK_087789C)	Left primer	CTCACCGTCTTCACCTTCTTC
	Right primer	TGCATTCTTCTCCAATTCTCC

REFERENCES

M.K. Aalto, E. Helenius, T. Kariola, V. Pennanen, P. Heino, H. Horak, I. Puzorjova,H.
Kollist, E.T. Palva, ERD15-An attenuator of plant ABA responses and stomatal aperture,
Plant Sci. 182 19–28. https://doi.org/10.1016/j.plantsci.2011.08.009 (2012).

Alvarez, J. M., Riveras, E., Vidal, E. A., Gras, D. E., Contreras-López, O., Tamayo, K. P.,
Aceituno, F., Gómez, I., Ruffel, S., Lejay, L., Jordana, X., Gutiérrez, R. A. Systems
approach identifies TGA1 and TGA4 transcription factors as important regulatory
components of the nitrate response of Arabidopsis thaliana roots. *Plant J*. 80(1), 1-13
https://doi.org/10.1111/tpj.12618 (2014).

Arce, A.L., Raineri, J., Capella, M., Cabello, J.V. , Chan, R.L. Uncharacterized conserved
motifs outside the HD-Zip domain in HD-Zip subfamily I transcription factors; a potential
source of functional diversity. *BMC Plant Biol.* 11(42). http://doi.org/10.1186/1471-
2229-11-42 (2011).

Ariel, F., Lucero, L., Christ, A., Mammarella, M. F., Jegu, T., Veluchamy, A., Mariappan, K.,
Latrasse, D., Blein, T., Liu, C., Benhamed, M., Crespi, M. R-Loop Mediated trans Action of
the   APOLO   Long   Noncoding   RNA.   *Mol.   Cell*   77(5),   1055-1065.
https://doi.org/10.1016/j.molcel.2019.12.015 (2020).

Barajas-Lopez JD, Tiwari A, Zarza X, Shaw MW, Pascual JS, Punkkinen M, Bakowska JC,
Munnik T, Fujii H. EARLY RESPONSE TO DEHYDRATION 7 Remodels Cell Membrane Lipid
Composition during Cold Stress in Arabidopsis. Plant Cell Physiol. 2021 Mar 25;62(1):80-
91. doi: 10.1093/pcp/pcaa139.

Bhosale, R., Giri, J., Pandey, B.K. *et al*. A mechanistic framework for auxin dependent
Arabidopsis root hair elongation to low external phosphate. *Nat. Commun.* 9, 1409.
https://doi.org/10.1038/s41467-018-03851-3 (2018).

Brady, S., Orlando, D., Lee, J. Y., Wang, J., Koch, J., Dinneny, J., Mace, D., Ohler, U.,
Benfey, P. A High-Resolution Root Spatiotemporal Map Reveals Dominant Expression
Patterns. *Science* 318(5851), 801-806. http://doi.org/10.1126/science.1146265 (2007).

Brooks, M., Juang, C., Katar,i M. Alvarez, J., Pasquino, A., Shih, H., Huang, J., Shanks, C.,
Cirrone, J., Coruzzi, G. ConnecTF: A platform to integrate transcription factor-gene
interactions   and   validate   regulatory   networks.   *Plant   Physiol.*   185(1),   49-66.
http://doi.org/10.1093/plphys/kiaa012 (2021).

Cabello, J. V., Arce, A. L., Chan, R. L. The homologous HD-Zip I transcription factors
HaHB1 and AtHB13 confer cold tolerance via the induction of pathogenesis related and
glucanase proteins. *Plant J.* 69(1), 141-153. http://doi.org/10.1111/j.1365-
313X.2011.04778.x (2012).

Capella, M., Ribone, P. A., Arce, A. L., Chan, R. L. Arabidopsis thaliana HomeoBox 1
(AtHB1), a homedomain-leucine zipper I (HD-Zip I) transcription factor, is regulated by
phytochrome-interacting factor 1 to promote hypocotyl elongation. *New Phytol.* 207(3),
669-682. http://doi.org/10.1111/nph.13401 (2015).

Casal, J. J., Estevez, J. M. Auxin–environment integration in growth responses to forage
for resources. *CSH Perspect. Biol*. 13(4), a040030.
http://doi.org/10.1101/cshperspect.a040030 (2021).

Cheng, M., Liao, P., Kuo, W., and Lin, T. (2013). The Arabidopsis ETHYLENE RESPONSE
FACTOR1 regulates abiotic stress-responsive gene expression by binding to different cis-
acting elements in response to different stress signals. Plant Physiol. 162, 1566–1582.
doi: 10.1104/pp.113.221911

Choi H., Jeong S., Kim D. S., Na H. J., Ryu J. S., Lee S. S., Nam H. G., Lim P. O., Woo H. R.
The homeodomain-leucine zipper ATHB23, a phytochrome B-interacting protein, is
important for phytochrome B-mediated red light signaling. *Plant Physiol.* 150(2), 308-
320. http://doi.org/10.1111/ppl.12087 (2014).

Datta, S., Prescott, H., Dolan, L. Intensity of a pulse of RSL4 transcription factor
synthesis determines Arabidopsis root hair cell size. *Nat. Plants* 1(15138), 1–6.
https://doi.org/10.1038/nplants.2015.138 (2015).

Denyer, T., Ma, X., Klesen, S., Scacchi, E., Nieselt, K., Timmermans, M. Spatiotemporal
Developmental Trajectories in the Arabidopsis Root Revealed Using High-Throughput
Single-Cell RNA Sequencing. *Dev. Cell* 48(6), 840-852.
https://doi.org/10.1016/j.devcel.2019.02.022 (2019).

Dodt, M., Roehr, J. T., Ahmed, R., Dieterich, C. FLEXBAR—Flexible Barcode and Adapter
Processing for Next-Generation Sequencing Platforms. *Biology* 1(3), 895-905.
https://doi.org/10.3390/biology1030895 (2012).

Doner NM, Seay D, Mehling M, Sun S, Gidda SK, Schmitt K, Braus GH, Ischebeck T,
Chapman KD, Dyer JM, Mullen RT. *Arabidopsis thaliana* EARLY RESPONSIVE TO
DEHYDRATION 7 Localizes to Lipid Droplets via Its Senescence Domain. Front Plant Sci.
2021 Apr 14;12:658961. doi: 10.3389/fpls.2021.658961.

Gao, D., Appiano, M., Huibers, R. P., Chen, X., Loonen, A. E. H. M., Visser, R. G. F., Wolters, B. Y. Activation tagging of ATHB13 in Arabidopsis thaliana confers broad-spectrum disease resistance, *Plant Mol. Biol.* 86(6), 641-653. https://doi.org/10.1007/s11103-014-0253-2 (2014).

Haling, R. E., Brown, L. K., Bengough, A.G., Young, I. M., Hallett, P. D., White, P. J., George, T. S. Root hairs improve root penetration, root–soil con tact, and phosphorus acquisition in soils of different strength. *J. Exp. Bot.* 64(12), 3711–3721. https://doi.org/10.1093/jxb/ert200 (2013).

Han, H., Wang, H., Han, Y., Hu, Z., Xin, M., Peng, H., Yao, Y., Sun, Q., Ni, Z. Altered expression of the *TaRSL2* gene contributed to variation in root hair length during allopolyploid wheat evolution. *Planta* 246(5), 1019–1028. https://doi.org/10.1007/s00425-017-2735-3 (2017).

Han, Y., Xin, M., Huang, K., Yuyun, X., Zhenshan, L., Zhaorong, H., Yingyin, Y., Huiru, P., Zhongfu, N., Qixin, S. Altered expression of *TaRSL4* gene by genome interplay shapes root hair length in allopolyploid wheat. *New Phytol.* 209(2), 721-732. https://doi.org/10.1111/nph.13615 (2016).

Hanson, J., Regan, S., Engstrom, P. The expression pattern of the homeobox gene ATHB13 reveals a conservation of transcriptional regulatory mechanisms between Arabidopsis and hybrid aspen. *Plant Cell Rep*. 21(1), 81-89. https://doi.org/10.1007/s00299-002-0476-6 (2002).

Henriksson, E., Olsson, A. S. B., Johannesson, H., Johansson, H., Hanson, J., Engström, P., Söderman, E. Homeodomain leucine zipper class I genes in Arabidopsis. Expression patterns and phylogenetic relationships. *Plant Physiol*. 139(1), 509–518. https://doi.org/10.1104/pp.105.063461 (2005).

Hwang, Y., Choi, H. S., Cho, H. M., & Cho, H. T. Tracheophytes contain conserved orthologs of a basic helix-loop-helix transcription factor that modulate ROOT HAIR SPECIFIC genes. *Plant Cell* 29(1), 39–53. https://doi.org/10.1105/tpc.16.00732 (2017).

Jia, Z., Giehl, R. F. H., Hartmann, A., Estevez, J. M., Bennett, M. J., von Wiren, N. A spatially concerted epidermal auxin signaling framework steers the root hair foraging response under low nitrogen. *Curr. Biol.* 33(18), 3926–3941. https://doi.org/10.1016/j.cub.2023.08.040 (2023).

1005   Jean-Baptiste, K., McFaline-Figueroa, J. L., Alexandre, C., Dorrity, M., Saunders, L., Bubb,
1006   K., Trapnell, C., Fields, S., Queitsch, C., Cuperus, J. Dynamics of Gene Expression in
1007   Single Root Cells of Arabidopsis thaliana. *Plant Cell* 31(5), 993–1011,
1008   https://doi.org/10.1105/tpc.18.00785 (2019).

1009

1010   Kim, C. M., Dolan, L. ROOT HAIR DEFECTIVE SIX-LIKE class I genes promote root hair
1011   development in the grass *Brachypodium distachyon*. *PLoS Genet.* 12(12), e1006480.
1012   https://doi.org/10.1371/journal.pgen.1006211 (2016).

1013

1014   Liao, Y., Smyth, G. K., Shi, W. The R package Rsubread is easier, faster, cheaper and
1015   better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res*.
1016   47(8). e47. https://doi.org/10.1093/nar/gkz114 (2019).

1017

1018   Lin, Q., Ohashi, Y., Kato, M., Tsuge, T., Gu, H., Qu, L. J., & Aoyama, T. GLABRA2 directly
1019   suppresses basic helix-loop-helix transcription factor genes with diverse functions in
1020   root hair development. *Plant Cell* *27*(10), 2894–2906.
1021   https://doi.org/10.1105/tpc.15.00607 (2015).

1022

1023   Lopez, L. E., Chuah Y. S., Encina, F., Carignani-Sardoy M., Berdion-Gabarain, V., Mutwil,
1024   M., Estevez, J. M. New molecular components that regulate the transcriptional hub in
1025   root hairs: coupling environmental signals with endogenous hormones to coordinate
1026   growth. *J. Exp. Bot*. erad419. https://doi.org/10.1093/jxb/erad419 (2023).

1027

1028   Maher, K. A., Bajic, M., Kajala, K., Reynoso, M., Pauluzzi, G., West, D. A., Zumstein, K.,
1029   Woodhouse, M., Bubb, K., Dorrity, M. W., Queitsch, C., Bailey-Serres, J., Sinha, N., Brady,
1030   S. M., Deal, R. B. Profiling of Accessible Chromatin Regions across Multiple Plant Species
1031   and Cell Types Reveals Common Gene Regulatory Principles and New Control Modules.
1032   *Plant Cell* 30(1), 15–36. https://doi.org/10.1105/tpc.17.00581 (2018).

1033

1034   Mangano, S., Denita-Juarez, S. P., Choi, H. S., Marzol, E., Hwang, Y., Ranocha, P.,
1035   Velasquez, S. M., Borassi, C., Barberini, M. L., Aptekmann, A. A., Muschietti, J. P., Nadra,
1036   A. D., Dunand, C., Cho, H. T., & Estevez, J. M. Molecular link between auxin and ROS-
1037   mediated polar growth. *Proc. Natl. Acad. Sci. U.S.A. 114*(20), 5289–5294.
1038   https://doi.org/10.1073/pnas.1701536114 (2017).

1039

Martínez-Pacheco, J., Song L., Kuběnová, L., Ovečka, M., Berdion-Gabarain, V., Peralta J.
M., Urzúa-Lehuedé, T., Ibeas, M. A., Ricardi, M. M., Zhu, S., Shen, Y., Schepetilnikov, M.,
Ryabova, L. A., Alvarez, J. M., Gutierrez, R. A., Grossmann, G., Šamaj, J., Yu, F., Estevez J.
M. Cell surface receptor kinase FERONIA linked to nutrient sensor TORC signaling
controls root hair growth at low temperature linked to low nitrate in Arabidopsis
thaliana. *New Phytol*. 238(1), 169-185.https://doi.org/10.1111/nph.18723 (2023).

1046

Martínez-Pacheco, J., Ranocha, P., Kasulin, L., Fusari, C. M., Servi, L., Aptekmann, A. A.,
Berdion-Gabarain, V., Peralta J. M., Borassi, C., Marzol, E., Rodríguez-Garcia D. R.,
Rondón-Guerrero, Y. C., Carignani-Sardoy, M., Ferrero, L., Botto, J. F., Meneses, C., Ariel,
F., Nadra, A. D., Petrillo, E., Dunand C., Estevez J.M. Apoplastic class III peroxidases
PRX62 and PRX69 promote Arabidopsis root hair growth at low temperature. *Nat.*
Commun. 13(1), 1310. https://doi.org/10.1038/s41467-022-28833-4 (2022).

1053

1054 Martínez Pacheco, J., Mansilla, N., Moison, M., Lucero, L., Berdion-Gabarain, V., Ariel, F.,
1055 & Estevez, J. M. The lncRNA APOLO and the transcription factor WRKY42 target
1056 common cell wall EXTENSIN encoding genes to trigger root hair cell elongation. *Plant*
1057 *Signal. Behav.* 16(8), 1920191. https://doi.org/10.1080/15592324.2021.1920191 (2021).

1058

1059 Menand, B., Yi, K., Jouannic, S., Hoffmann, L., Ryan, E., Linstead, P., Schaefer, D. G.,
1060 Dolan, L. An ancient mechanism controls the development of cells with a rooting
1061 function in land plants. *Science* 316(5830), 1477–1480.
1062 https://doi.org/10.1126/science.1142618 (2007).

1063

1064 Miguel, M. A., Postma, J. A., Lynch, J. P. Phene synergism between root hair length and
1065 basal root growth angle for phosphorus acquisition. *Plant Physiol* 167(4), 1430–1439.
1066 https://doi.org/10.1104/pp.15.00145 (2015).

1067

1068 Moison, M., Pacheco, J. M., Lucero, L., Fonouni-Farde, C., Rodríguez-Melo, J., Mansilla,
1069 N., Christ, A., Bazin, J., Benhamed, M., Ibañez, F., Crespi, M., Estevez, J. M., & Ariel, F.
1070 The lncRNA APOLO interacts with the transcription factor WRKY42 to trigger root hair
1071 cell expansion in response to cold. *Mol. Plant* 14(6), 937-948.
1072 https://doi.org/10.1016/j.molp.2021.03.008 (2021).

1073

1074 Moon, S., Cho, L. H., Kim, Y. J., *et al*. RSL Class II transcription factors guide the nuclear
1075 localization of RHL1 to regulate root hair development. *Plant Physiol* 179(2), 558–568.
1076 https://doi.org/10.1104/pp.18.01002 (2019).

1077

1078 Nolan TM, Vukašinović N, Hsu CW, Zhang J, Vanhoutte I, Shahan R, Taylor IW,
1079 Greenstreet L, Heitz M, Afanassiev A, Wang P, Szekely P, Brosnan A, Yin Y, Schiebinger G,
1080 Ohler U, Russinova E, Benfey PN. Brassinosteroid gene regulatory networks at cellular
1081 resolution in the *Arabidopsis* root. Science. 2023 Mar 31;379(6639):eadf4721. doi:
1082 10.1126/science.adf4721.

1083

1084 O'Malley, R. C., Shao-shan, C. H., Liang, S., Mathew, G. L., Bartlett, A., Nery, J. R., Galli,
1085 M., Gallavotti, A., Ecker J. R. Cistrome and Epicistrome Features Shape the Regulatory

1086   DNA Landscape. *Cell* 165(5), 1280-1292. http://dx.doi.org/10.1016/j.cell.2016.04.038
1087   (2016).

1088

1089   Perotti, M. F., Ribone, P. A., Chan, R. L. Plant transcription factors from the
1090   Homeodomain-Leucine Zipper family I. Role in development and stress responses.
1091   *IUBMB Life* 69(5), 280-289. https://doi.org/10.1002/iub.1619 (2017).

1092

1093   Perotti M.F., Ribone P.A., Cabello J.V., Ariel F.D., Chan R.L. AtHB23 participates in the
1094   gene regulatory network controlling root branching, and reveals differences between
1095   secondary and tertiary roots. *Plant J.* 100(6), 1224-1236.
1096   https://doi.org/10.1111/tpj.14511 (2019).

1097

1098   Perotti, M. F., Ariel, F. D., Chan, R. L. Lateral root development differs between main
1099   and secondary roots and depends on the ecotype. *Plant Signal. Behav.* 15(6), 1755504*.*
1100   https://doi.org/10.1080/15592324.2020.1755504 (2020).

1101

1102   Perotti, M. F., Arce, A. L., Chan, R. L. The underground life of homeodomain-leucine
1103   zipper transcription factors. *J. Exp. Bot*. 72(11), 4005–4021.
1104   https://doi.org/10.1093/jxb/erab112 (2021).

1105

1106   Perotti, M. F., Arce, A. L., Ariel, F. D., Figueroa, C. M., Chan, R. L. The transcription factor
1107   AtHB23 modulates starch turnover for root development and plant survival under
1108   salinity. *Environ. Exp. Bot.* 201(11), 104994.
1109   https://doi.org/10.1016/j.envexpbot.2022.104994 (2022).

1110

1111   Perotti, M. F., Arce, A. L., Chan, R. L. The underground life of homeodomain-leucine
1112   zipper transcription factors. J. Exp. Bot. 72(11), 4005–4021.
1113   https://doi.org/10.1093/jxb/erab112 (2021).

1114

1115   Proust, H., Honkanen, S., Jones, V. A., Morieri, G., Prescott, H., Kelly, S., Ishizaki, K.,
1116   Kohchi, T., Dolan, L. *RSL* class I genes controlled the development of epidermal
1117   structures in the common ancestor of land plants. *Curr. Biol.* 26(1), 93–99.
1118   https://doi.org/10.1016/j.cub.2015.11.042 (2016).

1119

1120   Rasmussen, S., Barah, P., Suarez-Rodriguez, M. C., Bressendorff, S., Friis, P., Costantino,
1121   P., et al. (2013). Transcriptome responses to combinations of stresses in Arabidopsis.
1122   Plant Physiol. 161, 1783–1794. doi: 10.1104/pp.112.21077

1123

1124   Ribone, P., Capella, M., Chan, R. Functional characterization of the homeodomain
1125   leucine zipper I transcription factor AtHB13 reveals a crucial role in Arabidopsis

development. *J. Exp. Bot.* 66(19), 5929-5943. https://doi.org/10.1093/jxb/erv302 (2015).

Riechmann, J. L., Heard, J., Martin, G. ,Reuber L., Jiang C.,Keddie J., Adam L., Pineda O.,Ratcliffe O.J., Samaha R.R., Creelman R., Pilgrim M., Broun P., Zhang J. Z., Ghandehari D., Sherman B.K. , Yu G. Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290(5499), 2105–2110. https://doi.org/10.1126/science.290.5499.2105 (2000).

Robinson, M. D., McCarthy, D. J., Smyth, G.K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1), 139-40. https://doi.org/10.1093/bioinformatics/btp616 (2010).

Rubio, V., Linhares, F., Solano, R., Martín, A. C., Iglesias, J., Leyva, A., Paz-Ares, J. A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes Dev*. 15(16), 2122–2133. https://doi.org/10.1101/gad.204401 (2001).

Ryu, K. H., Huang, L., Kang, H., Schiefelbein, J. Single-Cell RNA Sequencing Resolves Molecular Relationships Among Individual Plant Cells. *Plant Physiol*. 179(4), 1444-1456 https://doi.org/10.1104/pp.18.01482 (2019).

Silva, A. T., Ribone, P. A., Chan, R. L., Ligterink, W., Hilhorst, H. W. M. A predictive coexpression network identifies novel genes controlling the seed-to-seedling phase transition in Arabidopsis thaliana. *Plant Physiol*. 170(4), 2218–2231. https://doi.org/10.1104/pp.15.01704 (2016).

Silva-Navas, J., Moreno-Risueno, M. A., Manzano, C., Téllez-Robledo, B., Navarro-Neila, S., Carrasco, V., Pollmann, S., Gallego, F. J., del Pozo, J. C. Flavonols mediate root phototropism and growth through regulation of proliferation to-differentiation transition. *Plant Cell* 28(6), 1372–1387. https://doi.org/10.1105/tpc.15.00857 (2016).

Silva-Navas, J., Moreno-Risueno, M. A., Manzano, C., Pallero-Baena, M., Navarro-Neila, S., Téllez-Robledo, B., Garcia-Mina, J. M., Baigorri, R., Gallego, F. J., del Pozo, J. C. D-Root: a system for cultivating plants with the roots in darkness or under different light conditions. *Plant J.* 84(1), 244-255. https://doi.org/10.1111/tpj.12998 (2015).

Shibata, M., Breuer, C., Kawamura, A., Clark, N. M., Rymen, B., Braidwood, L., Morohashi, K., Busch, W., Benfey, P. N., Sozzani, R., Keiko, S. GTL1 and DF1 regulate root hair growth through transcriptional repression of ROOT HAIR DEFECTIVE 6-LIKE 4 in

Arabidopsis. *Development* 145(3), dev159707. https://doi.org/10.1242/dev.159707 (2018).

Shibata, M., Favero, D. S., Takebayashi, R., Takebayashi, A., Kawamura, A., Rymen, B., Hosokawa, Y., Sugimoto, K. Trihelix transcription factors GTL1 and DF1 prevent aberrant root hair formation in an excess nutrient condition. *New Phytol.* 235(4), 1426-1441. https://doi.org/10.1111/nph.18255 (2022).

Shulse, C. N., Cole, B. J., Ciobanu, D., Lin, J., Dickel, D. E. High-throughput single-cell transcriptome profiling of plant cell types. *Cell Rep.* 27(7), 2241-2247. https://doi.org/10.1016/j.celrep.2019.04.054 (2019).

Spies, F. P., Raineri, J., Miguel, V. N., Cho, Y., Hong, J. C., Chan, R. L. The Arabidopsis transcription factors AtPHL1 and AtHB23 act together promoting carbohydrate transport from pedicel-silique nodes to seeds. *Plant Sci*. 315, 111133. https://doi.org/10.1016/j.plantsci.2021.111133 (2022).

Spies, F. P., Perotti, M. F., Cho, Y., Ig Jo, C., Hong J. C., Chan, R. L. A complex tissue-specific interplay between the Arabidopsis transcription factors AtMYB68, AtHB23, and AtPHL1 modulates primary and lateral root development and adaptation to salinity. *Plant J.* 115(4), 952-966. https://doi.org/10.1111/tpj.16273 (2023).

Vijayakumar, P., Datta, S. & Dolan, L. ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes root hair elongation by transcriptionally regulating the expression of genes required for cell growth. *New Phytol.* 212(4), 944-953. https://doi.org/10.1111/nph.14095 (2016).

Viola, I. L., Gonzalez, D. H., Chapter 6. Structure and evolution of plant homeobox genes, in: D.H. Gonzalez (Ed.), Plant Transcription Factors, Elsevier, 2016, pp. 101–112. https://doi.org/10.1016/B978-0-12-800854-6.00006-3 (2016).

Wu, J., Kurten, E. L., Monshausen, G., Hummel, G. M., Gilroy, S., Baldwin, I. T. NaRALF, a peptide signal essential for the regulation of root hair tip apoplastic pH in *Nicotiana attenuata*, is required for root hair development and plant growth in native soils. *Plant J.* 52(5), 877-890. https://doi.org/10.1111/j.1365-313X.2007.03289.x (2007).

Yi, K., Menand, B., Bell, E., Dolan, L. A basic helix-loop-helix transcription factor controls cell growth and size in root hairs. *Nat. Genet.* 42(3), 264–267. https://doi.org/10.1038/ng.529 (2010).

1205   Yu, G., Wang, L. G., Han, Y., He, Q. Y. ClusterProfiler: an R package for comparing
1206   biological themes among gene clusters. *OMICS*. 16(5), 284-287.
1207   https://doi.org/10.1089/omi.2011.0118 (2012).

1208

1209   Zhang, T., Xu, Z.,Shang, G.,Wang, J. A Single-Cell RNA Sequencing Profiles the
1210   Developmental Landscape of Arabidopsis Root. *Mol. Plant* 12(5), 648-660.
1211   https://doi.org/10.1016/j.molp.2019.04.004 (2019).

1212

1213   Zhu, S., Estevez, J. M., Liao, H., Zhu, Y., Yang, T., Li, C., Wang, Y., Li, L., Liu, X., Martinez-
1214   Pacheco, J., Guo, H., Yu, F. The RALF1–FERONIA Complex Phosphorylates eIF4E1 to
1215   Promote Protein Synthesis and Polar Root Hair Growth. *Mol. Plant* 13(5), 698-716.
1216   https://doi.org/10.1016/j.molp.2019.12.014 (2020a).

1217

1218   Zhu, S., Martínez-Pacheco, J., Estevez, J. M. & Yu, F. Autocrine regulation of root hair
1219   size by the RALF-FERONIA-RSL4 signaling pathway. *New Phytol*. 227(1), 45-49.
1220   https://doi.org/10.1111/nph.16497 (2020b).