ScRNA-seq and scATAC-seq reveal that sertoli cell mediate spermatogenesis disorders through stage-specific communications in non-obstructive azoospermia

Shimin Wang[1,2,#,*], Hongxian Wang[3,#], Bicheng Jin[4], Hongli Yan[5], Qingliang Zheng[1,*], Dong Zhao[2,*]

[1]Prenatal Diagnosis Center, The Eighth Affiliated Hospital, Sun Yat-sen University, 3025# Shennan Road, Shenzhen 518000, China

[2]Department of Gynaecology and Obstetrics, Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 Zhizaoju Road, Shanghai 200011, China

[3]Department of Urology and Andrology, School of Medicine, Renji Hospital, Shanghai Jiao Tong University, Shanghai, China

[4]Department of Surgical subject, Guizhou Electric staff Hospital, Guiyang city, Guizhou province, 55000, China

[5]Reproductive medicine center, the Navy Medical University, Shanghai, China

[#]**Equal contributors**

*Corresponding author:

Shimin Wang, Email: wangsm_2003@163.com

Detailed address: The Eighth Affiliated Hospital, Sun Yat-sen University, 3025# Shennan Road, Shenzhen 518000, China. Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 Zhizaoju Road, Shanghai 200011, China.

Qingliang Zheng, Email: jackie4075@126.com

Detailed address: The Eighth Affiliated Hospital, Sun Yat-sen University, 3025# Shennan Road, Shenzhen 518000, China.

Dong Zhao, Email: hendryz@gmail.com

Detailed address: Ninth People's Hospital, Shanghai Jiao Tong University School of

Medicine, 639 Zhizaoju Road, Shanghai 200011, China.

Abstract

Non-obstructive azoospermia (NOA) belongs to male infertility due to

spermatogenesis failure. However, evidence for cell type-specific abnormalities of

spermatogenesis disorders in NOA remains lacking. We performed single-cell RNA

sequencing (scRNA-seq) and single-cell assay for transposase-accessible chromatin

sequencing (scATAC-seq) on testicular tissues from patients with obstructive

azoospermia(OA) and NOA. HE staining confirmed the structural abnormalities of

the seminiferous tubules in NOA patients. We identified 12 germ cell subtypes

(spermatogonial stem cell-0 (SSC0), SSC1, SSC2, diffing-spermatogonia

(Diffing-SPG), diffed-spermatogonia (Diffed-SPG), pre-leptotene (Pre-Lep),

leptotene-zygotene (L-Z), pachytene (Pa), diplotene (Di), spermatids-1 (SPT1), SPT2,

and SPT3) and 8 Sertoli cell subtypes (SC1-SC8). Among them, three novel Sertoli

cell subtypes phenotypes were identified, namely SC4/immature, SC7/mature, and

SC8/further mature Sertoli cells. For each germ or Sertoli cell subtype, we identified

unique new markers, among which immunofluorescence confirmed co-localization of

ST3GAL4, A2M, ASB9, and TEX19 and DDX4 (classical marker of germ cell).

PRAP1, BST2, and CCDC62 were co-expressed with SOX9 (classical marker of

Sertoli cell) in testes tissues also confirmed by immunofluorescence. The interaction

between germ cell subtypes and Sertoli cell subtypes exhibits stage-specific-matching

pattern, as evidenced by SC1/2/5/7 involving in SSC0-2 development, SC3

participating in the whole process of spermiogenesis, SC4/6 participating in Diffing

and Diffed-SPG development, and SC8 involving in the final stage of SPT3. This

pattern of specific interactions between subtypes of germ cell and Sertoli cell was confirmed by immunofluorescence of novel markers in testes tissues. The interaction was mainly contributed by Notch1/2/3 signaling. Our study profiled the single-cell transcriptome of human spermatogenesis and provided many potentials molecular markers for developing testicular puncture specific marker kits for NOA patients.

Keywords: Non-obstructive azoospermia, Single cell RNA-sequencing, Single cell ATAC-sequencing, Germ cell, Sertoli cell

Impact statement: The specific interactions between Sertoli cell subtypes and different stages of germ subgroups maintain normal development of germ cells. SC3 subtype absence may be the key factor leading to NOA.

Introduction

Infertility is a common disease affecting approximately 15% of couples, and 50% of cases in infertility is attributed to male factor infertility (Agarwal, Mulgund, Hamada, & Chyatte, 2015). Male infertility is mainly associated with sexual dysfunction, endocrine, varicocele, and reproductive system infection(Naz & Kamal, 2017).Azoospermia includes obstructive azoospermia (OA) and non-obstructive azoospermia (NOA). NOA accounts for about 1% of adult males and 10%-15% of infertile males (Fakhro et al., 2018). The etiology of NOA is varied, including cryptorchidism, post-pubertal mumps orchitis, and prior testicular torsion (Yao et al., 2022). At present, assisted reproductive technology, such as microdissection testicular sperm extraction and intracytoplasmic sperm injection, has made certain progress, providing help for patients to obtain biological offspring. At the same time, spermatogonial stem cell culture and transplantation technology and testicular tissue

transplantation technology has also made some progress, but the clinical application is

still not mature (Gassei & Orwig, 2016). At present, the etiology and mechanism of

many NOA patients are still unknown, which poses a challenge to treatment.

Therefore, it is particularly urgent to further study the molecular mechanisms

affecting spermatogenesis in NOA patients.

The formation of mature spermatozoa can be delineated into three stages: the

proliferation of spermatogonial cells, meiotic division of spermatocytes, and the

maturation of sperm cells (Jegou, 1993). The seminiferous tubule comprises three

distinct cell types: Sertoli cells, extending from the basal membrane to the tubular

lumen; Germ cells of various generations; Peritubular cells, surrounding sertoli cells

and germ cells, isolated from sertoli cells by the extracellular matrix. In most

mammals, germ cells undergo continuous renewal, while sertoli cells cease division

during the pubertal developmental period, forming the seminiferous epithelium. The

lining of the seminiferous epithelium is uniquely intricate within a complex tissue

structure. At any given point in a seminiferous tubule, several generations of germ

cells develop simultaneously in the process of contacting sertoli cells from the basal

epithelium to the apex. The evolution of each generation of germ cells is strictly

synchronized with others, resulting in the formation of specific cell associations or

stages at a particular segment of the tubule. The complete temporal sequence of these

cell associations or stages until the reappearance of the initial association in a defined

region of the tubule is referred to as the seminiferous epithelial cycle (Hess & Moore,

1993). Adult spermatogonial stem cells are distributed along the basal membrane at

the base of the seminiferous tubule, necessitating a delicate balance between

self-renewal and differentiation. Throughout this process, only sertoli cells

consistently maintain physical contact with sperm cells. Consequently, we posit that

the entire process of spermatogenesis inevitably involves intense information

exchange with the occurrence of sertoli cells. However, the specifics of this

interactive process remain incompletely understood at present.

Single-cell transcriptome sequencing (scRNA-seq) technology is used for

high-throughput molecular detection of a single cell and exploring the whole gene

expression profiles of a single cell (Tan, Song, & Wilkinson, 2020). It can delineate

the existence of rare cell subsets and the degree of cellular heterogeneity (Vertesy et

al., 2018). Liao et al. (2019) performed scRNA-seq analysis on mouse testicular germ

cells at 5.5 days after birth, revealing the heterogeneity of gene expression in

undifferentiated spermatogonial cells. Guo et al. (2018) performed scRNA-seq of

testicular cells from young adults and found that there were five different

spermatogonial states accompanying human spermatogonial differentiation. In

addition, scATAC-seq is a widely used method to obtain a genome-wide snapshot of

chromatin accessibility, signatures of active transcription and transcription factor (TF)

binding (Mimitou et al., 2021). Cellular identity is strongly affected by the epigenetic

wiring of the cell, which is measured by scATAC-seq (Pervolarakis et al., 2020).

Therefore, scRNA-seq and scATAC-seq can be used to analyze spermatogenesis.

In this study, we performed an integrative analysis of scRNA-seq and

scATAC-seq in testicular tissues from two OA and three NOA patients. Cellular

heterogeneity and marker gene identification results were analyzed, and the potential

functional mechanism was inferred. Altogether, our results provide comprehensive

understanding of the maturation of the spermatogenic microenvironment and the

mechanisms underlying pathogenesis, offering novel targets for NOA treatment

strategies.

Material and methods

Human sample acquisition

Testicular tissues were obtained from two OA patients (OA1 and OA2) and three NOA patients (NOA1, NOA2, NOA3). Notably, the sperm concentration of the three NOA patients varied in descending order (NOA1 > NOA2 > NOA3), with NOA3 being the patient with complete absence of sperm. Patient clinical and laboratory characteristics are presented in Table 1. This study was approved by the Ethics Committee of School of Medicine, Renji Hospital, Shanghai Jiao Tong University [approval number: KY2020-193]. All patients provided written informed consent.

Table 1 Patient characteristics in this study.

	OA1	OA2	NOA1	NOA2	NOA3
Demographics					
Age	26	30	32	29	26
BMI	20.2	25.8	24.2	28.7	25.4
Ob/Gyn History	No/0	Yes/one son	Yes/0	Yes/0	Yes/0
Smoking	Quit smoking for 3 months	10 years	No	No	Quit smoking for 6 months
Alcohol use	Yes	Yes	Yes	No	No
Hormone profile					
FSH (mIU/mL)	2.98	2.2	20.96	30.86	9.21
LH (mIU/mL)	6.86	4.3	7.64	8.72	5.13
PRL (mIU)	291	329	-	235.7	257.7
T (ng/mL)	5.3	13.57	3.46	4.18	1.53
E2 (pg/mL)	29.35	125.2	31	30.67	31.7

Volume of testis (mL)					
Left	15.1	13.3	6.9	3.9	5.43
Right	14.2	16.1	5.3	5.8	6.47
Spermatic cord					
Left	1.9 mm; regurgitation; 3 s	Non-regurgitation	1.5 mm; Non-regurgitation	2.5 mm; regurgitation; 5 s	1.5 mm; Non-regurgitation
Right	1.5 mm; Non-regurgitation	Non-regurgitation	1.5 mm; Non-regurgitation	Non-regurgitation	1.5 mm; Non-regurgitation
Other					
Ejaculatory duct	Widening (inner diameter approx. 2mm)	Normal	Normal	Normal	Normal
AZF	Normal	Normal	Normal	Normal	Normal

Sample dissociation

Testicular tissues were washed three times in Dulbecco's PBS (Gibco, Waltham, MA, USA) and the tunica albugineas of testes were removed. Testicular tissues were cut into small pieces of 1–2 mm and digested in 10 mL DMEM: F12 media (Gibco, Waltham, MA, USA) containing 1 mg/mL type IV collagenase for 15 min on a rotor at 37°C. Subsequently, tubules were washed again and incubated with 10 mL F12 media containing 500 μg/mL DNase I and 200 μg/mL trypsin for 15 minutes at 37°C. The digestion was stopped with the addition of fetal bovine serum (FBS) (Gibco, Waltham, MA, USA). The single cells were obtained by filtering through 40 μm cell-strainer. The cells were resuspended in 1 mL DPBS and 5 mL red blood cell lysis buffer to remove red blood cells. After that, the cells were washed with DPBS and

cryopreserved in DPBS with 1% FBS (Gibco, Waltham, MA, USA) until further use.

For scRNA-seq and scATAC-seq assay, the cryopreserved cells were thawed and

incubated with a Dead Cell Removal Kit (Miltenyi Biotec) to clear dead/stressed cells

in accordance with manufacturer's instruction.

scRNA-seq library preparation

scRNA-seq was carried out using 10x Chromium Single Cell 3' Reagent referring to

manufacturer's instruction. In brief, viable single cells were resuspended in PBS with

0.04% bovine serum albumin (BSA; Sigma) and counted using the hemocytometer.

Suspensions containing 5000 to 8000 cells cells per sample were mixed with RT-PCR

reaction and loaded into the Single Cell Chip B and processed through the 10x

controller for droplet production. After that, in-drop lysis and reverse transcription

occurs and mRNA transcripts from single cells were barcoded to determine the cell

origin. Following reverse transcription, barcoded cDNAs were purified, amplified by

12 cycles of PCR, end-repaired, and ligated with Illumina adapters. The final libraries

were sequenced on the Illumina Novaseq 6000 platform with paired end 150 bp

sequencing. Each sample was sequenced with depth of 100 G of raw reads.

scRNA-seq data processing and analysis

The scRNA-seq data produced by Illumina NovaSeq 6000 sequencing were processed

and mapped to human reference genome hg38. After mapping, the outputs were

processed using the Seurat package in R. To filter out low-quality cells, we referred to

the following criteria: cells with few genes per cell (<50) or plenty of molecules per

cell (>20,000); cells with over 30% of mitochondrial genes. Normalization was

carried out in accordance with the package manual (https://satij alab.org/seura t/v3.1/).

We identified cell clusters with Uniform Manifold Approximation and Projection

(UMAP). The differentially expressed genes were analyzed by the FindMarkers

function. Gene set enrichment analysis was carried out using clusterProfiler R

package. Regulatory gene network analysis was carried out by SCENIC.

Pseudo-time analysis

We applied the Single-Cell Trajectories analysis utilizing Monocle2

(http://cole-trapnell-lab.github.io/monocle-release) using DDR-Tree and default

parameter. Before Monocle analysis, we select marker genes of the Seurat clustering

result and raw expression counts of the cell passed filtering. Based on the pseudo-time

analysis, branch expression analysis modeling (BEAM Analysis) was applied for

branch fate determined gene analysis.

Cell communication analysis

To enable a systematic analysis of cell–cell communication molecules, we applied cell

communication analysis based on the CellPhoneDB, a public repository of ligands,

receptors and their interactions. Membrane, secreted and peripheral proteins of the

cluster of different time point was annotated. Significant mean and Cell

Communication significance (p-value<0.05) was calculated based on the interaction

and the normalized cell matrix achieved by Seurat Normalization.

SCENIC analysis

To assess transcription factor regulation strength, we applied the Single-cell

regulatory network inference and clustering (pySCENIC, v0.9.5) workflow, using the

20-thousand motifs database for RcisTarget and GRNboost.

scATAC-seq library preparation

scATAC-seq was conducted using 10x Chromium Single Cell ATAC Reagent (V1.1

chemistry) in accordance with manufacturer's instruction. In short, cells were

resuspended in PBS with 0.04% BSA (Sigma) and counted. Next, cells were cultured

with cold lysis buffer on ice for 5 min. Tn5 transposase reaction was performed using

the Tagment DNA Enzyme 1 (Illumina) at 37 °C for 30 min. Cells were processed for

targeting a recovery of 10,000 cells per sample and amplification were performed

with 12 cycles of PCR for library construction. The libraries were sequenced on

Illumina Novaseq 6000 platform and each sample was sequenced with depth of 200M

raw reads.

scATAC-seq data preprocessing and analysis

The scATAC-seq data analysis was performed as previously described. In brief, the

sequence reads were mapped to the human reference genome hg38. The R package

Signac was used to process the generated outputs. The Latent Semantic indexing

algorithm, T-distributed Stochastic Neighbor Embedding algorithm and Uniform

Manifold Approximation algorithm are used for dimension reduction and information

display. Each Motif of the cell can be annotated by associating identified open

chromatin sites with the JASPAR database. By using CHIPseeker, the distribution of

peak in different functional regions of the genome was annotated and statistically

analyzed. Prediction of results for each cell type in scATAC-seq can be achieved by

combining scRNA-seq data with scATAC-seq using KNN.

Immunofluorescence

The testicular tissues were fixed in 4% paraformaldehyde at room temperature for

30min, permeabilized with 0.5% Triton X-100, and blocked with 1% bovine serum

albumin (GIBCO, USA). The samples were incubated overnight at 4 °C with primary

antibodies, including ST3GAL4 (13546-1-AP, Proteintech), A2M (ab109422, Abcam),

DDX4 (ab270534, abcam), ASB9 (22728, SAB), TEX19 (AF6319, R&D), TSSK6

(H00083983-M02, Novus), PRAP1 (11932-1-AP, Proteintech), BST2

(H00000684-B02P, Novus), SOX9 (ab185966, Abcam), CCDC62 (25981-1-AP,

Proteintech),TF (ab185966, Abcam), SOX2 (ab92494, Abcam), ASPATS1

(NBP2-31037, Novus), C9orf57 (orb156207, Biorbyt), TSC21 (NBP1-91727, Novus),

BEND4 (24711-1-AP, Proteintech), and SMCP (NBP1-81252, Novus). After washing

with PBS, the samples were incubated with a secondary antibody (Abcam) for 1 h at

room temperature. DAPI (Cell Signaling, MA, USA) was used to stain cell nucleus.

After washing 3 time with PBS, the sections were performed for photograph under a

fluorescence microscope (Carl Zeiss, Oberkochen, Germany).

Results

scRNA-seq analysis of human testicular tissues

Testicular histology was examined by HE staining. As shown in Figure 1A, in

comparison with the OA group with normal structure of seminiferous epithelium, the

NOA groups had twisted and atrophied, sparsely arranged seminiferous tubules, and

almost all of seminiferous tubules were vacuolated in transverse section with a thin

layer of basal chambers containing supporting cells. To characterize the diversity of

testicular cells, we performed scRNA-seq on testicular samples from three NOA

patients and two OA patients (Figure 1B). After removing low quality cells, we

obtained 23889 cells for scRNA-seq (Table S1). Notably, the quantity of annotated

germ cells in patients categorized as NOA1, NOA2, and NOA3 gradually decreased

to zero (Table S1). This observation is in complete accordance with the clinical

features exhibited by the three patients, providing evidence for the reliability of our

scRNA-seq data. UMAP analysis of scRNA-seq revealed 12 major cell types,

including germ cells, Leydig cells, Sertoli cells, endothelial cells, peritubular myoid

cells (PMCs), smooth muscle, schwann cell, macrophage, mast cells, T cells, B cells,

and plasma cells (Figure 1C). A total of 17,571 high quality cells were obtained from

scATAC-seq analysis, and they were also annotated with B cells, endothelial cells,

PMCs, schwann cell, smooth muscle, T cells, germ cells, Leydig cells, macrophage, and Sertoli cells, which was similar to the scRNA-seq results (Figure 1D). Heatmap of marker genes in scRNA-seq data revealed that cell heterogeneity was obvious (Figure 1E). The well-known cell type markers were utilized to determine the cell clusters (Figure 1F). For instance, we observed that sertoli cells, in addition to specifically expressing classical markers AMH and SOX9 (Fröjdman, Harley, Pelliniemi, & biology, 2000; Lasala et al., 2011), also exhibited strong expression of APOA1 (Figure 1F). Furthermore, we identified that the PMCs not only expressed the classical marker MYH11 (Lottrup et al., 2014), which is concurrently highly expressed in smooth muscle cells, displaying limited specificity, but also demonstrated higher specificity in the expression of DPEP1, suggesting its potential as a specific marker for PMCs. These findings indicate that the scRNA-seq data from this study not only serve for cellular classification but also contribute to the identification of novel potential molecular markers for distinct cell types.

Germ cell subtypes reveal distinct molecular features

To explore the heterogeneity of germ cells during normal development, we re-clustered the germ cells and identified 12 subpopulations, including spermatogonial stem cell-0 (SSC0), SSC1, SSC2, and Diffing_SPG, Diffed_SPG, Pre-Leptotene, Leptotene_Zygotene, Pachytene, Diplotene, SPT1, SPT2, and SPT3 (Figure 2A). Heatmap of scRNA-seq results showed the differentially expressed markers seems cannot distinguish between different germ cell subpopulations (Figure S1A), due to the fact that some of the similar subtypes share the same markers with each other, as shown in violin plots (Figure S1B). For example, C19orf84, PIWIL4, ST3GAL4, UTF1, ID4, and A2M were expressed on SSC0 and SSC1 subtypes, ITGB1 and L1TD1 were expressed SSC0, SSC1, SSC2 (Figure S1B). To elucidate this

phenomenon, we attempted to characterize the sequential developmental relationships

among 12 germ cell subpopulations through pseudotime analysis, approaching this

issue from a developmental perspective. As shown in Figure 2B, germ cells could be

divided into three states, which was in accordance with the biological sequence of

continuous transition of spermatogenesis. States 2 and 3 primarily consist of

SSC-Diffed SPG cells, and these cells are not completely segregated in terms of the

cell states, particularly with SSC0-SSC3 positioned at two initial branches without

complete separation (Figure 2B, Figure S1C). This suggests a potential cyclical and

self-renewing state that is not entirely independent between SSC1 and SSC2. Cells

from the Pre-L to SPT3 stages all belong to state 1, following the developmental

chronological order of sperm generation. Therefore, the phenomenon of distinct

subpopulations existing independently while sharing markers with other

subpopulations suggests that germ cells from different subtypes may not have fully

differentiated into mature states during the course of evolution, remaining in an

intermediate state of continuous differentiation.

Furthermore, we presented the expression distribution of markers for different

subpopulations across three developmental stages in the chronological order of germ

cell maturation (Figure 2C). Interestingly, the positions of different subpopulations of

markers on UMAP show a trend of right-to-left panning as the arrows point to the

temporal developmental order of sperm cells, which was supported by RNA velocity

stream UMAP (Figure 2D). Among them, SPT1 cells can be clearly distinguished

from SPT2-3 cells using FAM24A, but between SPT2 and SPT3 cells, there is no

precise and specific marker that completely segregates these two cell types. We

speculate that this lack of differentiation may be due to the intense morphological

changes occurring in the sperm cells during this period, resulting in relatively minor

differences in gene expression.

Finally, to further validate the reliability of the identified new potential markers of

germ cells, we co-stained them with a classical marker DDX4 using

immunofluorescence (Figure 2E). We discovered that ST3GAL4, A2M, and DDX4

were co-expressed in testes tissues. ASB9 and TEX19 were also co-expressed with

DDX4 were in testes tissues. TSSK6 exhibited similar expression pattern to that of

DDX4 in testes tissues. In addition, closely resembled their expression patterns in

different subpopulations, we found that the region transcription start/promoter site to

A2M and ASB9 was accessible in the SSC0, SSC2, and Diffing_SPG but was not

accessible in other cell subtypes, with much higher levels of openness (Figure 2F).

The region transcription start site to TEX19 was accessible in the Diffed_SPG but was

not accessible in other cell subtypes, which also closely resembled TEX19 expression

pattern. Collectively, these results suggest that these new markers localize to testicular

tissue and have the same ability to label sperm cells as DDX4.

Cell cycle and transcriptional dynamics analysis of Germ cell differentiation

Most adult tissues are sustained by resident adult stem cells responsible for

maintaining tissue function and integrity. As aging or damaged cells undergo

apoptosis, resident adult stem cells are activated to generate new cells. Within various

tissues, a dichotomy exists between rapidly cycling (involved in tissue repair, EOMES

$^{-}$GFRA1^{+}) and quiescent stem cells (reserve cells, EOMES^{+}GFRA1^{+}) (Sharma et al.,

2018). To investigate the cellular developmental states of distinct sperm

subpopulations, we conducted an analysis of the cell cycle in sperm cells. As shown in

Figure 3A and 3B, SSC0/1 and SSC2 were quiescent stem cells and rapidly cycling

cells, respectively. Furthermore, to explore the biological function of germ cells, we

used transcriptional dynamics to analyze the important events and time points in the

whole transcriptional process of spermatogenesis. Maintenance of SSC pluripotency,

whether in the G0/G1 phase and G2/M arrest SSC0/SSC1, or rapid cycling SSC2, are

transcriptionally governed by the downregulation of ID4 (Figure 3C). The pivotal

driving genes promoting the transition of SSC2 into differentiating Diffing SPG cells

and progressing through mitosis are upregulated DMRT1 and PABPC4 (Figure 3C).

As BEND2 takes the central regulatory position, germ cell development officially

proceeds into meiosis, marking the transitioned cells (Diffed SPG) in a mixed

transitional state concurrently undergoing both mitosis and meiosis (Figure 3C). The

molecular markers distinguishing between the PreLep and SPT3 subpopulations

included C18orf63, MNS1, CCDC42, LDHC, TMIGD3, AC113189.2, and FLJ40194

(Figure 3C). Results showed that there is no distinctly discernible regulatory factor

between PreLep and L-Z, with C18orf63 being a crucial differentiator, upregulated in

PreLep and downregulated in L-Z. Finally, we summarized the spermatogenesis at

each stage and major events in the cell cycle (Figure 3D).

Subsequently, Beam analysis was performed to unveil the critical transcription

factors involved in sperm differentiation. Figure S2A hierarchically clustered the top

50 transcription factors into three groups, presenting an expression heatmap of

fate-determining genes associated with the differentiation from SSC0 to SPG state.

The expression pattern of transcription factors within cluster 1 gradually increased

along the temporal trajectory (Figure S2B), predominantly involved in cytoplasmic

translation GO functions (Figure S2C); Cluster 2 exhibited an initial increase

followed by a decline (Figure S2B), governing positive regulation of translation and

regulation of G2M transition of mitotic cell cycle (Figure S2C); transcription factors

within cluster 3 showed a sustained decrease in expression (Figure S2B) and were

enriched in cell cycle and meiotic cell cycle-related GO terms (Figure S2C). As

shown in Figure 3E, PIWIL4, C19orf84 and ST3GAL4 were specifically expressed in

SSC0 (belong to cluster 2). ID4, TCF3 and UTF1 were specifically expressed in

SSC1 (belong to cluster 2). DMRT1, ASB9 and NANOS3 were the main specifically

expressed genes in SSC2 (belong to cluster 3). DYNLL1, APRT and MKI67 were

specifically expressed in Diffing-SPG (belong to cluster 1/2/3). CTCFL, TEX19 and

SMC1B were specifically expressed in Diffed-SPG (belong to cluster 1). Heatmaps of

transcription factors showed that the transcription of SSC1 and SSC2 was very active

during reproductive development (Figure S2D). In contrast, SPT2/3 was almost

transcribed in silence. Diffing/ed served as a connecting link between the preceding

and the following, but its transcriptional state was closer to SSC (Figure S2D). In

conclusion, we obtained 50 transcription factors that are closely related to sperm

differentiation.

360　Sertoli cells from human testicular tissues are composed of heterogeneity

361　population

To explore the heterogeneity of Sertoli cells, we re-clustered the Sertoli cells and

identified 8 subclusters (Figure 4A). The top differentially expressed genes were

utilized to determine the Sertoli cell clusters (Figure 4B) using heatmap and violin

plots (Figure 4C). Of note, no cluster-specific distinctive features were identified in

SC1, as the marker genes (TF, SLC16A1 and CCNL1) of SC1 were also highly

expressed in other Sertoli cells subclusters (Figure 4C). SOX2, NF1B and COL27A1

were preferentially expressed in SC2. SPATS1, LINC01120 and LINC01206 were

specifically expressed in SC3. BEND2, PRAP1 and TEX1 were specifically

expressed in SC4. PLPP2, BEND4 and RHOXF1 were specifically expressed in SC5.

MLC1, SMCP and NUPR2 were highly expressed in SC6. FLRT2, CFH and BST2

were specifically expressed in SC7. TMIGD3, PGAM2 and CCDC62 were

specifically expressed in SC8.

Cell differentiation is accompanied by the expression of genes controlled by

cis-regulatory elements, which must be in an accessible state to function properly.

Therefore, scATAC-seq analysis was performed on the same Sertoli cells as those

used in scRNA-seq analysis, and a chromatin accessibility landscape for individual

cell marker was delineated (Figure 4D). Because the number of cells in the SC4-SC8

subpopulation was too small to calculate a peak plot, only the peak plots of markers in

the SC1-SC3 subpopulations are displayed. The promoter regions of TF were all

highly accessible in the SC1-SC3 subpopulations, and relatively higher openness was

observed for SOX2 in the SC2 subgroup. In addition to the high openness we

observed in the promoter region of the SPATS1, there were also multiple accessible

regions within the gene in the SC1-SC3 subpopulations.

Developmental trajectories of Sertoli cells and identified three novel

subpopulations

To analyze the origin and maturation process of Sertoli cells, pseudotime

trajectory analysis was carried out. The trajectory was determined to initiate with

State 3 as beginning and reached a terminally differentiate state of State 2 and State 1

(Figure 5A). According to Figure 5B, State 1 mainly included SC3/4/5 clusters, State

2 mainly included SC2/6/8 clusters, State 3 mainly included SC1/7 clusters. Figure 5C

demonstrated RNA velocity heatmap of dynamic evolutionary Sertoli cells on the

three States. Although most studies have categorized the development of Sertoli cells

after birth into two stages: immature and mature, the existence of an intermediate or

transitional state remains largely unknown. However, in this study, we identified three

distinct states of Sertoli cells, with State 1 expressing markers EGR3, CTSL, PCNA,

MK167, and KRT18 associated with immature Sertoli cells (Zhao et al., 2020), while

State 3 expressed markers, such as TF, HOPX, and DEF119, indicative of mature

Sertoli cells (Zhao et al., 2020). Consequently, we defined State 1 and State 3 as

immature and mature Sertoli cells, respectively. State 2, falling between immature and

mature states, does not belong to either category. Therefore, it is defined as a further

maturation Sertoli cells, aligning with previously reported findings in the literature

(Guo et al., 2021; Zhao et al., 2020). According to the latent time of RNA rate, the

differentiation process is SC3, SC4, SC5, SC2, SC6, SC8, SC1 and SC7 (Figure 5D).

Based on the above result, we identified three novel Sertoli cell subtypes (namely

SC4/7/8) and their specific markers, and defined SC4 (PRAP1) as immature, SC7

(BST2) as mature, and SC8 (CCDC62) as further mature (Figure 5D). To validate

these novel identified markers in Sertoli cells, we co-stained them with SOX9, a

classical marker of Sertoli cells. As shown in Figure 5E, we discovered that PRAP1,

BST2, and CCDC62 were co-expressed with SOX9 in testes tissues.

GO analysis of Sertoli cells in different states was shown in Figure S2E. Sertoli

cells in state 1 were mainly involved in cell cycle, meiotic cell cycle and

spermatogenesis. Sertoli cells in state 2 were mainly involved in platelet

degranulation, lipid metabolic process and steroid biosynthetic process. Sertoli cells in

state 3 were mainly involved in extracellular matrix organization, SRP-dependent

cotranslational protein targeting to membrane and viral transcription. Further Qusage

analysis was performed for Sertoli cells (Figure S2F). Sertoli cells in state 1 was

mainly involved in Cell cycle, DNA replication, and Mismatch repair. Sertoli cells in

state 2 was involved in D−Glutamine and D−glutamate metabolism, Ribosome, and

Valine, leucine and isoleucine biosynthesis. Sertoli cells in state 3 participated in

ECM−receptor interaction, and Protein digestion and absorption. Overall, Sertoli cells

performed different functions at different stages.

The existence of Sertoli cell subtypes is more crucial for spermatogenesis than its quantity

The highly interdependent structural relationship between Sertoli cells and germ cells has long been considered evidence of their close functional association (Griswold, 1995). For instance, some germ cells, due to their structural affinity with Sertoli cells, cannot be fully separated, when cultured independently in vitro, these germ cells exhibit very short survival times, however, the addition of Sertoli cells or the use of Sertoli cells-conditioned medium significantly improves the survival of germ cells (La et al., 2018; Mohammadi-Sardoo et al., 2021; Risley & Morse-Gaudio, 1992). To determine which subtype of Sertoli cells is more closely involved in spermatogenesis, we analyzed spermatogenesis and Sertoli cells distribution in NOA (experimental group) and OA (control group) groups (Figure 6). The number proportion of germ cell subsets in NOA1/2 and OA1/2 was shown in Figure 6A. The number of germ cells in NOA3 patients was 0. The proportion of Sertoli cell subsets in NOA1/2/3 and OA1/2 was shown in Figure 6B. It can be seen that the number of Sertoli cells in NOA2 samples is higher than that in NOA3 samples, but the number of Sertoli cells in three NOA samples were lower than that in OA samples, indicating that the number of Sertoli cells is somewhat correlated with spermatogenesis.

Moreover, interestingly, we observed a complete absence of immature Sertoli cells, especially SC3, in the testicular tissue of patients with NOA3 who exhibited a total absence of sperm, with only a small population of mature SC7 cells present (Figure 6C and 6D), suggesting that the absence of sperm in NOA3 patients may be associated with Sertoli cells SC3. For NOA2 samples, although the number of Sertoli cells was less than that of NOA3, SC3 was not missing in NOA2, so spermatogenesis was only partially affected (Figure 6C – 6F). In conclusion, these data suggested that

whether or not the critical subtype SC3 is missing is more important for

spermatogenesis than the number of Sertoli cells.

To determine the interaction between Sertoli cells and spermatogenesis, we applied

Cell-PhoneDB to infer cellular interactions according to ligand-receptor signalling

database. As shown in Figure S3A, compared with other cell types, germ cells were

mainly interacted with Sertoli cells. Cell Phone analysis showed that Notch1/2/3

signaling were involved in the interaction between Sertoli cells and germ cells (Figure

S3B). Interestingly, a chromatin accessibility landscape in scATAC-seq data showed

that promoter regions of Notch1 only accessible in the SC1-SC3 subpopulations not in

germ cells (Figure S3C), suggesting that Notch1 may be expressed by germ cells. The

openness of promoter region of Notch2 was observed not only in germ cells but also

in SSC2 and Pachytene celltypes. Notch3 accessibility was only observed within the

gene in different cell subtypes.

We also performed Spearman correlation analysis to determine the relationship

between Sertoli cells and germ cells. As shown in Figure 6D, State 1 SC3/4/5 were

tended to associated with PreLep, SSC0/1/2, and Diffing and Diffed-SPG sperm cells

(R > 0.72). Interestingly, SC3 was significantly positively correlated with all sperm

subpopulations (R > 0.5), again suggesting an important role for SC3 in

spermatogenesis and that SC3 is involved in the entire process of spermatogenesis.

In order to understand whether the functions of germ cells and Sertoli cells

correspond to each other, GO term enrichment analysis of germ cells and sertoli cells

was carried out (Figure S4, S5). We found that the functions could be divided into 8

categories, namely, material energy metabolism, cell cycle activity, the final stage of

sperm cell formation, chemical reaction, signal communication, cell adhesion and

migration, stem cells and sex differentiation activity, and stress reaction. These

different events were labeled with different colors in order to quickly capture the

important events occurring in the cells at each stage. As shown in Figure S4, we

discovered that SSC0/1/2 was involved in SRP-dependent cotranslational protein

targeting to membrane, and cytoplasmic translation; Diffing SPG was involved in cell

division and cell cycle; Diffied SPG was involved in cell cycle and RNA splicing;

Pre-Leptotene was involved in cell cycle and meiotic cell cycle; Leptotene_Zygotene

was involved in cell cycle and meiotic cell cycle; Pachytene was involved in cilium

assembly and spermatogenesis; Diplotene was involved in spermatogenesis and

cilium assembly; SPT1 was involved in cilium assembly and flagellated sperm

motility; SPT2 was involved in spermatid development and flagellated sperm motility;

SPT3 was involved in spermatid development and spermatogenesis.

As shown in Figure S5, SC1 were mainly involved in cell differentiation, cell

adhesion and cell communication; SC2 were involved in cell migration, and cell

adhesion; SC3 were involved in spermatogenesis, and meiotic cell cycle; SC4 were

involved in meiotic cell cycle, and positive regulation of stem cell proliferation; SC5

were involved in cell cycle, and cell division; SC6 were involved in obsolete

oxidation−reduction process, and glutathione derivative biosynthetic process; SC7

were involved in viral transcription and translational initiation; SC8 were involved in

spermatogenesis and sperm capacitation.

In summary, we found that while there is not a perfect one-to-one correspondence

between the differentiation stages of each Sertoli cell type and the sperm cell type,

there is a predominant trend-dependent supportive relationship between them (Figure

7). When we compared the GO terms of 8 Sertoli cell types with 12 sperm cell types,

we observed a close functional correlation between the two, particularly highlighting

the involvement of SC8 in supporting and participating in the formation of

spermatogenic cells at the SPT3 stage. SC5/6/7 primarily tend to be involved in SSC1/2, additionally, SC5/6 also exert their influence on Diffing and Diffed stages, while SC4 supports and participates in activities during the Diffed and Pre-Lep cell stages. SC1/2/8 are predominantly involved in SSC2 cell activities, with SC8 also showing a tendency to participate in the formation of L-Z and SPT3 cell stages. Notably, SC3 is almost universally involved in the entire process of spermatogenesis, playing a pivotal role throughout the process.

Co-localization of subpopulations of Sertoli cells and Germ cells

To further verify that Sertoli cell subtypes have "stage specificity" for each stage of sperm development, we performed immunofluorescence analysis using testicular tissues from OA and NOA2 samples. As shown in Figure 8, TF (SC1) and SOX2 (SC2) were co-expressed with ASB9 (SSC2) in testes tissues, indicating that SC1 and SC2 co-localized with SSC2 cells.

SC3 cells (marked by SPATS1) participated in the formation of multiple different stages of sperm cells, manifexted by co-localization with subpopulations of sperm cells in different periods, including SSC1 cells (marked by A2M), SSC2 cells (marked by ASB9), Pa cells (marked by C9orf57), SPT2 cells (marked by TSC21), SPT3 cells (marked by TSSK6).

SC4 cells (marked by PRAP1) co-localized with SSC2 cells (marked by ASB9) and Diffed cells (marked by TEX19).

BEND4 (SC5) was co-expressed with ST3GAL4 (SSC0), A2M (SSC1), ASB9 (SSC2), and TEX19 (Diffed) in testes tissues, indicating that SC5 co-localized with SSC0, SSC1, SSC2, Diffed cells.

SC6 (marked by SMCP) was co-localized with SSC1 cells (marked by A2M), SSC2 cells (marked by ASB9), and Diffed cells (marked by TEX19).

SC7 (BST2) was co-expressed with SSC1 (A2M) and SSC2 (ASB9) in testes tissues. SC8 (CCDC62) was co-expressed with SSC2 (ASB9), and SPT3 (TSSK6) in testes tissues.

Taken together, germ cells of a particular stage tended to co-localize with Sertoli cells of the corresponding stages. Germ cells and sertoli cells at each differentiation stage were functionally heterogeneous and stage-specific (Figure 7). This suggests that each stage of sperm development requires the assistance of sertoli cells to complete the corresponding stage of sperm development.

Leydig cell heterogeneity and development

Leydig cells are the main source of androgens and exist in the interstitial space between the seminiferous tubules (Shima et al., 2013). Studies have shown that Leydig cell differentiation undergoes immature - mature stage (Zhou et al., 2019). We also identified immature and mature Leydig cells. Re-clustering of Leydig cells generated twelve distinct cell clusters, including ten iLeydig cells and two Leydig cells (Figure S6A). Heatmap showed specific expression of markers in Leydig cells (Figure S6B). RSPO3 and MYOCD were highly expressed in iLeydig cells. CYP17A1 and CCDC69 were highly expressed in Leydig cells. Leydig cells in different samples are shown in Figure S6C. OA samples had more iLeydig cells than NOA samples (Figure S6D).

Peritubular myoid cells (PMCs) heterogeneity and development

PMCs are the key cellular components of the wall of seminiferous tubules and they are thought to be of great importance for the intratesticular transport of immotile sperm (Wang, Chen, & Liu, 2018). Re-clustering of PMCs revealed that there were 10 PMCs subclusters (Figure S7A). Heatmap showed specific expression of markers in PMCs (Figure S7B). PMCs cells in different samples are shown in Figure S7C. NOA

samples had more PMCs-1 than OA samples (Figure S7D). PMCs-1 might play an important role in spermatogenesis.

Macrophage heterogeneity and development

Macrophages play an important role in spermatogenesis and can produce 25-hydroxycholesterol within the testosterone biosynthetic pathway, thus contributing to the testosterone production process (Potter & DeFalco, 2017). Re-clustering of macrophage revealed that there were five macrophage subclusters (Figure S8A). Heatmap showed specific expression of markers in macrophage (Figure S8B). Macrophage cells in different samples are shown in Figure S8C. NOA samples had more macrophage-2 than OA samples (Figure S8D). Macrophage-2 might play a key part in spermatogenesis.

Discussion

NOA is due to pretesticular factors or testicular factors, usually abnormal spermatogenesis, which cannot be cured by surgery (Zarezadeh et al., 2021). Therefore, it is particularly important to study the etiology of NOA. Spermatogenesis is a complex developmental process that requires coordinated differentiation of multiple cell lines. In present study, we performed scRNA-seq and scATAC-seq to assess the heterogeneity of germ cells and somatic cells. We identified 11 subtypes for germ cells, 8 subtypes for Sertoli cells, 12 subtypes for Leydig cells, 10 subtypes for PMCs, 8 subtypes for macrophage and marker genes of specific cell type. The process of spermatogenesis was determined based on the cell trajectory analysis. Collectively, our results provide rich resources for exploring the potential mechanism of spermiogenesis.

In this study, some specific marker genes for germ cells were screened. We found

that UTF1 and ID4 were specifically expressed in SSC1, and DMRT1 was

specifically expressed in Diffing SPG. ID4 was a key regulator of SSC and ID4 also

marked SSCs in the mouse testis (Sun, Xu, Zhao, & Degui Chen, 2015). UTF1 was

reported to be a marker for SSCs in stallions (Jung, Roser, & Yoon, 2014). UTF1 was

discovered to play an important part in germ cell development, spermatogenesis, and

male fertility in mice (Raina, Dey, Thool, Sudhagar, & Thummer, 2021). DMRT1 is

necessary for male sexual development (Zhang, Oatley, Bardwell, & Zarkower, 2016).

The mechanism and expression pattern of these marker genes still need further

research.

SSC are adult stem cells in the testis of mammals that maintain spermatogenesis

and are essential for male fertility, but the mechanisms remain elusive. UTF1 is a

transcription factor expressed in SSC1, which plays a definite role in the proliferation

and differentiation of pluripotent stem cells (Tan & Wilkinson, 2019). Dmrt1 belongs

to a family of conserved transcriptional regulators that control several key processes

in mammalian testis, including germ cells and somatic cells. A recent discovery

suggested that DMRT1 was essential for the formation of SSC and had

gonadal-specific and amphoteric dimorphic expression patterns (Sohni et al., 2019).

NANOS3 is a member of the highly conservative NANOS family. The decrease in

NANOS3 expression could result in a decrease in the number of germ cells (Julaton &

Reijo Pera, 2011). In this study, UTF1 was specifically expressed in SSC1, and

DMRT1 and NANOS3 were the main specifically expressed genes in SSC2.

Therefore, UTF1, DMRT1 and NANOS3 might promote SSC proliferation and was

necessary for SSC differentiation.

Spermatogenesis consists of three processes: mitosis of spermatogonia, meiosis

of spermatocyte and deformation of spermatocyte. Abnormalities in any of these

stages can result in spermatogenesis disorders. During this process, the testicular

microenvironment in which spermatogenic cells are located plays an important role.

In addition to Sertoli cells, the testicular microenvironment also includes myoid cells

and mesenchymal cells and their secreted factors (Zhou et al., 2019). In this study, we

clustered four somatic cell types: Sertoli cells, Leydig cells, PMCs and macrophages.

Sertoli cells play an important role in normal spermatogenesis (Zomer & Reddi, 2020).

They provide the physical framework that supports the survival of germ cells and

secrete unique growth factors and cytokines to assist in the development of germ cells

(O'Donnell, Smith, & Rebourcet, 2022; Wu, Yan, Ge, & Cheng, 2020). In our study,

the NOA3 group had the least number of sperm cells due to the lack of SC3/8, and the

NOA2 group had a smaller number of SC3 cells than the normal OA group, so the

number of sperm cells in the NOA2 group was between the NOA3 and OA groups.

Although NOA2 did not have SC8 to participate in the formation of SPT3, this

process was compensated by SC3, which was involved in cell formation throughout

all stages of sperm development, and its function was particularly important. These

data suggested that the association between germ cells and Sertoli cells was

stage-specific, but this "specificity" is not entirely isolated.

Polycomb Group (PcG) is an important epigenetic modification gene, and PRC1

is one of the important complexes of PcG (Seif et al., 2020). PRC1 directs activation

of germline genes in good time during spermatogenesis. Inhibition of PRC1 in male

germ cells leads to the loss of a stem cell population little by little and serious

differentiation defects, thus causing male sterility (Maezawa et al., 2017). SCML1

was a member of PcG and is an essential component of the meiotic dense body

(Papanikos et al., 2017). Wang et al. (2018) reported that SCML1 was specifically

expressed in leptotene spermatocytes. These results indicated that SCML1 played an

important role in spermatogenesis.

Notch receptors have been reported to play an important role in cell fate

determination, maintenance and differentiation of stem cells (Huang, Rivas, &

Agoulnik, 2013). Notch signaling components have been discovered to express in

Sertoli and germ cells in the developing and mature testis (Garcia, DeFalco, Capel, &

Hofmann, 2013). In mice, a breakdown of Notch signaling can lead to abnormal cell

differentiation and early embryonic lethality (McCright et al., 2001; Swiatek, Lindsell,

del Amo, Weinmaster, & Gridley, 1994). Murta et al. (2014) also found notch

signaling disrupt results in abnormal spermatogenesis in the mouse. Notch signaling

was reported to be a key pathway regulating Sertoli cell physiology, and its changes

may disturb reaction of Sertoli cells to androgens (Kaminska et al., 2020). Notch

signaling in Sertoli cells is indirectly associated with germ cell development (Lu et al.,

2019). The upregulation of notch signaling in Sertoli cells induced the transformation

the quiescence to differentiation and meiosis in germ cells (Garcia & Hofmann, 2013).

In this study, we found Notch signaling were involved in the interaction between germ

cells and Sertoli cells. Therefore, Notch signaling might function in spermatogenesis.

Integrins are a kind of cell adhesion molecules in the integrin superfamily,

mediating cell to cell, cell to matrix or cell-matrix to cell adhesion and binding, and

participating in a series of important physiological and pathological processes such as

cell growth, differentiation and transmembrane signal transmission (Kuwar, Wen,

Zhang, & Sun, 2020). Previous studies have proved that there are specific integrin

subtypes and ligands in mammalian eggs and sperm, suggesting that this system may

be involved in the interaction between sperm and egg (Bowen & Hunt, 2000; Merc,

Frolikova, & Komrskova, 2021). In this study, we found integrins were involved in

the interaction between germ cells and Sertoli cells. Therefore, integrins might play a

key role in spermatogenesis.

In conclusion, our results revealed 11 germ cell subtypes and 8 Sertoli cell subtypes. We determined the process of spermatogenesis and found that SC3 subtypes (marked by SPATS1) of Sertoli cell played an important role in this process. The interaction between germ cells and sertoli cells at each differentiation stage were stage-specific. Notch1/2/3 signaling and integrin were discovered to be involved in germ cell-Sertoli cell interaction. Our results not only give us a comprehensive insight into human spermatogenesis, but also pave the way for determining molecules participated in the development of male germ cells, offering a powerful tool for further study on NOA.

Acknowledgements

Not applicable.

Author contributions

SW: Conceptualization, Data curation, Writing—original draft; HW: Conceptualization, Data curation, Writing—original draft; BJ: Visualization, Formal analysis; QZ: Visualization, Formal analysis; HY: Methodology, Project administration; DZ: Supervision, Project administration, Writing—review & editing. All authors read and approved the final manuscript.

Funding

The research was supported by ShenZhen Science and Technology Program (Grant No. JCYJ20230807111304010) and Futian Healthcare Research Project (Grant No. FTWS2023021).

Data Availability Statement

The data used to support the findings of this study are available from the

673 corresponding author upon request.

Ethics Approval and Consent to Participate

This study was approved by the Ethics Committee of Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine. All patients provided written informed consent.

Conflict of Interest

The authors have no relevant financial or non-financial interests to disclose.

References

Agarwal, A., Mulgund, A., Hamada, A., & Chyatte, M. R. (2015). **A unique view on male infertility around the globe**. *Reprod Biol Endocrinol, 13*, 37. https://doi.org/10.1186/s12958-015-0032-1

Bowen, J. A., & Hunt, J. S. (2000). **The role of integrins in reproduction**. *Proc Soc Exp Biol Med, 223*(4), 331-343. https://doi.org/10.1046/j.1525-1373.2000.22348.x

Fakhro, K. A., Elbardisi, H., Arafa, M., Robay, A., Rodriguez-Flores, J. L., Al-Shakaki, A., et al. (2018). **Point-of-care whole-exome sequencing of idiopathic male infertility**. *Genet Med, 20*(11), 1365-1373. https://doi.org/10.1038/gim.2018.10

Fröjdman, K., Harley, V. R., Pelliniemi, L. J. J. H., & biology, c. (2000). **Sox9 protein in rat sertoli cells is age and stage dependent**. *113*, 31-36.

Garcia, T. X., DeFalco, T., Capel, B., & Hofmann, M. C. (2013). **Constitutive activation of NOTCH1 signaling in Sertoli cells causes gonocyte exit from quiescence**. *Dev Biol, 377*(1), 188-201. https://doi.org/10.1016/j.ydbio.2013.01.031

Garcia, T. X., & Hofmann, M. C. (2013). **NOTCH signaling in Sertoli cells**

regulates gonocyte fate. *Cell Cycle, 12*(16), 2538-2545.

https://doi.org/10.4161/cc.25627

Gassei, K., & Orwig, K. E. (2016). **Experimental methods to preserve male**

fertility and treat male factor infertility. *Fertil Steril, 105*(2), 256-266.

https://doi.org/10.1016/j.fertnstert.2015.12.020

Griswold, M. D. (1995). **Interactions between germ cells and Sertoli cells in the**

testis. *Biol Reprod, 52*(2), 211-216. https://doi.org/10.1095/biolreprod52.2.211

Guo, J., Grow, E. J., Mlcochova, H., Maher, G. J., Lindskog, C., Nie, X., et al. (2018).

The adult human testis transcriptional cell atlas. *Cell Res, 28*(12), 1141-1157.

https://doi.org/10.1038/s41422-018-0099-2

Guo, J., Sosa, E., Chitiashvili, T., Nie, X., Rojas, E. J., Oliver, E., et al. (2021).

Single-cell analysis of the developing human testis reveals somatic niche cell

specification and fetal germline stem cell establishment. *Cell Stem Cell, 28*(4),

764-778 e764. https://doi.org/10.1016/j.stem.2020.12.004

Hess, R. A., & Moore, B. J. J. M. i. t. (1993). **Histological methods for evaluation of**

the testis. *3*(Part A), 86-94.

Huang, Z., Rivas, B., & Agoulnik, A. I. (2013). **NOTCH1 gain of function in germ**

cells causes failure of spermatogenesis in male mice. *PLoS One, 8*(7), e71213.

https://doi.org/10.1371/journal.pone.0071213

Jegou, B. (1993). **The Sertoli-germ cell communication network in mammals**. *Int*

Rev Cytol, 147, 25-96.

Julaton, V. T., & Reijo Pera, R. A. (2011). **NANOS3 function in human germ cell**

development. *Hum Mol Genet, 20*(11), 2238-2250. https://doi.org/10.1093/hmg/ddr114

Jung, H., Roser, J. F., & Yoon, M. (2014). **UTF1, a putative marker for spermatogonial stem cells in stallions**. *PLoS One, 9*(10), e108825. https://doi.org/10.1371/journal.pone.0108825

Kaminska, A., Pardyak, L., Marek, S., Wrobel, K., Kotula-Balak, M., Bilinska, B., et al. (2020). **Notch signaling regulates nuclear androgen receptor AR and membrane androgen receptor ZIP9 in mouse Sertoli cells**. *Andrology, 8*(2), 457-472. https://doi.org/10.1111/andr.12691

Kuwar, R., Wen, X., Zhang, N., & Sun, D. (2020). **An integrin-binding array platform identifies alphavbeta8 and alpha5beta1 integrins on rat primary cortical neurons to support their survival and growth**. *J Neurosci Methods, 339*, 108729. https://doi.org/10.1016/j.jneumeth.2020.108729

La, H. M., Mäkelä, J. A., Chan, A. L., Rossello, F. J., Nefzger, C. M., Legrand, J. M. D., et al. (2018). **Identification of dynamic undifferentiated cell states within the male germline**. *Nat Commun, 9*(1), 2819. https://doi.org/10.1038/s41467-018-04827-z

Lasala, C., Schteingart, H. F., Arouche, N., Bedecarrás, P., Grinspon, R. P., Picard, J.-Y., et al. (2011). **SOX9 and SF1 are involved in cyclic AMP-mediated upregulation of anti-Müllerian gene expression in the testicular prepubertal Sertoli cell line SMAT1**. *301*(3), E539-E547.

Liao, J., Ng, S. H., Luk, A. C., Suen, H. C., Qian, Y., Lee, A. W. T., et al. (2019).

Revealing cellular and molecular transitions in neonatal germ cell differentiation using single cell RNA sequencing. *Development, 146*(6). https://doi.org/10.1242/dev.174953

Lottrup, G., Nielsen, J., Maroun, L., Møller, L., Yassin, M., Leffers, H., et al. (2014). **Expression patterns of DLK1 and INSL3 identify stages of Leydig cell differentiation during normal development and in testicular pathologies, including testicular cancer and Klinefelter syndrome**. *29*(8), 1637-1650.

Lu, E., Feng, F., Wen, W., Tong, X., Li, X., Xiao, L., et al. (2019). **Notch signaling inhibition induces G0/G1 arrest in murine Leydig cells**. *Andrologia, 51*(10), e13413. https://doi.org/10.1111/and.13413

Maezawa, S., Hasegawa, K., Yukawa, M., Sakashita, A., Alavattam, K. G., Andreassen, P. R., et al. (2017). **Polycomb directs timely activation of germline genes in spermatogenesis**. *Genes Dev, 31*(16), 1693-1703. https://doi.org/10.1101/gad.302000.117

McCright, B., Gao, X., Shen, L., Lozier, J., Lan, Y., Maguire, M., et al. (2001). **Defects in development of the kidney, heart and eye vasculature in mice homozygous for a hypomorphic Notch2 mutation**. *Development, 128*(4), 491-502.

Merc, V., Frolikova, M., & Komrskova, K. (2021). **Role of Integrins in Sperm Activation and Fertilization**. *Int J Mol Sci, 22*(21). https://doi.org/10.3390/ijms222111809

Mimitou, E. P., Lareau, C. A., Chen, K. Y., Zorzetto-Fernandes, A. L., Hao, Y., Takeshima, Y., et al. (2021). **Scalable, multimodal profiling of chromatin**

accessibility, gene expression and protein levels in single cells. *Nat Biotechnol,*

39(10), 1246-1258. https://doi.org/10.1038/s41587-021-00927-2

Mohammadi-Sardoo, M., Mandegary, A., Nematollahi-Mahani, S. N., Moballegh

Nasery, M., Nabiuni, M., & Amirheidari, B. (2021). **Cytotoxicity of mancozeb on**

Sertoli-germ cell co-culture system: Role of MAPK signaling pathway. *Toxicol*

Ind Health, 37(11), 674-684. https://doi.org/10.1177/07482337211044028

Murta, D., Batista, M., Trindade, A., Silva, E., Henrique, D., Duarte, A., et al. (2014).

In vivo notch signaling blockade induces abnormal spermatogenesis in the mouse.

PLoS One, 9(11), e113365. https://doi.org/10.1371/journal.pone.0113365

Naz, M., & Kamal, M. (2017). **Classification, causes, diagnosis and treatment of**

male infertility: a review. *Oriental Pharmacy & Experimental Medicine, 17*(2),

1-21.

O'Donnell, L., Smith, L. B., & Rebourcet, D. (2022). **Sertoli cells as key drivers of**

testis function. *Semin Cell Dev Biol, 121*, 2-9.

https://doi.org/10.1016/j.semcdb.2021.06.016

Papanikos, F., Daniel, K., Goercharn-Ramlal, A., Fei, J. F., Kurth, T., Wojtasz, L., et al.

(2017). **The enigmatic meiotic dense body and its newly discovered component,**

SCML1, are dispensable for fertility and gametogenesis in mice. *Chromosoma,*

126(3), 399-415. https://doi.org/10.1007/s00412-016-0598-1

Pervolarakis, N., Nguyen, Q. H., Williams, J., Gong, Y., Gutierrez, G., Sun, P., et al.

(2020). **Integrated Single-Cell Transcriptomics and Chromatin Accessibility**

Analysis Reveals Regulators of Mammary Epithelial Cell Identity. *Cell Rep, 33*(3),

108273. https://doi.org/10.1016/j.celrep.2020.108273

Potter, S. J., & DeFalco, T. (2017). **Role of the testis interstitial compartment in**

spermatogonial stem cell function. *Reproduction, 153*(4), R151-R162.

https://doi.org/10.1530/REP-16-0588

Raina, K., Dey, C., Thool, M., Sudhagar, S., & Thummer, R. P. (2021). **An Insight**

into the Role of UTF1 in Development, Stem Cells, and Cancer. *Stem Cell Rev*

Rep, 17(4), 1280-1293. https://doi.org/10.1007/s12015-021-10127-9

Risley, M. S., & Morse-Gaudio, M. (1992). **Comparative aspects of spermatogenic**

cell metabolism and Sertoli cell function in Xenopus laevis and mammals. *J Exp*

Zool, 261(2), 185-193. https://doi.org/10.1002/jez.1402610209

Seif, E., Kang, J. J., Sasseville, C., Senkovich, O., Kaltashov, A., Boulier, E. L., et al.

(2020). **Phase separation by the polyhomeotic sterile alpha motif**

compartmentalizes Polycomb Group proteins and enhances their activity. *Nat*

Commun, 11(1), 5609. https://doi.org/10.1038/s41467-020-19435-z

Sharma, M., Srivastava, A., Fairfield, H. E., Bergstrom, D., Flynn, W. F., & Braun, R.

E. J. B. (2018). **Identification of slow-cycling germline stem cells and their**

regulation by PLZF. 477604.

Shima, Y., Miyabayashi, K., Haraguchi, S., Arakawa, T., Otake, H., Baba, T., et al.

(2013). **Contribution of Leydig and Sertoli cells to testosterone production in**

mouse fetal testes. *Mol Endocrinol, 27*(1), 63-73.

https://doi.org/10.1210/me.2012-1256

Sohni, A., Tan, K., Song, H. W., Burow, D., de Rooij, D. G., Laurent, L., et al. (2019).

The Neonatal and Adult Human Testis Defined at the Single-Cell Level. *Cell Rep,*

26(6), 1501-1517 e1504. https://doi.org/10.1016/j.celrep.2019.01.045

Sun, F., Xu, Q., Zhao, D., & Degui Chen, C. (2015). **Id4 Marks Spermatogonial**

Stem Cells in the Mouse Testis. *Sci Rep, 5*, 17594. https://doi.org/10.1038/srep17594

Swiatek, P. J., Lindsell, C. E., del Amo, F. F., Weinmaster, G., & Gridley, T. (1994).

Notch1 is essential for postimplantation development in mice. *Genes Dev, 8*(6),

707-719. https://doi.org/10.1101/gad.8.6.707

Tan, K., Song, H. W., & Wilkinson, M. F. (2020). **Single-cell RNAseq analysis of**

testicular germ and somatic cell development during the perinatal period.

Development, 147(3). https://doi.org/10.1242/dev.183251

Tan, K., & Wilkinson, M. F. (2019). **Human Spermatogonial Stem Cells**

Scrutinized under the Single-Cell Magnifying Glass. *Cell Stem Cell, 24*(2),

201-203. https://doi.org/10.1016/j.stem.2019.01.010

Vertesy, A., Arindrarto, W., Roost, M. S., Reinius, B., Torrens-Juaneda, V., Bialecka,

M., et al. (2018). **Parental haplotype-specific single-cell transcriptomics reveal**

incomplete epigenetic reprogramming in human female germ cells. *Nat Commun,*

9(1), 1873. https://doi.org/10.1038/s41467-018-04215-7

Wang, M., Liu, X., Chang, G., Chen, Y., An, G., Yan, L., et al. (2018). **Single-Cell**

RNA Sequencing Analysis Reveals Sequential Cell Fate Transition during

Human Spermatogenesis. *Cell Stem Cell, 23*(4), 599-614 e594.

https://doi.org/10.1016/j.stem.2018.08.007

Wang, Y. Q., Chen, S. R., & Liu, Y. X. (2018). **Selective deletion of WLS in**

peritubular myoid cells does not affect spermatogenesis or fertility in mice. *Mol*

Reprod Dev, 85(7), 559-561. https://doi.org/10.1002/mrd.22988

Wu, S., Yan, M., Ge, R., & Cheng, C. Y. (2020). **Crosstalk between Sertoli and**

Germ Cells in Male Fertility. *Trends Mol Med, 26*(2), 215-231.

https://doi.org/10.1016/j.molmed.2019.09.006

Yao, C., Xu, W., Liu, N., Zhi, E., Shi, C., Ji, Z., et al. (2022). **Preliminary experience**

with 3D digital image microscope system on the treatment of varicocele.

Andrologia, 54(7), 1592-1597. https://doi.org/10.1111/and.14423

Zarezadeh, R., Nikanfar, S., Oghbaei, H., Rastgar Rezaei, Y., Jafari-Gharabaghlou, D.,

Ahmadi, Y., et al. (2021). **Omics in Seminal Plasma: An Effective Strategy for**

Predicting Sperm Retrieval Outcome in Non-obstructive Azoospermia. *Mol*

Diagn Ther, 25(3), 315-325. https://doi.org/10.1007/s40291-021-00524-8

Zhang, T., Oatley, J., Bardwell, V. J., & Zarkower, D. (2016). **DMRT1 Is Required**

for Mouse Spermatogonial Stem Cell Maintenance and Replenishment. *PLoS*

Genet, 12(9), e1006293. https://doi.org/10.1371/journal.pgen.1006293

Zhao, L., Yao, C., Xing, X., Jing, T., Li, P., Zhu, Z., et al. (2020). **Single-cell analysis**

of developing and azoospermia human testicles reveals central role of Sertoli

cells. *Nat Commun, 11*(1), 5683. https://doi.org/10.1038/s41467-020-19414-4

Zhou, R., Wu, J., Liu, B., Jiang, Y., Chen, W., Li, J., et al. (2019). **The roles and**

mechanisms of Leydig cells and myoid cells in regulating spermatogenesis. *Cell*

Mol Life Sci, 76(14), 2681-2695. https://doi.org/10.1007/s00018-019-03101-9

Zomer, H. D., & Reddi, P. P. (2020). **Characterization of rodent Sertoli cell**

primary cultures. *Mol Reprod Dev,* *87*(8), 857-870.

Figure Legends

Figure 1. Overview of major cell types and cellular attributes of testes of OA and

NOA patients. (A) Representative images of HE staining of three groups of samples.

(**B**) Schematic of the experimental design for scRNA-seq and scATAC-seq. (**C**)

UMAP analysis of human testicular cells in scRNA-seq results. (**D**) UMAP analysis

of human testicular cells in scATAC-seq results. (**E**) Heatmap of expression of

markers for the 12 cell types. (**F**) UMAP analysis of testicular cell population.

Figure 2. Identification of germ cell subtypes. (**A**) UMAP analysis of 12 germ cell

subsets in scRNA-seq results. (**B**) Pseudotime analysis of 12 germ cell subpopulations.

(**C**) Gene expression patterns of marker genes on UMAP plots. (**D**) State of germ cell

subsets using pseudotime analysis. (**E**) Immunofluorescence double staining of

markers of newly identified spermatocyte subpopulations with the classical marker

DDX4. (**F**) Accessibility peak plots in scATAC-seq of markers for newly identified

spermatocyte subpopulations.

Figure 3. Analysis of germ cells at different stages. (**A**) Germ cell subsets at

different stages of the cell cycle. (**B**) Heatmap of cell cycle in germ cells. (**C**)

Transcriptional dynamics of driver genes of germ cell subtypes. (**D**) The main stage of

spermatogenesis. (**E**) Beam analysis of differentially expressed genes at different

stages of SSC.

Figure 4. Identification of Sertoli cell subtypes and epigenetic analysis. (**A**)

UMAP analysis showed 8 Sertoli cell subsets in scRNA-seq results. (**B**) Heatmap of

markers of differentially expressed genes in eight Sertoli cell subsets. (**C**) Gene

expression patterns of Sertoli cell subtype marker genes on violin plots. (**D**) Accessibility peak plots in scATAC-seq of markers for newly identified spermatocyte subpopulations.

Figure 5. The development of Sertoli cells. (**A**) State of Sertoli cells using pseudotime analysis. (**B**) Distribution of Sertoli cell subtypes in different states and SC state division. (**C**) Potential temporal differentiation of SC subgroups using RNA velocity. (**D**) Sertoli cells division diagram and representative markers. (**E**) Immunofluorescence of Sertoli cell subtype markers in testicular tissues from OA patients. The scale bar represents 50 μm.

Figure 6. Correlation analysis of germ and Sertoli cell expression in different groups. (**A**) Cell proportion of germ cell subtypes in different groups. (**B**) Cell proportion of Sertoli cell subtypes in different groups. (**B**) Pseudotime analysis of germ cell subtypes in different groups. (**D**) Pseudotime analysis of Sertoli cell subtypes in different groups. (**E**) Distribution map of germ cell subtypes in different samples. (**F**) Distribution map of Sertoli cell subtypes in different samples. (G) Spearman correlation between germ and Sertoli cells. (**D**) The interaction between SSC0/1/2 and Sertoli cells.

Figure 7. **Schematic diagram of spermatogenesis and stage specificity of sertoli cell subtypes.**

Figure 8. Immunofluorescence analysis of germ cell and Sertoli cell subtype markers in testicular tissues. Blue DAPI indicates nuclei. The colors of the markers and cells indicate the fluorescent color of the corresponding antibody. The scale bar represents 50 μm.

Supplementary Information

Figure S1. Analysis of germ cell markers in spermatogenesis. (**A**) Heatmap

showed marker gene expression of germ cell subtypes. (**B**) Gene expression patterns

of germ cell markers on violin plots. (**C**) Gene expression patterns of germ cell

markers on violin plots. Distribution diagram of germ cell subsets using pseudotime

analysis.

Figure S2. **Germ cell function analysis and epigenetic analysis.** (**A-C**) Analysis of

beam cluster transcription factors and its GO term. (**D**) Heatmap of transcription

factors in germ cells. (**E**) GO analysis of different states of Sertoli cells. (**F**) Qusage

analysis of different states of Sertoli cells.

Figure S3. Correlation analysis of germ and Sertoli cell expression. (**A**) The

interaction between germ cells and other cells. (**B**) The interaction between germ cells

and Sertoli cells through Notch signaling pathway. (**C**) Accessibility peak plots in

scATAC-seq of Notch1/2/3.

Figure S4. GO analysis of germ cells.

Figure S5. GO analysis of Sertoli cells.

Figure S6. Identification of Leydig cell types in the testis. (**A**) Clustering on the

Leydig cells revealed 12 different cell types. (**B**) Heatmap showed Leydig cell

subtype specific maker genes. (**C**) The distribution of Leydig cells in different

samples. (**D**) Proportion of Leydig cell subtypes in each sample.

Figure S7. Identification of PMCs types in the testis. (**A**) Clustering on the PMCs

revealed 8 different cell types. (**B**) Heatmap showed PMCs subtype specific maker

genes. (**C**) The distribution of PMCs in different samples. (**D**) Proportion of PMCs

subtypes in each sample.

Figure S8. Identification of macrophage types in the testis. (**A**) Clustering on the

macrophage revealed 6 different cell types. (**B**) Heatmap showed macrophage subtype

specific maker genes. (**C**) The distribution of macrophage in different samples. (**D**)

Proportion of macrophage subtypes in each sample.