Audio-visual concert performances synchronize an audience's heart rates.

Short title: Audio-visual performances synchronize heart rates.

Anna M. Czepiel*[1,2,3], Lauren K. Fink[2,4,5], Mathias Scharinger[6,7], Christoph Seibert[8], Melanie Wald-

Fuhrmann[2,5], & Sonja A. Kotz[3,9]

[1]Department of Psychology, University of Toronto Mississauga, Canada

[2]Department of Music, Max Planck Institute for Empirical Aesthetics, Germany

[3]Department of Neuropsychology and Psychopharmacology, Faculty of Psychology and Neuroscience,

Maastricht University, The Netherlands

[4] Department of Psychology, Neuroscience & Behaviour, McMaster Institute for Music & the Mind, McMaster

University

[5] Max Planck-NYU Center for Language, Music, and Emotion, Frankfurt am Main, Germany & New York

[6] Research Group Phonetics, Department of German Linguistics, University of Marburg, Germany

[7] Department of Language and Literature, Max Planck Institute for Empirical Aesthetics, Germany

[8] Institute for Music Informatics and Musicology, University of Music Karlsruhe, Germany

[9] Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, Germany

*Corresponding author

Email: a.czepiel@utoronto.ca

Anna M. Czepiel

Department of Psychology

University of Toronto Mississauga

3359 Mississauga Road, Mississauga

Canada

https://orcid.org/0000-0002-7101-945X

Abstract

People enjoy engaging with music. Live music concerts provide an excellent option to investigate

real-world music experiences, and at the same time, use neurophysiological synchrony to assess

dynamic engagement. In the current study, we assessed engagement in a live concert setting using

synchrony of cardiorespiratory measures, comparing inter-subject, stimulus-response, correlation,

and phase coherence. As engagement might be enhanced in a concert setting by *seeing* musicians

perform, we presented audiences with audio-only (AO) and audio-visual (AV) piano performances.

Only correlation measures were above chance level. In comparing time-averaged synchrony across

conditions, AV performances evoked higher inter-subject correlation of heart rate (ISC-HR).

However, self-reported engagement did not correspond to synchrony when averaged across music

pieces. On the other hand, time-resolved analyses show that synchronized deceleration-acceleration

heart rate (HR) patterns, typical of an 'orienting response' (an index of directed attention), occurred

within music pieces at salient events of section boundaries. That is, seeing musicians perform

heightened audience engagement at structurally important moments in Western classical music.

Overall, we could show that multisensory information shapes dynamic engagement. By comparing

different synchrony measures, we further highlight the advantages of timeseries analysis, specifically

ISC-HR, as a robust measure of holistic musical listening experiences in naturalistic concert settings.

Keywords: cardiorespiratory synchrony; engagement; inter-subject correlation; stimulus-response

correlation; music concerts.

1. Introduction

People enjoy engaging in music, especially in live concerts[1–3]. While concerts provide an excellent option to explore music listening experiences, assessing this experience dynamically is methodologically challenging. One way to explore musical engagement over longer time scales is with synchrony of peripheral responses either across participants or to perceived musical structures. However, many different analysis techniques can be employed. Here, we compared different synchrony analyses to gauge their robustness. Using cardiorespiratory synchrony, the current study assessed engagement, defined as a listener's real-time absorption in music[4–6]. As engagement might be enhanced in a concert setting by *seeing* musicians perform[7–10], we presented participants with audio-visual and audio-only musical performances in a concert context.

Musical engagement can describe either involvement in musical activities (e.g., instrumental practice) or the time-varying mental state when listening to music[11]. Here, we refer to engagement as an absorbed state of heightened attention and affect, which can change on a second-by-second basis while listening to music[5,6,11–13] (for a review, see[5]). Indeed, music is a particular driver of (regular) time-varying attentional focus due to its metrical structure/beat. Although musical engagement can be measured with continuous behavioural responses[12,13], these ratings might distract from directly experiencing music. A promising alternative to measure engagement in a non-distracting way is to assess neural and peripheral physiological responses. One potential index is the orienting response: the engagement of attention, which is reflected in increased skin conductance and deceleration-acceleration patterns of respiration/heart rate[14]. Another method to objectively assess engagement is through physiological synchrony (e.g.,[2,15–18]), which assumes that increased engagement with music is more likely to evoke common time-locked responses across participants[4,19]. Synchrony can be between a participant's response and a stimulus feature that represents how we process/respond to sound, like the auditory envelope (i.e., how sound intensity

changes over time). More recently, spectral flux (spectrum changes over time) has been used as a

more faithful representation of how humans respond to auditory stimuli[20–24] as it represents both

envelope and rhythmic information[24] and is related to perceived higher activity (e.g., faster music)

and fullness of sound[25]. Assessed by calculating the (Euclidean) distance between the spectrum of

successive frames, spectral flux can also be assessed within specific frequency bands (sub-bands),

which represent different rhythmic elements[26]. For example, in pop music, spectral flux of lower

frequency bands can represent kick drum and bass guitar rhythms, while higher frequency bands can

represent cymbals and hihat rhythms[27]. Synchronisation of participant responses to stimuli can be

assessed using correlation or phase coherence, i.e., rhythmic alignment[4,24]. In this paper, we refer to

these as stimulus-response correlation (SRC) and stimulus-response phase coherence (SRPC),

respectively. A related synchrony approach is to assess similarity across multiple time-locked

participant responses[28]. Similarly to stimulus-response synchrony, inter-subject synchrony can be

calculated with correlation[17,29] or using phase coherence; in this paper, we refer to these as inter-

subject correlation (ISC) and inter-subject phase coherence (ISPC), respectively.

Growing evidence has related synchrony of neural responses to a participant's engagement with

stimuli. In correlational measures, SRC and ISC were related to engagement with movies[17,30,31],

speech [32,33], and music[2,4,13,19]. In phase coherence measures, higher neural synchrony occurred in

engaging group discussions, indicating a potential marker for shared attention mechanisms[34].

Although phase synchrony in speech was attributed to stimulus intelligibility[35], it was postulated that

phase synchrony increases when engaging in stimuli. According to the Dynamic Attending Theory

(DAT), internal oscillations adapt to external rhythms so that attending energy is optimised at

expected time points[36,37]. Indeed, successful coordination in musical ensembles was related to phase

synchrony in neural[38] and heart[39] rhythms (see also[40,41]). Recent work shows coupling of cerebral

activity occurs when experiencing similar emotional response to live music concerts[15]. However,

most synchrony research focused exclusively on neural responses, such as those measured via

electroencephalography (EEG); more suitable approaches for a live concert audience could be to

explore synchrony in peripheral physiological responses[2,16,42].

Recent frameworks propose certain neural mechanisms – such as synchrony – might extend to

cardiac, respiratory, gut, and pupil rhythms[43–45]. Promising results show cardiorespiratory ISC

related to engagement with narratives and instructional videos[18,46]. In the phase domain, some

research indicates that the alignments of breathing/heartbeats to external rhythms aid the

processing of upcoming stimulus events[47,48]. In music, respiration has been shown to entrain to

musical beats[49,50], although heartbeats do not[51–53]. The current study aimed to further assess

cardiorespiratory synchrony as an index of engagement in response to music performances in a

concert audience.

Cardiorespiratory synchrony can be time-averaged across stimuli to assess engagement between

conditions (e.g., attended/unattended conditions[18]). However, music listening is dynamic, where

engagement, neural synchrony[12,13,54], and physiological synchrony[16] change over time. Previous

research reported that engagement tends to peak at musical climaxes or structural boundaries[55].

Indeed, it seems beneficial for attentional resources to 'latch onto' structural boundaries: the brain

naturally segments incoming information into more meaningful units to process them more

efficiently[56,57]. According to the Generative Theory of Tonal Music (GTTM[58]), we 'segment' auditory

events based on salient changes when several music features change simultaneously. The ability to

segment music at such salient moments has seen consistent empirical support in behavioural

ratings[59–62]. However, consciously rating engagement continuously distracts from a natural music

listening experience. Recent research showed that salient musical section boundaries evoke

neurophysiological synchrony, potentially reflecting heightened engagement[12,13,18]. Therefore, to

complement the current investigation of engagement during audio-only (AO) and audio-visual (AV)

music performance on a global level (time-averaged across pieces), we also tested time-varying

engagement – indexed by physiological synchrony – between AO and AV music performances during

salient moments, defined as section boundaries.

The current study

Audiences might find concerts engaging due to not only listening to music but also seeing a musician

perform[8,10]. To test this, concert audiences were presented with AO and AV performances of Western

classical piano pieces. Musical pieces were selected as they represent a range of genres (Baroque,

Classical-Romantic, 20th century music), and characters, as well as range of tempi, harmonic and

musical phrase structures. Presenting such a musical range was thought to evoke high engagement

levels. Additionally, pieces were chosen as they may be considered typical pieces heard in live

Western classical performances, though not known to the audience, thereby reducing familiarity

effects. Based on Kaneshiro et al.[4], dynamic engagement was assessed by calculating

cardiorespiratory synchrony using both correlation and phase coherence. For correlation, we

assessed heart and respiration rate, e.g., speed of breathing and heartbeats in terms of beats per

minute (BPM). For phase coherence, we assessed the phase alignment of heartbeat cycles (i.e.,

whether heartbeats align in time) and breathing cycles (i.e., whether breathing aligns in time).

Synchrony was then assessed both across (i.e., time-averaged synchrony) and within (i.e., time-

resolved synchrony) musical pieces. First, we tested that time-locked responses were synchronized

above chance level. Second, we hypothesised that time-averaged synchrony would be higher in AV

than in AO musical performances. Third, we expected time-resolved synchrony to be higher at section

boundaries (based on previous literature[12,13,16]). Further, we expect synchrony to be higher at section

boundaries in AV than AO at salient moments in the music (interaction effect), based on literature

showing that visual information enhances perceptual responses[63] and that visual imagery heightens

engagement[55].

2. Method

Participants, stimuli, and procedures are identical to Czepiel et al.[7] (see General Methods and

Experiment 2). Key details of the procedure are outlined below. Data and code are available on Open

Science Framework (OSF)[1].

Participants

The study was approved by the Ethics Council of the Max Planck Society and in accordance with the

Declaration of Helsinki. Participants gave their written informed consent. Twenty-six participants

attended the concerts. One participant was excluded due to missing physiological data, thus

behavioural and physiological data of twenty-five participants were analysed. Participants included

nine females, mean age of 51.08 years (SD = 15.48), who on average had 6.18 (SD = 8.20) years of

music lessons and attended on average 13 concerts per year (M = 19.97, SD = 20).

Stimuli

Participants were presented with AV and AO versions (see Figure 1a) of three Western classical piano

pieces: Johann Sebastian Bach: Prelude and Fugue in D major (Book Two from the Well-Tempered

Clavier, BWV 874), Ludwig van Beethoven: Sonata No. 7, Op. 10, No. 3, second movement (Largo e

mesto), and Olivier Messiaen: *Regard de l'Esprit de joie* (No. 10 from *Vingt Regards sur l'enfant Jésus*).

Both versions were performed by the same professional pianist, on the same piano, in the same

---

[1] Please note this repository is currently private and anonymous while this manuscript is under review; it will be made public upon acceptance of this manuscript.

concert hall. AO versions were recorded prior to the concert and were presented as recordings

through high-quality loudspeakers, while AV versions were performed live by the pianist during the

concert. A trained sound engineer checked that the sound level was equal across all stimuli;

additional analyses of acoustic features showed these performances were comparable across

concerts[7].

Procedure

Participants were invited to attend one of two concerts. Electrocardiography (ECG) and respiration

were collected using gelled self-adhesive Ag/AgCl electrodes and a respiration belt, respectively, and

continuously measured during the concert at a 1000 Hz sampling rate. During the concert, audiences

heard the three pieces both in AV and AO modalities. The order of modality was counterbalanced

across the two concerts. After each piece, participants rated items such as engagement. We used a

German word to reflect engagement as an absorbed state (see Introduction). Although it was

translated as 'absorption' in a previous study[7], it can also be used synonymously with engagement

(see also[64]).

Data Analysis

Musical analysis: spectral flux and section boundaries

To model how humans respond to music, previous studies have extracted the envelope from the

acoustic signal[4]. More recently, spectral flux has shown to be a better predictor of participants'

responses to an acoustic signal as it shares information of the acoustic envelope as well as

note/speech onset information[23,24,65]. Therefore, the continuous spectral flux signal was obtained

using the MIRToolbox[26] in Matlab (2019b), with a frame size of 25ms (as is appropriate for short-

term acoustic features[66]), with a 50% overlap (Figure 1b). To allow for comparison between

conditions, we confirmed that spectral flux was similar across AV and AO performances, with Pearson

correlations at $r > 0.92$ (see Supplementary Figure 1). To investigate the phase relationship between

spectral flux against the cardiac and respiration measures fairly, we assessed spectral flux within the

audience members' average heart (1.01 Hz) and respiration (0.30 Hz) frequencies. This approach

followed previous research that also selected spectral flux within specific frequencies (sub-bands) to

better ascertain rhythm hierarchy levels[25,27]. Although it is possible in the MIRToolbox to specify

these sub-bands, the default sub-bands were considered too wide for the current hypotheses of

heart/respiration frequencies. Therefore, to obtain rhythmic levels that coincide with heartbeat and

respiration rhythms, spectral flux was bandpass filtered at representative ranges of the interquartile

range of all heart (0.951 - 1.159 Hz) and respiration (0.268-0.358 Hz) frequencies (see

Supplementary Figure 2). The phase angle of filtered spectral flux was calculated from the real part

of the Hilbert envelope using MATLAB's *angle* function (Figure 1b, lower panel). The section

boundaries in the musical pieces were identified either by a double bar line or end repeat bar line in

the score, or by a change/repeat of thematic material (identified in the score by AMC, then confirmed

by a music theorist, see Supplementary Table 1).

Figure 1. Outline of the experimental design and analysis pipelines. **Panel A** shows the study design:

audiences were presented with music pieces in audio-visual (AV, orange box) and audio-only (AO,

blue box) conditions, while heart (ECG), respiration, and acoustic signal of the music (maroon) were

continuously recorded. **Panel B** shows how beats per minute (BPM) were extracted from the ECG

and respiration signals, which were then used for correlational measures (upper panel). **Panel B** also

shows that the angles of heartbeat and respiration cycles were extracted for phase coherence

measures, where start of cycle began at peaks of ECG and respiration signals (peaks marked as green

dots) (lower panel). **Panel C** shows the correlational measures (upper panel) and the phase-

coherence measures (lower panel) we extracted from the continuous measures. For the correlational

measures, we extracted heart and respiration rate (HR, RR, from cardiorespiratory signals) and

spectral flux (from acoustic signal). For the phase coherence measures, we extracted the phase angle

of ECG and respiration cycles (from cardiorespiratory signals), and angles of spectral flux band-pass

filtered at the audience's heart frequencies (0.951 - 1.159 Hz) and respiration frequencies (0.268-

0.358 Hz). **Panel D** shows the extracted synchrony measures: stimulus-subject (vertical, left) and

inter-subject (vertical, right) for correlation (horizontal, top) and phase coherence (horizontal,

bottom) measures.

Pre-processing of physiological data.

Raw heart and respiration signals were pre-processed in MatLab 2019b. Missing data from the raw

signals were first interpolated at the original sampling rate (all gaps in data were less than 60 ms).

Data were cut per piece and further pre-processed using Fieldtrip[67] and the *biosig* toolbox

(http://biosig.sourceforge.net). Respiration data were low pass filtered at 2 Hz, ECG data were band-

pass filtered between 0.6 and 20 Hz (Butterworth, 4th order), and both demeaned. Peaks in ECG and

respiration signals were extracted using, respectively, *nqrsdetect* function from biosignal and a

custom-made script that located when the respiration signal exceeded a peak threshold.

Computationally identified peaks were manually screened to ensure correct identification of R wave

(part of the Q-, R- and S-wave [QRS] complex of ECG) and respiration peak locations. Any peaks that

were not correctly identified were manually added, while falsely identified peaks were removed, for

example, if a T wave in the ECG was accidentally identified as a R wave. In rare cases where

participant data within a piece were too noisy to identify clear QRS and inspiration peaks, participant

data for this piece were rejected from further analysis (ECG = 14%, respiration = 7%). The pre-

processed heart (ECG) and respiration signals were then further analysed in two different ways to

calculate synchrony: heart *rate* and respiration *rate* (BPM) were calculated for correlational

synchrony, while *phase angle* of ECG (phase of heartbeat cycle) and respiration (phase of

inspiration/exhalation cycle) were calculated for phase synchrony (more details outlined below, see

Figure 1A-B).

Synchrony analyses

Four continuous synchrony measures were extracted: stimulus-response correlation (SRC), inter-

subject correlation (ISC), stimulus-responses phase coherence (SRPC), and inter-subject phase

coherence (ISPC) for each ECG and respiration (see Fig. 1D).

Correlational synchrony: ISC and SRC

Correlational synchrony measures (SRC, ISC) were calculated using heart rate (HR) and respiration

rate (RR). To do so, the heart (ECG) and respiration signals converted into rate (beats per minute,

BPM) by first obtaining the differential timing between peaks, i.e., inter-beat intervals (IBI) for ECG,

and inter-breath intervals-(IBrI) for respiration (see Figure 1B). IB(r)Is were then converted to BPM

and interpolated at the original sampling rate to obtain instantaneous HR and RR, which were

downsampled to 20 Hz[68]. SRC and ISC were then calculated across sliding windows. Although ISC and

SRC measures have previously been calculated within 5 second sliding-windows for EEG[4,12],

peripheral responses of heart and respiration are slower, where a typical evoked response can be up

to 10 seconds[14,16,69]. Therefore, we chose a 10 second sliding-time window with a one second overlap.

For SRC, each participants' HR and RR signal was correlated with the extracted spectral flux. For both

HR and RR signals, $p \times t$ matrices were created, where p was the HR or RR signal for each participant

over t timepoints. Signals were correlated in a pairwise fashion, i.e., between all possible participant

pairs within one concert. Values underwent Fisher-Z transformation, were averaged, and then

transformed back to r values (inverse Fisher-Z).

Phase coherence synchrony: ISPC and SRPC

Phase coherence measures were used to assess whether heartbeats and breaths aligned with the

spectral flux rhythms in music. For this, heart and respiration signals were transformed into the

phase domain. Here, we assessed changes of actual heartbeat and breath cycles that vary over time.

To account for these variations, phase was calculated from cycle peaks, see Figure 1B, green dots

marking the R peaks in the heart signals and inspiration peaks in the respiration signals. For the

phase angle of the heartbeat, a value of 0 denoted the R wave (peak of heartbeat cycle) and $\pm\pi$ the

midpoint of the cycle (see Figure 1B). For the phase angle of the respiration cycle, 0 denoted the

inspiration peak, while $\pm\pi$ represents the midpoint of the cycle or maximum exhalation (see Figure

1B). To obtain such values, a continuous sinusoidal wave was fitted to each IB(r)I of detrended and

normalised ECG and respiration signals based on[70] , using the following equation:

$$A \sin (2 \pi f_{(k)} t_{(k)} + \theta)$$

where A is the mean peak amplitude (i.e., average amplitude of signal at time points of

QRS/respiration peak), $f_{(k)}$ is the frequency calculated from IBI (i.e., $peak_{k+1}$-$peak_k$), converted to Hz

($\frac{1}{IBI(s)}$). Phase (θ) was optimised so the peak of the sine wave corresponded to the ECG/respiration

peak. Next, the phase angle was calculated from the real part of the Hilbert using MATLAB's *angle*

function.

SRPC of heartbeats was calculated by assessing coherence between the phase angles of heartbeat

cycles with the spectral flux phase angles corresponding to heart (bandpass filtered at 0.951 - 1.159

Hz), while SRPC of respiration was calculated by assessing coherence between phase angles of

respiration cycles with spectral flux phase angles corresponding to respiration (bandpass filtered

0.268-0.358 Hz) frequencies (see Figure 1C). Coherence was calculated based on the following

formula based on[71]:

292
$$\left| n^{-1} \sum_{r=1}^{n} e^{ik_{tr}} \right|$$

293
where n is the number of phase signals, e^{ik} is Eulers formula (i.e., the complex polar representation

of phase angle k for signal r at time point t). ISPC was assessed in a pairwise fashion between

heartbeat cycle and respiration cycle phase angles of all possible participant pairs within one concert

using the same coherence formula as above.

Adjusting for lags

To account for lags in responses to the stimuli for SRC and SRPC, we adjusted data with a constant

lag to optimally align the stimulus and the corresponding responses. This lag was informed by

previous literature that added lags to account for slight delays in stimulus responses[4,72–74]. To do this,

we first assessed lags on an individual basis. Lags between a stimulus and each participant's response

were calculated within the first 10 seconds after stimulus onset as this initial onset likely evokes the

most reliable responses. We calculated SRC and SRPC at lags up to each individual's full cardiac and

respiration cycle (based on[72]); that is, their mean inter-beat-interval for heart measures (on average:

990 ms) or inter-breath interval for respiration measures (on average: 3400 ms), as it might take up

to one heartbeat/respiration cycle to begin responding to a stimulus[14]. The positive correlations and

phase coherence values were obtained at the optimal lag[73]. We then averaged the individual optimum

lags[74]. This average lag was applied for every participant (i.e., each participant had the same lag for

each stimulus-response synchrony measure). For correlation values, the average optimal lags for HR

and RR were 579 ms and 1573 ms, respectively. For phase coherence values, the average optimal lag

for heart and respiration phase cycles were 500 ms and 1820 ms, respectively. These lags were

applied to all corresponding heart and respiration stimulus-response pairings for the correlation and

phase coherence. Therefore, for each stimulus-response synchrony measure, all participants had the

same lag.

Synchrony across (time-averaged) and within (time-resolved) pieces

Continuous synchrony measures for correlational measures (SRC-HR, ISC-HR, SRC-RR, ISC-RR)) and

phase measures (SRPC and ISPC of heartbeats, and SRPC and ISPC for respiration phase) were

calculated in a time-averaged or time-resolved fashion to test the different hypotheses. For the first

two hypotheses, we calculated time-averaged synchrony values. Rather than averaging synchrony

across the three music pieces - as three musical pieces were thought to yield too few observations -

we obtained time-averaged synchrony per piece section. The Bach piece was split into seven sections,

the Beethoven piece was split into nine sections, and the Messiaen piece was split into nine sections

(for details see Supplementary Table 1). Synchrony was then averaged per piece section, yielding 25

observations per participant, per condition, and per synchrony measure. Averaging across musical

sections was to account for naturalistic grouping as defined by the musical composition. As an

additional check, we also calculated average synchrony within 30 second bins, thereby controlling

for length and section changes (yielding 53 observations per participant, per condition, per

synchrony measure, see Supplementary Materials), and ran identical statistics. To address the third

hypothesis, we calculated time-resolved synchrony values by cutting epochs ±15 seconds relative to

section boundary onsets, to capture event-related respiration and heart responses[14,69], as well as any

anticipatory effects at musical events[75].

Significance-testing

Regarding the first hypothesis, we tested the assumption that responses time-locked to a stimulus

should evoke similar responses across participants[28]. Therefore, we created time-'unlocked' data by

circular shifting ECG and respiration signals. This was done 1000 times and time-averaged synchrony

measures were calculated as described above. For each of the permutated data sets, we calculated a

t-test statistic, creating a null distribution of permutation values. By assessing how many of these

permutation values were higher than or equal to the statistic of the true (time-locked) data, we

determined the chance of true synchrony (see Supplementary Figure 3). Values were corrected for

multiple comparisons using false discovery rate (FDR[76]).

AV versus AO: time-averaged synchrony

For the second hypothesis, time-averaged synchrony measures were compared between AV and AO.

Self-reported engagement was also compared between these conditions. Linear mixed models

(LMMs) with a fixed effect of modality were constructed for each of the synchrony measures as the

outcome variables. As the experimental design meant that data were clustered, the following random

effects were added to account for this non-independence: random intercepts for concert, piece, and

participants, where participants were nested within concerts, while participant and piece were

considered crossed effects. A random slope for participants was also included. This random-effect

structure represents maximal models[77,78]. If these maximal models generated convergence and/or

singularity fit errors, models were simplified following recommendations[77,79]. As output from models

generating errors should not to be trusted[77], we report simplified, error-free models in the results.

Maximal model outputs can be nonetheless found in Supplementary Materials, while all LMMS can

be found in accompanying code on OSF (see link/Footnote (1) above). LMMs were run using *lmer*

from the *lme4* packages[80,81]. Significance values and effect sizes were obtained from the *tab_model*

function from *sjPlot* package[82]. As a recommended sanity check, we also ran conventional *t*-tests

alongside LMMs[83]. This was done by a one sample t-test (or Wilcoxon if data were not normally

distributed), run on the difference between AO and AV values.

AV versus AO: epoched

To test the third hypothesis, synchrony at section boundaries was assessed between modalities

across time windows. Synchrony values in the 30-second epochs centred around section boundaries

were split into smaller time windows[69,84], yielding five 6-second time windows. For each synchrony

measure, linear mixed models (LMMs) with fixed effects of modality and time window were

constructed. We added random intercepts for concert, piece, and participants (participants nested

within concerts; participant and piece as crossed effects) and a random slope for participants. As

above, models generating errors were simplified; error-free models are reported (full models in

Supplementary Materials). Estimated marginal means (using *emmeans* package[85]) were used to

check pairwise comparisons with Bonferroni adjustments. As we predicted that synchrony would

occur at section boundaries, we expected the time window centred at the section boundary to have

significantly higher synchrony values than time windows prior to and after a section boundary. In

case synchrony was high, we investigated what responses were becoming synchronized. Thus, HR

and RR data at section boundaries were taken (i.e., HR and RR as in Figure 1, Panel B). Two LMMs

were constructed, each for aggregated HR and RR values, with fixed effects of time window and

random effects of participant id, piece, and concert, as above.

3. Results

Significance of synchrony measures

In assessing the significance of synchrony measures using a permutation approach and FDR

correction, we found that SRC-HR ($p < .001$), ISC-HR ($p = .048$), and ISC-RR ($p = .048$) showed

significant, above-chance synchrony (see Supplementary Figure 3). SRC-RR (p = .408), SRPC for

heartbeats (p = .412) and respiration (p = .594), and ISPC for heartbeats (p = .594) and respiration (p

= .594) were not above chance level. When synchrony values were averaged in 30 second bins,

significant results were replicated for SRC-HR (p < .001), but not for ISC-RR (p = .051) and ISC-HR (p

= .065). We therefore carried through SRC-HR, and (tentatively) ISC-RR and ISC-HR measures for

further modality comparisons.

Time-averaged synchrony between modalities

Figure 2 shows that compared to the AO condition, the AV condition evoked generally higher SRC-

HR, ISC-HR, and ISC-RR, but only for ISC-HR was this modality difference statistically significant (see

Table 1). This effect for ISC-HR occurred in both error-free and full models (see Supplementary Table

6). When checking conventional t-/Wilcoxon tests (depending on normality), results were similar to

LMM results, though ISC-HR was no longer significant (SRC-HR, $t(21)$ = 1.176, p = 0.253; ISC-HR, $t(21)$

= 1.890, p = .073; ISC-RR, $t(22)$ = .447, p = .659). Although self-reports of engagement were also higher

in AV (see Figure 2), this difference was not significant in the LMM (ß = 0.33, 95% CI [-0.10-0.76], p

= .129, $R^2_{(fixed)}$ = 0.007), nor in the traditional t-test ($t(24)$ = 1.901, p = 0.069). In checking the

behavioural relevance of synchrony as an engagement index, ISC-HR was not significantly correlated

(rho = -0.093, p = .295) nor a significant predictor of self-reported engagement (ß = -0.001, 95% CI

[-0.003-0.001], p = .464, $R^2_{(fixed)}$ = 0.005).

Figure 2. Raincloud plots showing modality differences of audio-only (blue) and audio-visual
(orange) conditions for time-averaged synchrony measures of SRC-HR (far right), ISC-HR (right), and
ISC-RR (left) as well as self-reported engagement (far left). Smaller points represent each data point,
while the larger point represents the mean; error bars represent standard error of the mean. The
shape of the distribution is shown by the half violin.

Table 1. Linear mixed models comparing synchrony between modalities

Predictors	SRC-HR			ISC-HR			ISC-RR		
	Estimates	CI	p	Estimates	CI	p	Estimates	CI	p
(Intercept)	0.050	$0.022 - 0.079$	**0.001**	0.012	$0.002 - 0.022$	**0.020**	0.021	$0.010 - 0.032$	**<0.001**
mod [AV]	0.009	$-0.006 - 0.024$	0.235	0.010	$0.001 - 0.019$	**0.028**	0.002	$-0.007 - 0.012$	0.633
Random Effects									
σ^2	0.01			0.00			0.01		
τ_{00}	0.00 mac:conc			0.00 piece			0.00 mac:conc		
	0.00 piece						0.00 piece		
τ_{11}	0.00 mac1.modAO			0.00 mac1.modAO					
	0.00 mac2.modAV			0.00 mac2.modAV					
ϱ_{01}									
ϱ_{01}									
ICC	0.08			0.05			0.01		
N	3 piece			3 piece			3 piece		
	16 mac			16 mac			16 mac		
	2 conc						2 conc		
Observations	1075			1075			1152		
Marginal R^2 / Conditional R^2	0.002 / 0.079			0.007 / 0.056			0.000 / 0.011		

Synchrony increases at section boundaries, depending on modality

LMMs showed that in time-resolved epochs centred around section boundaries, SRC-HR and ISC-HR were significantly higher in the AV condition, while ISC-RR was higher in AO (see Figure 3 and Table 2, see Supplementary Table 7 and code for maximal models, which replicated these effects).

LMMs additionally showed that time windows predicted synchrony measures at section boundaries. Synchrony measures were significantly higher in the time window centred at the section boundary compared to time windows before (10 seconds before: ISC-HR), and after (5 seconds after: SRC-HR, ISC-RR; 10 seconds after SRC-HR, ISC-HR, and ISC-RR) (see Figure 3 and Table 2).

In assessing the interaction between time window and modality, Figure 3 shows that synchrony was higher before and at section boundaries (window 0) compared to 5 and 10 seconds after section boundary in the AV condition for SRC-HR and ISC-HR. This latter effect was further confirmed by LMMs and by significant pairwise comparisons (Bonferroni corrected) for ISC-HR (Table 3). ISC-RR was higher in AV just after (5 second) section boundary, but this effect was not significant in Bonferroni corrected pairwise comparisons (Table 3).

Upon checking actual HR and RR responses at these time points[16], this synchrony corresponds to typical orienting responses (see Figure 4). HR follows a deceleration-acceleration pattern, where the estimate in HR in window -10 and window 10 is significantly higher than window 0 (intercept, see Table 4 and Supplementary Table 8). Although RR decelerated and accelerated similarly to HR (see Figure 4), these were not significant across time windows.

Figure 3. Synchrony values at 15 seconds prior to and after section boundaries for stimulus correlation of HR (left), inter-subject correlation of HR (middle) and inter-subject correlation of RR (right). The zero point represents the section boundary. Thick lines represent the average, and the ribbon represents the standard error of the mean for audio-only (AO, blue) and audio-visual (AV, orange) conditions. * represents a significant modality effect within that time window.

445 **Table 2.** Linear mixed-effects models for synchrony with predictors of time window and modality.

Linear mixed models

Predictors	SRC-HR			ISC-HR			ISC-RR		
	Estimates	*CI*	*p*	*Estimates*	*CI*	*p*	*Estimates*	*CI*	*p*
(Intercept)	0.04	0.02 – 0.06	**<0.001**	0.02	0.01 – 0.03	**<0.001**	0.05	0.03 – 0.07	**<0.001**
mod [AV]	0.02	0.00 – 0.04	**0.028**	0.02	0.01 – 0.03	**0.002**	-0.03	-0.06 – -0.00	**0.030**
win-10	-0.01	-0.03 – 0.01	0.401	-0.01	-0.02 – -0.00	**0.014**	-0.00	-0.03 – 0.02	0.786
win-5	0.01	-0.01 – 0.02	0.487	-0.00	-0.01 – 0.01	0.544	-0.00	-0.03 – 0.02	0.788
win [5]	-0.02	-0.04 – -0.00	**0.021**	-0.01	-0.02 – 0.00	0.063	-0.04	-0.06 – -0.01	**0.009**
win [10]	-0.02	-0.04 – -0.00	**0.021**	-0.01	-0.02 – -0.00	**0.007**	-0.03	-0.06 – -0.01	**0.018**
modAV:win-10	-0.00	-0.03 – 0.02	0.816	-0.01	-0.02 – 0.01	0.236	0.03	-0.01 – 0.07	0.153
modAV:win-5	0.00	-0.02 – 0.03	0.746	-0.00	-0.02 – 0.01	0.612	0.01	-0.02 – 0.05	0.493
mod [AV] × win [5]	-0.02	-0.05 – 0.00	0.089	-0.02	-0.04 – -0.01	**0.002**	0.04	0.00 – 0.08	**0.034**
mod [AV] × win [10]	-0.03	-0.06 – -0.01	**0.008**	-0.02	-0.04 – -0.01	**0.001**	0.04	-0.00 – 0.08	0.059
Random Effects									
σ^2	0.00			0.00			0.01		
τ_{00}	0.00 $_{piece}$			0.00 $_{mac:conc}$			0.00 $_{mac:conc}$		
	0.00 $_{conc}$								
τ_{11}	0.00 $_{mac.modAV}$			0.00 $_{mac.modAV}$					
	0.00 $_{mac1.modAO}$			0.00 $_{mac1.modAO}$					
	0.00 $_{mac2.modAV}$			0.00 $_{mac2.modAV}$					
ϱ_{01}									
ϱ_{01}									
ICC	0.08			0.06			0.03		
N	2 $_{conc}$			16 $_{mac}$			16 $_{mac}$		
	3 $_{piece}$			2 $_{conc}$			2 $_{conc}$		
	16 $_{mac}$								
Observations	645			645			690		
Marginal R^2 / Conditional R^2	0.120 / 0.191			0.117 / 0.167			0.024 / 0.055		

447 **Table 3.** Pairwise comparison (Bonferroni corrected) between modality (AO-AV) in each time

448 window.

	Contrast AO-AV time window	estimate	SE	df	T ratio	*p*
SRC-HR	Window -10	-0.018	0.010	154	-1.882	.308
	Window -5	**-0.025**	**0.010**	**154**	**-2.631**	**.047**
	Window 0	-0.021	0.010	154	-2.196	.148
	Window 5	0.001	0.010	154	0.094	1.000
	Window 10	0.013	0.010	154	1.402	.814
ISC-HR	Window -10	-0.007	0.005	216	-1.467	0.720
	Window -5	-0.012	0.005	216	-2.414	0.083
	Window 0	**-0.016**	**0.005**	**216**	**-3.123**	**0.010**
	Window 5	0.006	0.005	216	1.200	1.000
	Window 10	0.007	0.005	216	1.413	0.796
ISC-RR	Window -10	0.002	0.014	659	0.154	1.000
	Window -5	0.017	0.014	659	1.207	1.000
	Window 0	0.030	0.014	659	2.175	0.150
	Window 5	-0.012	0.014	659	-0.832	1.000
	Window 10	-0.007	0.014	659	-0.500	1.000

449
450

452 **Figure 4.** Heart rate (HR, left) and respiration rate (RR, right) BPM values at 15 seconds prior to and

453 after section boundaries. The zero point represents the section boundary. Thick lines denote the

454 average BPM, while ribbons around the average line represent the standard error.

456 **Table 4.** Linear mixed models for raw heart and respiration rate at epochs centred around section

457 boundaries

Predictors	Heart Rate			Respiration Rate		
	Estimates	*CI*	*p*	*Estimates*	*CI*	*p*
(Intercept)	61.84	52.52 – 71.17	**<0.001**	18.48	17.12 – 19.83	**<0.001**
win-10	0.46	0.07 – 0.85	**0.022**	0.15	-0.14 – 0.45	0.300
win-5	0.25	-0.14 – 0.64	0.207	-0.24	-0.53 – 0.06	0.115
win [5]	0.37	-0.02 – 0.76	0.061	0.10	-0.19 – 0.40	0.488
win [10]	0.53	0.14 – 0.92	**0.008**	0.07	-0.22 – 0.36	0.638
Random Effects						
σ^2	2.54			1.54		
τ_{00}	50.88 $_{mac:conc}$			7.24 $_{mac:conc}$		
	0.15 $_{piece}$			0.42 $_{piece}$		
	39.81 $_{conc}$			0.04 $_{conc}$		
τ_{11}	10.74 $_{mac1.modAO}$					
	20.67 $_{mac2.modAV}$					
ϱ_{01}						
ϱ_{01}						
ICC	0.97			0.83		
N	2 $_{conc}$			2 $_{conc}$		
	3 $_{piece}$			3 $_{piece}$		
	16 $_{mac}$			16 $_{mac}$		
Observations	645			690		
Marginal R^2 / Conditional R^2	0.000 / 0.973			0.002 / 0.833		

4. Discussion

An important aspect of (live) musical experiences is the listeners' engagement. Despite the methodological challenges in assessing such musical experiences, dynamic engagement is increasingly assessed with neurophysiological synchrony measures. By extending these techniques to a naturalistic setting in a concert hall, the current study pushed the boundaries in how to measure and analyse musical experiences. Importantly, we also compared approaches to gauge the robustness of these cardiorespiratory synchrony analyses. As engagement might be enhanced by seeing musicians perform[8,10], participants were presented with AO and AV performances of Western classical music in a concert setting while cardiorespiratory measures were continuously measured. We show that heart rate (i.e., *speed* of heartbeats in BPM) correlated with spectral flux of music above chance. Additionally, respiration rate and (tentatively) heart rate correlated across audience members above chance. Musical rhythms did not align with heartbeat rhythms nor breathing rhythms above chance. We also find that compared to the AO condition, AV music performances evoked significantly higher ISC-HR; this effect was more robust on a time-resolved, compared to time-averaged, level.

Audience heart rate and respiration rate correlation

HR correlated with spectral flux (SRC-HR) above chance level, indicating that the acceleration/deceleration of audiences' HR correlated with increases/decreases of spectral flux in the music. ISC-HR and ISC-RR measures were also above chance level, indicating that acceleration/deceleration of HR and RR correlated across audience members experiencing music simultaneously, supporting previous research showing that ISC of HR[18,46] and EEG[4] is evoked when participants simultaneously experience the same auditory stimuli. We note, however, that although

ISC-HR was significant in the main analysis of time-averaged synchrony within musical sections (see

Supplementary Table 1), when ISC-HR was time-averaged across 30-second bins this measure did

not meet the significance threshold after correcting for multiple comparisons. Therefore, it might be

important to assess synchrony related to musical structure and on a time-resolved level; see below

discussion on 'Time-resolved synchrony'. That ISC-RR was above chance level somewhat contrasts

findings of Madsen and Parra[46], who found ISC-RR not to be significant. One explanation could be due

to stimuli differences. Here, we used music stimuli, whereas Madsen and Parra used instructional

video (speech). More importantly, the setting of a concert hall might have increased the engagement

of the listeners compared to the laboratory setting. Our findings suggest that HR and RR (i.e., speed

of heartbeats/breaths) may become synchronized across groups of people experiencing auditory

stimuli (like music) simultaneously, but may be context dependent.

Heartbeat and breathing cycles do not rhythmically align to music

Both musical rhythm–heartbeat phase synchrony and musical rhythm–breath phase synchrony

(SRPC measures) were not significantly above chance level. In other words, musical rhythms aligned

with neither audiences' breathing cycles nor their heartbeat cycles. Although this lack of phase

synchrony was expected for heart rhythms[51-53], this contrasts previous findings by Etzel et al[49] and

Haas et al[50] that breathing aligns with musical beats. There are a few potential reasons for this

discrepancy between results. First, Etzel et al. and Haas et al. compared respiratory signals to the

beat of music, while the current study assesses spectral flux filtered at respiration frequencies.

Although our method is therefore not comparable with this previous research, we nonetheless

suggest that our method is more generalisable for future naturalistic research. It seemed like the

previous research chose music to roughly correspond to healthy breathing frequencies (see Haas et

al, Table 1, Figure 1 and 2; Etzel et al. Table 1). However, much music might not correspond to such

specific frequencies. Indeed, the musical choice for the current study was motivated by the need to

present a typical musical program, where the beat did not fall into the range of healthy

heartbeats/breathing (see Supplementary Table 1). It therefore seemed problematic to assess

synchrony of the actual beat with heartbeats/breaths. Assessing musical features that strongly

correlate with beat onsets (spectral flux[24]) nonetheless yields a somewhat comparable approach.

Additionally, extracting the energy at the natural frequency of heartbeats/breathing allows assessing

music that has beats falling outside of the natural breathing/heartbeat range, providing a useful

measure for future research. A second explanation for not replicating musical beat-respiration

synchrony, is that previous studies used music with relatively regular rhythmic and metric

structures[49,50], while the stimuli in the current study included one contemporary piece with

ambiguous rhythms. However, there were no respiration synchrony differences between pieces with

regular (Bach, Beethoven) and irregular (Messiaen) metric structures (see Supplementary Table 3).

A third explanation could be that respiration is more likely to align with beats in laboratory

experiments as the attention (per instruction) is mainly on music. Beyond complexities in musical

structure, a real-world setting presents several additional factors, such as the visual and social

aspects[10,86] , where attention may fluctuate a lot more between the music and other features. Thus,

although neural[72,74] and respiratory[49,50] rhythms may align to auditory beats in a laboratory setting,

the current results suggest that this may not occur to music with broader tempi in a music concert

setting, at least for this choice of Western Classical music.

Heartbeats and breaths likewise did not significantly align in phase between participants. One

explanation could be the lack of social interaction, which is a crucial aspect in higher-order phase

coupling[40]. Indeed, phase synchrony between participants has previously been found in dancing[87,88]

and musical ensembles [38,39,89,90] where there is a clear shared goal of a coordinated performance

(see[40,41]). Typical Western classical concert etiquette calls for enhanced focus on the performance,

thus reducing body movements and interaction between audience members[10]. Indeed, ratings for

'urge to move to the music' (see[7]) were heavily skewed to low ratings; even testing significance of

phase coherence in audience members who only had high 'urge to move' ratings did not yield

significant results (see Supplementary Table 5). Additionally, videos showed that the audience

members respected this etiquette and generally did not move too much[7]. Therefore, we suggest that

phase synchrony – typically seen in interactive settings[40] – may not occur in non-interactive settings.

Time-averaged synchrony may not relate to naturalistic engagement

Next, synchrony measures were compared between AV and AO conditions. ISC-HR was significantly

greater in AV performances. However, this result was not shown to be robust with a sanity check

using conventional t-test. Although our data are more appropriate for LMMs due to naturalistic

paradigm and physiological data[78], the fact that the significance is not replicated in a simpler t-test

thus allows only cautious interpretation[83]. Another reason to be cautious is that – contrary to what

was expected – ISC-HR was not significantly related to engagement that was self-reported after each

piece. The current findings contrast findings from Ki et al.[31] who found that neural ISC was stronger

in attend than non-attend conditions, especially in audiovisual stimuli. The current finding also

contrasts studies showing that HR[18] and neural[19] synchrony can reflect engagement. One explanation

for the discrepancy in results is that these previous studies had explicit attend/non-attend

conditions, where participants were instructed to count backwards in the non-attend conditions.

Here, participants experienced concert performances naturally with no formal instruction. This

suggests that while ISC (of HR) could be a marker in clearly defined engagement conditions, it might

not necessarily be generalisable to less controlled settings. Additionally, inter- and intra-individual

differences may influence synchrony and engagement[55,91,92] , and could be a potential venture for

future studies to explore individual versus group synchrony. Overall, we show time-averaged

synchrony does not reliably reflect overall self-reported engagement measurements. With this in

mind, we anticipated results for time-resolved synchrony might yield clearer results.

Time-resolved synchrony reveals synchronized heart rate orienting response

Engagement with music typically fluctuates across time[4,12,54]. Thus, we assessed synchrony changes

between modalities on a time-resolved scale. We 'zoomed in' on synchronized responses at salient

moments in the music, here defined as section boundaries in music (see Supplementary Table 1),

which are important structural locations in music. The current results show that both ISC-HR and

SRC-HR increased at such section boundaries, supporting and extending previous work on ISC[12,16] to

SRC. Although we did not have time-resolved self-reports to assess the relation of synchrony to

engagement, we assessed the continuous HR and RR. We show that synchronized responses reflected

significant deceleration-acceleration patterns in HR and RR. Such patterns have been long associated

with a typical orienting response, which marks an increase in attentiveness[93] to anticipate and

perceive events[14,94–96]. Thus, such increases of cardiorespiratory synchrony at section boundaries

suggest increased engagement with music at these time points, related to auditory cues indicating a

section is ending, such as a cadential ending (i.e., harmonic closure in Bach and Beethoven) and a

slight slowing down or change in tempo and/or material (see Supplementary Table 1). This

physiological synchrony at salient music events suggests a bottom-up capturing of engagement

during a shared musical experience[40]. We acknowledge that the definition of salient moments to

explore engagement are limited in the current study to section boundaries. However, humans also

respond to segmentation not only at section boundaries[4,12,61,97], but also at smaller phrasal[98] and

larger, movement[68] levels. Thus, the current work provides a starting point for exploring engagement

to salient moments; future work could compare 'non' salient moments, transitions, and segments at

lower and higher hierarchical musical levels.

Importantly, Figure 3 shows that ISC-HR significantly increased at section boundaries more so in the

AV condition. This increase suggests that visual information further increased engagement at section

boundaries. Indeed, previous performance studies have found that certain gestures cue and

anticipate structurally important locations, such as leaning forward towards a cadence(harmonic

closure)[99] , increased movement amplitude[100] , or finishing 'flourish' gestures[101] as well as deep

breaths and sweeping motions prior to onset of new phrases[102]. Indeed, videos show that the pianist

tended to lean forward at section boundaries (though see limitations regarding analysis of visual

features). Therefore, a combination of audio and visual aspects likely increases engagement with

music at a local level, as shown most robustly with ISC-HR.

5. Limitations

Although we show greater synchrony in the AV (compared to AO) condition, it is unclear whether

this synchrony increase might simply be related to increased information processing of both visual

and audio information, compared to simply audio information. For example, RR of audiences did not

synchronise to the spectral flux of the music (insignificant SRC-RR result), but RR did significantly

synchronise across audience members (ISC-RR). This suggests that the audiences' RR synchrony was

rather driven by something else than the acoustic signal, perhaps the musician movements (see[7]).

Indeed, we saw that SRC-RR was more significant in the AV than the AO condition (see

Supplementary Table 4). As it was not possible to extract reliable visual features in the current study

due to the camera location in the concert hall (see angle in Figure 1A), future work would therefore

need to assess responses not only to auditory features, but also visual features, such as facial

expressions[103], quantity of motion, and speed of movements[104] to shed more light on how much

variance in the audience experience can be explained by auditory and by visual features. Additionally,

we did not test a visual only condition which would allow us to disentangle the difference between

added information processing and actual engagement. Finally, we acknowledge that aspects of

'liveness' may have likewise contributed to the current results. For example, previous evidence

suggested that musicians have different perceived expressiveness when performing with no

audience (rehearsal condition) compared to performing in front of an audience (recital condition[105]).

However, as the literature is mixed as to whether liveness influences emotional audience
experience[106] (for further discussion see[7,107]), but the literature on audio-visual influences on music
appreciation is relatively robust[9], it seems more plausible that the current results are explained by
the modality difference rather than liveness.
612

6. Conclusion

The current study pushed the boundaries of how to measure and analyse music listening experiences.
One crucial aspect of such a musical experience is the listener's engagement, which can be assessed
by measuring neurophysiological synchrony. As engagement could be affected by modality, we
explored engagement differences between audio-visual and audio-only music performances in a
Western classical concert setting, using cardiorespiratory synchrony as a measure of engagement.
We extended studies investigating neural synchrony[4,17], by comparing the robustness of different
analyses to calculate peripheral physiological synchrony[18,46,54] in music concert settings[2,15,16]. We
showed that seeing musicians perform can heighten an audience's engagement, though this might be
context-dependent, and is best assessed by time-resolved (rather than time-averaged) inter-subject
correlation of HR (i.e., synchrony of heartbeat speed). Rhythmic alignment (phase synchrony) of
heartbeats/respiration with other audience members/music) were below chance level. However,
questions arose whether phase coherence might become more relevant in contexts where listeners
can move and interact, for example, at pop and rock concerts. More studies, involving a wider range
of genres and contexts are needed to further understand synchrony dynamics in real-world music
contexts. Importantly, the current study highlights the advantages of timeseries analysis in
quantifying dynamic engagement, which could be applied not only to data collected during concert
music listening, but in different naturalistic settings (e.g., open air concerts).

Acknowledgements

The authors would like to thank Lea Fink, our music theorist who advised on the musical sections. Thanks also go to Klaus Frieler and Örjan De Manzano for discussions about statistics. Many thanks also to the ArtLab team, who assisted in concert preparations.

Competing interests. The authors declare no competing interests.

7.   References

1.   Gabrielsson, A. & Wik, S. L. Strong Experiences Related to Music: A Descriptive System. *Music. Sci.* **7**, 157–217 (2003).

2.   Kragness, H. E. *et al.* An itsy bitsy audience: Live performance facilitates infants' attention and heart rate synchronization. *Psychol. Aesthet. Creat. Arts* (2023) doi:10.1037/aca0000597.

3.   Lamont, A. University students' strong experiences of music: Pleasure, engagement, and meaning. *Music. Sci.* **15**, 229–249 (2011).

4.   Kaneshiro, B., Nguyen, D. T., Norcia, A. M., Dmochowski, J. P. & Berger, J. Natural music evokes correlated EEG responses reflecting temporal structure and beat. *NeuroImage* **214**, 116559 (2020).

5.   Omigie, D. & Mencke, I. A model of time-varying music engagement. *Philos. Trans. R. Soc. B Biol. Sci.* **379**, 20220421 (2024).

6.   Schubert, E., Vincs, K. & Stevens, C. J. Identifying Regions of Good Agreement among Responders in Engagement with a Piece of Live Dance. *Empir. Stud. Arts* **31**, 1–20 (2013).

7.   Czepiel, A., Fink, L. K., Seibert, C., Scharinger, M. & Kotz, S. A. Aesthetic and physiological effects of naturalistic multimodal music listening. *Cognition* **239**, 105537 (2023).

8.   Dobson, M. C. New Audiences for Classical Music: The Experiences of Non-attenders at Live Orchestral Concerts. *J. New Music Res.* **39**, 111–124 (2010).

9.   Platz, F. & Kopiez, R. When the Eye Listens: A Meta-analysis of How Audio-visual Presentation Enhances

the Appreciation of Music Performance. *Music Percept.* **30**, 71–83 (2012).

10.   Wald-Fuhrmann, M. *et al.* Music Listening in Classical Concerts: Theory, Literature Review, and Research

Program. *Front. Psychol.* **12**, (2021).

11.   Olsen, K. N., Dean, R. T. & Stevens, C. J. A continuous measure of musical engagement contributes to

prediction of perceived arousal and valence. *Psychomusicology Music Mind Brain* **24**, 147–156 (2014).

12.   Dauer, T. *et al.* Inter-subject Correlation While Listening to Minimalist Music: A Study of

Electrophysiological and Behavioral Responses to Steve Reich's Piano Phase. *Front. Neurosci.* **15**, (2021).

13.   Kaneshiro, B., Nguyen, D. T., Norcia, A. M., Dmochowski, J. P. & Berger, J. Inter-subject correlation of

electroencephalographic and behavioural responses reflects time-varying engagement with natural

music. *Eur. J. Neurosci.* ejn.16324 (2024) doi:10.1111/ejn.16324.

14.   Bradley, M. M. & Lang, P. J. Affective reactions to acoustic stimuli. *Psychophysiology* **37**, 204–215 (2000).

15.   Chabin, T. *et al.* Interbrain emotional connection during music performances is driven by physical

proximity and individual traits. *Ann. N. Y. Acad. Sci.* **1508**, 178–195 (2022).

16.   Czepiel, A. *et al.* Synchrony in the periphery: inter-subject correlation of physiological responses during

live music concerts. *Sci. Rep.* **11**, 22457 (2021).

17.   Dmochowski, J. P., Sajda, P., Dias, J. & Parra, L. C. Correlated Components of Ongoing EEG Point to

Emotionally Laden Attention – A Possible Marker of Engagement? *Front. Hum. Neurosci.* **6**, (2012).

18.   Pérez, P. *et al.* Conscious processing of narrative stimuli synchronizes heart rate between individuals. *Cell

Rep.* **36**, 109692 (2021).

19.   Madsen, J., Margulis, E. H., Simchy-Gross, R. & Parra, L. C. Music synchronizes brainwaves across listeners

with strong effects of repetition, familiarity and training. *Sci. Rep.* **9**, 3576 (2019).

20.   Albouy, P., Benjamin, L., Morillon, B. & Zatorre, R. J. Distinct sensitivity to spectrotemporal modulation

supports brain asymmetry for speech and melody. *Science* **367**, 1043–1047 (2020).

21.   Albouy, P. *et al.* Spectro-temporal acoustical markers differentiate speech from song across cultures. *Nat.

Commun.* **15**, 4835 (2024).

22.   Obleser, J., Herrmann, B. & Henry, M. J. Neural Oscillations in Speech: Don't be Enslaved by the Envelope.

Front. Hum. Neurosci. **6**, (2012).

23. Schultz, B. G., Brown, R. M. & Kotz, S. A. Dynamic acoustic salience evokes motor responses. *Cortex* **134**,
320–332 (2021).

24. Weineck, K., Wen, O. X. & Henry, M. J. Neural synchronization is strongest to the spectral flux of slow music
and depends on familiarity and beat salience. *eLife* **11**, e75515 (2022).

25. Alluri, V. *et al.* Large-scale brain networks emerge from dynamic processing of musical timbre, key and
rhythm. *NeuroImage* **59**, 3677–3689 (2012).

26. Lartillot, O. & Toiviainen, P. A Matlab toolbox for musical feature extraction from audio. in *International
conference on digital audio effects* (Bordeaux, 2007).

27. Burger, B., Saarikallio, S., Luck, G., Thompson, M. R. & Toiviainen, P. Relationships Between Perceived
Emotions in Music and Music-induced Movement. *Music Percept.* **30**, 517–533 (2013).

28. Nastase, S. A., Gazzola, V., Hasson, U. & Keysers, C. Measuring shared responses across subjects using
intersubject correlation. *Soc. Cogn. Affect. Neurosci.* **14**, 667–685 (2019).

29. Hasson, U., Nir, Y., Levy, I., Fuhrmann, G. & Malach, R. Intersubject Synchronization of Cortical Activity
During Natural Vision. *Science* **303**, 1634–1640 (2004).

30. Kang, O. & Wheatley, T. Pupil dilation patterns spontaneously synchronize across individuals during
shared attention. *J. Exp. Psychol. Gen.* **146**, 569–576 (2017).

31. Ki, J. J., Kelly, S. P. & Parra, L. C. Attention Strongly Modulates Reliability of Neural Responses to Naturalistic
Narrative Stimuli. *J. Neurosci.* **36**, 3092–3101 (2016).

32. Irsik, V. C., Johnsrude, I. S. & Herrmann, B. Neural Activity during Story Listening Is Synchronized across
Individuals Despite Acoustic Masking. *J. Cogn. Neurosci.* **34**, 933–950 (2022).

33. Schmälzle, R., Häcker, F. E. K., Honey, C. J. & Hasson, U. Engaged listeners: shared neural processing of
powerful political speeches. *Soc. Cogn. Affect. Neurosci.* **10**, 1137–1143 (2015).

34. Dikker, S. *et al.* Brain-to-Brain Synchrony Tracks Real-World Dynamic Group Interactions in the Classroom.
Curr. Biol. **27**, 1375–1380 (2017).

35. Peelle, J. E., Gross, J. & Davis, M. H. Phase-Locked Responses to Speech in Human Auditory Cortex are
Enhanced During Comprehension. *Cereb. Cortex* **23**, 1378–1387 (2013).

36. Jones, M. R. & Boltz, M. Dynamic attending and responses to time. *Psychol. Rev.* **96**, 459–491 (1989).

37. Large, E. W. & Snyder, J. S. Pulse and Meter as Neural Resonance. *Ann. N. Y. Acad. Sci.* **1169**, 46–57 (2009).

38. Gugnowska, K. *et al.* Endogenous sources of interbrain synchrony in duetting pianists. *Cereb. Cortex* **32**,

4110–4127 (2022).

39. Müller, V. & Lindenberger, U. Cardiac and Respiratory Patterns Synchronize between Persons during Choir

Singing. *PLOS ONE* **6**, e24893 (2011).

40. Dumas, G. & Fairhurst, M. T. Reciprocity and alignment: quantifying coupling in dynamic interactions. *R.*

Soc. Open Sci. **8**, 210138 (2021).

41. Fairhurst, M. T., Janata, P. & Keller, P. E. Being and Feeling in Sync with an Adaptive Virtual Partner: Brain

Mechanisms Underlying Dynamic Cooperativity. *Cereb. Cortex* **23**, 2592–2600 (2013).

42. Tschacher, W. *et al.* Physiological synchrony in audiences of live concerts. *Psychol. Aesthet. Creat. Arts* **17**,

152–162 (2021).

43. Criscuolo, A., Schwartze, M. & Kotz, S. A. Cognition through the lens of a body–brain dynamic system. *Trends*

Neurosci. S0166223622001229 (2022) doi:10.1016/j.tins.2022.06.004.

44. Klimesch, W. The frequency architecture of brain and brain body oscillations: an analysis. *Eur. J. Neurosci.*

48, 2431–2453 (2018).

45. Parviainen, T., Lyyra, P. & Nokia, M. S. Cardiorespiratory rhythms, brain oscillatory activity and cognition:

review of evidence and proposal for significance. *Neurosci. Biobehav. Rev.* **142**, 104908 (2022).

46. Madsen, J. & Parra, L. C. Cognitive processing of a common stimulus synchronizes brains, hearts, and eyes.

PNAS Nexus **1**, pgac020 (2022).

47. Al, E. *et al.* Heart–brain interactions shape somatosensory perception and evoked potentials. *Proc. Natl.*

Acad. Sci. **117**, 10575–10584 (2020).

48. Grund, M. *et al.* Respiration, Heartbeat, and Conscious Tactile Perception. *J. Neurosci.* **42**, 643–656 (2022).

49. Etzel, J. A., Johnsen, E. L., Dickerson, J., Tranel, D. & Adolphs, R. Cardiovascular and respiratory responses

during musical mood induction. *Int. J. Psychophysiol.* **61**, 57–69 (2006).

50. Haas, F., Distenfeld, S. & Axen, K. Effects of perceived musical rhythm on respiratory pattern. *J. Appl. Physiol.*

61, 1185–1191 (1986).

51. Ellis, R. J. & Thayer, J. F. Music and Autonomic Nervous System (Dys)Function. *Music Percept.* **27**, 317–326

(2010).

52. Koelsch, S. & Jäncke, L. Music and the heart. *Eur. Heart J.* **36**, 3043–3049 (2015).

53. Mütze, H., Kopiez, R. & Wolf, A. The effect of a rhythmic pulse on the heart rate: Little evidence for

rhythmical 'entrainment' and 'synchronization'. *Music. Sci.* **24**, 377–400 (2020).

54. Fink, L. K., Hurley, B. K., Geng, J. J. & Janata, P. A linear oscillator model predicts dynamic temporal attention

and pupillary entrainment to rhythmic patterns. *J. Eye Mov. Res.* **11**, 10.16910/jemr.11.2.12 (2018).

55. Presicce, G. & Bailes, F. Engagement and visual imagery in music listening: An exploratory study.

Psychomusicology Music Mind Brain **29**, 136–155 (2019).

56. Sridharan, D., Levitin, D. J., Chafe, C. H., Berger, J. & Menon, V. Neural Dynamics of Event Segmentation in

Music: Converging Evidence for Dissociable Ventral and Dorsal Networks. *Neuron* **55**, 521–532 (2007).

57. Burunat, I., Levitin, D. J. & Toiviainen, P. Breaking (musical) boundaries by investigating brain dynamics of

event segmentation during real-life music-listening. *Proc. Natl. Acad. Sci.* **121**, e2319459121 (2024).

58. Lerdahl, F. & Jackendoff, R. An Overview of Hierarchical Structure in Music. *Music Percept. Interdiscip. J.* **1**,

229–252 (1983).

59. Clarke, E. F. & Krumhansl, C. L. Perceiving Musical Time. *Music Percept. Interdiscip. J.* **7**, 213–251 (1990).

60. Deliege, I. Grouping Conditions in Listening to Music: An Approach to Lerdahl & Jackendoff's Grouping

Preference Rules. *Music Percept. Interdiscip. J.* **4**, 325–359 (1987).

61. Phillips, M. *et al.* What Determines the Perception of Segmentation in Contemporary Music? *Front. Psychol.*

11, 1001 (2020).

62. Popescu, T., Widdess, R. & Rohrmeier, M. Western listeners detect boundary hierarchy in Indian music: a

segmentation study. *Sci. Rep.* **11**, 3112 (2021).

63. Davidson, J. W. Visual Perception of Performance Manner in the Movements of Solo Musicians. *Psychol.*

Music **21**, 103–113 (1993).

64. Herrmann, B. & Johnsrude, I. S. Absorption and Enjoyment During Listening to Acoustically Masked Stories.

Trends Hear. **24**, 233121652096785 (2020).

65. Kandylaki, K. D., Utroša-Škerjanec, M. & Kotz, S. Rhythmic Regularity in Speech is Modulated by Amplitude,

Pitch, Duration, and the Spectral Center of Gravity. *SSRN Electron. J.* (2022) doi:10.2139/ssrn.4213992.

66. Tzanetakis, G. & Cook, P. Musical genre classification of audio signals. *IEEE Trans. Speech Audio Process.* **10**,

293–302 (2002).

67.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software for Advanced Analysis

768   of MEG, EEG, and Invasive Electrophysiological Data. *Comput. Intell. Neurosci.* **2011**, 1–9 (2011).

769   68.   Merrill, J., Czepiel, A., Fink, L. T., Toelle, J. & Wald-Fuhrmann, M. The aesthetic experience of live concerts:

770   Self-reports and psychophysiology. *Psychol. Aesthet. Creat. Arts* **17**, 134–151 (2021).

771   69.   Sammler, D., Grigutsch, M., Fritz, T. & Koelsch, S. Music and emotion: Electrophysiological correlates of the

772   processing of pleasant and unpleasant music. *Psychophysiology* **44**, 293–304 (2007).

773   70.   Assaneo, M. F., Rimmele, J. M., Sanz Perl, Y. & Poeppel, D. Speaking rhythmically can shape hearing. *Nat.*

774   *Hum. Behav.* **5**, 71–82 (2021).

775   71.   Cohen. *Analyzing Neural Time Series Data: Theory and Practice*. (The MIT Press, Cambridge, Massachusetts,

776   2014).

777   72.   Harding, E. E., Sammler, D., Henry, M. J., Large, E. W. & Kotz, S. A. Cortical tracking of rhythm in music and

778   speech. *NeuroImage* **185**, 96–101 (2019).

779   73.   Luck, G., Toiviainen, P. & Thompson, M. R. Perception of Expression in Conductors' Gestures: A Continuous

780   Response Study. *Music Percept.* **28**, 47–57 (2010).

781   74.   Vanden Bosch der Nederlanden, C. M., Joanisse, M. F. & Grahn, J. A. Music as a scaffold for listening to

782   speech: Better neural phase-locking to song than speech. *NeuroImage* **214**, 116767 (2020).

783   75.   Wassiliwizky, E., Koelsch, S., Wagner, V., Jacobsen, T. & Menninghaus, W. The emotional power of poetry:

784   neural circuitry, psychophysiology and compositional principles. *Soc. Cogn. Affect. Neurosci.* **12**, 1229–

785   1240 (2017).

786   76.   Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under dependency.

787   *Ann. Stat.* **29**, (2001).

788   77.   Barr, D. J., Levy, R., Scheepers, C. & Tily, H. J. Random effects structure for confirmatory hypothesis testing:

789   Keep it maximal. *J. Mem. Lang.* **68**, 255–278 (2013).

790   78.   Page-Gould, E. Multilevel Modeling. in *Handbook of Psychophysiology* (eds. Cacioppo, J. T., Tassinary, L. G.

791   & Berntson, G. G.) 662–678 (Cambridge University Press, 2016). doi:10.1017/9781107415782.030.

792   79.   Barr, D. J. *Learning Statistical Models through Simulation in R: An Interactive Textbook*. (2021).

793   80.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using {lme4}. *J. Stat. Softw.*

794   **67**, 1–48 (2015).

81. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. {lmerTest} Package: Tests in Linear Mixed Effects

Models. *J. Stat. Softw.* **82**, 1–26 (2017).

82. Lüdecke, D. sjPlot: Data Visualization for Statistics in Social Science. (2023).

83. Arnqvist, G. Mixed Models Offer No Freedom from Degrees of Freedom. *Trends Ecol. Evol.* **35**, 329–335

(2020).

84. Dellacherie, D., Roy, M., Hugueville, L., Peretz, I. & Samson, S. The effect of musical experience on emotional

self-reports and psychophysiological responses to dissonance: Psychophysiology of musical emotion.

Psychophysiology **48**, 337–349 (2011).

85. Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Squares Means. (2021).

86. Tervaniemi, M. The neuroscience of music – towards ecological validity. *Trends Neurosci.* **46**, 355–364

(2023).

87. Hartmann, M. *et al.* Kinematics of perceived dyadic coordination in dance. *Sci. Rep.* **9**, 15594 (2019).

88. Toiviainen, P. & Carlson, E. Embodied Meter Revisited. *Music Percept.* **39**, 249–267 (2022).

89. Lange, E. B. *et al.* In touch: Cardiac and respiratory patterns synchronize during ensemble singing with

physical contact. *Front. Hum. Neurosci.* **16**, (2022).

90. Vickhoff, B. *et al.* Music structure determines heart rate variability of singers. *Front. Psychol.* **4**, (2013).

91. Forster, S. & Lavie, N. Distracted by your mind? Individual differences in distractibility predict mind

wandering. *J. Exp. Psychol. Learn. Mem. Cogn.* **40**, 251–260 (2014).

92. Wohltjen, S., Toth, B., Boncz, A. & Wheatley, T. Synchrony to a beat predicts synchrony with other minds.

Sci. Rep. **13**, 3591 (2023).

93. Barry, R. J. & Sokolov, E. N. Habituation of phasic and tonic components of the orienting reflex. *Int. J.*

Psychophysiol. **15**, 39–42 (1993).

94. Park, H.-D., Correia, S., Ducorps, A. & Tallon-Baudry, C. Spontaneous fluctuations in neural responses to

heartbeats predict visual detection. *Nat. Neurosci.* **17**, 612–618 (2014).

95. Proverbio, A. M. *et al.* Non-expert listeners show decreased heart rate and increased blood pressure (fear

bradycardia) in response to atonal music. *Front. Psychol.* **6**, (2015).

96. Stekelenburg, J. & Boxtel, A. V. Inhibition of pericranial muscle activity, respiration, and heart rate enhances

auditory sensitivity. *Psychophysiology* **38**, 629–641 (2001).

97. Czepiel, A. Multimodal_piano_periphys: Data and analysis code. (2023).

98. Teng, X., Larrouy-Maestri, P. & Poeppel, D. Segmenting and Predicting Musical Phrase Structure Exploits

Neural Gain Modulation and Phase Precession. *J. Neurosci.* e1331232024 (2024)

doi:10.1523/JNEUROSCI.1331-23.2024.

99. Davidson, J. W. Bodily movement and facial actions in expressive musical performance by solo and duo

instrumentalists: Two distinctive case studies. *Psychol. Music* **40**, 595–633 (2012).

100.Thompson & Luck, G. Exploring relationships between pianists' body movements, their expressive

intentions, and structural elements of the music. *Music. Sci.* **16**, 19–40 (2012).

101.Wanderley, M. M., Vines, B. W., Middleton, N., McKay, C. & Hatch, W. The Musical Significance of Clarinetists'

Ancillary Gestures: An Exploration of the Field. *J. New Music Res.* **34**, 97–113 (2005).

102.Vines, B. W., Wanderley, M. M., Krumhansl, C. L., Nuzzo, R. L. & Levitin, D. J. Performance Gestures of

Musicians: What Structural and Emotional Information Do They Convey? in *Gesture-Based*

Communication in Human-Computer Interaction (eds. Camurri, A. & Volpe, G.) 468–478 (Springer, Berlin,

Heidelberg, 2004). doi:10.1007/978-3-540-24598-8_43.

103.Waddell, G. & Williamon, A. Eye of the Beholder: Stage Entrance Behavior and Facial Expression Affect

Continuous Quality Ratings in Music Performance. *Front. Psychol.* **8**, 513 (2017).

104.Lillywhite, A. *et al.* A functional magnetic resonance imaging examination of audiovisual observation of a

point-light string quartet using intersubject correlation and physical feature analysis. *Front. Neurosci.* **16**,

921489 (2022).

105.Chang-Arana, Á. M., Mavrolampados, A., Pokki, N. & Thompson, M. R. When nerves hit: The effect of trait

anxiety, situational stress, and task mastery on the perception and interpersonal accuracy of musical

expressiveness. *Front. Psychol.* **13**, 1061922 (2023).

106.Belfi, A. M., Samson, D. W., Crane, J. & Schmidt, N. L. Aesthetic Judgments of Live and Recorded Music: Effects

of Congruence Between Musical Artist and Piece. *Front. Psychol.* **12**, (2021).

107.Czepiel, A. M. Assessing real-world music listening in concerts: Aesthetic experiences and peripheral

physiological responses. (maastricht university, 2023). doi:10.26481/dis.20231002ac.

849