1
2
3
4
5
6 Emergence of active contractile patterns alters monolayer force generation and
7 transmission in response to focal adhesion distribution
8
9

10 John Robert Davis[1,2‡*], Josephine Solowiej-Wedderburn[3,5,6‡], Sebastián L. Vega[7,8] Jason A.
11 Burdick[9], Carina Dunlop[4,10*], and Nicolas Tapon[1,10*]
12
[1] Apoptosis and Proliferation Control Laboratory, The Francis Crick Institute, London, UK
[2] Hungry Tissue Lab, Division of Molecular and Cellular Function, School of Biological
Sciences, Faculty of Biology Medicine Health, University of Manchester, Manchester, UK
[3] School of Mathematics and Physics, University of Surrey, Guildford, UK
[4] Department of Mathematics, University College London, London, UK
[5] Department of Mathematics and Mathematical Statistics, Umeå University, Umeå, Sweden
[6] Integrated Science Lab, Umeå University, Umeå, Sweden
[7] Department of Biomedical Engineering, Rowan University, Glassboro, NJ, USA
[8] Department of Orthopaedic Surgery, Cooper Medical School of Rowan University, Camden,
NJ, USA
[9] BioFrontiers Institute and Department of Chemical and Biological Engineering, College of
Engineering and Applied Science, University of Colorado, Boulder, USA
[10] Senior authors
‡ Authors contributed equally
* Correspondence: john-robert.davis@manchester.ac.uk (JRD), carina.dunlop@ucl.ac.uk
(CD), and nic.tapon@crick.ac.uk (NT)

Abstract

For tissue development, cells must generate contractile forces which are transmitted to their surrounding matrix or neighbouring cells via adhesion complexes. It is often envisaged that a simple linear counterbalance of cell generated stress with extracellular matrix (ECM) traction forces exists. However, experimental evidence indicates that modulating cell-ECM attachment does not necessarily lead to expected reciprocal changes in intercellular stresses. As ECM composition or mechanical properties are rarely uniform, it is important to understand the complexity of how focal adhesions alter stress transmission and the force-balance of a tissue. To address this, we confined monolayers on adhesive patterns altering focal adhesion distribution. Traction force microscopy and laser ablations of cell-cell junctions were used to examine stresses across epithelial monolayers whilst modulating substrate stiffness. We show that monolayers reach different force-balance states depending on focal adhesion distribution. Using an active matter model and confirmed experimentally, we reveal that a force-balance is generated by non-uniform patterns of cell contractility linked to adhesion patterning. This work highlights the importance of integrating the position of cell-ECM attachments into our vision of the mechanical landscape of living tissues.

Teaser
To infer a tissue's force-balance, positional information of focal adhesion distribution needs to be integrated due to the emergence of non-uniform patterns of cell contractility.

Introduction

A key aspect of many developmental, physiological, and pathological processes is the ability of cells to generate and transmit forces across a tissue. Cells transmit intracellular stresses to their environment, and this is essential for orchestrating cooperative behaviours such as morphogenesis[1], collective migration[2] and wound healing[3]. Stresses are transmitted through adhesion complexes[4]; either to the substrate generating traction stresses, or to neighbouring cells generating intercellular stresses. Furthermore, these adhesion complexes are important mechanotransduction nodes regulating cellular processes such as metabolism[5], gene transcription[6], as well as actin architecture[7]. Therefore, to gain biological insight linking tissue mechanics to molecular dynamics, there is a need to quantify local traction stresses and intercellular stresses across epithelial monolayers.

Various methodologies are available to interrogate stresses within tissues[8]. Traction force microscopy, where the movement of microbeads embedded within a substrate are tracked, is a proven method for examining cell-generated forces transmitted to the substrate[9]. As a fundamental principle of tissue mechanics is the necessity for monolayers to balance traction and intercellular stresses to maintain epithelial integrity traction force microscopy has been extended to infer intercellular stresses based on mechanical arguments [10–14]. This has been developed for a range of scenarios from cell doublets[14–17], to small cell clusters[13] to entire epithelial sheets[10,12,18,19]. A core assumption of these techniques is that intercellular stresses can be inferred from traction stresses through a passive elastic model, often diagrammatically evoked as a tug-of-war stalemate. This type of model leads to two conclusions: 1) that there is a linear relationship between the scalar sum of traction stresses and intercellular stresses within a monolayer[13,14], and 2) that intercellular stress would be greatest in the centre of the monolayer and lowest at the boundary[20,21]. There are some experimental data based on Förster resonance energy transfer (FRET) tension sensors which support these conclusions. Both E-Cadherin and ZO-2 tension sensors have been reported to decrease in FRET efficiency (increase in molecular tension) in monolayers cultured on stiffer substrates[22,23]. Furthermore, E-Cadherin-α-catenin tension sensors have also been reported to exhibit a reduction in FRET efficiency away from the edge of a wounded epithelium[24]. However, it has also been reported in cell doublets that cell-cell forces inferred from traction force data do not correlate with the FRET-efficiency of an E-Cadherin tension sensor, with the authors proposing a molecular

tension homeostasis mechanism to resolve these findings[16]. Furthermore, experiments

examining intercellular stresses through laser ablation have also made observations which

are not consistent with the conclusions drawn from passive elastic models linking traction

stresses and intercellular stresses.

Laser ablation is a widely employed technique to quantify intercellular stresses

within tissues and through rheological models infer tissue mechanical properties. This is

achieved by examining the relaxation of cells after individual or groups of junctions have

been cut[25,26]. By disrupting entire cell-cell junctions, this method provides a holistic

overview of cell mechanics incorporating both cortical cytoskeleton and all cell-cell adhesion

complexes. It is widely used in developmental systems that are not amenable to other types

of biomechanical techniques to measure or infer tissue mechanics but is also applied to cells

in vitro to examine both intracellular and intercellular stresses. Work in the *Drosophila*

abdominal epidermis showed that degradation of the basal ECM occurring at the pupal

stage led to an increase in intercellular tension[27]. Similarly, Madin-Darby canine kidney

(MDCK) cells cultured on large micropatterned disks on soft hydrogels showed lower

traction stresses but higher recoil velocities than those cultured on stiff hydrogels [28]. These

contradict the conclusions drawn from passive elastic models employed to link traction

stresses to intercellular stresses. In both studies, intercellular tension was greatest when

cell-ECM friction was reduced, in the case of *Drosophila* abdomen the ECM was degraded [27]

and for MDCK cells there was weaker cell-ECM attachments on softer hydrogels[28] which

leads to a reduction in cell-ECM friction[29]. This supports pioneering work in amnioserosa

cells during *Drosophila* embryonic dorsal closure showing that focal adhesions act as tethers

that reduce both intercellular tension and stress transmission[30]. Taken together, these

studies question the use of simple passive elastic models to infer intercellular stresses from

traction stress which do not explicitly model the active cell contractile behaviours, leading

to proposals to replace these types of models with active-matter models[31]. This approach

directly accounts for active contractility within the force balance enabling both active cell-

generated stress and the material stress generated in passive cell structures to be

decoupled, and this approach is now widely adopted in a range of contexts[32–37]. The benefit

of the active matter framework is that it is highly adaptable and capable of capturing a

range of the complexity in the cell-ECM force balance including ECM stiffness-induced

changes. Significantly, cell adhesions and adhesion patterning can be explicitly

described[38,39], which is significant given recent experimental findings that highlight that 1)

the distribution of focal adhesions is important in determining the length-scale that stresses

can be transmitted through the monolayer[30], and 2) that the compliance of the substrate

also influences the distance stresses can be transmitted through the substrate[40]. Indeed, it

is intuitively clear that examining the relationship between intercellular and traction

stresses across a monolayer requires an understanding of how both focal adhesion

distribution and substrate compliance alter the acquired force-balance. This is especially

true when considering that *in vivo* ECM composition, architecture and mechanical

properties are often heterogeneous[41].

To examine the relationship between traction and intercellular stresses we

combined traction force microscopy and annular laser ablation[25]. As traction forces are

known to be dependent on the size and geometry[34,38] of cells and tissues, we employed

micropatterning of hydrogels to standardise these factors. Furthermore, through

micropatterning we modulated cell-ECM interactions by culturing monolayers on disks or

rings to examine the role of focal adhesion distribution in determining monolayers

mechanical equilibrium. We also modulated substrate compliance which is known to alter

both substrate stress transmission[40] and traction stresses generated by cells[38]. When focal

adhesions are distributed throughout the monolayer, traction forces are dependent on the

substrate stiffness within our experimentally tested range. However, when focal adhesion

distribution is mainly restricted to the periphery, traction forces become independent of the

substrate. Furthermore, we observe that intercellular stress is inversely proportional to

substrate stiffness, so that intercellular stress reduces as hydrogel stiffness increases. We

combined this experimental approach with an active stress model based on a continuum

description, which explicitly modelled cell attachments and variation of cell activity within

the layer. This revealed that the relationship between substrate stiffness and intercellular

stress is elicited by the emergence of non-uniform contractility across the monolayer, with

greater contractility present at regions with greater adhesions.

Results

Adhesive surface area dictates focal adhesion distribution across a monolayer

To examine the relationship between traction and intercellular stresses, we cultured
non-tumorigenic mouse mammary epithelial cells (EpH4)[42] on methacrylated hyaluronic
acid (MeHA) hydrogels[43]. 3% w/v MeHA solutions were crosslinked with a range of
dithiothreitol (DTT) crosslinker concentrations to modulate the rigidity of hydrogels from
10.8 ± 2.5 kPa to 31.4 ± 2.8 kPa at intervals of ~5kPa (Fig. S1a-c). Whilst these values are
higher than what has been reported for mouse mammary tissue[44], they span reported
basement membrane values[41] and ranges employed in other systems examining stress
transmission[14,28,29,40]. To examine the role of focal adhesions in determining monolayers
mechanical equilibria, we modulated the adhesive surface area beneath monolayers by
micropatterning laminin, EpH4 cell's preferred ECM component[45]. MeHA hydrogels were
patterned with either disk (Fig. 1a) or ring (Fig. 1b) micropatterns, with an outer diameter of
200μm (approximately 10 – 20 cells wide) and the width of rings being 20μm
(approximately 1-2 cells wide). We conjugated an Alexa633 fluorophore to laminin to assess
if increasing levels of MeHA crosslinking increased the concentration of laminin on the
surface of hydrogels for both disk (Fig. 1c) and ring (Fig. 1d) patterns. Whilst there were
fluctuations in laminin intensity on different hydrogel stiffnesses (Fig. 1c,d) these did not
follow a particular pattern linked to substrate stiffness. Furthermore, whilst there were
radial differences across the disk pattern (Fig. 1c), these did not correlate with differences in
focal adhesions (Fig. 1g). Therefore, the concentration of laminin in patterned areas appears
to not alter the adhesions that cells can generate with the substrate.

To measure active focal adhesions on different patterns, we immunostained
monolayers for phosphorylated focal adhesion kinase (pFAK) (Fig. 1e,f) as previously
reported [add Pere paper]. Altering the adhesive surface did lead to changes in the
distribution of pFAK across the monolayer. On disks, pFAK was evenly distributed across the
entire monolayer with a slight peak at the periphery (Fig. 1g). On rings however, pFAK was
mainly focused on the perimeter of the adhesive area with peaks at the boundaries, and a
low-level present in the non-patterned area (Fig. 1h). As no laminin is present in the middle
of the ring micropatterns (Fig. 1d), the low-level of pFAK staining in this region is probably
due to nascent proteins secreted by EpH4 cells becoming bound to the substrate[46,47] and

not direct binding to the MeHA gel. Indeed, our hyaluronic modification rate was between

76 – 79% (Fig. S1a), far higher than the threshold for cells to bind to unmodified

disaccharides[48]. Furthermore, whilst cells cultured on disks were surrounded by pFAK

puncta (Fig. 1e) across the monolayer, on rings, cells in the centre tended to only have

isolated puncta of pFAK (Fig. 1f). Examining the total pFAK coverage beneath disks and rings

revealed that rings consistently had less total pFAK coverage than disks (Fig. S1d,e),

suggesting that monolayers cultured on ring micropatterns had reduced active cell-ECM

attachments. Surprisingly, given previous reports[29] both patterns showed a slight reduction

in total pFAK as substrate stiffness increased (Fig. S1d,e). The regulation of focal adhesion

dynamics is dependent on many factors[49], and there are contradictory reports on how

substrate rigidity alters focal adhesion coverage[50,51]. It is not clear why total pFAK coverage

reduces on stiffer hydrogels, however, cells have been reported to sense and respond to

strain energy[52], and as we are examining active focal adhesions it is possible that strain

energy is decreasing as stiffness increases in EpH4 cells across the range of rigidities

employed. In summary, we established experimental conditions that allow us to vary cell-

ECM adhesion distribution and substrate stiffness.

We next examined if altering the adhesive pattern or substrate rigidity altered cell-

cell adhesions or cell morphology across the monolayer. We observed no obvious

differences in the localisation of E-Cadherin for both patterns on all hydrogel stiffnesses,

with strong localisation along entire cell boundaries (Fig. 1e,f). Likewise, whilst there was a

distribution in the number of cells that composed each monolayer, the distribution was

similar for all patterns and substrate rigidities (Fig. 1i,j). Examining cell morphology, we

observed that cell height does not significantly change as substrate rigidity increases (Fig.

1k,l). However, we did observe that whilst on disks height is relatively constant (8.9 $\pm$

1.3μm; Fig. 1k), on rings height increases from 8.9 $\pm$ 1.8μm in the centre to 10.4 $\pm$ 1.8μm on

the adhesive area (Fig. 1l). Cell area tended to be slightly greater in the centre of the

patterns, especially for disks (Fig. S1f,g) and there tended to be a slight increase in

elongation at the perimeter for both patterns (Fig. S1h,i). However, these variations were

subtle between the different patterns suggesting that overall monolayer packing was similar

between patterns and on all substrate stiffnesses.

Finally, we examined whether there were junctional rearrangements over the

timescale of our movies. In our timelapse experiments, we imaged for one minute before

performing laser ablations, so subtracted the frame prior to laser ablation from the initial

frame and calculated the percentage pixel intensity of our membrane label that remained.

This analysis revealed a value close to zero on both patterns and on all hydrogel stiffnesses

(Fig. S1j,k). Furthermore, overlayed images of cell membranes for the same duration as our

ablation experiments (5 minutes apart), showed a high degree of overlap (Fig. S1l). We

therefore assume that junctional rearrangement is not a major contributing factor to the

force balance dynamics of the monolayer at the timescale of our experiments.

<u>Focal adhesion patterning alters dependence of traction forces on substrate stiffness</u>

218   We next wanted to assess how altering the adhesive area affected traction forces. To

measure traction forces, we embedded fluorescent microbeads within the MeHA hydrogels

and measured their displacement[53]. Following recent reporting of extensile forces

generated by MDCK cell layers[28], we first confirmed that all displacement vectors indicated

contraction towards the centre of the cell layer across all substrates (Fig. 2a,b, Fig. S2a-d).

We also observed the greatest displacements at the periphery of the monolayer (Fig. 2a,b,

Fig. S2e,f). As expected from previous reports[28,38], we observed a reduction in displacement

magnitude as substrate rigidity increased for both disks and rings (Fig. 2c,d, Fig. S2g-h). We

next examined the traction forces generated by the monolayers and noticed stark

differences between disk and ring patterns (Fig. 2e-h). On disk patterns we observe a

decrease in both the scalar sum of traction forces, which we term total traction, and the

maximum traction force as substrate stiffness increased (Fig. 2e,g). On rings, total traction

force remained almost constant (Fig. 2f), and maximum traction forces showed a weak

reduction (Fig. 2h). It has been reported in micropatterned single MDCK cells that strain

energy is not affected by substrate rigidity[38]. In the case of the monolayers considered here,

as displacement decreased for both patterns and traction forces decreased on disks and

remained constant on rings, we were interested in examining if our adhesive patterns

showed marked differences in strain energy. We observed that for both disks and rings

there is a decrease in strain energy as substrate stiffness increases, potentially explaining

the reduction in pFAK. However, this difference is stronger for disks than it is for rings (Fig.

2i,j), with a mean strain energy reduction between 10kPa to 30kPa for disks (25,242pJ)

being almost double that of rings (13,481pJ).

As the traction forces highlighted a clear difference in behaviour between disk and

rings on different substrates, we decided to examine whether this reflected a difference in

the spatial distribution of traction forces. Examining mean traction forces radially along the

epithelial clusters showed the highest traction forces at the periphery for both ring and disk

patterns on all substrates (Fig. 2k,l). Consistent with changes in total and maximum traction

forces, for disks we observe a reduction in traction magnitude with increasing stiffness

whilst on rings we observe virtually no change in traction force magnitude (Fig. 2k,l).

However, we noticed a slight change in the distribution of traction forces radially between

the patterns, with traction forces appearing to be more localised at the periphery on rings

than in disks especially on soft substrates (Fig. 2k,l). As our pFAK data shows on disks all cells

are surrounded by focal adhesions, this suggests that on compliant substrate, cells within

the monolayer locally contribute towards deforming the substrate, leading to a gradual

radial accumulation in traction forces. However, as the hydrogel becomes more rigid and

cells are less able to deform the substrate, traction forces become more localised to the

edge, as previously theoretically predicted[54]. This is altered on ring patterns where the cells

in the central non-adhesive region do not deform the substrate locally, so that the applied

traction forces are always localised to the periphery. Taken together our results suggest that

altering the adhesive surface leads to changes in the generation and distribution of traction

forces across the epithelium.

Intercellular stresses decrease with increasing substrate stiffness

To interrogate stress transmission within the monolayer, we combined theoretical

and experimental approaches. The theoretical model specifically describes the monolayer as

an active soft material in which the monolayer stresses and displacements are determined

from continuum elasticity theory; specifically, we include an isotropic active stress in the

relationship between stress and strain that accounts for the cell-derived contractility (Fig.

3a, see Supplemental Information Table 1 for model parameters) in contrast to passive

models for traction-force inference. Such active matter models are widely adopted to

describe active, contractile cells and tissues adhered to substrates[17,35,38,54]. Within our

framework, substrate adhesion is coupled into the active model in the force balance

equations through applied tractions, which may be restricted (as here) to act only where the

layer is adhered[54]. Based on our experimental observations with pFAK staining (Fig. 1i,j),

layers were either modelled as adhered uniformly to recapitulate the disk, or for the ring

pattern the model assumes adhesions concentrated at the ring but with an accompanying

underlying distribution of small adhesions sites everywhere else as observed (Fig. 3a). A

more detailed description of the model can be found in supplemental information and

methods section.

As we observe a reduction in total adhesion coverage as substrate stiffness increases

(Fig. S1d,e), we examined in our model the sensitivity of total traction forces to different

levels of adhesion coverage (Supplemental Information). This revealed that traction forces

remained relatively constant with different total levels of adhesion coverage above a 10%

coverage threshold (Supplemental Information Table 2). As the coverage of adhesions is

greater than 10% for all our conditions, we conclude that the reduction in adhesion

coverage we observe does not drastically limit the traction stresses generated by the

monolayer. Indeed, this analysis also shows that our results are insensitive to exact details

of the underlying distribution of adhesions distributed throughout the layer. As the

monolayer was stable at the timescales measured (Fig. S1k-m), we considered the system to

be in a stationary steady state and the forces balanced. The model can be used to fit both

the displacement data (Fig. 3b,c) and total traction (Fig. 3d,e) for each substrate stiffness,

with the cell-generated contractile stress as the fit parameter. In fitting this data, we

assume uniform contractility across the layer so that each cell is equally mechanically active.

Where active matter models are used to describe traction force-type experimental systems

the assumption of constant contractility is common[31–34,55]. Assuming that each cell across

the layer had the same uniform contractility was sufficient to reproduce the observed

traction force microscopy data, with small variations in this overall contractile activity on

each stiffness. As with monolayer stress inference techniques, we then extended this

analysis of traction forces to infer internal stress within the layer, akin to intercellular stress,

which can be determined from the force balance equations. As a reference point, we plot

the internal radial stress halfway across the radius (r=50μm; Fig. 3f,g). As might be

intuitively expected, the model predicts internal stresses that broadly follow the same

qualitative trends observed for total traction across both disks and rings. For the disks, we

predict a significant monotonic decrease as substrate stiffness increases, with the greatest

stresses being on the softer substrates (Fig. 3f). Whilst for the ring patterns, we see that the

stress remains broadly constant across the stiffnesses (Fig. 3g). We next sought to examine

experimentally whether the model's internal stress predictions were accurate by

considering both traction stresses using traction force microscopy and intercellular stresses

inferred from laser ablation.

We bisected the epithelial monolayers with an annular region of interest (ROI) (Fig.

3h) to infer bulk intercellular stresses[25], and analysed the recoil velocity (Fig. S3a,b). We

observed that the greatest recoil was near the ablation site, and that vectors were

orientated away from the cut highlighting that the monolayers were always under tension

(Fig. 3i,j, Fig. S3c-f). Measuring the maximum recoil velocity half-life revealed a similar

relaxation time of ~20 ± 10s for both patterns and all stiffnesses. Assuming that the cells'

Young's modulus is independent of substrate stiffness as previously reported[56], this suggests

that the viscoelastic properties of the monolayer were not altered by focal adhesion

patterning or substrate stiffness (Fig. S3g-j). We then examined the maximum recoil velocity

with respect to substrate stiffness for both patterns. For the disk pattern, the recoil velocity

showed a weak decrease (Fig. 3k) qualitatively similar to the model prediction, but with the

reduction in internal stress being more subtle than that predicted (Fig. 3m). On ring

patterns, while the model predicts a constant recoil velocity, we observed a consistent

decrease with increasing stiffness (Fig. 3l). This reduction, which is not intuitively expected

or predicted by the model, suggests that a model with uniform mechanical activity does not

describe the system fully and that additional factors must be at play.

Comparing traction forces before and after ablation also revealed a reduction,

highlighting mechanical coupling of stresses within the monolayer to the tractions applied

to the substrate (Fig3o,p). Extending this analysis to the strain energy, we also observed a

reduction in strain energy after ablation on both patterns (Fig. 3q,r), with the exception for

disks on the stiffest hydrogel where differences were negligible (Fig. 3q). As total traction

and strain energy vary for the different patterns and substrate stiffnesses, we examined the

percentage change to assess whether this reduction merely reflected the loss of cells in the

ablation region (Fig. 3o'-r'). This revealed that the percentage loss of traction forces and

strain energy (Fig. 3o'-r'), tended to be higher for rings than for disks. Taken together, this

supports the assumption that there is stress transmission from cell-cell interactions which

distributes forces into the substrate, and releasing intercellular stress through ablation leads

to a reduction in traction stresses generated by the monolayer.

<u>Theoretical modelling suggests non-uniformity of contractility</u>

To illustrate the incompatibility of a model with uniform contractility, where all cells

generate equal active forces across the monolayer, with the observed recoil velocities we

focused on the ring pattern which experimentally shows a constant total traction of

~53.7MPa (Fig. 2f) but a reduction in recoil velocity as substrate stiffness increased (Fig. 3l).

For uniform contractility (Fig. 4a) we used our model to examine how the constant

contractility ($P=a$) would have to be altered to maintain a constant internal stress

(represented in blue) or constant total traction force (represented in red) as substrate

stiffness increased. To orientate, we plotted an illustrative contour of constant internal

stress at r=0.5 (which corresponds to the ablation region, blue line) this enables us to

identify a region below the contour (shaded in blue) which has a lower internal stress at

r=0.5 (blue shaded region, Fig. 4a) as required. Likewise, we also plotted an illustrative

contour of constant total traction (red line) and shaded the region above the contour which

has greater traction forces (red shaded region, Fig. 4a). Note that with uniform contractility

the shaded regions do not generate a practically relevant overlap (Fig. 4a), so that we

cannot attain lower internal stress with constant traction, and a parameter sweep altering

the contractility of the layer demonstrates that this is true for all physically relevant regimes

(Supplemental information). This highlights that, as the system maintains a constant

traction force (red contour) it does not encroach into the lower internal stress regime (blue

shaded region). This would inevitably lead to a system whereby changes in traction forces

would lead to similar changes in internal stress as previously predicted by monolayer stress

inference techniques based on passive elastic models (Fig. 3f,g,[10,13,14,18]).

These phase plane plots do suggest a potential resolution by considering a non-

uniform distribution of mechanical contractility within the layer. By increasing contractility

at the edges, the system generates an overlap region where it is possible to maintain total

tractions but reduce internal stress (Fig. 4b). In the examples given, to maintain a constant

internal stress within the layer (blue line) traction forces must increase as the blue contour

now encroaches the shaded orange region, and similarly to maintain constant total traction

forces (red line) the internal stress must decrease as it now encroaches the blue shaded

region (Fig. 4b). As the latter example is similar to our experimental observations, we

explored whether parameterising the mechanical gradient could reconcile theoretical

predictions with experimental observations.

368   We introduced a new contractility function that generates monotonic increasing

369 contractility. Specifically, we take $P(r) = a(1 + br^5)$, as this enables analytical solutions in

370 restricted cases and simplifies the numerical solution (see Supplemental Information). Here

371 increasing b indicates more concentrated contractile activity at the edge ($b=0$ returns the

372 model to uniform cell contraction). By parameterising the total contractility (P_{Total}) of the

373 monolayer and distribution of contractility (b), we were able to fit both tractions and

374 relative changes in internal stress across the range of stiffnesses for both disk (Fig. 4c) and

375 ring patterns (Fig. 4h). Namely we find that the model can reproduce a large decrease in

376 traction and slight decrease in internal layer stress on disk patterns with increasing stiffness

377 (Fig. 4d-f), while also reproducing nearly constant total tractions and large decrease in

378 internal layer stress on the ring pattern as stiffness increases (Fig. 4i-k). Looking beyond the

379 cell traction and recoil data, we considered the strain energy of the system. Non-uniform

380 contractility within the model also reproduces the trends observed in the strain energy of

381 both adhesive patterns (Fig. 4g,l), successfully capturing the observed reduction in strain

382 energy for disks (Fig. 4g) but only qualitatively for ring patterns (Fig. 4l). However,

383 considering that ring patterns have lower adhesion coverage than disks, a deviation in strain

384 energy may be expected given the linear model of substrate resistance to deformation and

385 the contribution of non-local effects from the thickness of the hydrogel (see Supplementary

386 Information). Taken together, introducing non-uniform contractility into the model can

387 recapitulate experimental observations.

388   Significantly, we find a qualitative difference in behaviour between disk and ring

389 patterns for resolving experimental observations. For disk patterns the contractility must

390 become more uniformly distributed, with more cells sharing the work as stiffness increases

391 (Fig. 5a). However, on the ring pattern the opposite trend is required, with cells at the edge

392 of the layer being required to be more contractile compared with the rest of the layer as

393 stiffness increases, so that these edge cells do much of the work (Fig. 5a).

395 <u>Active myosin and F-Actin are not uniformly distributed across monolayers</u>

396   As our model suggested that non-uniformity of contractility could reconcile our

397 experimental observations, we decided to examine contractility within the monolayers. We

398 measured phosphorylated myosin light chain II (pMyoII) intensity across the monolayers as

399 a proxy for contractility[20], but also examined actin architecture which plays an essential role

in cell mechanics[7]. As traction and intercellular stresses are dependent on myosin

activity[20,57], we initially wanted to examine if the total intensity of pMyoII altered depending

on the pattern and substrate stiffness. To minimise experimental variation, we processed all

independent samples in parallel to allow direct comparison in intensity using widefield

imaging for speed. Measuring the intensity of DAPI (Fig. 5b,c), we observed consistent

results with our cell count data (Fig. 1i,j). Examining F-Actin revealed that on disks there is a

consistent level of F-Actin, that appears to be linked to cell number (Fig. 5d). However, on

rings we observe a slight decrease in total F-Actin intensity as substrate stiffness increases

(Fig. 5e). Next, we examined total intensity of pMyoII which for both patterns revealed a

decrease as substrate stiffness increases, but more pronounced for rings (Fig. 5f,g). This

agrees with our fitted model parameters which predicted a reduction in overall contractility

as substrate stiffness increased (Fig. 4c,h).

To examine the spatial distribution of pMyoII, we imaged samples with improved

optical resolution. Examining the total pMyoII intensity of z-sections across our monolayers

we did observe spatial variation in pMyoII to varying degrees on our monolayers (Fig. 5h-k).

For disk patterns, pMyoII intensity appeared more localised at the monolayer edge on soft

hydrogels, becoming more uniform on stiffer hydrogels (Fig. 5h,i). For rings, the opposite

trend in pMyoII intensity distribution was observed, with a more uniform distribution on

soft hydrogels becoming more localised to the edge on stiffer hydrogels (Fig. 5j,k). As

intensity was variable between samples, we normalised pixel intensity to the mean intensity

within the ablation ROI (40-50μm) to quantify the spatial distribution of pMyoII (Fig. 5l-q).

This revealed that no condition had a uniform localisation, with radial variation always

present (Fig. 5l-q), however, there was a redistribution of pMyoII for both patterns as

substrate stiffness increases. On disks there was a reduction in pMyoII intensity outside the

ablation ROI (Fig. 5p), whilst on rings an increase was observed (Fig. 5q). Measuring the

percentage difference in pMyoII intensity between the edge and central regions (Fig. 5r)

revealed a behaviour similar to that predicted by our model (Fig. 5a); as substrate stiffness

increased, on disk patterns there is a trend for pMyoII to become more uniform but on rings

patterns there is a trend to become more localised at the edge.

As there are minor differences in cell height on ring patterns (Fig. 1l), we wanted to

determine if cell height was an issue in this analysis. We therefore examined pMyoII

localisation across the cell height which revealed that it was predominantly localised basally

(Fig. S4a-c). Consequently, we examined actomyosin distribution across the monolayers

focusing only on the basal layer (Fig. S4d-l). Examining pMyoII we observed similar changes

in distribution across the basal layer as we did the whole monolayer (Fig. S4j-l), suggesting

that cell height wasn't confounding our analysis. Interestingly, this analysis revealed drastic

changes in the distribution of the actin cytoskeleton between patterns (Fig. S4d-i) which was

masked when examining total intensity, as F-actin tended to be more localised along cell

junctions (Fig. S4a-c). On disk patterns, cells at the periphery extended lamella outwardly

leading to a dense band of F-Actin at the edge (Fig. S4d,f,h). This was followed by a

reduction in F-Actin behind this band, as cell nuclei aligned close to the edge, with the more

centrally located cells exhibiting clear fibres that spanned the cell width and were

predominantly located near active focal adhesions (Fig. S4m, m'). On ring patterns, an F-

Actin bundle spanning the monolayer perimeter was observed, with a reduction in F-Actin

along the laminin ring, and another accumulation at the inner ring diameter but not as a

defined actin bundle (Fig. S4e,g,i). The architecture of the actin cytoskeleton is critical for

generating[58] and transmitting[17] stresses, and the differences between ring and disk patterns

highlights how altering focal adhesion distribution drastically alters the mechanical

equilibrium of the tissue.

Overall, reconciling the model and experimental observations, we find that the

relationship between traction and intercellular stress is not universal but changes in

response to both the distribution of focal adhesions and compliance of the substrate. This is

due to cellular contractility adapting, to generate a tissue-wide force balance.

Discussion

The control of a tissue's mechanical equilibrium is essential for the regulation of a wide array of physiological processes such as morphogenesis[1] or wound healing[3], and when disturbed can lead to disease progression[59]. Here we investigated the relationship between applied surface traction and intercellular stresses specifically focusing on the role of focal adhesions in the mechanical equilibrium of epithelial monolayers. We find that, for ring patterns where focal adhesion distribution is non-uniform, epithelial monolayer stresses do not conform to the intuition that the relationship between traction and intercellular stresses is linear (Fig. 3m,n). Instead, while traction forces do not change upon increased substrate stiffness (Fig. 2f,h), intercellular tension as measured by laser ablation dramatically drops (Fig. 3l). To reconcile experimental observations with the monolayers being at force-balance, we proposed and confirmed that spatial patterns in contractility emerge (Fig. 4, 5).

What is the origin of this apparently counter-intuitive behaviour? In comparison with single cell studies, for large clusters of cells, forces can be distributed differently between the cells and substrate according to the state of cell-cell junctions[55,60] and the distribution of cell-ECM attachments[16]. In particular, attachment to the ECM has been shown to effectively decrease the length-scale of stress transmission[30], so differences in substrate stiffness can potentially be transmitted to cell-cell junctions over longer distances if cells are not all anchored to the matrix. Conversely, as with the disk patterns where focal adhesions are widely distributed (Fig. 1g), often near cell-cell junctions (Fig 1e) and associated to the contractile acto-myosin network (Fig. S4m), cell-generated forces are most likely transmitted locally to the substrate. In this situation the mechanical properties of the substrate become important for stress transmission across the tissue, as has previously been reported for fibroblasts[61]. Indeed, we found that on disk patterns the monolayer was less able to deform stiffer hydrogels (Fig. 2c), and this was accompanied by a decrease in total traction (Fig. 2e). This is associated with a reduction in the overall total contractility of the layer on stiffer gels (lower pMyoII total intensity, Fig. 5f) as predicted by our theoretical model (Fig. 4c). On soft conditions however, mechanical stress propagates further across the monolayer as cells are more able to deform the substrate locally (Fig. S2e), and higher total contractility is observed (Fig. 5f). This is accompanied with contractile activity differentially localised most strongly to the edge of the layer (fig. 5l,p) where the highest deformations were measured (Fig. S2e).

In contrast, for ring patterns on increasing substrate stiffness, traction stress is

relatively stable (Fig. 2f,h) whilst intercellular tension decreases (Fig. 3l), highlighting that

different mechanical states are reached. For rings, it is only the cells at the edge of the

monolayer that can form focal adhesions and directly experience gel deformability (Fig. 1h).

The bulk of the monolayer can now only contribute predominantly to stress distribution

through cell-cell junctions. Similar to the disk patterns, there is still a decrease in

deformation on stiffer gels (Fig. 2d) and a corresponding decrease in total contractile

activity of the monolayer (lower pMyoII total intensity, Fig. 5g). However, the non-uniform

adhesion distribution generates a marked shift in the collective behaviour of the tissue.

With increasing stiffness, the mechanical activity becomes progressively more localised to

the edge of the monolayer on stiffer gels (Fig. 5r), thus generating lower internal stress as

observed and predicted (Fig. 4k). The stronger resistance to deformation of stiffer gels has

an anchoring effect on the outer ring, but now the balance of resistive forces with the

unadhered cells adjacent to these ring cells is altered. These results emphasise that the

processes contributing to force transmission throughout a monolayer cannot be isolated to

a single cell without considering their collective effects. This is highlighted by changes in F-

Actin architecture across the monolayer between disk and ring patterns (Fig. S4).

Changes in the spatial distribution of actomyosin contractility underpins the

response to differential adhesion on our ring and disk patterns (Fig. 4, 5). Spatial distribution

of contractility has been observed in single cells[62,63] linked to focal adhesion signalling[62],

however for epithelial monolayers, whilst radial patterns of focal adhesion distribution have

been reported[64], to our knowledge this has not been linked to non-uniform patterns of

contractility. Spatial variation in myosin have been observed *in vivo*, but these are normally

examined in the context of morphogenesis[65] or generating cell-population boundaries[66] and

not linked to cell-ECM attachments. We recently observed in the *Drosophila* abdominal

epidermis that a tissue-wide spatial pattern of myosin arises during the pupal stages as the

ECM is degraded[27] but whether this is important for the tissue's mechanical equilibrium is

not known. However, the emergence of spatial patterns of contractility in our monolayers

has important implications on the application of force inference techniques. To examine

intercellular stresses of a monolayer, inference methods such as monolayer force

microscopy (MSM)[10,11,18] or Bayesian inversion stress microscopy (BISM[12]) have been

developed. These techniques infer intercellular stresses based on traction forces measured

and can estimate spatial variation in intercellular stresses in a non-destructive manner,

unlike laser ablation, allowing long-term analysis to be made. The implementation of these

techniques often assumes a passive elastic model linking traction and intercellular

stresses[10,11,13,17,67]. For MDCK cell-doublets, where focal adhesions have been shown to

disassemble along cell-cell contacts[68], the direct balance of traction forces with cell-cell

adhesions potentially holds[14]. However, even in these simple cell-doublet systems the

distribution of focal adhesions can drastically alter the propagation of forces. For instance,

cardiomyocyte doublets seeded or cultured on 90kPa substrates generated focal adhesions

at the site of cell-cell junctions, impairing mechanical coupling and force propagation[15]. This

highlights the importance of examining the distribution of focal adhesions when inferring

stress transmission. Moreover, on our 200μm diameter disk micropatterns, we observed

spatial variation in pFAK with some regions having higher intensity than others (see Fig 1c).

Considering a larger tissue, the distribution of focal adhesions could vary drastically even on

a uniform substrate, altering the local mechanical equilibrium in that region of a tissue. This

would impact whether local intercellular stresses were positively correlated with traction

stress or not. Indeed, analysis of epithelial monolayers showed a "rugged landscape" of

traction forces[10], and spatial patterns of ECM composition and properties have been

observed in developmental and pathological situations[41]. Therefore, when employing

techniques such as MSM and BISM it is important to not only examine the variation in local

distribution of focal adhesions, but also spatial variation in contractility to confidently infer

intercellular stresses as a function of the mechanical activity of the tissue as a whole.

Methods
540
Cell culture

EpH4 cells with Cas9 inserted into the Rosa26 locus (kind gift from high throughput

screening STP, Francis Crick Institute) were cultured in 4.5g l^{-1} Dulbecco's modified Eagle

medium (Gibco, Thermo Fisher Scientific) containing 10% foetal bovine serum (Sigma-

Aldrich, Merck) and 1% penicillin-streptavidin (Gibco, Thermo Fisher Scientific). For seeding

on hydrogels, 180,000 cells in 2ml of culture medium were centrifuged at 80 r.c.f for 5

minutes. Cells were cultured for 2 hours and then extensively washed in PBS to remove non-

adherent cells and 2ml of fresh media was added. Cells were incubated for 48 hours, to

proliferate and cover the micropatterns. For live-imaging experiments, cells were washed in

PBS and then labelled for 15 minutes with 1.5μl CellMask Green (Thermo Fisher Scientific),

diluted and vortexed in 1.5ml FluoroBrite DMEM (Gibco, Thermo Fisher Scientific)

containing 1% penicillin-streptavidin but no FBS. Media was then replaced with 1.5ml

FluoroBrite DMEM media as described, ready for imaging.

Micropatterning of methacrylated hyaluronic acid hydrogels

3g of methacrylated hyaluronic acid (MeHA) was synthesised as previously described[43], with

a methacrylate modification of hyaluronic acid monomer between 76 – 79% as

characterised through ^{1}H NMR (Fig. S1a). MeHA hydrogels were formed at a concentration

of 3 wt% and crosslinked with increasing amounts of dithiothreitol (DTT) to modulate

substrate rigidity. Specifically, hydrogels were formed within a 250μm high silicon gasket

with a 16mm inner diameter and outer diameter of 18mm, sandwiched between a 19mm

micropatterned coverslip and a silanized glass surface. Hydrogels were incubated with 5%

BSA (Tocris Bioscience) in PBS for 2 hours before seeding cells.

Micropatterned coverslips were generated by plasma-activating 18mm coverslips and

coating in 0.1mg ml^{-1} poly-L-lysine poly(ethylene glycol) (PLL(20) – g[3.5]-PEG(2), SuSoS)

suspended in 10mM HEPES buffer at pH7.4. To pattern the coverslips, a quartz photomask

(Compugraphics) and UVO-cleaner (Jelight) were used to remove PLL-g-PEG from the

surface of the coverslip. Patterned coverslips were then incubated for 30 minutes with

laminin diluted in 100mM sodium bicarbonate buffer at pH7.4. Laminin consisted of either

20μg ml^{-1} laminin (Sigma-Aldrich, Merck) solution or 20μg ml^{-1} 2:1 mix of laminin and

laminin-labelled with alexa633 for live-imaging and immunostaining, respectively. To label

laminin, 500μl of 1mg ml^{-1} laminin solution was injected into a Slide-A-Lyzer (Thermo Fisher

Scientific) and dialysed two times in 1L of PBS at 4° C and stirred at a low shear rate for 24

hours. The dialysed laminin was then labelled with Alexa633 following the manufacturers

protocol (Thermo Fisher Scientific). Micropatterns were either repeating patterns of disks

with a diameter of 200μm, or rings with an outer diameter of 200μm and inner diameter of

160μm.

For silanized glass surfaces, either No. 1.5 glass bottom 35mm μ-dishes high (Ibidi) or 20mm

diameter uncoated 6-well plates (Mattek) were employed for live-imaging or

immunostainings, respectively. Glass surfaces were plasma-activated and silanized with (3-

aminopropyl) trimethoxysilane (VWR) and 5% glutaraldehyde (VWR).

Atomic force microscopy

Gels were generated as described but using non-patterned, passivated coverslips and No.

1.5 50mm dish with 30mm uncoated glass diameter (Mattek). Force curves were measured

on a JPK Nanowizard 4 using a CP-qp-CONT-Au-B-5 cantilever probe (sQube, NanoAndMore

GMBH). Measurements were taken in four regions per hydrogel (two at the periphery and

two central) with 50 indentations over a 100μm x 100μm grid. The number of hydrogels

quantified for each DTT concentration was 7 for 6.30 to 15.75mM and 6 for 18.9mM. Force

curves were made with an extend speed of 2μm s^{-1} and a setpoint (8 – 25nN, depending on

substrate stiffness) set to achieve a maximum indentation depth of ~500nm. The young's

modulus was calculated by fitting a Hertz model using the AFM_Youngs_modulus_fit

programme developed by Dr Stefania Marcotti (available on GitHub:

https://github.com/INSIGNEO/AFM_Youngs_modulus_fit), at different indentation depths

(Fig. S1b). A standard indentation depth of 250nm across all hydrogel conditions was used

to calculate the Young's modulus, as it is within the linear phase of the modulus versus

indentation depth for all hydrogel conditions and is 1/1000 of the hydrogel thickness.

Traction force microscopy

Crimson labelled 0.2μm carboxylate modified FluoSpheres (Thermo Fisher Scientific) were

mixed with MeHA hydrogel precursor solution with a 1:40 ratio prior to crosslinking with

DTT. Images were acquired as detailed below. After live-imaging, cells were treated with 10x

trypsin-EDTA, no-phenol red (Gibco) until cells detach from the hydrogels and a final

reference frame acquired.

Laser ablation

Images were acquired on a Zeiss LSM 780 system with an LD C-Apochromat 40x NA 1.1

water objective and 0.8 zoom, so that micropatterns could be acquired in a single 512 x 512

pixel tile. Z-stacks were acquired by measuring excitation by 480nm and 633nm lasers

simultaneously for z-slices every 1μm, spaced between 5 - 6μm below the hydrogel surface

and 10 - 12μm above. Z-stacks were acquired every 10 seconds for one minute before a

laser ablation and five minutes afterwards. An annulus ROI was generated using the

freehand tool and had an internal diameter of 80μm (154 pixels) – approximately 4 – 6 cells

in diameter – and external diameter of 100μm (193 pixels) – leaving 3 – 4 cells at the

periphery. Ablations were made in a z-plane positioned in the middle of the epithelial layer

(3 - 6μm from the hydrogel surface) with a Coherent Chameleon NIR tuneable laser set at

780nm and 40% - 60% power output and lasted ~6.7ms.

Immunostaining

Samples were fixed in 4%-paraformaldehyde (TAAB) for 15 minutes and permeabilised with

0.3% Triton X-100 for 5 minutes, covered from light. Samples were then blocked with 10%

neonatal goat serum (Generon) for an hour at room temperature and incubated with

primary antibodies diluted in 10% neonatal goat serum at 4° C overnight. Samples were

then incubated in secondary antibodies, Phalloidin-Alexa568 (Thermo Fisher Scientific,

#A12380: dilution 1:400) and DAPI (1000x stock solution; Sigma) for 1 hour at room

temperature. Samples were washed and stored in 0.1% Triton X-100. For batch processing

of all samples to examine total pMyoII levels, images were acquired on a Zeiss

Celldiscoverer 7 widefield system with a Zeiss Plan-Apochromat 20x NA 0.7 Autocorr

objective with 1x zoom and 2240 x 2254 pixels and a x-y resolution 0.1747602 for a single

plane. To examine spatial localisation of proteins of interest, samples were imaged on a

Nikon CSU-W1 spinning disk with an Apo LWD 40x WI λS DIC N2 NA 1.15 objective with

1200 x 1200 pixels and a x-y resolution of 0.2731993μm. Z-slices were acquired 0.4μm apart

for each channel sequentially. The following antibodies were used, mouse anti-phospho-

Myosin Light Chain 2 (Ser19) (Cell Signalling Technology, #3675: dilution 1:200), rabbit anti-

phospho-FAK (Tyr397) (Thermo Fisher Scientific, 44-624G: dilution 1:400), mouse anti-E-

Cadherin (BD Transduction #610182: dilution 1:400), Goat anti-mouse IgG H&L Alexa 405

(Abcam, #175660: dilution 1:400), and Goat anti-rabbit IgG H&L Alexa 488 (Thermo Fisher

Scientific, #A-11008: dilution 1:400). F-Actin was stained with Alexa Fluor 568 Phalloidin

(Thermo Fisher Scientific, #A-12380) along with the secondary antibodies.

Data analysis

All images were processed and analysed in Fiji, Mathworks MATLAB and Wolfram

Mathematica.

Traction force microscopy pipeline

Images were acquired as detailed above. After adding trypsin to detach cells, there is slight

stage movement in the in x, y and z directions. Therefore, all movies were registered with

the trypsin frame as a reference. For z registration, the surface of the hydrogel was

calculated by examining the total intensity across the image in 24x24 pixel blocks for each z-

slice, with the surface for each block set by the last slice with a signal. To project on a pixel-

by-pixel basis, a Gaussian Filter was employed across all blocks and then rounded, to

minimise any jumps in z-plane. A maximum projection of the beads was then calculated for

both the movie and trypsin reference frame from the surface to the same number of z-slices

below the surface set by the least number of frames available, normally 3 - 4 slices. Once z-

registration is complete the trypsin frame was added as the final frame of the movie and

registered in Fiji with the StackReg macro, using the last frame as the reference. Bead

displacement, traction forces and strain energy were calculated using the TFM package

developed by Gaudenz Danuser's group[53], using a 31x31 pixel template size and L-

parameterisation of 0.00000001.

Laser ablation analysis

Initially a projection of the cell mask was generated by identifying the ablation plane and

calculating a maximum projection between two slices above and below this frame. The

ablation plane was identified by measuring the loss of intensity within the ablation ROI for

each frame after the ablation and selecting the frame with the biggest loss. To measure the

recoil velocity, the Mathematica ImageFeatureTrack function was employed with specifying

features in every pixel and filtering pixels where no motion was detected. To measure

maximum recoil velocity for each movie, the recoil velocity value for the 95th percentile

across the entire field of view was taken (see Fig. S3).

<u>Cell morphology quantification</u>

To measure cell elongation and area, projections generated for analysing the recoil velocity

were segmented through Skeletor[27] and manually corrected in Fiji. Using a watershed

algorithm in Mathematica, cell area and elongation (1-width/length) were measured. To

measure cell height, all immunostaining experiments were combined, and an orthogonal

projection made using the Fiji macro with a box ROI 20 x 850 pixels centred on the

monolayer centroid in both the x and y direction. In Mathematica, cell height was measured

by combining F-Actin intensity with either pMyoII or E-Cad intensity for the orthogonal

projections, depending on the experiments, and binarizing based on pixels whose intensity

was greater than the mean plus four standard deviations of the entire projected image. A

FillingTransform and DeleteSmallComponents function were applied to clean up the

binarization so only the monolayer remained.

<u>Membrane intensity changes</u>

To measure the movement of the epithelium in our movies before the ablation to

determine their dynamics, the frame immediately before the ablation was subtracted from

the initial frame and the remaining pixel values summed and normalised to the original

summed pixel values. A value close to zero highlights very little movement within the

epithelium.

<u>Basal projection algorithm</u>

To quantify just the basal layer of the monolayers we used Laminin labelling to identify the

surface of the hydrogel. We selected a z-slice for each pixel that corresponded to the

highest laminin intensity, after noise-filtering the z-intensity profile with a LowpassFilter. We

then generated a contour map by smoothening the pixel values with a GaussianFilter,

before rounding to a whole number to correspond to a z-slice. Based on this contour map,

we then did a maximum projection over three z-slices, one above and one below our

contour map value.

<u>pFAK segmentation</u>

To quantify adhesion coverage and size, pFAK images were binarized based on pixel

intensity value in Mathematica. Due to variable background intensity across the field of

view, a MedianFilter was applied with a 61 pixel-by-pixel sliding window and subtracted

from the original image. A background threshold value was then calculated by measuring

the mean intensity plus two standard deviations of pixels outside the micropatterned region

which was determined based on the laminin micropattern channel. Any pixel which had a

higher intensity value than the background threshold was binarized. To de-noise the binary

image, isolated binarized pixels were removed using the DeleteSmallComponents function

with a value set to 1.

<u>Statistical analysis</u>

All statistics were performed in Mathematica using the LocationEquivalenceTest function

which tests the null hypothesis that populations are equal choosing the most powerful test

that applies to the data. The most common test applied was the Kruskal Wallis test as often

one condition would fail normality tests. If the LocationEquivalenceTest was significant a

post-test Mann-Whitney was performed comparing the value on the softest substrate to all

other substrate stiffnesses. For trend fitting, the LinearModelFit function was used on the

mean values for each population with a confidence interval of 95%. The model with the

greatest R^2-value was selected.

For all statistical values used in the analysis, see Tables S1&2 in Supplementary Information.

Theoretical modelling

We describe the theoretical and computational modelling briefly here, for a more detailed

description see the additional supplementary information.

Active matter model for the cell layer

730   An active matter model was used to predict the deformations and stresses in the cell layer.

731   This widely adopted approach incorporates an active cell-generated stress additively into

732   the constitutive relation of the cell material, see[36,37,39,54]. As such we assumed a constitutive

733   relation such that the stress $s = s_{\text{passive elastic}} + s_{\text{active}}$, as Fig. 3a specifically in tensor notation

734   this gives:

$$\sigma_{ij} = \frac{hE}{1+\nu}\left(\epsilon_{ij} + \frac{\nu}{1-\nu}\epsilon_{kk}\delta_{ij}\right) + \frac{hE}{2(1-\nu)}P(\mathbf{x})\delta_{ij},$$

736   where we use the summation convention and introduce the kronecker delta. The active contractility

737     $P(x)$ is assumed to act isotropically. The deformations are then determined from the force balance

$$\nabla \cdot \sigma - KT(\mathbf{x})\mathbf{u} = \mathbf{0},$$

Where K quantifies substrate resistance to deformation and $T(\mathbf{x})$ is an indicator function

that take the value of 1 where the cell is adhered and 0 otherwise. The contractility function

is assumed to be constant throughout the layer $P(\mathbf{x})=P_a$, for Fig. 3, with the addition of a

monotonic increasing contractility profile $P(\mathbf{x})=a(1+br^5)$ for Fig. 4. The polynomial

contractility profile is selected to enable analytical phase plane analysis. Where adhesion is

uniform analytical solutions of the force balance equation may be obtained by standard

techniques as the resulting equations are of Bessel type and solutions may be obtained by

variation of parameters. Where non-uniform adhesion is observed numerical solutions of

the force balance equation are obtained using finite element methods within the MATLAB

PDE Toolbox.

Adhesion geometry

For uniform adhesion the indicator function is either taken as unity throughout for the disk

geometry or as unity in the ring region. Where adhesion is informed by the pFAK data (Fig.

1) and modelled as non-uniform we take the resulting adhesion geometry is as in Fig. 1h.

Specifically, we consider two thin annuli of thickness r_t=0.06 at the edges of the

micropatterned ring corresponding to the regions with highest intensity pFAK. Additionally,

we set a distribution of adhered spots in the internal region and between the two annuli as

the pFAK intensity is non-zero in these regions; spots between the two annuli have a radius

of r_s=0.0167r_0 and in the internal region have radius r_s=0.018r_0. In these adhered regions

T(x)=1 and elsewhere we set T(x)=0.

Acknowledgements

We would like to thank the Ok-Ryul Song and Michael Howell from the High-throughput

screening STP at the Francis Crick Institute for sharing the EpH4-Cas9 cells. We are grateful

to Matt Renshaw from the Crick CALM STP, and Ravi Desai and Christina Dix from the Crick

Making STP for their assistance. We thank the bioAFM and bioImaging facilities within

FBMH at the University of Manchester. We are grateful to Stefania Marcotti, Richard

Thorogood, Nigel Hodson, Olivia Courbot and Alberto Elosegui-Artola for advice on the

analysis of AFM data. We thank Guillaume Charras and Alberto Elosegui-Artola for

comments on the manuscript. JRD was supported by a Wellcome Trust Sir Henry Wellcome

fellowship (201358/Z/16/Z) and University of Manchester Dean's Prize. JSW was supported

by postdoctoral EPSRC funding (EP/W522302/1). This work was supported by a Wellcome

Trust Investigator award (107885/Z/15/Z) to NT. Work in the Tapon lab is supported by the

Francis Crick Institute, which receives its core funding from Cancer Research UK (CC2138),

the UK Medical Research Council (CC2138), and the Wellcome Trust (CC2138). For Open

Access, the authors have applied a CC BY public copyright licence to any Author Accepted

Manuscript version arising from this submission.

1.   Clarke, D. N. & Martin, A. C. Actin-based force generation and cell adhesion in tissue
morphogenesis. *Current Biology* **31**, R667–R680 (2021).
2.   Pandya, P., Orgaz, J. L. & Sanz-Moreno, V. Actomyosin contractility and collective
migration: may the force be with you. *Curr Opin Cell Biol* **48**, 87–96 (2017).
3.   Lim, S. E., Vicente-Munuera, P. & Mao, Y. Forced back into shape: Mechanics of
epithelial wound repair. *Curr Opin Cell Biol* **87**, 102324 (2024).
4.   Mao, Y. & Baum, B. Tug of war-The influence of opposing physical forces on epithelial
cell morphology. *Dev Biol* **401**, 92–102 (2015).
5.   Romani, P., Valcarcel-Jimenez, L., Frezza, C. & Dupont, S. Crosstalk between
mechanotransduction and metabolism. *Nat Rev Mol Cell Biol* (2020)
doi:10.1038/s41580-020-00306-w.
6.   Dupont, S. & Wickström, S. A. Mechanical regulation of chromatin and transcription.
Nature Reviews Genetics vol. 23 624–643 Preprint at https://doi.org/10.1038/s41576-
022-00493-6 (2022).
7.   Sun, X. & Alushin, G. M. Cellular force-sensing through actin filaments. *FEBS Journal*
290, 2576–2589 (2023).
8.   Gómez-González, M., Latorre, E., Arroyo, M. & Trepat, X. Measuring mechanical
stress in living tissues. *Nature Reviews Physics* **2**, 300–317 (2020).
9.   Lekka, M., Gnanachandran, K., Kubiak, A., Zieliński, T. & Zemła, J. Traction force
microscopy – Measuring the forces exerted by cells. *Micron* **150**, (2021).
10.   Tambe, D. T. *et al.* Collective cell guidance by cooperative intercellular forces. *Nat
Mater* **10**, 469–75 (2011).
11.   Tambe, D. T. *et al.* Monolayer Stress Microscopy: Limitations, Artifacts, and Accuracy
of Recovered Intercellular Stresses. *PLoS One* **8**, e55172 (2013).
12.   Nier, V. *et al.* Inference of Internal Stress in a Cell Monolayer. *Biophys J* **110**, 1625–
1635 (2016).
13.   Ng, M. R., Besser, A., Brugge, J. S. & Danuser, G. Mapping the dynamics of force
transduction at cell-cell junctions of epithelial clusters. *Elife* **3**, e03282 (2014).
14.   Maruthamuthu, V., Sabass, B., Schwarz, U. S. & Gardel, M. L. Cell-ECM traction force
modulates endogenous tension at cell-cell contacts. *Proceedings of the National
Academy of Sciences* **108**, 4708–4713 (2011).
15.   McCain, M. L., Lee, H., Aratyn-Schaus, Y., Kléber, A. G. & Parker, K. K. Cooperative
coupling of cell-matrix and cell-cell adhesions in cardiac muscle. *Proc Natl Acad Sci U
S A* **109**, 9881–9886 (2012).
16.   Sim, J. Y. *et al.* Spatial distribution of cell-cell and cell-ECM adhesions regulates force
balance while main taining E-cadherin molecular tension in cell pairs. *Mol Biol Cell* **26**,
2456–2465 (2015).
17.   Ruppel, A. *et al.* Force propagation between epithelial cells depends on active
coupling and mechano-structural polarization. *Elife* **12**, 1–42 (2023).
18.   Trepat, X. *et al.* Physical forces during collective cell migration. *Nat Phys* **5**, 426–430
(2009).
19.   Balasubramaniam, L. *et al.* Investigating the nature of active forces in tissues reveals
how contractile cells can form extensile monolayers. *Nat Mater* **20**, 1156–1166
(2021).
20.   Vincent, R. *et al.* Active Tensile Modulus of an Epithelial Monolayer. *Phys Rev Lett*
115, (2015).

21. Serra-Picamal, X. *et al.* Mechanical waves during tissue expansion. *Nat Phys* **8**, 628–
634 (2012).

22. Haas, A. J. *et al.* Interplay between Extracellular Matrix Stiffness and JAM-A Regulates
Mechanical Load on ZO-1 and Tight Junction Assembly. *Cell Rep* **32**, 107924 (2020).

23. Mohan, A. *et al.* Spatial Proliferation of Epithelial Cells Is Regulated by E-Cadherin
Force. *Biophys J* **115**, 853–864 (2018).

24. Gayrard, C., Bernaudin, C., Déjardin, T., Seiler, C. & Borghi, N. Src- and confinement-
dependent FAK activation causes E-cadherin relaxation and β-catenin activity. *Journal*
of Cell Biology **217**, 1063–1077 (2018).

25. Bonnet, I. *et al.* Mechanical state, material properties and continuous description of
an epithelial tissue. *J R Soc Interface* **9**, 2614–2623 (2012).

26. Bambardekar, K., Clément, R., Blanc, O., Chardès, C. & Lenne, P.-F. Direct laser
manipulation reveals the mechanics of cell contacts in vivo. *Proceedings of the*
National Academy of Sciences **112**, 1416–1421 (2015).

27. Davis, J. R. *et al.* ECM degradation in the Drosophila abdominal epidermis initiates
tissue growth that ceases with rapid cell-cycle exit. *Current Biology* **32**, 1285-1300.e4
(2022).

28. Sonam, S. *et al.* Mechanical stress driven by rigidity sensing governs epithelial
stability. *Nat Phys* **19**, 132–141 (2023).

29. Vazquez, K., Saraswathibhatla, A. & Notbohm, J. Effect of substrate stiffness on
friction in collective cell migration. *Sci Rep* **12**, (2022).

30. Goodwin, K. *et al.* Basal Cell-Extracellular Matrix Adhesion Regulates Force
Transmission during Tissue Morphogenesis. *Dev Cell* **39**, 611–625 (2016).

31. Green, Y., Fredberg, J. J. & Butler, J. P. Relationship between velocities, tractions, and
intercellular stresses in the migrating epithelial monolayer. *Phys Rev E* **101**, (2020).

32. Friedrich, B. M. & Safran, S. A. How cells feel their substrate: Spontaneous symmetry
breaking of active surface stresses. *Soft Matter* **8**, 3223–3230 (2012).

33. Yang, W., Luo, M., Gao, Y., Feng, X. & Chen, J. Mechanosensing model of fibroblast
cells adhered on a substrate with varying stiffness and thickness. *J Mech Phys Solids*
171, (2023).

34. Mertz, A. F. *et al.* Scaling of Traction Forces with the Size of Cohesive Cell Colonies.
Phys Rev Lett **108**, 198101 (2012).

35. Banerjee, S. & Marchetti, M. C. Contractile Stresses in Cohesive Cell Layers on Finite-
Thickness Substrates. *Phys Rev Lett* **109**, 108101 (2012).

36. Edwards, C. M. & Schwarz, U. S. Force localization in contracting cell layers. *Phys Rev*
Lett **107**, 1–5 (2011).

37. Prost, J., Jülicher, F. & Joanny, J. F. Active gel physics. *Nat Phys* **11**, 111–117 (2015).

38. Oakes, P. W., Banerjee, S., Marchetti, M. C. & Gardel, M. L. Geometry regulates
traction stresses in adherent cells. *Biophys J* **107**, 825–833 (2014).

39. Solowiej-Wedderburn, J. & Dunlop, C. M. Sticking around: Cell adhesion patterning
for energy minimization and substrate mechanosensing. *Biophys J* **121**, 1777–1786
(2022).

40. Reinhart-King, C. A., Dembo, M. & Hammer, D. A. Cell-cell mechanical communication
through compliant substrates. *Biophys J* **95**, 6044–6051 (2008).

41. Kozyrina, A. N., Piskova, T. & Di Russo, J. Mechanobiology of Epithelia From the
Perspective of Extracellular Matrix Heterogeneity. *Front Bioeng Biotechnol* **8**, (2020).

42. Reichmann, E., Ball, R., Groner, B. & Friis, R. R. New mammary epithelial and
fibroblastic cell clones in coculture form structures competent to differentiate
functionally. *J Cell Biol* **108**, 1127–1138 (1989).

43. Guvendiren, M. & Burdick, J. A. Stiffening hydrogels to probe short- and long-term
cellular responses to dynamic mechanics. *Nat Commun* **3**, 792 (2012).

44. Lopez, J. I., Kang, I., You, W. K., McDonald, D. M. & Weaver, V. M. In situ force
mapping of mammary gland transformation. *Integrative Biology* **3**, 910–921 (2011).

45. Maschler, S. M. *et al.* Tumor cell invasiveness correlates with changes in integrin
expression and localization. *Oncogene* **24**, 2032–2041 (2005).

46. Loebel, C., Mauck, R. L. & Burdick, J. A. Local nascent protein deposition and
remodelling guide mesenchymal stromal cell mechanosensing and fate in three-
dimensional hydrogels. *Nat Mater* **18**, 883–891 (2019).

47. Loebel, C. *et al.* Metabolic labeling of secreted matrix to investigate cell–material
interactions in tissue engineering and mechanobiology. *Nature Protocols* vol. 17 618–
648 Preprint at https://doi.org/10.1038/s41596-021-00652-9 (2022).

48. Kwon, M. Y. *et al.* Influence of hyaluronic acid modification on CD44 binding towards
the design of hydrogel biomaterials. *Biomaterials* **222**, 119451 (2019).

49. Kechagia, J. Z., Ivaska, J. & Roca-Cusachs, P. Integrins as biomechanical sensors of the
microenvironment. *Nat Rev Mol Cell Biol* **20**, 457–473 (2019).

50. Han, S. J., Bielawski, K. S., Ting, L. H., Rodriguez, M. L. & Sniadecki, N. J. Decoupling
substrate stiffness, spread area, and micropost density: A close spatial relationship
between traction forces and focal adhesions. *Biophys J* **103**, 640–648 (2012).

51. Fusco, S., Panzetta, V. & Netti, P. A. Mechanosensing of substrate stiffness regulates
focal adhesions dynamics in cell. *Meccanica* **52**, 3389–3398 (2017).

52. Panzetta, V., Fusco, S. & Netti, P. A. Cell mechanosensing is regulated by substrate
strain energy rather than stiffness. *Proc Natl Acad Sci U S A* **116**, 22004–22013 (2019).

53. Han, S. J., Oak, Y., Groisman, A. & Danuser, G. Traction microscopy to identify force
modulation in subresolution adhesions. *Nat Methods* **12**, 653–656 (2015).

54. Solowiej-Wedderburn, J. & Dunlop, C. M. Cell-strain-energy costs of active control of
contractility. *Phys Rev E* **107**, (2023).

55. Mertz, A. F. *et al.* Cadherin-based intercellular adhesions organize epithelial cell-
matrix traction forces. *Proc Natl Acad Sci U S A* **110**, 842–847 (2013).

56. Rheinlaender, J. *et al.* Cortical cell stiffness is independent of substrate mechanics.
Nat Mater **19**, 1019–1025 (2020).

57. Valon, L., Marín-Llauradó, A., Wyatt, T., Charras, G. & Trepat, X. Optogenetic control
of cellular forces and mechanotransduction. *Nat Commun* **8**, (2017).

58. Doss, B. L. *et al.* Cell response to substrate rigidity is regulated by active and passive
cytoskeletal stress. *Proc Natl Acad Sci U S A* **117**, 12817–12825 (2020).

59. Ayad, N. M. E., Kaushik, S. & Weaver, V. M. Tissue mechanics, an important regulator
of development and disease. *Philosophical Transactions of the Royal Society B:
Biological Sciences* vol. 374 Preprint at https://doi.org/10.1098/rstb.2018.0215
(2019).

60. Bazellières, E. *et al.* Control of cell-cell forces and collective cell dynamics by the
intercellular adhesome. *Nat Cell Biol* **17**, 409–420 (2015).

61. Palmquist, K. H. *et al.* Reciprocal cell-ECM dynamics generate supracellular fluidity
underlying spontaneous follicle patterning. *Cell* **185**, 1960-1973.e11 (2022).

62.   De Mets, R. *et al.* Cellular tension encodes local Src-dependent differential β1 and β3
916   integrin mobility. *Mol Biol Cell* **30**, 181–190 (2019).

917   63.   Von Erlach, T. C., Hedegaard, M. A. B. & Stevens, M. M. High resolution Raman
918   spectroscopy mapping of stem cell micropatterns. *Analyst* **140**, 1798–1803 (2015).

919   64.   Pérez-González, C. *et al.* Active wetting of epithelial tissues. *Nat Phys* **15**, 79–88
920   (2019).

921   65.   Wang, Y. C. The origin and the mechanism of mechanical polarity during epithelial
922   folding. *Seminars in Cell and Developmental Biology* vol. 120 94–107 Preprint at
923   https://doi.org/10.1016/j.semcdb.2021.05.027 (2021).

924   66.   Sharrock, T. E. & Sanson, B. Cell sorting and morphogenesis in early Drosophila
925   embryos. *Semin Cell Dev Biol* **107**, 147–160 (2020).

926   67.   Chen, T., Saw, T. B., Mege, R. M. & Ladoux, B. Mechanical forces in cell monolayers. *J
927   Cell Sci* **131**, (2018).

928   68.   Yamada, S. & Nelson, W. J. Localized zones of Rho and Rac activities drive initiation
929   and expansion of epithelial cell-cell adhesion. *Journal of Cell Biology* **178**, 517–527
930   (2007).

Figure 1

Fig. 1. Modulating adhesive surface of EpH4 monolayers. a,b Example projection of laminin

micropatterns for disk (**a**) and ring (**b**). **c,d** Quantification of mean laminin intensity radially

across micropatterns. **e,f** Example projections of basal pFAK and apical E-Cadherin on disk

(**e**) and ring (**f**) micropatterns. **g,h** Quantification of mean pFAK coverage radially across

micropatterns. **i,j** Dot plot of cell number within each monolayer, bars highlight mean value

for each population. **k,l** Quantification of mean cell height radially across micropatterns. For

all quantification thick line are mean values and ribbons are S.E.M, grey shaded region

represents adhesive surface. All N numbers and statistics included in Table S1 in

Supplementary Information. All scale bars = 20μm.

Figure 2

Fig. 2. Monolayer traction stresses are altered by modulating adhesive surface. a,b Mean

displacement vector plots for monolayers on disks (**a**) and rings (**b**) on different substrate

stiffnesses. Red dashed line outlines the adhesive surface and scale bars represent 4μm. **c,d**

Quantification of maximum bead displacement by monolayers adhered to disks and rings on

different substrate stiffnesses. **e,f** Quantification of total traction forces generated by

monolayers adhered to disks and rings on different substrate stiffnesses. **g,h** Quantification

of maximum traction forces generated by monolayers adhered to disks and rings on

different substrate stiffnesses. **i,j** Quantification of strain energy transferred by monolayers

into substrate adhered to disks and rings on different substrate stiffnesses. For **c-j** black dots

highlight mean value, whiskers represent S.E.M and grey line is line of best fit. **k,l**

Quantification of mean traction forces radially across micropatterns. Thick lines are mean

values and ribbons are S.E.M, grey shaded region represents adhesive surface.

Figure 3

Fig. 3. Intercellular stress is not proportional to traction stress. a Schematic representation

of the major components that comprise the force balance model. Cells are considered as

active contractile agents (red arrows) bound together to form a monolayer continuum as

well as being bound to the substrate through adhesions (blue springs). Any stresses within

the monolayer are therefore a product of these active forces and passive elastic forces

transmitted through the monolayer (top equation). The active forces generated by the cells

are modelled as a pressure term (red equation) describing cell contractility, and for the

initial model we assumed a uniform distribution of contractility. Cell adhesions are

distributed throughout the monolayer based on experimental observations (blue

schematic), and at these points stresses within the monolayer are transmitted to the

substrate to deform it (bottom two equations). Schematic produced in part with

Biorender.com. **b,c** Total hydrogel displacement from experiments (same as Fig. S2g,h) were

used to parameterise the model. **d,e** Based on this parameterisation the model can

recapitulate the overall trend in experimental traction forces (same as Fig. 2e,f). **f,g** Model

predictions on intercellular stresses at the radial midpoint of the layer (r=0.5) normalised to

E=10kPa show changes as substrate stiffness increases. The model predicts intercellular

forces change in a manner similar to the total traction forces as substrate stiffness

increases. **h** Example time course projections of our laser ablation procedure for both

micropatterns. A ring ROI with outer radius of 50μm and inner radius of 40μm was ablated.

Merge shows overlay of 0s before and 60s after the ablation only to highlight deformations,

where white highlights regions of no deformation. Scale bar = 20μm. **i,j** Mean recoil velocity

vector plots for monolayers adhered on disks (**i**) and rings (**j**) on different substrate

stiffnesses. Red dashed line outlines the adhesive surface and scale bars represent

3.5μm/min. **k,l** Quantification of maximum recoil velocity after ablation for monolayers

adhered to disks and rings on different substrate stiffnesses. **m,n** Recoil velocity (same as

Fig. 3k,l) and model predictions of radial midpoint intercellular stresses (same as Fig. 3f,g)

have both been normalised to the value on the softest hydrogel (E=10kPa). The model is not

able to mimic experimental observations. **o-r** Quantification of the total loss in traction

forces (**o,p**) and strain energy (**q,r**) after ablations and once a new steady state is reached,

for monolayers adhered to disks (**o,q**) and rings (**p,r**) on different substrate stiffnesses.

Where **o'-r'** represent this value as a percentage of total traction forces (see Fig. 2e,f, **o',p'**)

and strain energy (see Fig. 2i,j, **q',r'**). For all quantification graphs, experimental data is in

black, with dots representing mean value, whiskers the S.E.M and grey line the line of best

fit. Model outputs are red circles, where included.

Figure 4

Fig. 4. Non-uniform contractility reconciles experimental observations. a,b Phase plane

plots of contractility in the centre of the monolayer against substrate stiffness for ring

monolayers only. The examples showcase situations where the spatial distribution of

contractility (b), represented with accompanying heatmaps, is either uniform (b =0, **a**) or

non-uniform (b =1, **b**). In both, solid lines represent situations where either intercellular

stress at the radial midpoint of the layer (r=0.5) is constant (blue lines) or total traction

forces generated by the monolayer is constant (red line). The shaded regions represent

either situations where the intercellular stress must be lower (blue shaded region) or total

traction forces must be greater (red shaded area). Note that on uniform contractility there is

no workable overlap of shaded regions meaning no workable parameters exist that allow

constant traction forces but lower intercellular stresses as substrate stiffness increases as

occurs experimentally. However, with non-uniform contractility an overlap region does

exist, highlighting a large parameter space which could reconcile experimental observations.

For **c-l** the model was parameterised to include non-uniform contractility for both disks (**c-g**)

and rings (**h-l**). **c,h** Schematic representations of the total and spatial distribution of

contractility parameters (b = spatial distribution of contractility; P = total contractility

normalised to softest substrate) used for disk (**c**) and rings (**h**) on different substrate

stiffnesses. **d,i** Total hydrogel displacement from experiments (same as Fig. S2g,h), **e,j** total

traction forces (same as Fig. 2e,f), and **f,k** normalised intercellular stress at the radial

midpoint of the layer (r=0.5) normalised to the softest substrate (same as Fig. 3m,n) were

used to parameterise the model. **g,l** Normalised strain energy to the softest substrate was

used as a final check to examine the power of the model to mimic all experimental

observations. Note that for disks (**g**) the model can reproducibly mimic experimental

observations but for rings (**l**) it mimics the trend but does not reproduce the actual

magnitude of change. For all quantification graphs, experimental data is in black, with dots

representing mean value, whiskers the S.E.M and grey line the line of best fit and model

outputs are red triangles.

1021

1022

Figure 5

Fig. 5. Contractility is non-uniform across micropatterned monolayers. a Schematic

representation of predictions based on parameterisation of the model. For disks the model

suggests that there needs to be a shift in contractility from focused to the edge on soft

substrates to more uniformity on stiffer substrates. On rings, the model suggests the

opposite needs to occur with contractility being more uniform on soft substrates and

becoming more localised to the edge on stiff substrates. **b-g** Dot plots of total intensity from

widefield images within each monolayer for DAPI (**b,c**), F-Actin (**d,e**) and pMyoII (**f,g**)

adhered to disks (**b,d,f**) and rings (**c,e,g**). Each dot represents one monolayer, and the bars

represent mean value for each population. **h-k** Example projections of total monolayer

intensity taken from confocal sections of pMyoII and F-Actin adhered to disk (**h,i**) and ring

(**j,k**) micropatterns on soft (**h,j**) and stiff (**i,k**) hydrogels. Scale bar = 20µm. **l-o** Heatmaps of

mean total monolayer pMyoII intensity normalised to the ablation ROI (40-50µm) for

monolayers adhered to disks (**l,m**) and rings (**n,o**) on soft (**l,n**) and stiff (**m,o**) hydrogels. **p,q**

Quantification of radial mean total pMyoII intensity normalised to the ablation ROI (40-

50µm) for monolayers adhered to disks (**p**) and rings (**q**) on different substrate stiffnesses. **r**

Quantification of percentage difference in normalised total monolayer pMyoII intensity

between edge (>60µm) and centre (<40µm) radii. Dots represent mean value, whiskers

S.E.M and shaded line the line of best fit, with disks in red and rings in black. Note that the

trend corroborates the model predictions (**a**).

1042

Supplemental Figure 1

Fig. S1. Characterisation of micropatterned MeHA on EpH4 cells. a ^{1}H NMR spectra of methacrylated hyaluronic acid (MeHA). **b** Young's moduli at different indentation depths on 3%(w/v) MeHA hydrogels cross-linked with different DTT concentrations. Dashed line is at 250nm which was used to define the Young's moduli for each DTT concentration. **c** Dot plot of Young's moduli for each concentration of DTT. Each dot is the average value from 50 force-curves taken for a 100μm x 100μm region. **d,e** Dot plot of total pFAK coverage within each monolayer, bars highlight mean value for each population. **f,g** Quantification of mean cell area radially across micropatterns. **h,i** Quantification of mean cell elongation radially across micropatterns. For **f-i** thick line are mean values and ribbons are S.E.M, grey shaded region represents adhesive surface. **j,k** Dot plot of % intensity change after subtracting the cell outline projection from the frame before an annular ablation from the initial frame 1 minute before. Lines represent mean value for each population. **l** Cell outline projections five minutes apart overlayed with the initial frame in green and last frame in Magenta. White represents pixels where there is no change in intensity. All N numbers and statistics included in Table S2 in Supplementary Information. All scale bars = 20μm.

Supplemental Figure 2

Fig. S2. Displacements of microbeads embedded within MeHA hydrogels by EpH4

monolayers. a-d Quantification of mean displacement along the x-axis (**a,b**) and y-axis (**c,d**).

Grey region represents adhesive surface. **e,f** Quantification of mean displacement radially

across micropatterns, where thick lines are mean values, ribbons are S.E.M, and grey

shaded region represents adhesive surface. **g,h** Quantification of total bead displacements

generated by monolayers adhered to disks and rings on different substrate stiffnesses. Dots

represent mean values, whiskers are S.E.M, and grey line is line of best fit. **i,j** Quantification

of the mean bead displacement around the perimeter of monolayers, with 0° along the

horizontal axis, where thick lines are mean values, ribbons are S.E.M, and grey shaded

region represents adhesive surface.

1070

1071

Supplemental Figure 3

Fig. S3. Recoil velocity dynamics and correlations after laser ablation. a,b Cumulative plots

of all recoil velocity values for each movie on both micropatterns. To calculate the maximum

recoil value the 95th percentile (grey dashed line) value was used, which was always located

near the edge of the ablation site. **c-f** Quantification of mean recoil velocity along the x-axis

(**c,d**) and y-axis (**e,f**). Grey region represents adhesive surface. **g,h** Quantification of the

mean maximum recoil velocity over the time course of our movies for both micropatterns.

i,j Dot plot of recoil half-life calculated from Fig. S3g,h. Each dot represents a movie, and the

line represents population mean value.

1080

1081

Supplemental Figure 4

Fig. S4. pMyoII localisation within monolayers. a Example orthogonal projections through the middle of a monolayer of pMyoII, F-Actin and Laminin on disk (left) and ring (right) micropatterns. **b,c** Quantification of mean orthogonal intensity of laminin, pMyoII and F-Actin normalised to the maximum value for the entire orthogonal projection (see **a**). Note that pMyoII maximum intensity is always near the laminin intensity highlighting basal accumulation of pMyoII, whilst F-actin peak is close to 50% of cell height where cell-cell junctions are located. **d,e** Example projections of basal monolayer intensity taken from confocal sections of pMyoII and F-Actin adhered to disk (**d**) and ring (**e**) micropatterns on 20kPa hydrogels. **f,g** Heatmaps of mean F-Actin intensity from basal projections normalised to the ablation ROI (40-50µm) for monolayers adhered to disks (**f**) and rings (**g**) on 20kPa hydrogels. **h,i** Quantification of radial mean F-Actin intensity from basal projections normalised to the ablation ROI (40-50µm) for monolayers adhered to disks (**h**) and rings (**i**) on different substrate stiffnesses. **j,k** Quantification of radial mean pMyoII intensity from basal projections normalised to the ablation ROI (40-50µm) for monolayers adhered to disks (**j**) and rings (**k**) on different substrate stiffnesses. For **h-k**, thick lines are mean values, ribbons are S.E.M, and grey shaded region represents adhesive surface. **l** Quantification of percentage difference in normalised pMyoII intensity from basal projections between edge (>60µm) and centre (<40µm) radii. Dots represent mean value, whiskers S.E.M and shaded line the line of best fit, with disks in red and rings in black. Note a similar trend to total projections seen in Fig. 5r). **m** Example basal projection of pFAK, F-Actin and E-Cadherin on disk micropattern with expanded insert (**m'**) to highlight F-Actin fibres that span epithelial cells, mainly between pFAK labelled adhesions. Scale bar = 20µm.