Gaze inversion effect: Sex invariance and difference in gaze direction sensitivity

Daisuke Matsuyoshi [1,2,3], Kana Kuraguchi [4,5], Hiroshi Ashida [5], Katsumi Watanabe [1]

[1] Faculty of Science and Engineering, Waseda University, 3-4-1 Ohkubo, Shinjuku, Tokyo 169-8555, Japan

[2] Araya Inc., 1-11 Kanda-sakumacho, Chiyoda, Tokyo 101-0025, Japan

[3] Advanced Neuroimaging Center, Institute for Quantum Medical Science, National Institutes for Quantum Science and Technology, 4-9-1 Anagawa, Inage, Chiba 263-8555, Japan

[4] Faculty of Social Sciences, Shitennoji University, 3-2-1 Gakuenmae, Habikino, Osaka 583-0868, Japan

[5] Department of Psychology, Graduate School of Letters, Kyoto University, Yoshida-honmachi, Sakyo, Kyoto 606-8501, Japan

Corresponding Author:

Daisuke MATSUYOSHI

Advanced Neuroimaging Center, Institute for Quantum Medical Science

National Institutes for Quantum Science and Technology

Phone: +81-43-206-3249   E-mail: matsuyoshi.daisuke@qst.go.jp

Abstract

Even though humans are adept at distinguishing various faces, inversion dramatically

spoils the ability. This face inversion effect is remarkably robust across observers;

however, whether inversion impairs the perception of facial parts (particularly the eye) is

inconsistent across studies. Some studies have shown that featural processing remains

preserved or even enhanced when faces are inverted, while others have shown that

inversion impairs featural processing. These mixed findings may result from confounding

factors such as a small number of participants, an imbalanced participant sex ratio, and

the varying nature of the tasks in previous reports. To address these issues, we assessed

the effect sizes of face orientation, sex, and their interaction in low-level psychophysical

discrimination of briefly presented gaze stimuli in a well-powered cohort with a balanced

sex ratio (N = 190). The results revealed a medium-to-large-sized effect of inversion (f =

0.35) and a small-to-medium-sized effect of sex (f = 0.17) on gaze direction

discrimination but no meaningful interaction (f = 0.05). Thus, our results demonstrated an

almost equivalent gaze inversion effect between females and males (sex invariance) and a

female superiority in overall gaze direction sensitivity (sex difference). We suggest that

inversion impairs sensitivity to gaze direction, but whether it leads to a change in explicit

behavioral performance may depend on specific task settings rather than the participant's

sex.

Keywords:

Gaze inversion effect; Face inversion effect; Gaze perception; Sex invariance; Sex

difference

Acknowledgements:

This study was supported by grants from the Japan Society for the Promotion of Science

(JP19H04433 to DM; JP22H00090 to KW). The photographed model acknowledges the

use of the face images in Figure 1 with written informed consent. The dataset analyzed

during the current study is available at GitHub (upon publication):

https://github.com/dicemt/matsuyoshi_gaze_inversion_effect

Abbreviation:

FIE: face inversion effect; GIE: gaze inversion effect

Competing interests:

None of the authors have any potential conflicts of interest.

Authors' contributions:

All authors designed the study and wrote the manuscript. DM and KK collected stimulus

materials. DM collected and analyzed the data.

1   **1. Introduction**

Humans are experts at distinguishing and recognizing faces under various

conditions. People can (albeit explicitly or implicitly) detect subtle differences and

distinguish between thousands of individuals even though typical human faces share

strikingly similar morphology (e.g., two eyes at the top, a nose in the middle, and a

mouth at the bottom of a frontal face). However, the case is remarkably different for

inverted faces (Yin, 1969). Turning a face upside-down makes it particularly difficult to

recognize, whereas inverting other types of objects do not affect recognition performance

as much.

This face inversion effect (FIE) is exceptionally robust across observers and

conditions. It has led researchers to speculate that there are *qualitative* differences between

upright and inverted face processing; upright face processing involves holistic or

configural processing of faces, whereas inverted face processing involves piece-meal or

parts-based processing of faces (Maurer et al., 2002). This holistic or configural

processing assumes that perceiving a face as a coherent whole (rather than as a collection

of separate elements) is an idiosyncratic feature that makes face recognition special (Farah

et al., 1998; Moscovitch et al., 1997). By contrast, recent studies have suggested that the

differences in processing upright and inverted faces are *quantitative* (i.e., a matter of

degree) rather than qualitative (i.e., distinct processing style) (Gold et al., 2012; Jiang et al.,

2006; Matsuyoshi et al., 2015; Sekuler et al., 2004). Sekuler et al. (2004) used a response

classification technique to reveal the most discriminative stimulus regions that affected

performance. They found that people used similar local regions of faces for discrimination

in both upright and inverted face conditions. In addition, Gold et al. (2012) showed that the

perception of a face is no better than would be predicted by the facial feature integration of

an optimal Bayesian integrator. If observers use a holistic face representation, they should

show superoptimal performance for a whole face (better than the sum of its parts); however,

the performance was comparable to the integrated prediction derived from individual facial

features presented in isolation. Contrary to the qualitative view, these results support the

quantitative view where humans use a similar recognition strategy between upright and

inverted faces. However, the two views are not mutually exclusive; dissociable neural

pathways may mediate the two mechanisms (Matsuyoshi et al., 2015).

The behavioral FIE itself is very strong, as rigorous replications have confirmed it;

however, it is unclear whether inversion impairs the perception of individual facial

features. In particular, whether the sensitivity to eye or gaze processing differs as a

function of face orientation remains to be clarified. Several studies have shown that

inversion has little effect on distinguishing faces that differ only in the shape of

individual features (Freire et al., 2000; Le Grand et al., 2001). For example, although

upright faces with different hairstyles and costumes tend to be perceived as different

individuals even though their internal features are identical (Sinha & Poggio, 1996),

composite face procedures (Murphy et al., 2017) have shown that inversion remarkably

improves the ability to identify them (Hole, 1994; Young et al., 1987). Furthermore, Xu

and Tanaka (2013) showed that inversion had little or no effect on discriminating featural

and configural differences in the eye region (while severely disrupting the perception of

changes in the lower region of the face). These results suggest that featural processing, at

least eye or gaze processing, is preserved or sometimes enhanced when faces are

upside-down. By contrast, numerous studies have shown the opposite: featural processing

becomes difficult when faces are inverted. One notable example is the Thatcher illusion

(Thompson, 1980). People fail to notice local feature changes in an inverted face, even

though the changes are evident in an upright face. Furthermore, studies have suggested

that an isolated feature (without a face context) alone can produce inversion effects

(Leder et al., 2001; Rakover & Teucher, 1997). These findings indicate that it is difficult

to process individual features when they are upside-down.

Because of the variation among studies (e.g., sample size, sex ratio, study design,

and task settings), these discrepancies are difficult to resolve. In particular, sample size

and sex ratio would be the most important factors in face processing studies. Insufficient

statistical power due to a small sample size likely leads to false positives and undermines

the reliability of findings (Button et al., 2013; Simmons et al., 2011). Furthermore,

although females often outperform males in face processing performance (Bayliss et al.,

2005; Goodman et al., 2012; Heisz et al., 2013; Matsuyoshi et al., 2014; Matsuyoshi &

Watanabe, 2021; McClure, 2000), studies have not always paid attention to the

participant sex ratio unless the study focuses on sex differences. Since participant sex can

interact with primary behavioral measures of interest (Cellerino et al., 2004; Lewin &

Herlitz, 2002; Matsuyoshi et al., 2014), an imbalanced sex ratio likely impinges on the

conclusion of whether inversion affects featural processing.

To address these issues, we examined the effects of inversion and sex on featural

processing in a well-powered cohort with a balanced sex ratio. We determined our sample

size based on an a priori power analysis, which allowed us to assess whether inversion

affects featural processing and whether sex interacts with it. In addition, we investigated

low-level psychophysical discrimination of gaze direction as it is crucial to determine the

effects of inversion on facial feature processing. The current approach allowed us to

estimate a more accurate effect size of inversion on featural processing while controlling

for possible confounding sources emanating from higher cognitive processes, a small

sample size, or an imbalanced sex ratio.

1   **2. Materials and Methods**

2.1 Participants

We assumed a medium-sized effect of sex on gaze direction discrimination based on

studies showing sex differences in face or gaze perception (Bayliss et al., 2005; Goodman

et al., 2012; Matsuyoshi et al., 2014; Matsuyoshi & Watanabe, 2021). To achieve a power

$(1 - \beta)$ of 0.95 and a medium effect size (Cohen's f = 0.25) while assuming a high

correlation between the measures (r = 0.7 between upright and inverted gaze processing), a

sample of 180 (90 for each group) is required to detect a significant difference (α = 0.05)

between participant sexes (Cohen, 1988; Faul et al., 2007). In addition, we conservatively

assumed the effect size of inversion as small-to-medium (Cohen's f = 0.15) because of the

mixed results of previous studies. A sample of 90 (45 for each group) is required to detect a

significant difference (α = 0.05, $1 - \beta$ = 0.95, r = 0.7) between upright and inverted faces

with a small-to-medium effect size, and a sample of 90 (45 for each group) is required to

detect a significant interaction (α = 0.05, $1 - \beta$ = 0.95, r = 0.7) between the face orientation

and sex with a small-to-medium effect size. Therefore, we decided to recruit a sample of

more than 200 participants to confirm small-to-medium-sized effects between conditions

and to overcome the potential loss of analyzable data, often caused by participants'

inappropriate responses or technical errors.

Two hundred and thirteen young adults participated. All had normal or

corrected-to-normal vision, and none reported a history of neurological or developmental

disorders. The procedures were made following the Declaration of Helsinki and approved

by the Committee of Ethics, Waseda University, Japan. Each participant gave written

informed consent before the experiment. Twenty-three participants were excluded from

further analysis because they did not perform the task correctly (their performance did

not reach the 60% level at an angle of 0°). After these exclusions, data from 190

participants (self-reported 92 females, 98 males; mean age: 21.3 years; range: 18-29

years) remained for analysis.

2.2 Stimuli

Face images were taken from three male and three female models, looking either

directly at a camera (0°) or 10°, 20°, or 30° to the left or right with their face square to the

camera, photographed by the authors (Matsuyoshi et al., 2014). Images were gray-scaled

and subtended approximately 12 by 17° of the visual angle. The mean brightness was

normalized across images. A mask stimulus was generated by grid-scrambling an

averaged image of the face stimuli.

2.3 Procedure

Participants viewed the stimuli on a 21-inch CRT monitor at a distance of 57 cm

with their heads resting on a chin-rest and judged whether the eyes were looking directly

at them (Matsuyoshi et al., 2014). All stimuli were presented against a black background.

Each trial began with an upright or inverted face stimulus (20 ms) followed by a brief

blank screen (20, 40, or 60 ms), a visual mask (100 ms), and finally, a cross (until

response) (Fig. 1). Face images were either direct gaze (0°), or left- or right-averted gaze

(10°, 20°, or 30°). The vertical positions of the eyes were aligned between upright and

inverted faces and located along the vertical center of the display. We used a variable

inter-stimulus interval (ISI) between the face and the mask to control task difficulty. Each

left or right angle was presented 12 times, except for 0° stimuli, which were presented 24

times, for each of the three ISIs for both face orientations, resulting in 576 trials in total

(6 gaze angles [left or right 10-30°] × 3 ISIs × 2 face orientations × 12 trials + 1 gaze

angle [0°] × 3 ISIs × 2 face orientations × 24 trials). The two face orientations were

presented separately in four alternating blocks of 144 trials each, with the order

counterbalanced across participants (i.e., either upright-inverted-upright-inverted or

inverted-upright-inverted-upright). Gaze directions and ISIs were randomized across

trials. Participants performed the main experiment after 30 practice trials.

2.4 Analysis

Responses from left- and right-averted gaze stimuli were pooled for each absolute

angle. In addition, responses from different ISIs were also pooled. The gaze threshold

was defined as the angle at which a 50% response was obtained, as estimated by a local

linear fit of a nonparametric psychometric function (Żychaluk & Foster, 2009) to each

observer's 'direct' responses as a function of gaze angle. A higher threshold was

considered to be less sensitive to gaze direction. We set the statistical thresholds to α =

0.05.

<div align="center">**3. Results**</div>

3.1 Psychometric functions

We performed a repeated-measures ANOVA (degrees of freedom were

Greenhouse-Geisser corrected in case of sphericity violation) on participants' 'direct'

responses with angle (0°, 10°, 20°, 30°) and face orientation (upright, inverted) as

within-subject factors and participant sex (male, female) as a between-subject factor (Fig.

2a). The main effects of the angle ($F_{(1.989,373.885)}$ = 4796.9115, p = 9.8544 × 10^{-270}, η_p^2 =

0.9636, ω_G^2 = 0.8809, Cohen's f = 2.7367), face orientation ($F_{(1,188)}$ = 158.2230, p =

1.0070 × 10^{-26}, η_p^2 = 0.4570, ω_G^2 = 0.0639, Cohen's f = 0.0909), and participant sex

($F_{(1,188)}$ = 9.7583, p = 2.0068 × 10^{-26}, η_p^2 = 0.0493, ω_G^2 = 0.0141, Cohen's f = 0.0436)

were significant. This demonstrated a decrease in 'direct' responses with increasing angle,

higher 'direct' responses in the inverted compared to the upright face condition, and

higher 'direct' responses in male compared to female participants, respectively.

We found significant interactions between the angle and orientation ($F_{2.143,402.849}$ =

276.8289, p = 9.4991 × 10^{-80}, η_p^2 = 0.5960, ω_G^2 = 0.1171, Cohen's f = 0.1270) and

between the angle and participant sex ($F_{1.989,373.885}$ = 6.8471, p = 0.0012, η_p^2 = 0.0351,

ω_G^2 = 0.0086, Cohen's f = 0.0350), while no significant interactions were found between

the orientation and sex ($F_{1,188}$ = 1.6217, p = 0.2044, η_p^2 = 0.0086, ω_G^2 = 0.0003,

Cohen's f = 0.0092) and between the angle, orientation, and sex ($F_{2.143,402.849}$ = 0.9617, p

= 0.3881, η_p^2 = 0.0051, ω_G^2 = −1.8401 × 10^{-5}, Cohen's f = 0.0074). Follow-up tests

revealed higher 'direct' responses in the upright compared with the inverted face

condition at an angle of 0° ($F_{1,188}$ = 68.2900, p = 2.4857 × 10^{-14}, η_p^2 = 0.2665, ω_G^2 =

0.0935, Cohen's f = 0.3251), and higher 'direct' responses in the inverted compared to the

upright face condition at angles between 10 and 30° ($F_{1,188}$ values > 65.8134, p values <

6.2740 × 10^{-14}, η_p^2 values > 0.2592, ω_G^2 values > 0.0835, Cohen's f values > 0.2978). In

addition, males showed higher 'direct' responses at angles between 10 and 30° than

females ($F_{1,188}$ values > 8.6726, p values < 0.0036, η_p^2 values > 0.0441, ω_G^2 values >

0.0311, Cohen's f values > 0.1651). By contrast, males and females showed similar

'direct' responses at an angle of 0° ($F_{1,188}$ = 0.2151, p = 0.6433, η_p^2 = 0.0011, ω_G^2 =

−0.0029, Cohen's f = 0.0271).

3.2 Gaze thresholds

A 2 (upright, inverted) × 2 (male, female) repeated-measures ANOVA on the 50%

gaze perception threshold (Fig. 2b) showed a significant main effect of face orientation

($F_{1,188}$ = 109.6661, p = 1.6503 × 10^{-20}, η_p^2 = 0.3684, ω_G^2 = 0.1112, Cohen's f = 0.3515)

with a larger threshold in the inverted compared to the upright face condition

(medium-to-large effect size), and a significant main effect of sex ($F_{1,188}$ = 7.4251, p =

0.0070, η_p^2 = 0.0380, ω_G^2 = 0.0257, Cohen's f = 0.1652) with a larger threshold for male

compared to female participants (small-to-medium effect size). However, there was no

significant interaction between the face orientation and participant sex ($F_{1,188}$ = 2.0730, p

= 0.1516, η_p^2 = 0.0109, ω_G^2 = 0.0012, Cohen's f = 0.0456), showing that females and

males suffered similarly from inversion (females, t_{91} = 6.6386, p = 2.2441 × 10^{-9},

Cohen's d = 0.6921 [95% CI: 0.3936, 0.9888]; males, t_{97} = 8.1743, p = 1.1419 × 10^{-12},

Cohen's d = 0.8257 [95% CI: 0.5330, 1.1165]; females − males, $t_{187.748}$ = 1.4443, p =

0.1503, Cohen's d = 0.2097 [95% CI: −0.0759, 0.4947]), whereas females had a higher

sensitivity than males for discriminating gaze direction irrespective of face orientation

(upright, $t_{170.178}$ = 2.3087, p = 0.0221, Cohen's d = 0.3351 [95% CI: 0.0481, 0.6212];

inverted, $t_{177.046}$ = 2.6104, p = 0.0098, Cohen's d = 0.3789 [95% CI: 0.0913, 0.6655]).

There was a significant strong correlation between the upright and inverted gaze

thresholds (Fig. 2c; overall, r = 0.6315, p = 1.5559 × 10^{-22}; males, r = 0.6933, p = 2.5310

× 10^{-15}; females, r = 0.4606, p = 3.8444 × 10^{-6}), indicating an overlap between upright

and inverted gaze processing. The correlation was significantly greater in males than in

females (z = 2.4154, p = 0.0157). Note, however, that the difference was not so strong

that excluding one male and one female outlier (who showed the largest inverted gaze

thresholds) resulted in a conventional marginal significance level (z = 1.8829, p =

0.0597).

1   **4. Discussion**

Our results demonstrated that inversion impaired (but did not preserve or enhance)

gaze direction sensitivity using a well-powered cohort with an a priori determined sample

size. The absence of a significant interaction between sex and face orientation indicates

that the gaze inversion effect (GIE) was not dependent on sex (i.e., sex invariant),

although the overall sensitivity was higher in females than in males (sex difference).

These results suggest that inversion does impair gaze direction sensitivity. Our results

support the quantitative view of the inversion effects, which assumes impaired featural

processing (Gold et al., 2012; Sekuler et al., 2004), and are inconsistent with the stringent

qualitative view, which postulates preserved or enhanced featural processing (Le Grand et

al., 2001; Leder & Bruce, 1998; Rossion, 2008; Tanaka & Sengco, 1997; Xu & Tanaka,

2013). Given that the quantitative and qualitative mechanisms underlying the inversion

effects can be dissociable in neural pathways (Matsuyoshi et al., 2015), our results do not

necessarily provide strong support for either view. Instead, our findings suggest a

nuanced perspective, providing a refutation of the strict quantitative view rather than the

ultimate proof of the quantitative view. One might argue that the GIE provides evidence

of qualitative differences in upright and inverted face processing, as it likely reflects

holistic processing within the eye for an upright face. However, our findings are (at least)

inconsistent with the strict qualitative view, which assumes preserved parts-based

processing for inverted faces.

4.1 Qualitative and quantitative views of the inversion effects

Several studies have suggested that inversion preserves featural processing while

severely impairing holistic or configural processing, such as the distance between

features (Freire et al., 2000; Le Grand et al., 2001; Rossion, 2008; Tanaka & Sengco,

1997; Young et al., 1987). The qualitative view predicts the comparable performance

between upright and inverted face processing because inversion prompts feature-based

encoding of faces (Farah et al., 1995). In particular, it has been suggested that the eye is

the least likely facial feature to be affected by inversion compared to other parts of the

face: Xu and Tanaka (2013) showed that inversion did not affect the perception of

changes in the eye region, whereas it degraded the perception of changes in the mouth

region (N = 22, 40% female). Although their results seem inconsistent with our results,

their task presented a face stimulus for 500 ms, which was significantly longer than our

experiment. The longer encoding duration likely alleviated or masked the lower

sensitivity to inverted gaze relative to upright gaze. Indeed, behavioral performance was

close to 90% in their study, perhaps reflecting a ceiling effect.

By contrast, the quantitative view predicts the opposite, assuming general

inefficiency for inverted relative to upright faces (Gold et al., 2012; Jiang et al., 2006;

Sekuler et al., 2004). Although using a between-participants design, a small population

(N = 6 each), and an unknown sex ratio, Jenkins and Langton (2003) reported that

inversion exacerbated gaze direction sensitivity. Furthermore, Rakover and Teucher

(1997) (N = 16, 73% female) and Leder et al. (2001) (N = 20, 90% female) found

difficulty in processing individual features when they were inverted. However, the small

sample sizes and unbalanced or unknown sex ratio made the results inconclusive.

Leveraging a well-powered cohort and a balanced sex ratio, we provide clear evidence

for an impairment of featural processing (at least gaze direction sensitivity) by inversion.

4.2 The effects of task settings on inversion effects

Although we demonstrated an impairment of featural processing by inversion, many

studies (Freire et al., 2000; Hole, 1994; Le Grand et al., 2001; Young et al., 1987),

including large-sample studies (N = 242–282) (Rezlescu et al., 2017; Susilo et al., 2013),

have confirmed the preserved (or sometimes better) ability to compare individual features

when faces are inverted. The apparent inconsistency would not result from low statistical

power or an imbalanced sex ratio as those (and our) studies included large populations,

and we (48% female) and Susilo et al. (2013) (52% female) achieved a balanced sex ratio.

We suggest that it is rather the task settings that have produced the apparent inconsistency.

Featural processing is sometimes preserved (or enhanced) despite the diminished

sensitivity, perhaps not because parts-based processing is dominant for inverted faces

(Farah et al., 1995; Tanaka & Farah, 1993), but because the tasks require discrimination

between exemplars from the same facial parts (e.g., comparing the eyes of different

people). Thus, even though perceptual sensitivity per se decreases, the ability to compare

two signals is relatively unlikely to deteriorate because sensitivity to both signals, but not

one or the other, decreases. Moreover, the apparent dominance of feature-based

processing for inverted faces may result from difficulty decomposing facial parts for

upright faces, rather than enhanced featural processing for inverted faces. Processing an

upright face likely involves tight integration of internal features, making it difficult to

parse the face into isolated features (Maurer et al., 2002); however, an inverted face is

unlikely subject to this integration. Therefore, whether inversion leads to a change in

explicit behavioral performance may depend on specific task settings rather than the

participant's sex.

Note that the effect size of the GIE was smaller than that of the typical FIE. Despite

a huge effect of the FIE in a face recognition task ($d = 1.298$) (Matsuyoshi et al., 2015),

we found a medium-to-large effect of the GIE in the present study ($d = 0.761$, all

participants combined). One reason for the tremendous difference in effect size might be

that a whole face accumulates the inversion effects of each feature (Gold et al., 2012).

Although the susceptibility to inversion may differ across parts (Rakover & Teucher,

1997; Xu & Tanaka, 2013), they may aggregate to produce a large effect size of the FIE.

Alternatively, the large effect size difference may simply be explained by the difference

in processing load required by each task. The present task encouraged observers to focus

on the eyes, whereas typical FIE tasks do not explicitly encourage observers to focus on a

local feature but rather on a whole face. Encoding a whole face might be more difficult

than encoding a single feature of the face.

4.3 Sex differences in face and gaze processing

Consistent with numerous studies showing female advantages in face processing

(Bayliss et al., 2005; Goodman et al., 2012; Heisz et al., 2013; Matsuyoshi et al., 2014;

Matsuyoshi & Watanabe, 2021; McClure, 2000), we found that females outperformed

males in gaze direction sensitivity. It is worth noting that the effect size observed for the

female advantage in upright gaze direction sensitivity (Cohen's $d = 0.3351$) was far

smaller than that for upright face recognition in a previous study (Cohen's $d = 0.6426$, N

= 180, 45% female) (Matsuyoshi & Watanabe, 2021). This indicates that featural

processing is likely to manifest weaker effects (i.e., inversion effects and female

advantage) than whole-face processing, as in the effect size difference between the GIE

and FIE.

What exactly causes the sex differences in face and gaze processing remains to be

investigated, although studies have suggested that not only genetic but also non-genetic

factors together contribute to sex differences in behavioral phenotypes (Bölte et al., 2023;

Kiesow et al., 2020; Kundakovic & Tickerhoof, 2024). For example, studies on sex

differences in nonverbal communication have suggested that sex differences may exist

because people possess stereotypes about them (Briton & Hall, 1995). We cannot, of

course, rule out the possibility that they are real sex differences tightly regulated by

genetic factors such as sex hormones; however, internalized social norms or stereotypes

may continue to influence us throughout our lives (Crandall et al., 2002; Gavrilets &

Richerson, 2017; Rutland et al., 2005). These social factors may also affect sex

differences in face and gaze differences.

A notable limitation of our study is that we treated sex/gender in a binary fashion

and findings solely depended on self-reported sex/gender. Outcomes can vary across

individuals of different birth-assigned sexes and gender identities (Bölte et al., 2023; Joel,

2021). The definitions of sex/gender are continuously evolving and sex/gender is now

considered a spectrum (Ainsworth, 2015; Kundakovic & Tickerhoof, 2024; Rosenthal,

2021). Future research taking human variation in the sex/gender spectrum into account

will further elucidate the roles sex/gender play in face and gaze processing.

4.4 Conclusion

In conclusion, our results provide clear evidence for an inversion-induced decrease

in the perception of gaze direction (GIE). We found both sex invariance (almost

equivalent GIE between females and males) and sex difference (small-to-medium-sized

female superiority) in gaze direction sensitivity. Although previous findings regarding the

effects of inversion on gaze direction processing are mixed, the a priori determined

sample size and the balanced sex ratio in our participants allowed us to characterize the

effect sizes for gaze direction sensitivity. Although our findings do not provide definitive

evidence to resolve the qualitative vs. quantitative debate over the mechanisms of FIE,

we suggest that it is difficult to accept the strict quantitative view of the FIE that assumes

preserved featural processing for inverted faces. It may be "apparently preserved,"

perhaps because some tasks did not require participants to process features in a

time-limited manner, and hence, there were redundant cues or processes to overcome the

impoverished perceptual sensitivity induced by inversion. Since we did not incorporate

tasks other than the discrimination of gaze direction, further research is required to

confirm the critical conditions that discriminate between preserved and impoverished

featural processing by inversion. Nevertheless, our results demonstrate that inversion

does decrease perceptual sensitivity, at least for gaze direction, using a well-powered

cohort with a balanced sex ratio.

References

Ainsworth, C. (2015). Sex redefined. *Nature*, *518*(7539), 288-291. https://doi.org/10.1038/518288a

Bayliss, A. P., di Pellegrino, G., & Tipper, S. P. (2005). Sex differences in eye gaze and symbolic cueing of attention. *The Quarterly Journal of Experimental Psychology Section A*, *58*(4), 631-650. https://doi.org/10.1080/02724980443000124

Bölte, S., Neufeld, J., Marschik, P. B., Williams, Z. J., Gallagher, L., & Lai, M.-C. (2023). Sex and gender in neurodevelopmental conditions. *Nature Reviews Neurology*, *19*(3), 136-159. https://doi.org/10.1038/s41582-023-00774-6

Briton, N. J., & Hall, J. A. (1995). Beliefs about female and male nonverbal communication. *Sex Roles*, *32*(1), 79-90. https://doi.org/10.1007/BF01544758

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafo, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience*, *14*(5), 365-376. https://doi.org/10.1038/nrn3475

Cellerino, A., Borghetti, D., & Sartucci, F. (2004). Sex differences in face gender recognition in humans. *Brain Research Bulletin*, *63*(6), 443-449. https://doi.org/10.1016/j.brainresbull.2004.03.010

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Lawrence Erlbaum.

Crandall, C. S., Eshleman, A., & O'Brien, L. (2002). Social norms and the expression and suppression of prejudice: The struggle for internalization. *Journal of Personality*

and Social Psychology, *82*(3), 359-378.
https://doi.org/10.1037/0022-3514.82.3.359

Farah, M. J., Tanaka, J. W., & Drain, H. M. (1995). What causes the face inversion effect? *Journal of Experimental Psychology: Human Perception and Performance*, *21*(3), 628-634. https://doi.org/10.1037/0096-1523.21.3.628

Farah, M. J., Wilson, K. D., Drain, M., & Tanaka, J. N. (1998). What is 'special' about face perception? *Psychological Review*, *105*(3), 482-498.
https://doi.org/10.1037/0033-295x.105.3.482

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*(2), 175-191.
https://doi.org/10.3758/BF03193146

Freire, A., Lee, K., & Symons, L. A. (2000). The face-inversion effect as a deficit in the encoding of configural information: Direct evidence. *Perception*, *29*(2), 159-170.
https://doi.org/10.1068/p3012

Gavrilets, S., & Richerson, P. J. (2017). Collective action and the evolution of social norm internalization. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(23), 6068-6073.
https://doi.org/doi:10.1073/pnas.1703857114

Gold, J. M., Mundy, P. J., & Tjan, B. S. (2012). The perception of a face is no more than the sum of its parts. *Psychological Science*, *23*(4), 427-434.
https://doi.org/10.1177/0956797611427407

Goodman, L. R., Phelan, H. L., & Johnson, S. A. (2012). Sex differences for the

recognition of direct versus averted gaze faces. *Memory*, *20*(3), 199-209. https://doi.org/10.1080/09658211.2011.651089

Heisz, J. J., Pottruff, M. M., & Shore, D. I. (2013). Females scan more than males: a potential mechanism for sex differences in recognition memory. *Psychological Science*, *24*(7), 1157-1163. https://doi.org/10.1177/0956797612468281

Hole, G. J. (1994). Configurational factors in the perception of unfamiliar faces. *Perception*, *23*(1), 65-74. https://doi.org/10.1068/p160747

Jenkins, J., & Langton, S. R. H. (2003). Configural processing in the perception of eye-gaze direction. *Perception*, *32*(10), 1181-1188. https://doi.org/10.1068/p3398

Jiang, X., Rosen, E., Zeffiro, T., VanMeter, J., Blanz, V., & Riesenhuber, M. (2006). Evaluation of a shape-based model of human face discrimination using fMRI and behavioral techniques. *Neuron*, *50*(1), 159-172. https://doi.org/10.1016/j.neuron.2006.03.012

Joel, D. (2021). Beyond the binary: Rethinking sex and the brain. *Neuroscience & Biobehavioral Reviews*, *122*, 165-175. https://doi.org/10.1016/j.neubiorev.2020.11.018

Kiesow, H., Dunbar, R. I. M., Kable, J. W., Kalenscher, T., Vogeley, K., Schilbach, L., Marquand, A. F., Wiecki, T. V., & Bzdok, D. (2020). 10,000 social brains: Sex differentiation in human brain anatomy. *Science Advances*, *6*(12), eaaz1170. https://doi.org/10.1126/sciadv.aaz1170

Kundakovic, M., & Tickerhoof, M. (2024). Epigenetic mechanisms underlying sex differences in the brain and behavior. *Trends in Neurosciences*, *47*(1), 18-35. https://doi.org/10.1016/j.tins.2023.09.007

Le Grand, R., Mondloch, C. J., Maurer, D., & Brent, H. P. (2001). Neuroperception: Early visual experience and face processing [10.1038/35073749]. *Nature*, *410*(6831), 890-890. https://doi.org/10.1038/35073749

Leder, H., & Bruce, V. (1998). Local and relational aspects of face distinctiveness. *The Quarterly Journal of Experimental Psychology Section A*, *51*(3), 449-473. https://doi.org/10.1080/713755777

Leder, H., Candrian, G., Huber, O., & Bruce, V. (2001). Configural features in the context of upright and inverted faces. *Perception*, *30*(1), 73-83. https://doi.org/10.1068/p2911

Lewin, C., & Herlitz, A. (2002). Sex differences in face recognition—Women's faces make the difference. *Brain and Cognition*, *50*(1), 121-128. https://doi.org/10.1016/S0278-2626(02)00016-7

Matsuyoshi, D., Kuraguchi, K., Tanaka, Y., Uchida, S., Ashida, H., & Watanabe, K. (2014). Individual differences in autistic traits predict the perception of direct gaze for males, but not for females. *Molecular Autism*, *5*(1), 12. https://doi.org/10.1186/2040-2392-5-12

Matsuyoshi, D., Morita, T., Kochiyama, T., Tanabe, H. C., Sadato, N., & Kakigi, R. (2015). Dissociable cortical pathways for qualitative and quantitative mechanisms in the face inversion effect. *The Journal of Neuroscience*, *35*(10), 4268-4279. https://doi.org/10.1523/JNEUROSCI.3960-14.2015

Matsuyoshi, D., & Watanabe, K. (2021). People have modest, not good, insight into their face recognition ability: a comparison between self-report questionnaires. *Psychological Research*, *85*(4), 1713-1723.

https://doi.org/10.1007/s00426-020-01355-8

Maurer, D., Le Grand, R., & Mondloch, C. J. (2002). The many faces of configural processing. *Trends in Cognitive Sciences*, *6*(6), 255-260. https://doi.org/10.1016/S1364-6613(02)01903-4

McClure, E. B. (2000). A meta-analytic review of sex differences in facial expression processing and their development in infants, children, and adolescents. *Psychological Bulletin*, *126*(3), 424-453. https://doi.org/10.1037/0033-2909.126.3.424

Moscovitch, M., Winocur, G., & Behrmann, M. (1997). What is special about face recognition? Nineteen experiments on a person with visual object agnosia and dyslexia but normal face recognition. *Journal of Cognitive Neuroscience*, *9*(5), 555-604. https://doi.org/10.1162/jocn.1997.9.5.555

Murphy, J., Gray, K. L. H., & Cook, R. (2017). The composite face illusion. *Psychonomic Bulletin & Review*, *24*(2), 245-261. https://doi.org/10.3758/s13423-016-1131-5

Rakover, S. S., & Teucher, B. (1997). Facial inversion effects: Parts and whole relationship. *Perception & Psychophysics*, *59*(5), 752-761. https://doi.org/10.3758/BF03206021

Rezlescu, C., Susilo, T., Wilmer, J. B., & Caramazza, A. (2017). The inversion, part-whole, and composite effects reflect distinct perceptual mechanisms with varied relationships to face recognition. *Journal of Experimental Psychology: Human Perception and Performance*, *43*(12), 1961-1973. https://doi.org/10.1037/xhp0000400

Rosenthal, S. M. (2021). Challenges in the care of transgender and gender-diverse youth:

an endocrinologist's view. *Nature Reviews Endocrinology*, *17*(10), 581-591. https://doi.org/10.1038/s41574-021-00535-9

Rossion, B. (2008). Picture-plane inversion leads to qualitative changes of face perception. *Acta Psychologica*, *128*(2), 274-289. https://doi.org/10.1016/j.actpsy.2008.02.003

Rutland, A., Cameron, L., Milne, A., & McGeorge, P. (2005). Social Norms and Self-Presentation: Children's Implicit and Explicit Intergroup Attitudes. *Child Development*, *76*(2), 451-466. https://doi.org/10.1111/j.1467-8624.2005.00856.x

Sekuler, A. B., Gaspar, C. M., Gold, J. M., & Bennett, P. J. (2004). Inversion leads to quantitative, not qualitative, changes in face processing. *Current Biology*, *14*(5), 391-396. https://doi.org/10.1016/j.cub.2004.02.028

Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological Science*, *22*(11), 1359-1366. https://doi.org/10.1177/0956797611417632

Sinha, P., & Poggio, T. (1996). I think I know that face [10.1038/384404a0]. *Nature*, *384*(6608), 404-404. https://doi.org/10.1038/384404a0

Susilo, T., Rezlescu, C., & Duchaine, B. (2013). The composite effect for inverted faces is reliable at large sample sizes and requires the basic face configuration. *Journal of Vision*, *13*(13), 14-14. https://doi.org/10.1167/13.13.14

Tanaka, J. W., & Farah, M. J. (1993). Parts and wholes in face recognition. *The Quarterly Journal of Experimental Psychology Section A*, *46*(2), 225-245. https://doi.org/10.1080/14640749308401045

Tanaka, J. W., & Sengco, J. A. (1997). Features and their configuration in face recognition. *Memory & Cognition*, *25*(5), 583-592. https://doi.org/10.3758/BF03211301

Thompson, P. (1980). Margaret Thatcher: a new illusion. *Perception*, *9*(4), 483-484. https://doi.org/10.1068/p090483

Xu, B., & Tanaka, J. W. (2013). Does face inversion qualitatively change face processing: An eye movement study using a face change detection task. *Journal of Vision*, *13(2)*, 22. https://doi.org/10.1167/13.2.22

Yin, R. K. (1969). Looking at upside-down faces. *Journal of Experimental Psychology*, *81*(1), 141-145. https://doi.org/10.1037/h0027474

Young, A. W., Hellawell, D., & Hay, D. C. (1987). Configurational information in face perception. *Perception*, *16*(6), 747-759. https://doi.org/10.1068/p160747

Żychaluk, K., & Foster, D. (2009). Model-free estimation of the psychometric function. *Attention, Perception, & Psychophysics*, *71*(6), 1414-1425. https://doi.org/10.3758/APP.71.6.1414

Figure Legends

Figure 1. Experimental Paradigm.

Schematic presentation of the experimental task. Upright and inverted faces were presented in different blocks (4 [2 upright + 2 inverted] blocks × 144 trials).

Figure 2. Experimental Results.

(a) Gaze angle psychometric functions. Mean percentage of 'direct' responses (y-axis), plotted as a function of gaze angle (x-axis) × participant sex (female = red, male = blue) × face orientation (upright face = filled circle with a solid line, inverted face = open circle with dashed line). Shaded areas represent 95% confidence intervals of the mean. (b) Mean gaze threshold (angle of 50% reported as direct, which was estimated from the psychometric function) as a function of participant sex (female = red, male = blue) × face orientation (upright face = filled, inverted face = open). Error bars represent 95% confidence intervals of the mean. (c) Scatter plot and density curves of individual upright and inverted gaze thresholds (female = red, male = blue). The gray dashed line indicates identical gaze thresholds between upright and inverted faces (i.e., no gaze inversion effect).

Figures

Figure 1

* The face image was deidentified for bioRxiv.

(a)

(b)

(c)

Figure 2