Deciphering hepatitis B virus phospho-signature in primary human hepatocytes

Florentin Pastor[1#], Emilie Charles[1#], Lucid Belmudes[2], Hélène Chabrolles [1], Marion Cescato[1], Michel Rivoire[3], Thomas Burger[2], Guillaume Passot[4], David Durantel[1], Julie Lucifora[1], Yohann Couté[2*], and Anna Salvetti[1*].

[1] International Center for Research in Infectiology (CIRI), INSERM U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS, Lyon, France

[2] Université Grenoble Alpes, CEA, INSERM, UA13 BGE, CEA, CNRS, FR2048, Grenoble, France

[3] Centre Léon Bérard (CLB), INSERM, U1032, Lyon, France

[4] Service de chirurgie générale et oncologique, Hôpital Lyon Sud, Hospices Civils de Lyon Et CICLY, EA3738, Université Claude Bernard Lyon 1, Lyon France

# Equal contribution and first authorship

* Last autorship

* Correspondence:

Anna Salvetti

anna.salvetti@inserm.fr

Yohann Couté

yohann.coute@cea.fr

Keywords: Hepatitis B virus, primary hepatocytes, phosphorylation, anti-viral response, DNA Damage Response, RNA-binding proteins.

Figures = 11

Supplemental Figures = 3

Supplemental Tables = 6

Word count:

Abstract: 350

Main text:

ABSTRACT

Phosphorylation is a major post-translation modification (PTM) of proteins, and small molecules,

which is finely tuned by the activity of several hundred kinases and phosphatases. It controls most if

not all cellular pathways including anti-viral responses. Accordingly, viruses often induce important

changes in the phosphorylation of host factors that can either help or counteract viral replication.

Among more than 500 kinases constituting the human kinome only few have been described as

important for the Hepatitis B virus (HBV) infectious cycle, and most of them intervene during early

or late infectious steps by phosphorylating the viral Core protein (HBc) protein. In addition, scarce

information is available on the consequences of HBV infection on the activity of cellular kinases.

The objective of this study was to investigate the global impact of HBV infection on the cellular

phosphorylation landscape at early after infection. For this, primary human hepatocytes (PHH) were

challenged or not with HBV, and a mass spectrometry (MS)-based quantitative phosphoproteomic

analysis was conducted two- and seven-days post-infection. The results indicated that while, as

expected, HBV infection only minimally modified the cell proteome, important changes were

observed on the phosphorylation level of several host proteins at both times points. Gene enrichment

and ontology analyses of up- and down- phosphorylated proteins revealed common and distinct

signatures induced following infection. In particular, HBV infection resulted in up-phosphorylation

of proteins involved in DNA damage signaling and repair, RNA metabolism, in particular splicing,

and cytoplasmic cell-signaling. Several down-phosphorylated factors, mostly involved in cell

signaling and communication, were also detected. Validation studies performed on some selected up-

phosphorylated proteins, revealed that HBV infection induced a DNA damage response characterized

by the appearance of 53BP1 foci, whose siRNA-mediated knock-down increased cccDNA levels. In

addition, among up-phosphorylated RBPs, SRRM2, a major scaffold of nuclear speckles behaved as

antiviral factor. In accordance with these findings, kinase prediction analysis indicated that HBV

infection upregulates the activity of major kinases involved in DNA repair. These results strongly

suggest that HBV infection triggers an intrinsic anti-viral response composed by DNA repair factors

and RBPs that contribute to reduce HBV replication in cell culture models.

1   INTRODUCTION

Despite the availability of an efficient preventive vaccine, nearly 300 million people worldwide are chronically infected with the Hepatitis B virus (HBV), making it a major contributor to liver diseases, especially cirrhosis and hepatocellular carcinoma (HCC) (https://www.who.int/news-room/fact-sheets/detail/hepatitis-b). Current antiviral treatments, primarily nucleotide analogs, can reduce viremia and the incidence of HCC. However, these treatments cannot completely eliminate the infection due to the persistence of a transcriptionally active viral genome in the nucleus of hepatocytes (Fanning et al., 2019).

The infection of human hepatocytes by HBV occurs through the recognition of its primary and secondary receptors on the cell surface, followed by the endocytosis of viral particles (Tsukuda and Watashi, 2020). After release into the cytosol, the capsid is directed towards the nucleus, where it binds to nuclear pores *via* its interaction with importins, before translocating through the nuclear channel and disassembling in the inner nuclear basket (Blondot et al., 2016;Diogo Dias et al., 2021). These early events result in the release of Core (HBc) protein, the unique structural component of the capsid, and the partially double-stranded (ds) circular genome known as relaxed circular DNA (rcDNA) to which the viral polymerase is covalently attached. The rcDNA is repaired and loaded with cellular histones and other transcriptional regulators to form an extra-chromosomal covalently-closed circular DNA (cccDNA) which constitutes the template for production of viral RNAs (Diogo Dias et al., 2021;Wei and Ploss, 2021). During these early phases, two viral proteins also associate with cccDNA: HBc, derived from incoming capsids (240 monomers per capsid) or *de novo* translation (Lucifora et al., 2021;Locatelli et al., 2022), and HBx, the major HBV regulatory protein (Diogo Dias et al., 2021;Wang et al., 2023). Notably, HBx production by the infected cell is necessary for continuous viral transcription due to its capacity to induce the degradation of the smc5/6 complex (Lucifora et al., 2011;Decorsiere et al., 2016;Murphy et al., 2016;Niu et al., 2017;Abdul et al., 2022). The late phase of the HBV cycle occurs in the cytoplasm and involves the assembly of newly formed capsids in which the viral pregenomic RNA (pgRNA) is packaged along with the viral polymerase required for its reverse-transcription into rcDNA. Subsequently, these capsids are enveloped within multi vesicular bodies and released into the extracellular space as infectious particles (Prange, 2022).

Most viral infections are characterized by the induction of strong intrinsic and innate anti-viral responses, against which viruses have developed sophisticated strategies to counteract them. In

particular, viral infections can trigger a cellular DNA damage response (DDR), due to the recognition

of viral genomes, which can prevent their replication (Weitzman and Fradet-Turcotte, 2018;Lopez et

al., 2022). Cellular RNA binding proteins (RBPs), participating at all steps of RNA metabolism, may

be also engaged in a conflictual relationship with viral replication (Garcia-Moreno et al.,

2018;Girardi et al., 2021;Lisy et al., 2021). In addition, cells can mount an innate response,

intimately linked to the intrinsic one, resulting in the production of interferons or inflammatory

cytokines which further amplify the anti-viral effect of both infected and neighboring cells

(Guillemin et al., 2021;Justice and Cristea, 2022;Lopez et al., 2022).

In contrast to most viruses, infection of primary hepatocytes with HBV was reported to neither

induce innate responses nor alter the transcription level of cellular genes and was, therefore, qualified

as "stealth" (Mutz et al., 2018;Suslov et al., 2018). Nevertheless, whether HBV infection can tune

cellular factors by acting at a post-translational step, is presently unknown. In particular little

information is available on the effect of HBV infection on the activity of cellular kinases or

phosphatases, and on their downstream consequences on the phosphorylation of host proteins. So far,

a unique study performed on infected hepatoma cells indicated that HBV can alter the host cell

phospho-proteome and, in particular, target proteins involved in RNA processing, further suggesting

that HBV infection can activate kinases involved in RBPs phosphorylation (Lim et al., 2022). Even

though interesting, these results may be biased by the cellular model used for these analyses, HepG2

cells, which, as most cancerous cell lines, have dysregulated pathways, in particular those involved in

viral sensing (Yang et al., 2014;Arzumanian et al., 2021).

The objective of this study was to investigate the global impact of HBV infection on the cellular

phosphorylation landscape of primary human hepatocytes (PHH). Our analyses indicate that HBV

infection triggers important changes on the host cell phospho-proteome, in particular up-

phosphorylation events targeting several RNA-binding proteins (RBPs) and some major factors

implicated in DNA damage signaling and repair. Among these factors, we identified a RBP, SRRM2,

and the DNA damage sensor, 53BP1, as having an antiviral activity which modulates the level of

viral RNAs and cccDNA, respectively. Altogether, these analyses indicate that HBV infection is

sensed by the host cell and triggers an anti-viral response that restrains viral replication.

2   MATERIAL AND METHODS

2.1   Cell culture and infection

HepaRG cells were cultured, differentiated, and infected by HBV as previously described (Gripon et al., 2002). Primary human hepatocytes (PHH) were freshly prepared from human liver resection obtained from the Centre Léon Bérard and Hôpital Lyon Sud (Lyon) with French ministerial authorizations (AC 2013-1871, DC 2013 – 1870, AFNOR NF 96 900 sept 2011) as previously described (Lecluyse and Alexandre, 2010). HBV genotype D inoculum (subtype ayw) was prepared from HepAD38 (Ladner et al., 1997) cell supernatant by polyethylene-glycol-MW-8000 (PEG8000, SIGMA) precipitation (8% final) as previously described (Luangsay et al., 2015). The titer of endotoxin free viral stocks was determined by qPCR. Cells were infected overnight in a media supplemented with 4% final of PEG, as previously described (Gripon et al., 2002). Measure of secreted HBs and HBe antigens was performed by CLIA (Chemo-Luminescent Immune Assay) following manufacturer's instructions (AutoBio, China). HepAD38 cell were cultured as previously described in the absence of tetracyclin (Tet) (Ladner et al., 1997).

2.2   AAV-HBV production

The AAV-HBV wild-type (wt) plasmid was obtained from ML Michel (Pasteur Institute) (Dion et al., 2013). It contains the 1.2 genome copies of HBV genotype D sequence inserted between the AAV2 inverted terminal repeats (ITRs). AAV serotype 3 vectors were produced by transient transfection of 293 cells and double purification on CsCl gradients by the Vector Core of the University of Nantes as previously described (Salvetti et al., 1998). Vectors were titrated by qPCR to determine the number of vector genomes (vg)/ml using primers on the AAV ITRs.

2.3   Cell extracts and mass spectrometry (MS)-based quantitative proteomic analyses

PHH purified from liver resections were plated in 15-cm dishes (3 plates per time point) and 24 to 48 hours later infected with HBV (Multiplicity of infection (MOI) of 500 viral genome equivalents (vge)/cell) or mock-infected overnight. The following day, cells were washed with media then incubated until 2- or 7-days post-infection (dpi). Two independent experiments were performed: one with PHH from one donor (PHH#1, 3 biological replicates per condition were prepared), and the other with PHH from four donors (PHH#4). To prepare protein extracts, the cells were washed once with cold 1XPBS and then directly lysed on the plate using a buffer containing 8M Urea in 50mM

Tris-HCl pH 8.0/75 mM NaCl/1mM EDTA supplemented with protease and phosphatase inhibitors (SIGMA). Cell lysates wee sonicated and frozen at -80°C.

Extracted proteins were reduced using 20mM of dithiothreitol for 1h at 37°C before alkylation with 55mM of iodoacetamide for 1h at 37°C in the dark. Samples were then diluted to ½ using ammonium bicarbonate and digested with LysC (Promega) at a ratio of 1:200 for 4 hours at 37 °C. Then they were diluted again to ¼ and digested overnight at 37 °C with sequencing grade-modified trypsin (Promega) at a ratio of 1:50. Resulting peptides were purified by C18 reverse phase chromatography (Sep-Pak C18, Waters) before drying down. Peptides were then labelled using an isobaric labelling-based approach, relying on tandem mass tags (TMT, (Thompson et al., 2003)) using the 16plex TMTpro isobaric Label Reagent kit (ThermoFisher Scientific) before mixing equivalent amounts and desalting using C18 reverse phase chromatography (Sep-Pak C18, Waters). An aliquot of labelled peptides was kept for total proteome analyses. Phosphopeptide enrichment was performed using titanium dioxide beads (TitanSphere, GL Sciences, Inc.) as previously described (Sonntag et al., 2017) before purification using C18 reverse phase chromatography (Marco SpinColumns, Harvard Apparatus). Isobaric-labelled peptides from total proteome and phosphoproteomes were then fractionated into eight fractions using the Pierce High pH Reversed-Phase Peptide Fractionation Kit (ThermoFisher Scientific) following the manufacturer's instructions, except for the total proteome analysis of samples prepared from PHH of 1 donor for which no fractionation was performed. The peptides were analyzed by online nanoliquid chromatography coupled to MS/MS (Ultimate 3000 RSLCnano and Q-Exactive HF, Thermo Fisher Scientific) using a 180 min gradient for fractions and a 480 min gradient if no fractionation was performed. For this purpose, the peptides were sampled on a precolumn (300 μm x 5 mm PepMap C18, Thermo Scientific) and separated in a 200 cm μPAC column (PharmaFluidics). The MS and MS/MS data were acquired by Xcalibur (version 2.9, Thermo Fisher Scientific). The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2022) partner repository with the dataset identifier PXD051216.

Peptides and proteins were identified and quantified using MaxQuant (version 1.6.0.17, (Tyanova et al., 2016)) searching in Uniprot databases (*Homo sapiens* and Hepatitis B virus taxonomies, 20210628 download) and in the database of frequently observed contaminants embedded in MaxQuant. Trypsin/P was chosen as the enzyme and 2 missed cleavages were allowed. Peptide modifications allowed during the search were: Carbamidomethyl (C, fixed), Acetyl (Protein N-term, variable), Oxidation (M, variable), and Phospho (STY, variable). Minimum peptide length and minimum number of razor+unique peptides were respectively set to seven amino acids and one.

Maximum false discovery rates - calculated by employing a reverse database strategy - were set to
0.01 at peptide-spectrum match, protein and site levels.

Statistical analysis of quantitative data was performed using Prostar (Wieczorek et al., 2017).
Peptides and proteins identified in the reverse and contaminant databases, and proteins only
identified by site were discarded. Only class I phosphosites (localization probability $\geq$ 0.75) and
proteins quantified in all replicates of at least one condition were further processed. After log2
transformation, extracted corrected reporter abundance values were normalized by the Variance
Stabilizing Normalization (vsn) method, before missing value imputation (DetQuantile algorithm).
Statistical testing was conducted with limma for results obtained with PHH from one donor and
limma with paired design for results obtained with PHH from four donors, whereby differentially
expressed proteins were selected using log2(Fold Change) and p-value cut-offs allowing to reach a
false discovery rate inferior to 5% according to the Benjamini-Hochberg estimator. Proteins and
phosphosites found differentially abundant but with imputed values in the condition in which they
were found to be more abundant were manually invalidated (p-value = 1).

2.4 Bionformatic analyses

Gene ontology were performed using Metascape (Zhou et al., 2019) and the Reactome Gene Sets
(http://reactome.org/), a minimum overlap of 3, a minimum enrichment of 1.5, a p value of 0.01 and
all genes as a background. Physical protein-protein interactions (PPI) were similarly analyzed by
Mescape with a minimum network size of 3. Protein networkformed by selected factors were also
analyzed by Cytoscape (v3.10.1) with automatic network weighting. Prediction of kinases involved
in the phosphorylation of differentially abundant phosphosites was performed using KinasePhos 3.0
(Ma et al., 2023).

2.5 siRNA transfection

dHepaRG or PHH cells seeded into a 24-well plate were transfected with 25 nM of siRNA using
Lipofectamine RNAiMax (Life Technologies), following manufacturer's instructions. SiRNA used
were the following: siHNRNPU (Dharmacon SmartPool L-03501-00), siSRRM2 (Dharmacon
SmartPool L-015368-00), si53BP1 (Dharmacon SmartPool L-003548-00), siRIF1 (Dharamcon
SmartPool L-027983) and siControl (Dharmacon D-001810).

2.6 Nucleic acid extractions and analysis

HBV RNAs and DNA were extracted from cells with the Nucleospin RNA (Macherey-Nagel) and
MasterPureTM Complexe Purification Kit (BioSearch Technologies) kit without digestion with
proteinase K (Allweiss et al., 2023), respectively, according to the manufacturer's instruction. RNA
reverse transcription was performed using Maxima First Strand cDNA Synthesis kit (Thermofisher).
Quantitative PCR for HBV were performed using HBV specific primers and normalized to PRNP
housekeeping gene as previously described (Lucifora et al., 2014). Pre-genomic RNA was quantified
using the TaqMan Fast Advanced Master Mix (Life Technologies) and normalized to GusB cDNA
levels. HBV cccDNA was quantified from total DNA by TaqMan qPCR analyses and normalized to
β-globin cDNA level, as previously described (Werle-Lapostolle et al., 2004) or by droplet digital
PCR using the "ddPCR Supermix for Probes (No dUTP)" (Bio-Rad) according to the manufacturer's
instruction. Droplets were generated using and the" QX200™ Droplet Generator" (Bio-Rad) and
analyzed after PCR with the "QX600 Droplet Reader" (Bio-Rad).

2.7   Western blot analyses

Proteins were resolved by SDS-PAGE and then transferred onto a nitrocellulose membrane.
Membranes were incubated with the primary antibodies corresponding to the indicated proteins.
Proteins were revealed by chemi-luminescence (Super Signal West Dura Substrate, Pierce) using a
secondary peroxydase-conjugated antibody (Dako) at a dilution of 1:10000. Primary antibodies used
were anti-53BP1 (Abcam 175933, 1:1000), HNRNPU (Santa-Cruz sc-32315, 1:2000), β-tubulin
(Abcam 6046,1:10000),

2.8   Immunofluorescence analyses

Analyses were performed as described previously using Alexa Fluor 555 secondary antibodies
(Molecular Probes) (Salvetti et al., 2016). Primary antibodies used were: anti-53BP1 (Abcam
175933; 1:250); anti-PML Santa-Cruz sc-966, 1:250); anti HBc (Thermo MA1-7607, 1/500). Nuclei
were stained with Hoescht 33258. Images were collected on a confocal NLO-LSM 880 microscope
(Zeiss). Further image processing was performed using ICY (de Chaumont et al., 2012).

2.9   Statistical analysis

Statistical analyses were performed using the GraphPad Prism 9 software and Mann-Whitney non-
parametric tests. A p value ≤ 0,05 was considered as significant. * correspond to p value ≤ 0.05; **
correspond to p value ≤ 0.01; *** correspond to p value ≤ 0.001.

3   RESULTS

3.1   Early impact of HBV infection on the host cell proteome and phosphoproteome

In order to examine the effect of HBV infection on the host cell phosphoproteome, we deployed a large-scale phosphoproteomic strategy. For this, HBV-infected PHH, derived from a single donor, were lysed at 2- and 7-dpi (Figure 1A and B show experimental outline and levels of infection) before MS-based quantitative analysis of total proteome and phosphoproteome after phosphopeptide enrichment. The analysis of the total proteome indicated that the proteome remains largely unaffected by HBV infection, with less than 30 proteins found as over- or under-expressed following infection at each time point, among the 3467 proteins identified and quantified (Figure 1C) (Supplemental Table 1). Interestingly, among proteins whose amount was increased at 2- and/or 7-dpi, figured, notably, fibronectin which was previously reported to be upregulated following HBV infection (Norton et al., 2004;Ren et al., 2016). Similarly, two HBV under-expressed proteins at 7-dpi, LSM7 and TRIM21, were previously reported to negatively interfere with HBV replication (Song et al., 2021;Rahman et al., 2022). The deep phosphoproteomic profiling of these samples allowed to identify and quantify 8308 phosphopeptides containing class I phosphosites (localization probability > 75%) from 3012 different proteins (Supplementary Table 2). Among them, 161 and 316 were differentially abundant in PHH infected by HBV compared to PHH mock-infected, at 2- and 7-dpi, respectively (Fig. 1D) (Supplemental Table 2). A part of these modulated phosphosites, 17 up-regulated and 54 down-regulated in HBV-infected PHH compared to mock-infected cells, were found differentially abundant at both time points (Supplementary Fig.1A). The comparison of total proteome and phosphoproteome results showed that only one protein, fibrinogen alpha chain (FIBA) but also its phosphorylated sites, were found to be more abundant and in HBV-infected cells compared to control cells at 7-dpi; it was therefore excluded from further analyses since the measured up-regulation of its phosphosites could be linked to the over-expression of the protein. Overall, these results indicated that, in contrast to minimal changes affecting the level of host proteins, HBV infection triggered more significant changes in the phosphoproteome, as soon as 2-dpi, which likely reflected the interplay between the virus and the host cell following attachment, internalization, and onset of viral replication.

3.2   Analysis of host pathways modulated by HBV-induced up or down-phosphorylation events

In order to identify pathways and biological processes targeted by these phosphorylation events, enrichment gene ontology analyses were first performed on up-phosphorylated proteins (Figure 2A).

This analysis indicated that HBV infection was characterized by up-phosphorylation of proteins involved in DNA repair pathways, notably non-homologous end-joining (NHEJ) and MAP kinase signaling, as soon as 2-dpi. DNA repair factors were also targeted at 7-dpi, a time point at which other significant pathways appeared, notably those related to RNA splicing, and cytoplasmic signaling by RhoGTPases. Physical protein-protein interaction analyses based on human interactome datasets indicated that phosphorylated factors at both time points formed a network of proteins which contained densely-connected complexes related to double-stranded DNA breaks (DSBs), in particular NHEJ, and to MAPK-signaling (Figure 2B). A higher number of pathways were enriched when down-phosphorylated proteins were similarly analyzed, highlighting biological processes related to cytoplasmic signal transduction, in particular Rho-GTPases, which formed an important network of interconnected host factors (Figure 3A and B). Notably, EGFR, a host cofactor during HBV internalization (Iwamoto et al., 2019;Fukano et al., 2021), was down-phosphorylated on serine 1166 and formed a major connected node at both time points (Figure 3B). Comparison of our data with those previously reported by Lim *et al.* (Lim et al., 2022), indicated that few up- and down-phosphorylated proteins were shared between the two studies (Supplemental Table 3). Many differences in the experimental set up may explain this low level of overlap, notably, the use of primary (PHH) versus transformed hepatocytes (HepG2). Comparative analyses performed with factors previously described as interacting with HBc, and HBx also retrieved few common proteins (Chabrolles et al., 2020;Van Damme et al., 2021). This latter observation suggests that most phosphorylation/dephosphorylations events observed in our study may not be associated to the capacity of these host factors to interact with viral proteins.

Altogether, these *in silico* analyses suggest that HBV infection triggers up- and down-phosphorylation events that target proteins involved in common pathways related to cytoplasmic signal transduction *via* Rho-GTPases as well as unique pathways, in particular related to DNA repair and RNA metabolism in the case of up-phosphorylated factors.

3.3 HNRNPU and SRRM2, two RBPs up-phosphorylated upon HBV infection, behave as anti-viral factor restricting HBV RNA production

To investigate the functional relevance of this phospho-proteomic analysis, we first focused on up-phosphorylated RBPs. Indeed, RBPs are tightly regulated by phosphorylation, a post-translational modification which is frequently tuned by viral infections (Pastor et al., 2021;Velazquez-Cruz et al.,

2021). In our analysis, several inter-connected RBPs were up-phosphorylated upon HBV infection. In particular, nine proteins (HNRNPH2, HNRNPU, SRRM2, PNN, NCL, BUD13, TP53RK, SET, and SENP13) constituted an enriched cluster of factors involved in mRNA splicing (Figure 4A). Among these, five RBPs, selected using the RBP2GO database (RBP2GO score above 40) (Caudron-Herger et al., 2021), formed an interconnected network (Figure 4B).

Functional validation analyses were performed on two RBPs: HNRNPU, also called SAF-A, (for Scaffold Attachment Factor A), is a DNA- and RNA-binding protein that was identified as a constituent of the nuclear matrix capable of binding to nuclear matrix/scaffold-attachment regions (S/MAR) (Kiledjian and Dreyfuss, 1992;Romig et al., 1992;Jenke et al., 2002;Jenke et al., 2004). More recent studies indicate that HNRNPU is a chromatin scaffold protein which, by oligomerizing with chromatin-associated RNA (caRNA), controls chromatin architecture and cellular gene expression (Nozawa et al., 2017;Fan et al., 2018;Xu et al., 2022). SRRM2, a member of the SR-related protein family, is involved in pre-mRNA maturation as a catalytic component of the spliceosome together with SRRM1, (Blencowe et al., 2000). Recent investigations also indicated that SRRM2, is a major component of nuclear speckles (Ilik et al., 2020;Xu et al., 2022). Both proteins, in particular SRRM2, are phosphorylated on multiple sites (http://www.phosphosite.org/). In this study, these proteins were up-phosphorylated at a discrete number of sites, mainly on serine residues (Supplemental Table 2). To investigate their role during the HBV life cycle, knock-down (KD) experiments were performed on HBV-infected PHH and differentiated HepaRG (dHepaRG), using siRNAs (Figure 5A, B and D). In both cell types, we found that knock-down of HNRNPU or SRRM2 resulted in an increase of HBV RNAs, both total RNAs and pgRNA, without significantly affecting cccDNA levels (Figure 5C and E). These results indicate that, as previously reported for some other cellular RBPs, these factors behave as anti-viral factor that exert their functions at transcriptional and/or post-transcriptional level (Sun et al., 2017;Yao et al., 2018;Yao et al., 2019;Chabrolles et al., 2020;Yao et al., 2023b).

3.4   HBV infection triggers an antiviral DNA damage response characterized by 53BP1 foci

DNA repair factors involved in NHEJ constituted another important cluster of proteins that were up-phosphorylated upon HBV infection (Figure 4A). Among these factors figured, notably, Ku70 (XRCC6), DDB2, Rad50, two sensors of DSBs (Kieffer and Lowndes, 2022), MDC1, and 53BP1 (Figure 4C). These proteins participate in a cascade of signaling events, including phosphorylation events, triggered by DNA damage, in particular by DSBs, that culminate with the recruitment of 53BP1 on chromatin, to form large foci that segregate damaged sites from the rest of the genome

(Shibata and Jeggo, 2020;Rass et al., 2022)). During this process, phosphorylation of 53BP1 by the ATM kinase, among others, is critical for its recruitment to the sites of DNA damage (Jowsey et al., 2007). The resulting 53BP1 foci are essential for the assembly of DNA repair complexes and the initiation of the DNA damage response pathway (Shibata and Jeggo, 2020).

To investigate whether HBV infection could trigger the formation of 53BP1 foci, immune-fluorescence (IF) analyses were performed on HBV-infected PHH. We found that numerous foci could be observed in HBV-infected cells whereas only few were visible in control non-infected cells. Many of these foci, which appeared as early as 1-dpi, were localized at the periphery of the nucleus (Figure 6A). Importantly, these foci were observed using PHH from different donors but were not observed, or only in few cells, when infection was performed in the presence of myrcludex, indicating that their formation may reflect a cell response to incoming HBV genomes (Figure 6B). Their presence was visible up to 7-dpi, when HBc staining was visible (Figure 6C). Altogether, these IF analyses strongly suggested that 53BP1 foci reflect a long-lasting host cell DNA damage response to incoming HBV genomes. In line with this hypothesis, we found that co-labeling of infected PHH with anti-53BP1 and anti-PML antibodies, indicated that most 53BP1 foci colocalize with PML signals which were previously associated to long-lasting 53BP1 foci (Figure 7A and B) (Vancurova et al., 2019). In particular, 53BP1 appeared to form a scaffold around the inner PML signal strongly suggesting that these foci may constitute a hub of antiviral factors (Figure 7C).

To determine whether 53BP1 may regulate the establishment of HBV infection, knock-down of this protein was performed before infecting hepatocytes with HBV (Figure 8A and B). Analyses performed at 7-dpi, indicated that depletion of 53BP1 resulted in a modest but significant increase in the level of cccDNA, strongly suggesting that this factor counteracts cccDNA establishment (Figure 8C). Recent studies have shown that upon binding to damaged DNA, 53BP1 recruits a shieldin complex which prevents long range resections required for DNA repair by homologous recombination (Setiaputra and Durocher, 2019). Recruitment of shieldin is mediated by RIF1 which recognizes phosphorylated residues on 53BP1 (Setiaputra et al., 2022). Interestingly, knock-down of RIF1 before HBV infection similarly increased the level of cccDNA and, downstream, of viral RNAs, as observed for 53BP1 (Supplemental Figure 2).

3.5   Extended phosphoproteomic analysis on several PHH donors and prediction of involved kinases

Our initial study of phosphosites modulated by HBV infection was performed on PHH derived from
a single donor. Therefore, an additional MS-based proteomic and phosphoproteomic analysis was
performed on PHH derived from 4 different donors (PHH#4). Strong donor-dependent variations in
the efficiency of infection were observed among these four PHH batches (Supplemental Figure 3A).
Not surprisingly, in these stringent conditions, of the 5731 proteins detected, very few were found to
be differentially abundant in HBV-infected cells compared to mock-infected ones at 2- and 7-dpi
(Supplemental Table 4 and supplemental Figure 4B). Concerning phosphosites, 9179 were
quantified, among which 54 and 60 were found modulated at 2- and 7-dpi, respectively
(Supplemental Table 5 and supplemental Figure 4 C). Several of them are from important proteins
found to be differentially phosphorylated upon HBV infection in the previous analysis using one
PHH donor, notably 53B and CP1 and SRRM2 (Supplemental Table 6). Enrichment analyses using
the phosphoproteomic dataset obtained from the four PHH donors indicated that HBV infection was
characterized by up-phosphorylation of proteins involved in cell division and signal transduction
(Figure 9A). PPI analysis indicated that most up-phosphorylated factors formed a connected network
of proteins (Figure 9B). All the peptides found significantly up-phosphorylated in this latter
experiment were analyzed with the KinasePhos3.0 software (Ma et al., 2023), to infer the most
probable kinases activated by HBV infection at both time points. Interestingly, among the top most
probable kinases involved in these modifications at both time points, figured, beyond the casein II
kinase (CK2) which is implicated in a plethora of pathways (Borgo et al., 2021), ATM, ATR, and
DNA-PK, the three major kinases activated upon DNA damage and which control the
phosphorylation of downstream effector proteins (Figure10) (Blackford, 2017).

Altogether, this second study performed using PHH from four different donors confirmed that HBV
infection of PHH induced the up-phosphorylation of a reduced but significant set of DNA repair
proteins, in particular 53BP1. This finding correlated with the emergence of 53BP1 foci reflecting the
induction of a DNA damage response. In addition, it also confirmed the up-phosphorylation of RBPs,
such as SRRM2, a major constituent of nuclear speckles.

4   **DISCUSSION**

The conflicting interactions between a virus and the host cell during the early steps of the viral cycle
are critical to determine the issue of the infectious process. In contrast to many other DNA and RNA
viruses, infection by HBV was not reported to induce significant changes in host gene expression or

innate responses, therefore leading to the assumption that HBV was able to enter the cell, and deposit
its genome within the hepatocyte nucleus, without being detected (Mutz et al., 2018;Suslov et al.,
2018). However, whether HBV infection may impact cell's functions by acting at a post-translational
level was still poorly documented. In particular, most viral infections can induce significant changes
in the phosphorylation of host factors, reflecting altered kinase/phosphatase activities hijacked or
induced upon viral entry, as shown for human imunodeficiency virus 1 (HIV-1) or severe acute
respiratory syndrome coronavirus 2 (SARS-COV2), among others (Wojcechowskyj et al.,
2013;Bouhaddou et al., 2020).

In this study we asked whether HBV infection can modify the phosphorylation landscape of primary
human hepatocytes which are growth-arrested and differentiated cells and, therefore, considered as
the gold standard to study HBV life cycle *in vitro* (Lucifora et al., 2020). We found that HBV
infection can trigger both up-phosphorylation and down-phosphorylation of several cellular factors
without significantly altering their expression level. As expected, many of these proteins were
involved in pathways related to cell signaling. In particular, factors involved in signaling by MAPK
were up-phosphorylated upon HBV infection. Even though signaling by these kinases are not
essential for HBV infection, induction of this pathway may be linked to the requirement of EGFR for
HBV entry (Iwamoto et al., 2019;Iwamoto et al., 2020). Interestingly, upon infection, EGFR was
down-phosphorylated on serine 1166, whose activation by phosphorylation was previously reported
to have negative impact on the EGFR activity (Assiddiq et al., 2012). In addition to MAPK-related
pathways, factors involved in Rho GTPAses signaling were also detected among up- and down-
phosphorylated proteins following infection, thus constituting an important signature. Rho GTPases
control the actin cytoskeleton and, therefore, cell mobility, shape, and migration (Lawson and Ridley,
2018). Interestingly, several effector proteins involved in the Rho GTPase signaling pathway and
related to cytoskeletal organization, were also found in the phosphoproteomic analysis of SARS-
CoV2-infected cells (Bouhaddou et al., 2020). A previous study, conducted by transfecting a plasmid
containing the HBV genome in dividing HepG2 cells, reported that viral replication can induce
morphological changes, activate Rac1 and, downstream, trigger the phosphorylation of ERK1 and
AKT (Tan et al., 2008). Whether similar modifications can occur in more physiological infectious
model is still unknown. In our analysis, neither AKT- nor ERK1-derived up-phosphorylated peptides
were detected. Nevertheless, the finding that Rho GTPases signaling pathway was also highlighted in
the more stringent analysis performed on four different PHH donors, strongly suggests that HBV
infection may have an impact on the hepatocyte cytoskeleton and cell morphology.

Many studies documented the interaction between HBV and a plethora of RBPs which played a role

at every step of viral RNAs production, from transcription to translation (Heise et al., 1999;Ricco and

Kanduc, 2010;Sun et al., 2017;Yao et al., 2018;Makokha et al., 2019;Yao et al., 2019;Chabrolles et

al., 2020;Kanzaki et al., 2022;Yao et al., 2022). The HBc protein itself, which has a positively-

charged, intrinsically disordered C-terminal domain similar to that found in many cellular RBPs, can

also interact with a network of RBPs, some of which possess anti-viral activities {Diab, 2018

#2450;Yao et al., 2023b ;Zhang et al., 2023). In the present study, several RBPs were up-

phosphorylated upon HBV infection. Many if not all RBPs are tightly regulated by their

phosphorylation level which controls their intra-cellular and intra-nuclear localization, their capacity

to from condensates by liquid-liquid phase separation, their interaction with other protein partners, as

well their affinity for RNA and DNA (Hofweber and Dormann, 2019;Velazquez-Cruz et al., 2021;He

et al., 2023). In our study, we focused on two RBPs: first, HNRNPU, a DNA- and RNA-binding

protein, which was recently described as chromatin scaffold factor able to regulate cellular gene

expression, in particular by interacting with chromatin-associated non-coding RNAs (Sakaguchi et

al., 2016;Nozawa et al., 2017;Fan et al., 2018), as well as a factor able to negatively regulate viral

gene expression (Valente and Goff, 2006;Cao et al., 2019;Liu et al., 2021;Cao et al., 2022;Yang et

al., 2022). Both up- and down-phosphorylated peptides of HNRNPU were found in our study

(Supplemental Table 2). In particular, serine 271 is described as a major target residue

(https://www.phosphosite.org), whose phosphorylation is overrepresented during mitosis (Sharp et

al., 2020). Interestingly, PLK1 which can phosphorylate HNRNPU and is activated upon infection

(Douglas et al., 2015;Diab et al., 2017) was also up-phosphorylated in our first analysis performed

with a single-donor PHH. As for HNRNPU, the absence of detectable up-phosphorylated peptides in

the second analysis, performed with PHH from four different donors, may be the consequence of

strong variations among infections levels and/or donor-dependent features, in particular

phosphorylation/dephosphorylation kinetics. The second selected RBP was SRRM2. Initially

described as a component of the spliceosome and a member of the SR family of proteins, SRRM2

was recently identified as a main scaffold of nuclear speckles (Ilik et al., 2020). In particular, SRRM2

can form liquid condensates in a kinase-controlled fashion (Rai et al., 2018;Xu et al., 2022). This

RBP was also up-phosphorylated in HIV-1-infected cells, in which it regulated alternative splicing of

viral RNAs, as well as following infection of human macrophages with influenza A virus

(Wojcechowskyj et al., 2013;Soderholm et al., 2016). Our validation studies indicate that both,

HNRNPU and SRRM2 behave as antiviral factors that down regulate the level of HBV RNAs

without affecting cccDNA. As observed for SRSF10, an RBP which is part of the HBc interactome

(Chabrolles et al., 2020), it is likely that the anti-viral effect of these proteins may vary according to their phosphorylation state. In particular, phosphorylation of speckles proteins, such SC35 and SRRM2 by DYRK1A, DYRK3, and probably other kinases can dissolve liquid condensates formed by these speckles proteins, in particular during the onset of mitosis (Alvarez et al., 2003;Rai et al., 2018;Xu et al., 2022). Interestingly, in this study, DYRK1A figured among the kinases predicted to be activated at day 2-pi. Future investigation should focus on the capacity of selected RBPs to interact with HBV cccDNA and/or RNAs to further gain an insight into the mechanism involved in this antiviral effect. In addition, as for SRSF10, it will be important to investigate how HBV modulates the phosphorylation of the RBPs and, in particular, identify the cellular kinases involved.

The most remarkable finding of this study was the detection of up-phosphorylated proteins involved in DNA damage response and repair. Notably, among these proteins, DDB2 was previously involved in cccDNA formation (Marchetti et al., 2022). In addition, some major DNA repair proteins were identified in both phosphoproteomics analyses, namely Rad50 and 53BP1. In particular, Rad50, a cohesin-like component of the MRN complex, was up-phosphorylated at serine 635 a major ATM target (Kinoshita et al., 2009;Gatei et al., 2011). In the case of 53BP1, a highly phosphorylated DNA damage sensor (www.phosphosite.org), two different serine residues, previously reported to be up-phosphorylated upon DNA damage induced by ionizing or UV radiation, were detected in our analyses (Matsuoka et al., 2007;Boeing et al., 2016). 53BP1 is a critical regulator of the cellular response to DSBs (Panier and Boulton, 2014). Indeed, by associating to damaged chromatin, 53BP1 constitutes a barrier to DNA resection via the recruitment of a shieldin complex and the prevention of homologous recombination (HR) to favor direct end-joining (Setiaputra and Durocher, 2019;Callen et al., 2020). When recruited to DSBs and phosphorylated, in particular by the ATM kinase, 53BP1 forms large foci which are a typical signature of an ongoing DDR (Anderson et al., 2001;Kilic et al., 2019;Shibata and Jeggo, 2020). In our study, detectable 53BP1 foci, most of them positioned at the nuclear periphery, were observed in HBV-infected PHH, as soon as 1-dpi, strongly suggesting that their formation was induced upon virus disassembly and rcDNA delivery at the inner face of the nuclear lamina (Rabe et al., 2009). Indeed, rcDNA harbors many features which represent danger signals for the cell, such as ssDNA breaks, ssDNA regions and a covalently attached polymerase (Schreiner and Nassal, 2017). Because most viral DNA genomes, possess unconventional features, their release within the nucleus, frequently results in a DDR (Weitzman and Fradet-Turcotte, 2018). In line with this hypothesis, we found that 53BP1 knock-down increased cccDNA levels, strongly suggesting that this factor may prevent repair of rcDNA to produce a closed, ds DNA episome.

Interestingly, knock-down of RIF1, which recognizes phosphorylated 53BP1 and recruits the shieldin complex, also increased cccDNA levels, strongly suggesting that recruitment of this complex prevents repair of rcDNA (Setiaputra et al., 2022). In our analyses, 53BP1 foci were observed until 7-dpi, indicating a persistent DDR. Even if cccDNA formation is achieved in approximately 3 days in infected PHH (Locatelli et al., 2022), a residual level of rcDNA may still be present and responsible for the presence of such foci. Alternatively, nuclear recycling of newly synthetized rcDNA in the nucleus may be responsible for the induction of these long-lasting DDR. A previous report indicated that PML bodies are recruited to persistent DNA damage lesions by 53BP1 (Vancurova et al., 2019). In our analyses, most of HBV-induced 53BP1 foci colocalized with PML signal. Interestingly, some studies indicated that, in the absence of HBx, PML bodies are important for HBV transcriptional silencing by recruiting inhibitory factors and viral genomes (Niu et al., 2017;Li et al., 2022;Yao et al., 2023a). It would be interesting to perform immune-FISH analyses, able to discriminate between rc and cccDNA, to determine whether PML/53BP1 signals also associate to viral DNA or whether they represent an alternative nuclear anti-viral structure.

Few information is available on the interplay between HBV and the DDR, in particular with its three major kinases, ATM, ATR and DNA-PK (Blackford and Jackson, 2017). Initial reports indicated that HBV infection may activate ATR and some of its downstream targets, therefore suggesting a possible recognition of rcDNA as a damaged molecule (Zhao et al., 2008a;Zhao et al., 2008b). Later on, both ATM and ATR were described as enhancing HBV replication (Kostyusheva et al., 2019), and the ATR/Check1 pathway shown to enhance cccDNA formation (Luo et al., 2020a). More recently, DNA-PK was described to up-regulate HBV transcription (Fan et al., 2022). In our analyses, these three kinases were predicted to be activated upon HBV infection, as soon as 2- and up to 7-dpi. As already explored in some studies, this finding opens the perspective to investigate the effect of inhibitors of these kinases on the HBV life cycle and, in particular during the early phase, on cccDNA formation. Interestingly, Lubyova *et al.* (Lubyova et al., 2021) reported that HBc may be phosphorylated by ATM kinase following a genotoxic stress (Lubyova et al., 2021). It will be interesting to determine how kinase inhibitors targeting ATM or other DDR kinases modify the capacity of capsid-derived HBc to associate with cccDNA and how this affects down-streams events. Besides DDR kinases, CDK2, which plays a major role in HBc phosphorylation and is recruited and packaged within HBV capsids, was also on the top-ranked kinases activated upon infection (Ludgate et al., 2012;Luo et al., 2020b).

In conclusion, our deep MS-based phosphoproteomic analysis strongly suggests that HBV infection

triggers an intrinsic anti-viral response composed by DNA repair factors and RBPs that contribute to

reduce HBV establishment and productive replication (Figure 11). Understanding how HBV may

evade or counteract some of these responses will be critical to further pinpoint factors able prevent

cccDNA establishment and expression, as well to investigate further antiviral strategies based on the

use of kinase inhibitors.

5   AUTHOR CONTRIBUTIONS

533   FP, EC, LB, HC and MC: Investigation, Methodology, Data curation. TB: Methodology. JL:

534   Investigation, Writing – review & editing. DD: Funding acquisition,Writing – review & editing. MR

535   and GP: YC : Funding acquisition, Investigation, Methodology, Software: Data curation, Funding

536   acquisition, Writing – review & editing. AS: Conceptualization, Funding acquisition, Project

537   administration, Writing – original draft, Writing – review & editing.

6   FUNDING

This work was funded by Institut National de la Santé et de la Recherche Médicale (INSERM), the

Centre National de la Recherche Scientifique (CNRS), and Université Claude Bernard Lyon 1

(UCBL). It was also supported by grants from the Agence Nationale de Recherche sur le Sida et les

hépatites virales (ANRS, ECTZ11892) and fellowship to FP (ANRS, ECTZ119385). The proteomic

experiments were partially supported by Agence Nationale de la Recherche under projects ProFI

(Proteomics French Infrastructure, ANR-10-INBS-08) and GRAL, a program from the Chemistry

Biology Health (CBH) Graduate School of University Grenoble Alpes (ANR-17-EURE-0003).

7   CONFLICT OF INTEREST

549   The authors declare that the research was conducted in the absence of any commercial or financial

550   relationships that could be construed as a potential conflict of interest.

8   ACKNOWLEDGMENTS

We would like to thank Christophe Vanbelle (Imaging platform of CRCL), Ema Bobocioiu
(PLATIM, imaging Platform of SFR Biosciences) for their help on confocal microscope analyses,
Maud Michelet and Anëlle Dubois for their help for PHH isolation, as well as the staff from Pr.
Michel Rivoire's and Dr Guillaume Passot's surgery rooms for providing us with liver resections.
We also thank the Plateforme de Thérapie Génique in Nantes (France) for the production of AAV-
HBV vectors.

9   **REFERENCES**

Abdul, F., Diman, A., Baechler, B., Ramakrishnan, D., Kornyeyev, D., Beran, R.K., Fletcher, S.P.,
and Strubin, M. (2022). Smc5/6 silences episomal transcription by a three-step function. *Nat*
Struct Mol Biol 29**,** 922-931.

Allweiss, L., Testoni, B., Yu, M., Lucifora, J., Ko, C., Qu, B., Lutgehetmann, M., Guo, H., Urban, S.,
Fletcher, S.P., Protzer, U., Levrero, M., Zoulim, F., and Dandri, M. (2023). Quantification of
the hepatitis B virus cccDNA: evidence-based guidelines for monitoring the key obstacle of
HBV cure. *Gut*.

Alvarez, M., Estivill, X., and De La Luna, S. (2003). DYRK1A accumulates in splicing speckles
through a novel targeting signal and induces speckle disassembly. *J Cell Sci* 116**,** 3099-3107.

Anderson, L., Henderson, C., and Adachi, Y. (2001). Phosphorylation and rapid relocalization of
53BP1 to nuclear foci upon DNA damage. *Mol Cell Biol* 21**,** 1719-1729.

Arzumanian, V.A., Kiseleva, O.I., and Poverennaya, E.V. (2021). The Curious Case of the HepG2
Cell Line: 40 Years of Expertise. *Int J Mol Sci* 22.

Assiddiq, B.F., Tan, K.Y., Toy, W., Chan, S.P., Chong, P.K., and Lim, Y.P. (2012). EGFR S1166
phosphorylation induced by a combination of EGF and gefitinib has a potentially negative
impact on lung cancer cell growth. *J Proteome Res* 11**,** 4110-4119.

Blackford, A.N., and Jackson, S.P. (2017). ATM, ATR, and DNA-PK: The Trinity at the Heart of the
DNA Damage Response. *Mol Cell* 66**,** 801-817.

Blencowe, B.J., Bauren, G., Eldridge, A.G., Issner, R., Nickerson, J.A., Rosonina, E., and Sharp,
P.A. (2000). The SRm160/300 splicing coactivator subunits. *RNA* 6**,** 111-120.

Blondot, M.L., Bruss, V., and Kann, M. (2016). Intracellular transport and egress of hepatitis B virus.
J Hepatol 64**,** S49-59.

Boeing, S., Williamson, L., Encheva, V., Gori, I., Saunders, R.E., Instrell, R., Aygun, O., Rodriguez-
Martinez, M., Weems, J.C., Kelly, G.P., Conaway, J.W., Conaway, R.C., Stewart, A.,
Howell, M., Snijders, A.P., and Svejstrup, J.Q. (2016). Multiomic Analysis of the UV-
Induced DNA Damage Response. *Cell Rep* 15**,** 1597-1610.

Borgo, C., D'amore, C., Sarno, S., Salvi, M., and Ruzzene, M. (2021). Protein kinase CK2: a
potential therapeutic target for diverse human diseases. *Signal Transduct Target Ther* 6**,** 183.

Bouhaddou, M., Memon, D., Meyer, B., White, K.M., Rezelj, V.V., Correa Marrero, M., Polacco,
B.J., Melnyk, J.E., Ulferts, S., Kaake, R.M., Batra, J., Richards, A.L., Stevenson, E., Gordon,
D.E., Rojc, A., Obernier, K., Fabius, J.M., Soucheray, M., Miorin, L., Moreno, E., Koh, C.,
Tran, Q.D., Hardy, A., Robinot, R., Vallet, T., Nilsson-Payant, B.E., Hernandez-Armenta, C.,

Dunham, A., Weigang, S., Knerr, J., Modak, M., Quintero, D., Zhou, Y., Dugourd, A., Valdeolivas, A., Patil, T., Li, Q., Huttenhain, R., Cakir, M., Muralidharan, M., Kim, M., Jang, G., Tutuncuoglu, B., Hiatt, J., Guo, J.Z., Xu, J., Bouhaddou, S., Mathy, C.J.P., Gaulton, A., Manners, E.J., Felix, E., Shi, Y., Goff, M., Lim, J.K., Mcbride, T., O'neal, M.C., Cai, Y., Chang, J.C.J., Broadhurst, D.J., Klippsten, S., De Wit, E., Leach, A.R., Kortemme, T., Shoichet, B., Ott, M., Saez-Rodriguez, J., Tenoever, B.R., Mullins, R.D., Fischer, E.R., Kochs, G., Grosse, R., Garcia-Sastre, A., Vignuzzi, M., Johnson, J.R., Shokat, K.M., Swaney, D.L., Beltrao, P., and Krogan, N.J. (2020). The Global Phosphorylation Landscape of SARS-CoV-2 Infection. *Cell* 182**,** 685-712 e619.

Callen, E., Zong, D., Wu, W., Wong, N., Stanlie, A., Ishikawa, M., Pavani, R., Dumitrache, L.C., Byrum, A.K., Mendez-Dorantes, C., Martinez, P., Canela, A., Maman, Y., Day, A., Kruhlak, M.J., Blasco, M.A., Stark, J.M., Mosammaparast, N., Mckinnon, P.J., and Nussenzweig, A. (2020). 53BP1 Enforces Distinct Pre- and Post-resection Blocks on Homologous Recombination. *Mol Cell* 77**,** 26-38 e27.

Cao, L., Liu, S., Li, Y., Yang, G., Luo, Y., Li, S., Du, H., Zhao, Y., Wang, D., Chen, J., Zhang, Z., Li, M., Ouyang, S., Gao, X., Sun, Y., Wang, Z., Yang, L., Lin, R., Wang, P., and You, F. (2019). The Nuclear Matrix Protein SAFA Surveils Viral RNA and Facilitates Immunity by Activating Antiviral Enhancers and Super-enhancers. *Cell Host Microbe* 26**,** 369-384 e368.

Cao, L., Luo, Y., Guo, X., Liu, S., Li, S., Li, J., Zhang, Z., Zhao, Y., Zhang, Q., Gao, F., Ji, X., Gao, X., Li, Y., and You, F. (2022). SAFA facilitates chromatin opening of immune genes through interacting with anti-viral host RNAs. *PLoS Pathog* 18**,** e1010599.

Caudron-Herger, M., Jansen, R.E., Wassmer, E., and Diederichs, S. (2021). RBP2GO: a comprehensive pan-species database on RNA-binding proteins, their interactions and functions. *Nucleic Acids Res* 49**,** D425-D436.

Chabrolles, H., Auclair, H., Vegna, S., Lahlali, T., Pons, C., Michelet, M., Coute, Y., Belmudes, L., Chadeuf, G., Kim, Y., Di Bernardo, A., Jalaguier, P., Cosset, F.L., Fusil, F., Rivoire, M., Arnold, L.D., Lopatin, U., Combet, C., Zoulim, F., Grierson, D., Chabot, B., Lucifora, J., Durantel, D., and Salvetti, A. (2020). Hepatitis B virus Core protein nuclear interactome identifies SRSF10 as a host RNA-binding protein restricting HBV RNA production. *PLoS Pathog* 16**,** e1008593.

De Chaumont, F., Dallongeville, S., Chenouard, N., Herve, N., Pop, S., Provoost, T., Meas-Yedid, V., Pankajakshan, P., Lecomte, T., Le Montagner, Y., Lagache, T., Dufour, A., and Olivo-Marin, J.C. (2012). Icy: an open bioimage informatics platform for extended reproducible research. *Nat Methods* 9**,** 690-696.

Decorsiere, A., Mueller, H., Van Breugel, P.C., Abdul, F., Gerossier, L., Beran, R.K., Livingston, C.M., Niu, C., Fletcher, S.P., Hantz, O., and Strubin, M. (2016). Hepatitis B virus X protein identifies the Smc5/6 complex as a host restriction factor. *Nature* 531**,** 386-389.

Diab, A., Foca, A., Fusil, F., Lahlali, T., Jalaguier, P., Amirache, F., N'guyen, L., Isorce, N., Cosset, F.L., Zoulim, F., Andrisani, O., and Durantel, D. (2017). Polo-like-kinase 1 is a proviral host factor for hepatitis B virus replication. *Hepatology* 66**,** 1750-1765.

Diogo Dias, J., Sarica, N., and Neuveut, C. (2021). Early Steps of Hepatitis B Life Cycle: From Capsid Nuclear Import to cccDNA Formation. *Viruses* 13.

Dion, S., Bourgine, M., Godon, O., Levillayer, F., and Michel, M.L. (2013). Adeno-associated virus-
mediated gene transfer leads to persistent hepatitis B virus replication in mice expressing
HLA-A2 and HLA-DR1 molecules. *J Virol* 87**,** 5554-5563.

Douglas, P., Ye, R., Morrice, N., Britton, S., Trinkle-Mulcahy, L., and Lees-Miller, S.P. (2015).
Phosphorylation of SAF-A/hnRNP-U Serine 59 by Polo-Like Kinase 1 Is Required for
Mitosis. *Mol Cell Biol* 35**,** 2699-2713.

Fan, H., Lv, P., Huo, X., Wu, J., Wang, Q., Cheng, L., Liu, Y., Tang, Q.Q., Zhang, L., Zhang, F.,
Zheng, X., Wu, H., and Wen, B. (2018). The nuclear matrix protein HNRNPU maintains 3D
genome architecture globally in mouse hepatocytes. *Genome Res* 28**,** 192-202.

Fan, Y., Liang, Y., Liu, Y., and Fan, H. (2022). PRKDC promotes hepatitis B virus transcription
through enhancing the binding of RNA Pol II to cccDNA. *Cell Death Dis* 13**,** 404.

Fanning, G.C., Zoulim, F., Hou, J., and Bertoletti, A. (2019). Therapeutic strategies for hepatitis B
virus infection: towards a cure. *Nat Rev Drug Discov*.

Fukano, K., Oshima, M., Tsukuda, S., Aizaki, H., Ohki, M., Park, S.Y., Wakita, T., Wakae, K.,
Watashi, K., and Muramatsu, M. (2021). NTCP Oligomerization Occurs Downstream of the
NTCP-EGFR Interaction during Hepatitis B Virus Internalization. *J Virol* 95**,** e0093821.

Garcia-Moreno, M., Jarvelin, A.I., and Castello, A. (2018). Unconventional RNA-binding proteins
step into the virus-host battlefront. *Wiley Interdiscip Rev RNA* 9**,** e1498.

Gatei, M., Jakob, B., Chen, P., Kijas, A.W., Becherel, O.J., Gueven, N., Birrell, G., Lee, J.H., Paull,
T.T., Lerenthal, Y., Fazry, S., Taucher-Scholz, G., Kalb, R., Schindler, D., Waltes, R., Dork,
T., and Lavin, M.F. (2011). ATM protein-dependent phosphorylation of Rad50 protein
regulates DNA repair and cell cycle control. *J Biol Chem* 286**,** 31542-31556.

Girardi, E., Pfeffer, S., Baumert, T.F., and Majzoub, K. (2021). Roadblocks and fast tracks: How
RNA binding proteins affect the viral RNA journey in the cell. *Semin Cell Dev Biol* 111**,** 86-
100.

Gripon, P., Rumin, S., Urban, S., Le Seyec, J., Glaise, D., Cannie, I., Guyomard, C., Lucas, J., Trepo,
C., and Guguen-Guillouzo, C. (2002). Infection of a human hepatoma cell line by hepatitis B
virus. *Proc Natl Acad Sci U S A* 99**,** 15655-15660.

Guillemin, A., Kumar, A., Wencker, M., and Ricci, E.P. (2021). Shaping the Innate Immune
Response Through Post-Transcriptional Regulation of Gene Expression Mediated by RNA-
Binding Proteins. *Front Immunol* 12**,** 796012.

He, S., Valkov, E., Cheloufi, S., and Murn, J. (2023). The nexus between RNA-binding proteins and
their effectors. *Nat Rev Genet* 24**,** 276-294.

Heise, T., Guidotti, L.G., and Chisari, F.V. (1999). La autoantigen specifically recognizes a predicted
stem-loop in hepatitis B virus RNA. *J Virol* 73**,** 5767-5776.

Hofweber, M., and Dormann, D. (2019). Friend or foe-Post-translational modifications as regulators
of phase separation and RNP granule dynamics. *J Biol Chem* 294**,** 7137-7150.

Ilik, I.A., Malszycki, M., Lubke, A.K., Schade, C., Meierhofer, D., and Aktas, T. (2020). SON and
SRRM2 are essential for nuclear speckle formation. *Elife* 9.

Iwamoto, M., Saso, W., Nishioka, K., Ohashi, H., Sugiyama, R., Ryo, A., Ohki, M., Yun, J.H., Park,
S.Y., Ohshima, T., Suzuki, R., Aizaki, H., Muramatsu, M., Matano, T., Iwami, S., Sureau, C.,
Wakita, T., and Watashi, K. (2020). The machinery for endocytosis of epidermal growth

factor receptor coordinates the transport of incoming hepatitis B virus to the endosomal
network. *J Biol Chem* 295**,** 800-807.

Iwamoto, M., Saso, W., Sugiyama, R., Ishii, K., Ohki, M., Nagamori, S., Suzuki, R., Aizaki, H., Ryo,
A., Yun, J.H., Park, S.Y., Ohtani, N., Muramatsu, M., Iwami, S., Tanaka, Y., Sureau, C.,
Wakita, T., and Watashi, K. (2019). Epidermal growth factor receptor is a host-entry cofactor
triggering hepatitis B virus internalization. *Proc Natl Acad Sci U S A* 116**,** 8487-8492.

Jenke, A.C., Stehle, I.M., Herrmann, F., Eisenberger, T., Baiker, A., Bode, J., Fackelmayer, F.O., and
Lipps, H.J. (2004). Nuclear scaffold/matrix attached region modules linked to a transcription
unit are sufficient for replication and maintenance of a mammalian episome. *Proc Natl Acad
Sci U S A* 101**,** 11322-11327.

Jenke, B.H., Fetzer, C.P., Stehle, I.M., Jonsson, F., Fackelmayer, F.O., Conradt, H., Bode, J., and
Lipps, H.J. (2002). An episomally replicating vector binds to the nuclear matrix protein SAF-
A in vivo. *EMBO Rep* 3**,** 349-354.

Jowsey, P., Morrice, N.A., Hastie, C.J., Mclauchlan, H., Toth, R., and Rouse, J. (2007).
Characterisation of the sites of DNA damage-induced 53BP1 phosphorylation catalysed by
ATM and ATR. *DNA Repair (Amst)* 6**,** 1536-1544.

Justice, J.L., and Cristea, I.M. (2022). Nuclear antiviral innate responses at the intersection of DNA
sensing and DNA repair. *Trends Microbiol* 30**,** 1056-1071.

Kanzaki, H., Chiba, T., Kaneko, T., Ao, J., Kan, M., Muroyama, R., Nakamoto, S., Kanda, T.,
Maruyama, H., Kato, J., Zen, Y., Kotani, A., Sekiba, K., Otsuka, M., Ohtsuka, M., and Kato,
N. (2022). The RNA-Binding Protein ELAVL1 Regulates Hepatitis B Virus Replication and
Growth of Hepatocellular Carcinoma Cells. *Int J Mol Sci* 23.

Kieffer, S.R., and Lowndes, N.F. (2022). Immediate-Early, Early, and Late Responses to DNA
Double Stranded Breaks. *Front Genet* 13**,** 793884.

Kiledjian, M., and Dreyfuss, G. (1992). Primary structure and binding activity of the hnRNP U
protein: binding RNA through RGG box. *EMBO J* 11**,** 2655-2664.

Kilic, S., Lezaja, A., Gatti, M., Bianco, E., Michelena, J., Imhof, R., and Altmeyer, M. (2019). Phase
separation of 53BP1 determines liquid-like behavior of DNA repair compartments. *EMBO J*
38**,** e101379.

Kinoshita, E., Van Der Linden, E., Sanchez, H., and Wyman, C. (2009). RAD50, an SMC family
member with multiple roles in DNA break repair: how does ATP affect function?
Chromosome Res 17**,** 277-288.

Kostyusheva, A., Brezgin, S., Bayurova, E., Gordeychuk, I., Isaguliants, M., Goptar, I., Urusov, F.,
Nikiforova, A., Volchkova, E., Kostyushev, D., and Chulanov, V. (2019). ATM and ATR
Expression Potentiates HBV Replication and Contributes to Reactivation of HBV Infection
upon DNA Damage. *Viruses* 11.

Ladner, S.K., Otto, M.J., Barker, C.S., Zaifert, K., Wang, G.H., Guo, J.T., Seeger, C., and King,
R.W. (1997). Inducible expression of human hepatitis B virus (HBV) in stably transfected
hepatoblastoma cells: a novel system for screening potential inhibitors of HBV replication.
Antimicrob Agents Chemother 41**,** 1715-1720.

Lawson, C.D., and Ridley, A.J. (2018). Rho GTPase signaling complexes in cell migration and
invasion. *J Cell Biol* 217**,** 447-457.

Lecluyse, E.L., and Alexandre, E. (2010). Isolation and culture of primary hepatocytes from resected human liver tissue. *Methods Mol Biol* 640**,** 57-82.

Li, Y., He, M., Gong, R., Wang, Z., Lu, L., Peng, S., Duan, Z., Feng, Y., Liu, Y., and Gao, B. (2022). Forkhead O Transcription Factor 4 Restricts HBV Covalently Closed Circular DNA Transcription and HBV Replication through Genetic Downregulation of Hepatocyte Nuclear Factor 4 Alpha and Epigenetic Suppression of Covalently Closed Circular DNA via Interacting with Promyelocytic Leukemia Protein. *J Virol* 96**,** e0054622.

Lim, Z., Mohd-Ismail, N.K.B., Png, E., Sze, C.W., Lin, Q., Hong, W., Lim, S.G., Tan, Y.J., and Gunaratne, J. (2022). Phosphoproteomics Unravel HBV Triggered Rewiring of Host Phosphosignaling Events. *Int J Mol Sci* 23.

Lisy, S., Rothamel, K., and Ascano, M. (2021). RNA Binding Proteins as Pioneer Determinants of Infection: Protective, Proviral, or Both? *Viruses* 13.

Liu, B.Y., Yu, X.J., and Zhou, C.M. (2021). SAFA initiates innate immunity against cytoplasmic RNA virus SFTSV infection. *PLoS Pathog* 17**,** e1010070.

Locatelli, M., Quivy, J.P., Chapus, F., Michelet, M., Fresquet, J., Maadadi, S., Aberkane, A.N., Diederichs, A., Lucifora, J., Rivoire, M., Almouzni, G., Testoni, B., and Zoulim, F. (2022). HIRA Supports Hepatitis B Virus Minichromosome Establishment and Transcriptional Activity in Infected Hepatocytes. *Cell Mol Gastroenterol Hepatol* 14**,** 527-551.

Lopez, A., Nichols Doyle, R., Sandoval, C., Nisson, K., Yang, V., and Fregoso, O.I. (2022). Viral Modulation of the DNA Damage Response and Innate Immunity: Two Sides of the Same Coin. *J Mol Biol* 434**,** 167327.

Luangsay, S., Gruffaz, M., Isorce, N., Testoni, B., Michelet, M., Faure-Dupuy, S., Maadadi, S., Ait-Goughoulte, M., Parent, R., Rivoire, M., Javanbakht, H., Lucifora, J., Durantel, D., and Zoulim, F. (2015). Early inhibition of hepatocyte innate responses by hepatitis B virus. *J Hepatol* 63**,** 1314-1322.

Lubyova, B., Tikalova, E., Krulova, K., Hodek, J., Zabransky, A., Hirsch, I., and Weber, J. (2021). ATM-Dependent Phosphorylation of Hepatitis B Core Protein in Response to Genotoxic Stress. *Viruses* 13.

Lucifora, J., Arzberger, S., Durantel, D., Belloni, L., Strubin, M., Levrero, M., Zoulim, F., Hantz, O., and Protzer, U. (2011). Hepatitis B virus X protein is essential to initiate and maintain virus replication after infection. *J Hepatol* 55**,** 996-1003.

Lucifora, J., Michelet, M., Rivoire, M., Protzer, U., Durantel, D., and Zoulim, F. (2020). Two-dimensional-cultures of primary human hepatocytes allow efficient HBV infection: Old tricks still work! *J Hepatol* 73**,** 449-451.

Lucifora, J., Pastor, F., Charles, E., Pons, C., Auclair, H., Fusil, F., Rivoire, M., Cosset, F.L., Durantel, D., and Salvetti, A. (2021). Evidence for long-term association of virion-delivered HBV core protein with cccDNA independently of viral protein production. *JHEP Rep* 3**,** 100330.

Lucifora, J., Xia, Y., Reisinger, F., Zhang, K., Stadler, D., Cheng, X., Sprinzl, M.F., Koppensteiner, H., Makowska, Z., Volz, T., Remouchamps, C., Chou, W.M., Thasler, W.E., Huser, N., Durantel, D., Liang, T.J., Munk, C., Heim, M.H., Browning, J.L., Dejardin, E., Dandri, M., Schindler, M., Heikenwalder, M., and Protzer, U. (2014). Specific and nonhepatotoxic degradation of nuclear hepatitis B virus cccDNA. *Science* 343**,** 1221-1228.

Ludgate, L., Ning, X., Nguyen, D.H., Adams, C., Mentzer, L., and Hu, J. (2012). Cyclin-dependent
kinase 2 phosphorylates s/t-p sites in the hepadnavirus core protein C-terminal domain and is
incorporated into viral capsids. *J Virol* 86**,** 12237-12250.

Luo, J., Luckenbaugh, L., Hu, H., Yan, Z., Gao, L., and Hu, J. (2020a). Involvement of Host ATR-
CHK1 Pathway in Hepatitis B Virus Covalently Closed Circular DNA Formation. *mBio* 11.

Luo, J., Xi, J., Gao, L., and Hu, J. (2020b). Role of Hepatitis B virus capsid phosphorylation in
nucleocapsid disassembly and covalently closed circular DNA formation. *PLoS Pathog* 16**,**
e1008459.

Ma, R., Li, S., Li, W., Yao, L., Huang, H.D., and Lee, T.Y. (2023). KinasePhos 3.0: Redesign and
Expansion of the Prediction on Kinase-specific Phosphorylation Sites. *Genomics Proteomics*
Bioinformatics 21**,** 228-241.

Makokha, G.N., Abe-Chayama, H., Chowdhury, S., Hayes, C.N., Tsuge, M., Yoshima, T., Ishida, Y.,
Zhang, Y., Uchida, T., Tateno, C., Akiyama, R., and Chayama, K. (2019). Regulation of the
Hepatitis B virus replication and gene expression by the multi-functional protein TARDBP.
Sci Rep 9**,** 8462.

Marchetti, A.L., Zhang, H., Kim, E.S., Yu, X., Jang, S., Wang, M., and Guo, H. (2022). Proteomic
Analysis of Nuclear Hepatitis B Virus Relaxed Circular DNA-Associated Proteins Identifies
UV-Damaged DNA Binding Protein as a Host Factor Involved in Covalently Closed Circular
DNA Formation. *J Virol* 96**,** e0136021.

Matsuoka, S., Ballif, B.A., Smogorzewska, A., Mcdonald, E.R., 3rd, Hurov, K.E., Luo, J.,
Bakalarski, C.E., Zhao, Z., Solimini, N., Lerenthal, Y., Shiloh, Y., Gygi, S.P., and Elledge,
S.J. (2007). ATM and ATR substrate analysis reveals extensive protein networks responsive
to DNA damage. *Science* 316**,** 1160-1166.

Murphy, C.M., Xu, Y., Li, F., Nio, K., Reszka-Blanco, N., Li, X., Wu, Y., Yu, Y., Xiong, Y., and Su,
L. (2016). Hepatitis B Virus X Protein Promotes Degradation of SMC5/6 to Enhance HBV
Replication. *Cell Rep* 16**,** 2846-2854.

Mutz, P., Metz, P., Lempp, F.A., Bender, S., Qu, B., Schoneweis, K., Seitz, S., Tu, T., Restuccia, A.,
Frankish, J., Dachert, C., Schusser, B., Koschny, R., Polychronidis, G., Schemmer, P.,
Hoffmann, K., Baumert, T.F., Binder, M., Urban, S., and Bartenschlager, R. (2018). HBV
Bypasses the Innate Immune Response and Does Not Protect HCV From Antiviral Activity of
Interferon. *Gastroenterology* 154**,** 1791-1804 e1722.

Niu, C., Livingston, C.M., Li, L., Beran, R.K., Daffis, S., Ramakrishnan, D., Burdette, D., Peiser, L.,
Salas, E., Ramos, H., Yu, M., Cheng, G., Strubin, M., Delaney, W.I., and Fletcher, S.P.
(2017). The Smc5/6 Complex Restricts HBV when Localized to ND10 without Inducing an
Innate Immune Response and Is Counteracted by the HBV X Protein Shortly after Infection.
PLoS One 12**,** e0169648.

Norton, P.A., Reis, H.M., Prince, S., Larkin, J., Pan, J., Liu, J., Gong, Q., Zhu, M., and Feitelson,
M.A. (2004). Activation of fibronectin gene expression by hepatitis B virus x antigen. *J Viral*
Hepat 11**,** 332-341.

Nozawa, R.S., Boteva, L., Soares, D.C., Naughton, C., Dun, A.R., Buckle, A., Ramsahoye, B.,
Bruton, P.C., Saleeb, R.S., Arnedo, M., Hill, B., Duncan, R.R., Maciver, S.K., and Gilbert, N.
(2017). SAF-A Regulates Interphase Chromosome Structure through Oligomerization with
Chromatin-Associated RNAs. *Cell* 169**,** 1214-1227 e1218.

Panier, S., and Boulton, S.J. (2014). Double-strand break repair: 53BP1 comes into focus. *Nat Rev*
Mol Cell Biol 15**,** 7-18.

Pastor, F., Shkreta, L., Chabot, B., Durantel, D., and Salvetti, A. (2021). Interplay Between CMGC
Kinases Targeting SR Proteins and Viral Replication: Splicing and Beyond. *Front Microbiol*
12**,** 658721.

Perez-Riverol, Y., Bai, J., Bandla, C., Garcia-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S.,
Kundu, D.J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., Walzer, M., Wang, S.,
Brazma, A., and Vizcaino, J.A. (2022). The PRIDE database resources in 2022: a hub for
mass spectrometry-based proteomics evidences. *Nucleic Acids Res* 50**,** D543-D552.

Prange, R. (2022). Hepatitis B virus movement through the hepatocyte: An update. *Biol Cell* 114**,**
325-348.

Rabe, B., Delaleau, M., Bischof, A., Foss, M., Sominskaya, I., Pumpens, P., Cazenave, C.,
Castroviejo, M., and Kann, M. (2009). Nuclear entry of hepatitis B virus capsids involves
disintegration to protein dimers followed by nuclear reassociation to capsids. *PLoS Pathog* 5**,**
e1000563.

Rahman, N., Sun, J., Li, Z., Pattnaik, A., Mohallem, R., Wang, M., Kazemian, M., Aryal, U.K., and
Andrisani, O. (2022). The cytoplasmic LSm1-7 and nuclear LSm2-8 complexes exert
opposite effects on Hepatitis B virus biosynthesis and interferon responses. *Front Immunol*
13**,** 970130.

Rai, A.K., Chen, J.X., Selbach, M., and Pelkmans, L. (2018). Kinase-controlled phase transition of
membraneless organelles in mitosis. *Nature* 559**,** 211-216.

Rass, E., Willaume, S., and Bertrand, P. (2022). 53BP1: Keeping It under Control, Even at a Distance
from DNA Damage. *Genes (Basel)* 13.

Ren, S., Wang, J., Chen, T.L., Li, H.Y., Wan, Y.S., Peng, N.F., Gui, X.E., and Zhu, Y. (2016).
Hepatitis B Virus Stimulated Fibronectin Facilitates Viral Maintenance and Replication
through Two Distinct Mechanisms. *PLoS One* 11**,** e0152721.

Ricco, R., and Kanduc, D. (2010). Hepatitis B virus and Homo sapiens proteome-wide analysis: A
profusion of viral peptide overlaps in neuron-specific human proteins. *Biologics* 4**,** 75-81.

Romig, H., Fackelmayer, F.O., Renz, A., Ramsperger, U., and Richter, A. (1992). Characterization of
SAF-A, a novel nuclear DNA binding protein from HeLa cells with high affinity for nuclear
matrix/scaffold attachment DNA elements. *EMBO J* 11**,** 3431-3440.

Sakaguchi, T., Hasegawa, Y., Brockdorff, N., Tsutsui, K., Tsutsui, K.M., Sado, T., and Nakagawa, S.
(2016). Control of Chromosomal Localization of Xist by hnRNP U Family Molecules. *Dev*
Cell 39**,** 11-12.

Salvetti, A., Coute, Y., Epstein, A., Arata, L., Kraut, A., Navratil, V., Bouvet, P., and Greco, A.
(2016). Nuclear Functions of Nucleolin through Global Proteomics and Interactomic
Approaches. *J Proteome Res* 15**,** 1659-1669.

Salvetti, A., Oreve, S., Chadeuf, G., Favre, D., Cherel, Y., Champion-Arnaud, P., David-Ameline, J.,
and Moullier, P. (1998). Factors influencing recombinant adeno-associated virus production.
Human Gene Therapy 9**,** 695-706.

Schreiner, S., and Nassal, M. (2017). A Role for the Host DNA Damage Response in Hepatitis B
Virus cccDNA Formation-and Beyond? *Viruses* 9.

Setiaputra, D., and Durocher, D. (2019). Shieldin - the protector of DNA ends. *EMBO Rep* 20.

Setiaputra, D., Escribano-Diaz, C., Reinert, J.K., Sadana, P., Zong, D., Callen, E., Sifri, C.,
Seebacher, J., Nussenzweig, A., Thoma, N.H., and Durocher, D. (2022). RIF1 acts in DNA
repair through phosphopeptide recognition of 53BP1. *Mol Cell* 82, 1359-1371 e1359.

Sharp, J.A., Perea-Resa, C., Wang, W., and Blower, M.D. (2020). Cell division requires RNA
eviction from condensing chromosomes. *J Cell Biol* 219.

Shibata, A., and Jeggo, P.A. (2020). Roles for 53BP1 in the repair of radiation-induced DNA double
strand breaks. *DNA Repair (Amst)* 93, 102915.

Soderholm, S., Kainov, D.E., Ohman, T., Denisova, O.V., Schepens, B., Kulesskiy, E., Imanishi,
S.Y., Corthals, G., Hintsanen, P., Aittokallio, T., Saelens, X., Matikainen, S., and Nyman,
T.A. (2016). Phosphoproteomics to Characterize Host Response During Influenza A Virus
Infection of Human Macrophages. *Mol Cell Proteomics* 15, 3203-3219.

Song, Y., Li, M., Wang, Y., Zhang, H., Wei, L., and Xu, W. (2021). E3 ubiquitin ligase TRIM21
restricts hepatitis B virus replication by targeting HBx for proteasomal degradation. *Antiviral
Res* 192, 105107.

Sonntag, E., Milbradt, J., Svrlanska, A., Strojan, H., Hage, S., Kraut, A., Hesse, A.M., Amin, B.,
Sonnewald, U., Coute, Y., and Marschall, M. (2017). Protein kinases responsible for the
phosphorylation of the nuclear egress core complex of human cytomegalovirus. *J Gen Virol
98, 2569-2581.

Sun, S., Nakashima, K., Ito, M., Li, Y., Chida, T., Takahashi, H., Watashi, K., Sawasaki, T., Wakita,
T., and Suzuki, T. (2017). Involvement of PUF60 in Transcriptional and Post-transcriptional
Regulation of Hepatitis B Virus Pregenomic RNA Expression. *Sci Rep* 7, 12874.

Suslov, A., Boldanova, T., Wang, X., Wieland, S., and Heim, M.H. (2018). Hepatitis B Virus Does
Not Interfere With Innate Immune Responses in the Human Liver. *Gastroenterology* 154,
1778-1790.

Tan, T.L., Fang, N., Neo, T.L., Singh, P., Zhang, J., Zhou, R., Koh, C.G., Chan, V., Lim, S.G., and
Chen, W.N. (2008). Rac1 GTPase is activated by hepatitis B virus replication--involvement
of HBX. *Biochim Biophys Acta* 1783, 360-374.

Thompson, A., Schafer, J., Kuhn, K., Kienle, S., Schwarz, J., Schmidt, G., Neumann, T., Johnstone,
R., Mohammed, A.K., and Hamon, C. (2003). Tandem mass tags: a novel quantification
strategy for comparative analysis of complex protein mixtures by MS/MS. *Anal Chem* 75,
1895-1904.

Tsukuda, S., and Watashi, K. (2020). Hepatitis B virus biology and life cycle. *Antiviral Res* 182,
104925.

Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. *Nat Protoc* 11, 2301-2319.

Valente, S.T., and Goff, S.P. (2006). Inhibition of HIV-1 gene expression by a fragment of hnRNP
U. *Mol Cell* 23, 597-605.

Van Damme, E., Vanhove, J., Severyn, B., Verschueren, L., and Pauwels, F. (2021). The Hepatitis B
Virus Interactome: A Comprehensive Overview. *Front Microbiol* 12, 724877.

Vancurova, M., Hanzlikova, H., Knoblochova, L., Kosla, J., Majera, D., Mistrik, M., Burdova, K.,
Hodny, Z., and Bartek, J. (2019). PML nuclear bodies are recruited to persistent DNA

damage lesions in an RNF168-53BP1 dependent manner and contribute to DNA repair. *DNA*
Repair (Amst) 78**,** 114-127.

Velazquez-Cruz, A., Banos-Jaime, B., Diaz-Quintana, A., De La Rosa, M.A., and Diaz-Moreno, I.
(2021). Post-translational Control of RNA-Binding Proteins and Disease-Related
Dysregulation. *Front Mol Biosci* 8**,** 658852.

Wang, F., Song, H., Xu, F., Xu, J., Wang, L., Yang, F., Zhu, Y., and Tan, G. (2023). Role of hepatitis
B virus non-structural protein HBx on HBV replication, interferon signaling, and
hepatocarcinogenesis. *Front Microbiol* 14**,** 1322892.

Wei, L., and Ploss, A. (2021). Mechanism of Hepatitis B Virus cccDNA Formation. *Viruses* 13.

Weitzman, M.D., and Fradet-Turcotte, A. (2018). Virus DNA Replication and the Host DNA
Damage Response. *Annu Rev Virol* 5**,** 141-164.

Werle-Lapostolle, B., Bowden, S., Locarnini, S., Wursthorn, K., Petersen, J., Lau, G., Trepo, C.,
Marcellin, P., Goodman, Z., Delaney, W.E.T., Xiong, S., Brosgart, C.L., Chen, S.S., Gibbs,
C.S., and Zoulim, F. (2004). Persistence of cccDNA during the natural history of chronic
hepatitis B and decline during adefovir dipivoxil therapy. *Gastroenterology* 126**,** 1750-1758.

Wieczorek, S., Combes, F., Lazar, C., Giai Gianetto, Q., Gatto, L., Dorffer, A., Hesse, A.M., Coute,
Y., Ferro, M., Bruley, C., and Burger, T. (2017). DAPAR & ProStaR: software to perform
statistical analyses in quantitative discovery proteomics. *Bioinformatics* 33**,** 135-136.

Wojcechowskyj, J.A., Didigu, C.A., Lee, J.Y., Parrish, N.F., Sinha, R., Hahn, B.H., Bushman, F.D.,
Jensen, S.T., Seeholzer, S.H., and Doms, R.W. (2013). Quantitative phosphoproteomics
reveals extensive cellular reprogramming during HIV-1 entry. *Cell Host Microbe* 13**,** 613-
623.

Xu, S., Lai, S.K., Sim, D.Y., Ang, W.S.L., Li, H.Y., and Roca, X. (2022). SRRM2 organizes splicing
condensates to regulate alternative splicing. *Nucleic Acids Res* 50**,** 8599-8614.

Yang, S.F., Chang, C.W., Wei, R.J., Shiue, Y.L., Wang, S.N., and Yeh, Y.T. (2014). Involvement of
DNA damage response pathways in hepatocellular carcinoma. *Biomed Res Int* 2014**,** 153867.

Yang, X., Zhao, X., Zhu, Y., Xun, J., Wen, Q., Pan, H., Yang, J., Wang, J., Liang, Z., Shen, X.,
Liang, Y., Lin, Q., Liang, H., Li, M., Chen, J., Jiang, S., Xu, J., Lu, H., and Zhu, H. (2022).
FBXO34 promotes latent HIV-1 activation by post-transcriptional modulation. *Emerg*
Microbes Infect 11**,** 2785-2799.

Yao, Q., Peng, B., Li, C., Li, X., Chen, M., Zhou, Z., Tang, D., He, J., Wu, Y., Sun, Y., and Li, W.
(2023a). SLF2 Interacts with the SMC5/6 Complex to Direct Hepatitis B Virus Episomal
DNA to Promyelocytic Leukemia Bodies for Transcriptional Repression. *J Virol* 97**,**
e0032823.

Yao, Y., Yang, B., Cao, H., Zhao, K., Yuan, Y., Chen, Y., Zhang, Z., Wang, Y., Pei, R., Chen, J.,
Hu, X., Zhou, Y., Lu, M., Wu, C., and Chen, X. (2018). RBM24 stabilizes hepatitis B virus
pregenomic RNA but inhibits core protein translation by targeting the terminal redundancy
sequence. *Emerg Microbes Infect* 7**,** 86.

Yao, Y., Yang, B., Chen, Y., Huang, D., Liu, C., Sun, H., Hu, X., Zhou, Y., Wang, Y., Chen, J., Pei,
R., Wen, Z., and Chen, X. (2022). RNA-Binding motif protein 38 (RBM38) mediates HBV
pgRNA packaging into the nucleocapsid. *Antiviral Res* 198**,** 105249.

Yao, Y., Yang, B., Chen, Y., Wang, H., Hu, X., Zhou, Y., Gao, X., Lu, M., Niu, J., Wen, Z., Wu, C.,
and Chen, X. (2019). RNA-Binding Motif Protein 24 (RBM24) Is Involved in Pregenomic

RNA Packaging by Mediating Interaction between Hepatitis B Virus Polymerase and the
Epsilon Element. *J Virol* 93.

Yao, Y.X., Chen, Y., Huang, D., Liu, C., Sun, H., Zhou, Y., Pei, R., Wang, Y., Wen, Z., Yang, B.,
and Chen, X. (2023b). RNA-binding motif protein 24 inhibits HBV replication in vivo. *J Med*
Virol 95**,** e28969.

Zhang, T., Zheng, H., Lu, D., Guan, G., Li, D., Zhang, J., Liu, S., Zhao, J., Guo, J.T., Lu, F., and
Chen, X. (2023). RNA binding protein TIAR modulates HBV replication by tipping the
balance of pgRNA translation. *Signal Transduct Target Ther* 8**,** 346.

Zhao, F., Hou, N.B., Song, T., He, X., Zheng, Z.R., Ma, Q.J., Li, L., Zhang, Y.H., and Zhong, H.
(2008a). Cellular DNA repair cofactors affecting hepatitis B virus infection and replication.
World J Gastroenterol 14**,** 5059-5065.

Zhao, F., Hou, N.B., Yang, X.L., He, X., Liu, Y., Zhang, Y.H., Wei, C.W., Song, T., Li, L., Ma, Q.J.,
and Zhong, H. (2008b). Ataxia telangiectasia-mutated-Rad3-related DNA damage checkpoint
signaling pathway triggered by hepatitis B virus infection. *World J Gastroenterol* 14**,** 6163-
6170.

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C., and
Chanda, S.K. (2019). Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* 10**,** 1523.

10 **FIGURE CAPTIONS**

Figure 1. A. Outline of the experimental procedure. PHH purified form liver resection were infected
with HBV (MOI=500 vge/cell) overnight. After washing with medium, cells were kept in culture for
2 and 7 days, before direct on plate lysis and processing for MS and phospho-MS analyses. The
experiment was performed in triplicate using PHH from a unique donor. **B.** Quantification of HBe
and HBs antigens (Ag) levels in the supernatant of cells at 7-dpi. **C.** Volcano plots of up- and down-
regulated proteins (proteomic analysis) at 2-(upper panel) and 7-dpi (lower panel). **D.** Volcano plots
of up- and down-regulated phosphopeptides and phosphoproteins at 2-(upper panel) and 7-dpi (lower
panel).

Figure 2. A. Gene ontology clusters of formed by statistically enriched up-phosphorylated proteins at
2- and 7-pi. **B**. Protein-protein interaction (PPI) networks formed by merged up-phosphorylated
proteins at 2- and 7-dpi. All protein-protein interactions among input genes were extracted from PPI
data sources using Metascape and a network MCODE algorithm was applied to identify
neighborhoods where proteins are densely connected. Each MCODE network is assigned a unique
color. Green:R-HSA-5693571|Nonhomologous End-Joining (NHEJ); Red: R-HSA-8939211|ESR-
mediated signaling. Blue: unassigned. Grey: other interacting factors of the network.

Figure 3. A. Gene ontology clusters of formed by statistically enriched down-phosphorylated
proteins at 2- and 7-dpi. **B**. PPI networks formed by down-phosphorylated proteins at 2- and 7-dpi.
All protein-protein interactions among input genes were extracted from PPI data sources using
Metascape and a network MCODE algorithm was then applied to this network to identify
neighborhoods where proteins are densely connected. Each MCODE network is assigned a unique
color. MCODE annotation: Red: R-HSA-199991|Membrane Trafficking|-6.2;R-HSA-
5653656|Vesicle-mediated transport|-6.1;R-HSA-6807878|COPI-mediated anterograde transport|-4.1.
Blue: R-HSA-1227990|Signaling by ERBB2 in Cancer|-9.3;R-HSA-1227986|Signaling by ERBB2|-
8.4;R-HSA-177929|Signaling by EGFR|-8.4. green: no annotation. Grey: other interacting factors of
the network.

Figure 4. A. Network of enriched ontology clusters formed by up-phosphorylated proteins upon
HBV infection. A subset of representative terms from the full cluster was converted into a network
layout. Each term is represented by a circle node, where its size is proportional to the number of
input genes that fall under that term, and its color represent its cluster identity (i.e., nodes of the same

color belong to the same cluster). Terms with a similarity score > 0.3 are linked by an edge (the thickness of the edge represents the similarity score). The network is visualized with Cytoscape with "force-directed" layout and with edge bundled for clarity. **B**. and **C**. Protein-protein interaction networks formed by up-phosphorylated,proteins at 2- and 7-dpi involved in RNA binding (left panel), selected using the RBP2GO database (RBP2GO score above 40) (Caudron-Herger et al., 2021)and DNA repair (right panel), belonging to the NHEJ cluster. The physical interactions among up-phosphorylated proteins (red circles) were retrieved using by Cytoscape (v3.10.1) and the GeneMania application. Grey circles represent missing nodes used to build the interactome network.

Figure 5. Effect of SRRM2 and HNRNPU knock-down on HBV life cycle. A. Experimental outline. PHH (B and C) or dHepaRG (D and E) were infected with HBV (moi:500 vge/cell) and seven days later transfected with siRNA targeting HNRNPU, SRRM2 or control (CTL) siRNA. RNA and DNA were extracted from cells seven days after transfection to check the knock-down of cellular mRNA targets (**B** and **D**) and to quantify viral (**C** and **E**) nucleic acids. Results are expressed as the mean normalized ratio +/- SD, of at least 3 independent experiments, each performed in triplicate. Results in PHH were obtained from four experiments performed with four different PHH donors.

Figure 6. Visualization of 53BP1 foci in HBV-infected PHH. A. HBV-infected (HBV) or mock (NI)-infected PHH were fixed at indicated time points and analyzed by immunofluorescence using an anti-53BP1 antibody (green signal), at 1- and 3-dpi. The nucleus was stained with Hoescht (blue signal). Scale bar : 20 µm. **B.** Staining was similarly performed on cells infected with HBV in the presence or not of myrcludex (Myr) at 2-dpi. Scale bar : 10 µm. **C.** HBV-infected PHH, in the presence or not of myrcludex, were stained at 7-dpi using an anti-53BP1 (green signal) and an anti-HBc (red signal) antibody. Scale bar : 10 µm.

Figure 7. Co-labeling of 53BP1 and PML in HBV-infected PHH. HBV-infected PHH were fixed at 2-dpi and analyzed by immunofluorescence using an anti-53BP1 (red signal) and anti-PML (green signal) antibodies. The nucleus was stained with Hoescht (blue signal). Scale bar:10 µm. **B.** Enlarged merged views. **C.** 3D view of an hepatocytocyte nucleus showing the red (53BP1) signal surrounding the inner PML core (green signal).

Figure 8. Effect of 53BP1 knock-down on HBV-infected dHepaRG. A. Experimental outline. **B.** Western blot validation of 53BP1 knock-down. **C.** Quantification of cccDNA and total HBV RNA levels. Results are expressed as the mean normalized ratio +/- SD, of three experiments, each performed in triplicate.

Figure 9. A. Ontology clusters formed by statistically enriched up-phosphorylated proteins at 2- and 7-dpi. (N=4, from four different PHH donors). **B.** Physical interactions formed by proteins up-phosphorylated upon HBV infection. All protein-protein interactions among input genes were extracted from PPI data sources using Metascape. Red node names indicate proteins that were also found in the first phosphoproteomic analysis performed with PHH from a single donor.

Figure 10. Cellular kinases predicted to be involved in up-phosphorylation events during HBV infection. Up-hosphorylated peptides found at d2- and d7pi were analyzed using KInasePhos3.0 software (Ma et al., 2023). The probability scores from 1 to 0.9, assigned to each kinases, were summed up to retrieve a list of the most probable kinases involved at each time point. Red arrow designated the three major kinases involved in DNA repair.

Figure 11. Hypothetical model describing the intervention of up-phosphorylated cellular DNA repair and RBPs during the HBV nuclear steps. Several host factors participate in cccDNA formation and viral gene expression (Diogo Dias et al., 2021;Wei and Ploss, 2021). Our study suggests that upon binding to the nuclear pore, and release of rcDNA at the inner side of the nuclear membrane, rcDNA is recognized as an abnormal molecule and induces a DDR characterized by ATM, ATR, activation and formation of 53BP1/PML foci. In addition, several RBPs, such HNRNPU and SRRM2, also regulated by phosphorylation, further counteract HBV gene expression.

Figure 1

A

R-HSA-5693571: Nonhomologous End-Joining (NHEJ)
R-HSA-9679191: Potential therapeutics for SARS
R-HSA-3371497: HSP90 chaperone cycle for steroid hormone receptors (SHR) in the presence of ligand
R-HSA-372708: p130Cas linkage to MAPK signaling for integrins
R-HSA-72163: mRNA Splicing - Major Pathway
R-HSA-8957275: Post-translational protein phosphorylation
R-HSA-68875: Mitotic Prophase
R-HSA-194315: Signaling by Rho GTPases
R-HSA-5653656: Vesicle-mediated transport
R-HSA-75105: Fatty acyl-CoA biosynthesis

Figure 2

Figure 3

Figure 4.

Figure 5

A

HBV_1-dpi

NI_1-dpi

HBV_3-dpi

NI_3-dpi

B

HBV_2-dpi

HBV + Myr_2-dpi

C

HBV_7-dpi

HBV + Myr_7-dpi

Figure 6

A

HBV_2-dpi

10 um   10 um   10 um

B   **C**

10 um   10 um

Figure 7

Figure 7

Figure 8

A

B

Figure 9

HBV_Up_2dpi   HBV_Up_7dpi

Figure 10

Figure 11