Title: Increased functional integration of emotional control network in

late adulthood

Authors: Leona Rahel Bätz[1*], Shuer Ye[1], Xiaqing Lan[1], Maryam Ziaei[1,2,3*]

[1] Kavli Institute for Systems Neuroscience, Norwegian University of Science and Technology, Trondheim,

Norway

[2] Queensland Brain Institute, University of Queensland, Brisbane, Australia

[3] K.G. Jebsen Centre for Alzheimer's disease, Norwegian University of Science and Technology, Trondheim,

Norway

*Corresponding Author:

Email: leona.r.batz@ntnu.no, maryam.ziaei@ntnu.no

Abbreviations:

Prefrontal Cortex (PFC), meta-analytic groupings (MAGs), functional magnetic resonance imaging

(fMRI), Human Connectome Project (HCP), Cambridge Center for Aging and Neuroscience (Cam-

CAN), participation coefficient (PC), General additive model (GAM), default mode network (DMN),

Highlights:

- We aimed to identify age-related differences in the functional integration of large-scale emotion regulation brain networks across the adult lifespan, and its implications for well-being.
- Frontal emotion control network showed increased while interoception network showed decreased functional integration with higher age in the HCP-Aging dataset.
- Increased integration of emotion control network was replicated and validated in the Cam-CAN dataset.

Abstract

Across the adult lifespan, emotion regulation ability remains stable or even improves. The corresponding effects, however, in the emotion regulation networks in the brain remain underexplored. By utilizing large-scale datasets such as the Human Connectome Project (HCP-Aging, N=621, 349 females) and Cambridge Centre for Ageing and Neuroscience (Cam-CAN, N=333, 155 females), we were able to investigate how emotion regulation networks' functional topography differs across the entire adult lifespan. Based on previous meta-analytic work that identified four large-scale functional brain networks involved in emotion generation and regulation, we investigated the association between the integration of these emotion regulation networks and measures of mental wellbeing with age in the HCP-Aging dataset. We found an increase in the functional integration of the emotional control network among older adults, which was replicated using the Cam-CAN data set. Further we found that the network that is mediating emotion generative and regulative processes, and carries our introspective and reflective functions, is less integrated in higher age. Our study highlights the importance of identifying topological changes in the functional emotion network architecture across the lifespan, as it allows for a better understanding of functional brain network changes that accompany emotional aging.

Keywords: Functional integration, emotional aging, brain network

1 Introduction

Growing old is often accompanied by a range of cognitive, physical and behavioral changes (Spreng & Turner, 2019). The perception and regulation of emotion are also subject to these changes in late adulthood (Charles, 2010; Mather, 2012). However, while cognitive capabilities tend to decline, emotion regulation capabilities are typically maintained or even improved with age (Livingstone and Isaacowitz, 2021; Reed and Carstensen, 2011; Charles, 2010b; Reuter-Lorenz and Lustig, 2005). Longitudinal studies have revealed that aging is associated with overall improved emotional well-being and greater emotional stability (Carstensen et al., 2011). Moreover, empirical evidence has suggested that older adults are better at regulating their negative emotions (Urry & Gross, 2010), can select situations that avoid conflict or negative affect better, (Birditt et al., 2005; Charles et al., 2009) deploy more attention to positive emotions (Isaacowitz et al., 2006), and prioritize effective emotion regulation more than their younger counterparts (English & Carstensen, 2014; Lawton et al., 1992; Nolen-Hoeksema & Aldao, 2011). Older adults, therefore, maintain an absence of negative affect more consistently and can recover more quickly from negative emotional states compared to younger adults (Carstensen et al., 2000; Hay & Diehl, 2011). The Socioemotional Selectivity Theory explains this bias by stating that older adults shift their focus from future-oriented goals to present-oriented ones, with positive stimuli holding a more immediate appeal (Carstensen et al., 1999) while the selective engagement hypothesis postulates that due to older adults' limited cognitive capabilities compared to younger adults, they are more astute in expending these finite resources (Hess, 2014). Hence, they allocate their resources more discerningly, leading to better performance on tasks with personal relevance, significance, or social implications (Carstensen et al., 2011).

Accompanied by these motivational changes and their reflection in behavior, several changes are occurring at the neural level (MacCormack et al., 2020; Mather, 2016). Investigating such changes in the brain will help to better understand the motivational and behavioral changes observed across the lifespan. For instance, older adults exhibit increased activity of the prefrontal cortex (PFC) and reduced activity of the amygdala in comparison to younger adults in several tasks that require emotional processing (Mather, 2012; Nashiro et al., 2012; Ziaei et al., 2017), reflecting the potential inhibitory or regulatory control from the PFC to the amygdala. Emotion regulation is a complex process, and while some of the improved emotion regulation in later life was attributed to the increased recruitment of frontal areas during this process (Pessoa, 2008; St. Jacques et al., 2010; Tomasi & Volkow, 2012), the impact of this observed recruitment of frontal areas on the rest of the brain has yet to be fully understood. The complex interplay between major nodes in the affect system, including the PFC and amygdala, insula, anterior cingulate cortex, and ventral striatum is critical for better understanding, experiencing, and regulating emotions (Ahmed et al., 2015; Ghashghaei et al., 2007). When investigating the emotional response specifically on a network level, a subsystem of four

brain networks that are specifically involved in emotion generation and regulation has been identified in a comprehensive meta-analytic study, known as meta-analytic groupings (MAGs) (Morawetz et al., 2020). To identify the MAGs, *K*-means clustering was performed on imaging data from 385 fMRI experiments from 107 published papers that investigated emotion generative processes vs emotion regulatory processes, with forward inference to establish a functional fingerprint of the MAGs. The first and second MAG (i.e., MAG1 and MAG2) comprise mostly cortical areas including parts of the PFC, while the third and fourth MAG (i.e., MAG3 and MAG4) also include subcortical areas like the bilateral amygdala. The networks' functional fingerprinting revealed that MAG1's role is predominantly in working memory, response inhibition, and reasoning, thus it can be referred to as exhibiting emotional control functions. MAG2 also shows a strong association with cognitive control, but it places a stronger focus on language compared to MAG1. On the other hand, MAG3 specializes in emotional reactivity and memory, while MAG4 has a more mixed functional fingerprint, being involved in both emotion generative and regulatory processes. MAG4's activation is associated with pain perception, interoception, and perception of the emotional stimulus in both generative and regulatory phases. Thus, MAG3 can be simplified to be an emotion generation network, while MAG4 can be defined as an interoception network.

Generally, besides the MAGs, other large-scale brain networks are not exempt from age-related changes, as overall higher network integration in these networks in later life stage has been reported (Chan et al., 2014; Ferreira & Busatto, 2013; Grady et al., 2016; L. He et al., 2020; Onoda & Yamaguchi, 2013). High functional integration of a brain network describes that it is strongly connected across the whole cortex while low functional integration describes high modularity of the network and richer within connections (Sporns, 2013). Increased integration of the frontoparietal control network, default mode network, and dorsal attention network predicts poorer working memory and decreased processing speed in late adulthood (Ng et al., 2016; Salami et al., 2018). However, higher integration may not only result in deficits regarding behavioral performance. Previous studies have shown that the age-related increase in integration of the salience network within the rest of the cortex (Voss et al., 2013), and enhanced integration between the executive control network and the rest of the cortex have beneficial effects on life satisfaction in older adults (Lyoo & Yoon, 2017). Increased integration of specifically right hemispheric networks was further associated with greater working memory capacity in older adults, but not in younger (Crowell et al., 2020). Indicating that cognition in the aged brain might to some degree depend on or profit from network integration. However, increased integration at the network level might also be detrimental to the system, as it implies that in which more neural systems are activated in parallel reflecting a more diffuse, and nonspecific recruitment of brain systems. Other findings indicate as a consequence of increased global integration in aging, older adults show slower cognitive processing (Imms et al., 2021; Xin Li et al., 2023). Generally, it can be inferred that changes at the network level can influence or even predict

observable behavioral changes, although with differential impact depending on the emotional or cognitive nature of the functions (Chen et al., 2022; Levakov et al., 2021; Wei Li et al., 2023; Yu Li et al., 2022). Changes occurring within the functional topography of emotion regulation networks across the lifespan could therefore provide valuable insights into why older adults are better at regulating negative affect (Mather, 2012) or showing higher mental health than young adults (Thomas et al., 2016). However, the connections between emotion regulation network functional integration and mental wellbeing across the aging lifespan is yet to be fully examined. Therefore, the present study aimed to investigate whether and how the emotion regulation networks' topography changes across the adult lifespan, and whether the changes in the networks' organization are associated with psychological outcomes, such as stress or loneliness. By using graph theory, specifically measures such as Participation Coefficient and modular interaction, we investigated the functional integration of the four large-scale networks (MAGs) across the adult lifespan with the available dataset of the human connectome project - Aging (HCP-Aging), and the Cambridge Centre for Ageing and Neuroscience (Cam-CAN) dataset.

Considering the increased involvement of frontal areas during emotional processing and emotion regulation specifically (Joseph U. Kim et al., 2019; Nashiro et al., 2012; Pessoa, 2008) as well as improved emotion regulation capacities in older adults (Mather, 2012; Schweizer et al., 2019), we speculated that the predominantly frontal emotional control networks (such as MAGs 1 and 2) would become more integrated in late adulthood. Specifically, we extracted participation coefficient (PC) and modular interaction and correlated them with age. This increased integration was expected to facilitate the inhibitory control required for emotion regulation, primarily exhibited by the frontal areas (Aron, 2007). Thus, we anticipated that frontal nodes of emotion regulation networks become more integrated with increasing age. Secondly, we hypothesized that differences in functional network integration would mediate age-related effects on perceived negative affects measured by psychological questionnaires. This was based on the idea that higher ability in regulating negative emotions could result in higher emotional well-being and lower stress (Extremera et al., 2020; Gross et al., 1997; Kraiss et al., 2020). Specifically, improved measures of emotional wellbeing (depression, anxiety, loneliness, stress) are mediated by network and nodal integration. Together, these analyses will provide additional insight into how functional emotion networks differ across the adult lifespan with specific details on the differences of functional integration, and association with psychological measures.

2 Material and methods

2.1 Participants

The data used for this study were obtained from the HCP-Aging (Bookheimer et al., 2019) and the Cam-CAN datasets (Shafto et al., 2014; Taylor et al., 2016). We utilized minimally processed imaging

data and demographic information, which included the age, gender and years of education of each subject. Additionally, questionnaire data, such as stress scale, loneliness scale, and depression and anxiety scales from the NIH Toolbox (Hodes et al., 2013)(Table 1), were used to assess brain-behavior relationship.

In total, after excluding participants with translation which exceeded 2 mm or rotation that exceeded 2°, and structural abnormalities, 621 healthy participants (349 females) between the ages of 36 and 100 years old (Mean age = 59.1 ± 15.1) were included in the analyses. Subject with suspected Alzheimer's disease and neurological disorders (e.g., brain tumors, Parkinson's disease, stroke) were excluded in both datasets. Additionally, HCP-Aging excluded participants who have been diagnosed and treated for major psychiatric disorders (e.g., schizophrenia, bipolar disorder) and individuals with major depression who had been treated for 12 months or more in the past five years (Bookheimer et al., 2019). All primary analyses were carried out on the HCP-Aging and for validation purposes, we subsequently performed the same analyses on data from 333 (155 Female, mean age = 58.6 ± 14.7) from the Cam-CAN dataset using the same pipelines. All participants in both the HCP-A and Cam-CAN datasets provided informed consent and their participation was voluntary.

Table 1: Descriptive Data from HCP-Aging and Cam-CAN for both younger and older adults.

	Mean	S.D.	Pearson-r (with age)	Significance level
HCP-A (N=621)				
Age	60.33	15.69	-	-
Anxiety Arousal	7.98	2.32	-0.06	0.16
Anxiety Affect	9.18	2.49	-0.03	0.48
Depression	9.05	2.64	0.05	0.24
Loneliness	9.23	3.83	-0.23	<0.001
Perceived Stress	22.42	5.80	-0.25	<0.001
Years of Education	17.53	2.14	-0.02	0.55
Head motion (FD)	0.17	0.06	0.12	0.002
Gender	272M/349F		-	-
Cam-CAN (N=333)				
Age	58.59	14.76	-	-
HADS Anxiety	4.59	2.97	-0.19	<0.001
HADS Depression	2.59	2.46	0.05	0.41
Years of Education	16.66	4.63	-0.36	<0.001
Head motion (FD)	0.24	0.09	0.39	<0.001
Gender	178M/155F		-	-

2.2 Acquisition and preprocessing of neuroimaging data

The brain images from HCP-A were acquired using a 32-channel head coil on a 3T Siemens Prisma System across multiple centers (Washington University, University of Minnesota, Massachusetts General Hospital, Harvard University, University of California Los Angeles, Oxford University). Acquisition protocols were unified across all centers. T1 structural images in 0.8 mm resolution were acquired using the multi-echo MPRAGE sequence (TR = 2500 ms; TI = 1000 ms; TE = 1.8/3.6/5.4/7.2 ms; and flip angle= 8°). Acquisition of functional data was done with a 2D multiband gradient echo planar imaging (EPI) sequence (TR = 800 ms; TE = 37 ms; 72 axial slices; voxel size = $2.0 \times 2.0 \times 2.0$ mm; flip angle = 52°; multiband factor = 8). All participants completed four sessions of 6 minutes and five seconds of eyes-open resting-state scans, total of 30.4 minutes of resting-state scan. Sessions were conducted either on the same or consecutive days (detail of scanning parameters for HCP and Cam-CAN can be found in **Supplementary table S1**).

We acquired the minimally processed data from HCP-A, which included spatial artifact/distortion removal using the field maps calculated based on the difference in distortion between the two-phase encoding directions, surface generation, cross-modal registration, and alignment to standard space (Glasser et al., 2013). CONN Toolbox version 19v was used to perform additional in-house preprocessing steps comprised of spatial smoothing with an isotropic Gaussian kernel of 4 mm full width at half maximum, bandpass filtering at 0.008-0.09 Hz, denoising including anatomical component-based noise correction procedure (aComCor), motion regression with 12 regressors (6 motion parameters and their first-order derivatives), scrubbing (removal of a variable number of noise components based on outlier scans that either showed framewise displacement larger than 0.9 or BOLD signal changes over 5 S.D.), and linear detrending (Nieto-Castanon, 2020).

Time courses of the 36 ROIs of the MAGs were extracted using masks provided by Morawetz et al (Table 2). The masks contain the exact voxels identified in the original paper that established the MAGs (Morawetz et al., 2020). We decided to use these masks due to the non-spherical nature of many of these ROIs, making the signal localization more precise.

2.3 Topological analyses of integration of MAG networks

We used resting state data from both HCP-Aging and Cam-CAN datasets to test our hypotheses. There is a high correspondence between task and resting-state functional connectivity (Cole et al., 2016; Killgore et al., 2017; Krylova et al., 2021) and previous research has linked functional connectivity in the MAGs to remission in major depressive disorder (Hang Wu et al., 2022), the reinforcement of moral foundations (Wei Li et al., 2023), and the estrous cycle in women (Derntl et al., 2024) during rest. Thus, we believe that resting state data could provide valuable insight into the emotion regulation network during the task. Here we defined the integration in the context of brain networks, which describes the concept of how different modules or communities within a brain network interact and integrate information. A module that is highly integrated exhibits extensive connections and interactions with other modules of the graph, enabling communication and coordination across the brain network. Two measurements of integration are given by the participation coefficient and intra- and inter-modular connectivity (this measurement will be referred to as modular interaction, (Jinhui Wang et al., 2010).

To address differences in integration of emotion regulation functional networks and their respective areas (Hypotheses 1 and 2) across the lifespan, we employed a graph theoretical approach. Four meta-analytical groupings were chosen with 36 regions of interest (ROI), and ROI-wise time series were extracted by averaging the signal across voxels within the ROI (coordinates of all ROIs are presented in Table 2). Time courses of each ROI were correlated using Pearson correlation and then the resulting correlation matrixes were r-to-z transformed. These z-transformed values were saved in 36×36 functional connectivity (FC) matrices. Subsequently, we constructed graphs from these FC matrices, where each node represented an ROI, and the edges represented functional connections. We specifically selected the top 30% highest positive values from the FC matrices, which were then represented as undirected unweighted edges in the graph. The choice of a 30% sparsity level was based on the argument of its biological plausibility (Dennis et al., 2012; Sporns & Betzel, 2016). Each of the MAGs represented a functional module within the graph. In addition, while the size of ROIs was different, correlating the mean time-courses of all voxels in each ROI to construct FC matrices ensures that there is no bias in the signal from each ROI affecting the results. To test our first two hypotheses, that functional integration is increased in higher age on the network and nodal level, participation coefficient (PC) and modular interaction were applied. The PC quantifies the relative connectivity of a node in its respective module compared to the entire graph. With the PC of node i being $PC_i = 1 - \sum_{m=1}^{M} \left(\frac{k_{im}}{k_i} \right)^2$, m corresponds to the respective module (in our case M = 4, given the four MAGs), k_i represents the total amount of edges a node has, and k_{im} represents the count of edges the node has within its own module (Sporns et al., 2007). That is, if all the node's edges are limited to its module, its participation coefficient is 0. Due to the sparsity of the graph, PC values will mostly be

either 0 or above 0.5, because when the number of total edges is low, few connections outside the
associated module will bring the PC into the higher ranges of 0.5 or more (Bloznelis, 2013).

To test the integration of the MAGs (Hypothesis 1) we calculated the modular PC by averaging the
nodal PC values of each module. In addition, to address Hypothesis 1, we computed the modular
interaction as the ratio of observed connections to the total possible connections. Due to the sparsity
of the graph, the correlations between the time courses of certain ROIs did not exceed the threshold
and were therefore not represented as an edge in the graph. Only edges that exceeded the sparsity
threshold of 0.3 are considered "observed connections". Thus, a modular interaction value of one
represents the state that every possible connection that one module can make either within itself or
with another module is also observed. To address which nodes of the network are more integrated in
higher age and thus contribute to differences in network integration (Hypothesis 2), nodal PC was
used.

Figure 1 Visualization of Workflow with ROIs of MAGs: A) The ROIs of each MAG. MAG1
(indicated in purple) consisted of the bilateral dorsolateral prefrontal cortex (dlPFC) and inferior
parietal lobule, right insula, cingulate gyrus, and precuneus. MAG2 (indicated in yellow) involved the
bilateral ventrolateral prefrontal cortex (vmPFC), left temporoparietal junction, and middle temporal
gyrus, and left dlPFC. MAG3 (indicated in red) was based on the bilateral amygdala and fusiform

gyrus, left parahippocampal gyrus, periaqueductal grey, and vmPFC. MAG4 (indicated in green)

consisted of the left superior parietal lobule, bilateral postcentral gyrus, left insula, precuneus,

thalamus and posterior cingulate cortex. Exact coordinates and volumes are presented in Table 2. B)

Flowchart of procedure 1) Participants were scanned at rest. 2) ROIs were masked, and their time

courses extracted. 3) Time courses of all 36 ROIs of the MAGs were correlated with one another. 4)

Graphs were derived from the correlation matrices with a sparsity of 0.3. 5) Mathematical measures

were derived from the graph, including Network participation coefficient (PC), nodal PC, and

network interactions. 6) Statistic analyses were performed on the graph measures, age, and behavior.

Table 2. MNI coordinates of 36 regions of interest from all four meta-analytic groupings

(MAGs)

Serial number	Hemisphere	Region	Volume (voxels)	MNI Coordinates		
				x	y	z
MAG1 - Emotional Control						
1	L	Superior Frontal Gyrus	1463	0	24	50
2	R	Middle Frontal Gyrus	1378	40	24	42
3	R	Inferior Parietal Lobule	1246	58	-52	38
4	L	Inferior Parietal Lobule	777	−58	−50	44
5	L	Middle Frontal Gyrus	583	−36	52	−2
6	L	Middle Frontal Gyrus	536	−42	14	48
7	R	Middle Frontal Gyrus	349	42	46	−8
8	R	Insula	250	36	16	6
9	R	Cingulate Gyrus	167	2	−22	30
10	R	Precuneus	118	10	−64	36
MAG2 - Emotional Control (language)						
11	L	Inferior Frontal Gyrus	2433	−46	24	−8
12	L	Superior Frontal Gyrus	2074	−4	10	62
13	R	Inferior Frontal Gyrus	857	50	28	−8
14	L	Superior Temporal Gyrus	838	−46	−52	28
15	L	Middle Temporal Gyrus	628	−54	−34	−2
16	L	Middle Frontal Gyrus	571	−44	6	50
17	L	Superior Frontal Gyrus	385	−30	48	26
18	L	Caudate	245	−16	10	12
19	R	Tuber	205	36	−60	−30
MAG3 - Emotion Generation						
20	L	Amygdala	1080	−22	−4	−16
21	R	Amygdala	814	24	−4	−18
22	R	Fusiform Gyrus	597	40	−46	−18
23	R	Thalamus	441	6	−26	0
24	L	Fusiform Gyrus	157	−38	−54	−14
25	L	Parahippocampal Gyrus	152	−22	−28	−4
26	B	Medial Frontal Gyrus	127	0	54	−10
27	L	Inferior Occipital Gyrus	114	−42	−76	−6
MAG4 - Interoception						
28	L	Postcentral Gyrus	520	−58	−22	32
29	L	Insula	469	−44	−4	10
30	L	Superior Parietal Lobule	280	−28	−52	56
31	R	Postcentral Gyrus	217	62	−22	30
32	L	Cuneus	153	−10	−76	22

33	L	Middle Occipital Gyrus	144	–48	–74	2
34	R	Thalamus	128	10	–26	–4
35	R	Precuneus	104	28	–60	38
36	R	Posterior Cingulate	104	16	–56	16

263 *Note: L=left; R=right; B=bilateral; MNI = Montreal Neurological Institute.*

2.4 Network properties and association with age and behavior

To test our first hypothesis regarding age-related differences in the functional integration of the
MAGs, partial correlations between age and modular PC for each module, as well as correlations
between age and each of the modular interaction values were performed. For the second hypothesis
regarding age-related differences in the functional integration of the ROIs in the MAGs, partial
correlations between age and nodal PC for each node were performed. During these analyses, the
effects of years of education, gender, and head motion were controlled for as partial regressors. Since
in large samples, common normality tests like the Shapiro Wilk test often fail we assessed normality
of each variable visually (Royston, 1982; Shapiro & Wilk, 1965). After visually assessing each
variable's normality, we determined whether parametric Pearson correlation or nonparametric
Spearman correlation will be performed (to see variables distributions see **Supplementary Figure**
S4). For this we normalized the data using the minmax algorithm. We applied Bonferroni correction
for multiple comparisons with $\alpha_{adjusted} = (0.05/ (4 \text{ modules} + 36 \text{ nodes} + 10 \text{ interactions})) = 0.001$. To
assess the relationship of age to network integration we fitted a generalized additive model to the data
to address the possibility of nonlinearity in the data. Generalized Additive Modeling (GAM) is a
method that incorporates smooth functions to represent the influence of variables. These functions can
exhibit nonlinearity, depending on the inherent data patterns (Hastie, Trevor J., 1992). When the
model involves nonlinear effects, such as possibly the impact of age on network integration, GAM
enables to detect and incorporate typical nonlinear patterns that a traditional linear model might
overlook. Our model contained one explanatory variable, age, which is captured by 7 penalized basis
splines. Gender, years of education, and head movement were treated as linear regressor terms using
Gaussian regression.

Additionally, for our third hypothesis regarding the association between network integration and
psychological measures, we examined whether there was an age-related effect on any of the
psychometric tests using partial correlation with subsequent linear regression. To assess whether the
integration of specific nodes or modules had any associations with alterations in psychometric scores,
we correlated all graph-derived variables with relevant psychometric score variables. To further
explore whether age-related differences in behavior might be mediated by differences in MAG
functional integration, we performed mediation analyses for the psychometric variables that showed
an age effect, as well as significant linear regression with MAG integration parameters. For only three
models all the linear associations between variables were significant. Thus, for the significance value
we chose $\alpha_{adjusted} = 0.05/3 = 0.017$. In these mediation analyses, age served as the independent

variable, the psychometric score as the dependent variable, and the network integration variable as the
mediator. All variables were z-scored for this analysis. All graph theoretical analyses were carried out
in the MATLAB-based toolbox GRETNA (J. Wang et al., 2015). All correlation analyses were
performed using Python 3.9.16 with the packages SciPy.stats, statsmodels (Seabold & Perktold,
2010), and pingouin (Seongho Kim, 2015). Visualization was carried out using Python's Seaborn and
MATLAB's BrainNetViewer (Xia et al., 2013). Mediation analysis was conducted using python
statsmodels.

2.5 Validation

To further ensure the robustness and generalizability of the results we performed all the same analyses
on resting-state fMRI data from the Cam-CAN for validation purposes, thereby confirming that the
conclusions drawn are not due to peculiarities or biases in a single dataset but are indeed reflective of
broader, reliable trends. The participant age ranges in Cam-CAN slightly differed from those in HCP-
Aging, with some participants being younger than the youngest participant in HCP-Aging. To ensure
consistency, the age range in Cam-CAN was adjusted to match that of HCP-Aging, by excluding
those participants in the cam-CAN dataset that were younger than the youngest participant in the
HCP-Aging dataset (HCP-Aging data set starts at 36 years old). Thus, analyses were performed on
333 subjects between the ages 36 and 88. Additional preprocessing was carried out with the CONN
default preprocessing pipeline which is identical to the processing of the HCP-Aging data. The
parameters for graph analysis remained the same, and the covariates used in the regressions were also
identical. Since Cam-CAN and HCP-Aging utilized different questionnaires, we used the behavioral
scales in Cam-CAN that most closely resembled the psychometric data from HCP-Aging. In this
instance, we found that the Hospital Anxiety and Depression Scales (HADSs) were the ones that
captured the most alike aspects of mood in terms of the behavioral scales used in HCP-Aging (Harms
et al., 2018; Hodes et al., 2013).

3 Results

3.1 Higher integration of MAG1 and lower integration of MAG4 in older age.

3.1.1 Network PC differences

Confirming our first hypothesis, the emotional control network (MAG1) showed a positive
association between age and network PC (r=0.260, p<0.001; **Figure 2A**). The GAM also revealed a
linear trend in all the splines.

On the other hand, we found age-related reduction in functional network integration of MAG4, which
is considered to be a mediator between the network that is involved in both the perception of the
emotional stimulus during the emotion generative process and during the regulation of emotional
responses, that did not confirm our hypothesis. Rather, our data showed a significant shift towards

lower PC of MAG4 over the time course of aging ($r = -0.285$, $p < 0.001$; **Figure 2B**). Together these findings partially support our first hypothesis, that brain networks involved in emotion regulation become more integrated in late adulthood. However, the GAM revealed that the integration of MAG4 followed a nonlinear pattern (pseudo $R^2 = 0.116$), as the PC values were constant until the age of 70, followed by a steep decrease in network integration. This implies that only roughly after 70 MAG4 functional integration starts to decrease. No significant association was found between the PC of MAG2 and MAG3 with age (MAG2: $r = -0.051$, $p = 0.207$. MAG3: $r = -0.042$, $p = 0.293$).

3.1.2 Age-related differences between MAGs are specific to MAGs 1 and 4

To further investigate Hypothesis 1, we examined the modular interaction between and within modules, which describes how many of the possible connections between all areas of one module to another are present. Modular interaction of one means all areas of both networks are connected, and zero means no functional connection. MAGs 1 and 2 seem to share most of the connections with one another in a functional community, and so do MAGs 3 and 4. However the interaction between the two communities (i.e., MAG1&2 with MAG3&4) is low (**Figure 2C**). Increasing integration of MAG1 with age was reflected in both increased within- and between- networks connections (**Figure 2D**). In the case of MAG1, this was indicated by the significant positive association between the modular interaction between MAG1 and both MAG2 ($r = 0.208$, $p < 0.001$) and MAG3 ($r = 0.315$, $p < 0.001$) and within itself ($r = 0.154$, $p < 0.001$) in later age. On the other hand, the decreased integration of MAG4 with age was mainly carried by a loss of connections to MAG2 ($r = -0.175$, $p < 0.001$) and MAG3 ($r = -0.242$, $p < 0.001$) and within itself ($r = -0.302$, $p < 0.001$). Further, MAG3 showed lower modular interaction within itself ($r = -0.146$, p<0.001).

Figure 2. Functional integration of emotion regulation networks: A) Generative additive model (GAM) fit of the integration of MAG1, the emotional control network, across age, showing a linear positive association between age and network PC. B) GAM fit of network integration to age to MAG4, the interoception network. The model reveals a decrease of integration after the age of 70. C) Modular interaction between the MAGs in percent, representing how many of all possible connections between nodes of the module are present. Darker color indicates high modular interaction. The modular interaction between the MAGs revealed a pattern of high connectivity between and within MAG1 and MAG2, and MAG3 and MAG4, however showed low interaction between them. D) Correlation values of this specific modular interaction with age. Blue fields indicate a negative association between modular interaction between these MAGs with age and red fields indicate a positive association between the interaction and age. Fields marked with a cross did not reveal a significant correlation with age.

3.2 Specific nodes within a network contribute to overall network integration.

To address Hypothesis 2, in which ROIs of the MAGs differ in functional integration during aging, we investigated the association of the integration of each individual ROI with age. The nodal PC of each node of the graph from each MAG was correlated with age.

We found that in MAG1, five ROIs showed a positive association between nodal PC and age (**Figure 3**). These ROIs included the bilateral middle frontal gyrus (right: $r = 0,150$, $p < 0.001$; left: $r = 0.155$, $p < 0.001$), left inferior parietal lobule ($r = 0.165$, $p<0.001$), right cingulate gyrus ($r = 0.288$, $p < 0.001$), and right precuneus ($r = 0.157$, $p < 0.001$). All these nodes showed a significant positive correlation between PC and age, indicating that these areas became more integrated globally with the other MAGs with increasing age.

A negative association between PC and age in three ROIs in MAG4 (**Figure 3**) were found in the left insula ($r = -0.198$, $p < 0.001$), left cuneus ($r = -0.157$, $p < 0.001$), as well as right thalamus ($r = -0.169$, $p < 0.001$; **Supplementary Figure 3**). These areas were less integrated in higher age, suggesting that they made proportionally less connections with the other MAGs with increasing age. No other significant correlations were found between age and any of the nodes from other MAGs. These findings altogether indicate that frontal ROIs showed positive association between nodal PC and age, more integrated with other MAGs with age, while there are also medial and lateral areas affected, becoming less integrated with age, confirming our second hypothesis.

Figure 3. Functional integration of specific brain areas associated with MAGs in aging: A) the r values for the ROIs that showed significant Pearson correlation with age demonstrated on the brain. Orange hues represent the areas that are more integrated in older adults while blue color indicate the areas that are less integrated with age. The areas that show increased integration correspond to the bilateral middle frontal gyrus, left inferior parietal lobule, right cingulate gyrus, and right precuneus, while the areas that show decreased integration correspond to the left insula, left cuneus and right thalamus. B) Glass-brain showing also the medial areas which show altered integration across the

3.3 Inter-MAG1 connectivity is associated with increased depression while integration of the left cuneus suppresses age effect on loneliness and perceived stress.

To test the third hypothesis that differences in emotional regulation could be attributed to differences in emotion network integration, partial correlations with the integration parameters of the nodes and modules and psychometric scores were performed. We found that the within modular interaction of MAG1 correlated positively with depression scores ($r = 0.142$, $p < 0.001$), indicating that across different ages, more functional connections between the ROIs of MAG1 indicate a higher tendency towards depressive behaviors.

To further test whether alterations in network integration could explain age-related mental-wellbeing changes, mediation analyses was conducted. First, linear associations between psychological questionnaires and age were assessed, to examine the association between the independent variable (i.e., age) and the dependent one (i.e., psychometric scores). Self-report questionnaires provided by the HCP-Aging included measures of anxiety, depression, stress, and loneliness. Of these, perceived stress and loneliness (**Figure 4A**) showed significant decrease with increasing age. Then we performed linear regression between functional integration of the MAGs as measured by modular/nodal PC and modular interaction, with psychometric scores, to check which network topographical index could constitute a mediating variable.

For the mediation analyses, age was used as the independent variable, perceived stress and loneliness as dependent variables in two separate mediations and network integration measures from all MAGS as mediators. Only one mediation model was significant, in the case of which it was a partial suppression (Smith et al., 1992). A suppressor variable does not have a direct effect on the dependent variable but, when included in the analysis, increases the relationship between the independent variable and the dependent variable. (for review on suppressor variables in psychology, see (Rucker et al., 2011). The direct effect from age on loneliness was negative (β = -263, $p < 0.001$) with the total effect (via cuneus integration) was slightly less negative (β = -0.240, $p = 0.014$), suggesting that the integration of the left cuneus of MAG4 suppressed the negative age effect on loneliness scores. Hence, for those older individuals that showed rather high cuneus integration even in older age, also exhibited increased loneliness scores. Thus, decreased PC of the cuneus suppressed the negative association of age and loneliness (**Figure 4D**). Since there was no other significant correlation between the integration of the MAGs and psychometric scores, those mediation models were not considered (for all mediation models see **Supplementary Table S6**).

Figure 4. Association between age, cuneus functional integration and perceived stress and loneliness. (A) Linear regression between age and loneliness scores showing negative association with age. (B) Linear regression between functional integration of the left cuneus and age, showing lower PC with higher age. (C) Linear regression between functional integration of the left cuneus and loneliness scores, indicating a small negative trend. (D) Mediation model between age and loneliness scores, with cuneus PC mediating the effect partially suppressing the negative age effect on perceived stress. The area marked in red are the respective voxels of the Cuneus.

3.4 Validation with Cam-CAN dataset confirms increased integration of MAG1

Following the same analyses steps as for the analysis on the HCP-Aging dataset, we found that the findings on the HCP-A could be partially validated by the Cam-CAN data. Regarding our first hypothesis, a positive association of the PC of MAG1 with age, in the Cam-CAN dataset was confirmed ($r = 0.119$, $p = 0.031$). Other findings regarding the nodal changes from each MAG were confirmed in the Cam-CAN data including increased PC of the precuneus of MAG1 ($r = 0.116$, $p = 0.036$) in later age, decreased modular interaction of connections between MAGs 3 and 4 ($r = -0.138$, $p = 0.012$).

As for the second hypothesis, a measure of positive or negative affect in later age Cam-CAN provided the HADS depression and anxiety scores of the subjects. We found a negative correlation between HADS anxiety scores and age ($r = -0.185$, $p < 0.001$). This might reflect the behavioral findings of decreased stress and loneliness in the HCP-Aging.

None of the findings in HCP-Aging were contradicted by the Cam-CAN data, however some associations between integration measures of the MAGs and age present in the HCP-Aging data were

448   not significant in the Cam-CAN dataset and vice versa. There were additional significant associations
449   between network integration and age. Details of these differences are presented in the supplementary
450   table (**Supplementary Table 2**). All nonlinear models of Cam-CAN and HCP-Aging data are
451   presented in **Figure S5**.

4 Discussion

In the present study, our objective was to explore age-related alterations in the functional network
architecture of emotion regulation networks. These alterations could provide valuable insights into
how a shift in the regulation of emotions occurs in aging. We have gathered evidence indicating that
across the adult lifespan alterations in large-scale brain networks associated with the emotional
response occur, which correlate to differences in psychological well-being in later age. Partially
confirming our hypothesis, we observed an increase in functional integration of the emotional control
network, namely MAG1, with increasing age. We also observe decreased integration of interoception
network, namely MAG 4, in late adulthood. Interestingly the regions in MAG1 that show significant
positive association with age are areas involved in high-order function, while the significant regions in
MAG4 are except the insula, mostly sensory areas (Huntenburg et al., 2018). Network integration has
previously been debated in the context of balancing metabolic costs and communication efficiency
(Hillary & Grafman, 2017). Thus, specifically in the older brain that is faced with metabolic changes
(Mattson & Arumugam, 2018), increased integration might reflect a compensatory mechanism for
metabolic changes. The current study was not able to directly determine in what way the aged brain is
advantaged or disadvantaged by the observed differences in emotion regulation networks. Therefore,
the following arguments represent a speculation. More generally, a recent study has shown that
cognitive ability is promoted by higher global integration, better crystallized intelligence, while on the
other hand processing speed and memory recruitment are associated with lower integration (Rong
Wang et al., 2021) The observed increased integration of MAG1 might reflect an improved emotion
regulation in later age due to integration-driven crystalline. Similarly, decreased integration of MAG4
could reflect a conservation of processing speed during emotion generation and reflection. A fast
emotional response is crucial to individuals irrespective of age, therefore this function might be
preserved into higher ages and reflected by the decreased integration of MAG4.

Increased integration of emotion control network

The MAG1 consists of mostly frontal regions, including the superior and middle frontal gyrus. This
finding aligns with previous studies that suggest the increased prefrontal recruitment in emotion
regulation tasks that is observed in older adults (Berboth & Morawetz, 2021; Nashiro et al., 2012).
Particularly the bilateral MFG, left IPL, precuneus, and cingulate nodes of this network showed
increased integration in later life, all of which have been demonstrated to be crucial in the regulation
of emotions and for emotional wellbeing and stability (Glinka et al., 2020; Tomasino et al., 2022;

Williams et al., 2006; Winecoff et al., 2011; Ziaei et al., 2017, 2021). On top of that, MAG1 is predominantly involved in the downregulation of negative emotions (Morawetz et al., 2020). We suggested that as MAG1 becomes more integrated in later life, there is an enhanced transmission of information to the other networks (Sporns, 2013) involved in generating and processing the emotional responses, leading to improved emotion regulation in late adulthood, which could in turn have beneficial effects on mood. Many of the ROIs belonging to MAG1 are also associated with other large-scale brain networks such as the DMN or frontoparietal network (FPN). It is specifically these areas of MAG1 that also exhibited increased nodal PC in higher age, including the MFG, inferior parietal lobule and cingulate gyrus. Thus, the observed increased integration of MAG1 in the current study is consistent with previous findings that there is increased network integration of the DMN and FPN in aging (Andrews-Hanna et al., 2007; Ng et al., 2016). In a study investigating the test-retest reliability of the MAGs at ultrahigh magnetic field strength, the first two MAGs showed high to excellent reliability in the context of emotion regulation tasks (Berboth et al., 2021), further supporting our findings.

Moreover, we found that there is a positive association between the inter-module connectivity of MAG1 and depression scores, regardless of the individual's age. Indeed, a depressive phenotype has previously been associated with an increased top-down processing from the prefrontal areas towards limbic areas (Challis & Berton, 2015; LeDuke et al., 2023; Mogg & Bradley, 2016). MAG1 consists mostly of frontal areas and exhibits a rather top-down emotional control function (Morawetz et al., 2020). We showed that the number of connections that MAG1 makes within itself is higher in subjects with higher depression scores. This finding could reflect a tendency towards increased top-down inhibition as many prefrontal areas that are part of MAG1 are more functionally connected. Moreover, since MAG1 shares several regions with the DMN, this finding might reflect the observed increased functional connectivity of the DMN in major depression (Gillespie et al., 2020; Bao-Juan Li et al., 2018). Our current finding replicates the involvement of the functional connectivity increase of certain nodes within the DMN in depression and extends it with the other MAG1 areas that are not part of the DMN to contribute to the altered functional connectivity that is involved in the neuropathy of depression.

It remains highly important to investigate the drivers behind changes in functional network integration. Potential candidates for investigation include changes in receptor expression for different transmitters (Kringelbach et al., 2020; Shine, 2019), or differences in white matter development (Stevens et al., 2009). Our findings underscore the significance of functional network integration in emotional wellbeing across the adult lifespan. However, with a deeper understanding of what precipitates these changes, it could become a target for therapeutic intervention aimed at enhancing emotional wellbeing in later life.

Decreased integration of interoception network in aging

We observed decreased integration of MAG4 after the age of 70, which has been identified to be
involved both in emotion generation and regulatory processes, particularly linking the two networks
via interoception. Interoception is vital to perceive our own body states and helps shape the feeling
and intensity of emotions (Craig, 2002). Existing research, however, provides limited information on
how age differences in interoception may affect emotional experience and general well-being. Our
study shed some light on the fact that the decreased integration of MAG4 in older age may contribute
to the decline in interoceptive capabilities. According to previous studies, some areas such as the
insula, which is a critical area for interoception, decrease in volume with age (Good et al., 2001). We
report significantly decreased integration of the left insula in late adulthood, which could be attributed
to decay in volume and could further explain difficulties in interoceptive processes as part of emotion
regulation networks in older adults. These results should be confirmed with targeted studies on
interoception in the future studies combining volumetric measures with large-scale network
architecture. It must be noted that the observed decrease in integration does not follow a fully linear
trend, but the rate of change seems to increase only after the age of 70. Previous studies find
accelerated white matter decay, after the age of 70 (Fjell & Walhovd, 2010; Resnick, 2000) which
could contribute to decreased integration of MAG4. This age is also characterized as the start of a
decline in interoceptive awareness, including awareness of visceral sensations such as esophageal
pain, gastric distension, and heartbeats (Rayner, MacIntosh, 2000; Khalsa et al., 2009; Lasch et al.,
1997). The brain areas within MAG4 that contributed the most to the overall decreased integration of
this network included the insula, the cuneus, and the right thalamus. These areas have been linked
with empathy and relating to another person's negative emotion or pain (Bufalari et al., 2007;
Nummenmaa et al., 2008; Ziaei et al., 2021). We speculate that lower integration of these areas and
consequent reduced signaling from them may indicate older adults' difficulty reading and processing
negative affect in others (Mill et al., 2009; Hayes et al., 2020; Stutesman and Frye, 2023). However, it
could also reflect a preservation of emotion and empathetic processes in later life. Taken together, the
increased integration of nodes in MAG1 that are mostly associated with emotional control or
cognitive control tasks (Friedman & Robbins, 2022; Koch et al., 2018) with the decreased integration
of nodes in MAG4 that are associated with empathy and relating to others (Arioli et al., 2021;
Braadbaart et al., 2014) could reflect that emotion processing in older adults relies more on crystalline
aspects of cognition like knowledge about oneself and the world, a process that has previously been
described in the cognitive domain as the crystallization of cognition (for review see Spreng and
Turner, 2019) and has also been suggested to affect the sociocognitive domain (Henry et al., 2023).
The combined findings of higher integration of the frontal emotion regulation network (i.e. MAG1)
with lower integration of the mediating emotion regulation and generation network (i.e. MAG4) in

later age could reflect the crystallization of cognition in later life. However, targeted research is necessary to establish this relationship.

Likewise, the question remains if the observed changes in network integration relate to improved mood in late adulthood. In the HCP-Aging dataset we found a decline in perceived stress as well as loneliness in later life, therefore, we were interested if one of the observed differences in brain network integration could mediate this effect. There was only one significant mediation model suggesting that the decreased integration of the left cuneus as a partial suppressor (Yongnam Kim, 2019) reverts age-related decrease in perceived loneliness. While the cuneus is mostly regarded as a visual area, it has also been associated with sensory integration and psychiatric conditions (Martel et al., 2017; Nyatega et al., 2021). Activity and volume of the left cuneus has been associated with symptom severity in depression and anxiety (Brühl et al., 2014; Dotson et al., 2022; Guo et al., 2012; Peng et al., 2019; Yoon et al., 2016) and the ability to adjust one's emotion (Tan et al., 2014) In healthy aging the activity of the visual and parietal areas tends to decrease (Springer et al., 2023). One could speculate that the Cuneus constitutes an important relay area integrating visual information (Xiaoxi He et al., 2021; Nyatega et al., 2021). Thus, this finding might indicate that the age-related decrease of cuneus integration might alter brain networks' functional connectivity in a way that is beneficial on the perception of one's own loneliness in later life.

Integration of emotion regulation network across datasets

To ensure the generalizability and reproducibility of our results, a critical step in our methodology involved the validation of our key findings from the HCP-Aging using the Cam-CAN dataset. Our main finding, the increased integration of the frontal emotional control network, was replicated in the Cam-CAN dataset. However, the decrease in integration of the interoceptive network, MAG4, did not survive in the Cam-CAN dataset. One reason for such difference between the two datasets could be attributable to different scanning protocols (including different scanner systems and different repetition times, details in **Supplementary Table S1**), difference in the duration of scan times and having participants closing their eyes or not (Han et al., 2023; Lei Wu et al., 2010). The replication of the finding that MAG1 is more integrated in later life in eyes-open and eyes-closed resting state supports that this finding is a general trend which is not condition-dependent. Another aspect that must be considered in the nature of the two datasets is that the countries of origin of the data are different, which is important while studying social-emotional processes across the lifespan that might be affecting these findings (Foulkes & Blakemore, 2018; Kivimäki et al., 2020; Tooley et al., 2021).

4.1 Limitations and future perspectives

Due to the nature of the method using mathematical graphs, the interpretation of the findings needs to be carefully evaluated. An increase in the integration of a functional network might indicate increased information transmission from one network to another but could also imply functional blurring and

the loss of specialized processing in the network (Sporns, 2013). In the context of aging, the increased

global integration across the whole brain has been associated with decreased performance in cognitive

demands tasks (Pedersen et al., 2021). Hence it cannot be concluded whether the increase in MAG1

integration in later age is in any way beneficial or detrimental for the whole system. A conservation of

the low integration of other MAGs in later life might therefore also be an indicator as to why

emotional processes are preserved longer in later life even though other cognitive processes decline.

Another limitation that would also count as an advantage is the size of the HCP-Aging and Cam-CAN

datasets. The large sizes of the data allow also small effects to be found, however, our findings remain

to be replicated in future studies using small samples. Our findings yield a great starting point for

further research to explore the role of emotion regulation networks not only in aging but across all

ages. Future studies could investigate whether these networks change across the lifespan occur in

longitudinal studies, employing other graph measures to investigate the functional topology of these

networks.

4.2 Conclusion

In the present study, we found alterations in the functional architecture of emotion regulation

networks across the adult lifespan. The emotional control network (i.e., MAG1) became more

integrated with the other networks, consistent across two large datasets, the HCP-A, and Cam-CAN.

Additionally, we discovered that the network mediating emotion regulation and generation, the

interoception network (i.e., MAG4), became more segregated. To link these findings to emotional

wellbeing in late adulthood we identified a mediating role on loneliness of the cuneus, an area that

only recently is understood to be more than a visual area but has impacts on more abstract cognitive

functions. Overall, identifying topological changes in emotion network architecture is of particular

interest as it allows for further understanding of emotional aging and late-life mood disorders.

Disclosure

The authors declare no competing interest.

Funding

Funded by the Norwegian center of excellence. Project number 332640

CRediT authorship contribution statement

Leona R. Bätz: Writing – original draft, Writing – review & editing, Methodology, Investigation, Formal analysis, Data curation, Visualization, Validation. **Shuer Ye:** Writing – review & editing, Software, Supervision, Methodology, Data curation, Resources, Methodology, Investigation, Conceptualization. **Xiaqing Lan:** Writing – review & editing, Resources, Data curation. **Maryam Ziaei:** Writing – review & editing, Supervision, Project administration, Conceptualization, Funding acquisition.

Acknowledgements

Research reported in this publication was supported by the National Institute on Aging of the National Institutes of Health under Award Number U01AG052564 and by funds provided by the McDonnell Center for Systems Neuroscience at Washington University in St. Louis. The HCP-Aging 2.0 Release data used in this report came from DOI: 10.15154/1520707.

Data collection and sharing for this project was provided by the Cambridge Centre for Ageing and Neuroscience (CamCAN). CamCAN funding was provided by the UK Biotechnology and Biological Sciences Research Council (grant number BB/H008217/1), together with support from the UK Medical Research Council and University of Cambridge, UK.

This work was partially supported by the Research Council of Norway through its Centres of Excellence scheme, project number 332640.

References

Ahmed, S. P., Bittencourt-Hewitt, A., & Sebastian, C. L. (2015). Neurocognitive bases of emotion

regulation development in adolescence. *Developmental Cognitive Neuroscience*, *15*, 11–25.

https://doi.org/10.1016/j.dcn.2015.07.006

Andrews-Hanna, J. R., Snyder, A. Z., Vincent, J. L., Lustig, C., Head, D., Raichle, M. E., & Buckner, R. L.

(2007). Disruption of Large-Scale Brain Systems in Advanced Aging. *Neuron*, *56*(5), 924–935.

https://doi.org/10.1016/j.neuron.2007.10.038

Arioli, M., Cattaneo, Z., Ricciardi, E., & Canessa, N. (2021). Overlapping and specific neural correlates

for empathizing, affective mentalizing, and cognitive mentalizing: A coordinate-based

meta-analytic study. *Human Brain Mapping*, *42*(14), 4777–4804.

https://doi.org/10.1002/hbm.25570

Aron, A. R. (2007). The Neural Basis of Inhibition in Cognitive Control. *The Neuroscientist*, *13*(3), 214–

228. https://doi.org/10.1177/1073858407299288

Berboth, S., & Morawetz, C. (2021). Amygdala-prefrontal connectivity during emotion regulation: A

meta-analysis of psychophysiological interactions. *Neuropsychologia*, *153*, 107767.

https://doi.org/10.1016/j.neuropsychologia.2021.107767

Berboth, S., Windischberger, C., Kohn, N., & Morawetz, C. (2021). Test-retest reliability of emotion

regulation networks using fMRI at ultra-high magnetic field. *NeuroImage*, *232*, 117917.

https://doi.org/10.1016/j.neuroimage.2021.117917

Birditt, K. S., Fingerman, K. L., & Almeida, D. M. (2005). Age Differences in Exposure and Reactions to

Interpersonal Tensions: A Daily Diary Study. *Psychology and Aging*, *20*(2), 330–340.

https://doi.org/10.1037/0882-7974.20.2.330

Bloznelis, M. (2013). Degree and clustering coefficient in sparse random intersection graphs. *The*

Annals of Applied Probability, *23*(3). https://doi.org/10.1214/12-AAP874

Bookheimer, S. Y., Salat, D. H., Terpstra, M., Ances, B. M., Barch, D. M., Buckner, R. L., Burgess, G. C.,

Curtiss, S. W., Diaz-Santos, M., Elam, J. S., Fischl, B., Greve, D. N., Hagy, H. A., Harms, M. P.,

Hatch, O. M., Hedden, T., Hodge, C., Japardi, K. C., Kuhn, T. P., … Yacoub, E. (2019). The

Lifespan Human Connectome Project in Aging: An overview. *NeuroImage*, *185*, 335–348.

https://doi.org/10.1016/j.neuroimage.2018.10.009

Braadbaart, L., De Grauw, H., Perrett, D. I., Waiter, G. D., & Williams, J. H. G. (2014). The shared

neural basis of empathy and facial imitation accuracy. *NeuroImage*, *84*, 367–375.

https://doi.org/10.1016/j.neuroimage.2013.08.061

Brühl, A. B., Delsignore, A., Komossa, K., & Weidt, S. (2014). Neuroimaging in social anxiety

disorder—A meta-analytic review resulting in a new neurofunctional model. *Neuroscience &*

Biobehavioral Reviews, *47*, 260–280. https://doi.org/10.1016/j.neubiorev.2014.08.003

Bufalari, I., Aprile, T., Avenanti, A., Di Russo, F., & Aglioti, S. M. (2007). Empathy for Pain and Touch in

the Human Somatosensory Cortex. *Cerebral Cortex*, *17*(11), 2553–2561.

https://doi.org/10.1093/cercor/bhl161

C. K. Rayner, C. G. MacIntosh, I. M. (2000). Effects of Age on Proximal Gastric Motor and Sensory

Function. *Scandinavian Journal of Gastroenterology*, *35*(10), 1041–1047.

https://doi.org/10.1080/003655200451153

Carstensen, L. L., Isaacowitz, D. M., & Charles, S. T. (1999). Taking time seriously: A theory of

socioemotional selectivity. *American Psychologist*, *54*(3), 165–181.

https://doi.org/10.1037/0003-066X.54.3.165

Carstensen, L. L., Pasupathi, M., Mayr, U., & Nesselroade, J. R. (2000). Emotional experience in

everyday life across the adult life span. *Journal of Personality and Social Psychology*, *79*(4),

644–655. https://doi.org/10.1037/0022-3514.79.4.644

Carstensen, L. L., Turan, B., Scheibe, S., Ram, N., Ersner-Hershfield, H., Samanez-Larkin, G. R., Brooks,

K. P., & Nesselroade, J. R. (2011). Emotional experience improves with age: Evidence based

on over 10 years of experience sampling. *Psychology and Aging*, *26*(1), 21–33.

https://doi.org/10.1037/a0021285

Challis, C., & Berton, O. (2015). Top-Down Control of Serotonin Systems by the Prefrontal Cortex: A

Path toward Restored Socioemotional Function in Depression. *ACS Chemical Neuroscience*,

6(7), 1040–1054. https://doi.org/10.1021/acschemneuro.5b00007

Chan, M. Y., Park, D. C., Savalia, N. K., Petersen, S. E., & Wig, G. S. (2014). Decreased segregation of

brain systems across the healthy adult lifespan. *Proceedings of the National Academy of*

Sciences, *111*(46). https://doi.org/10.1073/pnas.1415122111

Charles, S. T. (2010). Strength and vulnerability integration: A model of emotional well-being across

adulthood. *Psychological Bulletin*, *136*(6), 1068–1091. https://doi.org/10.1037/a0021232

Charles, S. T., Piazza, J. R., Luong, G., & Almeida, D. M. (2009). Now you see it, now you don't: Age

differences in affective reactivity to social tensions. *Psychology and Aging*, *24*(3), 645–653.

https://doi.org/10.1037/a0016673

Chen, J., Tam, A., Kebets, V., Orban, C., Ooi, L. Q. R., Asplund, C. L., Marek, S., Dosenbach, N. U. F.,

Eickhoff, S. B., Bzdok, D., Holmes, A. J., & Yeo, B. T. T. (2022). Shared and unique brain

network features predict cognitive, personality, and mental health scores in the ABCD study.

Nature Communications, *13*(1), 2217. https://doi.org/10.1038/s41467-022-29766-8

Cohen, Sheldon, Kessler, Ronald C., & Gordon, Lynn Underwood. (1997). *Measuring Stress: A Guide*

for Health and Social Scientists. Oxford University Press.

Cole, M. W., Ito, T., Bassett, D. S., & Schultz, D. H. (2016). Activity flow over resting-state networks

shapes cognitive task activations. *Nature Neuroscience*, *19*(12), 1718–1726.

https://doi.org/10.1038/nn.4406

Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the

body. *Nature Reviews Neuroscience*, *3*(8), 655–666. https://doi.org/10.1038/nrn894

Crowell, C. A., Davis, S. W., Beynel, L., Deng, L., Lakhlani, D., Hilbig, S. A., Palmer, H., Brito, A.,

Peterchev, A. V., Luber, B., Lisanby, S. H., Appelbaum, L. G., & Cabeza, R. (2020). Older adults

benefit from more widespread brain network integration during working memory.

NeuroImage, *218*, 116959. https://doi.org/10.1016/j.neuroimage.2020.116959

Cyranowski, J. M., Zill, N., Bode, R., Butt, Z., Kelly, M. A. R., Pilkonis, P. A., Salsman, J. M., & Cella, D.

(2013). Assessing social support, companionship, and distress: National Institute of Health

(NIH) Toolbox Adult Social Relationship Scales. *Health Psychology*, *32*(3), 293–301.

https://doi.org/10.1037/a0028586

Dennis, E. L., Jahanshad, N., Toga, A. W., McMahon, K. L., De Zubicaray, G. I., Martin, N. G., Wright, M.

J., & Thompson, P. M. (2012). Test-Retest Reliability of Graph Theory Measures of Structural

Brain Connectivity. In N. Ayache, H. Delingette, P. Golland, & K. Mori (Eds.), *Medical Image*

Computing and Computer-Assisted Intervention – MICCAI 2012 (Vol. 7512, pp. 305–312).

Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-33454-2_38

Derntl, B., Eber, C. H., Kogler, L., Rehbein, E., Sundstöm-Poromaa, I., & Morawetz, C. (2024). Estradiol

modulates changes in effective connectivity in emotion regulation networks.

Psychoneuroendocrinology, *167*, 107103. https://doi.org/10.1016/j.psyneuen.2024.107103

Dotson, V. M., Bogoian, H. R., Gradone, A. M., Taiwo, Z., & Minto, L. R. (2022). Subthreshold

depressive symptoms relate to cuneus structure: Thickness asymmetry and sex differences.

Journal of Psychiatric Research, *145*, 144–147.

https://doi.org/10.1016/j.jpsychires.2021.12.013

English, T., & Carstensen, L. L. (2014). Selective narrowing of social networks across adulthood is

associated with improved emotional experience in daily life. *International Journal of*

Behavioral Development, *38*(2), 195–202. https://doi.org/10.1177/0165025413515404

Extremera, N., Sánchez-Álvarez, N., & Rey, L. (2020). Pathways between Ability Emotional Intelligence

and Subjective Well-Being: Bridging Links through Cognitive Emotion Regulation Strategies.

Sustainability, *12*(5), 2111. https://doi.org/10.3390/su12052111

Ferreira, L. K., & Busatto, G. F. (2013). Resting-state functional connectivity in normal brain aging.

Neuroscience & Biobehavioral Reviews, *37*(3), 384–400.

https://doi.org/10.1016/j.neubiorev.2013.01.017

Fjell, A. M., & Walhovd, K. B. (2010). Structural Brain Changes in Aging: Courses, Causes and

Cognitive Consequences. *Reviews in the Neurosciences, 21*(3).

https://doi.org/10.1515/REVNEURO.2010.21.3.187

Foulkes, L., & Blakemore, S.-J. (2018). Studying individual differences in human adolescent brain

development. *Nature Neuroscience, 21*(3), 315–323. https://doi.org/10.1038/s41593-018-

0078-4

Friedman, N. P., & Robbins, T. W. (2022). The role of prefrontal cortex in cognitive control and

executive function. *Neuropsychopharmacology, 47*(1), 72–89.

https://doi.org/10.1038/s41386-021-01132-0

Ghashghaei, H. T., Hilgetag, C. C., & Barbas, H. (2007). Sequence of information processing for

emotions based on the anatomic dialogue between prefrontal cortex and amygdala.

NeuroImage, 34(3), 905–923. https://doi.org/10.1016/j.neuroimage.2006.09.046

Gillespie, C. F., Szabo, S. T., & Nemeroff, C. B. (2020). Unipolar depression. In *Rosenberg's Molecular*

and Genetic Basis of Neurological and Psychiatric Disease (pp. 613–631). Elsevier.

https://doi.org/10.1016/B978-0-12-813866-3.00036-9

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi,

S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The minimal

preprocessing pipelines for the Human Connectome Project. *NeuroImage, 80*, 105–124.

https://doi.org/10.1016/j.neuroimage.2013.04.127

Glinka, K., Staudinger, U. M., Voelcker-Rehage, C., & Godde, B. (2020). Neural processing of arousing

emotional information is associated with executive functioning in older adults. *Emotion*,

20(4), 541–556. https://doi.org/10.1037/emo0000560

Good, C. D., Johnsrude, I. S., Ashburner, J., Henson, R. N. A., Friston, K. J., & Frackowiak, R. S. J.

(2001). A Voxel-Based Morphometric Study of Ageing in 465 Normal Adult Human Brains.

NeuroImage, 14(1), 21–36. https://doi.org/10.1006/nimg.2001.0786

Grady, C., Sarraf, S., Saverino, C., & Campbell, K. (2016). Age differences in the functional interactions

among the default, frontoparietal control, and dorsal attention networks. *Neurobiology of*

Aging, 41, 159–172. https://doi.org/10.1016/j.neurobiolaging.2016.02.020

Gross, J. J., Carstensen, L. L., Pasupathi, M., Tsai, J., Götestam Skorpen, C., & Hsu, A. Y. C. (1997).

Emotion and aging: Experience, expression, and control. *Psychology and Aging, 12*(4), 590–

599. https://doi.org/10.1037/0882-7974.12.4.590

Guo, W., Liu, F., Xue, Z., Xu, X., Wu, R., Ma, C., Wooderson, S. C., Tan, C., Sun, X., Chen, J., Liu, Z., Xiao,

C., Chen, H., & Zhao, J. (2012). Alterations of the amplitude of low-frequency fluctuations in

treatment-resistant and treatment-response depression: A resting-state fMRI study. *Progress*

in Neuro-Psychopharmacology and Biological Psychiatry, 37(1), 153–160.

https://doi.org/10.1016/j.pnpbp.2012.01.011

Han, J., Zhou, L., Wu, H., Huang, Y., Qiu, M., Huang, L., Lee, C., Lane, T. J., & Qin, P. (2023). Eyes-Open

and Eyes-Closed Resting State Network Connectivity Differences. *Brain Sciences, 13*(1), 122.

https://doi.org/10.3390/brainsci13010122

Harms, M. P., Somerville, L. H., Ances, B. M., Andersson, J., Barch, D. M., Bastiani, M., Bookheimer, S.

Y., Brown, T. B., Buckner, R. L., Burgess, G. C., Coalson, T. S., Chappell, M. A., Dapretto, M.,

Douaud, G., Fischl, B., Glasser, M. F., Greve, D. N., Hodge, C., Jamison, K. W., … Yacoub, E.

(2018). Extending the Human Connectome Project across ages: Imaging protocols for the

Lifespan Development and Aging projects. *NeuroImage, 183,* 972–984.

https://doi.org/10.1016/j.neuroimage.2018.09.060

Hastie, Trevor J. (1992). Generalized Additive Models. In *Statistical Models in S* (1st ed.). Routledge.

Hay, E. L., & Diehl, M. (2011). Emotion complexity and emotion regulation across adulthood.

European Journal of Ageing, 8(3), 157–168. https://doi.org/10.1007/s10433-011-0191-7

He, L., Wang, X., Zhuang, K., & Qiu, J. (2020). Decreased Dynamic Segregation but Increased Dynamic

Integration of the Resting-state Functional Networks During Normal Aging. *Neuroscience,*

437, 54–63. https://doi.org/10.1016/j.neuroscience.2020.04.030

He, X., Li, X., Fu, J., Xu, J., Liu, H., Zhang, P., Li, W., Yu, C., Ye, Z., & Qin, W. (2021). The morphometry of

left cuneus mediating the genetic regulation on working memory. *Human Brain Mapping*,

42(11), 3470–3480. https://doi.org/10.1002/hbm.25446

Henry, J. D., Grainger, S. A., & Von Hippel, W. (2023). Determinants of Social Cognitive Aging:

Predicting Resilience and Risk. *Annual Review of Psychology*, *74*(1), 167–192.

https://doi.org/10.1146/annurev-psych-033020-121832

Hess, T. M. (2014). Selective Engagement of Cognitive Resources: Motivational Influences on Older

Adults' Cognitive Functioning. *Perspectives on Psychological Science*, *9*(4), 388–407.

https://doi.org/10.1177/1745691614527465

Hillary, F. G., & Grafman, J. H. (2017). Injured Brains and Adaptive Networks: The Benefits and Costs

of Hyperconnectivity. *Trends in Cognitive Sciences*, *21*(5), 385–401.

https://doi.org/10.1016/j.tics.2017.03.003

Hodes, R. J., Insel, T. R., Landis, S. C., & On behalf of the NIH Blueprint for Neuroscience Research.

(2013). The NIH Toolbox: Setting a standard for biomedical research. *Neurology*,

80(11_supplement_3). https://doi.org/10.1212/WNL.0b013e3182872e90

Huntenburg, J. M., Bazin, P.-L., & Margulies, D. S. (2018). Large-Scale Gradients in Human Cortical

Organization. *Trends in Cognitive Sciences*, *22*(1), 21–31.

https://doi.org/10.1016/j.tics.2017.11.002

Imms, P., Domínguez D, J. F., Burmester, A., Seguin, C., Clemente, A., Dhollander, T., Wilson, P. H.,

Poudel, G., & Caeyenberghs, K. (2021). Navigating the link between processing speed and

network communication in the human brain. *Brain Structure and Function*, *226*(4), 1281–

1302. https://doi.org/10.1007/s00429-021-02241-8

Isaacowitz, D. M., Wadlinger, H. A., Goren, D., & Wilson, H. R. (2006). Selective preference in visual

fixation away from negative images in old age? An eye-tracking study. *Psychology and Aging*,

21(1), 40–48. https://doi.org/10.1037/0882-7974.21.1.40

Khalsa, S. S., Rudrauf, D., & Tranel, D. (2009). Interoceptive awareness declines with age.

Psychophysiology, *46*(6), 1130–1136. https://doi.org/10.1111/j.1469-8986.2009.00859.x

Killgore, W. D. S., Smith, R., Olson, E. A., Weber, M., Rauch, S. L., & Nickerson, L. D. (2017). Emotional

intelligence is associated with connectivity within and between resting state networks. *Social*

Cognitive and Affective Neuroscience, *12*(10), 1624–1636.

https://doi.org/10.1093/scan/nsx088

Kim, J. U., Weisenbach, S. L., & Zald, D. H. (2019). Ventral prefrontal cortex and emotion regulation in

aging: A case for utilizing transcranial magnetic stimulation. *International Journal of Geriatric*

Psychiatry, *34*(2), 215–222. https://doi.org/10.1002/gps.4982

Kim, S. (2015). ppcor: An R Package for a Fast Calculation to Semi-partial Correlation Coefficients.

Communications for Statistical Applications and Methods, *22*(6), 665–674.

https://doi.org/10.5351/CSAM.2015.22.6.665

Kim, Y. (2019). The Causal Structure of Suppressor Variables. *Journal of Educational and Behavioral*

Statistics, *44*(4), 367–389. https://doi.org/10.3102/1076998619825679

Kivimäki, M., Batty, G. D., Pentti, J., Shipley, M. J., Sipilä, P. N., Nyberg, S. T., Suominen, S. B., Oksanen,

T., Stenholm, S., Virtanen, M., Marmot, M. G., Singh-Manoux, A., Brunner, E. J., Lindbohm, J.

V., Ferrie, J. E., & Vahtera, J. (2020). Association between socioeconomic status and the

development of mental and physical health conditions in adulthood: A multi-cohort study.

The Lancet Public Health, *5*(3), e140–e149. https://doi.org/10.1016/S2468-2667(19)30248-8

Koch, S. B. J., Mars, R. B., Toni, I., & Roelofs, K. (2018). Emotional control, reappraised. *Neuroscience*

& Biobehavioral Reviews, *95*, 528–534. https://doi.org/10.1016/j.neubiorev.2018.11.003

Kraiss, J. T., Ten Klooster, P. M., Moskowitz, J. T., & Bohlmeijer, E. T. (2020). The relationship between

emotion regulation and well-being in patients with mental disorders: A meta-analysis.

Comprehensive Psychiatry, *102*, 152189. https://doi.org/10.1016/j.comppsych.2020.152189

Kringelbach, M. L., Cruzat, J., Cabral, J., Knudsen, G. M., Carhart-Harris, R., Whybrow, P. C.,

Logothetis, N. K., & Deco, G. (2020). Dynamic coupling of whole-brain neuronal and

neurotransmitter systems. *Proceedings of the National Academy of Sciences*, *117*(17), 9566–

9576. https://doi.org/10.1073/pnas.1921475117

Krylova, M., Skouras, S., Razi, A., Nicholson, A. A., Karner, A., Steyrl, D., Boukrina, O., Rees, G.,

Scharnowski, F., & Koush, Y. (2021). Progressive modulation of resting-state brain activity

during neurofeedback of positive-social emotion regulation networks. *Scientific Reports*,

11(1), 23363. https://doi.org/10.1038/s41598-021-02079-4

Lasch, H., Castell, D. O., & Castell, J. A. (1997). Evidence for diminished visceral pain with aging:

Studies using graded intraesophageal balloon distension. *American Journal of Physiology-*

Gastrointestinal and Liver Physiology, *272*(1), G1–G3.

https://doi.org/10.1152/ajpgi.1997.272.1.G1

Lawton, M. P., Kleban, M. H., Rajagopal, D., & Dean, J. (1992). Dimensions of affective experience in

three age groups. *Psychology and Aging*, *7*(2), 171–184. https://doi.org/10.1037/0882-

7974.7.2.171

LeDuke, D. O., Borio, M., Miranda, R., & Tye, K. M. (2023). Anxiety and depression: A top-down,

bottom-up model of circuit function. *Annals of the New York Academy of Sciences*, *1525*(1),

70–87. https://doi.org/10.1111/nyas.14997

Levakov, G., Faskowitz, J., Avidan, G., & Sporns, O. (2021). Mapping individual differences across brain

network structure to function and behavior with connectome embedding. *NeuroImage*, *242*,

118469. https://doi.org/10.1016/j.neuroimage.2021.118469

Li, B., Friston, K., Mody, M., Wang, H., Lu, H., & Hu, D. (2018). A brain network model for depression:

From symptom understanding to disease intervention. *CNS Neuroscience & Therapeutics*,

24(11), 1004–1019. https://doi.org/10.1111/cns.12998

Li, W., Ye, S., Zhu, B., Hoffman, M., Zhou, J., & Yang, Q. (2023). Individual differences in harm-related

moral values are associated with functional integration of large-scale brain networks of

emotional regulation. *Journal of Neuropsychology*, *17*(2), 335–350.

https://doi.org/10.1111/jnp.12303

Li, X., Salami, A., & Persson, J. (2023). Hub architecture of the human structural connectome: Links to

aging and processing speed. *NeuroImage*, *278*, 120270.

https://doi.org/10.1016/j.neuroimage.2023.120270

Li, Y., Zhuang, K., Yi, Z., Wei, D., Sun, J., & Qiu, J. (2022). The trait and state negative affect can be

separately predicted by stable and variable resting-state functional connectivity.

Psychological Medicine, *52*(5), 813–823. https://doi.org/10.1017/S0033291720002391

Lyoo, Y., & Yoon, S. (2017). Brain Network Correlates of Emotional Aging. *Scientific Reports*, *7*(1),

15576. https://doi.org/10.1038/s41598-017-15572-6

MacCormack, J. K., Stein, A. G., Kang, J., Giovanello, K. S., Satpute, A. B., & Lindquist, K. A. (2020).

Affect in the Aging Brain: A Neuroimaging Meta-Analysis of Older Vs. Younger Adult Affective

Experience and Perception. *Affective Science*, *1*(3), 128–154.

https://doi.org/10.1007/s42761-020-00016-8

Martel, M., Harvey, M.-P., Houde, F., Balg, F., Goffaux, P., & Léonard, G. (2017). Unravelling the effect

of experimental pain on the corticomotor system using transcranial magnetic stimulation and

electroencephalography. *Experimental Brain Research*, *235*(4), 1223–1231.

https://doi.org/10.1007/s00221-017-4880-0

Mather, M. (2012). The emotion paradox in the aging brain. *Annals of the New York Academy of

Sciences*, *1251*(1), 33–49. https://doi.org/10.1111/j.1749-6632.2012.06471.x

Mather, M. (2016). The Affective Neuroscience of Aging. *Annual Review of Psychology*, *67*(1), 213–

238. https://doi.org/10.1146/annurev-psych-122414-033540

Mattson, M. P., & Arumugam, T. V. (2018). Hallmarks of Brain Aging: Adaptive and Pathological

Modification by Metabolic States. *Cell Metabolism*, *27*(6), 1176–1199.

https://doi.org/10.1016/j.cmet.2018.05.011

Mill, A., Allik, J., Realo, A., & Valk, R. (2009). Age-related differences in emotion recognition ability: A

cross-sectional study. *Emotion*, *9*(5), 619–630. https://doi.org/10.1037/a0016562

Mogg, K., & Bradley, B. P. (2016). Anxiety and attention to threat: Cognitive mechanisms and

treatment with attention bias modification. *Behaviour Research and Therapy*, *87*, 76–108.

https://doi.org/10.1016/j.brat.2016.08.001

Morawetz, C., Riedel, M. C., Salo, T., Berboth, S., Eickhoff, S. B., Laird, A. R., & Kohn, N. (2020).

Multiple large-scale neural networks underlying emotion regulation. *Neuroscience &*

Biobehavioral Reviews, *116*, 382–395. https://doi.org/10.1016/j.neubiorev.2020.07.001

Nashiro, K., Sakaki, M., & Mather, M. (2012). Age Differences in Brain Activity during Emotion

Processing: Reflections of Age-Related Decline or Increased Emotion Regulation.

Gerontology, *58*(2), 156–163. https://doi.org/10.1159/000328465

Ng, K. K., Lo, J. C., Lim, J. K. W., Chee, M. W. L., & Zhou, J. (2016). Reduced functional segregation

between the default mode network and the executive control network in healthy older

adults: A longitudinal study. *NeuroImage*, *133*, 321–330.

https://doi.org/10.1016/j.neuroimage.2016.03.029

Nieto-Castanon, A. (2020). *Handbook of functional connectivity magnetic resonance imaging*

methods in CONN. Hilbert Press.

Nolen-Hoeksema, S., & Aldao, A. (2011). Gender and age differences in emotion regulation strategies

and their relationship to depressive symptoms. *Personality and Individual Differences*, *51*(6),

704–708. https://doi.org/10.1016/j.paid.2011.06.012

Nummenmaa, L., Hirvonen, J., Parkkola, R., & Hietanen, J. K. (2008). Is emotional contagion special?

An fMRI study on neural systems for affective and cognitive empathy. *NeuroImage*, *43*(3),

571–580. https://doi.org/10.1016/j.neuroimage.2008.08.014

Nyatega, C. O., Qiang, L., Adamu, M. J., Younis, A., & Kawuwa, H. B. (2021). Altered Dynamic

Functional Connectivity of Cuneus in Schizophrenia Patients: A Resting-State fMRI Study.

Applied Sciences, *11*(23), 11392. https://doi.org/10.3390/app112311392

Onoda, K., & Yamaguchi, S. (2013). Small-worldness and modularity of the resting-state functional

brain network decrease with aging. *Neuroscience Letters*, *556*, 104–108.

https://doi.org/10.1016/j.neulet.2013.10.023

Pedersen, R., Geerligs, L., Andersson, M., Gorbach, T., Avelar-Pereira, B., Wåhlin, A., Rieckmann, A.,

Nyberg, L., & Salami, A. (2021). When functional blurring becomes deleterious: Reduced

system segregation is associated with less white matter integrity and cognitive decline in

aging. *NeuroImage*, *242*, 118449. https://doi.org/10.1016/j.neuroimage.2021.118449

Peng, W., Jia, Z., Huang, X., Lui, S., Kuang, W., Sweeney, J. A., & Gong, Q. (2019). Brain structural

abnormalities in emotional regulation and sensory processing regions associated with

anxious depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *94*,

109676. https://doi.org/10.1016/j.pnpbp.2019.109676

Pessoa, L. (2008). On the relationship between emotion and cognition. *Nature Reviews Neuroscience*,

9(2), 148–158. https://doi.org/10.1038/nrn2317

Pilkonis, P. A., Choi, S. W., Reise, S. P., Stover, A. M., Riley, W. T., Cella, D., & PROMIS Cooperative

Group. (2011). Item Banks for Measuring Emotional Distress From the Patient-Reported

Outcomes Measurement Information System (PROMIS®): Depression, Anxiety, and Anger.

Assessment, *18*(3), 263–283. https://doi.org/10.1177/1073191111411667

Resnick, S. M. (2000). One-year Age Changes in MRI Brain Volumes in Older Adults. *Cerebral Cortex*,

10(5), 464–472. https://doi.org/10.1093/cercor/10.5.464

Royston, J. P. (1982). An Extension of Shapiro and Wilk's W Test for Normality to Large Samples.

Applied Statistics, *31*(2), 115. https://doi.org/10.2307/2347973

Rucker, D. D., Preacher, K. J., Tormala, Z. L., & Petty, R. E. (2011). Mediation Analysis in Social

Psychology: Current Practices and New Recommendations. *Social and Personality Psychology*

Compass, *5*(6), 359–371. https://doi.org/10.1111/j.1751-9004.2011.00355.x

Salami, A., Rieckmann, A., Karalija, N., Avelar-Pereira, B., Andersson, M., Wåhlin, A., Papenberg, G.,

Garrett, D. D., Riklund, K., Lövdén, M., Lindenberger, U., Bäckman, L., & Nyberg, L. (2018).

Neurocognitive Profiles of Older Adults with Working-Memory Dysfunction. *Cerebral Cortex*.

https://doi.org/10.1093/cercor/bhy062

Schweizer, S., Stretton, J., Van Belle, J., Price, D., Calder, A. J., Cam-CAN, & Dalgleish, T. (2019). Age-

related decline in positive emotional reactivity and emotion regulation in a population-

derived cohort. *Social Cognitive and Affective Neuroscience, 14*(6), 623–631.

https://doi.org/10.1093/scan/nsz036

Seabold, S., & Perktold, J. (2010). *Statsmodels: Econometric and Statistical Modeling with Python. 9,*

92--96.

Shafto, M. A., Tyler, L. K., Dixon, M., Taylor, J. R., Rowe, J. B., Cusack, R., Calder, A. J., Marslen-Wilson,

W. D., Duncan, J., Dalgleish, T., Henson, R. N., Brayne, C., & Matthews, F. E. (2014). The

Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol: A cross-sectional,

lifespan, multidisciplinary examination of healthy cognitive ageing. *BMC Neurology, 14*(1),

204. https://doi.org/10.1186/s12883-014-0204-1

Shapiro, S. S., & Wilk, M. B. (1965). An Analysis of Variance Test for Normality (Complete Samples).

Biometrika, 52(3/4), 591. https://doi.org/10.2307/2333709

Shine, J. M. (2019). Neuromodulatory Influences on Integration and Segregation in the Brain. *Trends*

in Cognitive Sciences, 23(7), 572–583. https://doi.org/10.1016/j.tics.2019.04.002

Smith, R. L., Ager, J. W., & Williams, D. L. (1992). Suppressor Variables in Multiple

Regression/Correlation. *Educational and Psychological Measurement, 52*(1), 17–29.

https://doi.org/10.1177/001316449205200102

Sporns, O. (2013). Network attributes for segregation and integration in the human brain. *Current*

Opinion in Neurobiology, 23(2), 162–171. https://doi.org/10.1016/j.conb.2012.11.015

Sporns, O., & Betzel, R. F. (2016). Modular Brain Networks. *Annual Review of Psychology, 67*(1), 613–

640. https://doi.org/10.1146/annurev-psych-122414-033634

Sporns, O., Honey, C. J., & Kötter, R. (2007). Identification and Classification of Hubs in Brain

Networks. *PLoS ONE, 2*(10), e1049. https://doi.org/10.1371/journal.pone.0001049

Spreng, R. N., & Turner, G. R. (2019). The Shifting Architecture of Cognition and Brain Function in

Older Adulthood. *Perspectives on Psychological Science*, *14*(4), 523–542.

https://doi.org/10.1177/1745691619827511

Springer, S. D., Erker, T. D., Schantell, M., Johnson, H. J., Willett, M. P., Okelberry, H. J., Rempe, M. P.,

& Wilson, T. W. (2023). Disturbances in primary visual processing as a function of healthy

aging. *NeuroImage*, *271*, 120020. https://doi.org/10.1016/j.neuroimage.2023.120020

St. Jacques, P., Dolcos, F., & Cabeza, R. (2010). Effects of aging on functional connectivity of the

amygdala during negative evaluation: A network analysis of fMRI data. *Neurobiology of*

Aging, *31*(2), 315–327. https://doi.org/10.1016/j.neurobiolaging.2008.03.012

Stevens, M. C., Skudlarski, P., Pearlson, G. D., & Calhoun, V. D. (2009). Age-related cognitive gains are

mediated by the effects of white matter development on brain network integration.

NeuroImage, *48*(4), 738–746. https://doi.org/10.1016/j.neuroimage.2009.06.065

Stutesman, M. G., & Frye, D. A. (2023). Affective Theory of Mind in Late Adulthood: The Role of

Emotion Complexity and Social Relatedness. *Experimental Aging Research*, *49*(5), 472–500.

https://doi.org/10.1080/0361073X.2022.2137359

Tan, Y., Zhang, Q., Li, W., Wei, D., Qiao, L., Qiu, J., Hitchman, G., & Liu, Y. (2014). The correlation

between Emotional Intelligence and gray matter volume in university students. *Brain and*

Cognition, *91*, 100–107. https://doi.org/10.1016/j.bandc.2014.08.007

Taylor, J. R., Williams, N., Cusack, R., Auer, T., Shafto, M. A., Dixon, M., Tyler, L. K., Cam-CAN, &

Henson, R. N. (2016). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data

repository: Structural and functional MRI, MEG, and cognitive data from a cross-sectional

adult lifespan sample. *NeuroImage*, *144*, 262–269.

https://doi.org/10.1016/j.neuroimage.2015.09.018

Thomas, M. L., Kaufmann, C. N., Palmer, B. W., Depp, C. A., Martin, A. S., Glorioso, D. K., Thompson,

W. K., & Jeste, D. V. (2016). Paradoxical Trend for Improvement in Mental Health With Aging:

A Community-Based Study of 1,546 Adults Aged 21–100 Years. *The Journal of Clinical*

Psychiatry, *77*(08), e1019–e1025. https://doi.org/10.4088/JCP.16m10671

Tomasi, D., & Volkow, N. D. (2012). Aging and functional brain networks. *Molecular Psychiatry*, *17*(5),

549–558. https://doi.org/10.1038/mp.2011.81

Tomasino, B., Maggioni, E., Bonivento, C., Nobile, M., D'Agostini, S., Arrigoni, F., Fabbro, F., &

Brambilla, P. (2022). Effects of age and gender on neural correlates of emotion imagery.

Human Brain Mapping, *43*(13), 4116–4127. https://doi.org/10.1002/hbm.25906

Tooley, U. A., Bassett, D. S., & Mackey, A. P. (2021). Environmental influences on the pace of brain

development. *Nature Reviews Neuroscience*, *22*(6), 372–384.

https://doi.org/10.1038/s41583-021-00457-5

Urry, H. L., & Gross, J. J. (2010). Emotion Regulation in Older Age. *Current Directions in Psychological*

Science, *19*(6), 352–357. https://doi.org/10.1177/0963721410388395

Voss, M. W., Wong, C. N., Baniqued, P. L., Burdette, J. H., Erickson, K. I., Prakash, R. S., McAuley, E.,

Laurienti, P. J., & Kramer, A. F. (2013). Aging Brain from a Network Science Perspective:

Something to Be Positive About? *PLoS ONE*, *8*(11), e78345.

https://doi.org/10.1371/journal.pone.0078345

Wang, J., Wang, X., Xia, M., Liao, X., Evans, A., & He, Y. (2015). GRETNA: A graph theoretical network

analysis toolbox for imaging connectomics. *Frontiers in Human Neuroscience*, *9*.

https://doi.org/10.3389/fnhum.2015.00386

Wang, J., Zuo, X., & He, Y. (2010). Graph-based network analysis of resting-state functional MRI.

Frontiers in Systems Neuroscience. https://doi.org/10.3389/fnsys.2010.00016

Wang, R., Liu, M., Cheng, X., Wu, Y., Hildebrandt, A., & Zhou, C. (2021). Segregation, integration, and

balance of large-scale resting brain networks configure different cognitive abilities.

Proceedings of the National Academy of Sciences, *118*(23), e2022288118.

https://doi.org/10.1073/pnas.2022288118

Williams, L. M., Brown, K. J., Palmer, D., Liddell, B. J., Kemp, A. H., Olivieri, G., Peduto, A., & Gordon,

E. (2006). The Mellow Years?: Neural Basis of Improving Emotional Stability over Age. *Journal*

of Neuroscience, *26*(24), 6422–6430. https://doi.org/10.1523/JNEUROSCI.0022-06.2006

Winecoff, A., LaBar, K. S., Madden, D. J., Cabeza, R., & Huettel, S. A. (2011). Cognitive and neural

contributors to emotion regulation in aging. *Social Cognitive and Affective Neuroscience*,

6(2), 165–176. https://doi.org/10.1093/scan/nsq030

Wu, H., Liu, R., Zhou, J., Feng, L., Wang, Y., Chen, X., Zhang, Z., Cui, J., Zhou, Y., & Wang, G. (2022).

Prediction of remission among patients with a major depressive disorder based on the

resting-state functional connectivity of emotion regulation networks. *Translational*

Psychiatry, *12*(1), 391. https://doi.org/10.1038/s41398-022-02152-0

Wu, L., Eichele, T., & Calhoun, V. D. (2010). Reactivity of hemodynamic responses and functional

connectivity to different states of alpha synchrony: A concurrent EEG-fMRI study.

NeuroImage, *52*(4), 1252–1260. https://doi.org/10.1016/j.neuroimage.2010.05.053

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A Network Visualization Tool for Human Brain

Connectomics. *PLoS ONE*, *8*(7), e68910. https://doi.org/10.1371/journal.pone.0068910

Yoon, S., Shim, M., Kim, H. S., & Lee, S.-H. (2016). Enhanced Early Posterior Negativity to Fearful

Faces in Patients with Anxiety Disorder. *Brain Topography*, *29*(2), 262–272.

https://doi.org/10.1007/s10548-015-0456-0

Ziaei, M., Oestreich, L., Reutens, D. C., & Ebner, N. C. (2021). Age-related differences in negative

cognitive empathy but similarities in positive affective empathy. *Brain Structure and*

Function, *226*(6), 1823–1840. https://doi.org/10.1007/s00429-021-02291-y

Ziaei, M., Salami, A., & Persson, J. (2017). Age-related alterations in functional connectivity patterns

during working memory encoding of emotional items. *Neuropsychologia*, *94*, 1–12.

https://doi.org/10.1016/j.neuropsychologia.2016.11.012

Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica*

Scandinavica, *67*(6), 361–370. https://doi.org/10.1111/j.1600-0447.1983.tb09716.x

1040