Worldwide Soundscapes: a synthesis of passive acoustic monitoring across realms

Authors & affiliations

Kevin FA Darras[1,2,3C], Rodney Rountree[4,5], Steven Van Wilgenburg[6], Anna F Cord[3,7], Frederik Pitz[3], Youfang Chen[2], Lijun Dong[9], Amandine Gasc[10], Tzu-Hao Lin[11], Patrick Mauritz Diaz[12], Shih-Hung Wu[13], Marcus Salton[14], Sarah Marley[15], Laura Schillé[16], Paul Jacobus Wensveen[17], Camille Desjonquères[18,19], Orlando Acevedo-Charry[20,21], Matyáš Adam[22], Jacopo Aguzzi[23], Michel André[24], Alexandre Antonelli[25,26,27], Leandro Aparecido Do Nascimento[28], Giulliana Appel[29], Christos Astaras[30], Andrey Atemasov[31,32], Luc Barbaro[33], Fritjof Basan[34], Carly Batist[35,36], Adriá López Baucells[37], Julio Ernesto Baumgarten[38], Just T Bayle Sempere[39], Kristen Bellisario[40], Asaf Ben David[41], Oded Berger-Tal[42], Matthew G Betts[43], Iqbal Bhalla[44], Thiago Bicudo[35], Marta Bolgan[45], Sara Bombaci[46], Martin Boullhesen[47], Tom Bradfer-Lawrence[48,49], Robert A Briers[50], Michal Budka[51], Katie Burchard[52], Alice Calvente[53], Maite Cerezo-Araujo[54], Gunnar Cerwén[55], Maria Chistopolova[56], Christopher W Clark[57], Benjamin Cretois[58], Chapin Czarnecki[59], Luis P da Silva[60,61], Wigna da Silva[62], Laurence H De Clippele[63], David de la Haye[64], Ana Silvia de Oliveira Tissiani[65], Devin de Zwaan[66,67], Ricardo Díaz-Delgado[68], Pedro Diniz[69], Dorgival Diógenes Oliveira-Júnior[70,71], Thiago Dorigo[72], Saskia Dröge[73], Marina Duarte[74,75], Adam Duarte[76], Kerry Dunleavy[77], Robert Dziak[78], Simon Elise[79,80], Hiroto Enari[8], Haruka S Enari[8], Florence Erbs[24], Nina Ferrari[43], Luane Ferreira[62], Abram B Fleishman[77], Bárbara Freitas[81,82,83], Nick Friedman[84], Jérémy SP Froidevaux[48,85], Svetlana Gogoleva[86], Maria Isabel Gonçalves[38], Carolina Gonzaga[74], José Miguel González Correa[39], Eben Goodale[87], Benjamin Gottesman[57], Ingo Grass[88], Jack Greenhalgh[89], Jocelyn Gregoire[90], Jonas Hagge[91], William Halliday[92], Antonia Hammer[93], Tara Hanf-Dressler[94], Samara Haver[95], Daniel Hending[96], Jose Hernandez-Blanco[56], Thomas Hiller[88], Joe Chun-Chia Huang[97], Katie Lois Hutchinson[98], Janet Jackson[99], Alain Jacot[100], Olaf Jahn[101,102], Jasper Kanes^^, Ellen Kenchington[103], Sebastian Kepfer-Rojas[104], Justin Kitzes[59], Tharaka Kusuminda[105], Yael Lehnardt[42], Paula Leitman[72], José Leon[106], Cicero Simão Lima-Santos[70,71], Kyle John Lloyd[107,108,109], Audrey Looby[110,111], David López-Bosch[37], Tatiana Maeda[112], Christos Mammides[113], Gabriel Marcacci[100], Matthias Markolf[114,115], Marinez Isaac Marques[65], Charles W Martin[110], Dominic A Martin[116,117], Kathy Martin[66,118], Matthew McKown[77], Logan JT McLeod[119], Oliver Metcalf[120], Christoph Meyer[121], Grzegorz Mikusinski[122], Larissa Sayuri Moreira Sugai[57], Dave Morris[99], Sandra Müller[123], Kelsie A Murchy[124], Maria Mas Navarro[37], Rym Nouioua[125], Julian D Olden[126], Steffen Oppel[49,100], Anna N Osiecka[127], Miles Parsons[92], Filipa Isabel Pereira Samarra[17], Cristian Pérez-Granados[128], Liliana Piatti[129], Mauro Pichorim[70,71], Thiago Pinheiro[62], Jean-Nicolas Pradervand[100], Craig Radford[130], Emiliano Ramalho[35], Laura K Reynolds[131], Klaus Riede[101], Talen Rimmer[124], Ricardo Rocha[96], Luciana Rocha[62], Samuel RP-J Ross[132], Carolyn Rosten[133], Carlos Salustio-Gomes[70,71], Philip Samartzis[134], José Santos[62], Kevin Scharffenberg[135], Renée P Schoeman[136], Karl-Ludwig Schuchmann[65], Esther Sebastián-González[128], Fannie Shabangu[137,138], Taylor Shaw[123], Xiaoli Shen[139], David Singer[91], Jenni Stanley[140], Karolin Thomisch[141], Laura Torrent[37], Juan Traba[142,143], Junior A Tremblay[144], Sunny Tseng[145], Mao-Ning Tuanmu[146], Marisol Valverde[147], Ben Vernasco[76,148], Raiane Vital da Paz[70,71], Matthew Ward[149], Matthew Weldy[43,150], Julia Wiel[58], Jacob Willie[151], Heather Wood[152], Songhai Li[9C], Renata Sousa-Lima[62C], Thomas Cherico Wanger[2C]

C: corresponding authors

1: EFNO, ECODIV, INRAE, Domaine des Barres, Nogent-sur-Vernisson, Centre-Val de Loire, 45290, France

2: Sustainable Agricultural Systems & Engineering Lab, School of Engineering, Westlake University, No. 600 Dunyu Road, Sandun Town, Xihu District, Hangzhou, Zhejiang province, , China

3: Chair of Computational Landscape Ecology, Faculty of Environmental Sciences, Dresden University of Technology, Helmholtzstr. 10, Dresden, 01069, Germany

4: Biology Department, University of Victoria, Cunningham Building, Room 202, PO Box 1700 STN CSC, Victoria, BC, V8W 2Y2, Canada

5: The Fish Listener, , , 23 Joshua Lane, Waguoit, MA, 2536, U.S.A.

6: Terrestrial Unit, Prairie Region, Canadian Wildlife Service, Environment & Climate Change Canada, Prairie & Northern Wildlife Research Centre, 115 Perimeter Road, Saskatoon, Saskatchewan, S7N 0X4, Canada

7: Agro-ecological Modeling Group, Faculty of Agriculture, University of Bonn, Niebuhrstr. 1a, Bonn, 53113, Germany

8: Wildlife Ecology & Management, Faculty of Agriculture, Yamagata University, 1-23 Wakabamachi, Tsuruoka, Yamagata, 9978555, Japan

9: Marine Mammal and Marine Bioacoustics Laboratory, Department of Deep Sea Science, Institute of Deep-sea Science and Engineering Chinese Academy of Sciences, 28 Luhuitou Road, Sanya, Hainan, 572000, China

10: Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, , Aix Marseille Univ, Avignon Univ, CNRS, IRD, Campus Aix Technopôle de l'environnement Arbois Méditerranée Avenue Louis Philibert Bât Villemin, Aix-en-Provence, 13545, France

11: Marine Ecoacoustics and Informatics Lab, Biodiversity Research Center, Academia Sinica, 128 Academia Road, Section 2, Nankang, Taipei , , 11529, Taiwan

12: Climate Change and Biodiversity, , Hanns R. Neumann Stiftung Indonesia, , Muaradua, OKU Selatan, South Sumatera, 32212, Indonesia

13: Biodiversity Indicator Lab, Division of Habitats and Ecosystems, Endemic Species Research Institute, 1 Minsheng East Road, Jiji Town, Nantou County, Taiwan, 552005, Taiwan

14: Australian Antarctic Division, Science Branch, Department of Climate Change, Energy, Environment and Water, Channel Highway, Kingston, Tasmania, 7050, Australia

15: Scotland's Rural College, Craibstone Estate, Aberdeen, AB21 9YA, United Kingdom

16: BIOGECO, INRAE, University of Bordeaux, 69 route d'Arcachon, Cestas, 33610, France

17: Westman Islands Research Centre, , University of Iceland, Ægisgata 2, Vestmannaeyjar, 900, Iceland

18: Institut de Systématique, Évolution, Biodiversité, , Muséum National d'Histoire naturelle, Rue Buffon, Paris, 75005, France

19: Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, , , Grenoble, 38000, France

20: Colección de Sonidos Ambientales Mauricio Álvarez-Rebolledo - Colecciones Biológicas, Subdirección de Investigaciones, Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Claustro de San Agustín, Villa de Leyva, Boyacá, , Colombia

21: Quantitative Ecology Lab & Ordway Lab of Ecosystem Conservation, School of Natural Resources and Environment, Department of Wildlife Ecology and Conservation, & Florida Museum of Natural History, University of Florida, 715 Newell Dr, Building 866 & 1659 Museum Rd, Dickinson Hall 315, Gainesville, Florida, 32611, U.S.A.

22: Department of Environmental Security, Faculty of Logistics and Crisis Management, Tomas Bata University, Studentské nám. 1532, Uherské Hradiště, , 68601, Czech Republic

23: Marine Science Institute, ICM, CSIC, ICM, CSIC, Paseig Barceloneta, Barcelona, , Spain

24: Laboratory of Applied Bioacoustics, , Polytechnic University of Catalonia, BarcelonaTech, Rambla Exposicion, 24, Vilanova I la Geltrú, Barcelona, 8800, Spain

25: Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, United Kingdom, , Kew, Richmond, Surrey, TW9 3AE, United Kingdom

26: Gothenburg Global Biodiversity Centre, Department of Biological and Environmental Sciences, University of Gothenburg, Box 461, Göteborg, SE 40530, Sweden

27: Department of Plant, , University of Oxford, South Parks Road, Oxford, OX1 3RB, United
Kingdom
28: Bioacoustics Group, Science Department, Biometrio.earth, Dr. Schönemann Str. 38,
Saarbrücken, 66123, Germany
29: Biodiversidade e Serviços Ecossistêmicos, Instituto Tecnológico Vale, Boaventura da
Silva, 955, Belém, PA, 66055-090, Brazil
30: Wildlife Lab, Forest Research Institute, Hellenic Agricultural Organization - DIMITRA
(ELGO-DIMITRA), Vasilika, Thessaloniki, TK 57006, Greece
31: Department of Zoology and Animal Ecology, School of Biology, V.N.Karazin Kharkiv
National University, 4 Svobody Square, Kharkiv, 61022, Ukraine
32: Research Department, National Park 'Homilshanski Lisy', 27 Monastyrska Street,
Koropove village, Kharkiv region, 63437, Ukraine
33: DYNAFOR, University of Toulouse, INRAE, Chemin de Borde-Rouge, Castanet-Tolosan,
F-31326, France
34: Federal Maritime and Hydrographic Agency (BSH), , , Bernhard-Nocht Straße 78,
Hamburg, 20359, Germany
35: Rainforest Connection, Science department, , 440 Cobia Drive, Suite 1902, , TX, 77494,
U.S.A.
36: Arbimon, , , , , , U.S.A.
37: BiBio Research Group, , Natural Sciences Museum of Granollers, Av. Francesc Macià
51, Granollers, Catalonia, 08402, Spain
38: Applied Ecology & Conservation Lab, Biological Sciences, Universidade Estadual de
Santa Cruz, Rodovia Jorge Amado km 16, Salobrinho, Ilhéus, Bahia, 45662900, Brazil
39: Marine Biological Acoustic, Marine Sciences and Applied Biology , University of Alicante,
C/ San Vicente del Raspeig s/n, San Vicente del Raspeig, Alicante, 3690, Spain
40: HIFI Lab, John Martinson Honors College, Purdue University West Lafayette Indiana,
1101 Third Street, West Lafayette, Indiana , 47906, U.S.A.
41: Tamar Dayan's Lab, School of zoology, Steinhardt Museum of Natural History, Tel-Aviv
university, Klausner 12, Tel-Aviv, Tel-Aviv, 6997801, Israel
42: The Conservation Behavior Research Group, Mitrani Department of Desert Ecology,
Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev,
Midreshet Ben-Gurion, , 8499000, Israel
43: Forest Landscape Ecology Lab, Department of Forest Ecosystems and Society, Oregon
State University, , , , U.S.A.
44: School of Geography and the Environment, Oxford University, South Parks Rd, Oxford,
OX1 3QY, United Kingdom
45: Ocean Science Consulting Limited, , , Spott Road, Dunbar, EH42 1RR , United Kingdom
46: Equity in Ecological Research Lab, Department of Fish, Wildlife, and Conservation
Biology, Colorado State University, 1474 Campus Delivery, Fort Collins, Colorado, 80521,
U.S.A.
47: Instituto de Ecorregiones Andinas (INECOA-UNJu-CONICET), Facultad de Ingeniería,
Universidad Nacional de Jujuy, Gorriti 237, San Salvador de Jujuy, Jujuy, 4600, Argentina
48: Biological and Environmental Sciences, University of Stirling, Stirling, Stirlingshire, FK9
4LA, Scotland
49: Centre for Conservation Science, RSPB, 2 Lochside View, Edinburgh Park, Edinburgh, ,
EH12 9DH, United Kingdom
50: Centre for Conservation and Restoration Science, School of Applied Sciences,
Edinburgh Napier University, Sighthill Campus, Edinburgh, EH11 4BN, United Kingdom
51: Department of Behavioural Ecology, Faculty of Biology, Adam Mickiewicz University,
Uniwersytetu Poznańskiego 6, Poznań, 61614, Poland
52: Northeast Fisheries Science Center, Narragansett Laboratory, , 28 Tarzwell Drive,
Narragansett, RI, 2882, U.S.A.
53: Laboratório de Botânica Sistemática, Departamento de Botânica, Ecologia e Zoologia,
Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio
Grande do Norte, 59078-900, Brazil

54: South Iceland Research Centre, , University of Iceland, Lindarbraut 4, Laugarvatn, 840, Iceland

55: Sensola SLU Multisensory Outdoor Laboratory, Department of Landscape Architecture, Planning and Management, Swedish University of Agricultural Sciences, SLU, Sundsvägen 4-6, Alnarp, 23053, Sweden

56: , , , ,

57: K. Lisa Yang Center for Conservation Bioacoustics , Cornell Lab of Ornithology , Cornell University , 159 Sapsucker Woods Rd , New York , 14850, U.S.A.

58: Miljødata, Norwegian Institute for Nature Research, Høgskoleringen 9, Trøndelag, Trondheim, 7034, Norway

59: Department of Biological Sciences, University of Pittsburgh, Clapp Hall, Fifth and Ruskin Avenues, Pittsburgh, PA, 15237, U.S.A.

60: CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Campus de Vairão, Universidade do Porto-InBIO, Universidade do Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal

61: BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, University of Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal

62: EcoAcoustic Research Hub (EAR Hub), Biosciences Center, Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio Grande do Norte, 59078-900, Brazil

63: School of Biodiversity, One Health & Veterinary Medicine, University of Glasgow, Graham Kerr Building, Hillhead street 82, Scotland, United Kingdom, G12 8QQ, United Kingdom

64: School of Arts and Cultures, Newcastle University, , Newcastle upon Tyne, NE1 7RU, United Kingdom

65: INAU Pantanal Biodata Center, National Institute of Science and Technology in Wetlands (INAU), Computational Bioacoustics Research Unit (CO.BRA), Federal University of Mato Grosso (UFMT), Cuiabá, 2367 Fernando Correa da Costa Ave, Mato Grosso, 78060-900, Brazil

66: Forest and Conservation Sciences, University of British Columbia, 2424 Main Mall, Vancouver, British Columbia, V6T 1Z4, Canada

67: Department of Biology, Mount Allison University, 63B York Street, Sackville, New Brunswick, E4L 1E7, Canada

68: Estación Biológica de Doñana-CSIC, , CSIC, c/ Americo Vespucio 26, Sevilla, 41092, Spain

69: Laboratório de Comportamento Animal, Departamento de Zoologia, Universidade de Brasília, Instituto de Ciências Biológicas, Distrito Federal, 70910-900, Brazil

70: Laboratório de Ornitologia, Departamento de Botânica e Zoologia, Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, RN, 59078-900, Brazil

71: Programa de Pós-Graduação em Ecologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, RN, 59078-900, Brazil

72: Lifeplan Rio de Janeiro / Araçá project, , , Estrada Hans Garlipp, s/n, Macaé de Cima, Fazenda Bacchus, Nova Friburgo, 28605-270, Brazil

73: Forest, Nature and Landscape, KU Leuven, Celestijnenlaan 200e, Leuven, 3001, Belgium

74: ECOA - Laboratory of Bioacoustics, Post graduate Program in Vertebrate Biology / Museum of Natural Sciences, Pontifical Catholic University of Minas Gerais, , Minas Gerais, , Brazil

75: Environmental Studies Center (CEA), , Sao Paulo State University - UNESP, Av. 24a, 1515, Bela Vista, Rio Claro, Sao Paulo, 13506-900, Brazil

76: Pacific Northwest Research Station, U.S.D.A. Forest Service, 3625 93rd Ave. SW, Olympia, Washington, 98512, U.S.A.

77: Conservation Metrics, Inc., 145 McAllister Way, Santa Cruz, CA, 95060, U.S.A.

78: Acoustics Program, Pacific Marine Envrionmental Laboratory, National Oceanic and Atmospheric Administration, 2030 SE Marine Science Drive, Newport, OR, 97365, U.S.A.

79: Reef Pulse S.A.S., NA, NA, Allée Maureau, Sainte-Clotilde, Reunion Island, 97490,
France
80: UMR 9220 ENTROPIE, Faculté des sciences et technologies, University of Reunion
Island, Avenue René Cassin, Sainte-Clotilde, Reunion Island, 97490, France
81: National Museum of Natural Sciences, Spanish National Research Council, Calle José
Gutiérrez Abascal 2, Madrid, 28006, Spain
82: Centre de Recherche sur la Biodiversité et l'Environnement (UMR 5300 CNRS-IRD-
TINPT-UPS), Université Paul Sabatier, 118 route de Narbonne, Toulouse, 31062 Toulouse
Cedex 9, France
83: Facultad de Ciencias, Universidad Autónoma de Madrid, C. Darwin 2, Madrid, 28049,
Spain
84: Centre for Taxonomy and Morphology, Leibniz Institute for the Analysis of Biodiversity
Change, Museum of Nature Hamburg, Martin-Luther-King-Platz 3, Hamburg, 20146,
Germany
85: Centre d'Ecologie et des Sciences de la Conservation (CESCO, UMR 7204), Museum
National d'Histoire Naturelle, Place de la Croix, Concarneau, Finistère, 29900, France
86: Laboratory of tropical ecology, , A.N. Severtsov Institute of Ecology and Evolution of
Russian Academy of Sciences, Leninsky Prospect, 33, Moscow, 119071, Russia
87: Department of Health and Environmental Science, , Xi'an Jiaotong-Liverpool University,
No. 8 Chongwen Road, Suzhou Industrial Park, Suzhou, Jiangsu, 215123, China
88: Ecology of Tropical Agricultural Systems, Institute of Agricultural Sciences in the Tropics,
University of Hohenheim, Garbenstrasse 13, Stuttgart, 70599, Germany
89: School of Biological Sciences , Life Sciences Building, University of Bristol, 24 Tyndall
Ave, Bristol, BS8 1TQ, United Kingdom
90: Migratory Bird Unit, Northern Region, Canadian Wildlife Service, Environment and
Climate Change Canada, 5019 - 52nd Street, Yellowknife, Northwest Territories, X1A 2P7,
Canada
91: Species and Habitat Conservation, Forest Nature Conservation, Northwest German
Forest Research Institute, Prof.-Oelkers-Str. 6, Hann. Münden, Lower Saxony, 34346,
Germany
92: Wildlife Conservation Society Canada, 169 Titanium Way, Whitehorse, Yukon, Y1A 0E9,
Canada
93: Applied Zoology and Nature Conservation, Zoological Institute and Museum, University
of Greifswald, Loitzer Strasse 26, Greifswald, 17489, Germany
94: Bat Lab, Department of Evolutionary Ecology, Leibnitz Institute for Zoo and Wildlife
Research, Alfred-Kowalke-Straße 17, Berlin, 10315, Germany
95: Cooperative Institute for Marine Ecosystem and Resources Studies, , National Oceanic
and Atmospheric Administration Pacific Marine Environmental Laboratory and Oregon State
University, 2030 SE Marine Science Drive, Newport, OR, 97365, USA
96: Department of Biology, University of Oxford, , Oxford, OX1 3SZ, United Kingdom
97: Bat Squad Lab, Department of Life Science, National Taiwan Normal University, No. 88,
Section 4, Tingzhou Rd, Wenshan District, Taipei City, 116, Taiwan
98: Nature based Solutions Initiative, Department of Biology , University of Oxford, 11A
Mansfield Road, Oxford, OX13SZ, England
99: Centre for Northern Forest Ecosystem Research, Ministry of Northern Development,
Mines, Natural Resources and Forestry, Ontario Government, 103 – 421 James Street
South,Thunder Bay, Ontario, P7E 2V6, Canada
100: Swiss Ornithological Institute, Seerose 1, Sempach, 6204, Switzerland
101: Zoological Research Museum Alexander Koenig, , Leibniz Institute for the Analysis of
Biodiversity Change , Adenauerallee 160, Bonn, 53113, Germany
102: Museum für Naturkunde, , Leibniz-Institut für Evolutions- und Biodiversitätsforschung,
Invalidenstraße 43, Berlin, 10115, Germany
103: , , , ,
104: Ecology and Nature Management, Geosciences and Natural Resource Management,
University of Copenhagen, Rolighedsvej 23, Frederiksberg, 1958, Denmark

105: Department of Agricultural Biology, Faculty of Agriculture, University of Ruhuna,
Mapalana, Kamburupitiya, x, , Sri Lanka
106: Fundación de Conservación Jocotoco, Fundación de Conservación Jocotoco,
Fundación de Conservación Jocotoco, Valladolid N24-414 y Luis Cordero, Quito, 170525,
Ecuador
107: Landscape Conservation Programme, Conservation Division, BirdLife South Africa, ,
Gauteng, , South Africa
108: Afromontane Research Unit, Department of Zoology and Entomology, University of the
Free State, , Phuthaditjhaba, , South Africa
109: Centre for Statistics in Ecology, the Environment and Conservation, Department of
Statistical Sciences, University of Cape Town, , Cape Town, , South Africa
110: Nature Coast Biological Station, Institute of Food and Agricultural Sciences, University
of Florida, 552 1st Street, Cedar Key, Florida, 32625, U.S.A.
111: Fisheries and Aquatic Sciences, Institute of Food and Agricultural Sciences, University
of Florida, 136 Newins-Ziegler Hall, Florida, 32611, U.S.A.
112: Sound Forest Lab, The Nelson Institute of Environmental Studies and The Department
of Forest and Wildlife Ecologz, University of Wisconsin-Madison, 1650 Linden Drive,
Wisconsin, 53706, U.S.A.
113: Nature Conservation Unit, , Frederick University, 7, Yianni Frederickou Street, Nicosia,
1036, Cyprus
114: Madagascar Programme, , Chances for Nature, Brauweg 9a, Göttingen, 37073,
Germany
115: Cologne Zoo, Riehler Straße 173, Köln, Lower Saxony, 50735, Germany
116: Biodiversity, Macroecology and Biogeography, Faculty of Forest Sciences and Forest
Ecology, University of Goettingen, Büsgenweg 1, Göttingen, Niedersachsen, 37077,
Germany
117: Plant Ecology, Institute of Plant Sciences, University of Bern, Altenbergrain 21, Bern,
Bern, 3013, Switzerland
118: Pacific Wildlife Research Center, Environment and Climate Change Canada, 5421
Robertson Road, Delta, British Columbia, V4K 3Y3, Canada
119: Migratory Bird Unit, Northern Region, Canadian Wildlife Service, Environment and
Climate Change Canada, 91780 Alaska Highway, Whitehorse, Yukon Territory, Y1A 5B7,
Canada
120: Lancaster Environment Centre, Lancaster University, Library Ave, Bailrigg, Lancaster,
LA1 4YQ, United Kingdom
121: Environmental Research and Innovation Centre, School of Science, Engineering and
Environment, University of Salford, Peel Building, Salford, M5 4WT, United Kingdom
122: School for Forest Management, Swedish University of Agricultural Sciences SLU, Box
43, Skinnskatteberg, SE-739 21, Sweden
123: Chair of Geobotany, Chair of Geobotany, University of Freiburg, Schaenzlestr. 1,
Freiburg im Breisgau, 79104, Germany
124: Fisheries Ecology and Conservation Lab, Department of Biology , University of Victoria,
3800 Finnerty Rd., Victoria, British Columbia, V8P 5C2, Canada
125: Department of Botany and Biodiversity Research, Faculty of Life Sciences, University of
Vienna, Rennweg 14, Vienna, 1030, Austria
126: School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat
Street, Seattle, Washington, 98105, U.S.A.
127: Polar Ecology Group, Department of Vertebrate Ecology and Zoology, University of
Gdańsk, Poland, Wita Stwosza 59, Gdańsk, , 80-308, Poland
128: Department of Ecology, University of Alicante, Cra. San Vicente del raspeig sn,
Alicante, 3690, Spain
129: Laboratório de Ecologia, Instituto de Biociências, Universidade Federal de Mato Grosso
do Sul, , , , , , Brazil
130: Institute of Marine Science, , University of Auckland, Private Bag 92019. Auckland Mail
Centre, Auckland, 1142, New Zealand

131: Department of Soil, Water, and Ecosystem Sciences, Institute of Food and Agricultural
Sciences, University of Florida, 2181 McCarty Hall A, Florida, 32611, U.S.A.
132: Integrative Community Ecology Unit, , Okinawa Institute of Science and Technology
Graduate University, 1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa, 904-0495, Japan
133: Salmonid fishes, Norwegian Institute for Nature Research, Høgskoleringen 9,
Trøndelag, Trondheim, 7034, Norway
134: Cold Climate Research Lab, School of Art, RMIT University, 124 La Trobe Street,
Melbourne, Victoria, 3000, Australia
135: Fisheries and Oceans Canada, Freshwater Institute, 501 University Crescent,
Winnipeg, Manitoba, R3T 2N6, Canada
136: Centre for Marine Science and Technology, , Curtin University, , Perth, 6102, Western
Australia
137: Fisheries Management Branch, Department of Forestry, Fisheries and the
Environment, , Foreshore, Cape Town, 8000, South Africa
138: Mammal Research Institute Whale Unit, Department of Zoology and Entomology,
University of Pretoria, Hatfield, Pretoria, 28, South Africa
139: State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,
Chinese Academy of Sciences, , Beijing, 100093, China
140: Coastal Marine Field Station, School of Science, University of Waikato, 58 Cross Road,
Sulphur Point, Tauranga, 3110, New Zealand
141: Ocean Acoustics Group, , Alfred Wegener Institute Helmholtz Centre for Polar and
Marine Research, Klussmannstr. 3d, Bremerhaven, 27570, Germany
142: Department of Ecology, Universidad Autónoma de Madrid, Darwin 2, Madrid, 28049,
Spain
143: Centro de Investigación en Biodiversidad y Cambio Global, Universidad Autónoma de
Madrid, Darwin 2, Madrid, 28049, Spain
144: Wildlife Research Division, Environment and Climate Change Canada, 801-1550,
avenue d'Estimauville, Québec (Québec) , G1J 0C3, Canada
145: Natural Resources & Environmental Studies, University of Northern British Columbia,
3333 University Way, British Columbia, V2N 4Z9, Canada
146: Asian Soundscape Monitoring Network, Biodiversity Research Center, Academia
Sinica, 128 Academia Road, Section 2, Nankang, Taipei , , 11529, Taiwan
147: Department of Ecology and Evolutionary Biology & K. Lisa Yang Center for
Conservation Bioacoustics, , Cornell University, Corson Mudd, 215 Tower Road, Ithaca,
New York, 14853, U.S.A.
148: Department of Biology, Whitman College, 345 Boyer AVe, Walla Walla, Washington,
99362, U.S.A.
149: Talarak Foundation, Inc., Negros Forest Park, Bacolod City, Negros Occidental, 6100,
Philippines
150: Department of Forest Ecosystems and Society, Oregon State University, 3180 SW
Jefferson Way, Corvallis, Oregon, 97331, U.S.A.
151: Centre for Research and Conservation, Royal Zoological Society of Antwerp, , , , , ,
Belgium
152: Department of Physical Geography, Stockholm University, SE-106 91 Stockholm,
Stockholm, 10691, Sweden
371
372   [c]: corresponding authors
373

Abstract

The urgency for remote, reliable, and scalable biodiversity monitoring amidst mounting
human pressures on climate and ecosystems has sparked worldwide interest in Passive
Acoustic Monitoring (PAM), but there has been no comprehensive overview of its coverage

across realms. We present metadata from 358 datasets recorded since 1991 in and above
land and water constituting the first global synthesis of sampling coverage across spatial,
temporal, and ecological scales. We compiled summary statistics (sampling locations,
deployment schedules, focal taxa, and recording parameters) and used eleven case studies
to assess trends in biological, anthropogenic, and geophysical sounds. Terrestrial sampling
is spatially denser (42 sites/M·km^2) than aquatic sampling (0.2 and 1.3 sites/M·km^2 in
oceans and freshwater) with only one subterranean dataset. Although diel and lunar cycles
are well-covered in all realms, only marine datasets (65%) comprehensively sample all
seasons. Across realms, biological sounds show contrasting diel activity, while declining with
distance from the equator and anthropogenic activity. PAM can thus inform phenology,
macroecology, and conservation studies, but representation can be improved by widening
terrestrial taxonomic breadth, expanding coverage in the high seas, and increasing spatio-
temporal replication in freshwater habitats. Overall, PAM shows considerable promise to
support global biodiversity monitoring efforts.

Main text

Introduction

Sounds permeate all realms on Earth — terrestrial, freshwater, marine, and subterranean[1]. Passive Acoustic Monitoring (PAM) captures soundscapes that document soniferous life ("biophony"), human activities ("anthropophony"), and some geophysical events ("geophony")[2–6]. In ecoacoustics and soundscape ecology[7,8], PAM can measure levels and impacts of global change (e.g., climate change, urbanization, deep-sea mining)[9–11], monitor ecosystem health, recovery, and restoration[12–14], assess human-environment interactions (public health, cultural ecosystem services)[15,16] and guide conservation (protected area management, landscape planning)[17,18].

Despite the wide-ranging and increasing soundscape sampling effort, the data distribution remains undescribed. Currently soundscape-recording communities are only networked within realms and their methodologies differ. Previous reviews focused on single realms and were either systematic[19–22] or qualitative[3,23]. Marine scientist networks using PAM exist[24], but the freshwater community is nascent, and the terrestrial community grows faster than it can unite. Acoustic calibration and sound propagation modeling are advanced in aquatic studies[25] but seldom considered in terrestrial ones (except [26,27]). Artificial intelligence can identify increasing numbers of species on land[28] whereas aquatic sounds still challenge identification[29,30]. Overall, there is much to gain from sharing data, experience, and methods among PAM users.

Cross-realm PAM studies can yield new theoretical answers[31] and applied solutions: soundscapes track terrestrial and marine resilience to natural disasters[32]; sound and silence durations in multiple realms follow universal distributions[33]. Transnational sampling could form the basis for comprehensive soniferous biodiversity monitoring, just like community-initiated telemetry databases[34], collaborative camera trap surveys[35], individual animal observation networks[36], and invasive species control syntheses[37] advanced entire research fields. A global PAM network could establish historical biodiversity baselines, support systematic long-term and large-scale monitoring, and connect with the public through citizen science. Such information is critical to inform global biodiversity policies such as the Kunming-Montreal Global Biodiversity Framework.

We present the "Worldwide Soundscapes" project, the first global PAM meta-database and network. We use it to quantify the known state of PAM efforts, highlight apparent sampling gaps and biases, illustrate the potential of cross-realm PAM syntheses for research, and federate PAM users. The project currently holds 307 contributors who collated metadata about 358 passively-recorded, replicated soundscape datasets. Metadata describe the exact spatio-temporal coverage, sampled ecosystems (IUCN Global Ecosystem Typology: GET), transmission medium (air, water, or soil), focal taxa (IUCN Red list), recording settings, and data and publication availability. We inferred coverage within administrative (Global Administrative Database: GADM; International Hydrographic Organisation: IHO) and protected areas (World Database on Protected Areas: WDPA) from geographic locations. We selected recordings referenced in the meta-database to quantify soundscape components (biophony, anthropophony, geophony) across eleven ecosystems from all realms. We showcase their relevance to macroecology, conservation biology, and phenology research and identify opportunities to advance the global PAM network. The publicly-

437 accessible meta-database[38,39] continues to grow to enhance accessibility of data and
438 remains open for metadata contributions, facilitating future syntheses.

Results

Summary dataset statistics

To date, 358 validated soundscape meta-datasets (hereafter "datasets") have been
registered in our database from across the globe, dating back to 1991 (Fig. 1D). A dataset
gathers a team's metadata on a study or project. Metadata were validated by checking that
all fields were filled in and formatted correctly, and then cross-checked by their respective
contributors using site maps and recording timelines[39] on ecoSound-web[40]. Based on the
GET definition of four 'core' realms, our database includes 259, 80, 18, and 1 validated
datasets from the terrestrial, marine, freshwater, and subterranean realms, respectively. The
transmission medium was air for terrestrial and subterranean datasets and mostly water for
aquatic datasets, as three datasets with aerial above-water recordings were included in the
freshwater and marine realms. The majority of datasets (84%) include both spatial and
temporal replicates (Fig. 1C). Few datasets have openly-accessible recordings (10-15%, Fig.
S1). Presently, few terrestrial and freshwater datasets (27% and 22% respectively) are
associated with DOI-referenced publications in contrast to marine datasets (44%) (Fig. S1).

Spatial sampling coverage and density

The database contains 11 093 sampling sites, including 147 polar, 8 455 temperate, and 2
491 tropical sites (Fig 1C). On land, 10 387 sites are located within 79 (out of 263) GADM
level 0 areas (Fig. 2B), primarily in the Northern Hemisphere (Table S1). Most terrestrial
sites occur in Canada (28%), followed by the United States (18%), but a significant
proportion is widely distributed (23% do not belong to the top 10 GADM areas). Few
terrestrial sites (8%) are located in WDPA category Ia, Ib, or II areas. Our database currently
lacks data from vast areas in Russia, Greenland, the Antarctic, interior Australia, North
Africa, and Central Asia. Site elevations range from sea level up to 3 420 m (Fig. 2A), but
mountains above 4 000 m in the Northern Hemisphere, above 2 000 m in the Southern
Hemisphere, as well as the Transantarctic Mountains are currently not represented in the
data. At sea, 469 sites are located within 32 (out of 101) IHO sea areas. Most marine sites
occur in the North Pacific Ocean (22%), followed by Southeast Alaskan and British
Columbian coastal waters (17%), but a significant proportion is widespread (17% do not
belong to the top 10 IHO areas). Many sites are situated in WDPA high-protection category
Ia and II areas (14%). Our database currently lacks datasets from Arctic waters off Eurasia,
the Southeast Pacific, and Southeast Asian coastal areas. Sampling sites span ocean
depths from sea water surface to depths of 10 090 m, but tropical bathypelagic and Southern
benthic areas are poorly represented (Table S2). Few GADM areas (11) are represented in
the freshwater datasets. Spain holds most freshwater sites (56%). Few freshwater sites (2%)
are in WDPA category II or Ib areas. Freshwater bodies are sampled at elevations from sea
level up to 950 m. Mountain freshwater bodies and those in Africa, Asia, and Oceania are
currently not represented in our database. The database contains seven subterranean sites
situated in Brazil between 277 and 810 m.

Temporal sampling extent, coverage, and density

We compare sampling coverage (in sampled years summed over sites) across time windows
of the diel cycle (dawn, dusk, day, night), lunar phases (bright and dark), and the seasons

(spring, summer, autumn, winter - only for temperate sites) (Fig. 3A). Dawn and dusk diel
windows are shorter than day and night diel windows for most locations and less intensively
sampled. The lunar phase cycle is evenly covered across realms and comprehensively
covered within datasets (Fig. S2). In the terrestrial realm, daytime coverage surpasses
nighttime coverage (611 vs. 398 years), while 75% of datasets sampled all diel time
windows. Terrestrial datasets mostly sampled spring (415 years, 32%) and summer (564
years, 44%) while 24% sampled all seasons. Temporal coverage per site is highest in Japan
(272 days) and Taiwan (250 days). In the marine realm, temporal coverage among diel time
windows is even and 88% of marine datasets sampled all diel time windows. Marine
datasets have high and similar coverage for winter and spring (117 and 122 years, combined
59% of seasonal coverage) and 65% cover the full seasonal cycle. In the freshwater realm,
temporal coverage among diel time windows is even and 76% of freshwater datasets
sampled all diel time windows. Similarly, 24% sampled all seasons. The subterranean
tropical sites primarily covered the nighttime, across all seasons.

Sampling in ecosystems

Our database includes 76 of the 110 GET functional groups and all biomes except
anthropogenic shorelines and most subterranean biomes (Table S2). Sampling intensity
differed across ecosystem levels - realms, biomes, and functional groups, with transitional
realms representing the interface between core realms (Fig. 4A). The terrestrial realm has
the third-largest extent and the highest spatial sampling density among realms (42 sites per
Mkm^2), but temporal coverage is comparatively low (17% sampled out of mean extent per
site: 155 days). The most commonly sampled biome is temperate-boreal forests and
woodlands (59% of sites). The marine realm is the most extensive and spatial sampling
density is the lowest (0.2 sites per Mkm^2), but temporal sampling extent and coverage are
the highest among all realms (69% out of 497 days sampled). The most commonly sampled
biomes are the marine shelf and pelagic ocean waters (43% and 42% of sites respectively).
The freshwater realm has low spatial sampling densities (1.3 sites per Mkm^2) and high
temporal sampling densities (30% out of 115 days sampled). Rivers and streams are most
commonly sampled (47% of sites). The terrestrial-freshwater realm, representing 81% of the
area of non-subterranean realms, has the third-highest spatial sampling density (7.9 sites
per Mkm^2) and similar temporal sampling density to the terrestrial realm (16% days sampled
out of 230). The subterranean realm, though second-largest, includes seven tropical sites in
aerobic caves from one dataset.

Target taxa and frequency ranges

Most marine datasets do not target specific taxa (70%) and use wide frequency ranges from
0.01 to 30 kHz (mean bounds of frequency ranges across datasets, Fig. 4B). Marine
datasets that focus on single taxa comprise cetaceans (8%, 0.006 - 7 kHz) and fish (8%,
0.003 - 29 kHz). Similarly, most freshwater datasets are taxonomically unspecific (66%) and
cover frequencies from 1 to 22 kHz. Some datasets (20%) focus on ray-finned fish, covering
frequencies from 0.001 to 23 kHz. In contrast, terrestrial datasets mostly target single taxa
and have narrow frequency ranges. Bird-focused datasets are most common (44%),
spanning frequencies from 0.06 to 21 kHz, while bat-focused datasets are next (12%) and
range from 5 to 139 kHz. Taxonomically unspecific datasets account for 24% of datasets,
covering a broad range from 0.1 to 24 kHz. Generally, datasets targeting multiple taxa use
wider frequency ranges than those targeting single taxa.

Soundscape case studies

We analysed 150 recordings from eleven different ecosystems representing the diversity of soundscapes on Earth, spanning latitudes from 69 degrees South to 67 degrees North (Table S3, Fig. 5A). Biophony dominated with an average soundscape occupation of 30% across all ecosystems, notably: photic coral reefs (Okinawa, Japan) with snapping shrimps and grunting fish choruses (75% soundscape occupancy)[41]; tropical lowland rainforests (Jambi, Indonesia) with buzzing insects and echoing bird and primate songs (61%). Only marine island slopes (off Sanriku, Japan) and polar outcrops (Antarctic) contained no or very little (2%) biophony, respectively. Geophony was absent in most soundscape samples, with the exception of high wind noise in polar outcrops (14%) and some wind in montane tropical forests (5%). Anthropophony occupied on average 9% of the soundscapes. Cities (Jambi, Indonesia; Montreal, Canada) exhibited the highest anthropophony (45%) with prevalent engine noise and human voices, while deep-sea mining and vessel communication caused high anthropophony in marine island slopes (32%)[42]. Silence occupancy was highest in polar outcrops (82%) and large lowland rivers (78%).

The selected soundscapes reveal greater biological activity closer to the equator, a negative relationship between biophony and anthropophony, and variable phenology of soniferous organisms over the diel cycle (Fig. 5). All Bayesian beta regression models for biophony occupancy converged. We detected a negative correlation of biophony occupancy with increasing distance from the equator ($P_{negative}$=1) and with anthropophony occupancy ($P_{negative}$=1). The phenology model predicted biophony occupancy values for each diel time window and realm (Fig. 5B), revealing similar phenology for the terrestrial and marine realm, and opposed freshwater and freshwater-marine realm phenology.

Discussion

The "Worldwide Soundscapes" project has — to our knowledge — assembled the first global meta-database of PAM datasets across realms. We analysed its current content to quantify sampling extent, coverage, and density across spatiotemporal and ecological scales. We annotated soundscapes from eleven ecosystems to investigate macroecological, conservation biology, and phenological trends. The database remains open for contributions[39] and can be openly accessed to source datasets and initiate collaborative studies[38]. Next, we discuss the state and potential of PAM globally (Table 1).

Our results likely represent global PAM trends, even though spatial gaps reflect the background of the project contributors. Our database still misses some national programs (e.g., Australian Acoustic Observatory[43]), but otherwise our terrestrial coverage is similar to a recent systematic review[21]. The gaps in North Africa and Northeastern Europe correspond with the paucity of bioacoustic datasets for these regions[44]. Our database comprises 469 marine sampling locations while 991 were found in a recent systematic review[20], but most overlap (Fig S2). Marine tropical waters that are under-represented in our database reflect gaps found in the International Quiet Ocean Experiment network coverage[45]. Our marine and terrestrial database coverage is thus broadly comparable with published data. However, as the database originates from an active network of researchers, it represents the current availability of mostly as-yet unpublished data (Fig. S1). To our knowledge, no other spatially-explicit review of freshwater sampling exists for comparison, and no other work quantified temporal coverage.

Geographic coverage strikingly differs between realms: marine coverage is sparse but
widespread; terrestrial coverage is comparatively intensive in the Americas and Western
Europe; freshwater coverage is scattered. This partly reflects funding priorities for where
conservation and active management priorities lie, for instance in biodiversity hotspots and
densely-populated areas. Technical limits in extreme environments also drive geographic
patterns: high latitudes and elevations entail extremely cold temperatures that present
challenges for operation and maintenance, some of which can be solved with robust power
setups (solar panels, freeze-resistant batteries). Marine deployments are generally even
more constrained due to costly and demanding underwater work, but some deployments
reach the poles as water temperatures are buffered below the freezing point. Affordable
underwater recorders[46] may help to intensify sampling of marine coastal areas and close
gaps in freshwater coverage. By contrast, terrestrial monitoring is generally straightforward.
Northern temperate areas outside of Northeastern Europe are comparatively better covered,
and the tropics outside of Africa better represented. It appears that gaps in North Africa,
Central Asia, and Northeastern Europe arise from differential research means and priorities
between countries. Taken together, these gaps help to correct spatial biases[47] and to identify
high-priority, unique research areas that should be included in global assessments.

Currently, only marine studies achieve relatively even coverage of temporal cycles. Indeed,
offshore deployments — especially in the deep sea — are expensive and limited duration
deployments are not cost-effective[48]. Marine soundscapes fluctuate stochastically[49], but the
ocean buffers water temperatures so that animals retain a basal activity level year-round.
Although lunar phases affect marine life[50–52], we did not consider lunar tides, which affect
some ecosystems. In the terrestrial and freshwater realms, most deployments cover the
entire diel cycle but monitoring on land focuses either on diurnal birds or on nocturnal bats,
as found in the literature[21]. In contrast, although seasons drive activity cycles on land too[53,54],
spring- and summertime monitoring is disproportionately common and we lack a thorough
understanding of seasonal dynamics. Terrestrial deployments in particular may be short for
logistic reasons: in the cold, batteries struggle and access is harder; recorders are at risk of
theft; and limited numbers may cycle between sites[55]. Lunar phases — probably evenly
sampled by chance — also influence land animals[56] but should be explicitly considered.
Overall, we encourage longer-duration setups with regularly-spread sampling inside
temporal cycles to alleviate the higher expenses, energy consumption, storage, and traded-
off spatial coverage[57]. Global changes impact soundscapes in largely unpredictable ways
through changing species distributions and phenology, necessitating higher and unbiased
coverage across multiple time scales — including inter-annual ones — to successfully
monitor ongoing changes[58].

Re-use of soundscape datasets is restricted by their taxonomic focus. Admittedly,
taxonomically untargeted, long, and regular deployments in oceans, coupled to large
detection ranges, concurrently sample many taxa[59]. However, many soundscape recordings
sample particular frequencies, often in the human-audible range[60], although biophony ranges
from infrasound[61] to ultrasound[62]. For instance, studies of toothed whales or bats often use
triggers and high-pass filters to record purely ultrasonic recordings only when signals are
detected, resulting in spectrally-restricted and temporally-biased soundscape recordings.
Less-studied taxa such as anurans and insects could effectively be co-sampled by adjusting
ongoing deployments, so we encourage terrestrial researchers to maximise frequency
ranges to enhance interdisciplinary collaboration. These collaborations can help to mutualise

resources for mitigating potentially prohibitive power, storage, transportation, and post-
processing costs[63]. Emerging embedded-AI audio detectors may offer an alternative[64], but
soundscape recordings will remain essential for broader application.

In every realm, ecosystems await acoustic discovery. Except for one dataset from aerobic
caves, we lack data from all subterranean realms (anthropogenic voids, ground streams, sea
caves), while endolithic systems may be the only irrelevant ecosystem for biophony. Access
is usually challenging or restricted for non-specialists, but subterranean biodiversity shows
high spatial turnover[65]. Freshwater datasets were less rare, but several ones with
unreplicated sampling could not be included. Temporary, dynamic water bodies (seasonal,
episodic, and ephemeral ecosystems) are not yet well-studied[66] although they are accessible
from land. Notably, sounds can pass the boundary between recording media and be
captured in so-called holo-soundscapes in freshwater and shallow coastal areas[5,67,68].
Advances are imminent as the freshwater acoustic research community is growing
rapidly[69,70]. In the oceans, sound propagates far and multiple biomes can be sampled at once
(e.g., recorders on the seafloor sample pelagic waters) so that most ecosystems are
covered. Still, coverage gaps may exist in rhodolith/Maërl beds, upwelling zones, and
deepwater coastal inlets. On land, sampling is biased towards biodiverse forests, and our
database coverage suggests that the most challenging terrestrial ecosystems, such as arid
zones (sclerophyll hot deserts and semi-deserts, semi-desert steppes), rocky habitats
(young rocky pavements, lava flows and screes) but also some vegetated temperate
ecosystems (cool temperate heathlands, temperate pyric humid forests, temperate
subhumid grasslands) are poorly sampled. Within the IUCN GET framework, soil
soundscapes also belong to the terrestrial realm and to date no eligible datasets are in our
database, despite recent studies[71,72].

Our database highlights well-known global sampling biases[73] which could be resolved with
collaboration to remove cultural and socioeconomic barriers[74]. Technological progress for
more affordable equipment renders PAM more accessible in lower-income countries.
However, high- and deep-sea work remains considerably more expensive, and tropical
developing countries in particular often lack funding for marine programmes requiring large
vessels, underwater vehicles, or cabled stations on the seafloor[48]. Our network currently
consists of active, English-speaking members from 50 countries. Collaborative projects,
shared sea missions, and equipment loans should promote the establishment of soundscape
research communities[75]. Increased international collaboration with scientists and local
stakeholders supporting citizen-science[76] in heavily underrepresented regions would improve
not only data coverage, but also representation and dialogue within the field.

Collaborative soundscape research relies on interoperable data. We harmonised metadata
with a bottom-up approach leading to our global inventory, but comprehensive standards for
PAM do not currently exist, even though initiatives for the marine realm are ongoing[77,78]. Few
affordable solutions exist for sharing large audio data volumes[40], underlining the need for
distributed soundscape recording repositories[79]. Marine oil and gas industry projects
routinely upload data as part of their efforts to mitigate noise impacts on marine animals[80],
but these recordings often focus on frequencies relevant to seismic prospecting and access
may be restricted[81,82]. Furthermore, recording equipment requires calibration, sound
detection spaces need to be measured[26], and data privacy must be ensured on land[83]. In
parallel, species sound libraries[44,84–86] grow and continue to provide invaluable acoustic and
taxonomic references. International organisations such as GBIF will be key to roll-out

standards (e.g., Darwin Core) in a top-down manner. For the moment, we encourage early
planning of data archival. In the future, the Worldwide Soundscapes will interoperate with
other databases to close remaining coverage gaps.

A unified approach of ecology with PAM is now possible. More comprehensive coverage is
needed to decisively answer research, conservation, and management questions that PAM
can address, so we encourage prospective contributors to archive their metadata and join
our inclusive project. We show that a large portion of the PAM community is willing to
collaborate across realms and form a global network. We advocate for a bolder PAM effort to
inform the agenda of soundscape ecology[87], reaching out to places where no sound has
been recorded before as well as to urban settings[88]. The research community may open new
avenues to study environmental effects on acoustic activity[58], social species interactions[89],
human-wildlife relationships[41], function and phylogeny[90], soundscape effects on human
health[91], acoustic adaptation and niche hypotheses[92,93], macroecological patterns across
ecosystems[1], and initiate an integrated approach to noise impacts on wildlife.

PAM is now an established method that can be applied over large spatial and temporal
scales. Consistent, large-scale monitoring of the Earth's soundscapes is essential to
establish baselines for historical trends[94] and quantify rapid changes in biodiversity and
natural systems. International funding schemes should integrate PAM into biodiversity
monitoring platforms such as GBIF[95] and GEO BON[96,97]. Soundscapes are just starting to be
used in legislation as an ecosystem feature to be preserved[98]. Occupancy maps for
soniferous wildlife obtained from PAM would underpin the evaluation of progress towards
threat reduction and ecosystem service provision of the Kunming-Montreal Global
Biodiversity Framework[99]. By building collaborations around the knowledge frontiers
identified here, we can aim to comprehensively describe and understand the acoustic make-
up of the planet.

Methods

Ethics & Inclusion statement

The present study has involved people who carried out PAM-based studies as primary contributors. They could be corresponding authors for published studies, referred contacts for unpublished studies, or principal investigators, and were asked to identify further primary and secondary contributors of their study. Primary contributors could become co-authors, and secondary contributors are acknowledged here. Some primary contributors were invited as co-leads to be responsible for particular realms or biomes and are listed in the first-tier authors list. Primary contributors who additionally provided soundscape recordings are also listed in the first-tier authors list. All primary contributors were asked to identify further contacts to reach a comprehensive coverage for the database. Given the nature of the study (using passive acoustic monitoring methods), the research did not suffer from restrictions or prohibitions in the setting of the researchers. Necessary permits from landowners or environmental protection agencies, precautions to limit biological contaminations (e.g., biofilm on marine recorders) or for animal welfare, and local ethics committee reviews were the responsibility of the respective primary contributors. None of the PAM activities resulted in personal risk to participants; neither did they involve health, safety, security or other risk to researchers. Local and regional research relevant to our study were taken into account in citations throughout the main text.

Database construction

The database construction started in August 2021 within the frame of the Worldwide Soundscapes project[38] using collaborative, peer-driven metadata collation. It represents the current state of knowledge about publicly accessible meta-datasets for PAM within our network. We started contacting contributors that we personally know, and co-lead authors helped to contact potential contributors in the respective ecosystem types of their expertise (i.e. terrestrial, urban, freshwater, and marine). We conducted focal publication searches to actively plug coverage gaps by inviting the respective corresponding authors. We posted the call for contributors on specialised ecoacoustics platforms and social media, and will keep the project open for any contributor owning suitable soundscape recordings beyond the present publication. We communicated mostly in English but also in Spanish, French, Portuguese, German, Russian, and Chinese to gather metadata. We ended up including metadata from larger groups such as the Silent Cities project, Ocean Networks Canada, the Australian Acoustic Observatory, and Parks Canada.

Primary contributors provided the metadata that formed the basis for the database. Their willingness to be informed through a mailing list, responsibility for their metadata, approval for sharing the meta-data publicly, and willingness to participate in this study as co-authors were explicitly stated in an online form-based collaboration agreement. Primary contributors who became co-authors all fulfilled either data curation (e.g., as providers of structured metadata) or project administration (e.g., as principal investigators designing the corrresponding study) roles and additionally a manuscript revision role. Primary contributors cross-validated their metadata using maps and graphical timelines visualising their input. As a result, apart from basic coherence checks conducted by the co-leads and research assistants, the responsibility of the database content is borne by their contributors. The

database information page is integrated in our online collaborative ecoacoustics platform
ecoSound-web[100] that also hosts the soundscape recordings of the case study
(https://ecosound-web.de/ecosound_web/collection/show/49).

Our network consists of people reachable through email contact. Out of 588 contacted
contributors potentially involved in PAM, 298 (49%) provided some metadata. More datasets
exist but 24% of those contacted have not yet responded. This may be due to limitations in
sharing research data or, newcomers to PAM, or out-dated contact details. Older
soundscape datasets may be stored on analog media that are inaccessible over the internet,
but these are more often attended recordings[53] — thus not fulfilling our inclusion criteria (see
below). The missing share of datasets could be estimated with a systematic literature
search, but we posit that our dataset search was exhaustive, and we were able to fill gaps
more effectively due to focused calls for missing datasets in our large group of contributors.

Potential soundscape recording datasets were required to meet four criteria: 1) stationary —
mobile recorders would have complex, constantly changing spatial assignments, thus we
excluded recordings from cars, transect walks, or towed deployments; 2) passive —
obtained from automated, unattended recorders, without human presence near the acoustic
sensor; 3) ambient — with no particular recording direction or temporal selection, as
obtained from omnidirectional microphones and with non-triggered recordings; 4) from
spatially or temporally replicated study designs (Fig. 1). Datasets with spatial and temporal
replication are necessary to disentangle spatial and temporal effects from other soundscape
determinants. We considered datasets to have spatial replicates when several sites were
sampled simultaneously, and temporally replicated when one site was sampled over multiple
days at the same time of day. We chose sampling sites and days as the most elemental
units for defining replication in our unified analysis. However, we acknowledge that replicates
are defined differently in each study: for instance, sampling sites may only be spatial
replicates if they belong to the same category (i.e. habitat, management type). Temporal
replicates could also be defined at the scale of other solar and lunar cycles (e.g., multiple full
moons). Taken together, our requirements homogenise the dataset to enable statistical
analyses across datasets and future collaborative syntheses.

Time and space

Soundscapes are fundamentally determined by their time and location. Wildlife and human
activities are broadly determined by solar and lunar cycles[32,54,56,101], and geographical
positions on the planet relative to the poles or equator, or the land and water surface. For all
datasets, we determined their spatial coverage in terms of number of sites, and sampling
density as the number of sites within the realms' areal extent (as determined by IUCN GET
maps). Although spatial extent could be defined as the area bounded by the sites of a
dataset, challenges with calculating extents on the world sphere (especially for datasets with
very large extents spanning the globe), and unknown sampling areas covered in each site
led us to ignore spatial extents. Until sound detection spaces can be accurately measured,
our measure of spatial sampling density using points per area is thus provisional. Indeed,
acoustic sampling areas or volumes are extremely rarely measured in terrestrial sites and
seldom measured but sometimes simulated in marine environments[102]. They vary with sound
source intensity, frequency, directivity; recording medium temperature, currents, pressure,
humidity (for air); habitat structure such as topography and vegetation structure; ambient
sound level. Generally, detection ranges are greater in underwater environments due to the

higher density of the recording medium. For all datasets, we also determined their temporal
extent (i.e., the time range from the start of the first to the end of the last recording),
coverage (i.e., time sampled per site), and density (i.e., proportion of recorded time inside
the temporal extent).

We quantified the latitudinal and topographical coverage across realms by collecting latitude,
longitude and topography (above and below the sea level) data for each site. Contributors
could provide self-measured topography values for sites on land and gaps were filled in
using General Bathymetric Chart of the Oceans surface elevation data. For underwater sites,
depth values (below sea level for marine sites, below the surface for freshwater) were
provided by the contributors. We assigned sampling sites to administrative areas (GADM
divisions for freshwater and land, IHO for sea areas) and extracted their WDPA category.
Sites' climates were geographically classified into tropical (between -23.5° and 23.5°
latitude), polar (below -66.5° or above 6`.5° latitude), and temperate (between polar and
tropical regions).

Contributors were asked to specify the timing of their deployments using start and end dates
and times, operation modes (continuous, scheduled, periodical). For scheduled operation
schedules (daily start and end times or durations) were required, and duty cycles were
required for periodical operation. Duty cycles could additionally be indicated for scheduled
deployments. Deployments were assigned to single sampling sites. Subsequently added
sites and sampling repetitions were considered as additional deployments within datasets. A
temporal framework was used to quantify sampling coverage in three solar and lunar cycles.
Temporal information was inferred from the timing of sound recordings relative to time
events that structure life and meteorological events on Earth (Fig. 3). Seasonal cycles were
inferred only for temperate sites, by splitting the year cycle into four meteorological seasons
(winter: December-February, spring: March-May, summer: June-August, fall: September-
November). The daily cycle was split into four windows delimiting dawn (from astronomical
dawn start at -18° solar altitude until 18° solar altitude), day, dusk (from 18° solar altitude to
astronomical dusk end at -18° solar altitude), and night. The lunar illumination cycle was split
into two time windows centred on the full and new moon phases. It follows that the extrema
and ecotones in the temporal cycles define the relevant sampling time windows, noting that
in temperate zones, the equinoxes roughly correspond to thermal ecotones. Seasonal cycles
in tropical and polar regions arising from precipitation patterns were not considered in this
analysis.

Ecological characterisation

We assigned individual sampling locations to ecosystem types following the IUCN GET
(https://global-ecosystems.org). Correspondiongly, sites were assigned hierarchically to
realms (core or transitional ones), biomes, and functional groups. We calculated the major
and minor occurrence areas of all functional groups based on ecosystem maps[1,103,104]. We
used these data to quantify spatio-temporal extent, coverage, and sampling density within
realms and biomes, and to identify sampling gaps. To identify which taxa acoustic
deployments were designed for, deployments were linked to IUCN taxa (class, order, family,
or genus) when applicable — some datasets were collected without taxonomic focus. We
use these data to depict the acoustic frequency ranges depending on the target taxon.

Acoustic frequency ranges

Sound recordings store a representation of the original soundscape, so we needed to determine the spectral scope of the recording datasets. Microphones have variable frequency responses, and for digital recorders the spectral scope of the recording is limited by the sampling frequency. Contributors were asked to provide the audio parameters of their deployments (sampling rate, high-pass filter) as well as auxiliary metadata about the microphone and recorder model and brands. The frequency range is additionally determined by microphone roll-off frequencies at the low end and in general, by their frequency response, but we could not gather these detailed, often unpublished data for the high number of equipment types used here.

Soundscape case studies

To illustrate how the database can be used for macroecology, conservation biology, and phenology analyses, we selected 121 recordings across a variety of topographical, latitudinal, and anthropisation conditions — all fundamental gradients of assembly filters in both terrestrial and marine realms[1] — belonging to 10 functional groups. We aimed to select datasets containing four spatial replicates within the same functional group, with 10-minute recordings spanning all four diel time windows from the same date during the biologically active season, but the available data did not always allow this (Table S2). We extracted recordings starting at sunrise, solar noon, sunset, and solar midnight. Recordings covered the audible frequency range with sampling frequencies of at least 44.1 kHz in single or dual channel configuration. We acknowledge that this targeted, non-systematic selection is not statistically representative of global patterns but rather illustrative of the database potential.

Each recording's spectrogram (i.e., visualisation of sound intensity along time and frequency axes) was annotated with the three fundamental soundscape components: biophony, anthropophony, and geophony. Soundscape recordings were uploaded to ecoSound-web (https://ecosound-web.de/ecosound_web/collection/show/49)[40] for annotation: KD listened to them and visually inspected the spectrograms (Fast Fourier Transform window size of 1 024) at a visual density of 1 116 pixels per 10 minutes to create annotations. Annotations were rectangular tags on the spectrogram, encompassing only the annotated sound, with defined coordinates in the time and frequency dimensions. Annotations of different soundscape components could overlap if they were simultaneously visible or audible. Annotations of the same soundscape component were adjacent and non-overlapping to avoid double-counting. Soundscape components occurring above 22.05 kHz were excluded from the analysis as we focused on audible frequencies only, which were the frequencies targeted by most recordings. Sounds caused by microphone or recorder self-noise were excluded. All annotations were reviewed by the recording creators using the peer-review mode on ecoSound-web, and only accepted ones were used in the analysis. The annotations were exported and acoustic space occupancy for each soundscape component in each recording was calculated as the proportion of the spectro-temporal space[60] (i.e., duration multiplied by frequency range, divided by total area of spectrogram; range: 0-1).

All Bayesian beta regression models (4 chains of 1000 sampling iterations with 1000 warmup iterations) for biophony occupancy converged as determined by trace plots and R hat values equaling 1. The models using latitude and anthropophony as predictors included

862   the functional group as a random intercept. The phenology model used diel time windows

863   and realms, as well as their interactions, as predictors.

Display items

Table 1: An agenda towards comprehensive coverage and a global PAM network

Aims	Opportunities
Increasing spatial and ecosystemic coverage	- Affordable underwater recorders - Autonomous setups for extreme environments on land - Deploy in subterranean realm - Deploy in dynamic freshwater bodies - Deploy at high latitude and elevation on land
Increasing temporal and taxonomic coverage	- Longer deployments with duty cycles - Cover all seasons in freshwater and on land - Scale up to inter-annual cycles - Higher sampling frequencies - Disabling high-pass filters and triggers
Increasing collaboration	- Work interdisciplinarily with speleologists, urban ecologists, soil and deep-sea benthic scientists - Foster international missions to work at large scales - Collaborate, invest, and fund work with under-represented countries - Disseminate results in more languages
Interoperability	- Calibrate equipment and consider detection ranges - Build accessible and sustainable data repositories - Design and adopt standards - Integrate PAM into biodiversity and remote sensing databases

Figure 1: Overview of Worldwide Soundscapes meta-database: A) Framework used to define spatial and temporal replicates. B) Number of datasets in each core realm for the different replication levels. C) Spatial extent and coverage, based on sampling sites, split by core realm. Due to their higher representation and to avoid overlapping site clusters, terrestrial site densities were plotted on a 3 degree resolution raster (Interactive map: https://ecosound-web.de/ecosound_web/collection/index/106). D) Temporal extent and coverage, based on recorded days, split by core realm. An enlarged version of panel D without terrestrial sites can be found in Fig S3. For panels C and D, Sites from transitional realms were assigned to their parent core realm.

Figure 2: Spatial distribution of sampling sites. A) Latitudinal and topographic distribution of sampling sites across core realms. Due to their higher representation and to avoid overlapping site clusters, terrestrial sites are shown with transparency. The minimum (deepest seafloor) and maximum (highest elevation of land or sea level) topographical limits (dark grey lines) are shown against latitude, based on General Bathymetric Chart of the Oceans data[105]. Minimum topography above the sea level and maximum topography under the sea level were set to zero as the sea level represents the minimum and maximum in these cases. B) Number of sampling sites within different administrative regions (GADM

level 0 and IHO sea areas), split by core realm, across WDPA categories (Ia: strict nature reserve; Ib: wilderness area, II: national park). The areas that do not belong to the top 10 in terms of datasets have been aggregated under "others". One subterranean site in Brazil is not shown. Sites from transitional realms were assigned to their parent core realm.

Figure 3: Temporal sampling distribution. *A) Temporal sampling coverage across solar and lunar cycles for the three core realms. Cycles consist of solar (daily and seasonal) and lunar time cycles (lunar phase), subdivided in time windows. Seasons were only analysed in temperate regions. Sampling coverage is represented with sampling days in number labels. B) Mean number of sampling days per site within different administrative regions (GADM level 0, IHO areas, WDPA categories), split by core realm. WDPA categories (Ia: strict nature reserve; Ib: wilderness area, II: national park) are shown separately. One subterranean site in Brazil is not shown. Numbers to the right of bars indicate the number of sites the means were calculated from. Sites from transitional realms were assigned to their parent core realm.*

Figure 4: Sampling distribution across ecological scales. *A) Spatial extent of realms, based on major areas according to IUCN GET (coloured disk area proportional to area); spatial sampling density (in sites per Mkm^2) and coverage (in number of sites); temporal extent (mean range between first and last recording day), coverage (days sampled per site and density (proportion of days sampled per extent). B) Frequency ranges of datasets across realms (using Nyquist frequency i.e., actual recorded frequencies) for the main studied taxa. The dots at the ends of colored lines represent means of the lowest and highest recorded frequencies, and the range between the minimum and maximum of these values are indicated with black error bars. The limits of human hearing are indicated with dashed lines. Number of datasets indicated above lines - datasets can be counted several times if they contain deployments targeting different taxa.*

Figure 5: Soundscape components analysis. *A) Mean acoustic space occupancy of soundscape components (biophony, geophony, anthropophony) as calculated from annotations for the selected ecosystems, measured in proportion of spectro-temporal space used, across eleven ecosystems over 150 recordings covering the time windows of the diel cycle. Annotated recordings are accessible at https://ecosound-web.de/ecosound_web/collection/show/49. Sample spectrograms are shown in the background. B) Soundscape component occupancy data illustrate three research questions linked to macroecology (i.e., biophony with distance from equator relationship), conservation biology trade-offs (i.e., biophony with anthropophony relationship); and phenological trends (i.e., mean biophony along diel time windows for each realm). Gray ribbons indicate 95% credible intervals and numbers indicate probabilities of positive or negative relationships.*

Data availability

- [Database metadata](#) at Zenodo (version 2.0.0)
- [Demonstration collection](#)

Acknowledgements

We are thankful to our colleagues who supported our work: Carolina Gonzaga, Ellen Kenchington, Jinshan Xu, Abigail Seybert, Abram B. Fleishman, Adriana C. Acero-Murcia, Adrien Charbonneau, Adrià López-Baucells, Aklavik, Alain Paquette, Albert García, Alexander C. Lees, Alexandra Buitrago-Cardona, Alfonso Zúñiga, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Allan G Oliveira, Alpheous, Ambroise N. Zongo, Ana Rainho, Ana Širović, Anamaria Dal Molin, Andros T Gianuca, Annalea Beard, Anne Sourdril, Anthony Tan, Anushka Rege, Aran Mooney, Arne Wenzel, Astrid Brekke Skrindo, Australian Institute of Marine Science, Base de Estudos do Pantanal - UFMS, Bastien Castagneyrol, Bea Maas, Bibiana Gómez-Valencia, Bob Dziak, Bobbi J. Estabrook, Borja Milá, Brian Miller, Bridget Maher, Brittnie Spriel, C. Lisa Mahon, Carolina Ocampo-Ariza, Carolline Z Fieker, Catherine Potvin, Catrin Westphal, Charlotte Roemer, Chistophe Thébaud, Chong Chen, Christina Ieronymidou, Christine Cassin, Christine Erbe, Christopher Clark, Cicero Simao, Claire Attridge, Claudia A. Medina-Uribe, Clément Lemarchand, Colin Bates, Colin KC Wen, Conner Partaker, Cowichan Tribes Lulumexun department - Fisheries and Marine projects staff, Craig A Radford, Curtin University, Cécile Albert, Dadang Dwi Putra, Daisy Dent, Dan Warren, Daniel Mihai Toma, Danny Swainson , Dawit Yemane, Deanna Clement, Dedi Rahman, Dennis Higgs, Dennis Kühnel, Dexter Hodder, Diego Alejandro Gómez-Montes, Diego Balbuena, Diego Gil, Diego Pavón Jórdan, Diogo Provete, Dorgival Diogenes, Dustin Whalen, Edho Walesa Prabowo, Eduardo Rodríguez Martinez, Elizabeth Clingham, Eliziane Oliveira, Ellen Mc Arthur, Elma Kay, Eloy Revilla, Emanuel Stefan Baltag, Emilia Sokolowska, Emilio Tan, En-Hao Liu, Enoc Martinez, Eric Parmentier, Eric Rakotomalala, Erich Fischer, Erika Berenguer, Estância Mimosa Ecoturismo, Evan Economo, Fabio De Leo Cabrera, Faisal Ali Awanrali Khan, Fang-Yee Lin, Fannie W. Shabangu, Filipe F de Deus, Flávio Rodrigues, Francis Juanes, Frederic Sinniger, Frode Fossøy, Gabriel Leite, Gabriel leite, George Best, Gerard Bota, Germany, Giselle Mahung, Greg and Lisa Morgan, Groupe Chiroptères de Guyane, Guillermo McPherson, Hailey Davies, Hajar, Hannah Reyes, Hanneline Smit-Robinson, Harris Papadopoulos, Heriniaina Randrianarison, Hervé Jourdan, Hila shamon, Hiromi Kayama Watanabe, Holger Kreft, Ian Avery Bick, Ilias Karmiris, Irfan Fitriawan, Isabelle Côté, Italo A A Rondon, Jacques Keumo Kuenbou, Jakob Tougaard, Jamie Ratliff, Jamieson Atkinson, Janice Ser Huay Lee, Jason Gedamke, Jean Pierre Castro Namuche, Jean-Yves Royer, Jeff Robinson, Jennifer Clark, Jens-Georg Fischer, Jessica Deichmann, Jessica Qualley, Jinshan Xu, Joaquin del Rio, Johan Diepstraten, Johanna Järnegren, John Ewen, John Hildebrand, John Ryan, Jos Barlow, Jose Leon, Jose Manuel Ochoa-Quintero, José Luis Mena, José Nilson Santos, José Wagner, Juan Diego Tovar, Juliana Carolina Herran García, Julie Perrot, Julio Baumgarten, Junior Tremblay, Just T. Bayle Sempere, Justine Magbanua, Jérémy Anso, Jérémy Froidevaux, Jérémy Froideveaux, Jérôme Sueur, Karen H. Beard, Katie Huong, Katie Innes, Kelsie Murchie, Ken Otter, Ken P. Findlay, Kenneth Dudley, Kevin Darras,

Kevin G. Borja-Acosta, Kieran Cox, Kinga Buda, Kris Harmon, Laela Sayigh, Laure Desutter-
Grandcolas, Laurel Symes, Lauren Kuehne, Laurence H. De Clippele, Leeann Henry, Leila
Hatch, Lenka Brousset, Leonardo Trevelin, Liana Chesini Rossi, Linilson Padovese, Lisa
Loseto, Louise Emmerson, Luca Iacobucci, Lucia Di Iorio, Luciana Nacimento , Luis P. da
Silva, Mao-Ning Tuanmuu, Marc Lammers, Marcia Maia, Marconi Campos, Marconi Campos
Cerqueira, Marcos Vaira, Marcus Rowcliffe, Marlize Muller, Martim Melo, María Jesús
García Bianco, Matias Carandell, Matthew J. Weldy, Matthew K Pine, Matthew Slater,
Mauricio Akmentins, Max Ritts, Michael Scherer-Lorenzen, Mike van der Schaar, Milton
Cezar Ribeiro, Mina Anders, Muhammad Justi Makmun Jusrin, Nadia Pieretti, Niaina Nirina
Mahefa Andriamavosoloarisoa, Nicholas Brown, Nicholas Friedman, Nick Gardner, Nicolas
Farrugia, Nicolas Pajusco, Nikos Fakotakis, Noam Leader, Nuno Faria, Ocean Acoustics
Group, Ocean Networks Canada Society, Olaf Boebel, Omar Hiram Martinez Castillo, Pedro
B. Lopes, Peter Ellis, Peter Taylor, Pooja Choksi, Pousada Xaraés, Pshemek Zdroik,
Rainforest Connection (RFCx), Refúgio Ecológico Caiman, Refúgio da Ilha Ecolodge, Rex K.
Andrew, Richard Kinsey, Rob Rempel, Robert John Young, Robert McCauley, Robert
Spaan, Robert Young, Rouvah Andriafanomezantsoa, Saki Harii, Samuel Challéat, Samuel
Haché, Sandra Mueller, Sara Gonçalves, Sarab Sethi, Sarah Rowley, Sarika Khanwilkar,
Sean Martin, Shannon MacPhee, Shinsuke Kawagucci, Sierra Hunicutt, Sierra Jarriel, Sigal
Balshine, Silent cities consortium, Simon Childerhouse, Sina Weier, Sofie Van Parijs, Stan
Dosso, Stanislas Wroza, Stephanie Plön, Stephen Insley, Stuart Marsden, Sérgio Moreira,
Tatiana Atemasova, Tetsuya Miwa, The Hunters and Trappers Committees of Inuvik, The
Inuvialuit-Canada Fisheries Joint Management Committee, Thiago M Ventura, Thomas
Sattler, Todor Ganchev, Tomas Altamirano, Tomasz Stanislaw Osiejuk, Tomaz Melo,
Tomonari Akamatsu, Tristan Louth-Robins, Tómas Grétar Gunnarsson, Valentin Chevalier,
Valerie Linden, Valéria Tavares, Vijay Ramesh, Vu Dinh Thong, Vy Tram Nguyen, Wen-Ling
Tsai, Will Duguid, Yenifer Herrera-Varón, You-Fang Chen, Yuhang Song, Zehava Sigal,
Zuania Colón-Piñeiro, Zuzana Burivalova, and Tuktoyaktuk.

The use of trade or firm names in this publication is for reader information and does not
imply endorsement by the U.S. Government of any product or service. The findings and
conclusions in this publication are those of the authors and should not be construed to
represent any official U.S. Government determination or policy. We acknowledge the NFDI
Consortium Earth System Sciences — NFDI4Earth, coordinated by TU Dresden, funded by
the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — project
number: 460036893. BIOMON is funded by the European Union's Horizon Europe
programme under grant agreement 101090273. Larissa Sayuri M. Sugai and Liiana Piatti
acknowledge grant Fundect T.O.:95/2023; SIAFEM: 33112. This paper is NOAA-PMEL
contribution number 5948. Songhai Li acknowledges the National Natural Science
Foundation of China (Grant numbers 42225604). Anna Franziska Cord was supported by
the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under
Germany's Excellence Strategy – EXC 2070 – 390732324. This research was also
funded by a Westlake University Startup Fund (Thomas C Wanger).

Author contributions

Kevin FA Darras, Rodney Rountree did the conceptualisation. Kevin FA Darras, Steven Van
Wilgenburg, Frederik Pitz, Youfang Chen, Lijun Dong, Amandine Gasc, Tzu-Hao Lin, Patrick
Mauritz Diaz, Shih-Hung Wu, Marcus Salton, Sarah Marley, Laura Schillé, Paul Jacobus

Wensveen, Camille Desjonquères, Orlando Acevedo-Charry, Matyáš Adam, Michel André,
Alexandre Antonelli, Leandro Aparecido Do Nascimento, Giulliana Appel, Christos Astaras,
Andrey Atemasov, Luc Barbaro, Fritjof Basan, Carly Batist, Adriá López Baucells, Julio
Ernesto Baumgarten, Just T Bayle Sempere, Kristen Bellisario, Asaf Ben David, Oded
Berger-Tal, Matthew G Betts, Iqbal Bhalla, Thiago Bicudo, Marta Bolgan, Sara Bombaci,
Martin Boullhesen, Tom Bradfer-Lawrence, Robert A Briers, Michal Budka, Katie Burchard,
Alice Calvente, Maite Cerezo-Araujo, Gunnar Cerwén, Maria Chistopolova, Christopher W
Clark, Benjamin Cretois, Chapin Czarnecki, Luis P da Silva, Wigna da Silva, Laurence H De
Clippele, David de la Haye, Ana Silvia de Oliveira Tissiani, Devin de Zwaan, Ricardo Díaz-
Delgado, Pedro Diniz, Dorgival Diógenes Oliveira-Júnior, Thiago Dorigo, Saskia Dröge,
Marina Duarte, Adam Duarte, Kerry Dunleavy, Robert Dziak, Simon Elise, Hiroto Enari,
Haruka S Enari, Florence Erbs, Nina Ferrari, Luane Ferreira, Abram B Fleishman, Bárbara
Freitas, Nick Friedman, Jérémy SP Froidevaux, Svetlana Gogoleva, Maria Isabel Gonçalves,
Carolina Gonzaga, José Miguel González Correa, Eben Goodale, Benjamin Gottesman,
Ingo Grass, Jack Greenhalgh, Jocelyn Gregoire, Jonas Hagge, William Halliday, Antonia
Hammer, Tara Hanf-Dressler, Samara Haver, Daniel Hending, Jose Hernandez-Blanco,
Thomas Hiller, Joe Chun-Chia Huang, Katie Lois Hutchinson, Janet Jackson, Alain Jacot,
Olaf Jahn, Jasper Kanes, Ellen Kenchington, Sebastian Kepfer-Rojas, Tharaka Kusuminda,
Yael Lehnardt, Paula Leitman, José Leon, Cicero Simão Lima-Santos, Kyle John Lloyd,
Audrey Looby, David López-Bosch, Tatiana Maeda, Christos Mammides, Gabriel Marcacci,
Matthias Markolf, Marinez Isaac Marques, Charles W Martin, Dominic A Martin, Kathy
Martin, Logan JT McLeod, Oliver Metcalf, Christoph Meyer, Grzegorz Mikusinski, Larissa
Sayuri Moreira Sugai, Dave Morris, Sandra Müller, Kelsie A Murchy, Maria Mas Navarro,
Rym Nouioua, Julian D Olden, Steffen Oppel, Anna N Osiecka, Miles Parsons, Filipa Isabel
Pereira Samarra, Cristian Pérez-Granados, Liliana Piatti, Mauro Pichorim, Thiago Pinheiro,
Jean-Nicolas Pradervand, Craig Radford, Emiliano Ramalho, Laura K Reynolds, Klaus
Riede, Talen Rimmer, Ricardo Rocha, Luciana Rocha, Samuel RP-J Ross, Carlos Salustio-
Gomes, Philip Samartzis, José Santos, Kevin Scharffenberg, Renée P Schoeman, Karl-
Ludwig Schuchmann, Esther Sebastián-González, Fannie Shabangu, Taylor Shaw, Xiaoli
Shen, David Singer, Jenni Stanley, Karolin Thomisch, Laura Torrent, Juan Traba, Junior A
Tremblay, Sunny Tseng, Mao-Ning Tuanmu, Marisol Valverde, Ben Vernasco, Raiane Vital
da Paz, Matthew Ward, Matthew Weldy, Julia Wiel, Jacob Willie, Heather Wood, Renata
Sousa-Lima contributed to data curation. Kevin FA Darras did the formal analysis. Kevin FA
Darras, Anna F Cord, Anna F Cord, Thomas Cherico Wanger acquired funding. Kevin FA
Darras, Rodney Rountree, Steven Van Wilgenburg, Frederik Pitz, Youfang Chen, Lijun
Dong, Amandine Gasc, Songhai Li conducted the investigation. Kevin FA Darras designed
the methodology. Kevin FA Darras, Rodney Rountree, Steven Van Wilgenburg, Lijun Dong,
Amandine Gasc, Jacopo Aguzzi, Robert Dziak, Jose Hernandez-Blanco, Ellen Kenchington,
Justin Kitzes, Matthew McKown, Carolyn Rosten, Songhai Li, Renata Sousa-Lima
administered the project or sub-projects (i.e., collections). Kevin FA Darras provided
resources. Kevin FA Darras provided software. Kevin FA Darras, Anna F Cord, Anna F
Cord, Songhai Li, Renata Sousa-Lima, Thomas Cherico Wanger conducted supervision.
Kevin FA Darras, Frederik Pitz, Youfang Chen, Tzu-Hao Lin, Patrick Mauritz Diaz, Shih-
Hung Wu, Marcus Salton, Laura Schillé carried out validation. Kevin FA Darras designed
visualisations. Kevin FA Darras, Anna F Cord, Tom Bradfer-Lawrence, Miles Parsons,
Samuel RP-J Ross, Anna F Cord wrote the original draft and Kevin FA Darras, Rodney
Rountree, Steven Van Wilgenburg, Anna F Cord, Amandine Gasc, Tzu-Hao Lin, Patrick
Mauritz Diaz, Shih-Hung Wu, Sarah Marley, Laura Schillé, Paul Jacobus Wensveen, Camille

Desjonquères, Orlando Acevedo-Charry, Matyáš Adam, Jacopo Aguzzi, Michel André, Alexandre Antonelli, Leandro Aparecido Do Nascimento, Giulliana Appel, Christos Astaras, Andrey Atemasov, Luc Barbaro, Fritjof Basan, Carly Batist, Adriá López Baucells, Julio Ernesto Baumgarten, Just T Bayle Sempere, Kristen Bellisario, Asaf Ben David, Oded Berger-Tal, Matthew G Betts, Iqbal Bhalla, Thiago Bicudo, Marta Bolgan, Sara Bombaci, Martin Boullhesen, Tom Bradfer-Lawrence, Robert A Briers, Michal Budka, Katie Burchard, Alice Calvente, Maite Cerezo-Araujo, Gunnar Cerwén, Maria Chistopolova, Christopher W Clark, Benjamin Cretois, Chapin Czarnecki, Luis P da Silva, Wigna da Silva, Laurence H De Clippele, David de la Haye, Ana Silvia de Oliveira Tissiani, Devin de Zwaan, Ricardo Díaz-Delgado, Pedro Diniz, Dorgival Diógenes Oliveira-Júnior, Thiago Dorigo, Saskia Dröge, Marina Duarte, Adam Duarte, Kerry Dunleavy, Robert Dziak, Simon Elise, Hiroto Enari, Haruka S Enari, Florence Erbs, Nina Ferrari, Luane Ferreira, Abram B Fleishman, Bárbara Freitas, Nick Friedman, Jérémy SP Froidevaux, Svetlana Gogoleva, Maria Isabel Gonçalves, Carolina Gonzaga, José Miguel González Correa, Eben Goodale, Benjamin Gottesman, Ingo Grass, Jack Greenhalgh, Jocelyn Gregoire, Jonas Hagge, William Halliday, Antonia Hammer, Tara Hanf-Dressler, Samara Haver, Daniel Hending, Jose Hernandez-Blanco, Thomas Hiller, Joe Chun-Chia Huang, Katie Lois Hutchinson, Janet Jackson, Alain Jacot, Olaf Jahn, Jasper Kanes, Ellen Kenchington, Sebastian Kepfer-Rojas, Justin Kitzes, Tharaka Kusuminda, Yael Lehnardt, Paula Leitman, José Leon, Cicero Simão Lima-Santos, Kyle John Lloyd, Audrey Looby, David López-Bosch, Tatiana Maeda, Christos Mammides, Gabriel Marcacci, Matthias Markolf, Marinez Isaac Marques, Charles W Martin, Dominic A Martin, Kathy Martin, Matthew McKown, Logan JT McLeod, Oliver Metcalf, Christoph Meyer, Grzegorz Mikusinski, Larissa Sayuri Moreira Sugai, Dave Morris, Sandra Müller, Kelsie A Murchy, Maria Mas Navarro, Rym Nouioua, Julian D Olden, Steffen Oppel, Anna N Osiecka, Miles Parsons, Filipa Isabel Pereira Samarra, Cristian Pérez-Granados, Liliana Piatti, Mauro Pichorim, Thiago Pinheiro, Jean-Nicolas Pradervand, Craig Radford, Emiliano Ramalho, Laura K Reynolds, Klaus Riede, Talen Rimmer, Ricardo Rocha, Luciana Rocha, Samuel RP-J Ross, Carolyn Rosten, Carlos Salustio-Gomes, Philip Samartzis, José Santos, Kevin Scharffenberg, Renée P Schoeman, Karl-Ludwig Schuchmann, Esther Sebastián-González, Fannie Shabangu, Taylor Shaw, Xiaoli Shen, David Singer, Jenni Stanley, Karolin Thomisch, Laura Torrent, Juan Traba, Junior A Tremblay, Sunny Tseng, Mao-Ning Tuanmu, Marisol Valverde, Ben Vernasco, Raiane Vital da Paz, Matthew Ward, Matthew Weldy, Julia Wiel, Jacob Willie, Heather Wood, Anna F Cord, Renata Sousa-Lima edited it.

Competing interests

The authors declare to have no competing interests. All authors have seen and approved the manuscript; it hasn't been accepted or published elsewhere.

References

References

1. Keith, D. A. *et al.* A function-based typology for Earth's ecosystems. *Nature* 1–6 (2022)

doi:10.1038/s41586-022-05318-4.

2. Doohan, B., Fuller, S., Parsons, S. & Peterson, E. E. The sound of management:

Acoustic monitoring for agricultural industries. *Ecol. Indic.* **96**, 739–746 (2019).

3. Duarte, C. M. *et al.* The soundscape of the Anthropocene ocean. *Science* **371**, (2021).

4. Pensieri, S. *et al.* Underwater Acoustic Measurements to Estimate Wind and Rainfall in

the Mediterranean Sea. *Adv. Meteorol.* **2015**, e612512 (2015).

5. Rountree, R. A., Juanes, F. & Bolgan, M. Temperate freshwater soundscapes: A

cacophony of undescribed biological sounds now threatened by anthropogenic noise.

PLOS ONE **15**, e0221842 (2020).

6. Gibb, R., Browning, E., Glover-Kapfer, P. & Jones, K. E. Emerging opportunities and

challenges for passive acoustics in ecological assessment and monitoring. *Methods*

Ecol. Evol. **10**, 169–185 (2018).

7. Pijanowski, B. C., Farina, A., Gage, S. H., Dumyahn, S. L. & Krause, B. L. What is

soundscape ecology? An introduction and overview of an emerging new science.

Landsc. Ecol. **26**, 1213–1232 (2011).

8. Sueur, J. & Farina, A. Ecoacoustics: the Ecological Investigation and Interpretation of

Environmental Sound. *Biosemiotics* **8**, 493–502 (2015).

9. Kang, J. *et al.* Supportive soundscapes are crucial for sustainable environments. *Sci.*

Total Environ. **855**, 158868 (2023).

10. Sueur, J., Krause, B. & Farina, A. Climate Change Is Breaking Earth's Beat. *Trends*

Ecol. Evol. **34**, 971–973 (2019).

11. Williams, R. *et al.* Noise from deep-sea mining may span vast ocean areas. *Science*

377, 157–158 (2022).

12. Müller, J. *et al.* Soundscapes and deep learning enable tracking biodiversity recovery in

tropical forests. *Nat. Commun.* **14**, 6191 (2023).

13.   Ross, S. R. P.-J. *et al.* Divergent ecological responses to typhoon disturbance revealed

via landscape-scale acoustic monitoring. *Glob. Change Biol.* **30**, e17067 (2024).

14.   Sethi, S. S. *et al.* Characterizing soundscapes across diverse ecosystems using a

universal acoustic feature set. *Proc. Natl. Acad. Sci.* (2020)

doi:10.1073/pnas.2004702117.

15.   Alvarsson, J. J., Wiens, S. & Nilsson, M. E. Stress Recovery during Exposure to Nature

Sound and Environmental Noise. *Int. J. Environ. Res. Public. Health* **7**, 1036–1046

(2010).

16.   Chen, Z., Hermes, J., Liu, J. & von Haaren, C. How to integrate the soundscape

resource into landscape planning? A perspective from ecosystem services. *Ecol. Indic.*

141, 109156 (2022).

17.   Astaras, C. *et al.* Boots on the ground: the role of passive acoustic monitoring in

evaluating anti-poaching patrols. *Environ. Conserv.* **47**, 213–216 (2020).

18.   Haver, S. M. *et al.* Comparing the Underwater Soundscapes of Four U.S. National

Parks and Marine Sanctuaries. *Front. Mar. Sci.* **6**, 500 (2019).

19.   Greenhalgh, J. A., Genner, M. J., Jones, G. & Desjonquères, C. The role of freshwater

bioacoustics in ecological research. *WIREs Water* **7**, e1416 (2020).

20.   Havlik, M.-N., Predragovic, M. & Duarte, C. M. State of Play in Marine Soundscape

Assessments. *Front. Mar. Sci.* **9**, (2022).

21.   Sugai, L. S. M., Silva, T. S. F., Ribeiro, J. W. & Llusia, D. Terrestrial Passive Acoustic

Monitoring: Review and Perspectives. *BioScience* **69**, 15–25 (2019).

22.   Scarpelli, M. D. A., Ribeiro, M. C., Teixeira, F. Z., Young, R. J. & Teixeira, C. P. Gaps in

terrestrial soundscape research: It's time to focus on tropical wildlife. *Sci. Total Environ.*

707, 135403 (2020).

23.   Lindseth, A. V. & Lobel, P. S. Underwater Soundscape Monitoring and Fish

Bioacoustics: A Review. *Fishes* **3**, 36 (2018).

24.   Boyd, I. *et al.* An International Quiet Ocean Experiment. *Oceanography* **24**, 174–181

(2015).

25.　Wang, L. S. *et al. Review of Underwater Acoustic Propagation Models.*

https://eprintspublications.npl.co.uk/6340/ (2014).

26.　Darras, K. F. A., Pütz, P., Fahrurrozi, Rembold, K. & Tscharntke, T. Measuring sound

detection spaces for acoustic animal sampling and monitoring. *Biol. Conserv.* **201**, 29–

37 (2016).

27.　Haupert, S., Sèbe, F. & Sueur, J. Physics-based model to predict the acoustic

detection distance of terrestrial autonomous recording units over the diel cycle and

across seasons: Insights from an Alpine and a Neotropical forest. *Methods Ecol. Evol.*

n/a, (2022).

28.　Nieto-Mora, D. A., Rodríguez-Buritica, S., Rodríguez-Marín, P., Martínez-Vargaz, J. D.

& Isaza-Narváez, C. Systematic review of machine learning methods applied to

ecoacoustics and soundscape monitoring. *Heliyon* **9**, e20275 (2023).

29.　Looby, A. *et al.* Global inventory of species categorized by known underwater sonifery.

Sci. Data **10**, 1–10 (2023).

30.　Parsons, M. J. G. *et al.* A Global Library of Underwater Biological Sounds (GLUBS): An

Online Platform with Multiple Passive Acoustic Monitoring Applications. in *The Effects*

of Noise on Aquatic Life: Principles and Practical Considerations (eds. Popper, A. N.,

Sisneros, J., Hawkins, A. D. & Thomsen, F.) 1–25 (Springer International Publishing,

Cham, 2023). doi:10.1007/978-3-031-10417-6_123-1.

31.　Ross, S. R. P.-J. *et al.* Passive acoustic monitoring provides a fresh perspective on

fundamental ecological questions. *Funct. Ecol.* **37**, 959–975 (2023).

32.　Gottesman, B. L. *et al.* What does resilience sound like? Coral reef and dry forest

acoustic communities respond differently to Hurricane Maria. *Ecol. Indic.* **126**, 107635

(2021).

33.　de Sousa, I. P. *et al.* Scale-free distribution of silences. *Phys. Rev. E* **105**, 014107

(2022).

34.　Kays, R. *et al.* The Movebank system for studying global animal movement and

demography. *Methods Ecol. Evol.* **13**, 419–431 (2022).

35.   Kays, R. *et al.* SNAPSHOT USA 2020: A second coordinated national camera trap

1183   survey of the United States during the COVID-19 pandemic. *Ecology* **103**, e3775

1184   (2022).

1185   36.   Culina, A. *et al.* Connecting the data landscape of long-term ecological studies: The

1186   SPI-Birds data hub. *J. Anim. Ecol.* **90**, 2147–2160 (2021).

1187   37.   Spatz, D. R. *et al.* The global contribution of invasive vertebrate eradication as a key

1188   island restoration tool. *Sci. Rep.* **12**, 13391 (2022).

1189   38.   Darras, K. F. A. *et al.* Worldwide Soundscapes project meta-data. Zenodo

1190   https://doi.org/10.5281/zenodo.10598949 (2024).

1191   39.   Darras, K. F. Worldwide Soundscapes project. *Worldwide Soundscapes project on*

1192   *ecoSound-web* https://ecosound-web.de/ecosound_web/collection/index/106 (2024).

1193   40.   Darras, K. F. A. *et al.* ecoSound-web: an open-source, online platform for ecoacoustics.

1194   *F1000Research* vol. 9 1224 Preprint at https://doi.org/10.12688/f1000research.26369.2

1195   (2023).

1196   41.   Lin, T.-H., Sinniger, F., Harii, S. & Akamatsu, T. Using Soundscapes to Assess

1197   Changes in Coral Reef Social-Ecological Systems. *Oceanography* (2023)

1198   doi:10.5670/oceanog.2023.s1.7.

1199   42.   Chen, C., Lin, T.-H., Watanabe, H. K., Akamatsu, T. & Kawagucci, S. Baseline

1200   soundscapes of deep-sea habitats reveal heterogeneity among ecosystems and

1201   sensitivity to anthropogenic impacts. *Limnol. Oceanogr.* **66**, 3714–3727 (2021).

1202   43.   Roe, P. *et al.* The Australian Acoustic Observatory. *Methods Ecol. Evol.* **12**, 1802–1808

1203   (2021).

1204   44.   Xeno-canto Foundation. *Xeno-Canto: Sharing Bird Sounds from around the World*.

1205   (Xeno-canto Foundation Amsterdam, 2012).

1206   45.   IQOE. Acoustic Capabilities of Existing Observing Systems.

1207   https://www.iqoe.org/systems.

1208   46.   Lamont, T. A. C. *et al.* HydroMoth: Testing a prototype low-cost acoustic recorder for

1209   aquatic environments. *Remote Sens. Ecol. Conserv.* **8**, 362–378 (2022).

47. Beck, J., Böller, M., Erhardt, A. & Schwanghart, W. Spatial bias in the GBIF database

and its effect on modeling species' geographic distributions. *Ecol. Inform.* **19**, 10–15

(2014).

48. Rountree, R. *et al.* Towards an Optimal Design for Ecosystem-Level Ocean

Observatories. in 79–106 (2020). doi:10.1201/9780429351495-2.

49. Siddagangaiah, S., Chen, C.-F., Hu, W.-C. & Farina, A. The dynamical complexity of

seasonal soundscapes is governed by fish chorusing. *Commun. Earth Environ.* **3**, 1–11

(2022).

50. Hernández-León, S. *et al.* Zooplankton abundance in subtropical waters: is there a

lunar cycle? *Sci. Mar.* **65**, 59–64 (2001).

51. Simonis, A. E. *et al.* Lunar cycles affect common dolphin Delphinus delphis foraging in

the Southern California Bight. *Mar. Ecol. Prog. Ser.* **577**, 221–235 (2017).

52. Mougeot, F. & Bretagnolle, V. Predation risk and moonlight avoidance in nocturnal

seabirds. *J. Avian Biol.* **31**, 376–386 (2000).

53. Krause, B., Gage, S. H. & Joo, W. Measuring and interpreting the temporal variability in

the soundscape at four places in Sequoia National Park. *Landsc. Ecol.* **26**, 1247

(2011).

54. Grinfeder, E. *et al.* Soundscape dynamics of a cold protected forest: dominance of

aircraft noise. *Landsc. Ecol.* **37**, 567–582 (2022).

55. Sugai, L. S. M., Desjonquères, C., Silva, T. S. F. & Llusia, D. A roadmap for survey

designs in terrestrial acoustic monitoring. *Remote Sens. Ecol. Conserv.* **6**, 220–235

(2020).

56. Kronfeld-Schor, N. *et al.* Chronobiology by moonlight. *Proc. R. Soc. B Biol. Sci.* **280**,

20123088 (2013).

57. Wilgenburg, S. L. V. *et al.* Evaluating trade-offs in spatial versus temporal replication

when estimating avian community composition and predicting species distributions.

Avian Conserv. Ecol. **19**, (2024).

58. Desjonquères, C. *et al.* Acoustic species distribution models (aSDMs): A framework to

forecast shifts in calling behaviour under climate change. *Methods Ecol. Evol.* **13**,

2275–2288 (2022).

59. Lillis, I. V. & Boebel, O. Marine soundscape planning: Seeking acoustic niches for

anthropogenic sound. *J. Ecoacoustics* **2**, 1–1 (2018).

60. Luypaert, T. *et al.* A framework for quantifying soundscape diversity using Hill numbers.

Methods Ecol. Evol. **13**, 2262–2274 (2022).

61. Cummings, W. C. & Thompson, P. O. Underwater Sounds from the Blue Whale,

Balaenoptera musculus. *J. Acoust. Soc. Am.* **50**, 1193–1198 (1971).

62. Moir, H. M., Jackson, J. C. & Windmill, J. F. C. Extremely high frequency sensitivity in a

'simple' ear. *Biol. Lett.* **9**, 20130241 (2013).

63. Sousa-Lima, R. A Review and Inventory of Fixed Autonomous Recorders for Passive

Acoustic Monitoring of Marine Mammals. *Aquat. Mamm.* **39**, 23–53 (2013).

64. Höchst, J. *et al.* Bird@Edge: Bird Species Recognition at the Edge. in *Networked*

Systems: 10th International Conference, NETYS 2022, Virtual Event, May 17–19,

2022, Proceedings 69–86 (Springer-Verlag, Berlin, Heidelberg, 2022). doi:10.1007/978-

3-031-17436-0_6.

65. Zagmajster, M., Malard, F., Eme, D. & Culver, D. C. Subterranean Biodiversity Patterns

from Global to Regional Scales. in *Cave Ecology* (eds. Moldovan, O. T., Kováč, Ľ. &

Halse, S.) 195–227 (Springer International Publishing, Cham, 2018). doi:10.1007/978-

3-319-98852-8_9.

66. Messager, M. L. *et al.* Global prevalence of non-perennial rivers and streams. *Nature*

594, 391–397 (2021).

67. Murchy, K.A., Rountree, R.A., & Juanes, F. The role of the soundscape in the

behavioral ecology of stream-dwelling salmonids. in.

68. Schoeman, R. P., Erbe, C. & Plön, S. Underwater Chatter for the Win: A First

Assessment of Underwater Soundscapes in Two Bays along the Eastern Cape Coast

of South Africa. *J. Mar. Sci. Eng.* **10**, 746 (2022).

69.   Hill, A. P., Prince, P., Snaddon, J. L., Doncaster, C. P. & Rogers, A. AudioMoth: A low-

cost acoustic device for monitoring biodiversity and the environment. *HardwareX* **6**,

e00073 (2019).

70.   Linke, S. *et al.* Freshwater ecoacoustics as a tool for continuous ecosystem monitoring.

Front. Ecol. Environ. **16**, 231–238 (2018).

71.   Metcalf, O. C. *et al.* Listening to tropical forest soils. *Ecol. Indic.* **158**, 111566 (2024).

72.   Maeder, M., Guo, X., Neff, F., Mathis, D. S. & Gossner, M. M. Temporal and spatial

dynamics in soil acoustics and their relation to soil animal diversity. *PLOS ONE* **17**,

e0263618 (2022).

73.   Hughes, A. C. *et al.* Sampling biases shape our view of the natural world. *Ecography*

44, 1259–1269 (2021).

74.   Amano, T., González-Varo, J. P. & Sutherland, W. J. Languages Are Still a Major

Barrier to Global Science. *PLOS Biol.* **14**, e2000933 (2016).

75.   Reboredo Segovia, A. L., Romano, D. & Armsworth, P. R. Who studies where?

Boosting tropical conservation research where it is most needed. *Front. Ecol. Environ.*

18, 159–166 (2020).

76.   Newson, S. E., Evans, H. E. & Gillings, S. A novel citizen science approach for large-

scale standardised monitoring of bat activity and distribution, evaluated in eastern

England. *Biol. Conserv.* **191**, 38–49 (2015).

77.   Wall, C. C. *et al.* The Next Wave of Passive Acoustic Data Management: How

Centralized Access Can Enhance Science. *Front. Mar. Sci.* **8**, (2021).

78.   Roch, M. A. *et al.* Management of acoustic metadata for bioacoustics. *Ecol. Inform.* **31**,

122–136 (2016).

79.   Sugai, L. S. M. & Llusia, D. Bioacoustic time capsules: Using acoustic monitoring to

document biodiversity. *Ecol. Indic.* **99**, 149–152 (2019).

80.   Southall, B. L. *et al.* MARINE MAMMAL NOISE-EXPOSURE CRITERIA: INITIAL

SCIENTIFIC RECOMMENDATIONS. *Bioacoustics* **17**, 273–275 (2008).

81.   Haver, S. M. *et al.* Monitoring long-term soundscape trends in U.S. Waters: The

NOAA/NPS Ocean Noise Reference Station Network. *Mar. Policy* **90**, 6–13 (2018).

82.   Garibbo, S. *et al.* Low-frequency ocean acoustics – measurements from the Lofoten-

Vesterålen Ocean Observatory, Norway. *Proc. Meet. Acoust.* **40**, 040001 (2020).

83.   Cretois, B., Rosten, C. M. & Sethi, S. S. Voice activity detection in eco-acoustic data

enables privacy protection and is a proxy for human disturbance. *Methods Ecol. Evol.*

13, 2865–2874 (2022).

84.   Görföl, T. *et al.* ChiroVox: a public library of bat calls. *PeerJ* **10**, e12445 (2022).

85.   Parsons, M. J. G. *et al.* Sounding the Call for a Global Library of Underwater Biological

Sounds. *Front. Ecol. Evol.* **10**, (2022).

86.   Looby, A. *et al.* FishSounds version 1.0: A website for the compilation of fish sound

production information and recordings. *Ecol. Inform.* 101953 (2022)

doi:10.1016/j.ecoinf.2022.101953.

87.   Pijanowski, B. C. *et al.* Soundscape Ecology: The Science of Sound in the Landscape.

BioScience **61**, 203–216 (2011).

88.   Hao, Z. *et al.* Can urban forests provide acoustic refuges for birds? Investigating the

influence of vegetation structure and anthropogenic noise on bird sound diversity. *J.*

For. Res. **35**, 33 (2024).

89.   Briefer, E. F. *et al.* The power of sound: unravelling how acoustic communication

shapes group dynamics. *Philos. Trans. R. Soc. B Biol. Sci.* **379**, 20230182 (2024).

90.   Gasc, A. *et al.* Assessing biodiversity with sound: Do acoustic diversity indices reflect

phylogenetic and functional diversities of bird communities? *Ecol. Indic.* **25**, 279–287

(2013).

91.   Buxton, R. T., Pearson, A. L., Allou, C., Fristrup, K. & Wittemyer, G. A synthesis of

health benefits of natural sounds and their distribution in national parks. *Proc. Natl.*

Acad. Sci. **118**, (2021).

92.   Hart, P. J. *et al.* Timing Is Everything: Acoustic Niche Partitioning in Two Tropical Wet

Forest Bird Communities. *Front. Ecol. Evol.* **9**, (2021).

93. EY, E. & FISCHER, J. The "Acoustic Adaptation Hypothesis"—a Review of the

Evidence from Birds, Anurans and Mammals. *Bioacoustics* **19**, 21–48 (2009).

94. Pilotto, F. *et al.* Meta-analysis of multidecadal biodiversity trends in Europe. *Nat.*

Commun. **11**, 3486 (2020).

95. Global Biodiversity Information Facility (2024). Free and Open Access to Biodiversity

Data. What is GBIF? https://www.gbif.org/what-is-gbif (2024).

96. Gonzalez, A. *et al.* A global biodiversity observing system to unite monitoring and guide

action. *Nat. Ecol. Evol.* 1–5 (2023) doi:10.1038/s41559-023-02171-0.

97. Towards a Transnational Acoustic Biodiversity Monitoring Network (TABMON).

https://www.nina.no/english/TABMON.

98. Leiper, A. What does the consultancy industry need from academia? A soundscape

and planning perspective. in *ACOUSTICS 2020* (Institute of Acoustics, Virtual, 2020).

doi:10.25144/13347.

99. Batist, C. & Campos-Cerqueira, M. *Harnessing the Power of Sound and AI to Track*

Global Biodiversity Framework (GBF) Targets. (2023).

100. Darras, K. F. A., Pérez, N., -, M. & Hanf-Dressler, T. BioSounds: an open-source,

online platform for ecoacoustics. *F1000Research* **9**, 1224 (2020).

101. Frey, S., Fisher, J. T., Burton, A. C. & Volpe, J. P. Investigating animal activity patterns

and temporal niche partitioning using camera-trap data: challenges and opportunities.

Remote Sens. Ecol. Conserv. **3**, 123–132 (2017).

102. *Exploring Animal Behavior Through Sound: Volume 1: Methods*. (Springer International

Publishing, Cham, 2022). doi:10.1007/978-3-030-97540-1.

103. Keith, D. A. *et al.* Indicative distribution maps for Ecosystem Functional Groups - Level

3 of IUCN Global Ecosystem Typology. Zenodo

https://doi.org/10.5281/zenodo.4018314 (2020).

104. Keith, D. A., Ferrer-Paris, J. R., Nicholson, E. & Kingsford, R. T. *The IUCN Global*

Ecosystem Typology 2.0: Descriptive Profiles for Biomes and Ecosystem Functional

Groups. (IUCN, Gland, Switzerland, 2020).

105. GEBCO. Gridded bathymetry data (General Bathymetric Chart of the Oceans). *GEBCO*

https://www.gebco.net/data_and_products/gridded_bathymetry_data/.

A - Framework

Space

Spatial replicates

Temporal replicates

Time

B - Replication (dataset level)

Terrestrial

Subterranean

Marine

Freshwater

temporal   spatial   spatio temporal

C - Spatial extent and coverage of sampling sites

Terrestrial sites

200   400   600

D - Temporal extent and coverage (day level)

Sampled days

2000   2010   2020

Core realm: Freshwater   Marine   Subterranean   Terrestrial

A - Topographical and latitudinal distribution of sampling sites

IUCN core realms
- Freshwater
- Marine
- Subterranean
- Terrestrial

B - Top 10 administrative regions for spatial sampling coverage

WDPA categories
- Ia
- Ib
- II
- >II

A - Sampling coverage within time cycles

Freshwater diel | Marine diel | Terrestrial diel | Subterranean diel

Time windows: Dawn, Day, Dusk, Night

Freshwater lunar | Marine lunar | Terrestrial lunar | Subterranean lunar

Bright, Dark

Freshwater seasonal | Marine seasonal | Terrestrial seasonal | Subterranean seasonal

Spring, Summer, Autumn, Winter

B - Top 10 administrative regions for temporal sampling coverage

WDPA categories: Ia, Ib, II, NA

A - Case study soundscapes

Large lowland rivers

Riverine estuaries

Photic coral reefs

Island slopes

Bathypelagic ocean waters

Tropical lowland rainforests

Tropical montane rainforests

Plantations

Urban ecosystems

Polar outcrops

Aerobic caves

soundscape occupancy

Soundscape component: ■ anthropophony ■ biophony ■ geophony

B - Illustrative analyses

Macroecology

Conservation biology

Phenology

Biophony

Latitude

Biophony

Anthropophony

Biophony

Diel time windows

P(>0) = 1

P(<0) = 1

morning   noon   evening   night

Realms: ● Freshwater ● Freshwater-Marine ● Marine ● Subterranean ● Terrestrial