Title

Worldwide Soundscapes: a synthesis of passive acoustic monitoring across realms

Authors & affiliations

Kevin FA Darras[1,2,3C], Rodney Rountree[4,5], Steven Van Wilgenburg[6], Anna F Cord[3,7], Youfang Chen[2], Lijun Dong[8], Agnès Rocquencourt[1], Camille Desjonquères[9,10], Patrick Mauritz Diaz[11], Tzu-Hao Lin[12], Amandine Gasc[13], Sarah Marley[14], Marcus Salton[15], Laura Schillé[16], Paul Jacobus Wensveen[17], Shih-Hung Wu[18], Orlando Acevedo-Charry[19,20], Matyáš Adam[21], Jacopo Aguzzi[22], Irmak Akoglu[23], M Clara P Amorim[24], Michel André[25], Alexandre Antonelli[26,27,28], Leandro Aparecido Do Nascimento[29], Giulliana Appel[30], Stephanie Archer[31], Christos Astaras[32], Andrey Atemasov[33,34], Jamieson Atkinson[35], Joël Attia[36], Emanuel Baltag[37], Luc Barbaro[38], Fritjof Basan[39], Carly Batist[40,41], Adriá López Baucells[42], Julio Ernesto Baumgarten[43], Just T Bayle Sempere[44], Kristen Bellisario[45], Asaf Ben David[46], Oded Berger-Tal[47], Matthew G Betts[48], Iqbal Bhalla[49], Thiago Bicudo[40], Marta Bolgan[50], Sara Bombaci[51], Martin Boullhesen[52], Tom Bradfer-Lawrence[53,54], Robert A Briers[55], Michal Budka[56], Katie Burchard[57], Giuseppa Buscaino[58], Alice Calvente[59], Maria Ceraulo[58], Maite Cerezo-Araujo[60], Gunnar Cerwén[61], Maria Chistopolova[62], Christopher W Clark[63], Kieran D Cox[64], Benjamin Cretois[65], Chapin Czarnecki[66], Luis P da Silva[67,68], Wigna da Silva[69], Laurence H De Clippele[70], David de la Haye[71], Ana Silvia de Oliveira Tissiani[72], Devin de Zwaan[73,74], M. Eugenia Degano[75,76], Joaquin del Rio[77], Christian Devenish[78], Ricardo Díaz-Delgado[79], Pedro Diniz[80], Dorgival Diógenes Oliveira-Júnior[81,82], Thiago Dorigo[83], Saskia Dröge[84], Marina Duarte[85,86], Adam Duarte[87], Kerry Dunleavy[88], Robert Dziak[89], Simon Elise[90,91], Hiroto Enari[92], Haruka S Enari[92], Florence Erbs[25], Britas Klemens Eriksson[93], Pınar Ertör-Akyazi[23], Nina Ferrari[48], Luane Ferreira[69], Abram B Fleishman[88], Paulo Fonseca[94], Bárbara Freitas[95,96,97], Nick Friedman[98], Jérémy SP Froidevaux[53,99], Svetlana Gogoleva[100], Maria Isabel Gonçalves[43], Carolina Gonzaga[85], José Miguel González Correa[44], Eben Goodale[101], Benjamin Gottesman[63], Ingo Grass[102], Jack Greenhalgh[103], Jocelyn Gregoire[104], Jonas Hagge[105], William Halliday[106], Antonia Hammer[107], Tara Hanf-Dressler[108], Sylvain Haupert[109], Samara Haver[110], Daniel Hending[111], Jose Hernandez-Blanco[62], Thomas Hiller[102], Joe Chun-Chia Huang[112], Katie Lois Hutchinson[113], Carole Hyacinthe[114], Janet Jackson[115], Alain Jacot[116], Olaf Jahn[117,118], Francis Juanes[4], Ellen Kenchington[119], Sebastian Kepfer-Rojas[120], Justin Kitzes[66], Tharaka Kusuminda[121], Yael Lehnardt[47], Jialin Lei[122], Paula Leitman[83], José Leon[123], Deng Li[122], Cicero Simão Lima-Santos[81,82], Kyle John Lloyd[124,125,126], Audrey Looby[127,128], David López-Bosch[42], Tatiana Maeda[129], Franck Malige[130], Christos Mammides[131], Gabriel Marcacci[116], Matthias Markolf[132,133], Marinez Isaac Marques[72], Charles W Martin[127], Dominic A Martin[134,135], Kathy Martin[73,136], Matthew McKown[88], Logan JT McLeod[137], Oliver Metcalf[138], Christoph Meyer[139], Grzegorz Mikusinski[140], João Monteiro[141], Larissa Sayuri Moreira Sugai[63], Dave Morris[115], Sandra Müller[142], Sebastian Eduardo Muñoz-Duque[143], Kelsie A Murchy[144], Ivan Nagelkerken[145], Maria Mas Navarro[42], Rym Nouioua[146], Carolina Ocampo-Ariza[147,148], Julian D Olden[149], Steffen Oppel[54,116], Anna N Osiecka[150], Elena Papale[58], Miles Parsons[106], Julie Patris[130], João Pedro Marques[24], Filipa Isabel Pereira Samarra[17], Cristian Pérez-Granados[151], Liliana Piatti[152], Mauro Pichorim[81,82], Thiago Pinheiro[69], Jean-Nicolas Pradervand[116], John Quinn[153], Bernardo Quintella[24], Craig Radford[154], Xavier Raick[155], Ana Rainho[94], Emiliano Ramalho[40], Sylvie Rétaux[156], Laura K Reynolds[157], Klaus Riede[117], Talen Rimmer[144], Noelia Rios[158], Ricardo Rocha[111], Luciana Rocha[69], Paul Roe[159], Samuel RP-J Ross[160], Carolyn M Rosten[161], Carlos Salustio-Gomes[81,82], Philip Samartzis[162], José Santos[69], Kevin Scharffenberg[163], Renée P Schoeman[164], Karl-Ludwig Schuchmann[72], Esther Sebastián-González[151], Sebastian Seibold[165], Sarab Sethi[166], Fannie Shabangu[167,168], Taylor Shaw[142], Xiaoli Shen[169], David Singer[105], Ana Sirovic[170], Brittnie Spriel[64], Jenni Stanley[171], Valeria da Cunha Tavares[30,172,173], Karolin Thomisch[174], Jianfeng Tong[175], Laura Torrent[42], Juan Traba[176,177], Junior A

51 Tremblay[178], Leonardo Trevelin[30], Sunny Tseng[179], Mao-Ning Tuanmu[180], Théophile Turco[36],
52 Marisol Valverde[181], Ben Vernasco[87,182], Manuel Vieira[24], Raiane Vital da Paz[81,82], Matthew
53 Ward[183], Maryann Watson[93], Matthew Weldy[48,184], Julia Wiel[65], Jacob Willie[185], Heather
54 Wood[186], Jinshan Xu[119], Wenyi Zhou[187,188,189], Songhai Li[8C], Renata Sousa-Lima[69C], Thomas
55 Cherico Wanger[2C]
56
57 [C]: corresponding authors
58
1: EFNO, ECODIV, INRAE, Domaine des Barres, Nogent-sur-Vernisson, Centre-Val de
Loire, 45290, France
2: Sustainable Agricultural Systems & Engineering Lab, School of Engineering, Westlake
University, No. 600 Dunyu Road, Sandun Town, Xihu District, Hangzhou, Zhejiang province,
China
3: Chair of Computational Landscape Ecology, Faculty of Environmental Sciences, Dresden
University of Technology, Helmholtzstr. 10, Dresden, 01069, Germany
4: Biology Department, University of Victoria, Cunningham Building, Room 202, PO Box
1700 STN CSC, Victoria, BC, V8W 2Y2, Canada
5: The Fish Listener, 23 Joshua Lane, Waguoit, MA, 2536, U.S.A.
6: Terrestrial Unit, Prairie Region, Canadian Wildlife Service, Environment & Climate
Change Canada, Prairie & Northern Wildlife Research Centre, 115 Perimeter Road,
Saskatoon, Saskatchewan, S7N 0X4, Canada
7: Agro-ecological Modeling Group, Faculty of Agriculture, University of Bonn, Niebuhrstr.
1a, Bonn, 53113, Germany
8: Marine Mammal and Marine Bioacoustics Laboratory, Department of Deep Sea Science,
Institute of Deep-sea Science and Engineering Chinese Academy of Sciences, 28 Luhuitou
Road, Sanya, Hainan, 572000, China
9: Institut de Systématique, Évolution, Biodiversité, Muséum National d'Histoire naturelle,
Rue Buffon, Paris, 75005, France
10: Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, Grenoble,
38000, France
11: Climate Change and Biodiversity, Hanns R. Neumann Stiftung Indonesia, Muaradua,
OKU Selatan, South Sumatera, 32212, Indonesia
12: Biodiversity Research Center, Academia Sinica, 128 Academia Road, Section 2,
Nankang, Taipei , 11529, Taiwan
13: Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix Marseille
Univ, Avignon Univ, CNRS, IRD, Campus Aix Technopôle de l'environnement Arbois
Méditerranée Avenue Louis Philibert Bât Villemin, Aix-en-Provence, 13545, France
14: Scotland's Rural College, Craibstone Estate, Aberdeen, AB21 9YA, United Kingdom
15: Australian Antarctic Division, Science Branch, Department of Climate Change, Energy,
Environment and Water, Channel Highway, Kingston, Tasmania, 7050, Australia
16: BIOGECO, INRAE, University of Bordeaux, 69 route d'Arcachon, Cestas, 33610, France
17: Westman Islands Research Centre, University of Iceland, Ægisgata 2, Vestmannaeyjar,
900, Iceland
18: Biodiversity Indicator Lab, Division of Habitats and Ecosystems, Endemic Species
Research Institute, 1 Minsheng East Road, Jiji Town, Nantou County, Taiwan, 552005,
Taiwan
19: Colección de Sonidos Ambientales Mauricio Álvarez-Rebolledo - Colecciones
Biológicas, Subdirección de Investigaciones, Instituto de Investigación de Recursos
Biológicos Alexander von Humboldt, Claustro de San Agustín, Villa de Leyva, Boyacá,
Colombia
20: Quantitative Ecology Lab & Ordway Lab of Ecosystem Conservation, School of Natural
Resources and Environment, Department of Wildlife Ecology and Conservation, & Florida
Museum of Natural History, University of Florida, 715 Newell Dr, Building 866 & 1659
Museum Rd, Dickinson Hall 315, Gainesville, Florida, 32611, U.S.A.

21: Department of Environmental Security, Faculty of Logistics and Crisis Management,
Tomas Bata University, Studentské nám. 1532, Uherské Hradiště, 68601, Czech Republic
22: Marine Science Institute, ICM, CSIC, ICM, CSIC, Paseig Barceloneta, Barcelona, Spain
23: Institute of Environmental Sciences, Boğaziçi University, Hisar Campus, Bebek, İstanbul,
34342, Türkiye
24: MARE – Marine and Environmental Sciences Centre / ARNET - Aquatic Research
Network, Departamento de Biologia Animal, Faculdade de Ciências da Universidade de
Lisboa, Campo Grande, Lisbon, 1749-016 , Portugal
25: Laboratory of Applied Bioacoustics, Polytechnic University of Catalonia, BarcelonaTech,
Rambla Exposicion, 24, Vilanova I la Geltrú, Barcelona, 8800, Spain
26: Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, United Kingdom, Kew,
Richmond, Surrey, TW9 3AE, United Kingdom
27: Gothenburg Global Biodiversity Centre, Department of Biological and Environmental
Sciences, University of Gothenburg, Box 461, Göteborg, SE 40530, Sweden
28: Department of Plant, University of Oxford, South Parks Road, Oxford, OX1 3RB, United
Kingdom
29: Bioacoustics Group, Science Department, Biometrio.earth, Dr. Schönemann Str. 38,
Saarbrücken, 66123, Germany
30: Biodiversidade e Serviços Ecossistêmicos, Instituto Tecnológico Vale, Boaventura da
Silva, 955, Belém, PA, 66055-090, Brazil
31: Stephanie Archer, Louisiana Universities Marine Consortium, 8124 Highway 56,
Chauvin, LA, USA, 70344, USA
32: Wildlife Lab, Forest Research Institute, Hellenic Agricultural Organization - DIMITRA
(ELGO-DIMITRA), Vasilika, Thessaloniki, TK 57006, Greece
33: Department of Zoology and Animal Ecology, School of Biology, V.N.Karazin Kharkiv
National University, 4 Svobody Square, Kharkiv, 61022, Ukraine
34: Research Department, National Park 'Homilshanski Lisy', 27 Monastyrska Street,
Koropove village, Kharkiv region, 63437, Ukraine
35: Aquatic Research and Restoration Centre, British Columbia Conservation Foundation,
105 1885 Boxwood Rd, British Columbia, Canada
36: ENES Bioacoustics Research Laboratory, Centre de Recherche en Neurosciences de
Lyon (CRNL), University of Saint-Etienne, CNRS UMR 5292, 23 rue du docteur Paul
Michelon, Saint-Etienne, France, France
37: Marine Biological Station "Prof. Dr. Ioan Borcea", Agigea, Alexandru Ioan Cuza
University of Iasi, Nicolae Titulescu Street, no. 168, Agigea,, Constanta, Romania
38: DYNAFOR, University of Toulouse, INRAE, Chemin de Borde-Rouge, Castanet-Tolosan,
F-31326, France
39: Federal Maritime and Hydrographic Agency (BSH), Bernhard-Nocht Straße 78,
Hamburg, 20359, Germany
40: Rainforest Connection, Science department, 440 Cobia Drive, Suite 1902, TX, 77494,
U.S.A.
41: Arbimon, U.S.A.
42: BiBio Research Group, Natural Sciences Museum of Granollers, Av. Francesc Macià 51,
Granollers, Catalonia, 08402, Spain
43: Applied Ecology & Conservation Lab, Biological Sciences, Universidade Estadual de
Santa Cruz, Rodovia Jorge Amado km 16, Salobrinho, Ilhéus, Bahia, 45662900, Brazil
44: Marine Biological Acoustic, Marine Sciences and Applied Biology , University of Alicante,
C/ San Vicente del Raspeig s/n, San Vicente del Raspeig, Alicante, 3690, Spain
45: HIFI Lab, John Martinson Honors College, Purdue University West Lafayette Indiana,
1101 Third Street, West Lafayette, Indiana , 47906, U.S.A.
46: Tamar Dayan's Lab, School of zoology, Steinhardt Museum of Natural History, Tel-Aviv
university, Klausner 12, Tel-Aviv, Tel-Aviv, 6997801, Israel
47: The Conservation Behavior Research Group, Mitrani Department of Desert Ecology,
Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev,
Midreshet Ben-Gurion, 8499000, Israel

48: Forest Landscape Ecology Lab, Department of Forest Ecosystems and Society, Oregon
State University, U.S.A.
49: School of Geography and the Environment, Oxford University, South Parks Rd, Oxford,
OX1 3QY, United Kingdom
50: Ocean Science Consulting Limited, Spott Road, Dunbar, EH42 1RR , United Kingdom
51: Equity in Ecological Research Lab, Department of Fish, Wildlife, and Conservation
Biology, Colorado State University, 1474 Campus Delivery, Fort Collins, Colorado, 80521,
U.S.A.
52: Instituto de Ecorregiones Andinas (INECOA-UNJu-CONICET), Facultad de Ingeniería,
Universidad Nacional de Jujuy, Gorriti 237, San Salvador de Jujuy, Jujuy, 4600, Argentina
53: Biological and Environmental Sciences, University of Stirling, Stirling, Stirlingshire, FK9
4LA, Scotland
54: Centre for Conservation Science, RSPB, 2 Lochside View, Edinburgh Park, Edinburgh,
EH12 9DH, United Kingdom
55: Centre for Conservation and Restoration Science, School of Applied Sciences,
Edinburgh Napier University, Sighthill Campus, Edinburgh, EH11 4BN, United Kingdom
56: Department of Behavioural Ecology, Faculty of Biology, Adam Mickiewicz University,
Uniwersytetu Poznańskiego 6, Poznań, 61614, Poland
57: Northeast Fisheries Science Center, Narragansett Laboratory, 28 Tarzwell Drive,
Narragansett, RI, 2882, U.S.A.
58: BioacousticsLab Capo Granitola, National Research Council, Via del Mare, 6 - Torretta
Granitola, Campobello di Mazara (TP), 91021, Italy
59: Laboratório de Botânica Sistemática, Departamento de Botânica, Ecologia e Zoologia,
Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio
Grande do Norte, 59078-900, Brazil
60: South Iceland Research Centre, University of Iceland, Lindarbraut 4, Laugarvatn, 840,
Iceland
61: Sensola SLU Multisensory Outdoor Laboratory, Department of Landscape Architecture,
Planning and Management, Swedish University of Agricultural Sciences, SLU, Sundsvägen
4-6, Alnarp, 23053, Sweden
62: Behavior and Behavioral Ecology of Mammals, Servertsov Institute of Ecology and
Evolution of the Russian Academy of Sciences, Leninskiy prospekt, 33, Moscow, 119071,
Russia
63: K. Lisa Yang Center for Conservation Bioacoustics , Cornell Lab of Ornithology , Cornell
University , 159 Sapsucker Woods Rd , New York , 14850, U.S.A.
64: The Marine Ecology Lab, Department of Biological Sciences, Simon Fraser University,
8888 University Drive, British Columbia, V5A 1S6, Canada
65: Miljødata, Norwegian Institute for Nature Research, Høgskoleringen 9, Trøndelag,
Trondheim, 7034, Norway
66: Department of Biological Sciences, University of Pittsburgh, Clapp Hall, Fifth and Ruskin
Avenues, Pittsburgh, PA, 15237, U.S.A.
67: CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO
Laboratório Associado, Campus de Vairão, Universidade do Porto-InBIO, Universidade do
Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal
68: BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, University of
Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal
69: EcoAcoustic Research Hub (EAR Hub), Biosciences Center, Universidade Federal do
Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio Grande do Norte, 59078-900,
Brazil
70: School of Biodiversity, One Health & Veterinary Medicine, University of Glasgow,
Graham Kerr Building, Hillhead street 82, Scotland, United Kingdom, G12 8QQ, United
Kingdom
71: School of Arts and Cultures, Newcastle University, Newcastle upon Tyne, NE1 7RU,
United Kingdom

72: INAU Pantanal Biodata Center, National Institute of Science and Technology in Wetlands (INAU), Computational Bioacoustics Research Unit (CO.BRA), Federal University of Mato Grosso (UFMT), Cuiabá, 2367 Fernando Correa da Costa Ave, Mato Grosso, 78060-900, Brazil

73: Forest and Conservation Sciences, University of British Columbia, 2424 Main Mall, Vancouver, British Columbia, V6T 1Z4, Canada

74: Department of Biology, Mount Allison University, 63B York Street, Sackville, New Brunswick, E4L 1E7, Canada

75: Senckenberg Biodiversity and Climate Research Center, Georg-Voigt-Str. 14-16, Frankfurt am Main, 60325, Germany

76: Department of Biological Sciences, Goethe University, Senckenberganlage 31, Frankfurt am Main, 60325, Germany

77: SARTI-MAR Research Group, UPC-Vilanova, Universitat Politecnica de Catalunya, Rambla Exposició, 24, Vilanova i la Geltru, 8800, Spain

78: School of Geography, Geology and the Environment, Keele University, William Smith Building, Staffordshire, ST5 5BG, United Kingdom

79: Estación Biológica de Doñana-CSIC, CSIC, c/ Americo Vespucio 26, Sevilla, 41092, Spain

80: Laboratório de Comportamento Animal, Departamento de Zoologia, Universidade de Brasília, Instituto de Ciências Biológicas, Distrito Federal, 70910-900, Brazil

81: Laboratório de Ornitologia, Departamento de Botânica e Zoologia, Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, RN, 59078-900, Brazil

82: Programa de Pós-Graduação em Ecologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, RN, 59078-900, Brazil

83: Lifeplan Rio de Janeiro / Araçá project, Estrada Hans Garlipp, s/n, Macaé de Cima, Fazenda Bacchus, Nova Friburgo, 28605-270, Brazil

84: Forest, Nature and Landscape, KU Leuven, Celestijnenlaan 200e, Leuven, 3001, Belgium

85: ECOA - Laboratory of Bioacoustics, Post graduate Program in Vertebrate Biology / Museum of Natural Sciences, Pontifical Catholic University of Minas Gerais, Minas Gerais, Brazil

86: Environmental Studies Center (CEA), Sao Paulo State University - UNESP, Av. 24a, 1515, Bela Vista, Rio Claro, Sao Paulo, 13506-900, Brazil

87: Pacific Northwest Research Station, U.S.D.A. Forest Service, 3625 93rd Ave. SW, Olympia, Washington, 98512, U.S.A.

88: Conservation Metrics, Inc., 145 McAllister Way, Santa Cruz, CA, 95060, U.S.A.

89: Acoustics Program, Pacific Marine Envrionmental Laboratory, National Oceanic and Atmospheric Administration, 2030 SE Marine Science Drive, Newport, OR, 97365, U.S.A.

90: Reef Pulse S.A.S., NA, NA, Allée Maureau, Sainte-Clotilde, Reunion Island, 97490, France

91: UMR 9220 ENTROPIE, Faculté des sciences et technologies, University of Reunion Island, Avenue René Cassin, Sainte-Clotilde, Reunion Island, 97490, France

92: Wildlife Ecology & Management, Faculty of Agriculture, Yamagata University, 1-23 Wakabamachi, Tsuruoka, Yamagata, 9978555, Japan

93: Groningen Institute for Evolutionary Life-Sciences, GELIFES, University of Groningen, Nijenborgh 7, Groningen, 9747 AGH, The Netherlands

94: cE3c–Centre for Ecology, Evolution and Environmental Changes & CHANGE - Global Change and Sustainability Institute, Departamento de Biologia Animal. Faculdade de Ciências, Universidade de Lisboa, Portugal

95: National Museum of Natural Sciences, Spanish National Research Council, Calle José Gutiérrez Abascal 2, Madrid, 28006, Spain

96: Centre de Recherche sur la Biodiversité et l'Environnement (UMR 5300 CNRS-IRD-TINPT-UPS), Université Paul Sabatier, 118 route de Narbonne, Toulouse, 31062 Toulouse Cedex 9, France

97: Facultad de Ciencias, Universidad Autónoma de Madrid, C. Darwin 2, Madrid, 28049,
Spain
98: Centre for Taxonomy and Morphology, Leibniz Institute for the Analysis of Biodiversity
Change, Museum of Nature Hamburg, Martin-Luther-King-Platz 3, Hamburg, 20146,
Germany
99: Centre d'Ecologie et des Sciences de la Conservation (CESCO, UMR 7204), Museum
National d'Histoire Naturelle, Place de la Croix, Concarneau, Finistère, 29900, France
100: Laboratory of tropical ecology, A.N. Severtsov Institute of Ecology and Evolution of
Russian Academy of Sciences, Leninsky Prospect, 33, Moscow, 119071, Russia
101: Department of Health and Environmental Science, Xi'an Jiaotong-Liverpool University,
No. 8 Chongwen Road, Suzhou Industrial Park, Suzhou, Jiangsu, 215123, China
102: Ecology of Tropical Agricultural Systems, Institute of Agricultural Sciences in the
Tropics, University of Hohenheim, Garbenstrasse 13, Stuttgart, 70599, Germany
103: School of Biological Sciences , Life Sciences Building, University of Bristol, 24 Tyndall
Ave, Bristol, BS8 1TQ, United Kingdom
104: Wildlife & Habitat Assessment, Northern Region, Canadian Wildlife Service,
Environment and Climate Change Canada, 5019 - 52nd Street, Yellowknife, Northwest
Territories, X1A 2P7, Canada
105: Species and Habitat Conservation, Forest Nature Conservation, Northwest German
Forest Research Institute, Prof.-Oelkers-Str. 6, Hann. Münden, Lower Saxony, 34346,
Germany
106: Wildlife Conservation Society Canada, 169 Titanium Way, Whitehorse, Yukon, Y1A
0E9, Canada
107: Applied Zoology and Nature Conservation, Zoological Institute and Museum, University
of Greifswald, Loitzer Strasse 26, Greifswald, 17489, Germany
108: Bat Lab, Department of Evolutionary Ecology, Leibnitz Institute for Zoo and Wildlife
Research, Alfred-Kowalke-Straße 17, Berlin, 10315, Germany
109: Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire
naturelle, CNRS, Sorbonne Université, Ecole Pratique des Hautes Etudes, Université des
Antilles, 57 rue Cuvier, Paris, 75005, France
110: Cooperative Institute for Marine Ecosystem and Resources Studies, National Oceanic
and Atmospheric Administration Pacific Marine Environmental Laboratory and Oregon State
University, 2030 SE Marine Science Drive, Newport, OR, 97365, USA
111: Department of Biology, University of Oxford, Oxford, OX1 3SZ, United Kingdom
112: Bat Squad Lab, Department of Life Science, National Taiwan Normal University, No.
88, Section 4, Tingzhou Rd, Wenshan District, Taipei City, 116, Taiwan
113: Nature based Solutions Initiative, Department of Biology , University of Oxford, 11A
Mansfield Road, Oxford, OX13SZ, England
114: Tabin Lab, Department of Genetics, Harvard Medical School - Boston,
115: Centre for Northern Forest Ecosystem Research, Ministry of Northern Development,
Mines, Natural Resources and Forestry, Ontario Government, 103 – 421 James Street
South,Thunder Bay, Ontario, P7E 2V6, Canada
116: Swiss Ornithological Institute, Seerose 1, Sempach, 6204, Switzerland
117: Zoological Research Museum Alexander Koenig, Leibniz Institute for the Analysis of
Biodiversity Change , Adenauerallee 160, Bonn, 53113, Germany
118: Museum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung,
Invalidenstraße 43, Berlin, 10115, Germany
119: Ocean and Ecosystem Science Division, Department of Fisheries and Oceans, Bedford
Institute of Oceanography, 1 Challenger Drive, Dartmouth, Nova Scotia, B2Y42, Canada
120: Ecology and Nature Management, Geosciences and Natural Resource Management,
University of Copenhagen, Rolighedsvej 23, Frederiksberg, 1958, Denmark
121: Department of Agricultural Biology, Faculty of Agriculture, University of Ruhuna,
Mapalana, Kamburupitiya, x, Sri Lanka
122: Product R&D Department, Xueqing Road, Beijing, 100091, China

123: Fundación de Conservación Jocotoco, Fundación de Conservación Jocotoco,
Fundación de Conservación Jocotoco, Valladolid N24-414 y Luis Cordero, Quito, 170525,
Ecuador
124: Landscape Conservation Programme, Conservation Division, BirdLife South Africa,
Gauteng, South Africa
125: Afromontane Research Unit, Department of Zoology and Entomology, University of the
Free State, Phuthaditjhaba, South Africa
126: Centre for Statistics in Ecology, the Environment and Conservation, Department of
Statistical Sciences, University of Cape Town, Cape Town, South Africa
127: Nature Coast Biological Station, Institute of Food and Agricultural Sciences, University
of Florida, 552 1st Street, Cedar Key, Florida, 32625, U.S.A.
128: Fisheries and Aquatic Sciences, Institute of Food and Agricultural Sciences, University
of Florida, 136 Newins-Ziegler Hall, Florida, 32611, U.S.A.
129: Sound Forest Lab, The Nelson Institute of Environmental Studies and The Department
of Forest and Wildlife Ecologz, University of Wisconsin-Madison, 1650 Linden Drive,
Wisconsin, 53706, U.S.A.
130: DYNI, Laboratoire Informatique et Systèmes (LIS) CNRS UMR 7020, Université de
Toulon, Université d'Aix Marseille, La Garde, 83 130, France
131: Nature Conservation Unit, Frederick University, 7, Yianni Frederickou Street, Nicosia,
1036, Cyprus
132: Madagascar Programme, Chances for Nature, Brauweg 9a, Göttingen, 37073,
Germany
133: Cologne Zoo, Riehler Straße 173, Köln, Lower Saxony, 50735, Germany
134: Biodiversity, Macroecology and Biogeography, Faculty of Forest Sciences and Forest
Ecology, University of Goettingen, Büsgenweg 1, Göttingen, Niedersachsen, 37077,
Germany
135: Plant Ecology, Institute of Plant Sciences, University of Bern, Altenbergrain 21, Bern,
Bern, 3013, Switzerland
136: Pacific Wildlife Research Center, Environment and Climate Change Canada, 5421
Robertson Road, Delta, British Columbia, V4K 3Y3, Canada
137: Migratory Bird Unit, Northern Region, Canadian Wildlife Service, Environment and
Climate Change Canada, 91780 Alaska Highway, Whitehorse, Yukon Territory, Y1A 5B7,
Canada
138: Lancaster Environment Centre, Lancaster University, Library Ave, Bailrigg, Lancaster,
LA1 4YQ, United Kingdom
139: Environmental Research and Innovation Centre, School of Science, Engineering and
Environment, University of Salford, Peel Building, Salford, M5 4WT, United Kingdom
140: School for Forest Management, Swedish University of Agricultural Sciences SLU, Box
43, Skinnskatteberg, SE-739 21, Sweden
141: MARE – Marine and Environmental Sciences Centre / ARNET - Aquatic Research
Network, Agencia Regional para o Desenvolvimento da Investigação Tecnologia e Inovação
(ARDITI), Faculty of Life Sciences of University of Madeira, , Funchal, 9020-105, Portugal
142: Chair of Geobotany, Chair of Geobotany, University of Freiburg, Schaenzlestr. 1,
Freiburg im Breisgau, 79104, Germany
143: Fish Bioacoustics Lab, Departamento de Biologia Animal, Faculdade de Ciências da
Universidade de Lisboa, Campo Grande, Lisbon, 1749-016 , Portugal
144: Fisheries Ecology and Conservation Lab, Department of Biology , University of Victoria,
3800 Finnerty Rd., Victoria, British Columbia, V8P 5C2, Canada
145: Southern Seas Ecology Laboratories, School of Biological Sciences, The University of
Adelaide, DX 650 418, Adelaide, SA, 5005, Australia
146: Department of Botany and Biodiversity Research, Faculty of Life Sciences, University of
Vienna, Rennweg 14, Vienna, 1030, Austria
147: Agroecology, Department of Crop Sciences, University of Göttingen, Grisebachstrasse
6, Göttingen, Niedersachsen, 37077, Germany

148: Alliance Bioversity International - CIAT, Lima Office, NA, Av. La Molina 1084, La Molina, Lima, Peru

149: School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat Street, Seattle, Washington, 98105, U.S.A.

150: Polar Ecology Group, Department of Vertebrate Ecology and Zoology, University of Gdańsk, Poland, Wita Stwosza 59, Gdańsk, 80-308, Poland

151: Department of Ecology, University of Alicante, Cra. San Vicente del raspeig sn, Alicante, 3690, Spain

152: Laboratório de Ecologia, Instituto de Biociências, Universidade Federal de Mato Grosso do Sul, Brazil

153: CHESS Lab, Biology, Furman University, 3300 Poinsett Hwy, Greenville, SC, USA, USA

154: Institute of Marine Science, University of Auckland, Private Bag 92019. Auckland Mail Centre, Auckland, 1142, New Zealand

155: Freshwater and Oceanic Sciences Unit of Research, Deparment of Biology, Ecology, and Evolution, University of Liège, allée du six août B6c, Liège, Belgium

156: DECA, Neuropsi, CNRS - University Paris Saclay,

157: Department of Soil, Water, and Ecosystem Sciences, Institute of Food and Agricultural Sciences, University of Florida, 2181 McCarty Hall A, Florida, 32611, U.S.A.

158: MARE - Marine and Environmental Sciences Centre/ARNET - Aquatic Research Network, ISPA, Instituto Universitário, Lisbon, Portugal, Ecological behaviour, ISPA Instituto Universitario, Jardim do Tabaco 34, Lisbon, 1149-041, Portugal

159: QUT Ecoacoustics, Centre for the Environment, Queensland University of Technology, Brisbane, Queensland, 4001, Australia

160: Integrative Community Ecology Unit, Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa, 904-0495, Japan

161: Salmonid fishes, Norwegian Institute for Nature Research, Høgskoleringen 9, Trøndelag, Trondheim, 7034, Norway

162: Cold Climate Research Lab, School of Art, RMIT University, 124 La Trobe Street, Melbourne, Victoria, 3000, Australia

163: Fisheries and Oceans Canada, Freshwater Institute, 501 University Crescent, Winnipeg, Manitoba, R3T 2N6, Canada

164: Centre for Marine Science and Technology, Curtin University, Perth, 6102, Western Australia

165: Forest Zoology, TUD Dresden University of Technology, Pienner Str. 7, Tharandt, 1737, Germany

166: Ecosystem Sensing group, Department of Life Sciences, Imperial College London, White City Campus, 84 Wood Ln, London, W12 0BZ, United Kingdom

167: Fisheries Management Branch, Department of Forestry, Fisheries and the Environment, Foreshore, Cape Town, 8000, South Africa

168: Mammal Research Institute Whale Unit, Department of Zoology and Entomology, University of Pretoria, Hatfield, Pretoria, 28, South Africa

169: State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, 100093, China

170: Trondhjem Biological Station, Department of Biology, Norwegian University of Science and Technology, Bynesveien 46, Trondheim, 7018, Norway

171: Coastal Marine Field Station, School of Science, University of Waikato, 58 Cross Road, Sulphur Point, Tauranga, 3110, New Zealand

172: Museu Paraense Emílio Goeldi, Belém, PA, Brazil

173: Universidade Federal da Paraíba, João Pessoa, PB, Brazil

174: Ocean Acoustics Group, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Klussmannstr. 3d, Bremerhaven, 27570, Germany

175: College of Marine Living Resource Sciences and Management, Shanghai Ocean University, 999 Huchenghuan Road, Pudong New District, Shanghai, 201306, China

176: Department of Ecology, Universidad Autónoma de Madrid, Darwin 2, Madrid, 28049,
Spain
177: Centro de Investigación en Biodiversidad y Cambio Global, Universidad Autónoma de
Madrid, Darwin 2, Madrid, 28049, Spain
178: Wildlife Research Division, Environment and Climate Change Canada, 801-1550,
avenue d'Estimauville, Québec (Québec) , G1J 0C3, Canada
179: Natural Resources & Environmental Studies, University of Northern British Columbia,
3333 University Way, British Columbia, V2N 4Z9, Canada
180: Asian Soundscape Monitoring Network, Biodiversity Research Center, Academia
Sinica, 128 Academia Road, Section 2, Nankang, Taipei , 11529, Taiwan
181: Department of Ecology and Evolutionary Biology & K. Lisa Yang Center for
Conservation Bioacoustics, Cornell University, Corson Mudd, 215 Tower Road, Ithaca, New
York, 14853, U.S.A.
182: Department of Biology, Whitman College, 345 Boyer AVe, Walla Walla, Washington,
99362, U.S.A.
183: Talarak Foundation, Inc., Negros Forest Park, Bacolod City, Negros Occidental, 6100,
Philippines
184: Department of Forest Ecosystems and Society, Oregon State University, 3180 SW
Jefferson Way, Corvallis, Oregon, 97331, U.S.A.
185: Centre for Research and Conservation, Royal Zoological Society of Antwerp, Belgium
186: Department of Physical Geography, Stockholm University, SE-106 91 Stockholm,
Stockholm, 10691, Sweden
187: Ordway Lab of Ecosystem Conservation, Department of Biology, University of Florida,
876 Newell Dr, Gainesville, FL, 32611, USA
188: Ordway Lab of Ecosystem Conservation, Florida Museum of Natural History, University
of Florida, 1659 Museum Rd, Gainesville, FL, 32611, USA
189: State Key Laboratory of Biocontrol, School of Ecology, Sun Yat-sen University,
Shenzhen, China
458
459 [c]: corresponding authors
460

Ethics & Integrity

Authorship

The present study has involved people who carried out PAM-based studies as primary
contributors. Their willingness to be informed through a mailing list, responsibility for their
metadata, approval for sharing the meta-data publicly, and willingness to participate in this
study as co-authors were explicitly stated in an online form-based collaboration agreement.
Primary contributors who became co-authors all fulfilled either data curation (e.g., as
providers of structured metadata) or project administration (e.g., as principal investigators
designing the corresponding study) roles and additionally a manuscript revision role. They
could be corresponding authors for published studies, referred contacts for unpublished
studies, or principal investigators, and were asked to identify further primary and secondary
contributors of their study. Primary contributors could become co-authors, and secondary
contributors are acknowledged here. Some primary contributors were invited as co-leads to
expand the network for particular realms or biomes and are listed in the first-tier authors list.
Primary contributors who provided soundscape recordings additionally to metadata are also
listed in the first-tier authors list. All primary contributors were asked to identify further
contacts to reach a comprehensive coverage for the database.

Permits

Necessary permits from landowners or environmental protection agencies, precautions to limit biological contaminations (e.g., biofilm on marine recorders), approvals for animal welfare, and local ethics committee reviews were the responsibility of the respective primary contributors.

Data availability

The metadata used for all analyses except the case studies are archived in Zenodo: https://zenodo.org/records/12675124. We provide the R script for reproducing the metadata analysis and graphs, as well as the R script and data for reproducing the case study analysis and graph. The demonstration collection providing all case study recordings is hosted here: https://ecosound-web.de/ecosound_web/collection/show/49.

Conflict of interests

The authors declare to have no competing interests. All authors have seen and approved the manuscript; it hasn't been accepted or published elsewhere.

Funding

The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Government of any product or service. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official U.S. Government determination or policy. JGM was funded by Fundação para a Ciência e Tecnologia (FCT) under the Scientific Employment Stimulus - Institutional Call - (CEECINST/00037/2021). We acknowledge the NFDI Consortium Earth System Sciences — NFDI4Earth, coordinated by TU Dresden, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — project number: 460036893. BIOMON is funded by the European Union's Horizon Europe programme under grant agreement 101090273. Larissa Sayuri M. Sugai and Liiana Piatti acknowledge grant Fundect T.O.:95/2023; SIAFEM: 33112. This paper is NOAA-PMEL contribution number 5948. Songhai Li acknowledges the National Natural Science Foundation of China (Grant numbers 42225604). Anna Franziska Cord was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2070 – 390732324. This research was also funded by a Westlake University Startup Fund (Thomas C Wanger).

Acknowledgements

We are thankful to our colleagues who supported our work: Abigail Seybert, Abram B. Fleishman, Adriana C. Acero-Murcia, Adrien Charbonneau, Adrià López-Baucells, Aklavik, Alain Paquette, Albert García, Alexander C. Lees, Alexandra Buitrago-Cardona, Alfonso Zúñiga, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Allan G Oliveira, Almo Farina, Alpheous, Ambroise N. Zongo, Amy Oden, Ana Filipa Palmeirim, Ana Širović, Anamaria Dal Molin, Anastasia Viricheva, Andrea Gavio, Andrew N. Gweh, Andros T Gianuca, Annalea Beard, Anne Sourdril, Annebelle Kok, Anthony Tan, Anthony Truskinger, Anushka Rege, Aran Mooney, Arne Wenzel, Astrid Brekke Skrindo, Australian Institute of

Marine Science, Base de Estudos do Pantanal - UFMS, Bastien Castagneyrol, Bea Maas,
Benedictus Freeman, Bibiana Gómez-Valencia, Bob Dziak, Bobbi J. Estabrook, Borja Milá,
Brian Miller, Bridget Maher, C. Lisa Mahon, Carolline Z Fieker, Carolyn Rosten, Cat
Adshead, Catherine Potvin, Catrin Westphal, Chantal Huijbers, Charlie Griffiths, Charlotte
Roemer, Chistophe Thébaud, Chong Chen, Christina Ieronymidou, Christine Cassin,
Christine Erbe, Christopher Clark, Cicero Simao, Claire Attridge, Clara P. Amorim, Claudia
A. Medina-Uribe, Claudia Ascencio-Elizondo, Clément Lemarchand, Colin Bates, Colin KC
Wen, Conner Partaker, Cowichan Tribes Lulumexun department - Fisheries and Marine
projects staff, Craig A Radford, Curtin University, Cássio Alencar Nunes, Cécile Albert,
Cédric Gervaise, Dadang Dwi Putra, Daisy Dent, Dan Warren, Daniel Mihai Toma, Daniel
Rieker, Danny Swainson , David Lecchini, David Tucker, David Watson, Dawit Yemane,
Deanna Clement, Dedi Rahman, Denise Risch, Dennis Higgs, Dennis Kühnel, Dexter
Hodder, Didier Casane, Diego Alejandro Gómez-Montes, Diego Balbuena, Diego Gil, Diego
Pavón Jórdan, Diogo Provete, Dorgival Diogenes, Dustin Whalen, Edho Walesa Prabowo,
Eduardo Rodríguez Martinez, Elisa Schutz, Elizabeth Clingham, Eliziane Oliveira, Ellen Mc
Arthur, Elma Kay, Eloy Revilla, Emanuel Stefan Baltag, Emilia Sokolowska, Emilio Tan, En-
Hao Liu, Enoc Martinez, Eric Parmentier, Eric Rakotomalala, Erich Fischer, Erika Berenguer,
Estância Mimosa Ecoturismo, Evan Economo, Fabio De Leo Cabrera, Faisal Ali Awanrali
Khan, Fang-Yee Lin, Fannie W. Shabangu, Fernando Hidalgo, Filipe F de Deus, Flávio
Rodrigues, Frederic Sinniger, Frode Fossøy, Frédéric Bertucci, Frédéric Sèbe, Gabriel Leite,
Gabriel leite, George Best, Gerard Bota, Germany, Giselle Mahung, Gonzalo Pérez-
Rosales, Gonçalo Silva, Greg and Lisa Morgan, Groupe Chiroptères de Guyane, Guillermo
McPherson, Hailey Davies, Hajar, Hannah Reyes, Hanneline Smit-Robinson, Harris
Papadopoulos, Heriniaina Randrianarison, Hervé Jourdan, Hila shamon, Hiromi Kayama
Watanabe, Holger Kreft, Héloïse Rouzé, Ian Avery Bick, Ilias Karmiris, Irfan Fitriawan,
Isabelle Côté, Italo A A Rondon, Ivin Wilfred Raj, Jacques Keumo Kuenbou, Jakob
Tougaard, Jamie Ratliff, Janice Ser Huay Lee, Jason Gedamke, Jasper Kanes, Jean Pierre
Castro Namuche, Jean-Yves Royer, Jeff Robinson, Jennifer Clark, Jens-Georg Fischer,
Jessica Deichmann, Jessica Qualley, Jodanne Pereira, Johan Diepstraten, Johanna
Järnegren, John Ewen, John Hildebrand, John Ryan, Jos Barlow, Jose A. Hernandez-
Blanco, Jose Leon, Jose Manuel Ochoa-Quintero, José Luis Mena, José Nilson Santos,
José Wagner, João Gama Monteiro, Juan Diego Tovar, Juliana Carolina Herran García,
Julie Perrot, Julio Baumgarten, Junior Tremblay, Just T. Bayle Sempere, Justine Magbanua,
Jérémy Anso, Jérémy Froidevaux, Jérémy Froideveaux, Jérôme Sueur, Karen H. Beard,
Katie Huong, Katie Innes, Kelsie Murchie, Ken Otter, Ken P. Findlay, Kenneth Dudley, Kevin
Darras, Kevin G. Borja-Acosta, Kieran Cox, Kinga Buda, Kris Harmon, Laela Sayigh, Laetitia
Hédouin, Laure Desutter-Grandcolas, Laurel Symes, Lauren Kuehne, Laurence H. De
Clippele, Laurent Legendre, Leeann Henry, Leila Hatch, Lenka Brousset, Liana Chesini
Rossi, Lin Schwarzkopf, Linilson Padovese, Lisa Loseto, Louise Emmerson, Lourdes
Maigua Medrano, Luca Iacobucci, Lucia Di Iorio, Luciana Nacimento , Ludovic Crochard,
Luis P. da Silva, Lény Lego, M. Clara P. Amorim, Malika René-Trouillefou, Mao-Ning
Tuanmuu, Marc Fernandez, Marc Lammers, Marcia Maia, Marconi Campos, Marconi
Campos Cerqueira, Marcos Vaira, Marcus Rowcliffe, Maria Cielo Bazterrica, Marilyn
Beauchaud, Marina Puebla-Aparicio, Marlize Muller, Martim Melo, María Jesús García
Bianco, Mathias Igulu, Matias Carandell, Matthew J. Weldy, Matthew K Pine, Matthew
Slater, Mauricio Akmentins, Max Ritts, Michael Pashkevich, Michael Scherer-Lorenzen,
Michael Towsey, Miguel Pessanha Pais, Mike van der Schaar, Milton Cezar Ribeiro, Mina
Anders, Muhammad Justi Makmun Jusrin, Nadia Pieretti, Niaina Nirina Mahefa

Andriamavosoloarisoa, Nicholas Brown, Nicholas Friedman, Nick Gardner, Nicolas Farrugia,
Nicolas Hette-Tronquart, Nicolas Pajusco, Nikos Fakotakis, Noam Leader, Noelia Ríos,
Nuno Faria, Ocean Acoustics Group, Ocean Networks Canada Society, Olaf Boebel, Omar
Hiram Martinez Castillo, Paul McDonald, Paulo J. Fonseca, Pedro B. Lopes, Peter Ellis,
Peter Taylor, Phillip Eichinski, Pooja Choksi, Pousada Xaraés, Pshemek Zdroik, Rainforest
Connection (RFCx), Refúgio Ecológico Caiman, Refúgio da Ilha Ecolodge, Rex K. Andrew,
Ricardo Duarte, Richard Fuller, Richard Kinsey, Rob Rempel, Robert John Young, Robert
McCauley, Robert Spaan, Robert Young, Rodrigo Silva, Rouvah Andriafanomezantsoa, Saki
Harii, Samuel Challéat, Samuel Haché, Sandra Mueller, Sara Gonçalves, Sarah Rowley,
Sarika Khanwilkar, Sean Connell, Sean Martin, Sebastian Muñoz Duque, Shanghai Aquatic
Wildlife Conservation and Research Center, Shannon MacPhee, Shinsuke Kawagucci,
Sierra Hunicutt, Sierra Jarriel, Sigal Balshine, Silent cities consortium, Simon Childerhouse,
Sina Weier, Sofie Van Parijs, Soledad Gaston, Stan Dosso, Stanislas Wroza, Stephanie
Plön, Stephen Insley, Stuart Marsden, Stéphan Jacquet, Stéphane Père, Susanna Piovano,
Sérgio Moreira, Tatiana Atemasova, Tetsuya Miwa, The Hunters and Trappers Committees
of Inuvik, The Inuvialuit-Canada Fisheries Joint Management Committee, Thiago M Ventura,
Thomas Sattler, Todor Ganchev, Tomas Altamirano, Tomasz Stanislaw Osiejuk, Tomaz
Melo, Tomonari Akamatsu, Tristan Louth-Robins, Tulio Rossi, Tómas Grétar Gunnarsson,
Under The Pole Consortium, Valentin Chevalier, Valerie Linden, Valéria Tavares, Vijay
Ramesh, Vincent Médoc, Vu Dinh Thong, Vy Tram Nguyen, Wen-Ling Tsai, Will Duguid,
Yang Liu, Yenifer Herrera-Varón, You-Fang Chen, Yue Qiu, Yuhang Song, Zehava Sigal,
Zuania Colón-Piñeiro, Zuzana Burivalova, and Tuktoyaktuk, e, Çağlar Akçay, and Éric
Parmentier.

Main Text

Title

Worldwide Soundscapes: a synthesis of passive acoustic monitoring across realms

Running Title

passive acoustic monitoring synthesis

Abstract and keywords

Aim: The urgency for remote, reliable, and scalable biodiversity monitoring amidst mounting human pressures on ecosystems has sparked worldwide interest in Passive Acoustic Monitoring (PAM), which can track life underwater and on land. However, we lack a unified methodology to report this sampling effort and a comprehensive overview of PAM coverage to gauge its potential as a global research and monitoring tool. To remediate this, we created the Worldwide Soundscapes project, a collaborative network and growing database comprising metadata from 409 datasets across all realms (terrestrial, marine, freshwater, and subterranean).

Location: Worldwide, 12 200 sites, all ecosystems.

Time period: 1991 to present.

Major taxa studied: All soniferous taxa.

Methods: We synthesise sampling coverage across spatial, temporal, and ecological scales using metadata describing sampling locations, deployment schedules, focal taxa, and recording parameters. We explore global trends in biological, anthropogenic, and geophysical sounds based on 168 recordings from twelve ecosystems across all realms.

Results: Terrestrial sampling is spatially denser (45 sites/Mkm^2) than aquatic sampling (0.3 and 1.8 sites/Mkm^2 in oceans and fresh water) with only two subterranean datasets. Although diel and lunar cycles are well-covered in all realms, only marine datasets (56%) comprehensively sample all seasons. Across twelve ecosystems, biological sounds show contrasting diel patterns, decline with distance from the equator and negatively correlate with anthropogenic sounds.

Main conclusions: PAM can inform macroecology studies as well as global conservation and phenology syntheses, but representation can be improved by expanding terrestrial taxonomic scope, sampling coverage in the high seas and subterranean ecosystems, and spatio-temporal replication in freshwater habitats. Overall, this global PAM network holds promise to support global biodiversity research and monitoring efforts.

Keywords: ARU, automated sound recorder, biodiversity, conservation biology, ecoacoustics, ecology, IUCN GET realm, passive acoustic monitoring, phenology, soundscape ecology

Introduction

Sounds permeate all realms on Earth — terrestrial, freshwater, marine, and subterranean[1]. Passive Acoustic Monitoring (PAM) captures soundscapes that document soniferous (i.e., sound-producing) organisms, human activities, and some geophysical events (i.e., biophony, anthropophony, and geophony, respectively). In ecoacoustics and soundscape ecology[2,3], PAM can measure levels and impacts of global change (e.g., climate change, urbanization, deep-sea mining)[4–6]; monitor ecosystem health, recovery, and restoration[7–9]; assess human-environment interactions (e.g., public health, cultural ecosystem services)[10,11]; and guide environmental management and conservation policies (e.g., protected areas, landscape planning)[12,13].

Despite the wide-ranging and increasing soundscape sampling effort[14,15], its global distribution across realms remains undescribed. Soundscape-recording communities are currently only networked within realms, often differing methodologically. Previous reviews focused on single realms and were either systematic[14–17] or non-systematic qualitative[18,19] reviews that cannot describe data trends. The systematic reviews did not quantitatively address taxonomic focus in marine studies, nor ecosystemic terrestrial coverage, or spatio-temporal sampling distribution in the freshwater realm, while the subterranean realm has not been reviewed yet. Notably, all reviews so far used only published data and involved only a small part of their respective communities. Marine scientist networks using PAM exist[20], but the freshwater community is nascent, and the terrestrial community is fragmented among single taxa. Methodological differences are also striking: acoustic calibration and sound propagation modelling are advanced in aquatic studies[21] but seldom considered in terrestrial ones (but see[22,23]); artificial intelligence can identify increasing numbers of species on land[24], whereas most aquatic sounds are still challenging to identify[25,26].

Overall, building a united global PAM network can increase knowledge transfer resulting in more efficient and consistent methods, analyses, and cross-system syntheses[14]. Cross-realm PAM studies can yield new theoretical answers[27] and applied solutions: organisms' sound durations follow a common distribution across multiple realms[28]; soundscapes track terrestrial and marine resilience to disturbance regimes[29]. Transnational sampling could form the basis for comprehensive soniferous biodiversity monitoring, just as community-initiated telemetry databases[30] and collaborative camera trap surveys[31] have advanced entire research fields. A global PAM network could complement existing biodiversity-monitoring networks, establish historical biodiversity baselines, support systematic long-term and large-scale monitoring, and connect with the public through citizen science. Such information is critical to inform global biodiversity policies[32] such as the Kunming-Montreal Global Biodiversity Framework.

We present the "Worldwide Soundscapes" project, the first global PAM meta-database and network (https://ecosound-web.de/ecosound_web/collection/index/106). We used it to quantify the known state of PAM efforts, highlight apparent sampling gaps and biases, illustrate the potential of cross-realm PAM syntheses for research, and federate PAM users. The project currently comprises 351 active contributors who collated metadata from 409 passively-recorded, stationary, replicated soundscape datasets. Metadata describe the exact spatio-temporal coverage, sampled ecosystems (International Union for Conservation of Nature Global Ecosystem Typology: IUCN GET), transmission medium (air, water, or soil),

focal taxa (IUCN Red list), recording settings, as well as data and publication availability. We
inferred coverage within administrative (Global ADMinistrative Database: GADM;
International Hydrographic Organisation: IHO) and protected areas (World Database on
Protected Areas: WDPA) from geographic locations. We selected recordings referenced in
the meta-database to quantify soundscape components (biophony, anthropophony, and
geophony) across twelve ecosystems from all realms. We showcase their relevance to
exemplary macroecology, conservation biology, and phenology research questions and
identify opportunities to advance the global PAM network. The publicly-accessible meta-
database[33] continues to grow to enhance accessibility of data and remains open for
metadata contributions, facilitating future syntheses.

Methods

Database construction

The database construction started in August 2021 within the frame of the Worldwide
Soundscapes project using collaborative, peer-driven metadata collation[33]. It represents the
current state of knowledge of PAM within our network. We additionally conducted focal
publication searches to plug coverage gaps by inviting the respective corresponding authors.
We posted the call for contributors on specialised ecoacoustics platforms and social media,
and keep the project open for any contributor owning suitable soundscape recordings. We
communicated by emails in English, Spanish, French, Portuguese, German, Russian, and
Chinese to federate users in our network. Out of 592 contacted contributors potentially
involved in PAM, 63% already provided metadata, 23% have not yet responded, 12% are
pending, and 2% have declined. We included metadata from larger groups such as the
Silent Cities project, Ocean Networks Canada, and the Australian Acoustic Observatory.
Primary contributors provided the metadata and bear the responsibility of their accuracy.
Research assistants checked the coherence of the metadata input (beyond automated data
format checks) and primary contributors cross-validated metadata displayed as maps and
graphical timelines. The database information page and content is integrated in our online
collaborative ecoacoustics platform ecoSound-web[34] that also hosts the annotated
soundscape recordings of the case study
(https://ecosound-web.de/ecosound_web/collection/show/49).

Soundscape recording datasets were required to meet four criteria: 1) stationary — mobile
recorders have variable spatial assignments, thus we excluded recordings from cars,
transect walks, or towed deployments; 2) passive — obtained from unattended recorders; 3)
ambient — omni-directional, non-triggered recordings, under non-experimental conditions; 4)
spatially or temporally replicated (Fig. 1) — as to disentangle spatial and temporal effects
from other soundscape determinants. Datasets were defined as spatially replicated when
several sites were sampled simultaneously, and temporally replicated when a site was
sampled over multiple days at the same time of day. Sampling sites and days were our
elemental units for defining replication, however, in other contexts, spatial replicates may for
instance be required to be in the same habitat, and temporal replicates could be defined
across multiple full moon nights. Taken together, our requirements homogenise the dataset
to enable general, unified statistical analyses across datasets for future syntheses.

Time and space

Soundscapes result from geophysical phenomena as well as wildlife and human activities that are broadly determined by solar and lunar cycles, and geographical positions on the planet relative to the poles or equator, or the land and water surface. We defined and calculated spatial coverage as the number of sites, and spatial density as the number of sites per the containing realms' areal extent. Alternatively, spatial extent could have been defined as the area bounded by sites, but calculating extents on the world sphere is conceptually challenging for large extents. Further, the sampling area covered at each site is generally unknown: they are rarely measured in terrestrial sites but sometimes simulated in marine environments[35]. Detection spaces vary with sound source intensity, frequency, directivity; recording medium temperature, currents, pressure; habitat structure; and ambient sound level. Underwater, detection spaces are greater due to the higher density of the recording medium. Our measure of spatial sampling density using points per area is thus provisional. Temporal extent was defined and calculated as the time range from the start of the first to the end of the last recording, temporal coverage as the time sampled per site, and temporal density as the proportion of time sampled within the temporal extent.

We quantified latitudinal and topographical distribution by collecting coordinates, altitude, and depth data for each site. Topography values on land were either provided by the contributors or filled in using General Bathymetric Chart of the Oceans surface elevation data[36]. For underwater sites, depth values below the water surface were provided by the contributors, and for subterranean sites, depth below the land surface was recorded. We assigned sampling sites to administrative areas (GADM divisions for freshwater and land, IHO for sea areas) and extracted their WDPA category. Sites' climates were geographically classified into tropical (between −23.5° and 23.5° latitude), polar (below −66.5° or above 66.5° latitude), and temperate (between polar and tropical regions).

Primary contributors coded their exact recording times for each site using deployment start and end dates and times and operation modes (or combinations thereof). Continuous operations lasted from the deployment start to its end; scheduled operations had daily start and end times; periodical operations used duty cycles. A temporal framework was devised to quantify sampling coverage in three solar and lunar cycles using the timing of sound recordings relative to time events (Fig. 3). Seasonal coverage was inferred only for temperate sites, by splitting the year into four meteorological seasons (winter: December-February, spring: March-May, summer: June-August, fall: September-November, reversed for Southern latitude sites). The daily cycle was split into four diel windows delimiting dawn (from astronomical dawn start at -18° solar altitude until 18° solar altitude), day, dusk (from 18° solar altitude to astronomical dusk end at -18° solar altitude), and night. The lunar illumination cycle was split into two time windows centred on the full and new moon phases. Thus, extrema and ecotones in the temporal cycles define time windows, and in temperate zones, equinoxes roughly correspond to thermal ecotones. Seasonal cycles in tropical and polar regions arising from precipitation patterns were not considered in this analysis.

Ecological characterisation

We assigned individual sampling locations to ecosystem types following the IUCN GET (https://global-ecosystems.org). Sites were assigned hierarchically to realms (core or

transitional ones, which represent the interface between core realms), biomes, and
functional groups. We calculated major occurrence areas of all functional groups based on
ecosystem maps[37] to quantify spatio-temporal extent, coverage, and sampling density within
realms and biomes. Notably, only the result section "Sampling in ecosystems" separates
results among core and transitional realms. For the spatial and temporal result sections,
sites were assigned to their "parent" realm (i.e., the first mentioned realm in the compound
transitional realm names). Deployments were linked to IUCN Red List taxa (class, order,
family, or genus) when studies were designed for monitoring these taxa, and other
deployments could be collected without taxonomic focus.

Acoustic frequency ranges

Sound recordings store a representation of the original soundscape, so we needed to
determine their spectral scope. Microphones (including hydrophones and geophones) have
variable frequency responses, usually declining with frequencies above the human-audible
range. Additionally, digital recorders restrict the spectral scope of the recording with the
sampling frequency. Contributors provided audio parameters for their deployments: sampling
rate, high-pass filters, microphone and recorder models.

Soundscape case studies

To illustrate how the database can be used for macroecology, conservation biology, and
phenology analyses, we selected 168 recordings across a variety of topographical,
latitudinal, and anthropisation conditions — all fundamental gradients of assembly filters in
both terrestrial and marine realms[1] — belonging to 12 IUCN GET functional groups. We
aimed for four spatial replicates within the same functional group, with 10-minute audible
sound recordings (at least 44.1 kHz sampling frequency) starting at sunrise, solar noon,
sunset, and solar midnight from the same date during the biologically active season. The
available data did not always allow this (Table S2). We acknowledge that this targeted, non-
systematic selection is not statistically representative of global patterns but rather illustrative
of the database potential.

Each soundscape recording was analysed with respect to the three fundamental
soundscape components: biophony, anthropophony, and geophony. Soundscape recordings
were uploaded to ecoSound-web[34] for annotation
(https://ecosound-web.de/ecosound_web/collection/show/49): KD listened to them while
visually inspecting their spectrograms (Fast Fourier Transform window size of 1 024) at a
density of 1 116 pixels per 10 minutes. Annotations were rectangular boxes with defined
coordinates in the time and frequency dimensions on the spectrogram, bounding only the
annotated sound. Annotations of different soundscape components could overlap if they
were simultaneously visible or audible. The total expanse of annotations of the same
soundscape component was computed, excluding overlaps. Soundscape components above
22.05 kHz and sounds caused by microphone or recorder self-noise were excluded. All
annotations were reviewed by the recordists using the peer-review mode on ecoSound-web.
Acoustic space occupancy for each soundscape component in each recording was
calculated as the proportion of the sampled spectro-temporal space[38] (i.e., total annotation
area divided by total area of spectrogram; range: 0-1).

We asked whether biophony occupancy increases with proximity to the equator, whether
biophony was negatively correlated with anthropophony, and whether phenology patterns
differ across realms. Statistical models predicting biophony occupancy were Bayesian beta
regression models (4 chains of 1000 sampling iterations with 1000 warmup iterations) fitted
with the R package brms[39]. Models converged as determined by trace plots and R hat values
smaller than 1.1. The number of samples equaled the number of recordings (N=168). The
models using latitude and anthropophony were mixed-effect models including the functional
group as a random intercept. The phenology model used diel time windows and realms, as
well as their interactions, as predictors.

Results

Summary dataset statistics

To date, 409 validated soundscape meta-datasets (hereafter "datasets") have been
registered in our database from across the globe, dating back to 1991 (Fig. 1D). A dataset
gathers a team's metadata on a study or project. Based on the IUCN GET definition of four
'core' realms, our database includes 277, 104, 26, and 2 validated datasets from the
terrestrial, marine, freshwater, and subterranean realms, respectively. The transmission
medium was air for terrestrial datasets (except for soil in one dataset), mostly water for
aquatic datasets (eight above-water datasets mostly in freshwater realm). In the
subterranean realm, one dataset did aerial recordings, while the other did underwater
recordings. The majority of datasets (84%) include both spatial and temporal replicates (Fig.
1B). Few datasets have openly-accessible recordings (10-12% excluding subterranean
realm, Fig. S1). Presently, few terrestrial and freshwater datasets (27% and 19%,
respectively) are associated with DOI-referenced publications in contrast to marine datasets
(44%, excluding subterranean realm) (Fig. S1).

Spatial sampling coverage and density

The database contains 12 220 sampling sites, including 147 polar, 9 194 temperate, and 2
859 tropical sites (Fig 1C). On land, 11 229 sites are located within 85 (out of 263) GADM
level 0 areas (i.e., countries, Fig. 2B), primarily in the Northern Hemisphere (Table S1).
Administratively, most terrestrial sites occur in Canada (27%), followed by the United States
(18%), but a significant proportion is globally widely distributed (25% do not belong to the top
10 GADM areas). Few terrestrial sites (8%) are located in WDPA category Ia, Ib, or II areas,
corresponding to the highest protection levels. Our database currently lacks data from vast
areas in Russia, Greenland, the Antarctic, North Africa, and Central Asia. Site elevations
range from sea level up to 4 548 m (Fig. 2A), but mountains above 4 000 m in the Northern
Hemisphere, and above 2 000 m in the Southern Hemisphere (except for the Kilimanjaro),
as well as the Transantarctic Mountains, are currently not represented in the data. At sea,
634 sites are located within 35 (out of 101) IHO sea areas. Administratively, most marine
sites are widespread among IHO areas (23% do not belong to the top 10 IHO areas), but the
North Pacific Ocean and the Southeast Alaskan and British Columbian coastal waters
contain large parts of marine sites (16% and 13%, respectively). Many sites are situated in
WDPA high-protection category Ia and II areas (12%). Our database currently lacks datasets
from Arctic waters off Eurasia, and Southeast Asian coastal areas. Sampling sites span
ocean depths from sea water surface to depths of 10 090 m, but tropical bathypelagic and

Southern benthic areas are poorly represented (Table S2). Few GADM areas (16) are
represented in the 324 freshwater sites. Spain holds most freshwater sites (56%), followed
by China (21%), and few freshwater sites (2%) are in WDPA category II or Ib areas.
Freshwater bodies are sampled at elevations from sea level up to 3 770 m and were
sampled up to 25 m depth. Mountain freshwater bodies and those in Africa, Asia, and
Oceania are currently poorly represented in our database (only one site at 3770 m). The
database contains 13 subterranean sites situated in Brazil and Mexico between 277 and 810
m elevation, and up to a depth of 1 546 m (horizontal projection).

Temporal sampling extent, coverage, and density

We compare sampling coverage (in years sampled by recordings, summed over sites)
across time windows of the diel cycle, the lunar phases, and the seasons (Fig. 3A). Dawn
and dusk diel windows are shorter than day and night diel windows for most locations and
correspondingly less intensively sampled. The lunar phase cycle is evenly covered across
realms and comprehensively covered within datasets (Fig. S2). In the terrestrial realm,
daytime coverage surpasses nighttime coverage (606 vs. 395 years), while 74% of datasets
sampled all diel time windows. Terrestrial temperate datasets mostly sampled spring (411
years, 32%) and summer (559 years, 44%) while 26% sampled all seasons. Terrestrial
temporal coverage per site is highest in Australia (1541 days). In the marine realm, diel
coverage is even, as 89% of marine datasets sampled all diel time windows. Marine
temperate datasets have high and similar coverage for winter and spring (117 and 122
years, combined 59% of seasonal coverage) and 64% cover the full seasonal cycle. Marine
temporal coverage per site is highest in the Balearic (2771 days). In the freshwater realm,
temporal coverage among diel time windows is even and 81% of freshwater temperate
datasets sampled all diel time windows. Similarly, 13% sampled all seasons. Freshwater
temporal coverage per site is highest in Portugal (3 009 days). The subterranean tropical
sites primarily covered the nighttime.

Sampling in ecosystems

Our database includes 83 of the 110 IUCN GET functional groups and all biomes except
anthropogenic shorelines, the subterranean tidal biome, anthropogenic subterranean voids,
and anthropogenic subterranean freshwaters (Table S2). The terrestrial realm has the third-
largest extent and the highest spatial sampling density among realms (45 sites per Mkm^2
over entire temporal extent), but temporal coverage is comparatively low (30% sampled out
of 203 days of mean extent per site). The most commonly sampled terrestrial biome is the
temperate-boreal forests and woodlands biome (56% of sites). The marine realm is the most
extensive but spatial sampling density is the lowest (0.3 sites per Mkm^2), while temporal
sampling coverage is the highest among all realms (66% out of 379 days sampled). The
most commonly sampled marine biomes are the marine shelf and pelagic ocean waters
(56% and 33% of sites respectively). The freshwater realm has low spatial sampling
densities (1.8 sites per Mkm^2) and high temporal sampling densities (67% out of 238 days
sampled). Lakes are the most commonly sampled freshwater biome (48% of sites). The
terrestrial-freshwater realm, representing 81% of the area of non-subterranean realms, has
the third-highest spatial sampling density (7.9 sites per Mkm^2) and similar temporal sampling
density to the terrestrial realm (18% out of 233 days sampled). The marine-freshwater-
terrestrial realm (including 26 sites in coastal river deltas, saltmarshes, and intertidal forests)
has the largest temporal extent (465 days per site). The subterranean realm, though second-

largest, includes seven tropical sites (all in tropical aerobic caves) sampled with a low
temporal coverage (1% out of 129 days sampled), and the subterranean-freshwater realm
includes five sites in underground streams and pools, sampled with a high temporal
coverage (17% out of 507 days sampled).

Target taxa and frequency ranges

Most marine datasets do not target specific taxa (66%) and use wide frequency ranges from
0.01 to 31 kHz (mean bounds of frequency ranges across datasets, Fig. 4B). Marine
datasets that focus on single taxa comprise fish (12%, 0.002 - 20 kHz) and cetaceans (6%,
0.006 - 7 kHz). Similarly, most freshwater datasets are taxonomically unspecific (56%) and
cover frequencies from 1 to 29 kHz. Some datasets (14%) focus on ray-finned fish, covering
frequencies from 0.001 to 23 kHz. In contrast, terrestrial datasets mostly target single taxa
and have narrow frequency ranges. Bird-focused datasets are most common (44%),
spanning frequencies from 0.06 to 21 kHz, while bat-focused datasets are next (12%) and
range from 5 to 139 kHz. Taxonomically unspecific datasets account for 24% of terrestrial
datasets, covering a broad range from 0.2 to 23 kHz. Generally, datasets targeting multiple
taxa use wider frequency ranges than those targeting single taxa.

Soundscape case studies

We analysed 168 recordings from twelve ecosystems (corresponding to different IUCN GET
functional groups) representing the diversity of soundscapes on Earth, spanning latitudes
from 69 degrees South to 67 degrees North (Table S3, Fig. 5A). Biophony dominated with
an average soundscape occupation of 28% across all ecosystems. Notable examples
include the photic coral reefs in Okinawa, Japan, with snapping shrimps and grunting fish
choruses, and the tropical lowland rainforests in Jambi, Indonesia, with buzzing insects and
echoing bird and primate songs, showing soundscape occupancy of 75% and 61%,
respectively[40]. Only marine island slopes (off Sanriku, Japan) and polar outcrops (Antarctic)
contained no or very little (2%) biophony, respectively. Geophony was absent in most of our
soundscape samples, with the exception of high wind noise in polar outcrops (14%) and
some wind in montane tropical forests and urban ecosystems (both 5%). Anthropophony
occupied on average 11% of the soundscapes. Cities (Jambi, Indonesia; Montreal, Canada)
exhibited the highest anthropophony (43%) with prevalent engine noise and human voices,
while deep-sea mining and vessel communication signals caused high anthropophony in
marine island slopes (32%)[41]. Silence occupancy was highest in bathypelagic ocean water
(96%) and polar outcrops (82%).

The selected soundscapes reveal greater biological activity closer to the equator, a negative
relationship between biophony and anthropophony, and variable phenology patterns of
soniferous organisms over the diel cycle (Fig. 5). We detected a negative correlation of
biophony occupancy with increasing distance from the equator ($P_{negative}$=1) and with
anthropophony occupancy ($P_{negative}$=0.98). The phenology model predicted biophony
occupancy values for each diel time window and realm (Fig. 5B), revealing similar phenology
for the terrestrial and marine realm, and opposed freshwater and freshwater-marine realm
phenology.

Discussion

The "Worldwide Soundscapes" project has — to our knowledge — assembled the first global meta-database of PAM datasets across realms. We analysed its current content to quantify sampling extent, coverage, and density across spatiotemporal and ecological scales. We analysed soundscapes from twelve ecosystems to investigate macroecological, conservation biology, and phenological trends. The database remains open for contributions and can be openly accessed to source datasets and initiate collaborative studies[33]. Next, we discuss the state and potential of PAM globally (Table 1). While we focuse on large-scale research questions here, the database can also be used to address more focused questions, for instance using the finely-resolved taxonomic information that we recorded, or for within-realm syntheses.

Our results likely represent global PAM trends, even though they may be biased by the project contributors' background. Our terrestrial spatial coverage is similar to a recent systematic review[14]. Our database gaps in North Africa and Northeastern Europe correspond with the paucity of bioacoustic datasets for these regions in the Xeno-Canto bioacoustic repository[42]. Our database comprises 634 marine sampling locations, while a recent systematic review compiled 991[15] from published data. Although the latter review's locations are represented at the dataset level (which can comprise several sites), most overlap with our finer site-level locations (Fig S4). Marine tropical waters that are under-represented in our database reflect gaps found in the International Quiet Ocean Experiment network coverage[43]. Our marine and terrestrial database's spatial coverage is thus broadly comparable with published data, but it is more detailed as the exact sampling times and locations are known. Our database's temporal coverage is also more finely resolved and thus not directly comparable with previous work. To our knowledge, no other spatially-explicit review of freshwater sampling or synthesis of subterranean PAM coverage exists for comparison. Finally, as our database originates from an active network of researchers, it represents the current availability of mostly as-yet unpublished data (Fig. S1).

Geographic coverage strikingly differs among realms: marine coverage is sparse but widespread; terrestrial coverage is comparatively intensive in the Americas, Australia, and Western Europe; freshwater coverage is scattered. This partly reflects the gap between high-income countries, which have the resources to carry out conservation and active ecosystem management actions; and developing countries, which paradoxically contain the majority of biodiversity hotspots (and some of the most densely populated areas). Accessibility and technical limits in extreme environments also drive geographic patterns: high latitudes and elevations entail extremely cold temperatures that present challenges for operation and maintenance, some of which can be solved with robust power setups (solar panels, freeze-resistant batteries). Marine deployments are generally even more constrained due to costly and demanding underwater work, but some deployments reach the polar regions as water temperatures are buffered below the freezing point. Affordable underwater recorders[44] may help to intensify sampling of marine coastal areas and close gaps in freshwater coverage. By contrast, terrestrial monitoring is often straightforward. Northern temperate areas outside of Northeastern Europe are comparatively better covered, and the tropics outside of Africa better represented. It appears that gaps in North Africa, Central Asia, and Northeastern Europe arise from unequal research means and priorities between

countries. Taken together, these gaps will help to correct spatial biases[45] and to identify
high-priority, unique research areas that should be included in global assessments.

Currently, only marine studies achieve relatively even coverage of temporal cycles. Indeed,
offshore deployments — especially in the deep sea — are expensive and limited duration
deployments are not cost-effective[46]. Marine soundscapes fluctuate stochastically[47], but the
ocean buffers water temperatures so that animals retain a basal activity level year-round.
Although lunar phases affect marine life[48–50] and shape tidal ecosystems, we did not consider
lunar tides. In the terrestrial and freshwater realms, most deployments cover the entire diel
cycle but monitoring on land often focuses either on diurnal birds or on nocturnal bats, as
found in the literature[14]. In contrast, although seasons also drive acoustic activity cycles on
land[51,52], spring- and summertime monitoring is disproportionately common and we lack a
thorough understanding of seasonal dynamics. Terrestrial deployments in particular may be
short for logistic reasons: in the cold, batteries struggle and access is harder; in arid regions,
fire hazards complicate long-term deployments; recorders are generally at risk of theft; and
limited equipment may cycle between sites[53]. Lunar phases — probably evenly sampled by
chance — also influence land animals[54] but should be explicitly considered in study designs.
Overall, we encourage longer-duration setups with regularly-spread sampling inside
temporal cycles to alleviate the higher expenses for energy and storage, as well as trade-
offs with spatial coverage[55]. Global changes impact soundscapes in largely unpredictable
ways through changing species distributions and phenology, necessitating higher and
unbiased coverage across multiple time scales — including inter-annual ones — to
successfully monitor ongoing changes[56].

Re-use of soundscape datasets is restricted by their taxonomic focus. Admittedly,
taxonomically untargeted, long, and regular deployments in oceans, coupled to large
detection ranges, concurrently sample many taxa[57]. However, many soundscape recordings
sample particular frequencies, often in the human-audible range[38], although biophony ranges
from infrasound to ultrasound. For instance, studies of toothed whales or bats often use
triggers and high-pass filters to cope with high data storage demands by recording purely
ultrasonic recordings only when signals are detected, resulting in spectrally-restricted and
temporally-biased soundscape recordings. Less-studied taxa, such as anurans and insects,
could effectively be co-sampled by adjusting ongoing terrestrial deployments. We encourage
terrestrial researchers to maximise frequency ranges (and to broaden diel coverage — see
above) to enhance interdisciplinary collaboration. These collaborations can help to mutualise
resources for mitigating potentially prohibitive power, storage, transportation, and post-
processing costs[22]. Emerging embedded-AI audio detectors may offer an alternative to
continuous and broadband recording[58], but whole soundscape recordings will remain
essential for broader application.

In every realm, ecosystems await acoustic discovery. Except for one dataset from aerobic
caves, we currently lack data from all subterranean realms (anthropogenic voids, ground
streams, sea caves), while endolithic systems may be the only ecosystem for which PAM is
probably irrelevant. Access is usually challenging or restricted for non-specialists, but
subterranean biodiversity shows high spatial turnover[59]. Freshwater data were less rare, and
several datasets with unreplicated sampling could not be included. Temporary, dynamic
water bodies (seasonal, episodic, and ephemeral ecosystems), although prevalent[60], are not
yet well-studied (Table S2) although most are accessible from land. Notably, aquatic realms
feature important peculiarities: extraneous sounds from the air can be captured in so-called

holo-soundscapes in freshwater and shallow coastal areas[61], and particle motion (that
accompanies sound) impacts aquatic organisms[62]. Advances in soundscape research are
imminent as the freshwater acoustic research community is growing rapidly[63]. In the oceans,
sound propagates comparatively far and multiple biomes can be sampled at once (e.g.,
recorders on the seafloor sample pelagic waters too) so that most ecosystems are covered.
Still, coverage gaps may exist in rhodolith/Maërl beds. On land, sampling is biased towards
biodiverse forests, and rocky habitats (young rocky pavements, lava flows and screes) but
also some vegetated temperate ecosystems (cool temperate heathlands, temperate
subhumid grasslands) are poorly sampled. Within the IUCN GET framework, soil
soundscapes also belong to the terrestrial realm and to date only one spatio-temporally
replicated dataset is in our database, despite recent studies[64,65]. Soil and benthic habitats
may also be sampled with geophones to record infrasonic vibrations used to sense the
environment (e.g., by insects, frogs, elephants, interstitial fauna and benthic fish)[66].

Our database highlights well-known global sampling biases[67] which could be resolved with
collaboration and communication to remove cultural and socioeconomic barriers[68].
Technological progress for more affordable equipment renders PAM more accessible in
lower-income countries[44,69]. However, high- and deep-sea work remains considerably more
expensive, and tropical developing countries in particular often lack funding for marine
programmes requiring large vessels, underwater vehicles, or cabled stations on the
seafloor[46]. Our network currently consists of active members from 56 countries.
Collaborative projects, shared sea missions, and equipment loans should promote the
establishment of soundscape research communities[70]. However, equitable, collaborative
efforts will also require capacity-building so that local researchers can independently
process, analyse, and interpret PAM-derived data. Increased international collaboration with
scientists and local stakeholders supporting citizen-science[71] in heavily underrepresented
regions would improve not only data coverage, but also representation and dialogue within
the field.

Collaborative soundscape research relies on interoperable data. We harmonised metadata
with a bottom-up approach leading to our global inventory. Comprehensive standards for
PAM deployment, reporting, and data analysis are needed for enabling comparative, global
analyses. However, such standards do not currently exist, even though initiatives for the
marine realm are ongoing[72,73]. Few affordable solutions exist for sharing large audio data
volumes[34], underlining the need for distributed soundscape recording repositories[74]. Marine
oil and gas industry projects routinely upload data as part of their efforts to mitigate noise
impacts on marine animals[75], but these recordings often focus on frequencies relevant to
seismic prospecting and access may be restricted[76]. Furthermore, recording equipment
requires calibration, sound detection spaces need to be measured[23,77], and data privacy must
be ensured on land[78]. In parallel, species sound libraries[42,79–81] grow and continue to provide
invaluable acoustic and taxonomic references without which soundscape analysis for
soniferous organisms would be futile. International organisations such as the Global
Biodiversity Information Facility (GBIF) will be key to roll-out standards (e.g., Darwin Core) in
a top-down manner. For the moment, we encourage early planning of data archival. In the
future, the Worldwide Soundscapes will interoperate with other databases to close remaining
coverage gaps and enhance standardisation.

A unified approach to ecology with PAM is now possible. More comprehensive coverage is
needed to decisively answer research, conservation, and management questions that PAM

can address, so we encourage prospective contributors to curate their metadata and join our inclusive project. We show that a large portion of the PAM community is willing to collaborate across realms and form a global network. We advocate for a bolder PAM effort to inform the agenda of soundscape ecology[82], reaching out to places where no sound has been recorded before as well as to urban settings[83]. The research community may open new avenues to study environmental effects on acoustic activity[56], social species interactions[84], human-wildlife relationships[40], function and phylogeny[85], soundscape effects on human health[86], acoustic adaptation and niche hypotheses[87,88], macroecological patterns across ecosystems[1], and initiate an integrated approach to noise impacts on wildlife.

PAM is an established method that can be applied over large spatial and temporal scales. However, consistent, large-scale monitoring of the Earth's soundscapes is direly needed and essential to establish baselines for historical trends[89] and quantify rapid changes in biodiversity and natural systems. International funding schemes should integrate PAM into biodiversity monitoring platforms such as GBIF[90] and GEO BON[91,92]. Soundscapes are just starting to be used in legislation as an ecosystem feature to be preserved[93]. Occupancy maps for soniferous wildlife obtained from PAM would underpin the evaluation of progress towards threat reduction and ecosystem service provision of the Kunming-Montreal Global Biodiversity Framework[94]. By building collaborations around the knowledge frontiers identified here, we can aim to comprehensively describe and understand the acoustic make-up of the planet.

References

References

1. Keith, D. A. *et al.* A function-based typology for Earth's ecosystems. *Nature* 1–6 (2022)

doi:10.1038/s41586-022-05318-4.

2. Pijanowski, B. C., Farina, A., Gage, S. H., Dumyahn, S. L. & Krause, B. L. What is

soundscape ecology? An introduction and overview of an emerging new science.

Landsc. Ecol. **26**, 1213–1232 (2011).

3. Sueur, J. & Farina, A. Ecoacoustics: the Ecological Investigation and Interpretation of

Environmental Sound. *Biosemiotics* **8**, 493–502 (2015).

4. Kang, J. *et al.* Supportive soundscapes are crucial for sustainable environments. *Sci.*

Total Environ. **855**, 158868 (2023).

5. Sueur, J., Krause, B. & Farina, A. Climate Change Is Breaking Earth's Beat. *Trends*

Ecol. Evol. **34**, 971–973 (2019).

6. Williams, R. *et al.* Noise from deep-sea mining may span vast ocean areas. *Science*

377, 157–158 (2022).

7. Müller, J. *et al.* Soundscapes and deep learning enable tracking biodiversity recovery in

tropical forests. *Nat. Commun.* **14**, 6191 (2023).

8. Ross, S. R. P.-J. *et al.* Divergent ecological responses to typhoon disturbance revealed

via landscape-scale acoustic monitoring. *Glob. Change Biol.* **30**, e17067 (2024).

9. Sethi, S. S. *et al.* Characterizing soundscapes across diverse ecosystems using a

universal acoustic feature set. *Proc. Natl. Acad. Sci.* (2020)

doi:10.1073/pnas.2004702117.

10. Alvarsson, J. J., Wiens, S. & Nilsson, M. E. Stress Recovery during Exposure to Nature

Sound and Environmental Noise. *Int. J. Environ. Res. Public. Health* **7**, 1036–1046

(2010).

11. Chen, Z., Hermes, J., Liu, J. & von Haaren, C. How to integrate the soundscape

resource into landscape planning? A perspective from ecosystem services. *Ecol. Indic.*

141, 109156 (2022).

12. Haver, S. M. *et al.* Comparing the Underwater Soundscapes of Four U.S. National Parks
and Marine Sanctuaries. *Front. Mar. Sci.* **6**, 500 (2019).

13. Holgate, B., Maggini, R. & Fuller, S. Mapping ecoacoustic hot spots and moments of
biodiversity to inform conservation and urban planning. *Ecol. Indic.* **126**, 107627 (2021).

14. Sugai, L. S. M., Silva, T. S. F., Ribeiro, J. W. & Llusia, D. Terrestrial Passive Acoustic
Monitoring: Review and Perspectives. *BioScience* **69**, 15–25 (2019).

15. Havlik, M.-N., Predragovic, M. & Duarte, C. M. State of Play in Marine Soundscape
Assessments. *Front. Mar. Sci.* **9**, (2022).

16. Greenhalgh, J. A., Genner, M. J., Jones, G. & Desjonquères, C. The role of freshwater
bioacoustics in ecological research. *WIREs Water* **7**, e1416 (2020).

17. Scarpelli, M. D. A., Ribeiro, M. C., Teixeira, F. Z., Young, R. J. & Teixeira, C. P. Gaps in
terrestrial soundscape research: It's time to focus on tropical wildlife. *Sci. Total Environ.*
707, 135403 (2020).

18. Duarte, C. M. *et al.* The soundscape of the Anthropocene ocean. *Science* **371**, (2021).

19. Lindseth, A. V. & Lobel, P. S. Underwater Soundscape Monitoring and Fish
Bioacoustics: A Review. *Fishes* **3**, 36 (2018).

20. Boyd, I. *et al.* An International Quiet Ocean Experiment. *Oceanography* **24**, 174–181
(2015).

21. Wang, L. S. *et al. Review of Underwater Acoustic Propagation Models.*
https://eprintspublications.npl.co.uk/6340/ (2014).

22. Sousa-Lima, R. A Review and Inventory of Fixed Autonomous Recorders for Passive
Acoustic Monitoring of Marine Mammals. *Aquat. Mamm.* **39**, 23–53 (2013).

23. Haupert, S., Sèbe, F. & Sueur, J. Physics-based model to predict the acoustic detection
distance of terrestrial autonomous recording units over the diel cycle and across
seasons: Insights from an Alpine and a Neotropical forest. *Methods Ecol. Evol.* **n/a**,
(2022).

24. Nieto-Mora, D. A., Rodríguez-Buritica, S., Rodríguez-Marín, P., Martínez-Vargaz, J. D. &
Isaza-Narváez, C. Systematic review of machine learning methods applied to

ecoacoustics and soundscape monitoring. *Heliyon* **9**, e20275 (2023).

25. Looby, A. *et al.* Global inventory of species categorized by known underwater sonifery.

Sci. Data **10**, 1–10 (2023).

26. Parsons, M. J. G. *et al.* A Global Library of Underwater Biological Sounds (GLUBS): An

Online Platform with Multiple Passive Acoustic Monitoring Applications. in *The Effects of*

Noise on Aquatic Life: Principles and Practical Considerations (eds. Popper, A. N.,

Sisneros, J., Hawkins, A. D. & Thomsen, F.) 1–25 (Springer International Publishing,

Cham, 2023). doi:10.1007/978-3-031-10417-6_123-1.

27. Ross, S. R. P.-J. *et al.* Passive acoustic monitoring provides a fresh perspective on

fundamental ecological questions. *Funct. Ecol.* **37**, 959–975 (2023).

28. de Sousa, I. P. *et al.* Scale-free distribution of silences. *Phys. Rev. E* **105**, 014107

(2022).

29. Gottesman, B. L. *et al.* What does resilience sound like? Coral reef and dry forest

acoustic communities respond differently to Hurricane Maria. *Ecol. Indic.* **126**, 107635

(2021).

30. Kays, R. *et al.* The Movebank system for studying global animal movement and

demography. *Methods Ecol. Evol.* **13**, 419–431 (2022).

31. Kays, R. *et al.* SNAPSHOT USA 2020: A second coordinated national camera trap

survey of the United States during the COVID-19 pandemic. *Ecology* **103**, e3775 (2022).

32. Moersberger, H. *et al.* Biodiversity monitoring in Europe: User and policy needs.

Conserv. Lett. **n/a**, e13038.

33. Darras, K. F. A. *et al.* Worldwide Soundscapes project meta-data. Zenodo

https://doi.org/10.5281/zenodo.10598949 (2024).

34. Darras, K. F. A. *et al.* ecoSound-web: an open-source, online platform for ecoacoustics.

F1000Research vol. 9 1224 Preprint at https://doi.org/10.12688/f1000research.26369.2

(2023).

35. *Exploring Animal Behavior Through Sound: Volume 1: Methods*. (Springer International

Publishing, Cham, 2022). doi:10.1007/978-3-030-97540-1.

36. GEBCO. Gridded bathymetry data (General Bathymetric Chart of the Oceans). *GEBCO*

https://www.gebco.net/data_and_products/gridded_bathymetry_data/.

37. Keith, D. A. *et al.* Indicative distribution maps for Ecosystem Functional Groups - Level 3

of IUCN Global Ecosystem Typology. Zenodo https://doi.org/10.5281/zenodo.4018314

(2020).

38. Luypaert, T. *et al.* A framework for quantifying soundscape diversity using Hill numbers.

Methods Ecol. Evol. **13**, 2262–2274 (2022).

39. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat.*

Softw. **80**, 1–28 (2017).

40. Lin, T.-H., Sinniger, F., Harii, S. & Akamatsu, T. Using Soundscapes to Assess Changes

in Coral Reef Social-Ecological Systems. *Oceanography* (2023)

doi:10.5670/oceanog.2023.s1.7.

41. Chen, C., Lin, T.-H., Watanabe, H. K., Akamatsu, T. & Kawagucci, S. Baseline

soundscapes of deep-sea habitats reveal heterogeneity among ecosystems and

sensitivity to anthropogenic impacts. *Limnol. Oceanogr.* **66**, 3714–3727 (2021).

42. Xeno-canto Foundation. *Xeno-Canto: Sharing Bird Sounds from around the World*.

(Xeno-canto Foundation Amsterdam, 2012).

43. IQOE. Acoustic Capabilities of Existing Observing Systems.

https://www.iqoe.org/systems.

44. Lamont, T. A. C. *et al.* HydroMoth: Testing a prototype low-cost acoustic recorder for

aquatic environments. *Remote Sens. Ecol. Conserv.* **8**, 362–378 (2022).

45. Beck, J., Böller, M., Erhardt, A. & Schwanghart, W. Spatial bias in the GBIF database

and its effect on modeling species' geographic distributions. *Ecol. Inform.* **19**, 10–15

(2014).

46. Rountree, R. *et al.* Towards an Optimal Design for Ecosystem-Level Ocean

Observatories. in 79–106 (2020). doi:10.1201/9780429351495-2.

47. Siddagangaiah, S., Chen, C.-F., Hu, W.-C. & Farina, A. The dynamical complexity of

seasonal soundscapes is governed by fish chorusing. *Commun. Earth Environ.* **3**, 1–11

(2022).

48. Hernández-León, S. *et al.* Zooplankton abundance in subtropical waters: is there a lunar cycle? *Sci. Mar.* **65**, 59–64 (2001).

49. Simonis, A. E. *et al.* Lunar cycles affect common dolphin Delphinus delphis foraging in the Southern California Bight. *Mar. Ecol. Prog. Ser.* **577**, 221–235 (2017).

50. Mougeot, F. & Bretagnolle, V. Predation risk and moonlight avoidance in nocturnal seabirds. *J. Avian Biol.* **31**, 376–386 (2000).

51. Krause, B., Gage, S. H. & Joo, W. Measuring and interpreting the temporal variability in the soundscape at four places in Sequoia National Park. *Landsc. Ecol.* **26**, 1247 (2011).

52. Grinfeder, E. *et al.* Soundscape dynamics of a cold protected forest: dominance of aircraft noise. *Landsc. Ecol.* **37**, 567–582 (2022).

53. Sugai, L. S. M., Desjonquères, C., Silva, T. S. F. & Llusia, D. A roadmap for survey designs in terrestrial acoustic monitoring. *Remote Sens. Ecol. Conserv.* **6**, 220–235 (2020).

54. Kronfeld-Schor, N. *et al.* Chronobiology by moonlight. *Proc. R. Soc. B Biol. Sci.* **280**, 20123088 (2013).

55. Wilgenburg, S. L. V. *et al.* Evaluating trade-offs in spatial versus temporal replication when estimating avian community composition and predicting species distributions. *Avian Conserv. Ecol.* **19**, (2024).

56. Desjonquères, C. *et al.* Acoustic species distribution models (aSDMs): A framework to forecast shifts in calling behaviour under climate change. *Methods Ecol. Evol.* **13**, 2275–2288 (2022).

57. Lillis, I. V. & Boebel, O. Marine soundscape planning: Seeking acoustic niches for anthropogenic sound. *J. Ecoacoustics* **2**, 1–1 (2018).

58. Höchst, J. *et al.* Bird@Edge: Bird Species Recognition at the Edge. in *Networked Systems: 10th International Conference, NETYS 2022, Virtual Event, May 17–19, 2022, Proceedings* 69–86 (Springer-Verlag, Berlin, Heidelberg, 2022). doi:10.1007/978-3-031-

17436-0_6.

59. Zagmajster, M., Malard, F., Eme, D. & Culver, D. C. Subterranean Biodiversity Patterns

from Global to Regional Scales. in *Cave Ecology* (eds. Moldovan, O. T., Kováč, Ľ. &

Halse, S.) 195–227 (Springer International Publishing, Cham, 2018). doi:10.1007/978-3-

319-98852-8_9.

60. Messager, M. L. *et al.* Global prevalence of non-perennial rivers and streams. *Nature*

594, 391–397 (2021).

61. Rountree, R. A., Juanes, F. & Bolgan, M. Temperate freshwater soundscapes: A

cacophony of undescribed biological sounds now threatened by anthropogenic noise.

PLOS ONE **15**, e0221842 (2020).

62. Popper, A. N. & Hawkins, A. D. The importance of particle motion to fishes and

invertebrates. *J. Acoust. Soc. Am.* **143**, 470–488 (2018).

63. Linke, S. *et al.* Freshwater ecoacoustics as a tool for continuous ecosystem monitoring.

Front. Ecol. Environ. **16**, 231–238 (2018).

64. Metcalf, O. C. *et al.* Listening to tropical forest soils. *Ecol. Indic.* **158**, 111566 (2024).

65. Maeder, M., Guo, X., Neff, F., Mathis, D. S. & Gossner, M. M. Temporal and spatial

dynamics in soil acoustics and their relation to soil animal diversity. *PLOS ONE* **17**,

e0263618 (2022).

66. Šturm, R., López Díez, J. J., Polajnar, J., Sueur, J. & Virant-Doberlet, M. Is It Time for

Ecotremology? *Front. Ecol. Evol.* **10**, (2022).

67. Hughes, A. C. *et al.* Sampling biases shape our view of the natural world. *Ecography* **44**,

1259–1269 (2021).

68. Amano, T., González-Varo, J. P. & Sutherland, W. J. Languages Are Still a Major Barrier

to Global Science. *PLOS Biol.* **14**, e2000933 (2016).

69. Hill, A. P., Prince, P., Snaddon, J. L., Doncaster, C. P. & Rogers, A. AudioMoth: A low-

cost acoustic device for monitoring biodiversity and the environment. *HardwareX* **6**,

e00073 (2019).

70. Reboredo Segovia, A. L., Romano, D. & Armsworth, P. R. Who studies where? Boosting

tropical conservation research where it is most needed. *Front. Ecol. Environ.* **18**, 159–

166 (2020).

71. Newson, S. E., Evans, H. E. & Gillings, S. A novel citizen science approach for large-

scale standardised monitoring of bat activity and distribution, evaluated in eastern

England. *Biol. Conserv.* **191**, 38–49 (2015).

72. Wall, C. C. *et al.* The Next Wave of Passive Acoustic Data Management: How

Centralized Access Can Enhance Science. *Front. Mar. Sci.* **8**, (2021).

73. Roch, M. A. *et al.* Management of acoustic metadata for bioacoustics. *Ecol. Inform.* **31**,

122–136 (2016).

74. Sugai, L. S. M. & Llusia, D. Bioacoustic time capsules: Using acoustic monitoring to

document biodiversity. *Ecol. Indic.* **99**, 149–152 (2019).

75. Southall, B. L. *et al.* MARINE MAMMAL NOISE-EXPOSURE CRITERIA: INITIAL

SCIENTIFIC RECOMMENDATIONS. *Bioacoustics* **17**, 273–275 (2008).

76. Haver, S. M. *et al.* Monitoring long-term soundscape trends in U.S. Waters: The

NOAA/NPS Ocean Noise Reference Station Network. *Mar. Policy* **90**, 6–13 (2018).

77. Darras, K., Pütz, P., Fahrurrozi, Rembold, K. & Tscharntke, T. Measuring sound

detection spaces for acoustic animal sampling and monitoring. *Biol. Conserv.* **201**, 29–

37 (2016).

78. Cretois, B., Rosten, C. M. & Sethi, S. S. Voice activity detection in eco-acoustic data

enables privacy protection and is a proxy for human disturbance. *Methods Ecol. Evol.*

13, 2865–2874 (2022).

79. Görföl, T. *et al.* ChiroVox: a public library of bat calls. *PeerJ* **10**, e12445 (2022).

80. Parsons, M. J. G. *et al.* Sounding the Call for a Global Library of Underwater Biological

Sounds. *Front. Ecol. Evol.* **10**, (2022).

81. Looby, A. *et al.* FishSounds version 1.0: A website for the compilation of fish sound

production information and recordings. *Ecol. Inform.* 101953 (2022)

doi:10.1016/j.ecoinf.2022.101953.

82. Pijanowski, B. C. *et al.* Soundscape Ecology: The Science of Sound in the Landscape.

BioScience **61**, 203–216 (2011).

83. Hao, Z. *et al.* Can urban forests provide acoustic refuges for birds? Investigating the influence of vegetation structure and anthropogenic noise on bird sound diversity. *J. For. Res.* **35**, 33 (2024).

84. Briefer, E. F. *et al.* The power of sound: unravelling how acoustic communication shapes group dynamics. *Philos. Trans. R. Soc. B Biol. Sci.* **379**, 20230182 (2024).

85. Gasc, A. *et al.* Assessing biodiversity with sound: Do acoustic diversity indices reflect phylogenetic and functional diversities of bird communities? *Ecol. Indic.* **25**, 279–287 (2013).

86. Buxton, R. T., Pearson, A. L., Allou, C., Fristrup, K. & Wittemyer, G. A synthesis of health benefits of natural sounds and their distribution in national parks. *Proc. Natl. Acad. Sci.* **118**, (2021).

87. Hart, P. J. *et al.* Timing Is Everything: Acoustic Niche Partitioning in Two Tropical Wet Forest Bird Communities. *Front. Ecol. Evol.* **9**, (2021).

88. EY, E. & FISCHER, J. The "Acoustic Adaptation Hypothesis"—a Review of the Evidence from Birds, Anurans and Mammals. *Bioacoustics* **19**, 21–48 (2009).

89. Pilotto, F. *et al.* Meta-analysis of multidecadal biodiversity trends in Europe. *Nat. Commun.* **11**, 3486 (2020).

90. Global Biodiversity Information Facility (2024). Free and Open Access to Biodiversity Data. What is GBIF? https://www.gbif.org/what-is-gbif (2024).

91. Gonzalez, A. *et al.* A global biodiversity observing system to unite monitoring and guide action. *Nat. Ecol. Evol.* 1–5 (2023) doi:10.1038/s41559-023-02171-0.

92. Towards a Transnational Acoustic Biodiversity Monitoring Network (TABMON). https://www.nina.no/english/TABMON.

93. Leiper, A. What does the consultancy industry need from academia? A soundscape and planning perspective. in *ACOUSTICS 2020* (Institute of Acoustics, Virtual, 2020). doi:10.25144/13347.

94. Batist, C. & Campos-Cerqueira, M. *Harnessing the Power of Sound and AI to Track*

1303   *Global Biodiversity Framework (GBF) Targets*. (2023).

1304   95. Keith, D. A., Ferrer-Paris, J. R., Nicholson, E. & Kingsford, R. T. *The IUCN Global*

1305   *Ecosystem Typology 2.0: Descriptive Profiles for Biomes and Ecosystem Functional*

1306   *Groups*. (IUCN, Gland, Switzerland, 2020).

Tables

1307

1308   Table 1: An agenda towards a global PAM coverage and network

Aims	Opportunities
Increasing spatial and ecosystemic coverage	- Affordable underwater recorders - Autonomous setups for extreme environments on land - Deploy in subterranean realm - Deploy in dynamic freshwater bodies - Deploy at high latitude and elevation on land
Increasing temporal and taxonomic coverage	- Longer deployments with duty cycles on land - Cover all seasons in freshwater and on land - Scale up to inter-annual cycles - Higher sampling frequencies - Disabling high-pass filters and triggers
Increasing collaboration	- Work interdisciplinarily with speleologists, urban ecologists, soil and deep-sea benthic scientists - Foster international missions to work at large scales - Collaborate, invest, and fund work with under-represented countries - Disseminate results in more languages
Interoperability	- Calibrate equipment and consider detection ranges - Build accessible and sustainable data repositories - Design and adopt standards for deployment, reporting, and analysis - Integrate PAM into biodiversity and remote sensing databases

1309

Figure Legends

Figure 1: Overview of Worldwide Soundscapes meta-database. *A) Framework used to define spatial and temporal replicates. B) Number of datasets in each core realm for the different replication levels. C) Spatial extent and coverage, based on sampling sites, split by core realm. Due to their higher representation and to avoid overlapping site clusters, terrestrial site densities were plotted on a 3 degree resolution raster (Interactive map: https://ecosound-web.de/ecosound_web/collection/index/106). D) Temporal extent and coverage, based on recorded days, split by core realm. An enlarged version of panel D without terrestrial sites can be found in Fig S3. For panels C and D, sites from transitional realms were assigned to their parent core realm.*

Figure 2: Spatial distribution of sampling sites. *A) Latitudinal and topographic distribution of sampling sites across core realms. Due to their higher representation and to avoid overlapping site clusters, terrestrial sites are shown with transparency. The minimum (deepest seafloor) and maximum (highest elevation of land or sea level) topographical limits (dark grey lines) are shown against latitude, based on General Bathymetric Chart of the Oceans data[36]. Minimum topography above the sea level and maximum topography under the sea level were set to zero as the sea level represents the minimum and maximum in these cases. B) Number of sampling sites within different administrative regions (GADM level 0 and IHO sea areas), split by core realm, across WDPA categories (Ia: strict nature reserve; Ib: wilderness area, II: national park). The areas that do not belong to the top 10 in terms of datasets have been aggregated under "others". Six subterranean sites in Brazil are not shown. Sites from transitional realms were assigned to their parent core realm.*

Figure 3: Temporal sampling distribution. *A) Temporal sampling coverage across solar and lunar cycles for the three core realms. Cycles consist of solar (daily and seasonal) and lunar time cycles (lunar phase), subdivided in time windows. Seasons were only analysed in temperate regions. Sampling coverage is represented with sampling days in number labels. B) Mean number of sampling days per site within different administrative regions (GADM level 0, IHO areas, WDPA categories), split by core realm. WDPA categories (Ia: strict nature reserve; Ib: wilderness area, II: national park) are shown separately. Six subterranean sites in Brazil are not shown. Numbers to the right of bars indicate the number of sites the means were calculated from. Sites from transitional realms were assigned to their parent core realm.*

Figure 4: Sampling distribution across ecological scales. *A) Spatial extent of realms, based on major areas according to IUCN GET (coloured disk area proportional to area);*

spatial sampling density (in sites per Mkm2) and coverage (in number of sites); temporal extent (mean range between first and last recording day), coverage (days sampled per site and density (proportion of days sampled per extent). B) Frequency ranges of datasets across realms (using Nyquist frequency i.e., actual recorded frequencies) for the main studied taxa. The dots at the ends of coloured lines represent means of the lowest and highest recorded frequencies, and the range between the minimum and maximum of these values are indicated with black error bars. The limits of human hearing are indicated with dashed lines. Number of datasets indicated above lines - datasets can be counted several times if they contain deployments targeting different taxa. Data from transitional realms were assigned to their parent core realm in panel B.

Figure 5: Soundscape components analysis. *A) Mean acoustic space occupancy of soundscape components (biophony, geophony, anthropophony) as calculated from annotations for twelve selected ecosystems, measured in proportion of spectro-temporal space used, over 168 recordings covering the time windows of the diel cycle. Annotated recordings are accessible at https://ecosound-web.de/ecosound_web/collection/show/49. Sample spectrograms are shown in the background. B) Soundscape component occupancy data illustrate three research questions linked to macroecology (i.e., biophony with distance from equator relationship), conservation biology trade-offs (i.e., biophony with anthropophony relationship); and phenological trends (i.e., mean biophony along diel time windows for each realm). Gray ribbons indicate 95% credible intervals and numbers indicate probabilities of positive or negative relationships.*

A - Framework

Space

Spatial replicates

Temporal replicates

Time

B - Replication (dataset level)

	temporal	spatial	spatio-temporal
Terrestrial	21	8	247
Subterranean			2
Marine	27	1	76
Freshwater	9	9	18

C - Spatial extent and coverage of sampling sites

Terrestrial sites
600
400
200

D - Temporal extent and coverage (day level)

Sampled days
1500
1000
500
0

1990   2000   2010   2020

Core realm: Freshwater   Marine   Subterranean   Terrestrial

A - Topographical and latitudinal distribution of sampling sites

IUCN core realms
- Freshwater
- Marine
- Subterranean
- Terrestrial

B - Top 10 administrative regions for spatial sampling coverage

Freshwater
- Spain — 41%
- China — 21%
- United States — 10%
- United Kingdom — 7%
- Colombia — 4%
- France — 4%
- others (not top 10) — 3%
- Australia — 3%
- Guinea-Bissau — 2%
- Brazil — 2%
- Canada — 1%

Marine
- others (not top 10) — 23%
- North Pacific Ocean — 16%
- Southeast Alaska and British Columbia Coastal Waters — 13%
- South Pacific Ocean — 8%
- South Atlantic Ocean — 7%
- Southern Ocean — 7%
- Mozambique Channel — 6%
- Caribbean Sea — 6%
- The Northwestern Passages — 5%
- Norwegian Sea — 4%
- Indian Ocean — 4%

Terrestrial
- Canada — 27%
- others (not top 10) — 25%
- United States — 18%
- Germany — 8%
- Brazil — 5%
- France — 5%
- Portugal — 3%
- Indonesia — 2%
- Australia — 2%
- Spain — 2%
- Norway — 2%

WDPA categories
- Ia
- Ib
- II
- >II

A - Sampling coverage within time cycles

Freshwater diel | Marine diel | Terrestrial diel | Subterranean diel

Time windows: Dawn, Day, Dusk, Night

Freshwater lunar | Marine lunar | Terrestrial lunar | Subterranean lunar

Bright, Dark

Freshwater seasonal | Marine seasonal | Terrestrial seasonal | Subterranean seasonal

Spring, Summer, Autumn, Winter

Sampling years

B - Top 10 administrative regions for temporal sampling coverage

Days per site

WDPA categories: Ia, Ib, II, NA

A - Sampling intensity across realms

A - Case study soundscapes

Large lowland rivers

Small permanent freshwater lakes

Riverine estuaries

Photic coral reefs

Island slopes

Bathypelagic ocean waters

Tropical lowland rainforests

Tropical montane rainforests

Plantations

Urban ecosystems

Polar outcrops

Aerobic caves

soundscape occupancy

Soundscape component: ▇ anthropophony ▇ biophony ▇ geophony

B - Illustrative analyses

Macroecology

P(>0) = 1

Biophony

Latitude

Conservation biology

P(<0) = 0.98

Biophony

Anthropophony

Phenology

Biophony

Diel time windows: morning, noon, evening, night

Realms: ● Freshwater ● Freshwater-Marine ● Marine ● Subterranean ● Terrestrial