Worldwide soundscape ecology patterns across realms

Authors & affiliations

Kevin FA Darras[1,2C], Rodney Rountree[3,4], Steven Van Wilgenburg[5], Lijun Dong[6], Amandine Gasc[7], Frederik Pitz[8], Youfang Chen[2], Tzu-Hao Lin[9], Patrick Mauritz Diaz[10], Shih-Hung Wu[11], Marcus Salton[12], Sarah Marley[13], Anna F Cord[14,15], Orlando Acevedo-Charry[16,17], Matyáš Adam[18], Leandro Aparecido Do Nascimento[19], Christos Astaras[20], Andrey Atemasov[21,22], Luc Barbaro[23], Carly Batist[24,25], Adriá López Baucells[26], Just T Bayle Sempere[27], Kristen Bellisario[28], Asaf Ben David[29], Oded Berger-Tal[30], Matthew Betts[31], Iqbal Bhalla[32], Thiago Bicudo[24], Marta Bolgan[33], Sara Bombaci[34], Martin Boullhesen[35], Tom Bradfer-Lawrence[36,37], Robert A Briers[38], Michal Budka[39], Alice Calvente[40], Maite Cerezo-Araujo[41], Gunnar Cerwén[42], Camille Desjonquères[43,44], Pedro Diniz[45], Adam Duarte[46], Hiroto Enari[47], Haruka S Enari[47], Nina Ferrari[31], Bárbara Freitas[48,49], Nick Friedman[50], Jérémy Froidevaux[36,51], Svetlana Gogoleva[52], José Miguel González Correa[27], Eben Goodale[53], Jack Greenhalgh[54], Samuel Hache[55], Jonas Hagge[56], William Halliday[57], Tara Hanf-Dressler[58], Alain Jacot[59], Olaf Jahn[60,61], Sebastian Kepfer-Rojas[62], Yael Lehnardt[30], José Leon[63], Kyle John Lloyd[64,65,66], Audrey Looby[67,68], David López-Bosch[26], Christos Mammides[69], Gabriel Marcacci[59], Matthias Markolf[70,71], Marinez Isaac Marques[72], Charles W Martin[67], Dominic A Martin[73,74], Christoph Meyer[75], Sandra Müller[76], Maria Mas Navarro[26], Rym Nouioua[77], Steffen Oppel[37,59], Anna Osiecka[78], Miles Parsons[57], Filipa Isabel Pereira Samarra[79], Jean-Nicolas Pradervand[59], Cristian Pérez-Granados[80], Laura K Reynolds[81], Klaus Riede[60], Ricardo Rocha[82], Samuel RP-J Ross[83], Kevin Scharffenberg[84], Karl-Ludwig Schuchmann[72], Esther Sebastián-González[80], David Singer[56], Juan Traba[85,86], Junior A Tremblay[87], Sunny Tseng[88], Marisol Valverde[89], Ben Vernasco[46,90], Matthew Ward[91], Matthew Weldy[31,92], Paul Jacobus Wensveen[79], Heather Wood[93], Luis P da Silva[94,95], Ana Silvia de Oliveira Tissiani[72], Devin de Zwaan[96,97], Songhai Li[6C], Renata Sousa-Lima[98C], Thomas Cherico Wanger[2C]

[C]: corresponding authors

1: EFNO, ECODIV, INRAE, Domaine des Barres, Nogent-sur-Vernisson, Centre-Val de Loire, 45290, France
2: Sustainable Agricultural Systems & Engineering Lab, School of Engineering, Westlake University, No. 600 Dunyu Road, Sandun Town, Xihu District, Hangzhou, Zhejiang province, , China
3: , Biology Department, University of Victoria, Cunningham Building, Room 202, PO Box 1700 STN CSC, Victoria, BC, V8W 2Y2, Canada
4: The Fish Listener, , , 23 Joshua Lane, Waguoit, MA, 2536, USA
5: Terrestrial Unit, Prairie Region, Canadian Wildlife Service, Environment & Climate Change Canada, Prairie & Northern Wildlife Research Centre, 115 Perimeter Road, Saskatoon, Saskatchewan, S7N 0X4, Canada
6: Marine Mammal and Marine Bioacoustics Laboratory, Department of Deep Sea Science, Institute of Deep-sea Science and Engineering Chinese Academy of Sciences, 28 Luhuitou Road, Sanya, Hainan, 572000, China
7: Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, , Aix Marseille Univ, Avignon Univ, CNRS, IRD, Campus Aix Technopôle de l'environnement Arbois Méditerranée Avenue Louis Philibert Bât Villemin, Aix-en-Provence, 13545, France
8: Functional Agrobiodiversity, Department of Crop Sciences, University of Göttingen, Grisebachstrasse 6, Göttingen, Niedersachsen, 37077, Germany

9: Marine Ecoacoustics and Informatics Lab, Biodiversity Research Center, Academia Sinica, 128 Academia Road, Section 2, Nankang, Taipei , , 11529, Taiwan

10: Climate Change and Biodiversity, , Hanns R. Neumann Stiftung Indonesia, , Muaradua, OKU Selatan, South Sumatera, 32212, Indonesia

11: Biodiversity Indicator Lab, Division of Habitats and Ecosystems, Endemic Species Research Institute, 1 Minsheng East Road, Jiji Town, Nantou County, Taiwan, 552005, Taiwan

12: Australian Antarctic Division, Science Branch, Department of Climate Change, Energy, Environment and Water, Channel Highway, Kingston, Tasmania, 7050, Australia

13: , , Scotland's Rural College, Craibstone Estate, Aberdeen, AB21 9YA, UK

14: Chair of Computational Landscape Ecology, Faculty of Environmental Sciences, Dresden University of Technology, Helmholtzstr. 10, Dresden, 01069, Germany

15: Agro-ecological Modeling Group, Faculty of Agriculture, University of Bonn, Niebuhrstr. 1a, Bonn, 53113, Germany

16: Colección de Sonidos Ambientales Mauricio Álvarez-Rebolledo - Colecciones Biológicas, Subdirección de Investigaciones, Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Claustro de San Agustín, Villa de Leyva, Boyacá, , Colombia

17: Quantitative Ecology Lab & Ordway Lab of Ecosystem Conservation, School of Natural Resources and Environment, Department of Wildlife Ecology and Conservation, & Florida Museum of Natural History, University of Florida, 715 Newell Dr, Building 866 & 1659 Museum Rd, Dickinson Hall 315, Gainesville, Florida, 32611, USA

18: , Department of Environmental Security, Faculty of Logistics and Crisis Management, Tomas Bata University, Studentské nám. 1532, Uherské Hradiště, , 68601, Czech Republic

19: Bioacoustics Group, Science Department, Biometrio.earth, Dr. Schönemann Str. 38, Saarbrücken, 66123, Germany

20: Wildlife Lab, Forest Research Institute, Hellenic Agricultural Organization - DIMITRA (ELGO-DIMITRA), Vasilika, Thessaloniki, TK 57006, Greece

21: , Department of Zoology and Animal Ecology, School of Biology, V.N.Karazin Kharkiv National University, 4 Svobody Square, Kharkiv, 61022, Ukraine

22: , Research Department, National Park 'Homilshanski Lisy', 27 Monastyrska Street, Koropove village, Kharkiv region, 63437, Ukraine

23: DYNAFOR, University of Toulouse, INRAE, Chemin de Borde-Rouge, Castanet-Tolosan, F-31326, France

24: Rainforest Connection, Science departament, , 440 Cobia Drive, Suite 1902, , TX, 77494, USA

25: Arbimon, , , , , , USA

26: BiBio Research Group, , Natural Sciences Museum of Granollers, Av. Francesc Macià 51, Granollers, Catalonia, 08402, Spain

27: Marine Biological Acoustic, Marine Sciences and Applied Biology, University of Alicante, C/ San Vicente del Raspeig s/n, San Vicente del Raspeig, Alicante, 3690, Spain

28: HIFI Lab, John Martinson Honors College, Purdue University West Lafayette Indiana, 1101 Third Street, West Lafayette, Indiana , 47906, USA

29: Tamar Dayan's Lab, School of zoology, Steinhardt Museum of Natural History, Tel-Aviv university, Klausner 12, Tel-Aviv, Tel-Aviv, 6997801, Israel

30: The Conservation Behavior Research Group, Mitrani Department of Desert Ecology, Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Midreshet Ben-Gurion, , 8499000, Israel

31: Forest Landscape Ecology Lab, Department of Forest Ecosystems and Society, Oregon State University, , , ,

32: , School of Geography and the Environment, Oxford University, South Parks Rd, Oxford, OX1 3QY, UK

33: Ocean Science Consulting Limited, , , Spott Road, Dunbar, EH42 1RR, United Kingdom

34: Equity in Ecological Research Lab, Department of Fish, Wildlife, and Conservation Biology, Colorado State University, 1474 Campus Delivery, Fort Collins, Colorado, 80521,

USA
35: Instituto de Ecorregiones Andinas (INECOA-UNJu-CONICET), Facultad de Ingeniería,
Universidad Nacional de Jujuy, Gorriti 237, San Salvador de Jujuy, Jujuy, 4600, Argentina
36: , Biological and Environmental Sciences, University of Stirling, Stirling, Stirlingshire, FK9
4LA, Scotland
37: , Centre for Conservation Science, RSPB, 2 Lochside View, Edinburgh Park,
Edinburgh, , EH12 9DH, UK
38: Centre for Conservation and Restoration Science, School of Applied Sciences,
Edinburgh Napier University, Sighthill Campus, Edinburgh, EH11 4BN, UK
39: Department of Behavioural Ecology, Faculty of Biology, Adam Mickiewicz University,
Uniwersytetu Poznańskiego 6, Poznań, 61614, Poznań
40: Laboratorio de Botanica Sistematica, Departamento de Botanica, Ecologia e Zoologia ,
Universidade Federal do Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio
Grande do Norte, 59078-900, Brazil
41: South Iceland Research Centre, , University of Iceland, Lindarbraut 4, Laugarvatn, 840,
Iceland
42: Sensola SLU Multisensory Outdoor Laboratory, Department of Landscape Architecture,
Planning and Management, Swedish University of Agricultural Sciences, SLU, Sundsvägen
4-6, Alnarp, 23053, Sweden
43: Institut de Systématique, Évolution, Biodiversité, , Muséum National d'Histoire naturelle,
Rue Buffon, Paris, 75005, France
44: , , Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, LECA, , , Grenoble,
38000, France
45: Laboratório de Comportamento Animal, Departamento de Zoologia, Universidade de
Brasília, Instituto de Ciências Biológicas, Distrito Federal, 70910-900, Brazil
46: , Pacific Northwest Research Station, U.S.D.A. Forest Service, 3625 93rd Ave. SW,
Olympia, Washington, 98512, U.S.A.
47: Wildlife Ecology & Management, Faculty of Agriculture, Yamagata University, 1-23
Wakabamachi, Tsuruoka, Yamagata, 9978555, Japan
48: , , National Museum of Natural Sciences, Spanish National Research Council, Calle José
Gutiérrez Abascal 2, Madrid, 28006, Spain
49: , Centre de Recherche sur la Biodiversité et l'Environnement (UMR 5300 CNRS-IRD-
TINPT-UPS), Université Paul Sabatier, 118 route de Narbonne, Toulouse, 31062 Toulouse
Cedex 9, France
50: Centre for Taxonomy and Morphology, Leibniz Institute for the Analysis of Biodiversity
Change, Museum of Nature Hamburg, Martin-Luther-King-Platz 3, Hamburg, 20146,
Germany
51: , Centre d'Ecologie et des Sciences de la Conservation (CESCO, UMR 7204), Museum
National d'Histoire Naturelle, Place de la Croix, Concarneau, Finistère, 29900, France
52: Laboratory of tropical ecology, , A.N. Severtsov Institute of Ecology and Evolution of
Russian Academy of Sciences, Leninsky Prospect, 33, Moscow, 119071, Russia
53: Behavioral and Community Ecology, Conservation Biology, College of Forestry, Guangxi
University, No. 100 Daxue Road, Nanning , Guangxi, 530004, People's Republic of China
54: School of Biological Sciences, Life Sciences Building, University of Bristol, 24 Tyndall
Ave, Bristol, BS8 1TQ, UK
55: Migratory Bird Unit, Northern Region, Canadian Wildlife Service, Environment and
Climate Change Canada, 5019 - 52nd Street, Yellowknife, Northwest Territories, X1A 2P7,
Canada
56: Species and Habitat Conservation, Forest Nature Conservation, Northwest German
Forest Research Institute, Prof.-Oelkers-Str. 6, Hann. Münden, Lower Saxony, 34346,
Germany
57: , , Wildlife Conservation Society Canada, 169 Titanium Way, Whitehorse, Yukon, Y1A
0E9, Canada
58: Bat Lab, Department of Evolutionary Ecology, Leibnitz Institute for Zoo and Wildlife
Research, Alfred-Kowalke-Straße 17, Berlin, 10315, Germany

59: , , Swiss Ornithological Institute, Seerose 1, Sempach, 6204, Switzerland

60: Zoological Research Museum Alexander Koenig, , Leibniz Institute for the Analysis of Biodiversity Change, Adenauerallee 160, NRW, 53113, Germany

61: Museum für Naturkunde, , Leibniz-Institut für Evolutions- und Biodiversitätsforschung, Invalidenstraße 43, Berlin, 10115, Germany

62: Ecology and Nature Management, Geosciences and Natural Resource Management, University of Copenhagen, Rolighedsvej 23, Frederiksberg, 1958,

63: Fundación de Conservación Jocotoco, Fundación de Conservación Jocotoco, Fundación de Conservación Jocotoco, Valladolid N24-414 y Luis Cordero, Quito, 170525, Ecuador

64: Landscape Conservation Programme, Conservation Division, BirdLife South Africa, , Gauteng, , South Africa

65: Afromontane Research Unit, Department of Zoology and Entomology, University of the Free State, , Phuthaditjhaba, , South Africa

66: Centre for Statistics in Ecology, the Environment and Conservation, Department of Statistical Sciences, University of Cape Town, , Cape Town, , South Africa

67: Nature Coast Biological Station, Institute of Food and Agricultural Sciences, University of Florida, 552 1st Street, Cedar Key, Florida, 32625, USA

68: Fisheries and Aquatic Sciences, Institute of Food and Agricultural Sciences, University of Florida, 136 Newins-Ziegler Hall, Florida, 32611, USA

69: Nature Conservation Unit, , Frederick University, 7, Yianni Frederickou Street, Nicosia, 1036, Cyprus

70: Madagascar Programme, , Chances for Nature, Brauweg 9a, Göttingen, 37073, Germany

71: , , Cologne Zoo, Riehler Straße 173, Köln, Lower Saxony, 50735, Germany

72: INAU Pantanal Biodata Center, National Institute of Science and Technology in Wetlands (INAU), Computational Bioacoustics Research Unit (CO.BRA), Federal University of Mato Grosso (UFMT), Cuiabá, 2367 Fernando Correa da Costa Ave, Mato Grosso, 78060-900, Brazil

73: Biodiversity, Macroecology and Biogeography, Faculty of Forest Sciences and Forest Ecology, University of Goettingen, Büsgenweg 1, Göttingen, Niedersachsen, 37077, Germany

74: Plant Ecology, Institute of Plant Sciences, University of Bern, Altenbergrain 21, Bern, Bern, 3013, Switzerland

75: Environmental Research and Innovation Centre, School of Science, Engineering and Environment, University of Salford, Peel Building, Salford, M5 4WT, UK

76: Chair of Geobotany, Chair of Geobotany, University of Freiburg, Schaenzlestr. 1, Freiburg im Breisgau, 79104, Germany

77: Department of Botany and Biodiversity Research, Faculty of Life Sciences, University of Vienna, Rennweg 14, Vienna, 1030, Austria

78: Polar Ecology Group, Department of Vertebrate Ecology and Zoology, University of Gdańsk, Poland, Wita Stwosza 59, Gdańsk, , 80-308, Poland

79: Westman Islands Research Centre, , University of Iceland, Ægisgata 2, Vestmannaeyjar, 900, Iceland

80: , Department of Ecology, University of Alicante, Cra. San Vicente del raspeig sn, Alicante, 3690, Spain

81: Department of Soil, Water, and Ecosystem Sciences, Institute of Food and Agricultural Sciences, University of Florida, 2181 McCarty Hall A, Florida, 32611, USA

82: , Department of Biology, University of Oxford, , Oxford, OX1 3SZ, United Kingdom

83: Integrative Community Ecology Unit, , Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Kunigami-gun, Okinawa, 904-0495, Japan

84: , Fisheries and Oceans Canada, Freshwater Institute, 501 University Crescent, Winnipeg, Manitoba, R3T 2N6, Canada

85: , Department of Ecology, Universidad Autónoma de Madrid, Darwin 2, Madrid, 28049, Spain

86: , Centro de Investigación en Biodiversidad y Cambio Global, Universidad Autónoma de
Madrid, Darwin 2, Madrid, 28049, Spain
87: , Wildlife Research Division, Environment and Climate Change Canada, 801-1550,
avenue d'Estimauville, Québec (Québec), G1J 0C3, CANADA
88: , Natural Resources & Environmental Studies, University of Northern British Columbia,
3333 University Way, British Columbia, V2N 4Z9, Canada
89: Department of Ecology and Evolutionary Biology & K. Lisa Yang Center for Conservation
Bioacoustics, , Cornell University, Corson Mudd, 215 Tower Road, Ithaca, New York, 14853,
USA
90: , Department of Biology, Whitman College, 345 Boyer AVe, Walla Walla, Washington,
99362, U.S.A.
91: , , Talarak Foundation, Inc., Negros Forest Park, Bacolod City, Negros Occidental, 6100,
Philippines
92: , Department of Forest Ecosystems and Society, Oregon State University, 3180 SW
Jefferson Way, Corvallis, Oregon, 97331, U.S.A.
93: , Department of Physical Geography, Stockholm University, SE-106 91 Stockholm,
Stockholm, 10691, Sweden
94: CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO
Laboratório Associado, Campus de Vairão, Universidade do Porto-InBIO, Universidade do
Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal
95: BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Univeristy of
Porto, Campus de Vairão, Rua Padre Armando Quintas, Porto, 4485-661 Vairão, Portugal
96: , Forest and Conservation Sciences, University of British Columbia, 2424 Main Mall,
Vancouver, British Columbia, V6T 1Z4, Canada
97: , Department of Biology, Mount Allison University, 63B York Street, Sackville, New
Brunswick, E4L 1E7, Canada
98: EcoAcoustic Research Hub (EAR Hub), Biosciences Center, Universidade Federal do
Rio Grande do Norte, Av. Sen. Salgado Filho 3000, Natal, Rio Grande do Norte, 59078-900,
Brazil

Abstract

The urgent need for remote, reliable, and scalable biodiversity monitoring amidst mounting
human pressures on ecosystems and changing climate has sparked interest in Passive
Acoustic Monitoring (PAM) worldwide. PAM holds potential for supporting global
sustainability goals by aiding conservation efforts, but so far, there is no comprehensive
overview of its coverage. Here we present metadata from 293 PAM datasets recorded since
1991 in the first global synthesis of ecoacoustic sampling coverage across spatial, temporal,
and ecological scales. We report data on sampling sites, deployment schedules, focal taxa,
and recording parameters, and quantify biological, anthropogenic, and geophysical
soundscape components across nine terrestrial and aquatic ecosystems. We found that
terrestrial sampling is spatially denser (33 sites/Mkm2) compared to aquatic realms, with
significant data gaps in subterranean realms. While diel and lunar cycles are well-sampled,
seasonal coverage is lacking in freshwater and terrestrial ecosystems, while 57% of marine
datasets cover all seasons. Opportunities for improvement include broader taxonomic
sampling on land, expanding coverage in the high seas, and increasing spatial replication in
freshwater environments. Additionally, we highlight nine case studies showcasing how PAM-
based soundscape ecology can contribute to macroecology, conservation, and phenology
studies, illustrating its potential to support global sustainability efforts both on land and
underwater.

Main text

Introduction

Sounds permeate all realms on Earth — terrestrial, freshwater, marine, and subterranean (Keith et al. 2022). In soundscape ecology (i.e., ecoacoustics), acoustic cues are used to study relationships between organisms and their environment (Pijanowski, Farina, et al. 2011; Sueur and Farina 2015). Much of this relies on passive acoustic monitoring (PAM) of soniferous organisms in water and on land. PAM can be used to track the three main soundscape components (Farina and Gage 2017; Pijanowski, Farina, et al. 2011): biophony (soniferous wildlife), anthropophony (human-related sounds), and geophony (sounds of geophysical origin) (Astaras et al. 2020; Doohan et al. 2019; Duarte et al. 2021; Kuşku et al. 2018; Pensieri et al. 2015; R. A. Rountree, Juanes, and Bolgan 2020; Scarpelli et al. 2020). Soundscape recordings can be used to non-invasively document the global state of soniferous life and the environment, and to derive metrics that measure its responses to transformations such as climate change, land use change, urbanisation, or restoration (Kang et al. 2023; Müller et al. 2023; Stowell and Sueur 2020; Sueur, Krause, and Farina 2019). PAM can thus inform conservation actions, biodiversity stewardship, protected area management (Astaras et al. 2020; Haver et al. 2019; Van Doren et al. 2023), and potentially soundscape-based policies. As such, it can support the COP15 biodiversity monitoring targets within the Global Biodiversity Framework and track progress towards UN Sustainable Development Goals.

Similar ecoacoustic concepts and methods are employed in aquatic and terrestrial systems, but their differences in application could enrich the disciplines of the respective realms. For instance, as sound travels fast and far in water, complex physics-based models of acoustic propagation were developed by marine engineers to account for great and variable detection ranges underwater. This underpins marine studies focused on signal detection, characterisation, localisation and density estimation. In contrast, terrestrial studies still mostly ignore these aspects of sound transmission (Kevin Felix Arno Darras et al. 2016; Haupert, Sèbe, and Sueur 2022), and population density estimation is in its infancy (Marques et al. 2013; Pérez-Granados and Traba 2021). In water, the identification of sound sources is challenging as sound sources cannot be intuitively identified and visual validation is rarely possible. As a result, many aquatic sound sources are unknown (Parsons et al. 2022), and while most fauna is expected to produce sound underwater, most animal sounds await documentation (Greenhalgh et al. 2020; Looby et al. 2023). This taxonomic haziness partly explains why, despite the terrestrial origins of the soundscape concept (Pijanowski, Farina, et al. 2011), marine ecologists pioneered studies assessing soundscape components (Halliday et al. 2020; R. A. Rountree, Juanes, and Bolgan 2020). On land, species-level automated identification is relatively advanced for some taxa (Nieto-Mora et al. 2023), and algorithms could potentially be re-purposed for aquatic species. Historically, underwater acoustics also benefited from technological developments in military passive surveillance and the need to evaluate and manage noise emissions in an environment that is not easily accessible to humans. It follows that aquatic standardisation efforts, especially with regard to noise level measurement (Haver et al. 2018), are relatively advanced (Abrahams, Desjonquères, and Greenhalgh 2021; ISO 2022; Roch et al. 2016), even though standards have not yet been reached (van Geel, Risch, and Wittich 2022). On land, calibrated noise measurements are performed in urban planning but rarely for soundscape ecology (Buxton

et al. 2017) although bioacoustic studies more frequently analyse noise impacts on wildlife
(Shannon et al. 2016). Correspondingly, terrestrial recording standards are still missing,
even though general recommendations have been given (Metcalf et al. 2023; Sugai et al.
2020).

The state of ecoacoustic research has only been reviewed separately for each realm for now
(Duarte et al. 2021; Greenhalgh et al. 2020; Havlik, Predragovic, and Duarte 2022; Lindseth
and Lobel 2018; Sugai et al. 2019), but cross-realm syntheses bear potential to answer new
questions. PAM can help to address fundamental questions in ecology (Ross et al. 2023).
We could investigate the general relationship between anthropophony and biophony (Lin et
al. 2023), but also specifically monitor impactful human activities (e.g., illegal hunting, bottom
trawling, habitat restoration) (Sethi et al. 2020) to model human impact on biodiversity and
ecosystem services, and to support law enforcement and conservation planning. One aspect
of this relationship is pervasive noise pollution effects on wildlife (Slabbekoorn et al. 2018),
that have been extensively studied in the marine and terrestrial realms (Boyd et al. 2015;
Buxton et al. 2017; Cox et al. 2018; Francis, Ortega, and Cruz 2009; Jerem and Mathews
2021; Solé et al. 2023) and increasingly so for freshwater (Mickle and Higgs 2018). We
could quantitatively assess the threshold at which noise levels become detrimental for
biological activity to design recommendations and designate "Quiet land, freshwater and
marine" areas for conservation (but see (Bayne, Habib, and Boutin 2008)). A global PAM
network could also support the identification of priority areas for conservation and
restoration, in line with the goals of the Kunming-Montreal Global Biodiversity Framework
(Deng 2023), while connecting with the wider public through publicly-available, audible
experiences and citizen science. Further, we could appraise the effects of the different
soundscape components on human health (Buxton et al. 2021; De Coensel, Sun, and
Botteldooren 2017; Van Renterghem 2019). Investigating macro-ecological theories related
to acoustic space competition, acoustic niche formation, and adaptation to noise is currently
possible, and the responses of biodiversity to global ecosystem assembly filters (topography,
latitude, water availability) (Keith et al. 2022) can be quantified. Despite the wide-ranging
and increasing soundscape sampling effort, the availability and distribution of datasets
remain undescribed, which makes it challenging to initiate large-scale syntheses. However,
the foundational role of large, inclusive data centralisation initiatives should not be
overlooked: community-initiated telemetry databases (Davidson et al. 2020; Hindell et al.
2020; Kays et al. 2022; Sequeira et al. 2018), individual animal observation networks (Culina
et al. 2021), and invasive species control syntheses (Spatz et al. 2022) spurred important
advances for entire research fields.

Here we assembled the first global PAM meta-database in the frame of the "WorldWide
SoundScapes" project and present a global synthesis of the state of ecoacoustic research.
We included standardized metadata from 294 passively recorded soundscape datasets with
spatial and/or temporal replication, considering only passively-recorded soundscape
datasets with spatial and/or temporal replication. The metadata identified the spatio-temporal
scope and coverage, ecosystems and focal taxa (IUCN GET - Global Ecosystem Typology
and IUCN Red list), protected status (WDPA - World Database on Protected Areas),
administrative status (IHO - International Hydrographic Organisation and GADM - Global
administrative Database (GADM 2022; Keith et al. 2022) and audio recording settings. We
used the meta-database to quantitatively assess a snapshot of the current state of sampling
efforts, and highlighted gaps and sampling biases. We identified opportunities to advance

the research agenda of soundscape ecology (Pijanowski, Villanueva-Rivera, et al. 2011) on a global scale, across realms. Lastly, to demonstrate the utility and potential of our database, we conducted a joint analysis of soundscape components (biophony, anthropophony, geophony) across nine ecosystems from terrestrial, marine, and freshwater realms, using sound recordings collated using the meta-database. We illustrate global soundscape patterns in relation to macroecology, conservation biology, and phenology. Although the database is inevitably biased by its contributing members, the project holds 258 contributors from around the world and represents the most comprehensive summary of PAM datasets across time and space to date. The resulting publicly accessible database continues to grow and remains open for metadata contributions that will facilitate future syntheses.

Results

Summary statistics

To date, 294 validated soundscape meta-datasets (hereafter "datasets") have been registered in our online database from across the globe, dating back to 1991 (Kevin F. A. Darras et al. 2024) (Fig. 1). A dataset was defined as a collection of metadata gathered by a single team on a coherent topic (generally corresponding to an individual study); subsequently added sites and yearly repetitions were coded as additional deployments within datasets. Metadata were validated by KD, YC, and FP by checking that all fields were filled and correctly formatted, and then cross-checked by their respective contributors who visualized site maps and recording timelines online on ecoSound-web (Kevin F. A. Darras et al. 2023) (https://ecosound-web.de/ecosound_web/collection/index/106). Based on the IUCN GET (Keith et al. 2022) definition of four 'core' realms, our database covers 215 terrestrial, 53 marine, 14 freshwater, and no subterranean, validated datasets. Further, sampling sites could be assigned to transitional realms, which sit at the interface between core realms. The majority of datasets were both spatially and temporally replicated for each core realm (Fig. 1). Few freshwater and terrestrial datasets had associated, openly-accessible recordings (15% for freshwater, 9% for terrestrial datasets) or peer-reviewed publications at the time of writing (21% for freshwater, 33% for terrestrial datasets) (Fig. S1), but marine datasets fared better (24% with available recordings, 58% published).

Spatial sampling extent, coverage, and density

The datasets contain 8189 terrestrial and 113 freshwater sites situated in 60 (out of 263) GADM level 0 administrative areas, and 254 marine sites situated in 28 (out of 101) IHO sea areas. Marine sites in particular were sourced from a variety of sea areas, as the bulk of sites (34) was not included in the top ten IHO areas, followed by the South Atlantic Ocean (32 sites) (Fig. 2). Thirty-two marine sites were situated in WDPA high-protection category Ia and II areas. However, our database lacked data from Arctic waters as well as from those off the coast of South Asian and Southeast Asian countries. Canada was the top contributor for sampling sites in the terrestrial realm (2996 sites), followed by 1829 terrestrial sites again sourced from a variety of regions outside the top ten (Fig. 2). 704 terrestrial sites were situated in WDPA category Ia, Ib, or II areas. Some areas had poor sampling coverage or missing data: the database currently does not include data from Russia, Greenland, and terrestrial Australia, as well as Saharan and several Central Asian countries. Spain was the top contributor for freshwater (20 sites), followed by the UK (16 sites). Four freshwater sites were situated in WDPA category II or Ib areas. Most GADM areas were not represented in the freshwater database subset.

The topographical analysis revealed that high-elevation mountains, polar caps, marine
tropical bathypelagic and benthic areas, and high-altitude freshwater bodies constituted gaps
in our database (Fig. 2). We did not receive contributions from sites above 4000 m in the
Northern Hemisphere or 2000 m in the Southern Hemisphere that are, to our knowledge,
currently acoustically unexplored. Terrestrial soundscape recordings from the northern polar
ice caps and the Transantarctic mountains were completely missing in our database. In
tropical marine areas, we compiled only scattered shallow-water soundscape datasets, and
benthic recordings were missing in the Southern Hemisphere. The Northern Hemisphere
was well represented in the distribution of terrestrial recordings (mean latitude across sites:
37°, Table S2).

Temporal sampling extent, coverage, and density

We analysed the diel, the seasonal cycle (only for temperate regions with four well-defined
seasons) and the lunar phase cycle, and split them into time windows (Fig. 3). The temporal
sampling coverage across time windows, measured in sampling years for the different solar
and lunar cycles, showed unequal contributions depending on the realm. Dawn and dusk
diel windows were shorter in duration than the day and night time windows, which reflects
the stronger representation of low latitude sites, where these intermediate time windows are
shorter. Daytime coverage was markedly higher than nighttime coverage in the terrestrial
realm (378 vs. 279 years), while dusk and dawn were similarly covered across all realms.
Terrestrial datasets had the lowest proportion of datasets that sampled all diel time windows
(73%). For temperate regions, seasons were defined as equally long windows, but their
coverage markedly differed across realms: in the marine realm, coverage was equal and
highest in winter and spring (both 44 years, together: 70% of coverage); in freshwater,
summer coverage was markedly higher (4.5 years — 38% of coverage); and in the terrestrial
realm, spring and summer were covered best (respectively 324 and 260 years, accounting
for 40% and 32% of coverage). Here, while 57% of the marine datasets completely covered
the seasonal cycle's time windows, only respectively 29% and 19% of the temperate
freshwater and terrestrial datasets sampled all seasons. The lunar phase cycle, being
categorised as a simple cycle with two time windows, was evenly-covered across realms
and comprehensively-covered within datasets.

Sampling in ecosystems

We quantified the extent, coverage, and density of the spatial and temporal acoustic
sampling effort across ecosystems (realms, biomes, functional types, as defined by IUCN
GET, Fig. 4). The terrestrial core realm has the third largest extent (239 Mkm^2), and spatial
sampling density was the highest across realms (33.52 sites per Mkm^2). Temporal sampling
density was relatively low (14% of the average 202 days' extent covered). The most
commonly-sampled biomes were temperate-boreal forests and woodlands (4855 sites),
tropical-subtropical forests (1503 sites), and the intensive land-use biome (1064 sites) (Table
S1). The marine core realm, being the most extensive (2028 Mkm^2), had the lowest spatial
sampling densities (0.11 sites per Mkm^2). However, it reached the highest temporal sampling
extent, coverage, and density (39% of the average 625 days' extent covered). The marine
shelf (107 sites) and pelagic ocean waters (68 sites) were the most-commonly sampled
marine biomes. The freshwater realm's spatial and temporal sampling densities were
intermediate (respectively 0.48 sites per Mkm^2 for an extent of 144 Mkm^2, and 70% of the
average 179 days' extent covered). Lakes (40 sites) and artificial wetlands (19 sites) were
the most commonly sampled biomes. Finally, to date there are no datasets that met our

inclusion criteria for the subterranean realm, even though it is second-largest in extent (541
Mkm^2). The terrestrial-freshwater realm was the best-studied transitional realm, and the
anthropogenic shorelines biome was the only non-subterranean biome that was not included
in our database. Only 24 out of the IUCN GET's 108 ecosystem functional types (i.e., 22%)
are not currently represented in our database (Table S1).

Taxonomic and spectral coverage
Distinct frequency ranges were covered for the different target taxa, while taxonomically
unspecific datasets tended to have a wider spectral coverage (i.e., average frequency range;
Fig. 4). Most aquatic datasets were taxonomically unspecific (78%) and some focused on
multiple taxa (13%), and both of them covered large frequency ranges extending below and
beyond the human-audible range (14 Hz — 26 kHz). All target taxa in the marine realm were
mammals, and the ones that did not target Carnivora rarely exceeded 1 kHz. Similarly,
freshwater datasets were mostly taxonomically untargeted (65%), covering a range from 32
Hz to 30 kHz, or focused on Actinopterygii (4 datasets). Terrestrial datasets were dominated
by bird-focused deployments (42%) from 82 Hz to 20 kHz, while taxonomically unspecific
datasets made up 27% of the total, covering a frequency range of 59 Hz to 23 kHz. Next,
datasets targeting multiple taxa (2 Hz — 45 kHz) and bat-focused datasets (6208 Hz — 119
kHz) respectively made up 13% and 9% of all datasets.

Soundscape analysis
In order to explore the soundscape patterns across different realms, we annotated the
biophony, geophony, and anthropophony in 116 recordings over nine ecosystem types (Fig.
5). We used the annotation-derived soundscape components to illustrate the potential of
soundscape data for studies about macroecology, conservation biology, and phenology. Our
analysis illustrates the contributions of various sound sources to the soundscape. On the
continental slopes off Sanriku (Japan), the acoustic space (i.e. the extent in frequency and
time) contained no detectable biological activity over the analysed evening time intervals, but
considerable human activity due to deep-sea mining and research vessel communication.
Photic coral reefs in Okinawa (Japan) had the highest biological activities due to snapping
shrimps and fish choruses, transient human activity due to passing vessels, and geophysical
activity due to breaking waves. In river estuaries in Western Australia, passing vessels,
dolphin whistles and snapping shrimp caused high levels of biophony and anthropophony,
while insects formed the bulk of the biophony in French large lowland river arms. Our Asian
soundscape samples from protected tropical rainforest areas were dominated by birds,
primates, and insects, while the montane tropical rainforests had less biological activity but
also included some human and geophysical sounds. The acoustic space of tree plantations
had a lower biological activity level than the tropical lowland rainforests and also included
some human activity from passing vehicles, while cities had an even higher anthropophony
and slightly lower biophony. Polar screes had very low levels of biophony, but high
geophony levels due to wind. Taken together, the soundscape components from the nine
ecosystems was summarised to illustrate the increase of biological activity near the equator
across large latitudinal gradients, the negative relationship between biological and
anthropogenic activity across aquatic and terrestrial systems, and the variable phenology of
soniferous organisms during the diel time windows (Fig. 5).

Discussion

The "WorldWide SoundScapes" project assembled the first meta-database describing soundscape datasets that cover the main realms and regions of the globe since 1991. While the database will remain open for contributions, we analysed its metadata to quantify the current state of PAM sampling extent, coverage, and density across spatio-temporal and ecological scales. We used its comprehensive information to quantify the sampling biases and gaps that may be representative of PAM on the entire globe. We further analysed 116 soundscapes across nine ecosystems to exemplify the potential use of the database for macroecology, conservation biology, and phenology studies across realms.

Previous quantitative syntheses were typically systematic reviews focusing on single realms (Greenhalgh et al. 2020; Havlik, Predragovic, and Duarte 2022; Sugai et al. 2019), freshwater syntheses were rather focused on bioacoustics, and the only cross-realm synthesis known to us was a qualitative review (Lindseth and Lobel 2018). However, recent case studies underline the potential of ecoacoustics for cross-realm syntheses: the discovery of universal distributions of sound and silence durations for freshwater, marine, and terrestrial realms (de Sousa et al. 2022) and the demonstration that soundscapes can track resilience and recovery of terrestrial and marine animals from natural disasters (Gottesman et al. 2021). Our database mainly includes yet unpublished datasets (Fig. S1), thus offering unique and novel insights into the state of ecoacoustic sampling. Spatio-temporal sampling effort was assessed using a unified framework, enabling quantitative evaluations of sampling extent, coverage, and density. More importantly, anyone can access our continuously growing, open database (CC-BY license) to source datasets and start collaborative ecoacoustic studies (https://ecosound-web.de/ecosound_web/collection/index/106) (Kevin F. A. Darras et al. 2024).

The frontiers of the current ecoacoustic spatial coverage are determined by removable technical and cultural barriers. On the one hand, our analysis reflects the technical difficulty in obtaining soundscape datasets at high latitudes and high elevations. Remoteness and extremely cold ambient temperatures present challenges for battery operation and replacement that can be solved with technical innovations. However, some marine deployments are able to reach the poles as water temperatures are buffered near the freezing point. On the other hand, large swathes of Russia, Central Asia, and Africa are still not represented in our database. This may reflect genuine disciplinary gaps as indicated by some of these regions' low bioacoustic coverage (Xeno-canto Foundation 2012) but may also result from cultural barriers (Amano, González-Varo, and Sutherland 2016) and limited hardware accessibility. Marine tropical waters are under-represented in our database, similarly to the IQOE marine networks' coverage (IQOE, n.d.). Indeed, particularly tropical countries may lack funding for setting up costly marine ecoacoustic programmes that require large vessels to reach high-sea areas, underwater vehicles or cabled stations on the seafloor (R. Rountree et al. 2020). By contrast, terrestrial monitoring is relatively inexpensive, and even tropical datasets are well-represented despite technical challenges under high humidity. Overall, while we currently miss some national programs in our database (e.g., Australian Acoustic Observatory (Roe et al. 2021)), we generally achieved a widespread distribution of terrestrial and marine datasets, and the northern hemisphere's temperate areas are well-covered. Affordable existing and future automated acoustic recorders deployed in the aquatic realm (Hill et al. 2019; Lamont et al. 2022) may close gaps in the limited spatial coverage of freshwater PAM and help to intensify spatial sampling of

marine areas. While these technical advances could make PAM more accessible in lower-
income countries, collaborative projects and equipment loans should promote the
establishment of soundscape research communities (Reboredo Segovia, Romano, and
Armsworth 2020). An increased collaboration with scientists and supporting citizen-science
projects in heavily underrepresented regions would thus improve not only the data fullness,
but also the representation and dialogue within the field. We recommend targeted site
selections to correct spatial biases and facilitate re-use of the data for global analyses (Beck
et al. 2014), and suggest that our geographic compilation allows the identification of high-
priority research areas.

Currently, only marine studies achieve a satisfactory and relatively even coverage of the
major solar and lunar cycles within their datasets, while other realms are limited by specific
technical constraints and taxonomic specialisation. Indeed, marine deployments offshore are
so expensive, especially in the deep sea (R. Rountree et al. 2020), that it makes little sense
to run them for limited durations. The ocean also buffers water temperatures so that despite
seasonal variations, animals will retain a basal activity throughout the seasons, even though
soundscapes may vary non-linearly and stochastically (Siddagangaiah et al. 2022).
Furthermore, lunar phases have a well-recognised effect on marine life such as zooplankton
and their associated predators (Hernández-León et al. 2001; Simonis et al. 2017; R. S.
Sousa-Lima and Clark 2008) that is usually accounted for. However, here we do not
consider more complex, ecosystem-specific cycles such as lunar tides. In the terrestrial
realm, we observe a marked bias towards avian monitoring during the day, which is
balanced by bat monitoring at night (Sugai et al. 2019). Often, terrestrial deployments have
shorter durations due to recorders cycling between sites (Sugai et al. 2020) or because
recorders can more easily be reached and are thus at risk of theft or vandalism. In temperate
regions, spring soundscapes are also over-represented in contrast to winter during which
birds have reduced activity, and when batteries may fail at low temperatures. We generally
don't have a good understanding of seasonal changes of vocal activities (even for birds in
the most studied regions). We encourage longer-duration setups to the limits of tenable
spatially representative designs: continuously-powered devices can increase temporal
coverage, but their complexity and cost (large batteries and/or solar panels) represents a
tradeoff against operating more recorders for greater spatial coverage. PAM offers a unique
opportunity here to conduct year-round sampling (Grinfeder et al. 2022) as seasonal cycles
determine activity cycles on land too (Krause, Gage, and Joo 2011; Pérez-Granados,
Schuchmann, and Marques 2020; Boullhesen et al. 2021), especially for migrant birds. The
lunar cycle may have a balanced coverage in terrestrial studies as its influence is
established for seabirds (Mougeot and Bretagnolle 2000), but it could also have been
randomly evenly sampled. Nonetheless, the cycle should be explicitly considered for more
taxa, especially for nocturnal or crepuscular ones which exhibit behavioral plasticity
depending on lunar illumination (Kronfeld-Schor et al. 2013; Appel et al. 2017; Lima et al.
2021; Ferreira et al. 2020; Schaaf, Boullhesen, and Akmentins 2023). Climate and land use
change will impact soundscapes in unpredictable ways through changing species
distributions and phenology, thereby justifying the need for better coverage across multiple
time scales — including inter-annual ones — to monitor changes in the ecosystems
(Desjonquères et al. 2022).

Re-use of soundscape datasets is sometimes restricted by their focus on certain target taxa,
which may hinder the analysis of other sound classes. For instance, studies focusing on

toothed whales or bats often need to trigger ultrasound recordings due to higher storage
demands, thereby resulting in species-focused, unrepresentative soundscape recordings.
Although there is no simple solution to this problem, we encourage researchers to record
audio files over frequency ranges that support potential re-use and interdisciplinary
collaboration (Lin and Tsao 2020). Technical limitations for making broadband recordings
are often slight, even though not all recorders offer high sampling rates and some
microphones have non-removable high-pass filters or low sensitivity at higher frequencies.
The downsides are higher power and memory consumption (R. Sousa-Lima 2013), as well
as potential post-processing, which altogether can make such endeavors prohibitively costly.
Analogously, re-use of data requires centralised soundscape repositories and species sound
libraries (e.g., Xeno-Canto, ChiroVox). New repositories should be created and existing ones
made interoperable (Pijanowski, Farina, et al. 2011; Parsons et al. 2022; Looby et al. 2022;
Kevin F. A. Darras et al. 2023). Fundamental bioacoustics continue to be needed to
establish the taxonomic base: While vocalizations of marine and freshwater mammals are
rather well studied, the sounds of most other aquatic wildlife still await proper documentation
(R. A. Rountree, Juanes, and Bolgan 2020). Although soundscape recording repositories
present tremendous data management challenges, established hybrid commercial solutions
are available to store recordings on low-cost, long-term archival media with temporary
transfers on fast hard disk space for regularly-accessed data (c.f. WildTrax
https://www.wildtrax.ca). Governmental and industrial marine projects from the oil and gas
industry also routinely upload data, sometimes as part of mitigation efforts. However,
recordings may be restricted to low frequencies (Haver et al. 2018), or be access-limited. We
encourage planning soundscape data archival at an early stage to safeguard data in the
long term and to enable public availability within ethical and legal boundaries.

New frontiers in soundscape research appear upon close scrutiny of the sampled ecosystem
types and their soundscapes. To our knowledge, all core and transitional subterranean
realms, such as aerobic caves, anthropogenic voids, underground streams or sea caves,
await acoustic exploration. Although the subterranean realm is the second-largest, access is
usually challenging or restricted. One may expect that subterranean soundscapes comprise
substrate-borne geophony and anthropophony, while the biophony of aerobic caves would
comprise social calls of cave-dwelling birds and bats. Notably, freshwater ecoacoustic
studies were also too rare to draw robust conclusions from. They suffer from a lack of spatio-
temporal replication arising from opportunistic sampling: several studies could not be
included in our database for this very reason. Dynamic water bodies (seasonal,
episodic/ephemeral, freeze/thaw, and rice paddies) are not yet well-studied (Messager et al.
2021). This underlines that the freshwater realm is relatively accessible but poorly
represented, even though that realm also comprises above-water, aerial recordings. We
expect geophony to increase with elevation in moving water, prevalent anthropogenic noise
in large flowing lowland water bodies, and the underwater biophony to come mainly from
insects and fish (R. A. Rountree, Juanes, and Bolgan 2020). Luckily the freshwater
community grows rapidly (Hill et al. 2019; Linke et al. 2018).

The relatively better-explored marine and terrestrial realms also leave plenty of opportunities
for acoustic exploration. Marine research is dominated by the omnipresent impact of noise
on wildlife (Duarte et al. 2021), as borne out by the high prevalence of anthropophony in our
case study's selected soundscapes with deep-sea mining sound and vessel noise in coral
reefs. At the same time, our reef soundscapes exhibited the highest biophony occupancy in

snapping shrimp aggregations. Importantly, sound in the ocean propagates so far that
multiple biomes can be sampled at once, i.e., recorders on the seafloor will also record the
soundscape of the pelagic waters above. Still, our database suggests coverage gaps exist in
marine sessile animal and algae habitats (kelp, shellfish, seagrass, animal forests). On land,
the terrestrial realm is biased towards sampling forests, which in our case study exhibited a
high biophony and low geophony, with more intensively-managed systems such as
plantations or cities having increasing anthropophony levels. In our case study soundscapes,
biophony declined with elevation and almost disappeared in less hospitable high-latitude
areas as well as arid habitats such as polar screes and deserts. Our database coverage
suggested that the most challenging terrestrial functional groups were yet unsampled (lava
flows, polar tundra, and sclerophyll hot deserts). Finally, as most datasets focused on single
realms, transitional realms were also under-represented: these habitats are more
challenging to assess, but by being at the interface between realms, potentially very dynamic
and diverse. To add to this complexity, soundscapes can be recorded in aquatic realms
above the water surface — and in terrestrial systems, below the ground (Maeder et al. 2022;
Inyang et al. 2019). The interactions between realms, at the boundary between recording
media, are rarely considered but can be captured in holo-soundscapes (R. A. Rountree,
Juanes, and Bolgan 2020; Murchy, K.A., Rountree, R.A., and Juanes, F., n.d.).

A unified, interdisciplinary approach to soundscape science is now possible. We need to
come up with a consistent, broad-scale monitoring of the Earth soundscape in a period of
major population increase and climate change which has the potential to cause rapid
changes in biodiversity and natural systems. Sound can be used across the globe to
measure fundamental qualities of the environment (Ross et al. 2023). Although more
comprehensive coverage is needed to answer research questions decisively, our database
grows as it remains open for future dataset inclusions. We encourage prospective
contributors to archive their metadata sustainably and to join our inclusive project. We
showed that a large portion of the ecoacoustic community is willing to publicly share their
work across realm-specific disciplines and to collaborate. We advocate a bolder ecoacoustic
research agenda, reaching out to challenging places where no sound has been recorded
before as well as to populated places — while accounting for data privacy issues.
Ecoacoustic monitoring should be integrated into long-term ecosystem monitoring platforms
(i.e., eLTER (Mirtl 2018)) and come to aid to support the necessary evaluation of progress
towards the ambitious goals and targets of the newly established Kunming-Montreal Global
Biodiversity Framework. Ideally, it should be included in the future resolutions of the next
Convention on Biological Diversity's Conference Of Parties. Soundscape recording
standardisation should be initiated and general soundscape repositories be created to
sustainably archive data like "time capsules" (Sugai and Llusia 2019) and to make it
available for citizen science. Together with more taxonomically inclusive recording practices,
we would obtain a treasure of general-purpose datasets that maximise potential re-use and
facilitate comparability for large-scale syntheses. Further, international collaborations should
accelerate plugging the remaining coverage gaps, and future studies should be framed
within a broader context by collaborating with a variety of disciplinary experts across socio-
economic groups. By building collaborations around the research frontiers identified here, we
can aim to comprehensively describe and understand the acoustic make-up of the planet.

Methods

Ethics & Inclusion statement

The present study has involved all local researchers who carried out the studies forming the basis for the database. Local researchers were the respective study leads — usually corresponding authors for published studies, and referred contacts for unpublished studies — who were asked to identify further primary and secondary contributors of their study. Primary contributors were invited as co-authors, and secondary contributors are acknowledged at the end of the main text. As such, the research was locally relevant as it had already been initiated and performed by the primary contributors. The responsibility of primary contributors regarding their metadata and their implication as co-authors were explicitly agreed in a collaboration agreement. Primary contributors all fulfil data curation and manuscript revision roles, and were invited to participate more actively. Selected contributors were invited as co-leads to be responsible for particular realms or biomes, and are all listed in the first-tier authors list. All contributors were asked to identify further contacts to reach a comprehensive coverage for the database.

By the nature of the study (using passive acoustic monitoring methods), the research did not suffer from restrictions or prohibitions in the setting of the researchers. Necessary permits from landowners or environmental protection agencies, precautions to limit biological contaminations (biofilm on marine recorders) or for animal welfare, and local ethics committee reviews were the responsibility of the respective dataholders. None of the passive acoustic monitoring activities resulted in personal risk to participants; neither did they involve health, safety, security or other risk to researchers; nor did they require benefit-sharing measures as nothing was transferred out of any country. Local and regional research relevant to our study were taken into account in citations throughout the main text.

Database construction

The database was constructed within the frame of the Worldwide Soundscapes project (Kevin F. A. Darras et al. 2024) using a collaborative, peer-driven metadata collation. It represents the current state of publicly accessible meta-datasets. We started contacting contributors that we personally know, and designating co-lead authors who supported contacting potential contributors in the respective ecosystem types of their expertise (i.e. terrestrial, urban, freshwater, and marine). We posted the call for contributors on specialised ecoacoustics platforms and social media and constantly keep the project open for any contributor owning suitable soundscape recordings beyond the present publication. We communicated mostly in English but also in Spanish, French, Portuguese, German, and Chinese to gather metadata; we tracked metadata completion and cross-validated them with graphical map and timeline checks by the contributors. As a result, apart from basic coherence checks conducted by the co-leads and research assistants, the responsibility of the database content is borne by their individual contributors, who were all invited as co-authors. The database information page is integrated on our online collaborative ecoacoustics platform ecoSound-web (Kevin Felix Arno Darras et al. 2020). It already hosts selected soundscape recordings for the present soundscape component analysis (https://ecosound-web.de/ecosound_web/collection/show/49).

At the time of writing, our database achieves a global coverage of soundscape datasets. Its scope is limited to datasets accessible over the internet, through email contact with their creators and through publications recommended by peers. Out of 420 contacted contributors potentially involved in PAM, 259 (62%) provided metadata, but the database remains open for future contributions. More datasets exist but are not accessible despite valid contact details, as reflected by the 23% percentage of contacts that have not yet responded. This may correspond to peers with limited availability to share research data. There may be further datasets that could not be obtained yet due to no longer reachable contacts. Older soundscape datasets may be stored on analog media that are inaccessible over the internet, but these are more often attended recordings (e.g. (Krause, Gage, and Joo 2011)) — thus not fulfilling our inclusion criteria (see below). The missing share of datasets could be estimated with a systematic literature search, but we posit that our dataset search was exhaustive enough and importantly, able to find gaps more effectively due to focused calls for missing datasets in our large group of contributors. Moreover, our database is open for new contributions and constantly growing to be more comprehensive.

We made sure that datasets have as little bias as possible and are clearly representative of spatio-temporal patterns. We only considered soundscape recording datasets that are: 1) stationary — mobile recorders would have constantly changing and shifting acoustic detection spaces, we excluded recordings from cars, transect walks, or towed deployments; 2) passive — obtained from automated, unattended recorders, without human presence near the acoustic sensor; 3) ambient — with no particular recording direction, as obtained from omnidirectional microphones; 4) from spatially or temporally replicated study designs (Fig. 1). Datasets with spatial and temporal replication are necessary to disentangle spatial and temporal effects from other soundscape determinants: we considered that datasets were spatially replicated when several sites were sampled simultaneously, and temporally replicated when one site was sampled over multiple days at the same daytime. We chose sampling sites and days as the most elemental units for defining replication in our unified analysis. However, we acknowledge that replicates are defined differently in each study: sampling sites could only be spatial replicates if they belong to the same category (i.e. habitat, management type). Temporal replicates could also be defined at the scale of other solar and lunar cycles (i.e. multiple full moons). Taken together, our requirements homogenize the dataset to enable statistical analyses across datasets and future collaborative syntheses.

Time and space

Soundscapes are fundamentally determined by their time and location. Wildlife and human activity is broadly determined by events in the solar and lunar cycles (Frey et al. 2017; Gottesman et al. 2021; Grinfeder et al. 2022; Kronfeld-Schor et al. 2013), as well as geographical positions on the planet relative to the poles, equator, or the land and water surface. For both space and time, we determined the extent (area of realms or time range), coverage (number of sites or number of days per site), and density (sites per area or recorded time per time range) of the respective units under consideration (datasets, sites, deployments). Notably, the true, areal coverage of sites is unknown in the vast majority of cases and generally larger in marine environments than terrestrial ones. Our measure of spatial sampling density using points per area is thus a provisional one.

For each sampling site, we gathered numerical latitude and longitude and topography
(above and below the sea level), and further assigned them to administrative areas: GADM
(Global administrative areas database) divisions for freshwater and land, and IHO
(International Hydrographic Organisation) for sea areas. We also extracted the WDPA
protected area category each sampling site belonged to. We use these metadata to quantify
the latitudinal and topographical sampling site coverage across realms.

Contributors were asked to code the timing of their deployments using start and end dates
and times, operation modes (continuous, periodical, scheduled), schedules if applicable
(daily starting times and end times or durations) as well as duty cycles if applicable.
Deployments were assigned to single sampling sites. Temporal information was inferred
from the timing of sound recordings relative to time events that structure life and
meteorological events on Earth (Fig. 3): the yearly cycle was split into four seasons
approximately centered on solstices and equinoxes (only temperate regions); the daily cycle
was split into four windows delimiting dawn (from astronomical dawn start at -18° solar
altitude until 18° solar altitude), day, dusk (from 18° solar altitude to astronomical dusk end
at -18° solar altitude), and night; the lunar illumination cycle was split into two time windows
centered around the full and new moon phases. It follows that the extrema and ecotones in
the temporal cycles define the relevant sampling time windows, noting that in temperate
zones, the equinoxes roughly correspond to thermal ecotones. However, seasonal cycles in
tropical, subtropical, and arid regions that are due to precipitation variations were not
considered in this analysis. The resulting temporal framework was used to quantify sampling
coverage in these three solar and lunar cycles.

Ecology and frequency range

We assigned individual sampling locations to ecosystem types such as realms (core and
transitional ones), biomes, and functional groups as determined by the IUCN GET
(https://global-ecosystems.org), and calculated the major and minor occurrence areas of all
functional groups (Keith et al. 2022; Keith, Ferrer-Paris, Nicholson, Bishop, et al. 2020;
Keith, Ferrer-Paris, Nicholson, and Kingsford 2020). We used these metadata to quantify
spatio-temporal extent, coverage, and sampling density within ecosystem types.
Deployments were also linked to their main studied IUCN taxa (class, order, family, or
genus) when applicable — some datasets were collected without taxonomic focus. We use
these data to depict the sampled frequency ranges depending on the target taxon.

Sound recordings store a representation of the original soundscape. Microphones have
variable frequency responses, and the spectral scope of the recording is limited by the
sampling frequency of the recorder (nowadays, overwhelmingly digital). Contributors were
asked to provide the audio parameters of their deployments (sampling rate, high-pass filter)
as well as auxiliary metadata about the microphones' and recorders' makes and brands. The
frequency range is additionally determined by microphone roll-off frequencies at the low end
and their frequency response at high frequencies, but we could not gather these detailed,
often unpublished data for the high number of equipment types used here. Together, the
available metadata allowed us to determine the spectral scope of the soundscape recording
datasets and to potentially correct for variable detection space areas.

Soundscape component case studies

To illustrate how the database can be used for macroecology, conservation biology, and phenology analyses, we chose to acoustically sample a selection of ecosystems across a variety of topographical, humid, latitudinal, and anthropisation conditions — all fundamental gradients of assembly filters in both terrestrial and marine realms (Keith et al. 2022). We aimed to select datasets containing at least four spatial replicates within the same functional group, with recordings spanning all four diel time windows from the same date. Our sampling sites spanned -68 to 39 degrees of numerical latitude and were recorded during the Northern Hemisphere's summer. The recordings covered the audible frequency range with sampling rates of 44.1 kHz in mono or stereo. We extracted 10-minute recordings starting at sunrise, solar noon, sunset, and solar midnight from the chosen datasets and sampling sites. Our selection resulted in 16 recordings for each of the functional groups (tropical-subtropical forests biome: tropical/subtropical lowland rainforests, tropical/subtropical montane rainforests; intensive land use biome: plantations). We could not find temporally replicated recordings for the deepest marine functional group (four solar noon recordings from continental and island slopes — deep sea floors biome), and only three spatial replicates for the shallowest marine functional group (12 recordings from photic coral reefs — marine shelf biome). Overall, we obtained 116 recordings.

Each recording's spectrogram (i.e., a visual representation of sound intensity along the time and frequency axes) was annotated with the three fundamental soundscape components: biophony, anthropophony, and geophony. KD resampled recordings to 44.1 kHz when higher sampling frequencies were used, and uploaded them on ecoSound-web (https://ecosound-web.de/ecosound_web/collection/show/49) (Kevin F. A. Darras et al. 2023), where all recordings were listened to while visually inspecting the spectrograms (FFT window size of 1024) at 15 pixels per second, and creating annotations at a visual density of 1116 pixels per 10 minutes on the left channel. Sounds caused by microphone or recorder self-noise were not annotated. Annotations were rectangular tags on the spectrogram, encompassing only the annotated sound, with defined coordinates in the time and frequency dimensions. Annotations of different soundscape components could overlap if they were simultaneously visible or audible. Annotations of the same soundscape component were adjacent and non-overlapping to avoid double-counting. All annotations were reviewed by the contributors who created the recordings, using the peer-review mode on ecoSound-web, and only accepted ones were used in the analysis. The annotations were then exported and acoustic space occupancy for each soundscape component was calculated as the proportion of the spectro-temporal space (duration multiplied by frequency range and divided by total area of spectrogram, range: 0-1. Similar to (Luypaert et al. 2022)).

838 # Display items

839 Table 1: An agenda for enabling global soundscape ecology research.

Research gap or bias	Solution
Poor global coverage, acceptance, and standardisation	- Build general data repositories - Invest in low-income country cooperation, capacity building, equitable funding access - Disseminate results in local languages - Get explicit agreements for data collection
Poor spatial coverage in marine realm	- Cheaper recorders for coastal recordings (Hydromoth) - More collaboration for equipment sharing and vessel use for high seas (World Ocean Day)
Null coverage in subterranean realm	- Collaboration with speleologists, chiropterologists, urban ecologists
Biased spatial coverage in terrestrial realm	- Deploy more recorders in extreme and dynamic biomes
Biased temporal coverage and selection in terrestrial realm	- Cover all seasons and whole diel - Include geophony and anthropophony periods into analysis
Biased taxonomic coverage in terrestrial realm	- Avoid taxonomic focus by covering wide frequency range: disable high-pass filters, use high sampling rates
Missing standardisation in terrestrial realms	- Calibrate equipment - Measure detection spaces - Estimate animal densities
Poor coverage in freshwater realm	- Increase collaboration with scientists from other realms - Record soundscapes linking terrestrial and freshwater realm
Poor coverage of transitional realms	- More dynamic/time-series analyses (Gottesman et al. 2021) to acknowledge temporal and spatial dynamics of complex systems
Differing workflows across realms	- Adopt common standards and increase interoperability of software tools to foster interdisciplinarity and integration into biodiversity and remote sensing databases
Unknown geophony-biophony interactions	- Collaboration with meteorological networks to include acoustic monitoring

Acknowledgements

We are thankful to our colleagues who supported the work underlying the database creation: Abigail Seybert, Adrien Charbonneau, Alexandra Buitrago-Cardona, Alfonso Zúñiga, Allan G Oliveira, Ambroise N. Zongo, Annalea Beard, Anthony Tan, Australian Institute of Marine Science, Base de Estudos do Pantanal - UFMS, Bea Maas, Bibiana Gómez-Valencia, Borja Milá, Carolline Z Fieker, Chistophe Thébaud, Chong Chen, Claudia A. Medina-Uribe, Colin KC Wen, Conner Partaker, Curtin University, Cécile Albert, Daisy Dent, Dan Warren, Dedi Rahman, Dexter Hodder, Diego Alejandro Gómez-Montes, Dustin Whalen, ECO²SCAPE Project Group, Elizabeth Clingham, Emilia Sokolowska, Emilio Tan, En-Hao Liu, Eric Rakotomalala, Estância Mimosa Ecoturismo, Evan Economo, Fang-Yee Lin, Filipe F de Deus, Francis Juanes, Frederic Sinniger, Greg and Lisa Morgan, Guillermo McPherson, Hannah Reyes, Hanneline Smit-Robinson, Harris Papadopoulos, Hervé Jourdan, Hiromi Kayama Watanabe, Holger Kreft, Ilias Karmiris, Ingo Grass, Italo A A Rondon, Jacques Keumo Kuenbou, Jakob Tougaard, Janice Ser Huay Lee, Jeff Robinson, Johan Diepstraten, John Hildebrand, Jose Manuel Ochoa-Quintero, José Luis Mena, José Wagner, Juan Diego Tovar, Justine Magbanua, Jérémy Anso, Jérôme Sueur, Karen H. Beard, Katie Huong, Ken Otter, Kenneth Dudley, Kinga Buda, Laela Sayigh, Laure Desutter-Grandcolas, Leeann Henry, Lenka Brousset, Lisa Loseto, Luca Iacobucci, Marcos Vaira, Marlize Muller, Martim Melo, Mauricio Akmentins, Michael Scherer-Lorenzen, Nicholas Brown, Nick Gardner, Nikos Fakotakis, Noam Leader, Nuno Faria, Peter Taylor, Pousada Xaraés, Rainforest Connection (RFCx), Refúgio Ecológico Caiman, Refúgio da Ilha Ecolodge, Rob Rempel, Robert Spaan, Rouvah Andriafanomezantsoa, Saki Harii, Sandra Mueller, Saskia Dröge, Shannon MacPhee, Shinsuke Kawagucci, Sierra Hunicutt, Sigal Balshine, Silent cities consortium, Sina Weier, Stephen Insley, Tetsuya Miwa, Thiago M Ventura, Thiago Pinheiro, Todor Ganchev, Tomasz Stanislaw Osiejuk, Tomonari Akamatsu, Valentin Chevalier, Vy Tram Nguyen, Wen-Ling Tsai, Wigna da Silva, Yenifer Herrera-Varón, Yuhang Song, Zehava Sigal, Zuania Colón-Piñero, Zuzana Burivalova. The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Government of any product or service. The findings and conclusions in this publication are those of the authors and should not be construed to represent any official U.S. Government determination or policy. We acknowledge the NFDI Consortium Earth System Sciences — NFDI4Earth, coordinated by TU Dresden, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — project number: 460036893. BIOMON is funded by the European Union's Horizon Europe programme under grant agreement 101090273.

Competing interests

The authors declare to have no competing interests. All authors have seen and approved the manuscript; it hasn't been accepted or published elsewhere.

References

Abrahams, Carlos, Camille Desjonquères, and Jack Greenhalgh. 2021. "Pond Acoustic Sampling Scheme: A Draft Protocol for Rapid Acoustic Data Collection in Small Waterbodies." *Ecology and Evolution* 11 (12): 7532–43. https://doi.org/10.1002/ece3.7585.

Amano, Tatsuya, Juan P. González-Varo, and William J. Sutherland. 2016. "Languages Are Still a Major Barrier to Global Science." *PLOS Biology* 14 (12): e2000933. https://doi.org/10.1371/journal.pbio.2000933.

Appel, Giulliana, Adrià López-Baucells, William Ernest Magnusson, and Paulo Estefano D. Bobrowiec. 2017. "Aerial Insectivorous Bat Activity in Relation to Moonlight Intensity." *Mammalian Biology* 85 (July): 37–46. https://doi.org/10.1016/j.mambio.2016.11.005.

Astaras, Christos, Joshua M. Linder, Peter Wrege, Robinson Orume, Paul J. Johnson, and David W. Macdonald. 2020. "Boots on the Ground: The Role of Passive Acoustic Monitoring in Evaluating Anti-Poaching Patrols." *Environmental Conservation* 47 (3): 213–16. https://doi.org/10.1017/S0376892920000193.

Bayne, Erin M., Lucas Habib, and Stan Boutin. 2008. "Impacts of Chronic Anthropogenic Noise from Energy-Sector Activity on Abundance of Songbirds in the Boreal Forest." *Conservation Biology: The Journal of the Society for Conservation Biology* 22 (5): 1186–93. https://doi.org/10.1111/j.1523-1739.2008.00973.x.

Beck, Jan, Marianne Böller, Andreas Erhardt, and Wolfgang Schwanghart. 2014. "Spatial Bias in the GBIF Database and Its Effect on Modeling Species' Geographic Distributions." *Ecological Informatics* 19 (January): 10–15. https://doi.org/10.1016/j.ecoinf.2013.11.002.

Boullhesen, M., M. Vaira, R. M. Barquez, and M. S. Akmentins. 2021. "Evaluating the Efficacy of Visual Encounter and Automated Acoustic Survey Methods in Anuran Assemblages of the Yungas Andean Forests of Argentina." *Ecological Indicators* 127 (August): 107750. https://doi.org/10.1016/j.ecolind.2021.107750.

Boyd, Ian, George Frisk, Ed Urban, Peter Tyack, Jesse Ausubel, Sophie Seeyave, Doug Cato, et al. 2015. "An International Quiet Ocean Experiment." *Oceanography* 24 (2): 174–81. https://doi.org/10.5670/oceanog.2011.37.

Buxton, Rachel T., Megan F. McKenna, Daniel Mennitt, Kurt Fristrup, Kevin Crooks, Lisa Angeloni, and George Wittemyer. 2017. "Noise Pollution Is Pervasive in U.S. Protected Areas." *Science* 356 (6337): 531–33. https://doi.org/10.1126/science.aah4783.

Buxton, Rachel T., Amber L. Pearson, Claudia Allou, Kurt Fristrup, and George Wittemyer. 2021. "A Synthesis of Health Benefits of Natural Sounds and Their Distribution in National Parks." *Proceedings of the National Academy of Sciences* 118 (14). https://doi.org/10.1073/pnas.2013097118.

Cox, Kieran, Lawrence P. Brennan, Travis G. Gerwing, Sarah E. Dudas, and Francis Juanes. 2018. "Sound the Alarm: A Meta-Analysis on the Effect of Aquatic Noise on Fish Behavior and Physiology." *Global Change Biology* 24 (7): 3105–16. https://doi.org/10.1111/gcb.14106.

Culina, Antica, Frank Adriaensen, Liam D. Bailey, Malcolm D. Burgess, Anne Charmantier, Ella F. Cole, Tapio Eeva, et al. 2021. "Connecting the Data Landscape of Long-Term Ecological Studies: The SPI-Birds Data Hub." *Journal of Animal Ecology* 90 (9): 2147–60. https://doi.org/10.1111/1365-2656.13388.

Darras, Kevin F. A., Noemí Pérez, Mauladi -, Liu Dilong, Tara Hanf-Dressler, Matthias Markolf, and Thomas C. Wanger. 2023. "ecoSound-Web: An Open-Source, Online Platform for Ecoacoustics." *F1000Research*. https://doi.org/10.12688/f1000research.26369.2.

Darras, Kevin F. A., Rodney Rountree, Steven Van Wilgenburg, Amandine Gasc, Songhai Li, Lijun Dong, Yuhang Song, Youfang Chen, and Thomas Cherico Wanger. 2024. "Worldwide Soundscapes Project Meta-Data." Zenodo.

934 https://doi.org/10.5281/zenodo.10598949.

Darras, Kevin Felix Arno, Noemí Pérez, Mauladi -, and Tara Hanf-Dressler. 2020.
 "BioSounds: An Open-Source, Online Platform for Ecoacoustics." *F1000Research* 9
 (October): 1224. https://doi.org/10.12688/f1000research.26369.1.

Darras, Kevin Felix Arno, Peter Pütz, Fahrurrozi, Katja Rembold, and Teja Tscharntke. 2016.
 "Measuring Sound Detection Spaces for Acoustic Animal Sampling and Monitoring."
 Biological Conservation 201 (September): 29–37.
 https://doi.org/10.1578/AM.39.1.2013.23.

Davidson, Sarah C., Gil Bohrer, Eliezer Gurarie, Scott LaPoint, Peter J. Mahoney, Natalie T.
 Boelman, Jan U. H. Eitel, et al. 2020. "Ecological Insights from Three Decades of
 Animal Movement Tracking across a Changing Arctic." *Science* 370 (6517): 712–15.
 https://doi.org/10.1126/science.abb7080.

De Coensel, Bert, Kang Sun, and Dick Botteldooren. 2017. "Urban Soundscapes of the
 World: Selection and Reproduction of Urban Acoustic Environments with
 Soundscape in Mind." *INTER-NOISE and NOISE-CON Congress and Conference
 Proceedings* 255 (2): 5407–13.

Deng, Iris. 2023. "Harnessing the Power of Sound and AI to Track Global Biodiversity
 Framework (GBF) Targets."

Desjonquères, Camille, Sara Villén-Pérez, Paulo De Marco, Rafael Márquez, Juan F.
 Beltrán, and Diego Llusia. 2022. "Acoustic Species Distribution Models (aSDMs): A
 Framework to Forecast Shifts in Calling Behaviour under Climate Change." *Methods
 in Ecology and Evolution* 13 (10): 2275–88. https://doi.org/10.1111/2041-
 210X.13923.

Doohan, B., S. Fuller, S. Parsons, and E. E. Peterson. 2019. "The Sound of Management:
 Acoustic Monitoring for Agricultural Industries." *Ecological Indicators* 96 (January):
 739–46. https://doi.org/10.1016/j.ecolind.2018.09.029.

Duarte, Carlos M., Lucille Chapuis, Shaun P. Collin, Daniel P. Costa, Reny P. Devassy,
 Victor M. Eguiluz, Christine Erbe, et al. 2021. "The Soundscape of the Anthropocene
 Ocean." *Science* 371 (6529). https://doi.org/10.1126/science.aba4658.

Farina, Almo, and Stuart H. Gage. 2017. "Ecoacoustics: A New Science." In *Ecoacoustics*,
 1–11. John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119230724.ch1.

Ferreira, Luane S., Luciana H.S. Rocha, Danielly Duarte, Edvaldo Neto, Júlio E.
 Baumgarten, Flávio H.G. Rodrigues, and Renata S. Sousa-Lima. 2020. "Temporal
 and spatial patterns of the long-range calls of maned wolves (Chrysocyon
 brachyurus)." *Mastozoología Neotropical* 27 (1): 081–095.
 https://doi.org/10.31687/saremMN.20.27.1.0.24.

Francis, Clinton D., Catherine P. Ortega, and Alexander Cruz. 2009. "Noise Pollution
 Changes Avian Communities and Species Interactions." *Current Biology* 19 (16):
 1415–19. https://doi.org/10.1016/j.cub.2009.06.052.

Frey, Sandra, Jason T. Fisher, A. Cole Burton, and John P. Volpe. 2017. "Investigating
 Animal Activity Patterns and Temporal Niche Partitioning Using Camera-Trap Data:
 Challenges and Opportunities." *Remote Sensing in Ecology and Conservation* 3 (3):
 123–32. https://doi.org/10.1002/rse2.60.

GADM. 2022. "Global Administrative Areas." GADM database of Global Administrative
 Areas. http://www.gadm.org/home.

GEBCO. n.d. "Gridded Bathymetry Data (General Bathymetric Chart of the Oceans)."
 GEBCO. Accessed August 27, 2022.
 https://www.gebco.net/data_and_products/gridded_bathymetry_data/.

Geel, Nienke C. F. van, Denise Risch, and Anja Wittich. 2022. "A Brief Overview of Current
 Approaches for Underwater Sound Analysis and Reporting." *Marine Pollution Bulletin*
 178 (May): 113610. https://doi.org/10.1016/j.marpolbul.2022.113610.

Gottesman, Benjamin L., Jack C. Olson, Soohyun Yang, Orlando Acevedo-Charry, Dante
 Francomano, Felix A. Martinez, Richard S. Appeldoorn, Doran M. Mason, Ernesto
 Weil, and Bryan C. Pijanowski. 2021. "What Does Resilience Sound like? Coral Reef
 and Dry Forest Acoustic Communities Respond Differently to Hurricane Maria."

989   *Ecological Indicators* 126 (July): 107635.
990   https://doi.org/10.1016/j.ecolind.2021.107635.
Greenhalgh, Jack A., Martin J. Genner, Gareth Jones, and Camille Desjonquères. 2020.
"The Role of Freshwater Bioacoustics in Ecological Research." *WIREs Water* 7 (3):
e1416. https://doi.org/10.1002/wat2.1416.
Grinfeder, Elie, Sylvain Haupert, Manon Ducrettet, Julien Barlet, Marie-Pierre Reynet,
Frédéric Sèbe, and Jérôme Sueur. 2022. "Soundscape Dynamics of a Cold
Protected Forest: Dominance of Aircraft Noise." *Landscape Ecology* 37 (2): 567–82.
https://doi.org/10.1007/s10980-021-01360-1.
Halliday, William D., Matthew K. Pine, Xavier Mouy, Piia Kortsalo, R. Casey Hilliard, and
Stephen J. Insley. 2020. "The Coastal Arctic Marine Soundscape near Ulukhaktok,
Northwest Territories, Canada." *Polar Biology* 43 (6): 623–36.
https://doi.org/10.1007/s00300-020-02665-8.
Haupert, Sylvain, Frédéric Sèbe, and Jérôme Sueur. 2022. "Physics-Based Model to Predict
the Acoustic Detection Distance of Terrestrial Autonomous Recording Units over the
Diel Cycle and across Seasons: Insights from an Alpine and a Neotropical Forest."
Methods in Ecology and Evolution n/a (n/a). https://doi.org/10.1111/2041-
210X.14020.
Haver, Samara M., Michelle E. H. Fournet, Robert P. Dziak, Christine Gabriele, Jason
Gedamke, Leila T. Hatch, Joseph Haxel, et al. 2019. "Comparing the Underwater
Soundscapes of Four U.S. National Parks and Marine Sanctuaries." *Frontiers in
Marine Science* 6: 500. https://doi.org/10.3389/fmars.2019.00500.
Haver, Samara M., Jason Gedamke, Leila T. Hatch, Robert P. Dziak, Sofie Van Parijs,
Megan F. McKenna, Jay Barlow, et al. 2018. "Monitoring Long-Term Soundscape
Trends in U.S. Waters: The NOAA/NPS Ocean Noise Reference Station Network."
Marine Policy 90 (April): 6–13. https://doi.org/10.1016/j.marpol.2018.01.023.
Havlik, Michelle-Nicole, Milica Predragovic, and Carlos Manuel Duarte. 2022. "State of Play
in Marine Soundscape Assessments." *Frontiers in Marine Science* 9.
https://www.frontiersin.org/articles/10.3389/fmars.2022.919418.
Hernández-León, Santiago, C. Almeida, L. Yebra, J. Arístegui, M. L. Fernández de Puelles,
and J. García-Braun. 2001. "Zooplankton Abundance in Subtropical Waters: Is There
a Lunar Cycle?" *Scientia Marina* 65 (S1): 59–64.
https://doi.org/10.3989/scimar.2001.65s159.
Hill, Andrew P., Peter Prince, Jake L. Snaddon, C. Patrick Doncaster, and Alex Rogers.
2019. "AudioMoth: A Low-Cost Acoustic Device for Monitoring Biodiversity and the
Environment." *HardwareX* 6 (October): e00073.
https://doi.org/10.1016/j.ohx.2019.e00073.
Hindell, Mark A., Ryan R. Reisinger, Yan Ropert-Coudert, Luis A. Hückstädt, Philip N.
Trathan, Horst Bornemann, Jean-Benoît Charrassin, et al. 2020. "Tracking of Marine
Predators to Protect Southern Ocean Ecosystems." *Nature* 580 (7801): 87–92.
https://doi.org/10.1038/s41586-020-2126-y.
Inyang, Edidiong I., Raymond L. Hix, Violeta Tsolova, Barukh B. Rohde, Omotola Dosunmu,
and Richard W. Mankin. 2019. "Subterranean Acoustic Activity Patterns of Vitacea
Polistiformis (Lepidoptera: Sesiidae) in Relation to Abiotic and Biotic Factors."
Insects 10 (9): 267. https://doi.org/10.3390/insects10090267.
IQOE. n.d. "Acoustic Capabilities of Existing Observing Systems." Accessed March 15,
2023. https://www.iqoe.org/systems.
ISO. 2022. "ISO 18405:2017: Underwater Acoustics — Terminology." TC 43/SC 3
Underwater acoustics.
Jerem, Paul, and Fiona Mathews. 2021. "Trends and Knowledge Gaps in Field Research
Investigating Effects of Anthropogenic Noise." *Conservation Biology* 35 (1): 115–29.
https://doi.org/10.1111/cobi.13510.
Kang, Jian, Francesco Aletta, Tin Oberman, Andrew Mitchell, Mercede Erfanian, Huan Tong,
Simone Torresin, Chunyang Xu, Tingting Yang, and Xiaochao Chen. 2023.
"Supportive Soundscapes Are Crucial for Sustainable Environments." *Science of The*

Total Environment 855 (January): 158868.
https://doi.org/10.1016/j.scitotenv.2022.158868.

Kays, Roland, Sarah C. Davidson, Matthias Berger, Gil Bohrer, Wolfgang Fiedler, Andrea Flack, Julian Hirt, et al. 2022. "The Movebank System for Studying Global Animal Movement and Demography." *Methods in Ecology and Evolution* 13 (2): 419–31. https://doi.org/10.1111/2041-210X.13767.

Keith, David A., Jose R. Ferrer-Paris, Emily Nicholson, Melanie J. Bishop, Beth A. Polidoro, Eva Ramirez-Llodra, Mark G. Tozer, et al. 2020. "Indicative Distribution Maps for Ecosystem Functional Groups - Level 3 of IUCN Global Ecosystem Typology." Zenodo. https://doi.org/10.5281/zenodo.4018314.

Keith, David A., José R. Ferrer-Paris, Emily Nicholson, Melanie J. Bishop, Beth A. Polidoro, Eva Ramirez-Llodra, Mark G. Tozer, et al. 2022. "A Function-Based Typology for Earth's Ecosystems." *Nature*, October, 1–6. https://doi.org/10.1038/s41586-022-05318-4.

Keith, David A., José R. Ferrer-Paris, Emily Nicholson, and Richard T. Kingsford. 2020. *The IUCN Global Ecosystem Typology 2.0: Descriptive Profiles for Biomes and Ecosystem Functional Groups*. Gland, Switzerland: IUCN.

Krause, Bernie, Stuart H. Gage, and Wooyeong Joo. 2011. "Measuring and Interpreting the Temporal Variability in the Soundscape at Four Places in Sequoia National Park." *Landscape Ecology* 26 (9): 1247. https://doi.org/10.1007/s10980-011-9639-6.

Kronfeld-Schor, Noga, Davide Dominoni, Horacio de la Iglesia, Oren Levy, Erik D. Herzog, Tamar Dayan, and Charlotte Helfrich-Forster. 2013. "Chronobiology by Moonlight." *Proceedings of the Royal Society B: Biological Sciences* 280 (1765): 20123088. https://doi.org/10.1098/rspb.2012.3088.

Kuşku, Halit, Murat Yiğit, Sebahattin Ergün, Ümüt Yiğit, and Nic Taylor. 2018. "ACOUSTIC NOISE POLLUTION FROM MARINE INDUSTRIAL ACTIVITIES: EXPOSURE AND IMPACTS." *AQUATIC RESEARCH*, 148–61. https://doi.org/10.3153/AR18017.

Lamont, Timothy A. C., Lucille Chapuis, Ben Williams, Sasha Dines, Tess Gridley, Guilherme Frainer, Jack Fearey, et al. 2022. "HydroMoth: Testing a Prototype Low-Cost Acoustic Recorder for Aquatic Environments." *Remote Sensing in Ecology and Conservation* 8 (3): 362–78. https://doi.org/10.1002/rse2.249.

Lima, Mauro S. C. S., Jonas Pederassi, Ulisses Caramaschi, Kléssia Denise S. S. Sousa, and Carlos Alberto S. Souza. 2021. "Frog Vocalization Is Influenced by Moon Phases: Brazilian Frogs Tend to Prefer Low-Albedo Phases." *Web Ecology* 21 (1): 1–13. https://doi.org/10.5194/we-21-1-2021.

Lin, Tzu-Hao, Frederic Sinniger, Saki Harii, and Tomonari Akamatsu. 2023. "Using Soundscapes to Assess Changes in Coral Reef Social-Ecological Systems." *Oceanography*. https://doi.org/10.5670/oceanog.2023.s1.7.

Lin, Tzu-Hao, and Yu Tsao. 2020. "Source Separation in Ecoacoustics: A Roadmap towards Versatile Soundscape Information Retrieval." *Remote Sensing in Ecology and Conservation* 6 (3): 236–47. https://doi.org/10.1002/rse2.141.

Lindseth, Adelaide V., and Phillip S. Lobel. 2018. "Underwater Soundscape Monitoring and Fish Bioacoustics: A Review." *Fishes* 3 (3): 36. https://doi.org/10.3390/fishes3030036.

Linke, Simon, Toby Gifford, Camille Desjonquères, Diego Tonolla, Thierry Aubin, Leah Barclay, Chris Karaconstantis, Mark J Kennard, Fanny Rybak, and Jérôme Sueur. 2018. "Freshwater Ecoacoustics as a Tool for Continuous Ecosystem Monitoring." *Frontiers in Ecology and the Environment* 16 (4): 231–38. https://doi.org/10.1002/fee.1779.

Looby, Audrey, Christine Erbe, Santiago Bravo, Kieran Cox, Hailey L. Davies, Lucia Di Iorio, Youenn Jézéquel, et al. 2023. "Global Inventory of Species Categorized by Known Underwater Sonifery." *Scientific Data* 10 (1): 1–10. https://doi.org/10.1038/s41597-023-02745-4.

Looby, Audrey, Sarah Vela, Kieran Cox, Amalis Riera, Santiago Bravo, Hailey L. Davies, Rodney Rountree, et al. 2022. "FishSounds Version 1.0: A Website for the

Compilation of Fish Sound Production Information and Recordings." *Ecological Informatics*, December, 101953. https://doi.org/10.1016/j.ecoinf.2022.101953.

Luypaert, Thomas, Anderson S. Bueno, Gabriel S. Masseli, Igor L. Kaefer, Marconi Campos-Cerqueira, Carlos A. Peres, and Torbjørn Haugaasen. 2022. "A Framework for Quantifying Soundscape Diversity Using Hill Numbers." *Methods in Ecology and Evolution* 13 (10): 2262–74. https://doi.org/10.1111/2041-210X.13924.

Maeder, Marcus, Xianda Guo, Felix Neff, Doris Schneider Mathis, and Martin M. Gossner. 2022. "Temporal and Spatial Dynamics in Soil Acoustics and Their Relation to Soil Animal Diversity." *PLOS ONE* 17 (3): e0263618. https://doi.org/10.1371/journal.pone.0263618.

Marques, Tiago A., Len Thomas, Stephen W. Martin, David K. Mellinger, Jessica A. Ward, David J. Moretti, Danielle Harris, and Peter L. Tyack. 2013. "Estimating Animal Population Density Using Passive Acoustics." *Biological Reviews* 88 (2): 287–309. https://doi.org/10.1111/brv.12001.

Messager, Mathis Loïc, Bernhard Lehner, Charlotte Cockburn, Nicolas Lamouroux, Hervé Pella, Ton Snelder, Klement Tockner, Tim Trautmann, Caitlin Watt, and Thibault Datry. 2021. "Global Prevalence of Non-Perennial Rivers and Streams." *Nature* 594 (7863): 391–97. https://doi.org/10.1038/s41586-021-03565-5.

Metcalf, Oliver, Carlos Abrahams, Bob Ashington, Ed Baker, Tom Bradfer-Lawrence, Ella Browning, Jonathan Carruthers-Jones, et al. 2023. "Good Practice Guidelines for Long-Term Ecoacoustic Monitoring in the UK." Report. The UK Acoustics Network. https://acoustics.ac.uk/.

Mickle, Megan F., and Dennis M. Higgs. 2018. "Integrating Techniques: A Review of the Effects of Anthropogenic Noise on Freshwater Fish." *Canadian Journal of Fisheries and Aquatic Sciences* 75 (9): 1534–41. https://doi.org/10.1139/cjfas-2017-0245.

Mirtl, Michael. 2018. "eLTER, European Long-Term Ecosystem and Socio-Ecological Research Infrastructure, H2020." *Impact* 2018 (8): 30–32. https://doi.org/10.21820/23987073.2018.8.30.

Mougeot, François, and Vincent Bretagnolle. 2000. "Predation Risk and Moonlight Avoidance in Nocturnal Seabirds." *Journal of Avian Biology* 31 (3): 376–86. https://doi.org/10.1034/j.1600-048X.2000.310314.x.

Müller, Jörg, Oliver Mitesser, H. Martin Schaefer, Sebastian Seibold, Annika Busse, Peter Kriegel, Dominik Rabl, et al. 2023. "Soundscapes and Deep Learning Enable Tracking Biodiversity Recovery in Tropical Forests." *Nature Communications* 14 (1): 6191. https://doi.org/10.1038/s41467-023-41693-w.

Murchy, K.A., Rountree, R.A., and Juanes, F. n.d. "The Role of the Soundscape in the Behavioral Ecology of Stream-Dwelling Salmonids." In .

Nieto-Mora, D. A., Susana Rodríguez-Buritica, Paula Rodríguez-Marín, J. D. Martínez-Vargaz, and Claudia Isaza-Narváez. 2023. "Systematic Review of Machine Learning Methods Applied to Ecoacoustics and Soundscape Monitoring." *Heliyon* 9 (10): e20275. https://doi.org/10.1016/j.heliyon.2023.e20275.

Parsons, Miles J. G., Tzu-Hao Lin, T. Aran Mooney, Christine Erbe, Francis Juanes, Marc Lammers, Songhai Li, et al. 2022. "Sounding the Call for a Global Library of Underwater Biological Sounds." *Frontiers in Ecology and Evolution* 10. https://doi.org/10.3389/fevo.2022.810156.

Pensieri, Sara, Roberto Bozzano, Jeffrey A. Nystuen, Emmanouil N. Anagnostou, Marios N. Anagnostou, and Renzo Bechini. 2015. "Underwater Acoustic Measurements to Estimate Wind and Rainfall in the Mediterranean Sea." *Advances in Meteorology* 2015 (April): e612512. https://doi.org/10.1155/2015/612512.

Pérez-Granados, Cristian, Karl-L. Schuchmann, and Marinez I. Marques. 2020. "Vocal Behavior of the Undulated Tinamou (Crypturellus Undulatus) over an Annual Cycle in the Brazilian Pantanal: New Ecological Information." *Biotropica* 52 (1): 165–71. https://doi.org/10.1111/btp.12742.

Pérez-Granados, Cristian, and Juan Traba. 2021. "Estimating Bird Density Using Passive Acoustic Monitoring: A Review of Methods and Suggestions for Further Research."

Ibis 163 (3). https://doi.org/10.1111/ibi.12944.

Pijanowski, Bryan C., Almo Farina, Stuart H. Gage, Sarah L. Dumyahn, and Bernie L. Krause. 2011. "What Is Soundscape Ecology? An Introduction and Overview of an Emerging New Science." *Landscape Ecology* 26 (9): 1213–32. https://doi.org/10.1007/s10980-011-9600-8.

Pijanowski, Bryan C., Luis J. Villanueva-Rivera, Sarah L. Dumyahn, Almo Farina, Bernie L. Krause, Brian M. Napoletano, Stuart H. Gage, and Nadia Pieretti. 2011. "Soundscape Ecology: The Science of Sound in the Landscape." *BioScience* 61 (3): 203–16. https://doi.org/10.1525/bio.2011.61.3.6.

Reboredo Segovia, Ana L, Donato Romano, and Paul R Armsworth. 2020. "Who Studies Where? Boosting Tropical Conservation Research Where It Is Most Needed." *Frontiers in Ecology and the Environment* 18 (3): 159–66. https://doi.org/10.1002/fee.2146.

Roch, Marie A., Heidi Batchelor, Simone Baumann-Pickering, Catherine L. Berchok, Danielle Cholewiak, Ei Fujioka, Ellen C. Garland, et al. 2016. "Management of Acoustic Metadata for Bioacoustics." *Ecological Informatics* 31 (January): 122–36. https://doi.org/10.1016/j.ecoinf.2015.12.002.

Roe, Paul, Philip Eichinski, Richard A. Fuller, Paul G. McDonald, Lin Schwarzkopf, Michael Towsey, Anthony Truskinger, David Tucker, and David M. Watson. 2021. "The Australian Acoustic Observatory." *Methods in Ecology and Evolution* 12 (10): 1802–8. https://doi.org/10.1111/2041-210X.13660.

Ross, Samuel R. P.-J., Darren P. O'Connell, Jessica L. Deichmann, Camille Desjonquères, Amandine Gasc, Jennifer N. Phillips, Sarab S. Sethi, Connor M. Wood, and Zuzana Burivalova. 2023. "Passive Acoustic Monitoring Provides a Fresh Perspective on Fundamental Ecological Questions." *Functional Ecology* 37 (4): 959–75. https://doi.org/10.1111/1365-2435.14275.

Rountree, Rodney A., Francis Juanes, and Marta Bolgan. 2020. "Temperate Freshwater Soundscapes: A Cacophony of Undescribed Biological Sounds Now Threatened by Anthropogenic Noise." *PLOS ONE* 15 (3): e0221842. https://doi.org/10.1371/journal.pone.0221842.

Rountree, Rodney, Jacopo Aguzzi, Simone Marini, Emanuela Fanelli, Fabio De Leo, Joaquin del Rio, and Francis Juanes. 2020. "Towards an Optimal Design for Ecosystem-Level Ocean Observatories." In , 79–106. https://doi.org/10.1201/9780429351495-2.

Scarpelli, Marina D. A., Milton Cezar Ribeiro, Fernanda Z. Teixeira, Robert J. Young, and Camila P. Teixeira. 2020. "Gaps in Terrestrial Soundscape Research: It's Time to Focus on Tropical Wildlife." *Science of The Total Environment* 707 (March): 135403. https://doi.org/10.1016/j.scitotenv.2019.135403.

Schaaf, Alejandro Alberto, Martín Boullhesen, and Mauricio Sebastián Akmentins. 2023. "Vocal Activity of the Rufous Nightjar (*Antrostomus Rufus*) in Subtropical Yungas Andean Forests of Argentina." *Neotropical Biodiversity* 9 (1): 59–63. https://doi.org/10.1080/23766808.2023.2185963.

Sequeira, A. M. M., J. P. Rodríguez, V. M. Eguíluz, R. Harcourt, M. Hindell, D. W. Sims, C. M. Duarte, et al. 2018. "Convergence of Marine Megafauna Movement Patterns in Coastal and Open Oceans." *Proceedings of the National Academy of Sciences* 115 (12): 3072–77. https://doi.org/10.1073/pnas.1716137115.

Sethi, Sarab S., Nick S. Jones, Ben D. Fulcher, Lorenzo Picinali, Dena Jane Clink, Holger Klinck, C. David L. Orme, Peter H. Wrege, and Robert M. Ewers. 2020. "Characterizing Soundscapes across Diverse Ecosystems Using a Universal Acoustic Feature Set." *Proceedings of the National Academy of Sciences*, July. https://doi.org/10.1073/pnas.2004702117.

Shannon, Graeme, Megan F. McKenna, Lisa M. Angeloni, Kevin R. Crooks, Kurt M. Fristrup, Emma Brown, Katy A. Warner, et al. 2016. "A Synthesis of Two Decades of Research Documenting the Effects of Noise on Wildlife: Effects of Anthropogenic Noise on Wildlife." *Biological Reviews* 91 (4): 982–1005. https://doi.org/10.1111/brv.12207.

Siddagangaiah, Shashidhar, Chi-Fang Chen, Wei-Chun Hu, and Almo Farina. 2022. "The Dynamical Complexity of Seasonal Soundscapes Is Governed by Fish Chorusing." *Communications Earth & Environment* 3 (1): 1–11. https://doi.org/10.1038/s43247-022-00442-5.

Simonis, Anne E., Marie A. Roch, Barbara Bailey, Jay Barlow, Rachel E. S. Clemesha, Sam Iacobellis, John A. Hildebrand, and Simone Baumann-Pickering. 2017. "Lunar Cycles Affect Common Dolphin Delphinus Delphis Foraging in the Southern California Bight." *Marine Ecology Progress Series* 577 (August): 221–35. https://doi.org/10.3354/meps12247.

Slabbekoorn, Hans, Robert J. Dooling, Arthur N. Popper, and Richard R. Fay, eds. 2018. *Effects of Anthropogenic Noise on Animals*. Vol. 66. Springer Handbook of Auditory Research. New York, NY: Springer. https://doi.org/10.1007/978-1-4939-8574-6.

Solé, Marta, Kenzo Kaifu, T. Aran Mooney, Sophie L. Nedelec, Frédéric Olivier, Andrew N. Radford, Mirella Vazzana, et al. 2023. "Marine Invertebrates and Noise." *Frontiers in Marine Science* 10. https://www.frontiersin.org/articles/10.3389/fmars.2023.1129057.

Sousa, Ivandson P. de, Gustavo Zampier dos Santos Lima, Eliziane G. Oliveira, Marina Henriques Lage Duarte, José A. Alves-Gomes, Lara C. Lopes, Luane S. Ferreira, Renata S. Sousa-Lima, and Gilberto Corso. 2022. "Scale-Free Distribution of Silences." *Physical Review E* 105 (1): 014107. https://doi.org/10.1103/PhysRevE.105.014107.

Sousa-Lima, Renata. 2013. "A Review and Inventory of Fixed Autonomous Recorders for Passive Acoustic Monitoring of Marine Mammals." *Aquatic Mammals* 39 (1): 23–53. https://doi.org/10.1578/AM.39.1.2013.23.

Sousa-Lima, Renata S., and Christopher W. Clark. 2008. "Modeling the Effect of Boat Traffic on the Fluctuation of Humpback Whale Singing Activity in the Abrolhos National Marine Park, Brazil." *Canadian Acoustics* 36 (1): 174–81.

Spatz, Dena R., Nick D. Holmes, David J. Will, Stella Hein, Zachary T. Carter, Rachel M. Fewster, Bradford Keitt, et al. 2022. "The Global Contribution of Invasive Vertebrate Eradication as a Key Island Restoration Tool." *Scientific Reports* 12 (1): 13391. https://doi.org/10.1038/s41598-022-14982-5.

Stowell, Dan, and Jérôme Sueur. 2020. "Ecoacoustics: Acoustic Sensing for Biodiversity Monitoring at Scale." *Remote Sensing in Ecology and Conservation* 6 (3): 217–19. https://doi.org/10.1002/rse2.174.

Sueur, Jérôme, and Almo Farina. 2015. "Ecoacoustics: The Ecological Investigation and Interpretation of Environmental Sound." *Biosemiotics* 8 (3): 493–502. https://doi.org/10.1007/s12304-015-9248-x.

Sueur, Jérôme, Bernie Krause, and Almo Farina. 2019. "Climate Change Is Breaking Earth's Beat." *Trends in Ecology & Evolution* 34 (11): 971–73. https://doi.org/10.1016/j.tree.2019.07.014.

Sugai, Larissa Sayuri Moreira, Camille Desjonquères, Thiago Sanna Freire Silva, and Diego Llusia. 2020. "A Roadmap for Survey Designs in Terrestrial Acoustic Monitoring." *Remote Sensing in Ecology and Conservation* 6 (3): 220–35. https://doi.org/10.1002/rse2.131.

Sugai, Larissa Sayuri Moreira, and Diego Llusia. 2019. "Bioacoustic Time Capsules: Using Acoustic Monitoring to Document Biodiversity." *Ecological Indicators* 99 (April): 149–52. https://doi.org/10.1016/j.ecolind.2018.12.021.

Sugai, Larissa Sayuri Moreira, Thiago Sanna Freire Silva, José Wagner Ribeiro, and Diego Llusia. 2019. "Terrestrial Passive Acoustic Monitoring: Review and Perspectives." *BioScience* 69 (1): 15–25. https://doi.org/10.1093/biosci/biy147.

Van Doren, Benjamin M., Vincent Lostanlen, Aurora Cramer, Justin Salamon, Adriaan Dokter, Steve Kelling, Juan Pablo Bello, and Andrew Farnsworth. 2023. "Automated Acoustic Monitoring Captures Timing and Intensity of Bird Migration." *Journal of Applied Ecology* 60 (3): 433–44. https://doi.org/10.1111/1365-2664.14342.

Van Renterghem, Timothy. 2019. "Towards Explaining the Positive Effect of Vegetation on the Perception of Environmental Noise." *Urban Forestry & Urban Greening*, Urban

1264   green infrastructure – connecting people and nature for sustainable cities, 40 (April):
1265   133–44. https://doi.org/10.1016/j.ufug.2018.03.007.
1266 Xeno-canto Foundation. 2012. *Xeno-Canto: Sharing Bird Sounds from around the World*.
1267   Xeno-canto Foundation Amsterdam.

A - Framework

Space

Spatial
replicates

Temporal
replicates

Time

B - Replication of datasets

	temporal	spatial	spatio-temporal
Terrestrial	12	9	188
Marine	13		36
Freshwater	3	2	10

C - Spatial extent and coverage (sites)

Terrestrial sites   100 200 300 400

D - Temporal extent and coverage (days)

1990   2000   2010   2020

Core realm   Freshwater   Marine   Terrestrial

A - Topographical and latitudinal distribution across realms (sites)

IUCN realms
(core and transitional)

- ○ Freshwater
- ○ Freshwater-Marine
- ○ Marine
- ○ Marine-Freshwater-Terrestrial
- ○ Marine-Terrestrial
- ○ Terrestrial
- ○ Terrestrial-Freshwater

B - Top 10 administrative regions of core realms (sites)

WDPA
protected areas

- Ia
- Ib
- II
- >II

Freshwater

Spain	22%
United Kingdom	18%
United States	16%
Colombia	13%
Australia	9%
Brazil	8%
France	7%
Taiwan	3%
Japan	1%
Canada	1%

Marine

others (not top 10)	17%
South Atlantic Ocean	16%
Southern Ocean	15%
North Pacific Ocean	11%
The Northwestern Passages	9%
North Atlantic Ocean	7%
Coral Sea	7%
South China Sea	5%
Indian Ocean	5%
Caribbean Sea	4%
Beaufort Sea	3%

Terrestrial

Canada	26%
others (not top 10)	25%
Germany	14%
Brazil	7%
United States	6%
France	5%
Portugal	5%
Spain	4%
Panama	3%
Puerto Rico	3%
Burkina Faso	2%

A - Sampling intensity across realms

	Spatial extent [M km2]	Spatial coverage and density	Temporal extent, coverage, density
Marine	2028	9.1	0.44
Subterranean	541	0	
Terrestrial	239	32.6	0.16
Freshwater	144	6.6	0.4
Subterranean-Freshwater	79	0	
Terrestrial-Freshwater	54	7.4	0.17
Marine-Terrestrial	6	5	0.05
Freshwater-Marine	5	5.7	0.12
Subterranean-Marine	2	0	
Marine-Freshwater-Terrestrial	1	1	0.17

B - Spectral range used for focal taxa in core realms (datasets)