The role of FraI in cell-cell communication and differentiation in the hormogonia-forming cyanobacterium Nostoc punctiforme

Ana Janović [1], Iris Maldener [1,C*], Claudia Menzel [1], Gabriel A. Parrett [2], Douglas D. Risser [2,C*]

[1] Interfaculty Institute of Microbiology and Infection Medicine Tübingen, Organismic Interactions, University of Tübingen, Auf der Morgenstelle 28, 72076 Tübingen, Germany

[2] Department of Biology, University of Colorado, Colorado Springs, USA

[*] Correspondence: drisser@uccs.edu; iris.maldener@uni-tuebingen.de

Keywords

Cell-cell communication, hormogonia, filamentous cyanobacteria, *Nostoc punctiforme,* motility, septal junctions, NpF4142, FraI

Abstract

Multicellular cyanobacteria, like *Nostoc punctiforme,* rely on septal junctions for cell-cell communication, which is crucial for coordinating various physiological processes including differentiation of N_2-fixing heterocysts, spore-like akinetes and hormogonia - short, motile filaments important for dispersal. In this study we functionally characterize a protein, encoded by gene *NpF4142*, which in a random mutagenesis approach, initially showed a motility-related function. The reconstructed *NpF4142* knockout mutant exhibits further distinct phenotypic traits, including altered hormogonia formation with significant reduced motility, inability to differentiate heterocysts and filament fragmentation. For that reason, we named the protein FraI (fragmentation phenotype). The mutant displays severely impaired cell-cell communication, due to almost complete absence of the nanopore array in the septal cell wall, which is an essential part of the septal junctions. Despite lack of communication, hormogonia in the Δ*fraI* mutant maintain motility and phototactic behaviour, even though less pronounced than the wild type. This suggests an alternative mechanism for coordinated movement beyond septal junctions. Our study underscores the significance of FraI in nanopore formation and cell differentiation and provides additional evidence for the importance of septal junction formation and communication in various differentiation traits of cyanobacteria. The findings contribute to a deeper understanding of the regulatory networks governing multicellular cyanobacterial behaviour, with implications for broader insights into microbial multicellularity.

Importance

The filament-forming cyanobacterium *Nostoc punctiforme* serves as a valuable model for studying cell differentiation, including the formation of nitrogen-fixing heterocysts and hormogonia. Hormogonia filaments play a crucial role in dispersal and plant colonization, providing a nitrogen source through atmospheric nitrogen fixation, thus holding promise for fertilizer-free agriculture. The coordination among the hormogonia cells enabling uniform movement towards the positive signal remains poorly understood. This study investigates the role of septal junction-mediated communication in hormogonia differentiation and motility, by studying a Δ*fraI* mutant with significantly impaired communication. Surprisingly, impaired communication does not abolish synchronized filament movement, suggesting an alternative coordination mechanism. These findings deepen our understanding of cyanobacterial biology and have broader implications for multicellular behavior in prokaryotes.

Introduction

The filamentous cyanobacterium *Nostoc punctiforme* ATCC 29133/PCC73102 is a true multicellular bacterium, able to form specialized cells which enable the bacterium to adapt to various environmental conditions. Heterocysts are nitrogen-fixing cells that differentiate from semi-regularly positioned vegetative cells upon nitrogen starvation to provide a microooxic environment for the oxygen-labile nitrogenase[1,2]. Hormogonia (see below) are short filaments specialized as motile units allowing dispersal. Finally, unicellular akinetes form under various stress conditions enabling survival as spore-like resistant cells [3–5].

Within a filament, cells interact through a remarkable cell-cell communication system, involving so called septal junctions. These multi-protein complexes connect the individual cells by traversing the septal peptidoglycan layer through nanopores[6,7]. On each cytoplasmic side, every septal junction contains a flexible cap and plug structure by which the intercellular connection closes in a reversible manner. This gating mechanism of halting communication is crucial under stress conditions, such as UV-C light exposure, cytoplasm leakage due to filament breakage, or the isolation of senescent cells from the rest of the filament [7–9]. Mutants with impaired cell-cell communication also exhibit deficiencies in heterocyst development[1,10–14], emphasizing the significance of intercellular signaling in cellular differentiation [1,2,10–15].

Formation of septal nanopores requires the activity of cell wall amidases. In unicellular bacteria amidases play a pivotal role in cell division, by cleaving the newly synthesized peptidoglycan to facilitate separation of daughter cells [16]. In multicellular cyanobacteria, cell wall amidases play a role in cell-cell communication, by perforating the septal peptidoglycan[13]. Several amidases were identified in multicellular cyanobacteria [12,13,17,18]. A knockout mutant of the

AmiC2 enzyme in *N. punctiforme* showed severe impairment in cell division, filament

morphology and cell-cell communication, caused by the complete absence of nanopores in the

septal peptidoglycan [17]. The *amiC2* mutant is unable to differentiate heterocysts, hormogonia

and akinetes[12,17]. The AmiC2 protein primarily localizes along the septal plane of nascent septa

and is typically absent in fully developed septa [12]. AmiC1, another amidase encoded in the

same gene operon, is likely essential for *Nostoc punctiforme,* since the *knockout* of this gene

was reported to be impossible to create [12,17]. In the closely related cyanobacterium *Nostoc* sp.

PCC 7120 mutants in both orthologues, *amiC1* and *amiC2* as well as double *amiC1amiC2*

knockouts were created. All mutants exhibited aberrant nanopore shape and numbers and

lower rates of cell-cell communication compared to the wild type [13]. Furthermore,

overexpression of AmiC1 in *Nostoc* sp. 7120 resulted in a lethal phenotype, in contrast to the

AmiC2 overexpression, which has no significant effect on growth. The LytM factor Alr3353 is

an activator of AmiC1 of *Nostoc* sp. 7120 and its absence leads to a reduction in nanopore

numbers and the rate of cell-cell communication [19]. To date, no other regulators of amidase´s

nanopore-drilling activity have been identified.

Transiently formed hormogonia, 10-20 cell-long filaments of *Nostoc punctiforme*, play a crucial

role in dispersal and symbiotic plant colonization [20]. After colonization of the host plant, the

hormogonia develop back to vegetative filaments containing heterocysts for N_2 fixation. This

symbiosis is of potential agricultural importance as it enables fertilizer-free cultivation providing

nitrogen sources for the crop, as shown for *Oryza sativa* [21]. Induction of hormogonia

differentiation from vegetative filaments can be triggered by various signals such as nitrogen

stepdown, red-light exposure or HIF (*hormogonium-inducing factor*) exposure. Following the

induction signal, vegetative cells differentiate into hormogonia within 18-24 hours through rapid

cell divisions not accompanied by overall biomass growth. These filaments demonstrate photo

and chemo-tactic movement, facilitated by a type IV pili machinery on their outer cell's surface.

The movement is additionally supported by the export of an exopolysaccharide sheath known

as hormogonia polysaccharide (HPS) [22]. Directionality is determined by the presence of the

HmpF protein at one pole of each cell, positioned uniformly at the leading poles of each cell in

motile filaments. Repositioning of the HmpF protein occurs rapidly and synchronously along

the filament, typically within about 30 seconds, in response to changes in light conditions,

leading to a change in the filament's course. [23,24]. A putative amidase encoded by *the tftA* gene

was shown to be important for the tapered cell phenotype at the filament's termini, likely by the

degradation of the terminal peptidoglycan. Further, a *tftA* knockout showed reduced migration

ability through agar [25]. However, the role of cell-cell communication in the synchronized

movement of the entire filament and during the change of movement's direction in hormogonia

remained uninvestigated. In this study, we identified a mutant lacking the NpF4142 protein,

herein designated FraI, that presents decreased hormogonia motility. Characterization of this

mutant strain allowed us to show that FraI plays a key role in nanopore formation and hormogonia differentiation. Furthermore, this study showed that transmission of the signal for direction of movement in hormogonia probably does not involve communication through the septal junctions.

Material and methods

Cultivation of bacterial strains

As standard medium, *N. punctiforme* and the mutant strains were cultivated in AA/4 medium according to Allen and Arnon [26] in a fourfold dilution. To supply a source of fixed nitrogen, the medium was supplemented with 2.5 mM NH_4Cl and 5 mM MOPS buffer, pH 7.8. In experiments performed in Tübingen the cyanobacteria were cultivated in either liquid BG11 medium [27] or AA/4 medium supplemented with 2.5 mM KNO_3 and 2.5 mM $NaNO_3$. Liquid cultures were shaken at 120 rpm, agar plates contained 1% Difco agar. Growth occurred at 28 °C with constant illumination at 25-40 $\mu E\ m^{-2}\ s^{-1}$. Sucralose was added to suppress hormogonia development in concentrations mentioned in the text. Neomycin was used at a concentration of 50 $\mu g/ml$ in appropriate strains.

Plasmid and strain construction

For a detailed description of the plasmids, strains, and oligonucleotides used in this study refer to supplemental material, Tables S1. All constructs were sequenced to insure fidelity.

To construct plasmid pDDR558 for in-frame deletion of *fraI* (Npun_F4142), approximately 900 bp of flanking DNA on either side of the gene and several codons at the beginning and end of the gene were amplified via overlap extension PCR using primers NpF4142-5'-F, NpF4142-5'-R, NpF4142-3'-F and NpF4142-3'-R and cloned into pRL278[28] as a BamHI-SacI fragment using restriction sites introduced on the primers.

To construct plasmid pGAP109 for replacement of the chromosomal allele of *fraI* with a C-terminal *gfpuv*-tagged variant, approximately 900 bp of DNA downstream of the stop codon were amplified via PCR using primers NpF4142-gfp-3'-F and NpF4142-3'-R and cloned into pSCR569 [29] , as an SpeI-SacI fragment using restriction sites introduced on the primers. The coding region and approximately 900 bp of DNA upstream of the start codon were then amplified via PCR using primers NpF4142-5'-F and NpF4142-gfp-5'-R and cloned into this plasmid as a BamHI-SmaI fragment using restriction sites introduced on the primers.

Generation of transposon mutants and identification of transposon insertion sites was performed as previously described [30] using plasmid pRL1063a[31]. Gene deletions and allelic replacements were performed as previously described[32] with *N. punctiforme* cultures supplemented with 4 mM sucralose to inhibit hormogonium development and enhance

140   conjugation efficiency [30,33]. To construct UCCS103, plasmid pDDR558 was introduced into wild

141   type *N. punctiforme*. To create UCCS113, plasmid pGAP109 was introduced into UCCS103.

Motility and phototaxis assays

143   Plate and time-lapse motility and phototaxis assays were performed as previously described[34]

144   and imaged using a Leica SD9 dissecting microscope equipped with a Flexcam C3 camera

145   and controlled by Leica X LAS X software. The students T-test was applied to determine

146   statistical significance in colony spreading assays.

Light- and fluorescence microscopy

A Leica DM2500 B microscope equipped with a Leica DFC420C camera, or an Evos M5000

microscope was employed for light microscopy. Fluorescence microscopy utilized a Leica

DM5500 B microscope with a 100x/1.3 oil objective lens, connected to a Leica DFC360FX

camera. GFP and chlorophyll fluorescence were induced using BP470/40 nm or BP535/50 nm

filters, and emission was detected with BP525/50 nm or BP610/75 nm filters, respectively.

Septal visualization was performed using Vancomycin-FL (Bodipy-FL, Invitrogen) fluorescent

stains that specifically stains septal peptidoglycan by the previously described procedure[35].

FRAP and CCCP-FRAP

A three-day-old liquid culture of vegetative filaments from both the *N. punctiforme* wild type

and the Δ*fraI* mutant strains was utilized for CCCP-FRAP, following previously established

protocols [8,36]. Briefly, 2 ml of culture underwent triple washing with BG11 medium and were

then resuspended in 500 µL of medium. Subsequently, the cells were incubated with 10 – 15

µL of 1 mg/ml calcein in DMSO (Sigma-Aldrich) for 90 minutes in the dark at 28 °C with

shaking. Following this incubation period, the cells underwent three additional washes with

BG11 medium and were then either resolved in medium alone or in medium containing

CCCP (from a 50 mM stock solution in DMSO), achieving a final concentration of 50 µM in

the culture. After another 90-minute incubation with shaking at 28 °C in the dark, the

filaments were spotted onto BG11 agar plates (1%).

The FRAP procedure was conducted by employing a laser with 488 nm excitation at 0.2%

intensity on a Zeiss LSM 800 confocal microscope equipped with a 63x/1.4 oil-immersion

objective. The ZEN 2.9 (blue edition) software was used for the experiment. Simultaneous

detection of calcein fluorescence emission (400–530 nm) and chlorophyll autofluorescence

emission (650–700 nm) was carried out. To bleach a specific cell, the laser intensity was briefly

increased to 3.5% after capturing 5 pre-bleached images. Subsequently, fluorescence

recovery in the bleached cell was observed at 1-second intervals over a period of 30–

120 seconds. Image analysis was performed using ImageJ (version 1.51j) and GraphPad
Prism 10, as detailed in previous descriptions.

For hormogonia FRAP, a log-phase culture grown with 4 mM sucralose in BG11 medium was
washed twice with BG11 for hormogonia induction. After 18-20 hours following hormogonia
induction the cells were harvested (3-5 ml) and used for FRAP as described above. The
stained cells (25 µL) were put onto a plain microscopic slide and covered with a cover slip. The
measurement was only conducted after the hormogonia ceased movement upon becoming
trapped between coverslip and slide due to evaporation.

Akinete induction

182 Akinetes were induced essentially as described[37]. In short, liquid cultures of WT and NpF4142
183 mutant were grown in BG11 medium with nitrate until early log-phase. Akinete formation was
184 induced by washing the culture twice followed by inoculation in phosphate-free BG11 medium
185 and incubation with low shaking (50 rpm) for 4 weeks.

Isolation of septal peptidoglycan and transmission electron microscopy

Septal peptidoglycan was isolated as described[36]. First, vegetative filaments of *Nostoc*
punctiforme WT and Δ*fraI* grew for 5 days in liquid BG11 medium supplemented with 4 mM
sucralose to supress hormogonia development. For hormogonia induction, a log-phase culture
was induced by washing twice with BG11 medium and incubated in fresh BG11 medium for
20 h. The vegetative and hormogonia cultures were washed and resuspended in 700 µL 0.1 M
Tris-HCl pH 6. Briefly, cells were sonicated (Branson Sonier 250), then boiled in SDS for
30 min. Then, the samples were sonified in a water bath for 30 min, followed by washing and
incubation in 50 mM Na_3PO_4 pH 6.8 with 300 µf of α-chymotrypsin. The chymotrypsin was
readded and the next day the cells were again sonicated for 30-60 s. Purified septa (10 µL)
were added onto an UV-irradiated (16 h) formvar/carbon film-coated copper grid (Electron
Microscopy Sciences) for 30 min, stained with 1% (w/v) uranyl acetate and imaged with a
Philips Tecnai10 electron microscope at 80 kV equipped with a Rio Camera (Gatan). Images
were analysed using ImageJ 2.9. to calculate the number of nanopores per septa and the
septum's diameter.

Transmission electron microscopy of ultrathin sections

To examine hormogonia via transmission electron microscopy, we induced their formation by
washing of the cells previously cultivated in AA/4 media with 2.5 mM of sucralose. After 20
hours, hormogonia were harvested by centrifugation and ultrathin sections were prepared
following standard procedure[38]. Prior to sectioning, the cells underwent fixation with
glutaraldehyde and potassium permanganate. Subsequently, the ultrathin sections were

stained with uranyl acetate and lead citrate. Finally, the samples were analysed using a Philips

Tecnai10 microscope operating at 80 kV.

Immunolocalization of AmiC2

Immunolocalization of AmiC2 in *Nostoc* strains was performed based on two protocols[17,39].

Nostoc punctiforme cultures were cultivated in AA/4 medium with supplemented NaNO$_3$, KNO$_3$

and 2.5 mM sucralose, washed once and resuspended in 1 mL PBS. The cells were fixed with

4% paraformaldehyde PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$,

pH 7.4) on ice for one hour, followed by three rounds of washing with PBS and final incubation

in GTE buffer (50 mM glucose, 25 mM Tris-Cl, pH 8.0, 10 mM EDTA). Then, 200 µL of the cell

suspension were put onto Epredia™TM Polysine adhesion slides (Epredia, Netherlands) and

let to dry. The cells were then fixed with -20 °C methanol for 5 min followed by incubation with

-20 °C acetone for 30 s. Cells were dried at RT and rehydrated with 200 µL PBS for 5 min.

After blocking with 200 µL 2 % w/v BSA in PBS for 30 min, cells were incubated overnight in a

wet chamber at 4 °C with 1:5 anti-AmiC1 [17] in 2 % w/v BSA BSA-PBS. The slides were washed

three times with PBS, followed by a two-hour incubation at room temperature in darkness with

FITC-coupled α-rabbit antibodies (diluted at 1:200 in BSA-PBS from Sigma Aldrich). After

subsequent washing and drying, a drop of Vectashield Mounting Medium H-1200 (Vector

Laboratories, USA) was added. The preparation was covered with a coverslip. Fluorescence

imaging was conducted with a DM5500B Leica microscope and a DFC360FX monochrome

camera. Autofluorescence was measured as mentioned above, while FITC-fluorescence was

captured using a BP470/40 nm excitation filter and a BP525/50 nm emission filter. Z-stacks

with 0.20 µm intervals were acquired and 3D-deconvolution using Leica ASF built-in function

was performed.

Test on significance

231   Statistical analysis was performed with GraphPad Prism version 10. Comparison of one group

232   with multiple other groups was performed via Welch ANOVA test followed by Brown-Forsythe

233   followed by Dunnett's T3 multiple comparison test. Significance *P* is indicated with asterisks:

234   *: *P*≤0.05; **: *P*≤0.01; ***: *P*≤0.001; ****: *P*≤0.0001; ns (not significant): *P*>0.05.

Results and Discussion

fraI deletion affects hormogonia motility, filament length and diazotrophy

Using a transposon mutagenic screen[30] we identified a mutant strain with reduced motility containing a transposon insertion in open reading frame Npun_F4142 (Table S1). Its homologue in *Nostoc sp.* 7120 that we previously named FraI[40] (Kieninger 2023) was shown to be highly abundant in a co-immunoprecipitation experiment against septal junction proteins FraD and SepN [36] and displayed a fragmentation phenotype[40]. Based on this, and the data presented below, we have designated the gene name for Npun_F4142 as *fraI.* To confirm that the transposon insertion in *fraI* was responsible for the motility defect, a strain with an in-frame deletion of *fraI* was constructed (Δ*fraI*). Compared to the wild type, the Δ*fraI* strain exhibited a significant reduction in colony spreading in plate motility assays (Fig. 1A). Moreover, there was a qualitative difference in the appearance of motile colonies for the Δ*fraI* strain, with filaments spreading in collective masses rather than dispersed filaments as observed in the wild type (Fig. 1A). In time-lapse motility assays of individual hormogonium filaments (Video S1) the Δ*fraI* strain exhibited motility of individual filaments, but the filaments were much shorter than those of the wild type, and many of the shortest filaments were non-motile.

Figure 1. Deletion of *fraI* affects motility. **(A)** Images of plate motility assays (left) and quantification of colony spreading (right) of strains as indicated. For images on left, black boxes in top panel correspond to closeup images of colony periphery shown in bottom panel. For quantification, n=3, error bars =+/- 1 S.D., *=p-value <0.01 vs wild type. **(B)** Images of phototaxis assays for strains as indicated. Red circle indicates the initial site of colony at beginning of assay. Direction of light indicated with arrow.

In an attempt to visualize the localization of FraI, a *fraI-gfp* allele was re-introduced at the native chromosomal locus of the Δ*fraI* strain. Introduction of *fraI-gfp* into the deletion strain restored wild-type motility indicating this allele is functional (Fig. 1A), and further demonstrating that deletion of *fraI* was responsible for the observed phenotype. However, no fluorescence signal could be visualized, indicating that the abundance of FraI-GFP in the cell may be below the limit of detection, or that fusion of GFP to the C-terminus of FraI may affect GFP folding or stability. Attempts were made to introduce either the native, or a *gfp*-tagged allele of *fraI* into

the deletion strain on a replicative shuttle vector, but in both cases these efforts failed to yield

viable colonies, possibly implying that increased expression of *fraI* from a multicopy plasmid is

lethal in *N. punctiforme*. Given the motility defect in the Δ*fraI* strain, its capacity for phototaxis

was also investigated (Fig. 1B). Although migration was reduced compared to the wild type,

the Δ*fraI* strain exhibited directional movement in response to light, indicating the mutant strain

retains the ability to perceive and respond to light signals.

In liquid cultures supplemented with ammonia, the Δ*fraI* strain was more dispersed than the

wild type (Fig. 2A). Under these standard growth conditions, the wild-type strain produces a

mixture of vegetative and hormogonium filaments (Fig. 2B). The Δ*fraI* strain also produced a

mixture of both filament types, but in general, the filaments were much shorter (Fig. 2B).

Moreover, we frequently observed what appeared to be hybrid filaments containing a mixture

of vegetative and hormogonium cell types in the same filament (Fig. 2B). Although difficult to

quantify given the tendency of *N. punctiforme* to aggregate into dense clumps, this phenotype

was never observed in the wild-type strain, nor have we ever observed such a phenotype in

previous studies with either the wild type or any of the mutants we have characterized. These

results indicate that while *fraI* is dispensable for hormogonium development and motility, it may

be critical for cells within a filament to synchronize hormogonium differentiation.

Figure 2. Influence of *fraI* on diazotrophy and development. **(A)** Images of cultures (strains as indicated)
in presence (NH$_4^+$) or absence (N$_2$) of a source of fixed nitrogen. **(B)** Light micrographs of individual
filaments (strains as indicated) in presence (NH$_4^+$) or absence (N$_2$) of a source of fixed nitrogen. V =
vegetative filament, H = hormogonia, M = mixed filament containing both hormogonium and vegetative
cells, > = heterocysts. Bar = 10 um. **(C)** Light micrographs (strains as indicated) of filaments following
induction of akinetes by removal of phosphate from the growth medium. A = akinete, Bar = 10 um.

In liquid cultures lacking a source of fixed nitrogen, conditions where the wild type develops

nitrogen-fixing heterocysts to support growth on N$_2$, the Δ*fraI* strain became chlorotic and failed

to grow (Fig. 2A). Under these conditions the fragmentation phenotype became more severe,

and there was no obvious indication that morphologically distinct heterocysts were present

(Fig. 2B). As with motility, the introduction of *fraI-gfp* into the deletion strain restored

diazotrophy (Fig. 2A). Upon phosphate starvation the Δ*fraI* strain was able to develop akinetes

although of aberrant shape and not in consecutive manner when compared to the WT[37] (Fig.

2C). Collectively, the absence of heterocysts and fragmentation phenotype of the Δ*fraI* mutant

resembles that reported for mutants in other genes known to be involved in formation of the

septal junctions, most of which have been characterized primarily in the closely related

filamentous cyanobacterium *Nostoc* sp. PCC 7120.

The reduced motility phenotype motivated us to examine the ultrastructure of hormogonia

filaments using transmission electron microscopy (TEM). In Δ*fraI* we could observe short

hormogonia that were completely lacking the surrounding exopolysaccharide sheath (Fig. 3),

which may be the cause of the of observed reduced motility (Fig. 1). In addition, no septal cell-

cell connections were visible in Δ*fraI* hormogonia (Fig. 3). Furthermore, in contrast to the WT,

the thylakoid membranes were aberrantly arranged. This could be explained by the presence

of numerous glycogen granula, which may disturb the organization of the membranes.

WT hormogonia

Δ *fraI* hormogonia

Figure 3. The hormogonia of Δ*fraI* have no cell-cell connections and are lacking the HPS. After induction of hormogonia, ultra-thin sections were prepared and the ultrastructure investigated by transmission electron microscopy. Shown are representative electron micrographs of each strain. Blue rectangles signify the zoomed region shown below (WT) or at the right (Δ*fraI*). Red arrows point to the cell-cell junctions visible in the WT. Yellow arrows point to the exopolysaccharide sheath present only in the WT hormogonia. G – glycogen granules, TM – thylakoid membrane. Scale bar – 1 μm.

FraI is related to known septal proteins

To further explore the relationship between FraI and septal junctions, we investigated the expression pattern of *fraI* as well as the orthologs of other septal junction genes during hormogonium and heterocyst development in *N. punctiforme* using previously published transcriptomic data sets[29,41,42]. Additionally, the evolutionary co-occurrence of this gene set in cyanobacteria was determined. During hormogonium development all the identified septal junction genes are transcriptionally upregulated in wild-type *N. punctiforme*, with expression typically peaking between 6 to 12 h post induction (Fig. S1A). This observation is consistent with the fact that developing hormogonia undergo a round of cell division, necessitating the construction of large numbers of new septal junctions at the new septa. For most of these genes, enhanced transcription in hormogonia is independent of the hormogonium sigma factor cascade[41], as deletion of *sigJ*, *sigC*, or *sigF* has no obvious effect on expression. In contrast, upregulation of *fraI*, *sepI* and *sjcF1* in hormogonia is dependent on both *sigJ* and *sigC*. Given that expression of *sigC* is also dependent on *sigJ*, these results imply that these three genes are most directly regulated by *sigC*. Transcriptional start site mapping via Cappable seq (CAPseq) indicated the presence of two transcriptional start sites (-265 bp and -200 bp of the *fraI* start codon) in the *fraI* promoter region, with the more distal closely corresponding to total read coverage from RNAseq data (Fig. S1B). However, this TSS only showed a very minor decrease in abundance in the Δ*sigC* strain. In developing heterocysts there was less pronounced upregulation of this gene set, with only a few genes displaying marked upregulation (Fig. S1A). Co-occurrence analysis indicated that FraI was found almost exclusively in heterocyst-forming cyanobacteria, similar to the conservation pattern of SepN, FraD, and FraC, although orthologs of these genes are found occasionally in non-heterocyst forming filamentous cyanobacteria (Fig. S2). Overall, the expression and co-occurrence patterns of *fraI* in comparison to other septal junction genes support a role for *fraI* in the formation of septal junctions.

The Δ*fraI* mutant is severely impaired in cell-cell communication

The relation to septal junction proteins and the Δ*fraI* phenotype [10,11,43], prompted us to investigate cell-cell communication and septal junction gating ability in the Δ*fraI* mutant of *N. punctiforme*[8,36]. The study of cell-cell communication was performed by FRAP experiments

[10,44], where filaments were first stained with a fluorescent dye, subsequently, a single cell is

laser-bleached, and finally the recovery of fluorescence, resulting from the diffusion of the dye

through septal junctions, was monitored over time[10,44]. Compared to the wild type of *N.*

punctiforme, the vegetative cells of the Δ*fraI* mutant were highly impaired in cell-cell

communication, with 39% of the cells showing no communication at all, and 57% of the cells

showing a very low communication rate ($R < 0.02$ s^{-1}) (Fig. 4A). To investigate gating ability of

septal junctions, the cells were treated with the protonophore CCCP which induces closure of

SJ[8,36]. Incubation with CCCP caused almost complete halt of communication (86% of the cells)

in the WT and in the Δ*fraI* mutant (77%) (Fig. 4A). The recovery rates were calculated for

"normal recovery" and "slow increase" cells. Here, recovery rates in vegetative filaments are

0.053 s^{-1} and 0.012 s^{-1} for WT and Δ*fraI* mutant respectively.

Figure 4: The Δ*fraI* mutant is highly impaired in cell-cell communication in both vegetative filaments
and hormogonia. CCCP-FRAP analysis of *Anabaena* PCC 7120, *N. punctiforme* WT (*N. pun* WT) and
Δ*fraI* mutant vegetative filaments **(A)** and hormogonia **(B)**. FRAP-responses of calcein-stained untreated
or CCCP-treated (90 min, 50 µm) were assigned to one of four groups indicated by the color scheme.
The numbers within the bars indicate the number of analyzed cells (n) from different filaments. Recovery
rates (*R*) of communicating cells **(C)**. Mean is shown. One-way ANOVA followed by Brown-Forsythe
multiple comparison test with the WT was performed. The numbers above the mean indicate the number
of cells used for calculation of *R. horm* – hormogonia. *N.pun WT horm* * excludes cells in hormogonia
of the WT with measured R > 0.19 (see main text for explanation). Cumulated results from at least two
independent cultures are shown. **, p<0.01****, p<0.0001; ns, not significant

Since FRAP experiments haven't yet been described in hormogonia, we wanted to measure

the rate of communication between the cells in hormogonia and the cell's ability to close septal

junctions upon treatment with the protonophore CCCP (Fig. 4B,C). On average, WT

hormogonia showed much higher rates of communication ($R = 0.104$ s^{-1}) compared to

vegetative cells ($R = 0.054$ s^{-1}) (Fig. 4C), with a fraction of the cells (23%) showing very high

recovery rates ($R > 0.19$ s^{-1}). Many cells undergo cell-division more or less simultaneously

during hormogonia development and hence in many cases the new septum may have not yet

completely closed, explaining the large number of cells with such unusual high communication

rates in FRAP analysis [45,46]. This idea is also supported by the fact that only 55% of the WT

hormogonia cells completely stopped communication after CCCP treatment, in contrast to 86%

among vegetative cells (Fig. 4B). Also, 76% of Δ*fraI* mutant hormogonia cells showed a

communicating phenotype, compared to 61% of the vegetative cells, because, like in the WT,
cell division was not yet completed and the fluorescent dye could easily diffuse between the
future daughter cells (Fig. 4B).

The inability of the Δ*fraI* mutant to differentiate heterocysts can be explained by the lack of cell-
cell communication. In contrast, the Δ*fraI* mutant can still differentiate akinetes despite
decreased cell-cell communication". This is in line with our previous observation with the
related species *Trichormus* (*Anabaena*) *variabilis* ATCC29413. There, we showed that during
akinete formation molecular transfer decreases and seems not to play a role in this spore
differentiation process[47].

The Δ*fraI* mutant lacks the septal nanopore array

To investigate whether aberrant nanopore array formation is causing the impaired
communication in the Δ*fraI* mutant, the septal peptidoglycan of vegetative cells and
hormogonia of the WT and the mutant were isolated and analyzed by TEM. The Δ*fraI* mutant
showed very few or almost no nanopores, both in PG from vegetative filaments as well as from
hormogonia (Fig. 5A,B). Interestingly, the Δ*fraI* mutant also had a significantly larger septa
diameter of vegetative filaments compared to the WT (Fig. 5B). A similar phenotype, namely
the lack of nanopores in septa and enlarged septa was previously observed for an *amiC2*
mutant of *N. punctiforme*[35]. Interestingly, the *amiC2* mutant displayed delocalized septa
leading to a severe morphological phenotype[12], which is not observed for the Δ*fraI* mutant
using Vancomycin-FL (Figure S4). In contrast, the pores in the lateral wall just around the
septal discs, were present in the Δ*fraI* mutant and the WT (Fig. 5A,B), implying that FraI is not
involved in formation of these lateral pores. In the center of the septal discs of the WT and
Δ*fraI* mutant hormogonia, we could occasionally observe few very large nanopores (Figure
S5), which may also explain higher rates of communication measured in some hormogonia
cells (see above).

Figure 5: The Δ*fraI* mutant completely lacks the septal nanopore array and has greater septa diameter. Septal peptidoglycan in *N. punctiforme* WT and Δ*fraI* mutant vegetative filaments **(A)** and hormogonia **(B)**. Representative TEM images are shown. Scale bar – 200 nm. **(C)** Number of nanopores and septum diameter of *N. punctiforme* vegetative and hormogonia cells in WT and Δ*fraI* mutant, which has very few or no nanopores and a greater septa diameter. One-way ANOVA followed by Brown-Forsythe multiple comparison test. Median is shown. $p<0.05$ ** $p<0.01$, **** $p<0.0001$, ns $p>0.05$.

Despite almost complete lack of nanopores, the AmiC2 protein could be detected in the septa of Δ*fraI* mutant using anti-AmiC2 antibodies (Figure S3). Hence, FraI is unlikely to be involved in septal localization of AmiC2. Moreover, FraI is predicted to have three transmembrane helixes with a cytoplasmic C-terminus and an N-terminal signal sequence of just 20 amino acids residing in the periplasm (Phobius), which makes a direct protein-protein interaction with the periplasmic amidase unlikely. However, at the moment we cannot exclude that a regulatory protein, like the LytM factor, is influenced by FraI. In line with the idea of a FraI-AmiC2 relation is the fact that *fraI* expression is the highest during hormogonia development (Figure S1), when many cells are undergoing cell division and the need for AmiC2 activity is the highest[48]. The lethality of overexpressed FraI (see above), is another indication that this protein is involved in regulation of the cell wall lytic enzyme AmiC, since the over-expression of the *amiC* orthologues in *Nostoc* sp. PCC 7120 was also lethal[13].

It is interesting that the impaired cell-cell communication rate in the Δ*fraI* mutant did not prevent hormogonia development. However, the motility of the hormogonia was reduced (Fig. 1A). Nevertheless, the Δ*fraI* mutant can move towards the source of light, indicating that cell-cell

communication through the septal junctions is not crucial for hormogonia motility and
synchronized movement along the filament in response to a positive external signal.
Additionally, the Δ*fraI* mutant was able to change the direction of movement following a
change of light conditions (data not shown). Therefore, there may exist another way of
communication, perhaps through the common periplasm, which enables synchronized
movement of the filament. As an alternative explanation, each single cell may decide the
direction of the movement based on the sensing of external factors such as light and
chemoattractant gradient. It has already been speculated that a change in an electromotive
force may be a signal for translocation of HmpF protein of the pili machinery to the opposite
side of the cell, which may be sensed by each cell in hormogonia[49]. Furthermore, the observed
reduction in motility in the Δ*fraI* mutant could be attributed to the diminished presence of the
exopolysaccharide sheath surrounding hormogonia filaments (Fig. 3). Previous studies have
demonstrated reductions in motility in mutants with impaired exopolysaccharide synthesis and
export pathways [32,34]. Why the absence of *fraI* gene leads to lower presence of hormogonia-
specific exopolysaccharide remains unknown. It has been speculated that HPS is exported
through the periplasm close to the cell junctions in hormogonia[50].

Alternatively, despite being significantly impaired in communication, 81% of Δ*fraI* hormogonia
cells (Fig. 4) still exhibit some communication, possibly facilitated by the presence of remaining
nanopores or incomplete septa formation. This residual communication may still be sufficient
for the transport of signal necessary for the uniform polar distribution of pili machinery enabling
directionality of movement along the filament in Δ*fraI* mutant.

Conclusion

In this study we discovered FraI, a new player in cell-cell communication and differentiation.
Mutation of *fraI* leads to a low-communicating phenotype in both vegetative cells and
hormogonia of *Nostoc punctiforme,* which can be explained by a nearly complete absence of
nanopores in the septa of neighboring cells. Despite the low levels of communication, the Δ*fraI*
strain forms motile hormogonia capable of synchronized movement, albeit in a different way
than the WT. The filaments also preform phototaxis, suggesting that motility and directionality
of movement do not require communication through the septal junctions. Lower motility
observed in Δ*fraI* is more likely to be explained by the absence of the HPS sheath. In contrast,
the presence of filaments containing a combination of vegetative and hormogonium cells in the
Δ*fraI* mutant, a phenotype never observed in the wild type, implies that communication through
the septal junctions may play a critical role in coordinating the development of hormogonia. As
in other mutants of septal nanopore array formation, severely impaired cell-cell communication
of the Δ*fraI* mutant is affecting heterocyst differentiation and function and hence diazotrophic

growth. Nonetheless, akinete development is still possible in the Δ*fraI* mutant. FraI may play a role in indirectly controlling amidase activity through an unknown mechanism. Additional experiments focusing on the FraI protein will be necessary to support this claim. Finally, characterization of orthologues of known septal junction proteins from *Nostoc* sp. 7120 in *N. punctiforme* will provide additional understanding of the role of septal junction-mediated communication in the process of hormogonium development and motility.

Author contribution

AJ: designed and performed experiments, analyzed and interpreted data, drafted the work and wrote the manuscript.

CM: performed TEM experiments and sample preparation for ultra-thin sections.

DDR: designed and performed experiments, analyzed and interpreted data, made considerable contributions to the design of the study, interpreted data, wrote the manuscript.

GAP: designed and performed experiments, analyzed and interpreted data

IM: designed and supervised the research in Tübingen, made considerable contributions to the design of the study, interpreted data, revised the manuscript critically.

All authors approved the final manuscript.

Data Availability

Data available from authors upon reasonable request.

Acknowledgment

Conflict of Interest

The authors declare no competing interests.

References

1.   Zeng, X. & Zhang, C. C. The Making of a Heterocyst in Cyanobacteria. *Annu Rev Microbiol* **76**, 597–618 (2022).

2.   Herrero, A., Stavans, J. & Flores, E. The multicellular nature of filamentous heterocyst-forming cyanobacteria. *FEMS Microbiol Rev* **40**, 831–854 (2016).

3.   Sukenik, A., Rücker, J. & Maldener, I. Dormant Cells (Akinetes) of Filamentous Cyanobacteria Demonstrate a Great Variability in Morphology, Physiology, and Ecological Function. in *Cyanobacteria: From Basic Science to Applications* 65–77 (Academic Press, 2019). doi:10.1016/B978-0-12-814667-5.00004-0.

4. Garg, R. & Maldener, I. The Formation of Spore-Like Akinetes: A Survival Strategy of
Filamentous Cyanobacteria. *Research Article Microb Physiol* **31**, 296–305 (2021).

5. Maldener, I., Summers, M. L. & Sukenik, A. Cellular differentiation in filamentous
cyanobacteria. in *Cell Biology of Cyanobacteria* (eds. F. Enrique & H. Antonia) 263–291
(Caister Academic Press, Norfolk, 2014). doi:10.21775/9781910190739.

6. Flores, E., Nieves-Morión, M. & Mullineaux, C. W. Cyanobacterial Septal Junctions: Properties
and Regulation. *Life* **9**, 1–14 (2018).

7. Kieninger, A.-K. & Maldener, I. Cell–cell communication through septal junctions in
filamentous cyanobacteria. *Curr Opin Microbiol* **61**, 35–41 (2021).

8. Weiss, G. L., Kieninger, A. K., Maldener, I., Forchhammer, K. & Pilhofer, M. Structure and
Function of a Bacterial Gap Junction Analog. *Cell* **178**, 374-384.e15 (2019).

9. Kieninger, A. K., Tokarz, P., Janović, A., Pilhofer, M., Weiss, G. L. & Maldener, I. SepN is a
septal junction component required for gated cell–cell communication in the filamentous
cyanobacterium Nostoc. *Nat Commun* **13**, (2022).

10. Merino-Puerto, V., Schwarz, H., Maldener, I., Mariscal, V., Mullineaux, C. W., Herrero, A. &
Flores, E. FraC/FraD-dependent intercellular molecular exchange in the filaments of a
heterocyst-forming cyanobacterium, *Anabaena sp. Mol Microbiol* **82**, 87–98 (2011).

11. Ramos-León, F., Arévalo, S., Mariscal, V. & Flores, E. Specific mutations in the permease
domain of septal protein SepJ differentially affect functions related to multicellularity in the
filamentous cyanobacterium *Anabaena*. *Microbial Cell* **5**, 555 (2018).

12. Lehner, J., Zhang, Y., Berendt, S., Rasse, T. M., Forchhammer, K. & Maldener, I. The
morphogene AmiC2 is pivotal for multicellular development in the cyanobacterium *Nostoc*
punctiforme. *Mol Microbiol* **79**, 1655–1669 (2011).

13. Berendt, S., Lehner, J., Zhang, Y. V., Rasse, T. M., Forchhammer, K. & Maldener, I. Cell Wall
Amidase AmiC1 Is Required for Cellular Communication and Heterocyst Development in the
Cyanobacterium *Anabaena* PCC 7120 but Not for Filament Integrity. *J Bacteriol* **194**, 5218–
5227 (2012).

14. Nieves-Morión, M., Mullineaux, C. W. & Flores, E. Molecular Diffusion through Cyanobacterial
Septal Junctions. *mBio* (2017) doi:10.1128/mBio.01756-16.

15. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in
filamentous cyanobacteria. *Nat Rev Microbiol* **8**, 39–50 (2010).

16. Vermassen, A., Leroy, S., Talon, R., Provot, C., Popowska, M. & Desvaux, M. Cell wall
hydrolases in bacteria: Insight on the diversity of cell wall amidases, glycosidases and
peptidases toward peptidoglycan. *Front Microbiol* **10**, 418687 (2019).

17. Büttner, F. M., Faulhaber, K., Forchhammer, K., Maldener, I. & Stehle, T. Enabling cell-cell
communication via nanopore formation: structure, function and localization of the unique cell
wall amidase AmiC2 of *Nostoc punctiforme*. *FEBS* **283**, 1336–1350 (2016).

18. Zheng, Z., Omairi-Nasser, A., Li, X., Dong, C., Lin, Y., Haselkorn, R. & Zhao, J. An amidase is
required for proper intercellular communication in the filamentous cyanobacterium
Anabaena sp. PCC 7120. *PNAS* **114**, E1405–E1412 (2017).

19.   Bornikoel, J., Staiger, J., Madlung, J., Forchhammer, K. & Maldener, I. LytM factor Alr3353
535   affects filament morphology and cell-cell communication in the multicellular cyanobacterium
536   *Anabaena sp.* PCC 7120. *Mol Microbiol* **108**, 187–203 (2018).

537   20.   Rai, A. N., Söderbäck, E. & Bergman, B. Cyanobacterium - Plant symbioses. *New Phytologist*
538   **147**, 449–481 (2000).

539   21.   Álvarez, C., Navarro, J. A., Molina-Heredia, F. P. & Mariscal, V. Endophytic Colonization of Rice
540   (*Oryza sativa* L.) by the Symbiotic Strain *Nostoc punctiforme* PCC 73102. *Mol Plant Microbe*
541   *Interact* **33**, (2020).

542   22.   Zuniga, E. G., Boateng, K. K. A., Bui, N. U., Kurnfuli, S., Muthana, S. M. & Risser, D. D.
543   Identification of a hormogonium polysaccharide-specific gene set conserved in filamentous
544   cyanobacteria. *Mol Microbiol* **114**, 597–608 (2020).

545   23.   Cho, Y. W., Gonzales, A., Harwood, T. V., Huynh, J., Hwang, Y., Park, J. S., Trieu, A. Q., Italia, P.,
546   Pallipuram, V. K. & Risser, D. D. Dynamic localization of HmpF regulates type IV pilus activity
547   and directional motility in the filamentous cyanobacterium *Nostoc punctiforme*. *Mol Microbiol*
548   **106**, 252–265 (2017).

549   24.   Harwood, T. V., Zuniga, E. G., Kweon, H. J. & Risser, D. D. The cyanobacterial taxis protein
550   HmpF regulates type IV pilus activity in response to light. *Proc Natl Acad Sci U S A* **118**, 1–12
551   (2021).

552   25.   Parrett, G. A., Brones, P. D., Jenkins, G. M., Mounts, S. M., Nguyen, A. & Risser, D. D.
553   Identification of a morphogene required for tapered filament termini in filamentous
554   cyanobacteria. *Microbiology (N Y)* **169**, (2023).

555   26.   Allen, M. B. & Arnon, D. I. Studies on Nitrogen-Fixing Blue-Green Algae. I. Growth and
556   Nitrogen Fixation by *Anabaena Cylindrica* Lemm. *Plant Physiol* **30**, 366–372 (1955).

557   27.   Rippka, E., Deruelles, J. & Waterbury, N. B. Generic Assignments, Strain Histories and
558   Properties of Pure Cultures of Cyanobacteria. *J Gen Microbiol* **111**, 1–61 (1979).

559   28.   Black, T. A., Cai, Y. & Wolk, C. P. Spatial expression and autoregulation of hetR, a gene
560   involved in the control of heterocyst development in *Anabaena*. *Mol Microbiol* **9**, 77–84
561   (1993).

562   29.   Risser, D. D., Wong, F. C. Y. & Meeks, J. C. Biased inheritance of the protein PatN frees
563   vegetative cells to initiate patterned heterocyst differentiation. *Proc Natl Acad Sci U S A* **109**,
564   15342–15347 (2012).

565   30.   Khayatan, B., Bains, D. K., Cheng, M. H., Cho, Y. W., Huynh, J., Kim, R., Omoruyi, O. H., Pantoja,
566   A. P., Park, J. S., Peng, J. K., Splitt, S. D., Tian, M. Y. & Risser, D. D. A putative O-linked β-
567   Nacetylglucosamine transferase is essential for hormogonium development and motility in
568   the filamentous cyanobacterium *Nostoc punctiforme*. *J Bacteriol* **199**, 1–14 (2017).

569   31.   Wolkt, C. P., Cait, Y. & Panoff, J.-M. Use of a transposon with luciferase as a reporter to
570   identify environmentally responsive genes in a cyanobacterium. *Proc. Natl. Acad. Sci. USA* **88**,
571   5355–5359 (1991).

572   32.   Risser, D. D. & Meeks, J. C. Comparative transcriptomics with a motility-deficient mutant leads
573   to identification of a novel polysaccharide secretion system in *Nostoc punctiforme*. *Mol*
574   *Microbiol* **87**, 884–893 (2013).

575   33.   Splitt, S. D. & Risser, D. D. The non-metabolizable sucrose analog sucralose is a potent
576   inhibitor of hormogonium differentiation in the filamentous cyanobacterium *Nostoc*
577   *punctiforme*. *Arch Microbiol* **198**, 137–147 (2016).

578   34.   Khayatan, B., Meeks, J. C. & Risser, D. D. Evidence that a modified type IV pilus-like system
579   powers gliding motility and polysaccharide secretion in filamentous cyanobacteria. *Mol*
580   *Microbiol* **98**, 1021–1036 (2015).

581   35.   Lehner, J., Berendt, S., Dörsam, B., Pérez, R., Forchhammer, K. & Maldener, I. Prokaryotic
582   multicellularity: A nanopore array for bacterial cell communication. *FASEB Journal* **27**, 2293–
583   2300 (2013).

584   36.   Kieninger, A. K., Tokarz, P., Janović, A., Pilhofer, M., Weiss, G. L. & Maldener, I. SepN is a
585   septal junction component required for gated cell–cell communication in the filamentous
586   cyanobacterium *Nostoc*. *Nature Communications 2022 13:1* **13**, 1–15 (2022).

587   37.   Perez, R., Forchhammer, K., Salerno, G. & Maldener, I. Clear differences in metabolic and
588   morphological adaptations of akinetes of two Nostocales living in different habitats.
589   *Microbiology (N Y)* **162**, 214–223 (2016).

590   38.   Fiedler, G., Arnold, M., Hannus, S. & Maldener, I. The DevBCA exporter is essential for
591   envelope formation in heterocysts of the cyanobacterium *Anabaena* sp. strain PCC 7120. *Mol*
592   *Microbiol* **27**, 1193–1202 (1998).

593   39.   Risser, D. D., Chew, W. G. & Meeks, J. C. Genetic characterization of the hmp locus, a
594   chemotaxis-like gene cluster that regulates hormogonium development and motility in *Nostoc*
595   *punctiforme*. *Mol Microbiol* **92**, 222–233 (2014).

596   40.   Kieninger, A.-K. Identification and Characterization of the Cell-Cell Communication System of a
597   Multicellular Cyanobacterium. Dissertation. (University of Tübingen, Tübingen, 2021).

598   41.   Gonzalez, A., Riley, K. W., Harwood, T. V, Zuniga, E. G., Risser, D. D. & Ellermeier, C. D. A
599   Tripartite, Hierarchical Sigma Factor Cascade Promotes Hormogonium Development in the
600   Filamentous Cyanobacterium *Nostoc punctiforme*. *mSphere* **4**, 1–15 (2019).

601   42.   Harwood, T. V & Risser, D. D. The primary transcriptome of hormogonia from a filamentous
602   cyanobacterium defined by cappable-seq. *Microbiology (N Y)* **167**, 1–9 (2021).

603   43.   Nürnberg, D. J., Mariscal, V., Bornikoel, J., Nieves-Morión, M., Krauß, N., Herrero, A.,
604   Maldener, I., Flores, E. & Mullineaux, C. W. Intercellular diffusion of a fluorescent sucrose
605   analog via the septal junctions in a filamentous cyanobacterium. *mBio* **6**, 1–12 (2015).

606   44.   Mullineaux, C. W., Mariscal, V., Nenninger, A., Khanum, H., Herrero, A., Flores, E. & Adams, D.
607   G. Mechanism of intercellular molecular exchange in heterocyst-forming cyanobacteria.
608   *EMBO Journal* **27**, 1299–1308 (2008).

609   45.   Meeks, J. C. & Elhai, J. Regulation of Cellular Differentiation in Filamentous Cyanobacteria in
610   Free-Living and Plant-Associated Symbiotic Growth States. *Microbiology and Molecular*
611   *Biology Reviews* **66**, 94–121 (2002).

612   46.   Maldener, I., Summers, M. L. & Sukenik, A. *Cellular Differentiation in Filamentous*
613   *Cyanobacteria in The Cell Biology of Cyanobacteria*. (Caister Academic Press, 2014).
614   doi:10.21775/9781910190739.

47. Garg, R., Luckner, M., Berger, J., Hipp, K., Wanner, G., Forchhammer, K. & Maldener, I.
Changes in Envelope Structure and Cell–Cell Communication during Akinete Differentiation
and Germination in Filamentous Cyanobacterium *Trichormus variabilis* ATCC 29413. *Life* **12**,
1–12 (2022).

48. Lehner, J., Berendt, S., Dörsam, B., Pérez, R., Forchhammer, K. & Maldener, I. Prokaryotic
multicellularity: a nanopore array for bacterial cell communication. *The FASEB Journal •*
Research Communication **27**, 2293–2300 (2013).

49. Harwood, T. V., Zuniga, E. G., Kweon, H. J. & Risser, D. D. The cyanobacterial taxis protein
HmpF regulates type IV pilus activity in response to light. *Proc Natl Acad Sci U S A* **118**, 1–12
(2021).

50. Hoiczyk, E. & Baumeister, W. The junctional pore complex, a prokaryotic secretion organelle,
is the molecular motor underlying gliding motility in cyanobacteria. *Current Biology* **8**, 1161–
1168 (1998).

629