The role of FraI in cell-cell communication and differentiation in the

hormogonia-forming cyanobacterium Nostoc punctiforme

Ana Janović [1], Iris Maldener [1,*], Claudia Menzel [1], Gabriel A. Parrett [2], Douglas D. Risser [2,*]

[1] Interfaculty Institute of Microbiology and Infection Medicine Tübingen, Organismic Interactions,

University of Tübingen, Auf der Morgenstelle 28, 72076 Tübingen, Germany

[2] Department of Biology, University of Colorado, Colorado Springs, USA

[*] Correspondence: drisser@uccs.edu; iris.maldener@uni-tuebingen.de

Keywords

12 Cell-cell communication, hormogonia, filamentous cyanobacteria, *Nostoc punctiforme,*

13 motility, septal junctions, NpF4142, FraI

Abstract

Multicellular cyanobacteria, like *Nostoc punctiforme,* rely on septal junctions for cell-cell

communication, which is crucial for coordinating various physiological processes including

differentiation of N_2-fixing heterocysts, spore-like akinetes and hormogonia - short, motile

filaments important for dispersal. In this study we functionally characterize a protein, encoded

by gene *NpF4142*, which in a random mutagenesis approach, initially showed a motility-

related function. The reconstructed *NpF4142* knockout mutant exhibits further distinct

phenotypic traits, including altered hormogonia formation with significant reduced motility,

inability to differentiate heterocysts and filament fragmentation. For that reason, we named

the protein FraI (fragmentation phenotype). The mutant displays severely impaired cell-cell

communication, due to almost complete absence of the nanopore array in the septal cell

wall, which is an essential part of the septal junctions. Despite lack of communication,

hormogonia in the ∆*fraI* mutant maintain motility and phototactic behaviour, even though less

pronounced than the wild type. This suggests an alternative mechanism for coordinated

movement beyond septal junctions. Our study underscores the significance of FraI in

nanopore formation and cell differentiation and provides additional evidence for the

importance of septal junction formation and communication in various differentiation traits of

cyanobacteria. The findings contribute to a deeper understanding of the regulatory networks

governing multicellular cyanobacterial behaviour, with implications for broader insights into

microbial multicellularity.

Importance

The filament-forming cyanobacterium *Nostoc punctiforme* serves as a valuable model for studying cell differentiation, including the formation of nitrogen-fixing heterocysts and hormogonia. Hormogonia filaments play a crucial role in dispersal and plant colonization, providing a nitrogen source through atmospheric nitrogen fixation, thus holding promise for fertilizer-free agriculture. The coordination among the hormogonia cells enabling uniform movement towards the positive signal remains poorly understood. This study investigates the role of septal junction-mediated communication in hormogonia differentiation and motility, by studying a Δ*fraI* mutant with significantly impaired communication. Surprisingly, impaired communication does not abolish synchronized filament movement, suggesting an alternative coordination mechanism. These findings deepen our understanding of cyanobacterial biology and have broader implications for multicellular behavior in prokaryotes.

Introduction

The filamentous cyanobacterium *Nostoc punctiforme* ATCC 29133/PCC73102 is a true multicellular bacterium, able to form specialized cells which enable the bacterium to adapt to various environmental conditions. Heterocysts are nitrogen-fixing cells that differentiate from semi-regularly positioned vegetative cells upon nitrogen starvation to provide a microoxic environment for the oxygen-labile nitrogenase[1,2]. Hormogonia (see below) are short filaments specialized as motile units allowing dispersal. Finally, unicellular akinetes form under various stress conditions enabling survival as spore-like resistant cells [3–5].

Within a filament, cells interact through a remarkable cell-cell communication system, involving so called septal junctions. These multi-protein complexes connect the individual cells by traversing the septal peptidoglycan layer through nanopores[6,7]. On each cytoplasmic side, every septal junction contains a flexible cap and plug structure by which the intercellular connection closes in a reversible manner. This gating mechanism of halting communication is crucial under stress conditions, such as UV-C light exposure, cytoplasm leakage due to filament breakage, or the isolation of senescent cells from the rest of the filament [7–9]. Mutants with impaired cell-cell communication also exhibit deficiencies in heterocyst development[1,10–14], emphasizing the significance of intercellular signaling in cellular differentiation [1,2,10–15].

Formation of septal nanopores requires the activity of cell wall amidases[16]. In unicellular bacteria amidases play a pivotal role in cell division, by cleaving the newly synthesized peptidoglycan to facilitate separation of daughter cells [17]. In multicellular cyanobacteria, cell wall amidases play a role in cell-cell communication, by perforating the septal peptidoglycan[13,18,19]. A knockout mutant of the AmiC2 enzyme in *N. punctiforme* showed

severe impairment in cell division, filament morphology and cell-cell communication, caused

by the complete absence of nanopores in the septal peptidoglycan[16]. The *amiC2* mutant is

unable to differentiate heterocysts, hormogonia and akinetes[12]. The AmiC2 protein primarily

localizes along the septal plane of nascent septa and is typically absent in fully developed

septa [12]. AmiC1, another amidase encoded in the same gene operon, is likely essential for

Nostoc punctiforme, since the *knockout* of this gene was reported to be impossible to create

[12,20]. In the closely related cyanobacterium *Nostoc* sp. PCC 7120 mutants in both

orthologues, *amiC1* and *amiC2* as well as double *amiC1amiC2* knockouts were created. All

mutants exhibited aberrant nanopore shape and numbers and lower rates of cell-cell

communication compared to the wild type[13,21]. Furthermore, overexpression of AmiC1 in

Nostoc sp. 7120 resulted in a lethal phenotype, in contrast to the AmiC2 overexpression,

which has no significant effect on growth. The LytM factor Alr3353 is an activator of AmiC1 of

Nostoc sp. 7120 and its absence leads to a reduction in nanopore numbers and the rate of

cell-cell communication[22]. Recently, transcriptional factor FurC was identified as direct

regulator of *amiC1* and *amiC2* genes and further proteins of cell wall related proteins[19].

Transiently formed hormogonia, 10-20 cell-long filaments of *Nostoc punctiforme*, play a

crucial role in dispersal and symbiotic plant colonization [23]. After colonization of the host

plant, the hormogonia develop back to vegetative filaments containing heterocysts for N_2

fixation. This symbiosis is of potential agricultural importance as it enables fertilizer-free

cultivation providing nitrogen sources for the crop, as shown for *Oryza sativa* [24]. Induction of

hormogonia differentiation from vegetative filaments can be triggered by various signals such

as nitrogen stepdown, red-light exposure or HIF (*hormogonium-inducing factor*) exposure.

Following the induction signal, vegetative cells differentiate into hormogonia within 18-24

hours through rapid cell divisions not accompanied by overall biomass growth. These

filaments demonstrate photo and chemo-tactic movement, facilitated by a type IV pili

machinery on their outer cell's surface. The movement is additionally supported by the export

of an exopolysaccharide sheath known as hormogonia polysaccharide (HPS) [25].

Directionality is determined by the presence of the HmpF protein at one pole of each cell,

positioned uniformly at the leading poles of each cell in motile filaments. Repositioning of the

HmpF protein occurs rapidly and synchronously along the filament, typically within about 30

seconds, in response to changes in light conditions, leading to a change in the filament's

course [26,27]. A putative amidase encoded by *the tftA* gene was shown to be important for the

tapered cell phenotype at the filament's termini, likely by the degradation of the terminal

peptidoglycan. Further, a *tftA* knockout showed reduced migration ability through agar [28].

However, the role of cell-cell communication in the synchronized movement of the entire

filament and during the change of movement's direction in hormogonia remained

uninvestigated. In this study, we identified a mutant lacking the NpF4142 protein, herein

designated FraI, that presents decreased hormogonia motility. Characterization of this mutant strain allowed us to show that FraI plays a key role in nanopore formation and hormogonia differentiation. Furthermore, this study showed that transmission of the signal for direction of movement in hormogonia probably does not involve communication through the septal junctions.

Material and methods

Cultivation of bacterial strains

As standard medium, *N. punctiforme* and the mutant strains were cultivated in AA/4 medium according to Allen and Arnon [29] in a fourfold dilution. To supply a source of fixed nitrogen, the medium was supplemented with 2.5 mM NH_4Cl and 5 mM MOPS buffer, pH 7.8. In experiments performed in Tübingen the cyanobacteria were cultivated in either liquid BG11 medium [30] or AA/4 medium supplemented with 2.5 mM KNO_3 and 2.5 mM $NaNO_3$. Liquid cultures were shaken at 120 rpm, agar plates contained 1% Difco agar. Growth occurred at 28 °C with constant illumination at 25-40 $\mu E\ m^{-2}\ s^{-1}$. Sucralose was added to suppress hormogonia development in concentrations mentioned in the text. Neomycin was used at a concentration of 50 μg/ml in appropriate strains.

Plasmid and strain construction

For a detailed description of the plasmids, strains, and oligonucleotides used in this study refer to supplemental material, Tables S1. All constructs were sequenced to insure fidelity.

To construct plasmid pDDR558 for in-frame deletion of *fraI* (Npun_F4142), approximately 900 bp of flanking DNA on either side of the gene and several codons at the beginning and end of the gene were amplified via overlap extension PCR using primers NpF4142-5'-F, NpF4142-5'-R, NpF4142-3'-F and NpF4142-3'-R and cloned into pRL278[31] as a BamHI-SacI fragment using restriction sites introduced on the primers.

To construct plasmid pGAP109 for replacement of the chromosomal allele of *fraI* with a C-terminal *gfpuv*-tagged variant, approximately 900 bp of DNA downstream of the stop codon were amplified via PCR using primers NpF4142-gfp-3'-F and NpF4142-3'-R and cloned into pSCR569 [32], as an SpeI-SacI fragment using restriction sites introduced on the primers. The coding region and approximately 900 bp of DNA upstream of the start codon were then amplified via PCR using primers NpF4142-5'-F and NpF4142-gfp-5'-R and cloned into this plasmid as a BamHI-SmaI fragment using restriction sites introduced on the primers.

Generation of transposon mutants and identification of transposon insertion sites was performed as previously described [33] using plasmid pRL1063a[34]. Gene deletions and allelic replacements were performed as previously described[35] with *N. punctiforme* cultures

142   supplemented with 4 mM sucralose to inhibit hormogonium development and enhance

143   conjugation efficiency [33,36]. To construct UCCS103, plasmid pDDR558 was introduced into

144   wild type *N. punctiforme*. To create UCCS113, plasmid pGAP109 was introduced into

145   UCCS103.

Motility and phototaxis assays

147   Plate and time-lapse motility and phototaxis assays were performed as previously

148   described[37] and imaged using a Leica SD9 dissecting microscope equipped with a Flexcam

149   C3 camera and controlled by Leica X LAS X software. The students T-test was applied to

150   determine statistical significance in colony spreading assays.

Light- and fluorescence microscopy

A Leica DM2500 B microscope equipped with a Leica DFC420C camera, or an Evos M5000

microscope was employed for light microscopy. Fluorescence microscopy utilized a Leica

DM5500 B microscope with a 100x/1.3 oil objective lens, connected to a Leica DFC360FX

camera. GFP and chlorophyll fluorescence were induced using BP470/40 nm or

BP535/50 nm filters, and emission was detected with BP525/50 nm or BP610/75 nm filters,

respectively.

Septal visualization was performed using Vancomycin-FL (Bodipy-FL, Invitrogen) fluorescent

stains that specifically stains septal peptidoglycan by the previously described procedure[18].

FRAP and CCCP-FRAP

A three-day-old liquid culture of vegetative filaments from both the *N. punctiforme* wild type

and the Δ*fraI* mutant strains was utilized for CCCP-FRAP, following previously established

protocols [8,9]. Briefly, 2 ml of culture underwent triple washing with BG11 medium and were

then resuspended in 500 µL of medium. Subsequently, the cells were incubated with 10 – 15

µL of 1 mg/ml calcein in DMSO (Sigma-Aldrich) for 90 minutes in the dark at 28 °C with

shaking. Following this incubation period, the cells underwent three additional washes with

BG11 medium and were then either resolved in medium alone or in medium containing

CCCP (from a 50 mM stock solution in DMSO), achieving a final concentration of 50 µM in

the culture. After another 90-minute incubation with shaking at 28 °C in the dark, the

filaments were spotted onto BG11 agar plates (1%).

The FRAP procedure was conducted by employing a laser with 488 nm excitation at 0.2%

intensity on a Zeiss LSM 800 confocal microscope equipped with a 63x/1.4 oil-immersion

objective. The ZEN 2.9 (blue edition) software was used for the experiment. Simultaneous

detection of calcein fluorescence emission (400–530 nm) and chlorophyll autofluorescence

emission (650–700 nm) was carried out. To bleach a specific cell, the laser intensity was

briefly increased to 3.5% after capturing 5 pre-bleached images. Subsequently, fluorescence
recovery in the bleached cell was observed at 1-second intervals over a period of 30–
120 seconds. Image analysis was performed using ImageJ (version 1.51j) and GraphPad
Prism 10, as detailed in previous descriptions.

For hormogonia FRAP, a log-phase culture grown with 4 mM sucralose in BG11 medium
was washed twice with BG11 for hormogonia induction. After 18-20 hours following
hormogonia induction the cells were harvested (3-5 ml) and used for FRAP as described
above. The stained cells (25 μL) were put onto a plain microscopic slide and covered with a
cover slip. The measurement was only conducted after the hormogonia ceased movement
upon becoming trapped between coverslip and slide due to evaporation.

Akinete induction

Akinetes were induced essentially as described[38]. In short, liquid cultures of WT and
NpF4142 mutant were grown in BG11 medium with nitrate until early log-phase. Akinete
formation was induced by washing the culture twice followed by inoculation in phosphate-free
BG11 medium and incubation with low shaking (50 rpm) for 4 weeks.

Isolation of septal peptidoglycan and transmission electron microscopy

Septal peptidoglycan was isolated as described[9]. First, vegetative filaments of *Nostoc*
punctiforme WT and Δ*fraI* grew for 5 days in liquid BG11 medium supplemented with 4 mM
sucralose to supress hormogonia development. For hormogonia induction, a log-phase
culture was induced by washing twice with BG11 medium and incubated in fresh BG11
medium for 20 h. The vegetative and hormogonia cultures were washed and resuspended in
700 μL 0.1 M Tris-HCl pH 6. Briefly, cells were sonicated (Branson Sonier 250), then boiled
in SDS for 30 min. Then, the samples were sonified in a water bath for 30 min, followed by
washing and incubation in 50 mM Na_3PO_4 pH 6.8 with 300 μf of α-chymotrypsin. The
chymotrypsin was readded and the next day the cells were again sonicated for 30-60 s.
Purified septa (10 μL) were added onto an UV-irradiated (16 h) formvar/carbon film-coated
copper grid (Electron Microscopy Sciences) for 30 min, stained with 1% (w/v) uranyl acetate
and imaged with a Philips Tecnai10 electron microscope at 80 kV equipped with a Rio
Camera (Gatan). Images were analysed using ImageJ 2.9. to calculate the number of
nanopores per septa and the septum's diameter.

Transmission electron microscopy of ultrathin sections

To examine hormogonia via transmission electron microscopy, we induced their formation by
washing of the cells previously cultivated in AA/4 media with 2.5 mM of sucralose. After 20
hours, hormogonia were harvested by centrifugation and ultrathin sections were prepared

following standard procedure[39]. Prior to sectioning, the cells underwent fixation with

glutaraldehyde and potassium permanganate. Subsequently, the ultrathin sections were

stained with uranyl acetate and lead citrate. Finally, the samples were analysed using a

Philips Tecnai10 microscope operating at 80 kV.

Immunolocalization of AmiC2

Immunolocalization of AmiC2 in *Nostoc* strains was performed based on two protocols[20,40].

Nostoc punctiforme cultures were cultivated in AA/4 medium with supplemented $NaNO_3$,

KNO_3 and 2.5 mM sucralose, washed once and resuspended in 1 mL PBS. The cells were

fixed with 4% paraformaldehyde PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM

KH_2PO_4, pH 7.4) on ice for one hour, followed by three rounds of washing with PBS and final

incubation in GTE buffer (50 mM glucose, 25 mM Tris-Cl, pH 8.0, 10 mM EDTA). Then,

200 µL of the cell suspension were put onto Epredia™TM Polysine adhesion slides (Epredia,

Netherlands) and let to dry. The cells were then fixed with -20 °C methanol for 5 min followed

by incubation with -20 °C acetone for 30 s. Cells were dried at RT and rehydrated with

200 µL PBS for 5 min. After blocking with 200 µL 2 % w/v BSA in PBS for 30 min, cells were

incubated overnight in a wet chamber at 4 °C with 1:5 anti-AmiC1 [20] in 2 % w/v BSA BSA-

PBS. The slides were washed three times with PBS, followed by a two-hour incubation at

room temperature in darkness with FITC-coupled α-rabbit antibodies (diluted at 1:200 in

BSA-PBS from Sigma Aldrich). After subsequent washing and drying, a drop of Vectashield

Mounting Medium H-1200 (Vector Laboratories, USA) was added. The preparation was

covered with a coverslip. Fluorescence imaging was conducted with a DM5500B Leica

microscope and a DFC360FX monochrome camera. Autofluorescence was measured as

mentioned above, while FITC-fluorescence was captured using a BP470/40 nm excitation

filter and a BP525/50 nm emission filter. Z-stacks with 0.20 µm intervals were acquired and

3D-deconvolution using Leica ASF built-in function was performed.

Test on significance

Statistical analysis was performed with GraphPad Prism version 10. Comparison of one

group with multiple other groups was performed via Welch ANOVA test followed by Brown-

Forsythe followed by Dunnett's T3 multiple comparison test. Significance *P* is indicated with

asterisks: *: *P*≤0.05; **: *P*≤0.01; ***: *P*≤0.001; ****: *P*≤0.0001; ns (not significant): *P*>0.05.

Results and Discussion

fraI deletion affects hormogonia motility, filament length and diazotrophy

Using a transposon mutagenic screen[33] we identified a mutant strain with reduced motility containing a transposon insertion in open reading frame Npun_F4142 (Table S1). Its homologue in *Nostoc sp.* 7120 that we previously named FraI[41] was shown to be highly abundant in a co-immunoprecipitation experiment against septal junction proteins FraD and SepN[9] and displayed a fragmentation phenotype[41] . Based on this, and the data presented below, we have designated the gene name for Npun_F4142 as *fraI.* To confirm that the transposon insertion in *fraI* was responsible for the motility defect, a strain with an in-frame deletion of *fraI* was constructed (Δ*fraI*). Compared to the wild type, the Δ*fraI* strain exhibited a significant reduction in colony spreading in plate motility assays (Fig. 1A). Moreover, there was a qualitative difference in the appearance of motile colonies for the Δ*fraI* strain, with filaments spreading in collective masses rather than dispersed filaments as observed in the wild type (Fig. 1A). In time-lapse motility assays of individual hormogonium filaments (Video S1) the Δ*fraI* strain exhibited motility of individual filaments, but the filaments were much shorter than those of the wild type, and many of the shortest filaments were non-motile.

Figure 1. Deletion of *fraI* affects motility. **(A)** Images of plate motility assays (left) and quantification of colony spreading (right) of strains as indicated. For images on left, black boxes in top panel correspond to closeup images of colony periphery shown in bottom panel. For quantification, n=3, error bars =+/- 1 S.D., *=p-value <0.01 vs wild type. **(B)** Images of phototaxis assays for strains as indicated. Red circle indicates the initial site of colony at beginning of assay. Direction of light indicated with arrow.

In an attempt to visualize the localization of FraI, a *fraI-gfp* allele was re-introduced at the native chromosomal locus of the Δ*fraI* strain. Introduction of *fraI-gfp* into the deletion strain restored wild-type motility indicating this allele is functional (Fig. 1A), and further demonstrating that deletion of *fraI* was responsible for the observed phenotype. However, no fluorescence signal could be visualized, indicating that the abundance of FraI-GFP in the cell

may be below the limit of detection, or that fusion of GFP to the C-terminus of FraI may affect
GFP folding or stability. Attempts were made to introduce either the native, or a *gfp*-tagged
allele of *fraI* into the deletion strain on a replicative shuttle vector, but in both cases these
efforts failed to yield viable colonies, possibly implying that increased expression of *fraI* from
a multicopy plasmid is lethal in *N. punctiforme*. Given the motility defect in the Δ*fraI* strain, its
capacity for phototaxis was also investigated (Fig. 1B). Although migration was reduced
compared to the wild type, the Δ*fraI* strain exhibited directional movement in response to
light, indicating the mutant strain retains the ability to perceive and respond to light signals.

In liquid cultures supplemented with ammonia, the Δ*fraI* strain was more dispersed than the
wild type (Fig. 2A). Under these standard growth conditions, the wild-type strain produces a
mixture of vegetative and hormogonium filaments (Fig. 2B). The Δ*fraI* strain also produced a
mixture of both filament types, but in general, the filaments were much shorter (Fig. 2B).
Moreover, we frequently observed what appeared to be hybrid filaments containing a mixture
of vegetative and hormogonium cell types in the same filament (Fig. 2B). Although difficult to
quantify given the tendency of *N. punctiforme* to aggregate into dense clumps, this
phenotype was never observed in the wild-type strain, nor have we ever observed such a
phenotype in previous studies with either the wild type or any of the mutants we have
characterized. These results indicate that while *fraI* is dispensable for hormogonium
development and motility, it may be critical for cells within a filament to synchronize

hormogonium differentiation.

Figure 2. Influence of *fraI* on diazotrophy and development. **(A)** Images of cultures (strains as
indicated) in presence (NH_4^+) or absence (N_2) of a source of fixed nitrogen. **(B)** Light micrographs of
individual filaments (strains as indicated) in presence (NH_4^+) or absence (N_2) of a source of fixed
nitrogen. V = vegetative filament, H = hormogonia, M = mixed filament containing both hormogonium
and vegetative cells, > = heterocysts. Bar = 10 um. **(C)** Light micrographs (strains as indicated) of

filaments following induction of akinetes by removal of phosphate from the growth medium. A =
akinete, Bar = 10 um.

In liquid cultures lacking a source of fixed nitrogen, conditions where the wild type develops
nitrogen-fixing heterocysts to support growth on N_2, the $\Delta fraI$ strain became chlorotic and
failed to grow (Fig. 2A). Under these conditions the fragmentation phenotype became more
severe, and there was no obvious indication that morphologically distinct heterocysts were
present (Fig. 2B). As with motility, the introduction of *fraI-gfp* into the deletion strain restored
diazotrophy (Fig. 2A). Upon phosphate starvation the $\Delta fraI$ strain was able to develop
akinetes although of aberrant shape and not in consecutive manner when compared to the
WT[38] (Fig. 2C). Collectively, the absence of heterocysts and fragmentation phenotype of the
$\Delta fraI$ mutant resembles that reported for mutants in other genes known to be involved in
formation of the septal junctions, most of which have been characterized primarily in the
closely related filamentous cyanobacterium *Nostoc* sp. PCC 7120.

The reduced motility phenotype motivated us to examine the ultrastructure of hormogonia
filaments using transmission electron microscopy (TEM). In $\Delta fraI$ we could observe short
hormogonia that were completely lacking the surrounding exopolysaccharide sheath (Fig. 3),
which may be the cause of the of observed reduced motility (Fig. 1). In addition, no septal
cell-cell connections were visible in $\Delta fraI$ hormogonia (Fig. 3). Furthermore, in contrast to the
WT, the thylakoid membranes were aberrantly arranged. This could be explained by the
presence of numerous glycogen granula, which may disturb the organization of the
membranes.

Figure 3. The hormogonia of Δ*fraI* have no cell-cell connections and are lacking the HPS. After induction of hormogonia, ultra-thin sections were prepared and the ultrastructure investigated by transmission electron microscopy. Shown are representative electron micrographs of each strain. Blue rectangles signify the zoomed region shown below (WT) or at the right (Δ*fraI*). Red arrows point to the cell-cell junctions visible in the WT. Yellow arrows point to the exopolysaccharide sheath present only in the WT hormogonia. G – glycogen granules, TM – thylakoid membrane. Scale bar – 1 µm.

FraI is related to known septal proteins

To further explore the relationship between FraI and septal junctions, we investigated the expression pattern of *fraI* as well as the orthologs of other septal junction genes during hormogonium and heterocyst development in *N. punctiforme* using previously published transcriptomic data sets[32,42,43]. Additionally, the evolutionary co-occurrence of this gene set in cyanobacteria was determined. During hormogonium development all the identified septal junction genes are transcriptionally upregulated in wild-type *N. punctiforme*, with expression typically peaking between 6 to 12 h post induction (Fig. S1A). This observation is consistent with the fact that developing hormogonia undergo a round of cell division, necessitating the construction of large numbers of new septal junctions at the new septa. For most of these

genes, enhanced transcription in hormogonia is independent of the hormogonium sigma

factor cascade[42], as deletion of *sigJ*, *sigC*, or *sigF* has no obvious effect on expression. In

contrast, upregulation of *fraI*, *sepI* and *sjcF1* in hormogonia is dependent on both *sigJ* and

sigC. Given that expression of *sigC* is also dependent on *sigJ*, these results imply that these

three genes are most directly regulated by *sigC*. Transcriptional start site (TSS) mapping via

Cappable seq (CAPseq) indicated the presence of two transcriptional start sites (-265 bp and

-200 bp of the *fraI* start codon) in the *fraI* promoter region, with the more distal closely

corresponding to total read coverage from RNAseq data (Fig. S1B). However, this TSS only

showed a very minor decrease in abundance in the Δ*sigC* strain. In developing heterocysts

there was less pronounced upregulation of this gene set, with only a few genes displaying

marked upregulation (Fig. S1A). Co-occurrence analysis indicated that *fraI* was found almost

exclusively in heterocyst-forming cyanobacteria, similar to the conservation pattern of *sepN*,

fraD, and *fraC*, although orthologs of these genes are found occasionally in non-heterocyst

forming filamentous cyanobacteria (Fig. S2). Overall, the expression and co-occurrence

patterns of *fraI* in comparison to other septal junction genes support a role for *fraI* in the

formation of septal junctions.

352 **The Δ*fraI* mutant is severely impaired in cell-cell communication**

The relation to septal junction proteins together with the Δ*fraI* phenotype [10,11,44] prompted us

to investigate cell-cell communication and septal junction gating ability in the Δ*fraI* mutant of

N. punctiforme[8,9]. The study of cell-cell communication was performed by FRAP experiments

[10,45], where filaments were first stained with a fluorescent dye, subsequently, a single cell is

laser-bleached, and finally the recovery of fluorescence, resulting from the diffusion of the

dye through septal junctions, was monitored over time[10,45]. Compared to the wild type of *N.*

punctiforme, the vegetative cells of the Δ*fraI* mutant were highly impaired in cell-cell

communication, with 39% of the cells showing no communication at all, and 57% of the cells

showing a very low communication rate ($R < 0.02$ s^{-1}) (Fig. 4A). To investigate gating ability

of septal junctions, the cells were treated with the protonophore CCCP which induces closure

of SJ [8,9]. Incubation with CCCP caused almost complete halt of communication (86% of the

cells) in the WT and in the Δ*fraI* mutant (77%) (Fig. 4A). The recovery rates were calculated

for "normal recovery" and "slow increase" cells. Here, recovery rates in vegetative filaments

are 0.053 s^{-1} and 0.012 s^{-1} for WT and Δ*fraI* mutant respectively.

Figure 4: The Δ*fraI* mutant is highly impaired in cell-cell communication in both vegetative filaments and hormogonia. CCCP-FRAP analysis of *Anabaena* PCC 7120, *N. punctiforme* WT (*N. pun* WT) and Δ*fraI* mutant vegetative filaments **(A)** and hormogonia **(B)**. FRAP-responses of calcein-stained untreated or CCCP-treated (90 min, 50 µm) were assigned to one of four groups indicated by the color scheme. The numbers within the bars indicate the number of analyzed cells (n) from different filaments. Recovery rates (*R*) of communicating cells **(C)**. Mean is shown. One-way ANOVA followed by Brown-Forsythe multiple comparison test with the WT was performed. The numbers above the mean indicate the number of cells used for calculation of *R*. *horm* – hormogonia. *N.pun WT horm* * excludes cells in hormogonia of the WT with measured R > 0.19 (see main text for explanation). Cumulated results from at least two independent cultures are shown. **, p<0.01****, p<0.0001; ns, not significant

Since FRAP experiments haven't yet been described in hormogonia, we wanted to measure the rate of communication between the cells in hormogonia and the cell's ability to close septal junctions upon treatment with the protonophore CCCP (Fig. 4B,C). On average, WT hormogonia showed much higher rates of communication (R = 0.104 s^{-1}) compared to vegetative cells (R = 0.054 s^{-1}) (Fig. 4C), with a fraction of the cells (23%) showing very high recovery rates (R > 0.19 s^{-1}). Many cells undergo cell-division more or less simultaneously during hormogonia development and hence in many cases the new septum may have not yet completely closed, explaining the large number of cells with such unusual high communication rates in FRAP analysis [5,46]. This idea is also supported by the fact that only 55% of the WT hormogonia cells completely stopped communication after CCCP treatment, in contrast to 86% among vegetative cells (Fig. 4B). Also, 76% of Δ*fraI* mutant hormogonia cells showed a communicating phenotype, compared to 61% of the vegetative cells, because, like in the WT, cell division was not yet completed and the fluorescent dye could easily diffuse between the future daughter cells (Fig. 4B).

The inability of the Δ*fraI* mutant to differentiate heterocysts can be explained by the lack of cell-cell communication. In contrast, the Δ*fraI* mutant can still differentiate akinetes despite decreased cell-cell communication". This is in line with our previous observation with the related species *Trichormus* (*Anabaena*) *variabilis* ATCC29413. There, we showed that during akinete formation molecular transfer decreases and seems not to play a role in this spore differentiation process[47].

399   **The Δ*fraI* mutant lacks the septal nanopore array**

400   To investigate whether aberrant nanopore array formation is causing the impaired

401   communication in the Δ*fraI* mutant, the septal peptidoglycan of vegetative cells and

402   hormogonia of the WT and the mutant were isolated and analyzed by TEM. The Δ*fraI* mutant

403   showed very few or almost no nanopores, both in PG from vegetative filaments as well as

404   from hormogonia (Fig. 5A,B). Interestingly, the Δ*fraI* mutant also had a significantly larger

405   septa diameter of vegetative filaments compared to the WT (Fig. 5B). A similar phenotype,

406   namely the lack of nanopores in septa and enlarged septa was previously observed for an

407   *amiC2* mutant of *N. punctiforme*[18]. Interestingly, the *amiC2* mutant displayed delocalized

408   septa leading to a severe morphological phenotype[12], which is not observed for the Δ*fraI*

409   mutant using Vancomycin-FL (Figure S4). In contrast, the pores in the lateral wall just around

410   the septal discs, were present in the Δ*fraI* mutant and the WT (Fig. 5A,B), implying that FraI

411   is not involved in formation of these lateral pores. In the center of the septal discs of the WT

412   and Δ*fraI* mutant hormogonia, we could occasionally observe few very large nanopores

413   (Figure S5), which may also explain higher rates of communication measured in some

414   hormogonia cells (see above).

415

416   **Figure 5:** The Δ*fraI* mutant completely lacks the septal nanopore array and has greater septa

417   diameter. Septal peptidoglycan in *N. punctiforme* WT and Δ*fraI* mutant vegetative filaments **(A)** and

418   hormogonia **(B)**. Representative TEM images are shown. Scale bar – 200 nm**. (C)** Number of

419   nanopores and septum diameter of *N. punctiforme* vegetative and hormogonia cells in WT and Δ*fraI*

420   mutant, which has very few or no nanopores and a greater septa diameter. One-way ANOVA followed

by Brown-Forsythe multiple comparison test. Median is shown. p<0.05 ** p<0.01, **** p<0.0001, ns
p>0.05.

Despite almost complete lack of nanopores, the AmiC2 protein could be detected in the
septa of Δ*fraI* mutant using anti-AmiC2 antibodies (Figure S3). Hence, FraI is unlikely to be
involved in septal localization of AmiC2. Moreover, FraI is predicted to have three
transmembrane helixes with a cytoplasmic C-terminus and an N-terminal signal sequence of
just 20 amino acids residing in the periplasm (Phobius), which makes a direct protein-protein
interaction with the periplasmic amidase unlikely. However, at the moment we cannot
exclude that a regulatory protein, like the LytM factor[22], is influenced by FraI. In line with the
idea of a FraI-AmiC2 relation is the fact that *fraI* expression is the highest during hormogonia
development (Figure S1), when many cells are undergoing cell division and the need for
AmiC2 activity is the highest[18]. The lethality of overexpressed FraI (see above), is another
indication that this protein is involved in regulation of the cell wall lytic enzyme AmiC, since
the over-expression of the *amiC* orthologues in *Nostoc* sp. PCC 7120 was also lethal[13].

It is interesting that the impaired cell-cell communication rate in the Δ*fraI* mutant did not
prevent hormogonia development. However, the motility of the hormogonia was reduced
(Fig. 1A). Nevertheless, the Δ*fraI* mutant can move towards the source of light, indicating that
cell-cell communication through the septal junctions is not crucial for hormogonia motility and
synchronized movement along the filament in response to a positive external signal.
Additionally, the Δ*fraI* mutant was able to change the direction of movement following a
change of light conditions (data not shown). Therefore, there may exist another way of
communication, perhaps through the common periplasm, which enables synchronized
movement of the filament. As an alternative explanation, each single cell may decide the
direction of the movement based on the sensing of external factors such as light and
chemoattractant gradient. It has already been speculated that a change in an electromotive
force may be a signal for translocation of HmpF protein of the pili machinery to the opposite
side of the cell, which may be sensed by each cell in hormogonia[48]. Furthermore, the
observed reduction in motility in the Δ*fraI* mutant could be attributed to the diminished
presence of the exopolysaccharide sheath surrounding hormogonia filaments (Fig. 3).
Previous studies have demonstrated reductions in motility in mutants with impaired
exopolysaccharide synthesis and export pathways [35,37]. Why the absence of *fraI* gene leads
to lower presence of hormogonia-specific exopolysaccharide remains unknown. It has been
speculated that HPS is exported through the periplasm close to the cell junctions in
hormogonia[49].

Alternatively, despite being significantly impaired in communication, 81% of Δ*fraI* hormogonia
cells (Fig. 4) still exhibit some communication, possibly facilitated by the presence of

remaining nanopores or incomplete septa formation. This residual communication may still be sufficient for the transport of signal necessary for the uniform polar distribution of pili machinery enabling directionality of movement along the filament in Δ*fraI* mutant.

Conclusion

In this study we discovered FraI, a new player in cell-cell communication and differentiation. Mutation of *fraI* leads to a low-communicating phenotype in both vegetative cells and hormogonia of *Nostoc punctiforme,* which can be explained by a nearly complete absence of nanopores in the septa of neighboring cells. Despite the low levels of communication, the Δ*fraI* strain forms motile hormogonia capable of synchronized movement, albeit in a different way than the WT. The filaments also perform phototaxis, suggesting that motility and directionality of movement do not require communication through the septal junctions. Lower motility observed in Δ*fraI* is more likely to be explained by the absence of the HPS sheath. In contrast, the presence of filaments containing a combination of vegetative and hormogonium cells in the Δ*fraI* mutant, a phenotype never observed in the wild type, implies that communication through the septal junctions may play a critical role in coordinating the development of hormogonia. As in other mutants of septal nanopore array formation, severely impaired cell-cell communication of the Δ*fraI* mutant is affecting heterocyst differentiation and function and hence diazotrophic growth. Nonetheless, akinete development is still possible in the Δ*fraI* mutant. FraI may play a role in indirectly controlling amidase activity through an unknown mechanism. Additional experiments focusing on the FraI protein will be necessary to support this claim. Finally, characterization of orthologues of known septal junction proteins from *Nostoc* sp. 7120 in *N. punctiforme* will provide additional understanding of the role of septal junction-mediated communication in the process of hormogonium development and motility.

Author contribution

AJ: designed and performed experiments, analyzed and interpreted data, drafted the work and wrote the manuscript.

CM: performed TEM experiments and sample preparation for ultra-thin sections.

DDR: designed and performed experiments, analyzed and interpreted data, made considerable contributions to the design of the study, interpreted data, wrote the manuscript.

GAP: designed and performed experiments, analyzed and interpreted data

IM: designed and supervised the research in Tübingen, made considerable contributions to the design of the study, interpreted data, revised the manuscript.

491   All authors approved the final manuscript.

Data Availability

493   Data available from authors upon reasonable request.

Acknowledgment

Conflict of Interest

496   The authors declare no competing interests.

References

1.   Zeng, X. & Zhang, C. C. The Making of a Heterocyst in Cyanobacteria. *Annu Rev Microbiol* **76**,
597–618 (2022).

2.   Herrero, A., Stavans, J. & Flores, E. The multicellular nature of filamentous heterocyst-forming
cyanobacteria. *FEMS Microbiol Rev* **40**, 831–854 (2016).

3.   Sukenik, A., Rücker, J. & Maldener, I. Dormant Cells (Akinetes) of Filamentous Cyanobacteria
Demonstrate a Great Variability in Morphology, Physiology, and Ecological Function. in
Cyanobacteria: From Basic Science to Applications 65–77 (Academic Press, 2019).
doi:10.1016/B978-0-12-814667-5.00004-0.

4.   Garg, R. & Maldener, I. The Formation of Spore-Like Akinetes: A Survival Strategy of
Filamentous Cyanobacteria. *Research Article Microb Physiol* **31**, 296–305 (2021).

5.   Maldener, I., Summers, M. L. & Sukenik, A. Cellular differentiation in filamentous
cyanobacteria. in *Cell Biology of Cyanobacteria* (eds. Flores, E. & Herrero, A.) 263–291 (Caister
Academic Press, Norfolk, 2014). doi:10.21775/9781910190739.

6.   Flores, E., Nieves-Morión, M. & Mullineaux, C. W. Cyanobacterial Septal Junctions: Properties
and Regulation. *Life* **9**, 1 (2018).

7.   Kieninger, A.-K. & Maldener, I. Cell–cell communication through septal junctions in
filamentous cyanobacteria. *Curr Opin Microbiol* **61**, 35–41 (2021).

8.   Weiss, G. L., Kieninger, A. K., Maldener, I., Forchhammer, K. & Pilhofer, M. Structure and
Function of a Bacterial Gap Junction Analog. *Cell* **178**, 374-384.e15 (2019).

9.   Kieninger, A. K., Tokarz, P., Janović, A., Pilhofer, M., Weiss, G. L. & Maldener, I. SepN is a
septal junction component required for gated cell–cell communication in the filamentous
cyanobacterium Nostoc. *Nat Commun* **13**, 7486 (2022).

10.   Merino-Puerto, V., Schwarz, H., Maldener, I., Mariscal, V., Mullineaux, C. W., Herrero, A. &
Flores, E. FraC/FraD-dependent intercellular molecular exchange in the filaments of a
heterocyst-forming cyanobacterium, *Anabaena sp. Mol Microbiol* **82**, 87–98 (2011).

11.   Ramos-León, F., Arévalo, S., Mariscal, V. & Flores, E. Specific mutations in the permease
domain of septal protein SepJ differentially affect functions related to multicellularity in the
filamentous cyanobacterium *Anabaena*. *Microbial Cell* **5**, 555–565 (2018).

12. Lehner, J., Zhang, Y., Berendt, S., Rasse, T. M., Forchhammer, K. & Maldener, I. The
morphogene AmiC2 is pivotal for multicellular development in the cyanobacterium *Nostoc*
punctiforme. *Mol Microbiol* **79**, 1655–1669 (2011).

13. Berendt, S., Lehner, J., Zhang, Y. V., Rasse, T. M., Forchhammer, K. & Maldener, I. Cell Wall
Amidase AmiC1 Is Required for Cellular Communication and Heterocyst Development in the
Cyanobacterium *Anabaena* PCC 7120 but Not for Filament Integrity. *J Bacteriol* **194**, 5218–
5227 (2012).

14. Nieves-Morión, M., Mullineaux, C. W. & Flores, E. Molecular Diffusion through Cyanobacterial
Septal Junctions. *mBio* **8**, e01756-16 (2017).

15. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in
filamentous cyanobacteria. *Nat Rev Microbiol* **8**, 39–50 (2010).

16. Kieninger, A.-K., Forchhammer, K. & Maldener, I. A nanopore array in the septal peptidoglycan
hosts gated septal junctions for cell-cell communication in multicellular cyanobacteria.
International Journal of Medical Microbiology **309**, 151303 (2019).

17. Vermassen, A., Leroy, S., Talon, R., Provot, C., Popowska, M. & Desvaux, M. Cell wall
hydrolases in bacteria: Insight on the diversity of cell wall amidases, glycosidases and
peptidases toward peptidoglycan. *Front Microbiol* **10:331**, (2019).

18. Lehner, J., Berendt, S., Dörsam, B., Pérez, R., Forchhammer, K. & Maldener, I. Prokaryotic
multicellularity: A nanopore array for bacterial cell communication. *FASEB Journal* **27**, 2293–
2300 (2013).

19. Sarasa-Buisan, C., Nieves-Morión, M., Arévalo, S., Helm, R. F., Sevilla, E., Luque, I. & Fillat, M.
F. FurC (PerR) contributes to the regulation of peptidoglycan remodeling and intercellular
molecular transfer in the cyanobacterium *Anabaena* sp. strain PCC 7120. *mBio* **15**, e03231-23
(2024).

20. Büttner, F. M., Faulhaber, K., Forchhammer, K., Maldener, I. & Stehle, T. Enabling cell-cell
communication via nanopore formation: structure, function and localization of the unique cell
wall amidase AmiC2 of *Nostoc punctiforme*. *FEBS* **283**, 1336–1350 (2016).

21. Bornikoel, J., Carrión, A., Fan, Q., Flores, E., Forchhammer, K., Mariscal, V., Mullineaux, C. W.,
Perez, R., Silber, N., Wolk, C. P. & Maldener, I. Role of Two Cell Wall Amidases in Septal
Junction and Nanopore Formation in the Multicellular Cyanobacterium *Anabaena* sp. PCC
7120. *Front Cell Infect Microbiol* **7:386**, (2017).

22. Bornikoel, J., Staiger, J., Madlung, J., Forchhammer, K. & Maldener, I. LytM factor Alr3353
affects filament morphology and cell-cell communication in the multicellular cyanobacterium
Anabaena sp. PCC 7120. *Mol Microbiol* **108**, 187–203 (2018).

23. Rai, A. N., Söderbäck, E. & Bergman, B. Cyanobacterium - Plant symbioses. *New Phytologist*
147, 449–481 (2000).

24. Álvarez, C., Navarro, J. A., Molina-Heredia, F. P. & Mariscal, V. Endophytic Colonization of Rice
(*Oryza sativa* L.) by the Symbiotic Strain *Nostoc punctiforme* PCC 73102. *Mol Plant Microbe*
Interact **33**, 1040–1045 (2020).

25. Zuniga, E. G., Boateng, K. K. A., Bui, N. U., Kurnfuli, S., Muthana, S. M. & Risser, D. D.
Identification of a hormogonium polysaccharide-specific gene set conserved in filamentous
cyanobacteria. *Mol Microbiol* **114**, 597–608 (2020).

26. Cho, Y. W., Gonzales, A., Harwood, T. V., Huynh, J., Hwang, Y., Park, J. S., Trieu, A. Q., Italia, P.,
Pallipuram, V. K. & Risser, D. D. Dynamic localization of HmpF regulates type IV pilus activity
and directional motility in the filamentous cyanobacterium *Nostoc punctiforme*. *Mol Microbiol*
106, 252–265 (2017).

27. Harwood, T. V., Zuniga, E. G., Kweon, H. J. & Risser, D. D. The cyanobacterial taxis protein
HmpF regulates type IV pilus activity in response to light. *Proc Natl Acad Sci U S A* **118**, (2021).

28. Parrett, G. A., Brones, P. D., Jenkins, G. M., Mounts, S. M., Nguyen, A. & Risser, D. D.
Identification of a morphogene required for tapered filament termini in filamentous
cyanobacteria. *Microbiology (N Y)* **169:001416**, (2023).

29. Allen, M. B. & Arnon, D. I. Studies on Nitrogen-Fixing Blue-Green Algae. I. Growth and
Nitrogen Fixation by *Anabaena Cylindrica* Lemm. *Plant Physiol* **30**, 366–372 (1955).

30. Rippka, E., Deruelles, J. & Waterbury, N. B. Generic Assignments, Strain Histories and
Properties of Pure Cultures of Cyanobacteria. *J Gen Microbiol* **111**, 1–61 (1979).

31. Black, T. A., Cai, Y. & Wolk, C. P. Spatial expression and autoregulation of hetR, a gene
involved in the control of heterocyst development in *Anabaena*. *Mol Microbiol* **9**, 77–84
(1993).

32. Risser, D. D., Wong, F. C. Y. & Meeks, J. C. Biased inheritance of the protein PatN frees
vegetative cells to initiate patterned heterocyst differentiation. *Proc Natl Acad Sci U S A* **109**,
15342–15347 (2012).

33. Khayatan, B., Bains, D. K., Cheng, M. H., Cho, Y. W., Huynh, J., Kim, R., Omoruyi, O. H., Pantoja,
A. P., Park, J. S., Peng, J. K., Splitt, S. D., Tian, M. Y. & Risser, D. D. A putative O-linked β-
Nacetylglucosamine transferase is essential for hormogonium development and motility in
the filamentous cyanobacterium *Nostoc punctiforme*. *J Bacteriol* **199**, e00075-17 (2017).

34. Wolk, C. P., Cai, Y. & Panoff, J.-M. Use of a transposon with luciferase as a reporter to identify
environmentally responsive genes in a cyanobacterium. *Proc. Natl. Acad. Sci. USA* **88**, 5355–
5359 (1991).

35. Risser, D. D. & Meeks, J. C. Comparative transcriptomics with a motility-deficient mutant leads
to identification of a novel polysaccharide secretion system in *Nostoc punctiforme*. *Mol
Microbiol* **87**, 884–893 (2013).

36. Splitt, S. D. & Risser, D. D. The non-metabolizable sucrose analog sucralose is a potent
inhibitor of hormogonium differentiation in the filamentous cyanobacterium *Nostoc
punctiforme*. *Arch Microbiol* **198**, 137–147 (2016).

37. Khayatan, B., Meeks, J. C. & Risser, D. D. Evidence that a modified type IV pilus-like system
powers gliding motility and polysaccharide secretion in filamentous cyanobacteria. *Mol
Microbiol* **98**, 1021–1036 (2015).

38. Perez, R., Forchhammer, K., Salerno, G. & Maldener, I. Clear differences in metabolic and
morphological adaptations of akinetes of two Nostocales living in different habitats.
Microbiology (N Y) **162**, 214–223 (2016).

39. Fiedler, G., Arnold, M., Hannus, S. & Maldener, I. The DevBCA exporter is essential for
envelope formation in heterocysts of the cyanobacterium *Anabaena* sp. strain PCC 7120. *Mol
Microbiol* **27**, 1193–1202 (1998).

40. Risser, D. D., Chew, W. G. & Meeks, J. C. Genetic characterization of the hmp locus, a
chemotaxis-like gene cluster that regulates hormogonium development and motility in *Nostoc*
punctiforme. *Mol Microbiol* **92**, 222–233 (2014).

41. Kieninger, A.-K. Identification and Characterization of the Cell-Cell Communication System of a
Multicellular Cyanobacterium. Dissertation. (University of Tübingen, Tübingen, 2021).

42. Gonzalez, A., Riley, K. W., Harwood, T. V, Zuniga, E. G., Risser, D. D. & Ellermeier, C. D. A
Tripartite, Hierarchical Sigma Factor Cascade Promotes Hormogonium Development in the
Filamentous Cyanobacterium *Nostoc punctiforme*. *mSphere* **4**, e00231-219 (2019).

43. Harwood, T. V & Risser, D. D. The primary transcriptome of hormogonia from a filamentous
cyanobacterium defined by cappable-seq. *Microbiology (Reading)* **167**, (2021).

44. Nürnberg, D. J., Mariscal, V., Bornikoel, J., Nieves-Morión, M., Krauß, N., Herrero, A.,
Maldener, I., Flores, E. & Mullineaux, C. W. Intercellular diffusion of a fluorescent sucrose
analog via the septal junctions in a filamentous cyanobacterium. *mBio* **6**, e02109 (2015).

45. Mullineaux, C. W., Mariscal, V., Nenninger, A., Khanum, H., Herrero, A., Flores, E. & Adams, D.
G. Mechanism of intercellular molecular exchange in heterocyst-forming cyanobacteria.
EMBO Journal **27**, 1299–1308 (2008).

46. Meeks, J. C. & Elhai, J. Regulation of Cellular Differentiation in Filamentous Cyanobacteria in
Free-Living and Plant-Associated Symbiotic Growth States. *Microbiology and Molecular*
Biology Reviews **66**, 94–121 (2002).

47. Garg, R., Luckner, M., Berger, J., Hipp, K., Wanner, G., Forchhammer, K. & Maldener, I.
Changes in Envelope Structure and Cell–Cell Communication during Akinete Differentiation
and Germination in Filamentous Cyanobacterium *Trichormus variabilis* ATCC 29413. *Life* **12**,
429 (2022).

48. Harwood, T. V., Zuniga, E. G., Kweon, H. J. & Risser, D. D. The cyanobacterial taxis protein
HmpF regulates type IV pilus activity in response to light. *Proc Natl Acad Sci USA* **118**,
e2023988118 (2021).

49. Hoiczyk, E. & Baumeister, W. The junctional pore complex, a prokaryotic secretion organelle,
is the molecular motor underlying gliding motility in cyanobacteria. *Current Biology* **8**, 1161–
1168 (1998).

641