Proteomic profile analysis of plasma and aqueous humor from glaucoma and non-glaucomatous patients

Carmen L. Pessuti[1], Chia-Ling Huang[2], Angela Banks[2], Nhi Vo[2], Lori Jennings[2], Joseph Loureiro[2], Kleber S. Ribeiro[1], Deise Fialho Costa[1], Heloisa Nascimento[1], Cristina Muccioli[1], Ivan Maynart Tavares[1], Alessandra G. Commodaro[1], Rubens Belfort Jr[1], Christopher W. Wilson[2], Amy Chen[2], Ganesh Prasanna[2], VijayKrishna Raghunathan[2].

[1]Department of Ophthalmology, Universidade Federal de Sao Paulo, Sao Paulo, Brazil

[2]BioMedical Research, Novartis, Cambridge, MA, USA

Corresponding author(s):

VijayKrishna Raghunathan, Ph.D., Ophthalmology, BioMedical Research, Novartis, 22 Windsor St, Cambridge, MA, Massachusetts, 02139. _Email_: Vijay.Raghunathan@novartis.com

Ganesh Prasanna, Ph.D., Ophthalmology, BioMedical Research, Novartis, 22 Windsor St, Cambridge, MA, Massachusetts, 02139. _Email_: ganesh.prasanna@novartis.com

Running title: Proteomic analysis of glaucomatous donor biofluids

Keywords: Aqueous humor, plasma, intraocular pressure, glaucoma, cup-to-disc ratio, SomaScan, proteomics

ABSTRACT

Purpose: Glaucoma, a multifactorial ocular neuropathy, can lead to irreversible vision loss.

Diagnosis involves assessing optic cupping (increased cup-to-disc ratios) and structural

changes (like retinal nerve fiber layer thinning) through clinical imaging. Elevated intraocular

pressure (IOP) is commonly associated with glaucoma, but not always. However, understanding

disease progression is hindered by limited access to donor ocular tissue and consistent clinical

data. We hypothesize that the proteome of aqueous humor and plasma may be altered in

disease and correlates with clinical parameters such as IOP and cup-to-disc ratios.

Methods: Aqueous humor (AH) and plasma samples were collected from 36 glaucoma patients

(17 male, 19 female), and 35 non-glaucomatous control patients (16 male, 19 female)

undergoing cataract surgery. The protein profile was compared using the SOMAscan® assay

system for proteome profiling. From glaucomatous donors, correlations between IOP and cup-

to-disc ratios to proteome differences were determined.

Results: Overall proteomics profiles between both AH and plasma were compared by

combining all samples (glaucoma and non-glaucoma) and then performing correlation analyses.

This study revealed similar protein abundance in the two biological fluids. Additionally, it

identified different abundance of proteins in plasma and AH between glaucoma and non-

glaucoma samples. The differential proteins identified were involved in pathways related to

vascular integrity, inflammation, immune response, cell adhesion, and complement activation.

Generally, glaucomatous AH showed higher protein levels. Neurofilament light chain (NEFL)

protein correlated with elevated IOP and inflammatory markers, but not with cup-to-disc ratios.

Conclusions: Together, our data demonstrate that the proteins identified in this study from

glaucomatous donors correspond to markers of neurodegeneration and those that may inhibit

cell proliferation or disrupt vascular integrity.

INTRODUCTION

Primary open angle glaucoma (POAG) is one of the leading causes of irreversible blindness, estimated to affect 111.8 million people worldwide by 2040 [1,2]. Although the etiology and progression of POAG is multifactorial, the only modifiable causal risk factor for managing visual field loss is to reduce intraocular pressure (IOP), which is demonstrated to slow the progression of damage to the optic nerve and death of retinal ganglion cells [3]. IOP is largely determined by a balance between aqueous humor production and resistance to outflow [4,5]. In the human eye, approximately 75% of all aqueous humor flows from the anterior chamber through the trabecular meshwork (TM) and Schlemm's canal (SC) [6,7] and is thus the primary egress for aqueous humor [8]. Increased resistance to aqueous drainage due to changes in TM cells and extracellular matrix (ECM) is considered a major contributor to ocular hypertension (OHT) [9,10]. Differential profiles and levels of soluble factors have been identified in the AH, which have the potential to affect cells in the tissues (outflow tract) that they traverse through, and thus influence OHT.

AH is secreted into the posterior chamber by the non-pigmented epithelial cells of the ciliary processes. Though the total amount of protein in the anterior chamber is less than 1% (w/v) due to the functional blood aqueous barrier [4], the aqueous humor is enriched with proteins secreted by intraocular structures from both the anterior and posterior segments [11]. It is thus reasonable to hypothesize that systemic disorders, genetic conditions, or perturbations to ocular integrity and health may all adversely influence expression of proteins in the aqueous humor and other ocular tissues [12]. Disease associations of protein compositions of tear films, AH and vitreous humors have been observed with dry eye, glaucoma, age related macular degeneration, or other pathologies [13-18], although very few studies have correlated these with clinically relevant parameters.

Several studies have demonstrated differences between metabolites or proteomes in the AH of patients with primary open angle glaucoma and have subsequently posited that these may serve as biomarkers of dysfunction in the outflow apparatus [16,19-29]. Proteomic analysis of

aqueous humor reveals significant protein expression differences between cataract and primary open angle glaucoma (POAG) patients. Multiple studies report that dozens to hundreds of proteins are differentially expressed in POAG, involving pathways such as oxidative stress, inflammation, lipid metabolism, extracellular matrix regulation, and neural degeneration [30-41]. Key findings from these various studies identify several proteins in the complement cascade, apolipoproteins, and antioxidant defense suggesting immune involvement and metabolic stress in POAG.

For the aforementioned investigations, complementary approaches were used to quantify protein expression in AH from glaucomatous donors, including ELISA, multiplexed immunoassays, LC-MS/MS, or antibody microarrays. Limitations in quantifying proteins/metabolites include the small volume and low protein concentration in AH, alongside the need for validated antibody-based methods, which restricts biomarker identification. Thus, more sensitive methods are needed to broadly identify AH proteins to find disease progression biomarkers when comparing pre- and post-treatment stages. Discovering new proteins in the AH of patients could indeed shed light on the molecular mechanisms behind outflow resistance in OHT and glaucoma. Overall, while these proteomic changes provide insight into POAG mechanisms and support biomarker development, limited studies[42] have performed direct comparisons between plasma and aqueous humor or with clinical parameters such as IOP or cup-to-disc ratios. Others have suggested that differences in proteome profiling within POAG populations may enable subgrouping to identify POAG severity[43], though these studies lacked clinical correlation. We hypothesize that proteins secreted into the AH and plasma of patients with glaucoma differ from non-glaucomatous individuals and are associated with clinically relevant parameters. Here, we utilize the SOMAscan® assay [44], a highly multiplexed, aptamer-based technology, to detect and assess relative abundance of a broad range of proteins from

small volumes of human blood alongside AH from patients undergoing glaucoma surgery, as

compared to controls patients subject to cataract surgery.

MATERIALS AND METHODS

Study population

The study was approved by the ethics committee investigational review board (CEP 1.209.725)

and adhered to the principles of the Declaration of Helsinki. Informed consent was obtained

from all participants. Aqueous humor (AH) and plasma samples were collected from 32

glaucoma patients (16 male, 16 female), and 35 cataract control patients (16 male, 19 female)

undergoing cataract surgery (**Figure 1**). The control cohort underwent clinical examinations

(e.g. elevated IOP, optic nerve head changes, or uveitis) to rule out any glaucoma-related

findings as part of inclusion/exclusion criteria. Within the glaucoma patient group one underwent

trabeculectomy, two were pseudophakic, and AH was collected at the beginning of the surgery.

Clinical diagnosis of glaucoma was based on elevated IOP (IOP > 21 mmHg), documented

progressive cupping or thinning of the neuroretinal rim. All glaucoma patients were treated with

medication to control IOP. Patients with other eye diseases or associated with systemic

diseases were not included. Our study is exploratory in nature, and the primary focus is on the

proteomic analysis of plasma and aqueous humor samples from the study participants. As such,

the data about the glaucoma patients in this study will be limited to the existing inclusion criteria

and general demographic information. Additional longitudinal clinical data including summary

statistics of HVF, glaucoma phenotype, number of medications, etc. were unavailable and thus

not being reported here for this study. All donor meta data including IOP, cup to disc ratios and

visual acuity are included as **Supplementary Table 1**.

Aqueous humor (AH) and plasma sample collection

AH samples were collected from patients following a previously validated method [45] at the

beginning of the surgery. Briefly, after anesthetic and antiseptic eyedrops administration, a 1mL

tuberculin syringe (27 gauge, ½ inch) was used to aspirate the AH. Approximately, 0.1 ml was

collected, and transferred to two labeled Eppendorf tubes. Peripheral blood (10 mL) was

collected from an arm of the patient in EDTA tubes during the surgery. The tubes were

centrifuged for 10 min at 1900 x g, and plasma was collected. Total protein concentrations in the

AH samples were determined using the BCA Protein Assay Kit (Pierce, Rockford, IL).

SOMAscan® analysis

All AH and plasma samples were analyzed in parallel using the SOMAscan® proteome profiling

platform that provides a broad and relative abundant assessment of selected proteins in

biological fluid. The SOMAscan® platform, using affinity-based slow off-rate modified aptamers

(SOMAmers), has been described extensively elsewhere [44,46,47]. The custom SOMAscan®

version used in our study covered 5,080 SOMAmers, including 4,785 human-specific

SOMAmers targeting 4,161 human proteins. Evidence supporting SOMAmer endogenous

antigen annotation, including SOMAmer validation by proteogenetic evidence, mass

spectrometry detection, or orthogonal assay concordance, where known, are provided in the

referenced studies[48-66]. The targeted proteins include transmembrane receptors, secreted

proteins, kinases, transcription regulators and signal transducers. The samples (55 µl) were

aliquoted into 2D barcoded tubes and were shipped on dry ice to SomaLogic for analysis.

Statistical analysis

The relative abundance of captured SOMAmers were quantified by microarray hybridization as

relative fluorescence units (RFU). The annotations for the SOMAmers and $\log_2$(RFU) for all

samples analyzed are provided in **Supplementary table 2**. Quality control and sample outlier

detection was conducted using the arrayQualityMetrics[67], and by running principal component

analysis (PCA) and Spearman's correlation analysis with custom R scripts. RFU were log2-

transformed and normalized by smooth quantile normalization using R package qsmooth [68]. In

total, 5 samples (2 non-glaucomatous, 3 glaucoma) were removed from the AH cohort and 2

non-glaucomatous samples were removed from the plasma cohort, which left 62 samples in the

AH cohort and 65 samples in the AH cohort. Exclusion of 5 samples were based on an

unbiased statistical outlier analysis (**Supplemental Figure 1**) and are thus not presented in

subsequent analysis. We applied the limma R package (v3.50.3) in R 4.1.0 environment to

determine differentially expressed proteins between groups[69]. Pathway enrichment analysis was

performed with pre-ranked GSEA using R package fgsea v1.20.0 and Gene Ontology Biological

Process category (GOBP) in MSigDB[70-73].

Correlation analysis

We conducted Elastic Net, a machine learning based method, to identify proteins correlated with

the total protein concentration. For this analysis, we utilized the cv.glmnet function from the R

package glmnet v4.1-3. We employed a 10-fold cross-validation approach to optimize the alpha

and lambda parameters[74,75]. Additionally, we performed a 1000-fold bootstrapping using the

optimized alpha and lambda values to determine the frequency at which each protein was

selected as a top predictor for estimating the total protein concentration. To enhance the

performance of our analysis, we only included proteins with an absolute Pearson's correlation

coefficient greater than 0.4 with the total protein concentration.

RESULTS

Donor demographics

Donor demographics for the patient samples used in this study are provided in **Figure 1**. The

average age of donors with cataract was 66.9 ± 10.9 while those with glaucoma was 68.41 ±

12.84. No significant differences (p=0.60, t-test; **Figure 1**) were seen in donor ages between the

groups *in toto* or when evaluated by sex. Amongst these donors, we note that one donor had

primary angle closure glaucoma while the remainder had POAG. The mean IOP of glaucoma

donors (IOP controlled with medication) was 17.94 ± 8.28 mmHg with no differences seen

between male or female donors. Only one POAG donor had an IOP > 24 mmHg. A statistically

significant difference in total protein concentration in AH was observed between cataract and

glaucoma groups (Cataract: 0.383 ± 0.505 mg/ml; Glaucoma: 1.17 ± 1.48 mg/ml). Visual acuity

and cup-to-disc ratios of POAG donors are listed in donor metadata within **Supplementary**

Table 1.

Differential analysis of SOMAScan® profiling data

Principal component analysis of the cataract and glaucoma samples revealed heterogeneity

between samples, including outliers that were subsequently excluded from further analysis.

Subsequent analysis revealed few differences in protein levels between non-glaucomatous and

glaucomatous aqueous humor and plasma (**Supplementary Fig. 1**). Nevertheless, we

observed that samples were clustered by tissue types with reasonable correlation ($r > 0.5$)

between AH and plasma samples (**Figure 2a**). The histogram (**Figure 2b**) demonstrates that

the distribution of Pearson's correlation coefficients between AH and plasma samples from

same donor demonstrates some variability. Considering all donors, a reasonable correlation ($r >$

0.5) was observed comparing protein levels between AH and plasma samples.

Subsequent to multivariate analysis, we performed differential protein analysis for both

plasma and aqueous samples. Expression values that were $|log2(RFU)| > 0.5$ and where the

adjusted p-value was < 0.05 are highlighted as significant (**Figure 3**). Within glaucomatous AH

(Figure 3A), 7 SOMAmer probes representing 6 target proteins were observed to be significantly

down regulated, while 49 SOMAmer probes representing 49 target proteins were upregulated.

Examples of top up-regulated proteins include complement proteins (C3, C7, C1QTNF3),

Apolipoprotein (APOE, APOF), matrix proteins (endostatin/Col18A1), macrophage associated

proteins (CSF1R, CD163), while top down-regulated proteins include growth factors such as

FGF9, FGF20 and CCK. We observed that in plasma samples, the majority of the differential

proteins were upregulated in glaucoma. Some of the upregulated proteins of interest include

ENO1, NEK7, AKT, PRKCA, PRKCB, JAK2, MAP kinase-activated protein kinase 2

(MAPKAPK2), whereas the top down-regulated proteins include APOC3, GDF2, DCP1B and

EDIL3. A comprehensive list of proteins altered in their levels comparing glaucomatous and

non-glaucomatous plasma and AH are provided in **Supplementary table 3**. Although protein

levels from AH and plasma from the same donor showed a positive correlation, we only

observed SNX4 to be significantly up regulated in both AH and plasma samples from

glaucomatous donors.

Pathway enrichment analysis

Next, we asked if the differentially enriched proteins were enriched in common biological

pathways, by performing pathway analysis (**Supplementary table 4**). We identified 51 and 559

significantly up-regulated pathways (adjusted p-value < 0.05) in AH and plasma samples from

glaucoma donors, respectively. 16 pathways were up-regulated in both AH and plasma,

including inflammatory response, IL6 production and immune response related pathways.

Correlation analysis of protein levels with total protein concentration

While individual protein changes may help identify signaling pathways perturbed in disease,

total protein concentration of the AH may help ascertain the overall health of the blood-aqueous

barrier and the outcomes of IOP lowering medications. Thus, normalizing individual proteins to

total protein concentrations assists in determining if observed changes reflect pathway changes

underpinning disease, or are merely a consequence of structural failure or drug treatment(s).

We conducted an Elastic Net regression analysis to identify proteins that can accurately predict

the total protein concentration (**Figure 4a**). Our analysis specifically focused on AH proteins, as

proteins from these samples may provide a better reflection of changes in glaucoma disease

compared to plasma samples. Through our analysis, we successfully identified 5 proteins

(KNG1, IGFBO6, MAP2K4, H6PD, and C3) that demonstrated the highest predictive capability

for the total protein concentration collected from the patients (**Figure 4a**).

Correlation analysis of protein levels with IOP

We conducted Elastic net regression analysis to identify 13 proteins to be associated with IOP

values reported in glaucomatous patients (**Figure 4b**). We note that a glaucoma sample with

extremely high IOP measurement was excluded from the analysis. Among the proteins of

interest in glaucoma, the protein with maximal correlation with IOP levels in glaucomatous

patients was neurofilament light chain (NEFL; **Figure 4c, Supplementary Fig. 2**). Linear

regression analysis validated this finding and demonstrated a net correlation with IOP

measurements (R^2 = 0.3067). Since NEFL is a widely accepted marker for several

neurodegenerative diseases including glaucoma [76-85], we then decided to further investigate if

levels of NEFL correlated with other SOMAmers identified in glaucomatous donors. We

calculated the Pearson's correlation coefficient of NEFL and rest of the proteins and identified

29 proteins that are positively or negatively correlated with NEFL ($|r > 0.4|$), including TEAD3,

CNTF, C4A, C4B and CCL27 (**Figure 5**).

Changes in protein levels as a function of cup-to-disc ratios in glaucoma

Finally, we sought to determine if changes in relative abundance of protein levels in AH

correlated with cup-to-disc ratios of glaucomatous patients via Pearsons's correlation analysis

(**Figure 6**). For patients where cup-to-disc ratios were available for both eyes, we used the

average measurements from both eyes. We note that among the 29 glaucomatous samples

obtained, cup-to-disc ratios were available for 26 donors. Figure 6 shows the top 6 positive

(KLRG2, C8G, PENK, ALB, APOA1, and KNG1) and negative (IL12A/B, LRRC3, ZAP70,

CTSG, DEFB115, and NSG1) correlated proteins with cup-to-disc ratios.

DISCUSSION

Monitoring glaucoma in the clinic typically involves measuring IOP, gonioscopy, and advanced

imaging of the retina/optic nerve head, which require regular visits and patient compliance.

Without a detailed medical history, it is challenging to understand the molecular mechanisms of

disease progression or identify biomarkers to ascertain treatment effectiveness. Since collecting

intraocular tissues from living patients is not feasible, analyzing biofluids such as AH and blood

can provide critical insights into disease mechanisms. In this study, we examined protein levels

in AH and plasma from patients with and without glaucoma undergoing cataract surgery using a

SomaLogic platform and correlating these with IOP and cup-to-disc ratios measured

immediately prior to the surgery.

Comparative analysis of plasma and aqueous humor proteomes

The ages and sex of glaucomatous and non-glaucomatous donors in this dataset were

comparable. We observed significant changes in relative abundance of a subset of proteins

identified both in AH and plasma. Protein levels from AH and plasma from the same donor

sample showed a positive correlation (r > 0.5 for all of samples). Interestingly, when comparing

protein levels and correlations across all fluids and disease states, we only observed a

significant upregulation of SNX4 in glaucomatous AH and plasma when compared with those

from non-glaucomatous donors. SNX4 is a synaptic & endosomal membrane trafficking protein

in the secretory pathway and is also implicated as a mitochondrial recycling quality protein in

aging [86-88]. The lack of greater correlation between plasma and AH proteins may be due to

several reasons: turnover rates of AH *vs* plasma, tissues & cell types involved in protein

secretions, systemic *vs* local transport of proteins. It is yet unclear if glaucoma is purely a

disease of the eye or if systemic factors exist that regulate its etiology and progression, although

co-morbidities have been reported [89-93]. Further, Miguel Coca-Prados et al[94] have identified

several unique proteins and peptides in the AH attributed to the neuroendocrine nature of ciliary

non-pigmented epithelial cells in addition to plasma proteins. It is thus unsurprising that

differences in uniquely identified proteins exists between the two biofluids. Nevertheless, in the

plasma, 48 proteins were upregulated while 4 (EDIL3, DCP1B, GDF2, APOC3) were

downregulated when comparing non-glaucomatous and glaucomatous donors. Gene ontology

analyses of proteins altered in the plasma of glaucomatous donors were related to multiple

pathways corresponding to protein & macromolecule localization, intracellular transport, kinase

activity pathways, post translational / post transcriptional regulation of proteins / genes,

cytoskeletal reorganization and GPCR related pathways. On the other hand, 7 (ISLR2, CCK,

INSL4, FGF9, FGF20, BMPR1A) proteins were significantly downregulated and 49 were

significantly upregulated in the AH of glaucomatous donors. Gene ontology analyses of proteins

identified as altered in the AH of glaucomatous donors were related to multiple pathways

corresponding to inflammatory response, humoral & adaptive immune response, cell-cell & cell-

matrix adhesion, complement activation, and wound healing responses. These together suggest

that though overall concordance in proteins may be observed between AH and plasma, some

differences exist and thus proximity of biofluid to relevant tissue being studied (i.e. ciliary

process, ciliary muscle and trabecular meshwork/Schlemm's canal) may be important to

consider. As such, deeper analysis in this study was focused on the ocular specific biofluid (AH)

proteins.

Effect of disease state on aqueous humor proteome

All changes in protein levels were normalized to total protein concentration and as such

these changes are independent of mean protein concentration. Interestingly, regression

modeling demonstrated that total protein concentrations trended towards being significantly

greater in AH of glaucomatous compared with non-glaucomatous donors. While this is

suggestive of a breakdown in the blood-aqueous barrier [4], whether this is due to the chronic

nature of the disease or due to IOP lowering medications (e.g. prostaglandin analogs), other

systemic medications or other co-morbidities is unclear. Nevertheless, the changes in proteins

appear to be directly related to pathways with the potential for consequential effects on IOP

regulation and outflow homeostasis. When comparing our overall findings with that from

previous publications using other technologies, it was noteworthy that proteins related to

complement and immune-related proteins (e.g. C3b, IL1, C1Q, C7 etc), apolipoproteins (APOE,

APOF etc), heat shock/oxidative stress (HSPD1/Hsp60), matrix and serpin (SERPINA1,

SERPING1, BGN etc) remained consistently differentially expressed between glaucomatous

and non-glaucomatous AH[34,41,95-101]. These suggest that the proteomics methodology used in

our study is sensitive and robustly identifies certain proteins, in glaucoma, regardless of study

design or detection method. This highlights that certain pathways and proteins may have a

functional or disease-correlative role that required further studies to establish their role in

disease mechanisms. Nonetheless, several unique proteins were also identified in our study.

For example, Col18A1, which was upregulated in AH of glaucomatous donors, is expressed

ubiquitously in ocular tissues except in photoreceptors, and protein fragments that contain

endostatin (from Col18A1 cleavage) has been observed to accumulate in ocular fluid samples

[102]. Interestingly, endostatin has been reported to promote expression and release of

thrombospondin-1, implicated in outflow resistance & glaucoma [103-105], in non-ocular endothelial

cells [106]. Conversely, endostatin has also been posited to crosstalk with Rho/ROCK, TGFβ, NF-

κB, PDGF, and autophagy pathways to yield anti-fibrotic effects [107]. SVEP1, another

overexpressed secreted ECM protein in AH & a disease modifier allele for congenital glaucoma

[108], is known to be a binding partner for TIE1 and can thus regulate signaling outcomes in

lymphatics and vasculature [109]. Since AH predominantly drains into the Schlemm's canal, a

unique vessel with both vasculature and lymphatic properties [110], studying SVEP1 in the context

of glaucoma warrants further investigation. Of particular interest, VCAM1, CDH11, and BGN

were other proteins that were overexpressed in the AH of glaucomatous patients. SVEP1,

CDH11 and VCAM1 were all identified as genes relevant to POAG pathogenesis in a genome-

wide meta-analysis across ancestries [111]. CDH11 can interact reciprocally with fibronectin

binding protein syndecan-4 to facilitate cell migration and adhesion, partake in EMT, and

modulate proliferation [112-115]. In non-ocular cells, CDH11 was shown to mediate adhesion of

macrophages to fibroblasts promoting transdifferentiation into myofibroblasts and a self-

sustaining profibrotic niche [116]. Biglycan (BGN) is an extracellular proteoglycan, expressed in

the trabecular meshwork [117], is posited as a prognostic marker for cancer aggressiveness [118],

fibrosis & inflammation [119] and is predicted to be a circulating "messenger" for triggering

inflammation and/or autophagy [120]. Whether biglycan in the AH serves as a biomarker, a

signaling molecule to trigger autophagy or inflammatory phenotypes in the trabecular meshwork

to mediate subsequent outflow regulation requires mechanistic studies. High levels of circulatory

VCAM1 in the blood/plasma was found to correlate with ventricular hypertrophy, hypertension

and is suggested to thus serve as a predictive soluble biomarker for cardiovascular disease and

inflammation [121-123]. In this study, whether VCAM1 expression is a result of chronic elevated

IOP, drugs to lower IOP, or due to undiagnosed/unknown underlying systemic vascular

conditions is unknown. Regardless, increased presence of the aforementioned proteins may

serve as 'proteins-of-interest' in mechanistic understanding of outflow regulation for as potential

biomarkers for dysregulation in outflow homeostasis. Further studies and orthogonal methods

are needed to validate these findings.

Protein changes identified as a function of clinical parameters in glaucoma

Since proteins with associations with 'hypertension' were identified, we next sought to

determine if protein changes were a function of IOP levels through additional regression

modeling. Two proteins notably were found to be of importance: DLL4 and NEFL. DLL4, a

Notch ligand, is a key regulator of vascular morphogenesis, vessel maturation, and function.

Secreted/soluble DLL4 has previously been reported to significantly reduce hydraulic

conductivity, vascular permeability, and disrupt endothelial barrier function in non-ocular

vasculature [124,125]. Further, DLL4 has been demonstrated to be critical in developing retinal

vasculature, increased endothelial cell proliferation, and angiogenic sprouting [126]. Interestingly,

DLL4 was also shown to inhibit inflammatory choroidal neovascularization despite opposing

effects seen in endothelial cells (anti-angiogenic) and macrophages (pro-angiogenic) [127].

Collectively, we speculate that DLL4, found upregulated with IOP, may act to increase outflow

resistance via a yet unknown mechanism. Further, whether a dichotomous function for the DLL4

ligand or the Notch pathway exists in outflow regulation remains to be seen. A recent study

suggests that Notch pathway protein expression may differ between segmental flow regions of

healthy and glaucomatous donors, at least *in vitro* [128]. Thus, ligands circulating in the aqueous

humor could differentially impact TM cell function necessitating further studies.

NEFL is a well-established marker of neurodegeneration [77,79-82,129] and similar to our finding,

has previously been reported to be elevated in glaucomatous AH in humans and animal models

[76,78,130]. In fact, NEFL had the highest correlation with IOP in all donors in this study, providing

additional confidence that NEFL in the AH may indeed be a suitable biomarker for

glaucomatous neurodegeneration and ocular hypertension. However, we advise caution that AH

samples utilized in this study were from patients identified for glaucoma filtration surgery.

Therefore, it is likely that the disease stage may be advanced and thus markers of axonal

degeneration are expected within the AH. Whether NEFL would serve as an early marker of

disease is unclear and thus requires comprehensive longitudinal natural history investigations.

Further the source of the neurofilament protein, determined in the aqueous humor, is unclear

(i.e. whether it is due to degeneration of the optic nerve, secretion from cells of neural crest

origin (developmentally) in the anterior segment, or Schwann/nerve cells in the anterior segment

remains unknown). Interestingly, while NEFL levels correlated with IOP, its levels did not appear

to correlate with cup-to-disc ratios of glaucoma donors (**Supplementary Fig. 3**), though other

proteins that did correlate were identified. Interestingly, cup-to-disc ratio was independent of

IOP values as well (**Supplementary Fig. 3**). Inflammatory proteins (such as interleukins and

cathepsins) corresponded with low cup-to-disc ratio, while apolipoprotein A1 (APOA1), albumin,

kininogen-1 (KNG1) and complement C8G correlated with a higher cup-to-disc ratio suggesting

proteins altered in the AH may differ based on disease severity. KNG1 is an antiangiogenic

molecule that has been suggested to be a marker of neurodegeneration [131] and can impair the

proliferation of endothelial cells [132]. To the best of our knowledge, the role of KNG1 in POAG

and/or ocular hypertension has not been studied, although its cleaved nonapeptide, Bradykinin

(BK), has been the target for IOP lowering investigations [133-138]. A prior study reported a

decrease in C8G [139] in glaucomatous AH conflicting with our results. However, Kim et al [140]

previously reported C8G to act as a neuroinflammation inhibitor; our result correlating C8G

levels with high cup-to-disc ratios may thus reflect the advanced stage of
neurodegeneration/neuroinflammation in glaucoma.

Consistent with our study, elevated levels of APOA1 were reported in AH of POAG donors
[141]. It is important to note that the direct role of APOA1 in IOP homeostasis is not known.
However, APOA1 plays a critical role in reverse cholesterol transport pathway via direct
interactions with the ABCA1 gene whose variants are implicated in POAG [142]. Cholesterol is
itself a risk factor for elevated IOP [143]. Elevation in albumin levels in glaucomatous AH is also
reported in several studies, though attributed to administration of IOP lowering drugs [144]. Finally,
a correlation analysis of NEFL to other proteins identified in glaucomatous aqueous humor
revealed that NEFL levels may be associated with proteins regulating apoptosis, complement
activation, proliferation, and cytoskeletal reorganization. Together, the proteins identified in this
study correspond to both markers of neurodegeneration and those that may inhibit proliferation
or vascular integrity.

LIMITATIONS

This study is not without limitations. One of the major limitations is that the biofluids obtained in
this study were obtained at the time of ocular surgery thus representing a singular snapshot in
time and no information was gleaned about the stage of the disease; although, clinically
measured IOP measurements and cup-to-disc ratios are available at this stage. We understand
that glaucoma can be asymmetric, and in future studies, we will consider incorporating cup-to-
disc ratios from objective approaches such as OCT to validate or refine these findings. Also,
while cup-to-disc ratios are a clinical parameter, and protein levels (which potentially are
dynamic) are a molecular parameter, it is important to note that these variables are
independent, and that the kinetics of structural changes may differ from the kinetics of molecular
turnover. The independent parameters were nevertheless compared to ascertain if the clinical
parameter investigated had any association with the proteome content in the AH and if this

could perhaps reflect the current state of the disease in the donor. Under the current study

design, it is important to note that any causal or correlational relationship is hard to conclude

since there are practical limitations on frequency of aqueous humor sampling. Further,

comprehensive and longitudinal patient histories of systemic co-morbidities or ocular diseases

and medications that may affect systemic or ocular hypertension including but limited to

intraocular pressure were unavailable. Since IOP in glaucoma patients was controlled with IOP

lowering medication, the impact of these therapeutics on proteome changes cannot be

ascertained. Further functional measurements for visual field or additional structural deficits

including longitudinal and pre-surgical fundus photography or OCT measurements (nerve fiber

layer thinning, rim width, ganglion cell analysis etc.) are not known for these patients. We

speculate that additional information that may be obtained from *longitudinal prospective studies*

with disease state factored in could further enable identification of determining factors driving

changes in the proteome. We did not perform any genetic linkage or association analysis to

identify any polygenic risk assessment from the POAG patients enrolled. Future studies may be

needed to identify correlations between genetic causes, structure-function changes, and

molecular profiling approaches in a longitudinal manner if feasible. Since the entirety of the

samples collected in this study were utilized for proteomics, attempts to validate the results

using orthogonal or secondary methodologies were not undertaken. As such, since mass

spectrometry (a commonly used profiling technique for proteomics) was not part of the current

study's scope, we recommend future research from independent investigators to include such

methods, and to compare SOMAscan findings with previously published aqueous humor

proteomic profiles and/or methods. Thus, we anticipate that the approach and data presented

here will enable the design of validation studies for future investigations within the glaucoma

community.

ACKNOWLEDGEMENTS

The authors would like to thank the patient donors for the biological fluid samples without whose

consent these experiments would be impossible.

FUNDING STATEMENT

This work was sponsored by Novartis Biomedical Research, and CNPq (Ministry of Science,

Technology, and Innovation of Brazil).

AUTHORS' DISCLOSURE

C.H, A.B, N.V, L.J, J.L, C.W.W, A.C, G.P, V.R are all employees of Novartis Biomedical

Research.

CONTRIBUTION STATEMENT

C.L.S, K.S.R, D.F.C, H.N, C.M, I.M.T, A.G.C, R.B.J: clinical design, sample collection, data

interpretation, critical review, revisions, and approval of article. C.H, A.B: data analysis, data

interpretation, critical review, revisions, and approval of article. A.C, G.P: Conceptualization of

study, data interpretation, critical review, revisions, and approval of article. N.V, L.J, J.L: sample

preparation & analysis, data analysis, data interpretation, critical review, revisions, and approval

of article. C.W.W: data interpretation, critical review, revisions, and approval of article. V.R: data

analysis, data interpretation, critical review, revisions, and approval of article.

FIGURE LEGENDS

Figure 1: Patient demographics demonstrate study participants were age and sex matched.

Intraocular pressures and protein concentrations from AH are reported for both glaucomatous

and non-glaucomatous donors.

Figure 2: Proteins identified in aqueous humor and plasma clustered by type of biofluid, and appear to correlate reasonably well as evidenced by Spearman's correlation. **(A)** Heatmap, and **(B)** histogram of Spearman's correlation index demonstrates that the correlation, while evident, is not absolute i.e. $0.5 < R^2 < 0.7$.

Figure 3: Principal component (PCA) and differential expression analysis demonstrates differences in proteins identified in **(A)** aqueous humor and **(B)** plasma comparing glaucomatous and non-glaucomatous donors. PCA demonstrates large variance between cataract (non-glaucoma) and glaucomatous donor samples with no discernible separation between the two disease groups. Volcano plots demonstrate that no significant differences were seen comparing cataract (non-glaucoma) and glaucomatous donors for most proteins. Few proteins whose levels were significantly altered are represented in the bar plot.

Figure 4: Elastic net regression analysis (aqueous humor only) demonstrated **(A)** increased protein concentration is observed in aqueous humor of donors with glaucoma, with no specific sex related differences observed, **(B)** changes in some proteins correlate significantly with IOP independent of sex of donor. **(C)** Scatter plots and linear regression of proteins ($R^2 > 0.25$) as a function of IOP indicate NEFL, DLL4, and NFE2L1 to most significantly altered proteins in the aqueous humor.

Figure 5: Heatmap demonstrates correlation of NEFL to top 30 proteins identified in the aqueous humor of glaucomatous donors. Pearson's correlation analysis comparing NEFL with rest of the proteins identified ~30 proteins that either positively (red) or negatively (blue) correlated with NEFL ($|r > 0.4|$). In general, more proteins that negatively correlated with NEFL levels were identified than those that exhibited positive correlation.

Figure 6: Heatmap identifies top 6 proteins either up or down regulated in the aqueous humor

of glaucomatous donors as a function of cup-to-disc ratios.

SUPPLEMENTARY INFORMATION

Supplementary figure 1: Outlier and principal component analysis identifying donors to

exclude from data set in an unbiased manner.

Supplementary figure 2: Scatter plots of proteins identified in glaucomatous AH as a function

of IOP demonstrate large variability between samples.

Supplementary figure 3: (A) Scatter plot of NEFL expression levels as a function of cup-to-

disc ratios of glaucomatous donors demonstrate no significant relationship ($R^2 = 0.037$) between

the two parameters. **(B)** Cup-to-disc ratios of glaucomatous donors show no significant

correlation ($R^2 = 0.008$) with intraocular pressures reported.

Supplementary table 1: Donor metadata including age, sex, disease state, sample ID, IOP,

cup-to-disc ratios & visual acuity.

Supplementary table 2: List of SOMAmer annotations and $\log_2$(RFU) for all samples analyzed

from plasma and aqueous humor for both glaucoma and non-glaucoma donors.

Supplementary table 3: List of differentially expressed proteins in aqueous humor and plasma

comparing glaucomatous and non-glaucomatous donors.

Supplementary table 4: GSEA pathway analysis of proteins identified in aqueous humor and

plasma comparing glaucomatous and non-glaucomatous donors.

REFERENCES

1. Tham Y-C, Li X, Wong TY, et al. Global Prevalence of Glaucoma and Projections of Glaucoma
Burden through 2040: A Systematic Review and Meta-Analysis. Ophthalmology 2014;121(11):2081-2090,
doi:https://doi.org/10.1016/j.ophtha.2014.05.013
2. Allison K, Patel D, Alabi O. Epidemiology of Glaucoma: The Past, Present, and Predictions for the
Future. Cureus 2020, doi:10.7759/cureus.11686
3. Glaucoma. In: Moorfields Manual of Ophthalmology. (Jackson TL. ed.) Mosby: Edinburgh; 2008;
pp. 274-330.
4. Freddo TF. A contemporary concept of the blood-aqueous barrier. Prog Retin Eye Res
2013;32(181-95, doi:10.1016/j.preteyeres.2012.10.004
5. Freddo TF, Johnson M. Chapter 6 Aqueous Humor Outflow Resistance. In: The Eye's Aqueous
Humor. 2008; pp. 161-192.
6. Toris CB, Yablonski ME, Wang Y-L, et al. Aqueous humor dynamics in the aging human eye.
American journal of ophthalmology 1999;127(4):407-412
7. Toris CB, Koepsell SA, Yablonski ME, et al. Aqueous humor dynamics in ocular hypertensive
patients. Journal of glaucoma 2002;11(3):253-258
8. Stamer WD, Acott TS. Current understanding of conventional outflow dysfunction in glaucoma.
Curr Opin Ophthalmol 2012;23(2):135-43, doi:10.1097/ICU.0b013e32834ff23e
9. Acott TS, Kelley MJ. Extracellular matrix in the trabecular meshwork. Exp Eye Res
2008;86(4):543-61, doi:10.1016/j.exer.2008.01.013
10. Vranka JA, Kelley MJ, Acott TS, et al. Extracellular matrix in the trabecular meshwork: intraocular
pressure regulation and dysregulation in glaucoma. Exp Eye Res 2015;133(112-25,
doi:10.1016/j.exer.2014.07.014
11. Yorio T, Prasanna G, Coca-Prados M. CHAPTER 4 - The Ciliary Body: A Potential Multifaceted
Functional Neuroendocrine Unit. In: Ocular Therapeutics. (Yorio T, Clark AF, Wax MB. eds.) Academic
Press: London; 2008; pp. 69-85.
12. Ahmad MT, Zhang P, Dufresne C, et al. The Human Eye Proteome Project: Updates on an
Emerging Proteome. PROTEOMICS 2018;18(5-6):1700394, doi:10.1002/pmic.201700394
13. Baek J-H, Lim D, Park KH, et al. Quantitative proteomic analysis of aqueous humor from patients
with drusen and reticular pseudodrusen in age-related macular degeneration. BMC Ophthalmology
2018;18(1), doi:10.1186/s12886-018-0941-9
14. Aghamollaei H, Parvin S, Shahriary A. Review of proteomics approach to eye diseases affecting
the anterior segment. Journal of Proteomics 2020;225(103881,
doi:https://doi.org/10.1016/j.jprot.2020.103881
15. Skeie JM, Roybal CN, Mahajan VB. Proteomic Insight into the Molecular Function of the Vitreous.
PLOS ONE 2015;10(5):e0127567, doi:10.1371/journal.pone.0127567
16. Kang GY, Bang JY, Choi AJ, et al. Exosomal proteins in the aqueous humor as novel biomarkers
in patients with neovascular age-related macular degeneration. J Proteome Res 2014;13(2):581-95,
doi:10.1021/pr400751k
17. Dammeier S, Martus P, Klose F, et al. Combined Targeted Analysis of Metabolites and Proteins
in Tear Fluid With Regard to Clinical Applications. Translational Vision Science & Technology
2018;7(6):22-22, doi:10.1167/tvst.7.6.22
18. De Almeida Borges D, Alborghetti MR, Franco Paes Leme A, et al. Tear proteomic profile in three
distinct ocular surface diseases: keratoconus, pterygium, and dry eye related to graft-versus-host
disease. Clinical Proteomics 2020;17(1), doi:10.1186/s12014-020-09307-5
19. Agnifili L, Pieragostino D, Mastropasqua A, et al. Molecular biomarkers in primary open-angle
glaucoma: from noninvasive to invasive. Prog Brain Res 2015;221(1-32, doi:10.1016/bs.pbr.2015.05.006
20. Babizhayev MA. Biomarkers and special features of oxidative stress in the anterior segment of
the eye linked to lens cataract and the trabecular meshwork injury in primary open-angle glaucoma:
challenges of dual combination therapy with N-acetylcarnosine lubricant eye drops and oral formulation of
nonhydrolyzed carnosine. Fundam Clin Pharmacol 2012;26(1):86-117, doi:10.1111/j.1472-
8206.2011.00969.x
21. Bollinger KE, Crabb JS, Yuan X, et al. Quantitative proteomics: TGFbeta(2) signaling in
trabecular meshwork cells. Invest Ophthalmol Vis Sci 2011;52(11):8287-94, doi:10.1167/iovs.11-8218

22.   Burgos-Blasco B, Vidal-Villegas B, Saenz-Frances F, et al. Tear and aqueous humour cytokine
profile in primary open-angle glaucoma. Acta Ophthalmol 2020;98(6):e768-e772, doi:10.1111/aos.14374
23.   Castany M, Jordi I, Catala J, et al. Glaucoma patients present increased levels of diadenosine
tetraphosphate, Ap4A, in the aqueous humour. Experimental Eye Research 2011;92(3):221-226,
doi:http://dx.doi.org/10.1016/j.exer.2010.12.004
24.   Csosz E, Deak E, Toth N, et al. Comparative analysis of cytokine profiles of glaucomatous tears
and aqueous humour reveals potential biomarkers for trabeculectomy complications. FEBS Open Bio
2019;9(5):1020-1028, doi:10.1002/2211-5463.12637
25.   Garweg JG, Zandi S, Pfister IB, et al. Comparison of cytokine profiles in the aqueous humor of
eyes with pseudoexfoliation syndrome and glaucoma. PLoS One 2017;12(8):e0182571,
doi:10.1371/journal.pone.0182571
26.   Guo T, Guo L, Fan Y, et al. Aqueous humor levels of TGFbeta2 and SFRP1 in different types of
glaucoma. BMC Ophthalmol 2019;19(1):170, doi:10.1186/s12886-019-1183-1
27.   Igarashi N, Honjo M, Asaoka R, et al. Aqueous autotaxin and TGF-betas are promising diagnostic
biomarkers for distinguishing open-angle glaucoma subtypes. Sci Rep 2021;11(1):1408,
doi:10.1038/s41598-021-81048-3
28.   Lin JB, Sheybani A, Santeford A, et al. Increased Aqueous Humor GDF15 Is Associated with
Worse Visual Field Loss in Pseudoexfoliative Glaucoma Patients. Transl Vis Sci Technol 2020;9(10):16,
doi:10.1167/tvst.9.10.16
29.   Sun C, Zhang H, Jiang J, et al. Angiogenic and inflammatory biomarker levels in aqueous humor
and vitreous of neovascular glaucoma and proliferative diabetic retinopathy. Int Ophthalmol
2020;40(2):467-475, doi:10.1007/s10792-019-01207-4
30.   Kaeslin MA, Killer HE, Fuhrer CA, et al. Changes to the Aqueous Humor Proteome during
Glaucoma. PLOS ONE 2016;11(10):e0165314, doi:10.1371/journal.pone.0165314
31.   Kaur I, Kaur J, Sooraj K, et al. Comparative evaluation of the aqueous humor proteome of
primary angle closure and primary open angle glaucomas and age-related cataract eyes. International
Ophthalmology 2019;39(1):69-104, doi:10.1007/s10792-017-0791-0
32.   Sharma S, Bollinger KE, Kodeboyina SK, et al. Proteomic Alterations in Aqueous Humor From
Patients With Primary Open Angle Glaucoma. Investigative Opthalmology & Visual Science
2018;59(6):2635, doi:10.1167/iovs.17-23434
33.   Rejas-González R, Montero-Calle A, Pastora Salvador N, et al. Unraveling the nexus of oxidative
stress, ocular diseases, and small extracellular vesicles to identify novel glaucoma biomarkers through in-
depth proteomics. Redox Biology 2024;77(103368, doi:10.1016/j.redox.2024.103368
34.   Saccà SC, Centofanti M, Izzotti A. New Proteins as Vascular Biomarkers in Primary Open Angle
Glaucomatous Aqueous Humor. Investigative Opthalmology & Visual Science 2012;53(7):4242,
doi:10.1167/iovs.11-8902
35.   Beutgen VM, Graumann J. Advances in aqueous humor proteomics for biomarker discovery and
disease mechanisms exploration: a spotlight on primary open angle glaucoma. Frontiers in Molecular
Neuroscience 2024;17(doi:10.3389/fnmol.2024.1397461
36.   Adav SS, Wei J, Terence Y, et al. Proteomic Analysis of Aqueous Humor from Primary Open
Angle Glaucoma Patients on Drug Treatment Revealed Altered Complement Activation Cascade. Journal
of Proteome Research 2018;17(7):2499-2510, doi:10.1021/acs.jproteome.8b00244
37.   Nikhalashree S, George R, Shantha B, et al. Detection of Proteins Associated with Extracellular
Matrix Regulation in the Aqueous Humour of Patients with Primary Glaucoma. Current Eye Research
2019;44(9):1018-1025, doi:10.1080/02713683.2019.1608261
38.   Izzotti A, Longobardi M, Cartiglia C, et al. Proteome Alterations in Primary Open Angle Glaucoma
Aqueous Humor. Journal of Proteome Research 2010;9(9):4831-4838, doi:10.1021/pr1005372
39.   Liu H. Proteome alterations in aqueous humour of primary open angle glaucoma patients.
International Journal of Ophthalmology 2020;13(1):176-179, doi:10.18240/ijo.2020.01.24
40.   Funke S, Perumal N, Bell K, et al. The potential impact of recent insights into proteomic changes
associated with glaucoma. Expert Review of Proteomics 2017;14(4):311-334,
doi:10.1080/14789450.2017.1298448
41.   Kiełbus M, Kuźmiuk D, Skrzyniarz AM, et al. Changes in the Protein Composition of the Aqueous
Humor in Patients with Glaucoma: An Update Review. International Journal of Molecular Sciences
2025;26(7):3129, doi:10.3390/ijms26073129

42.   Kodeboyina SK, Lee TJ, Bollinger K, et al. Aqueous Humor Proteomic Alterations Associated with
Visual Field Index Parameters in Glaucoma Patients: A Pilot Study. Journal of Clinical Medicine
2021;10(6):1180, doi:10.3390/jcm10061180
43.   Hubens WHG, Mohren RJC, Liesenborghs I, et al. The aqueous humor proteome of primary open
angle glaucoma: An extensive review. Exp Eye Res 2020;197(108077, doi:10.1016/j.exer.2020.108077
44.   Gold L, Ayers D, Bertino J, et al. Aptamer-Based Multiplexed Proteomic Technology for
Biomarker Discovery. PLoS ONE 2010;5(12):e15004, doi:10.1371/journal.pone.0015004
45.   Belfort R, Jr., Moura NC, Mendes NF. T and B lymphocytes in the aqueous humor of patients with
uveitis. Arch Ophthalmol 1982;100(3):465-7, doi:10.1001/archopht.1982.01030030467019
46.   Mehan MR, Ostroff R, Wilcox SK, et al. Highly multiplexed proteomic platform for biomarker
discovery, diagnostics, and therapeutics. Adv Exp Med Biol 2013;735(283-300, doi:10.1007/978-1-4614-
4118-2_20
47.   Gold L, Walker JJ, Wilcox SK, et al. Advances in human proteomics at high scale with the
SOMAscan proteomics platform. N Biotechnol 2012;29(5):543-9, doi:10.1016/j.nbt.2011.11.016
48.   Zhang J, Dutta D, Kottgen A, et al. Plasma proteome analyses in individuals of European and
African ancestry identify cis-pQTLs and models for proteome-wide association studies. Nat Genet
2022;54(5):593-602, doi:10.1038/s41588-022-01051-w
49.   Surapaneni A, Schlosser P, Zhou L, et al. Identification of 969 protein quantitative trait loci in an
African American population with kidney disease attributed to hypertension. Kidney Int 2022;102(5):1167-
1177, doi:10.1016/j.kint.2022.07.005
50.   Katz DH, Robbins JM, Deng S, et al. Proteomic profiling platforms head to head: Leveraging
genetics and clinical traits to compare aptamer- and antibody-based methods. Sci Adv
2022;8(33):eabm5164, doi:10.1126/sciadv.abm5164
51.   Gudjonsson A, Gudmundsdottir V, Axelsson GT, et al. A genome-wide association study of
serum proteins reveals shared loci with common diseases. Nat Commun 2022;13(1):480,
doi:10.1038/s41467-021-27850-z
52.   Emilsson V, Gudmundsdottir V, Gudjonsson A, et al. Coding and regulatory variants are
associated with serum protein levels and disease. Nat Commun 2022;13(1):481, doi:10.1038/s41467-
022-28081-6
53.   Pietzner M, Wheeler E, Carrasco-Zanini J, et al. Mapping the proteo-genomic convergence of
human diseases. Science 2021;374(6569):eabj1541, doi:10.1126/science.abj1541
54.   Ferkingstad E, Sulem P, Atlason BA, et al. Large-scale integration of the plasma proteome with
genetics and disease. Nat Genet 2021;53(12):1712-1721, doi:10.1038/s41588-021-00978-w
55.   Stanley S, Vanarsa K, Soliman S, et al. Comprehensive aptamer-based screening identifies a
spectrum of urinary biomarkers of lupus nephritis across ethnicities. Nat Commun 2020;11(1):2197,
doi:10.1038/s41467-020-15986-3
56.   Raffield LM, Dang H, Pratte KA, et al. Comparison of Proteomic Assessment Methods in Multiple
Cohort Studies. Proteomics 2020;20(12):e1900278, doi:10.1002/pmic.201900278
57.   Cuvelliez M, Vandewalle V, Brunin M, et al. Circulating proteomic signature of early death in heart
failure patients with reduced ejection fraction. Sci Rep 2019;9(1):19202, doi:10.1038/s41598-019-55727-1
58.   Bodewes ILA, van der Spek PJ, Leon LG, et al. Fatigue in Sjogren's Syndrome: A Search for
Biomarkers and Treatment Targets. Front Immunol 2019;10(312, doi:10.3389/fimmu.2019.00312
59.   Yao C, Chen G, Song C, et al. Genome-wide mapping of plasma protein QTLs identifies
putatively causal genes and pathways for cardiovascular disease. Nat Commun 2018;9(1):3268,
doi:10.1038/s41467-018-05512-x
60.   Sun BB, Maranville JC, Peters JE, et al. Genomic atlas of the human plasma proteome. Nature
2018;558(7708):73-79, doi:10.1038/s41586-018-0175-2
61.   Giudice V, Biancotto A, Wu Z, et al. Aptamer-based proteomics of serum and plasma in acquired
aplastic anemia. Exp Hematol 2018;68(38-50, doi:10.1016/j.exphem.2018.09.008
62.   Forster K, Sass S, Ehrhardt H, et al. Early Identification of Bronchopulmonary Dysplasia Using
Novel Biomarkers by Proteomic Screening. Am J Respir Crit Care Med 2018;197(8):1076-1080,
doi:10.1164/rccm.201706-1218LE
63.   Emilsson V, Ilkov M, Lamb JR, et al. Co-regulatory networks of human serum proteins link
genetics to disease. Science 2018;361(6404):769-773, doi:10.1126/science.aaq1327
64.   Suhre K, Arnold M, Bhagwat AM, et al. Connecting genetic risk to disease end points through the
human blood plasma proteome. Nat Commun 2017;8(14357, doi:10.1038/ncomms14357

65.   Sasayama D, Hattori K, Ogawa S, et al. Genome-wide quantitative trait loci mapping of the
human cerebrospinal fluid proteome. Hum Mol Genet 2017;26(1):44-51, doi:10.1093/hmg/ddw366
66.   Carayol J, Chabert C, Di Cara A, et al. Protein quantitative trait locus study in obesity during
weight-loss identifies a leptin regulator. Nat Commun 2017;8(1):2084, doi:10.1038/s41467-017-02182-z
67.   Kauffmann A, Gentleman R, Huber W. arrayQualityMetrics--a bioconductor package for quality
assessment of microarray data. Bioinformatics 2009;25(3):415-6, doi:10.1093/bioinformatics/btn647
68.   Hicks SC, Okrah K, Paulson JN, et al. Smooth quantile normalization. Biostatistics
2018;19(2):185-198, doi:10.1093/biostatistics/kxx028
69.   Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for RNA-
sequencing and microarray studies. Nucleic Acids Res 2015;43(7):e47, doi:10.1093/nar/gkv007
70.   Aleksander SA, Balhoff J, Carbon S, et al. The Gene Ontology knowledgebase in 2023.
GENETICS 2023;224(1), doi:10.1093/genetics/iyad031
71.   Ashburner M, Ball CA, Blake JA, et al. Gene Ontology: tool for the unification of biology. Nature
Genetics 2000;25(1):25-29, doi:10.1038/75556
72.   Liberzon A, Subramanian A, Pinchback R, et al. Molecular signatures database (MSigDB) 3.0.
Bioinformatics 2011;27(12):1739-1740, doi:10.1093/bioinformatics/btr260
73.   Liberzon A, Birger C, Thorvaldsdóttir H, et al. The Molecular Signatures Database Hallmark Gene
Set Collection. Cell Systems 2015;1(6):417-425, doi:10.1016/j.cels.2015.12.004
74.   Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. J Stat Softw 2010;33(1):1-22, doi:10.18637/jss.v033.i01
75.   Tay JK, Narasimhan B, Hastie T. Elastic Net Regularization Paths for All Generalized Linear
Models. Journal of Statistical Software 2023;106(1), doi:10.18637/jss.v106.i01
76.   Lin JB, Pitts KM, El Helwe H, et al. Neurofilament Light Chain in Aqueous Humor as a Marker of
Neurodegeneration in Glaucoma. Clinical Ophthalmology 2023;Volume 17(2209-2217,
doi:10.2147/opth.s417664
77.   Kaeser SA, Lehallier B, Thinggaard M, et al. A neuronal blood marker is associated with mortality
in old age. Nature Aging 2021;1(2):218-225, doi:10.1038/s43587-021-00028-4
78.   Sasaki Y, Engber TM, Hughes RO, et al. cADPR is a gene dosage-sensitive biomarker of SARM1
activity in healthy, compromised, and degenerating axons. Experimental Neurology 2020;329(113252,
doi:https://doi.org/10.1016/j.expneurol.2020.113252
79.   Baka P, Steenken L, Escolano-Lozano F, et al. Studying serum neurofilament light chain levels
as a potential new biomarker for small fiber neuropathy. Eur J Neurol 2024;e16192,
doi:10.1111/ene.16192
80.   Fissolo N, Benkert P, Sastre-Garriga J, et al. Serum biomarker levels predict disability
progression in patients with primary progressive multiple sclerosis. J Neurol Neurosurg Psychiatry 2023,
doi:10.1136/jnnp-2023-332251
81.   Frank A, Bendig J, Schnalke N, et al. Serum neurofilament indicates accelerated
neurodegeneration and predicts mortality in late-stage Parkinson's disease. NPJ Parkinsons Dis
2024;10(1):14, doi:10.1038/s41531-023-00605-x
82.   Franthal S, Khalil M, Kern D, et al. Elevated serum neurofilament light chain protein in patients
with essential tremor. Eur J Neurol 2024;31(2):e16143, doi:10.1111/ene.16143
83.   Novobilsky R, Bartova P, Licha K, et al. Serum neurofilament light chain levels in patients with
cognitive deficits and movement disorders: comparison of cerebrospinal and serum neurofilament light
chain levels with other biomarkers. Front Hum Neurosci 2023;17(1284416,
doi:10.3389/fnhum.2023.1284416
84.   Sano T, Masuda Y, Yasuno H, et al. Plasma neurofilament light chain as a potential biomarker of
neurodegeneration in murine brain. Toxicol Res (Camb) 2023;12(5):751-755, doi:10.1093/toxres/tfad063
85.   Yang Y, Kim WS, Michaelian JC, et al. Predicting neurodegeneration from sleep related biofluid
changes. Neurobiol Dis 2024;190(106369, doi:10.1016/j.nbd.2023.106369
86.   Seaman MNJ. Membrane traffic in the secretory pathway. Cellular and Molecular Life Sciences
2008;65(18):2842-2858, doi:10.1007/s00018-008-8354-1
87.   Vazquez-Sanchez S, Gonzalez-Lozano MA, Walfenzao A, et al. The endosomal protein sorting
nexin 4 is a synaptic protein. Scientific Reports 2020;10(1), doi:10.1038/s41598-020-74694-6
88.   Henkel V, Schürmanns L, Brunner M, et al. Role of sorting nexin PaATG24 in autophagy, aging
and development of Podospora anserina. Mechanisms of Ageing and Development 2020;186(111211,
doi:10.1016/j.mad.2020.111211

89.   Dada T, Verma S, Gagrani M, et al. Ocular and Systemic Factors Associated with Glaucoma. J
Curr Glaucoma Pract 2022;16(3):179-191, doi:10.5005/jp-journals-10078-1383
90.   Zhang T, Xie X, Lu F. Primary open-angle glaucoma: Neuroendocrine–immune disorder? Medical
Hypotheses 2014;83(4):514-515, doi:10.1016/j.mehy.2014.08.017
91.   Salim S, Shields MB. Glaucoma and Systemic Diseases. Survey of ophthalmology
2010;55(1):64-77, doi:10.1016/j.survophthal.2009.03.006
92.   Pache M, Flammer J. A Sick Eye in a Sick Body? Systemic Findings in Patients with Primary
Open-angle Glaucoma. Survey of ophthalmology 2006;51(3):179-212,
doi:10.1016/j.survophthal.2006.02.008
93.   Nucci C, Martucci A, Cesareo M, et al. Links among glaucoma, neurodegenerative, and vascular
diseases of the central nervous system. In: Elsevier: 2015; pp. 49-65.
94.   Coca-Prados M, Escribano J. New perspectives in aqueous humor secretion and in glaucoma:
the ciliary body as a multifunctional neuroendocrine gland. Prog Retin Eye Res 2007;26(3):239-62,
doi:10.1016/j.preteyeres.2007.01.002
95.   Auler N, Tonner H, Pfeiffer N, et al. Antibody and Protein Profiles in Glaucoma: Screening of
Biomarkers and Identification of Signaling Pathways. Biology 2021;10(12):1296,
doi:10.3390/biology10121296
96.   Samelska K, Zaleska-Żmijewska A, Bałan B, et al. Immunological and molecular basics of the
primary open angle glaucoma pathomechanism. Central European Journal of Immunology
2021;46(1):111-117, doi:10.5114/ceji.2021.104328
97.   Vernazza S, Oddone F, Tirendi S, et al. Risk Factors for Retinal Ganglion Cell Distress in
Glaucoma and Neuroprotective Potential Intervention. International Journal of Molecular Sciences
2021;22(15):7994, doi:10.3390/ijms22157994
98.   Kodeboyina SK, Lee TJ, Churchwell L, et al. The Constitutive Proteome of Human Aqueous
Humor and Race Specific Alterations. Proteomes 2020;8(4):34, doi:10.3390/proteomes8040034
99.   Beutgen VM, Perumal N, Pfeiffer N, et al. Autoantibody Biomarker Discovery in Primary Open
Angle Glaucoma Using Serological Proteome Analysis (SERPA). Frontiers in Immunology
2019;10(doi:10.3389/fimmu.2019.00381
100.   Williams E, Altman J, Ahmed S, et al. Unveiling the levels and significance of different serpin
family proteins in aqueous humor dynamics. BMC Ophthalmology 2025;25(1), doi:10.1186/s12886-025-
04119-3
101.   De Maria A, Zientek KD, David LL, et al. Compositional Analysis of Extracellular Aggregates in
the Eyes of Patients With Exfoliation Syndrome and Exfoliation Glaucoma. Invest Ophthalmol Vis Sci
2021;62(15):27, doi:10.1167/iovs.62.15.27
102.   Määttä M, Heljasvaara R, Pihlajaniemi T, et al. Collagen XVIII/endostatin shows a ubiquitous
distribution in human ocular tissues and endostatin-containing fragments accumulate in ocular fluid
samples. Graefe's Archive for Clinical and Experimental Ophthalmology 2006;245(1):74-81,
doi:10.1007/s00417-006-0281-y
103.   Haddadin RI, Oh DJ, Kang MH, et al. Thrombospondin-1 (TSP1)-null and TSP2-null mice exhibit
lower intraocular pressures. Invest Ophthalmol Vis Sci 2012;53(10):6708-17, doi:10.1167/iovs.11-9013
104.   Shan S-W, Do C-W, Lam TC, et al. Thrombospondin-1 mediates Rho-kinase inhibitor-induced
increase in outflow-facility. Journal of Cellular Physiology 2021;236(12):8226-8238,
doi:https://doi.org/10.1002/jcp.30492
105.   Wirtz MK, Sykes R, Samples J, et al. Identification of Missense Extracellular Matrix Gene
Variants in a Large Glaucoma Pedigree and Investigation of the N700S Thrombospondin-1 Variant in
Normal and Glaucomatous Trabecular Meshwork Cells. Curr Eye Res 2022;47(1):79-90,
doi:10.1080/02713683.2021.1945109
106.   Goyanes AM, Moldobaeva A, Marimoutou M, et al. Functional Impact of Human Genetic Variants
of COL18A1/Endostatin on Pulmonary Endothelium. American Journal of Respiratory Cell and Molecular
Biology 2020;62(4):524-534, doi:10.1165/rcmb.2019-0056oc
107.   Zhang Z, Liu X, Shen Z, et al. Endostatin in fibrosis and as a potential candidate of anti-fibrotic
therapy. Drug Delivery 2021;28(1):2051-2061, doi:10.1080/10717544.2021.1983071
108.   Young TL, Whisenhunt KN, Jin J, et al. SVEP1 as a Genetic Modifier of TEK-Related Primary
Congenital Glaucoma. Invest Ophthalmol Vis Sci 2020;61(12):6, doi:10.1167/iovs.61.12.6

109.   Hußmann M, Schulte D, Weischer S, et al. Svep1 is a binding ligand of Tie1 and affects specific
aspects of facial lymphatic development in a Vegfc-independent manner. eLife 2023;12(e82969,
doi:10.7554/eLife.82969
110.   Kizhatil K, Ryan M, Marchant JK, et al. Schlemm's Canal Is a Unique Vessel with a Combination
of Blood Vascular and Lymphatic Phenotypes that Forms by a Novel Developmental Process. PLoS
Biology 2014;12(7):e1001912, doi:10.1371/journal.pbio.1001912
111.   Gharahkhani P, Jorgenson E, Hysi P, et al. Genome-wide meta-analysis identifies 127 open-
angle glaucoma loci with consistent effect across ancestries. Nat Commun 2021;12(1):1258,
doi:10.1038/s41467-020-20851-4
112.   Langhe RP, Gudzenko T, Bachmann M, et al. Cadherin-11 localizes to focal adhesions and
promotes cell-substrate adhesion. Nat Commun 2016;7(10909, doi:10.1038/ncomms10909
113.   Schneider DJ, Wu M, Le TT, et al. Cadherin-11 contributes to pulmonary fibrosis: potential role in
TGF-β production and epithelial to mesenchymal transition. The FASEB Journal 2012;26(2):503-512,
doi:10.1096/fj.11-186098
114.   Passanha FR, Divinagracia ML, Lapointe VLS. Cadherin-11 Regulates Cell Proliferation via the
PDGFRβ-ERK1/2 Signaling Pathway in Human Mesenchymal Stem Cells. Stem Cells 2022;40(2):165-
174, doi:10.1093/stmcls/sxab019
115.   Passanha FR, Geuens T, LaPointe VLS. Cadherin-11 Influences Differentiation in Human
Mesenchymal Stem Cells by Regulating the Extracellular Matrix Via the TGFβ1 Pathway. Stem Cells
2022;40(7):669-677, doi:10.1093/stmcls/sxac026
116.   Lodyga M, Cambridge E, Karvonen HM, et al. Cadherin-11–mediated adhesion of macrophages
to myofibroblasts establishes a profibrotic niche of active TGF-β. Science Signaling
2019;12(564):eaao3469, doi:10.1126/scisignal.aao3469
117.   Wirtz MK, Bradley JMB, Xu H, et al. Proteoglycan expression by human trabecular meshworks.
Current Eye Research 1997;16(5):412-421, doi:10.1076/ceyr.16.5.412.7040
118.   Pinto F, Santos-Ferreira L, Pinto M, et al. The Extracellular Small Leucine-Rich Proteoglycan
Biglycan Is a Key Player in Gastric Cancer Aggressiveness. Cancers 2021;13(6):1330,
doi:10.3390/cancers13061330
119.   Schulz M, Diehl V, Trebicka J, et al. Biglycan: A regulator of hepatorenal inflammation and
autophagy. Matrix Biol 2021;100-101(150-161, doi:10.1016/j.matbio.2021.06.001
120.   Poluzzi C, Nastase MV, Zeng-Brouwers J, et al. Biglycan evokes autophagy in macrophages via
a novel CD44/Toll-like receptor 4 signaling axis in ischemia/reperfusion injury. Kidney Int 2019;95(3):540-
562, doi:10.1016/j.kint.2018.10.037
121.   Kuroda Y. Vascular cell adhesion molecule-1 as a biochemical marker of left ventricular mass in
the patients with hypertension. American Journal of Hypertension 2001;14(9):868-872,
doi:10.1016/s0895-7061(01)02139-2
122.   Troncoso MF, Ortiz-Quintero J, Garrido-Moreno V, et al. VCAM-1 as a predictor biomarker in
cardiovascular disease. Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease
2021;1867(9):166170, doi:10.1016/j.bbadis.2021.166170
123.   Papayianni A, Alexopoulos E, Giamalis P, et al. Circulating levels of ICAM-1, VCAM-1, and
MCP-1 are increased in haemodialysis patients: association with inflammation, dyslipidaemia, and
vascular events. Nephrology Dialysis Transplantation 2002;17(3):435-441, doi:10.1093/ndt/17.3.435
124.   Boardman R, Pang V, Malhi N, et al. Activation of Notch signaling by soluble Dll4 decreases
vascular permeability via a cAMP/PKA-dependent pathway. American Journal of Physiology-Heart and
Circulatory Physiology 2019;316(5):H1065-H1075, doi:10.1152/ajpheart.00610.2018
125.   Scehnet JS, Jiang W, Kumar SR, et al. Inhibition of Dll4-mediated signaling induces proliferation
of immature vessels and results in poor tissue perfusion. Blood 2007;109(11):4753-60,
doi:10.1182/blood-2006-12-063933
126.   Lobov IB, Renard RA, Papadopoulos N, et al. Delta-like ligand 4 (Dll4) is induced by VEGF as a
negative regulator of angiogenic sprouting. Proc Natl Acad Sci U S A 2007;104(9):3219-24,
doi:10.1073/pnas.0611206104
127.   Camelo S, Raoul W, Lavalette S, et al. Delta-like 4 inhibits choroidal neovascularization despite
opposing effects on vascular endothelium and macrophages. Angiogenesis 2012;15(4):609-622,
doi:10.1007/s10456-012-9290-0
128.   Dhamodaran K, Baidouri H, Nartey A, et al. Endogenous expression of Notch pathway molecules
in human trabecular meshwork cells. Exp Eye Res 2022;216(108935, doi:10.1016/j.exer.2022.108935

129.   Khalil M, Pirpamer L, Hofer E, et al. Serum neurofilament light levels in normal aging and their
association with morphologic brain changes. Nature Communications 2020;11(1), doi:10.1038/s41467-
020-14612-6
130.   Taniguchi T, Shimazawa M, Hara H. Alterations in neurofilament light in optic nerve in rat kainate
and monkey ocular hypertension models. Brain Research 2004;1013(2):241-248,
doi:http://dx.doi.org/10.1016/j.brainres.2004.04.023
131.   Markaki I, Bergström S, Tsitsi P, et al. Cerebrospinal Fluid Levels of Kininogen-1 Indicate Early
Cognitive Impairment in Parkinson's Disease. Movement Disorders 2020;35(11):2101-2106,
doi:10.1002/mds.28192
132.   Liu Y, Cao DJ, Sainz IM, et al. The inhibitory effect of HKa in endothelial cell tube formation is
mediated by disrupting the uPA-uPAR complex and inhibiting its signaling and internalization. Am J
Physiol Cell Physiol 2008;295(1):C257-67, doi:10.1152/ajpcell.00569.2007
133.   Sharif NA, Katoli P, Scott D, et al. FR-190997, a Nonpeptide Bradykinin B2-Receptor Partial
Agonist, is a Potent and Efficacious Intraocular Pressure Lowering Agent in Ocular Hypertensive
Cynomolgus Monkeys. Drug Development Research 2014;75(4):211-223,
doi:https://doi.org/10.1002/ddr.21174
134.   Sharif NA, Li L, Katoli P, et al. Preclinical pharmacology, ocular tolerability and ocular hypotensive
efficacy of a novel non-peptide bradykinin mimetic small molecule. Experimental Eye Research
2014;128(0):170-180, doi:http://dx.doi.org/10.1016/j.exer.2014.10.008
135.   Sharif NA, Wang Y, Katoli P, et al. Human Non-Pigmented Ciliary Epithelium Bradykinin B2-
Receptors: Receptor Localization, Pharmacological Characterization of Intracellular Ca2+ Mobilization,
and Prostaglandin Secretion. Current Eye Research 2014;39(4):378-389,
doi:10.3109/02713683.2013.816324
136.   Sharif NA, Xu SX. Pharmacological Characterization of Bradykinin Receptors Coupled to
Phosphoinositide Turnover in SV40-Immortalized Human Trabecular Meshwork Cells. Experimental Eye
Research 1996;63(6):631-637, doi:http://dx.doi.org/10.1006/exer.1996.0157
137.   Webb JG, Shearer TW, Yates PW, et al. Bradykinin enhancement of PGE2 signalling in bovine
trabecular meshwork cells. Experimental Eye Research 2003;76(3):283-289,
doi:http://dx.doi.org/10.1016/S0014-4835(02)00313-5
138.   Webb JG, Yang X, Crosson CE. Expression of the kallikrein/kinin system in human anterior
segment. Experimental Eye Research 2009;89(2):126-132,
doi:http://dx.doi.org/10.1016/j.exer.2009.02.016
139.   Vashishtha A, Maina SW, Altman J, et al. Complement System Proteins in the Human Aqueous
Humor and Their Association with Primary Open-Angle Glaucoma. Journal of Personalized Medicine
2023;13(9):1400, doi:10.3390/jpm13091400
140.   Kim J-H, Afridi R, Han J, et al. Gamma subunit of complement component 8 is a
neuroinflammation inhibitor. Brain 2021;144(2):528-552, doi:10.1093/brain/awaa425
141.   Patel PA, Lee TJ, Kodeboyina SK, et al. Intra-population differences of apolipoproteins in the
aqueous humor. Lipids in Health and Disease 2021;20(1), doi:10.1186/s12944-021-01555-0
142.   Gharahkhani P, Burdon KP, Fogarty R, et al. Common variants near ABCA1, AFAP1 and GMDS
confer risk of primary open-angle glaucoma. Nature Genetics 2014;46(10):1120-1125,
doi:10.1038/ng.3079
143.   Stewart WC, Sine C, Sutherland S, et al. Total Cholesterol and High-Density Lipoprotein Levels
as Risk Factors for Increased Intraocular Pressure. American Journal of Ophthalmology 1996;122(4):575-
577, doi:10.1016/s0002-9394(14)72121-x
144.   Stur M, Grabner G, Dorda W, et al. The effect of timolol on the concentrations of albumin and IgG
in the aqueous humor of the human eye. Am J Ophthalmol 1983;96(6):726-9, doi:10.1016/s0002-
9394(14)71914-2

Condition	Gender	# of patients	Age	IOP (mmHg)
Cataract	Male	16	66.7 ± 10.2	12.4 ± 2.5
	Female	19	67.7 ± 12.2	14.3 ± 2.2
Glaucoma	Male	17	71.2 ± 6.3	19.6 ± 10.7
	Female	17	71.2 ± 9.9	15.4 ± 4.9