Proteomic profile analysis of plasma and aqueous humor from glaucoma and non-glaucomatous patients

Carmen L. Pessuti[1], Chia Ling Huang[2], Angela Banks[2], Nhi Vo[2], Lori Jennings[2], Joseph Loureiro[2], Kleber S. Ribeiro[1], Deise Fialho Costa[1], Heloisa Nascimento[1], Cristina Muccioli[1], Ivan Maynart Tavares[1], Alessandra G. Commodaro[1], Rubens Belfort Jr[1], Christopher W. Wilson[2], Amy Chen[2], Ganesh Prasanna[2], VijayKrishna Raghunathan[2].

[1]Department of Ophthalmology, Universidade Federal de Sao Paulo, Sao Paulo, Brazil

[2]BioMedical Research, Novartis, Cambridge, MA, USA

Corresponding author(s):

VijayKrishna Raghunathan, Ph.D., Ophthalmology, BioMedical Research, Novartis, 22 Windsor St, Cambridge, MA, Massachusetts, 02139. *Email*: Vijay.Raghunathan@novartis.com

Ganesh Prasanna, Ph.D., Ophthalmology, BioMedical Research, Novartis, 22 Windsor St, Cambridge, MA, Massachusetts, 02139. *Email*: ganesh.prasanna@novartis.com

Abstract

Purpose: Glaucoma, a multifactorial ocular neuropathic and age associated disease, can lead

to irreversible vision loss. Diagnosis involves assessing optic cupping (increased cup-to-disc

ratios) and structural changes (like retinal nerve fiber layer thinning) through clinical imaging.

Elevated intraocular pressure (IOP) is commonly associated with glaucoma, but not always.

However, understanding disease progression is hindered by limited access to donor ocular

tissue and consistent clinical data. Our study focuses on live patient samples, analyzing their

proteome for potential biomarkers to enhance precise diagnosis and monitor glaucoma

progression.

Methods: Aqueous humor (AH) samples were collected from 36 glaucoma patients (17 male,

19 female), and 35 non-glaucomatous control patients (16 male, 19 female) undergoing cataract

surgery. The protein profile was compared using the SOMAscan® assay system for proteome

profiling. From glaucomatous donors, significant correlations between IOP and cup-to-disc

ratios to proteome differences were identified.

Results: Correlations in proteins between plasma and AH were identified. These proteins were

enriched in pathways related to vascular integrity, inflammatory response, humoral & adaptive

immune response, cell-cell & cell-matrix adhesion, and complement activation. Glaucomatous

AH exhibited increased protein levels in general. Neurofilament light chain (NEFL) protein

correlated with elevated IOP and inflammatory markers, but not with cup-to-disc ratios.

Conclusions: Together, our data demonstrate that the proteins identified in this study from

glaucomatous donors correspond to both markers of neurodegeneration and those that may

inhibit cell proliferation or disrupt vascular integrity.

Translational Relevance: Proteome analysis may lead to potential biomarker identification

during disease progression.

Precis: Clinical glaucoma diagnosis is by assessing structure-function relationship which

doesn't yield molecular mechanisms underlying disease. Here, we analyzed proteome from

aqueous and plasma of glaucomatous patients. Proteins correspond to markers of

neurodegeneration, pressure and cup-to-disc-ratios.

1. INTRODUCTION

Primary open angle glaucoma (POAG) is one of the leading causes of irreversible blindness, estimated to affect 111.8 million people worldwide by 2040 [1, 2]. Although the etiology and progression of POAG is multifactorial, the only modifiable causal risk factor for managing visual field loss is to reduce intraocular pressure (IOP), which is demonstrated to slow the progression of damage to the optic nerve and death of retinal ganglion cells [3]. IOP is largely determined by a balance between aqueous humor production and resistance to outflow [4, 5]. In the human eye, approximately 75% of all aqueous humor flows from the anterior chamber through the trabecular meshwork (TM) and Schlemm's canal (SC) [6, 7] and is thus the primary egress for aqueous humor [8]. Increased resistance to aqueous drainage due to changes in TM cells and extracellular matrix (ECM) is considered a major contributor to ocular hypertension (OHT) [9, 10]. Differential profiles and levels of soluble factors have been identified in the AH, which have the potential to affect cells in the tissues (outflow tract) that they traverse through, and thus influence OHT.

AH is secreted into the posterior chamber by the non-pigmented epithelial cells of the ciliary processes, and serves to supply critical nutrients and signaling molecules, and to remove metabolic waste from the cornea, lens, and trabecular meshwork. Though the total amount of protein in the anterior chamber is less than 1% (w/v) due to the functional blood aqueous barrier [4], the aqueous humor is enriched with proteins secreted by intraocular structures from both the anterior and posterior segments [11]. It is also thus feasible to hypothesize that systemic disorders, genetic conditions, or perturbations to ocular integrity and health may all adversely influence expression of proteins in the aqueous humor and other ocular tissues [12]. Disease associations of protein compositions of tear films, AH and vitreous humors have been observed with dry eye, glaucoma, age related macular degeneration, or other pathologies [13-18].

Several studies have demonstrated differences between metabolites or proteomes in the AH of patients with primary open angle glaucoma and have subsequently posited that these may serve as biomarkers of dysfunction in the outflow apparatus [16, 19-29]. Complementary approaches

were used to quantify protein expression in AH from glaucomatous donors, including ELISA, multiplexed immunoassays, LC-MS/MS, or antibody microarrays. A few limitations in using the above methods to comprehensively quantify proteins/metabolites are small volume of and limited protein concentration within the AH, as well as the requirement of validated antibody-based approaches, which limits the number of biomarkers that can be identified. There is thus a need to identify the AH proteins more broadly with increased sensitivity, in order to identify biomarkers of disease progression when comparing pre- and post- therapeutic intervention. Further discovery of previously unreported proteins in the AH of patients may provide new insight into the molecular mechanisms underlying outflow resistance in OHT and glaucoma. Here, we utilize the SOMAscan® assay [30], a multiplexed, aptamer-based technology, to detect and assess relative abundance of a broad range of proteins from small volumes of human blood alongside AH from patients undergoing glaucoma surgery, as compared to controls patients subject to cataract surgery.

2. MATERIALS AND METHODS

2.1 Study population

The study was approved by the ethics committee investigational review board (CEP 1.209.725) and adhered to the principles of the Declaration of Helsinki. Informed consent was obtained from all participants. Aqueous humor (AH) and plasma samples were collected from 36 glaucoma patients (17 male, 19 female), and 35 cataract control patients (16 male, 19 female) undergoing cataract surgery (**Figure 1**). Within the glaucoma patient group one underwent trabeculectomy, two were pseudophakic, and AH was collected at the beginning of the surgery. Clinical diagnosis of glaucoma was based on elevated IOP (IOP > 21 mmHg), documented progressive cupping or thinning of the neuroretinal rim. All glaucoma patients were treated with medication to control IOP. Patients with other eye diseases or associated with systemic diseases were not included.

2.2 Aqueous humor (AH) and plasma sample collection

AH samples were collected from patients following a previously validated method [31] at the

beginning of the surgery. Briefly, after anesthetic and antiseptic eyedrops administration, a 1mL

tuberculin syringe (27 gauge, ½ inch) was used to aspirate the AH. Approximately, 0.1 ml was

collected, and transferred to two labeled Eppendorf tubes. Peripheral blood (10 mL) was

collected from an arm of the patient in EDTA tubes during the surgery. The tubes were

centrifuged for 10 min at 1900 x g, and plasma was collected. Total protein concentrations in the

AH samples were determined using the BCA Protein Assay Kit (Pierce, Rockford, IL).

2.3 SOMAscan® analysis

All AH and plasma samples were analyzed in parallel using the SOMAscan® proteome profiling

platform that provides a broad and relative abundant assessment of selected proteins in

biological fluid. The SOMAscan® platform, using affinity-based slow off-rate modified aptamers

(SOMAmers), has been described extensively elsewhere [30, 32, 33]. The custom SOMAscan®

version used in our study covered 5,080 SOMAmers, including 4,785 human-specific somamers

targeting 4,161 human proteins. The targeted proteins include transmembrane receptors,

secreted proteins, kinases, transcription regulators and signal transducers. The samples were

aliquoted into 2D barcoded tubes and were shipped on dry ice to SomaLogic for analysis.

2.4 Statistical analysis

The relative abundance of captured SOMAmers were quantified by microarray hybridization as

relative fluorescence units (RFU). We focused on SOMAmers detecting human proteins in this

study which focuses on 4,785 SOMAmers for the follow up analysis. Quality control and sample

outlier detection was conducted using the arrayQualityMetrics[34], and by running principal

component analysis (PCA) and Spearman's correlation analysis with custom R scripts. RFU

were log2-transformed and normalized by smooth quantile normalization using R package

qsmooth [35]. In total, 5 samples (2 non-glaucomatous, 3 glaucoma) were removed from the AH

cohort and 2 non-glaucomatous samples were removed from the plasma cohort, which left 62

samples in the AH cohort and 65 samples in the AH cohort. We applied the limma R package

(v3.50.3) in R 4.1.0 environment to determine differentially expressed proteins between

groups[36]. Pathway enrichment analysis was performed with pre-ranked GSEA using R package

fgsea v1.20.0 and Gene Ontology Biological Process category (GOBP) in MSigDB[37-40].

2.5 Correlation analysis

We conducted Elastic Net, a machine learning based method, to identify proteins correlated with

the total protein concentration. For this analysis, we utilized the cv.glmnet function from the R

package glmnet v4.1-3. We employed a 10-fold cross-validation approach to optimize the alpha

and lambda parameters. Additionally, we performed a 1000-fold bootstrapping using the

optimized alpha and lambda values to determine the frequency at which each protein was

selected as a top predictor for estimating the total protein concentration. To enhance the

performance of our analysis, we only included proteins with an absolute Pearson's correlation

coefficient greater than 0.4 with the total protein concentration.

3. RESULTS

3.1 Donor demographics

Donor demographics for the patient samples used in this study are provided in **Figure 1**. The

average age of donors with cataract was 66.9 ± 10.9 while those with glaucoma was 68.41 ±

12.84. No significant differences (p=0.60, t-test; **Figure 1**) were seen in donor ages between the

groups *in toto* or when evaluated by sex. Amongst these donors, we note that one donor had

primary angle closure glaucoma while the remainder had POAG. The mean IOP of glaucoma

donors (IOP controlled with medication) was 17.94 ± 8.28 mmHg with no differences seen

between male or female donors. Only one POAG donor had an IOP > 24 mmHg. A statistically

significant difference in total protein concentration in AH was observed between cataract and

glaucoma groups.

3.2 Differential analysis of SOMAScan® profiling data

Principal component analysis of the cataract and glaucoma samples revealed heterogeneity

between samples, including outliers that were subsequently excluded from further analysis.

Subsequent analysis revealed few differences in protein levels between non-glaucomatous and

glaucomatous aqueous humor and plasma (**Supplementary Fig. 1**). Nevertheless, we

observed that samples were clustered by tissue types with reasonable correlation (r > 0.5)

between AH and plasma samples (**Figure 2a**). The histogram (**Figure 2b**) demonstrates that

the distribution of Pearson's correlation coefficients between AH and plasma samples from

same donor demonstrates some variability. Considering all donors, a reasonable correlation (r >

0.5) was observed comparing protein levels between AH and plasma samples.

Subsequent to multivariate analysis, we performed differential protein analysis for both

plasma and aqueous samples. Expression values that were |log2(RFU)| > 0.5 and where the

adjusted p-value was < 0.05 are highlighted as significant (**Figure 3**). Within glaucomatous AH

(Figure 3A), 7 SOMAmer probes representing 6 target proteins were observed to be significantly

down regulated, while 49 SOMAmer probes representing 49 target proteins were upregulated.

Examples of top up-regulated proteins include complement proteins (C3, C7, C1QTNF3),

Apoliprotein (APOE, APOF), matrix protein (endostatin/Col18A1), macrophage associated

proteins (CSF1R, CD163), while top down-regulated proteins include growth factors such as

FGF9, FGF20 and CCK. We observed that in plasma samples, the majority of the differential

proteins were upregulated in glaucoma. Some of the upregulated proteins of interest include

ENO1, NEK7, AKT, PRKCA, PRKCB, JAK2, MAP kinase-activated protein kinase 2

(MAPKAPK2), whereas the top down-regulated proteins include APOC3, GDF2, DCP1B and

EDIL3. A comprehensive list of proteins altered in their levels comparing glaucomatous and

non-glaucomatous plasma and AH are provided in **supplementary table 1**. Although protein

levels from AH and plasma from the same donor showed a positive correlation, we only

observed SNX4 to be significantly up regulated in both AH and plasma samples from

glaucomatous donors.

3.3 Pathway enrichment analysis

Next, we asked if the differentially enriched proteins were enriched in common biological pathways, by performing pathway analysis (**supplementary table 2**). We identified 51 and 559 significantly up-regulated pathways (adjusted p-value < 0.05) in AH and plasma samples from glaucoma donors, respectively. 16 pathways were up-regulated in both AH and plasma, including inflammatory response, IL6 production and immune response related pathways.

3.4 Correlation analysis of protein levels with total protein concentration

While individual protein changes may help identify signaling pathways perturbed in disease, total protein concentration of the AH may help ascertain the overall health of the blood-aqueous barrier and the outcomes of IOP lowering medications. Thus, normalizing individual proteins to total protein concentrations assists in determining if observed changes reflect pathway changes underpinning disease, or are merely a consequence of structural failure or drug treatment(s). We conducted an Elastic Net regression analysis to identify proteins that can accurately predict the total protein concentration (**Figure 4a**). Our analysis specifically focused on AH proteins, as proteins from these samples may provide a better reflection of changes in glaucoma disease compared to plasma samples. Through our analysis, we successfully identified 5 proteins (KNG1, IGFBO6, MAP2K4, H6PD, and C3) that demonstrated the highest predictive capability for the total protein concentration collected from the patients (**Figure 4a**).

3.5 Correlation analysis of protein levels with IOP

We conducted Elastic net regression analysis to identify 13 proteins to be associated with IOP values reported in glaucomatous patients (**Figure 4b**). We note that a glaucoma sample with extremely high IOP measurement was excluded from the analysis. Among the proteins of interest in glaucoma, the protein with maximal correlation with IOP levels in glaucomatous patients was neurofilament light chain (NEFL; **Figure 4c, Supplementary Fig. 2**). Linear regression analysis validated this finding and demonstrated a net correlation with IOP

measurements (R^2 = 0.3067). Since NEFL is a widely accepted marker for several

neurodegenerative diseases including glaucoma [41-50], we then decided to further investigate if

levels of NEFL correlated with other SOMAmers identified in glaucomatous donors. We

calculated the Pearson's correlation coefficient of NEFL and rest of the proteins and identified

29 proteins that are positively or negatively correlated with NEFL (|r > 0.4|), including TEAD3,

CNTF, C4A, C4B and CCL27 (**Figure 5**).

3.6 Changes in protein levels as a function of cup-to-disc ratios in glaucoma

Finally, we sought to determine if changes in relative abundance of protein levels in AH

correlated with cup-to-disc ratios of glaucomatous patients via Pearsons's correlation analysis

(**Figure 6**). For patients where cup-to-disc ratios were available for both eyes, we used the

average measurements from both eyes. We note that among the 29 glaucomatous samples

obtained, cup-to-disc ratios were available for 26 donors. Figure 6 shows the top 6 positive

(KLRG2, C8G, PENK, ALB, APOA1, and KNG1) and negative (IL12A/B, LRRC3, ZAP70,

CTSG, DEFB115, and NSG1) correlated proteins with cup-to-disc ratios.

4. Discussion

Diagnosis and prognosis of glaucoma disease is clinically monitored by IOP, and structure

function measurements (advanced imaging, and visual field testing etc). However, these require

longitudinal measurements supported by patient compliance, adherence to medications and

regular in-patient visits. While these are critically valuable to ascertain the stage of glaucoma, in

the absence of longitudinal ocular and systemic medical history, very little information is gleaned

about molecular mechanisms underlying the disease or help in identifying a soluble biomarker

to ascertain therapeutic engagement and efficacy. Since obtaining intraocular tissues, such as

retina and optic nerve, from live human donors is not possible, reliance on sampling biofluids

(such as AH and blood) to ascertain changes in the secretome that may further our standing of

homeostasis/disease, is favored. In this study, we determined the abundance of proteins in both

the AH and blood plasma from patients (with and without clinically diagnosed glaucoma)

undergoing cataract surgery and correlated them with disease-relevant clinical measures: IOP

and cup-to-disc ratio.

The ages and sex of glaucomatous and non-glaucomatous donors in this dataset were

comparable. We observed significant changes in relative abundance of a subset of proteins

identified both in AH and plasma. Protein levels from AH and plasma from the same donor

sample showed a positive correlation (r > 0.5 for all of samples). Interestingly, when comparing

protein levels and correlations across all fluids and disease states, we only observed a

significant upregulation of SNX4 , a synaptic & endosomal membrane trafficking in the secretory

pathway and mitochondrial recycling quality protein in aging [51-53], upregulated in glaucomatous

AH and plasma when compared with those from non-glaucomatous donors. The lack of greater

correlation between plasma and AH proteins may be due to several reasons: turnover rates of

AH *vs* plasma, tissues & cell types involved in protein secretions, systemic *vs* local transport of

proteins. It is yet unclear if glaucoma is purely a disease of the eye or if systemic factors exist

that regulate its etiology and progression, although co-morbidities have been reported [54-58].

Nevertheless, in the plasma, 48 proteins were upregulated while 4 (EDIL3, DCP1B, GDF2,

APOC3) were downregulated when comparing non-glaucomatous and glaucomatous donors.

Gene ontology analyses of proteins altered in the plasma of glaucomatous donors were related

to multiple pathways corresponding to protein & macromolecule localization, intracellular

transport, kinase activity pathways, post translational / post transcriptional regulation of proteins

/ genes, cytoskeletal reorganization and GPCR related pathways. On the other hand, 7 (ISLR2,

CCK, INSL4, FGF9, FGF20, BMPR1A) proteins were significantly downregulated and 49 were

significantly upregulated in the AH of glaucomatous donors. Gene ontology analyses of proteins

identified as altered in the AH of glaucomatous donors were related to multiple pathways

corresponding to inflammatory response, humoral & adaptive immune response, cell-cell & cell-

matrix adhesion, complement activation, and wound healing responses. These together suggest

that though overall concordance in proteins may be observed between AH and plasma, some

differences exist and thus proximity of biofluid to relevant tissue being studied (i.e. ciliary

process, ciliary muscle and trabecular meshwork/Schlemm's canal) may be important to

consider. As such, deeper analysis in this study was focused on the ocular specific biofluid (AH)

proteins.

All changes in protein levels were normalized to total protein concentration and as such

these changes are independent of mean protein concentration. However, interestingly

regression modeling demonstrated that total protein concentrations trended towards being

significantly greater in AH of glaucomatous compared with non-glaucomatous donors. While this

is suggestive of a breakdown in the blood-aqueous barrier [4], whether this is due to the chronic

nature of the disease or due to IOP lowering medications, other systemic medications or other

co-morbidities is unclear. Nevertheless, the changes in proteins appear to be directly related to

pathways with the potential for consequential effects on IOP regulation and outflow

homeostasis.

For example, endostatin (Col18A1), which was upregulated in AH of glaucomatous donors,

is expressed ubiquitously in ocular tissues except in photoreceptors, and protein fragments that

contain endostatin has been observed to accumulate in ocular fluid samples [59]. Interestingly,

endostatin has been reported to promote expression and release of thrombospondin-1,

implicated in outflow resistance & glaucoma [60-62], in non-ocular endothelial cells [63]. Conversely,

endostatin has also been posited to crosstalk with Rho/ROCK, TGFβ, NF-κB, PDGF, and

autophagy pathways to yield anti-fibrotic effects [64]. SVEP1, another overexpressed secreted

ECM protein in AH & a disease modifier allele for congenital glaucoma [65], is known to be a

binding partner for Tie1 and can thus regulate signaling outcomes in lymphatics and vasculature

[66]. Since AH predominantly drains into the Schlemm's canal, a unique vessel with both

vasculature and lymphatic properties [67], studying SVEP1 in the context of glaucoma warrants

further investigation. Of particular interest, VCAM1, CDH11, and BGN were other proteins that

were overexpressed in the AH of glaucomatous patients. SVEP1, CDH11 and VCAM1 were all

identified as genes relevant to POAG pathogenesis in a genome-wide meta-analysis across

ancestries [68]. CDH11 can interact reciprocally with fibronectin binding protein syndecan-4 to

facilitate cell migration and adhesion, partake in EMT, and modulate proliferation [69-72]. In non-

ocular cells, CDH11 was shown to mediate adhesion of macrophages to fibroblasts promoting

transdifferentiation into myofibroblasts and a self-sustaining profibrotic niche [73]. Biglycan (BGN)

is an extracellular proteoglycan, expressed in the trabecular meshwork [74], is posited as a

prognostic marker for cancer aggressiveness [75], fibrosis & inflammation [76] and is predicted to be

a circulating "messenger" for triggering inflammation and/or autophagy [77]. Whether biglycan in

the AH serves as a biomarker, a signaling molecule to trigger autophagy or inflammatory

phenotypes in the trabecular meshwork to mediate subsequent outflow regulation requires

mechanistic studies. High levels of circulatory VCAM1 in the blood/plasma was found to

correlate with ventricular hypertrophy, hypertension and is suggested to thus serve as a

predictive soluble biomarker for cardiovascular disease and inflammation [78-80]. In this study,

whether VCAM1 expression is a result of chronic elevated IOP, drugs to lower IOP, or due to

undiagnosed/unknown underlying systemic vascular conditions is unknown. Regardless,

increased presence of the aforementioned proteins may serve as 'proteins-of-interest' in

mechanistic understanding of outflow regulation for as potential biomarkers for dysregulation in

outflow homeostasis.

Since proteins with associations with 'hypertension' were identified, we next sought to

determine if protein changes were a function of IOP levels through additional regression

modeling. Two proteins notably were found to be of importance: DLL4 and NEFL. DLL4, a

Notch ligand, is a key regulator of vascular morphogenesis, vessel maturation, and function.

Secreted/soluble DLL4 has previously been reported to significantly reduce hydraulic

conductivity, vascular permeability, and disrupt endothelial barrier function in non-ocular

vasculature [81, 82]. Further, DLL4 has been demonstrated to be critical in developing retinal

vasculature, increased endothelial cell proliferation, and angiogenic sprouting [83]. Interestingly,

DLL4 was also shown to inhibit inflammatory choroidal neovascularization despite opposing

effects seen in endothelial cells (anti-angiogenic) and macrophages (pro-angiogenic) [84].

Collectively, we speculate that DLL4, found upregulated with IOP, may act to increase outflow

resistance via a yet unknown mechanism. Further, whether a dichotomous function for the DLL4

ligand or the Notch pathway exists in outflow regulation remains to be seen. A recent study

suggests that Notch pathway protein expression may differ between segmental flow regions of

healthy and glaucomatous donors, at least *in vitro* [85]. Thus, ligands circulating in the aqueous

humor could differentially impact TM cell function necessitating further studies.

NEFL is a well-established marker of neurodegeneration [42, 44-47, 86] and similar to our finding,

has previously been reported to be elevated in glaucomatous AH in humans and animal models

[41, 43, 87]. In fact, NEFL had the highest correlation with IOP in all donors in this study, providing

additional confidence that NEFL in the AH may indeed be a suitable biomarker for

glaucomatous neurodegeneration and ocular hypertension. However, we advise caution that AH

samples utilized in this study were from patients identified for glaucoma filtration surgery.

Therefore, it is likely that the disease stage may be advanced and thus markers of axonal

degeneration are expected within the AH. Whether NEFL would serve as an early marker of

disease is unclear and thus requires comprehensive longitudinal natural history investigations.

Further the source of the neurofilament protein, determined in the aqueous humor, is unclear

(i.e. whether it is due to degeneration of the optic nerve, secretion from cells of neural crest

origin (developmentally) in the anterior segment, or Schwann/nerve cells in the anterior segment

remains unknown). Interestingly, while NEFL levels correlated with IOP, its levels did not appear

to correlate with cup-to-disc ratios of glaucoma donors (**Supplementary Fig. 3**), though other

proteins that did correlate were identified. Interestingly, cup-to-disc ratio was independent of

IOP values as well (**Supplementary Fig. 3**). Inflammatory proteins (such as interleukins and

cathepsins) corresponded with low cup-to-disc ratio, while apolipoprotein A1 (APOA1), albumin,

kininogen-1 (KNG1) and complement C8G correlated with a higher cup-to-disc ratio suggesting

proteins altered in the AH may differ based on disease severity. KNG1 is an antiangiogenic

molecule that has been suggested to be a marker of neurodegeneration [88] and can impair the

proliferation of endothelial cells [89]. To the best of our knowledge, the role of KNG1 in POAG

and/or ocular hypertension has not been studied, although its cleaved nonapeptide, Bradykinin

(BK), has been the target for IOP lowering investigations [90-95]. A prior study reported a decrease

in C8G [96] in glaucomatous AH conflicting with our results. However, Kim et al [97] previously

reported C8G to act as a neuroinflammation inhibitor; our result correlating C8G levels with high

cup-to-disc ratios may thus reflect the advanced stage of neurodegeneration/neuroinflammation

in glaucoma.

Consistent with our study, elevated levels of APOA1 were reported in AH of POAG donors

[98]. It is important to note that whether a direct role of APOA1 exists in IOP homeostasis is not

known. However, APOA1 plays a critical role in reverse cholesterol transport pathway via direct

interactions with the ABCA1 gene whose variants are implicated in POAG [99]. Cholesterol is itself

a risk factor for elevated IOP [100]. Elevation in albumin levels in glaucomatous AH is also

reported in several studies, though attributed to administration of IOP lowering drugs [101]. Finally,

a correlation analysis of NEFL to other proteins identified in glaucomatous aqueous humor

revealed that NELF levels may be associated with proteins regulating apoptosis, complement

activation, proliferation, and cytoskeletal reorganization. Together, the proteins identified in this

study correspond to both markers of neurodegeneration and those that may inhibit proliferation

or vascular integrity.

5. Summary

In this study, we demonstrate the ability of the SOMAScan® proteomics platform to identify

novel and previously reported proteins in the AH and plasma of glaucomatous donors. We

report that proteins identified in AH in part significantly correlate directly with those identified in

plasma. However, we also note that POAG patients were treated with IOP lowering drugs, and

as such the effect of medication on proteins identified cannot be delineated. Further, we

identified aqueous humor proteins that correlated with either IOP elevation or with reported cup-

to-disc ratios. Finally, our data comprehensively demonstrate that proteins identified in diseased

AH may be indicative of markers of inflammation, neurodegeneration, and dysregulation of

extracellular matrix. Together, we speculate that proteins identified in this study may be useful

to the glaucoma scientific community to help identify new proteins/genes of interest and further

our understanding in the identification of molecular biomarkers and the mechanisms underlying

ocular hypertension and glaucoma.

6. Limitations

This study is not without limitations. One of the major limitations is that the biofluids obtained in

this study were obtained at the time of ocular surgery thus representing a singular snapshot in

time and no information is gleaned about the stage of the disease; although, IOP measurements

and cup-to-disc ratios are available at this stage. Further a comprehensive and longitudinal

history of systemic co-morbidities or ocular diseases and medications that may affect systemic

or ocular hypertension including but limited to intraocular pressure is unknown. Since IOP in

glaucoma patients were controlled with IOP lowering medication, the effect of drugs from

disease on proteome changes cannot be adequately unraveled. Further functional

measurements for visual field or additional structural deficits including longitudinal and pre-

surgical fundus photography or OCT measurements (nerve fiber layer thinning, rim width,

ganglion cell analysis etc.) are not known for these patients.

7. Acknowledgements

The authors would like to thank the patient donors for the biological fluid samples without whose

consent these experiments would be impossible. This work was sponsored by Novartis

Biomedical Research, and CNPq (Ministry of Science, Technology, and Innovation of Brazil).

Figure Legends

Figure 1: Patient demographics demonstrate study participants were age, sex matched. Intraocular pressures are reported only for glaucomatous donors.

Figure 2: Proteins identified in aqueous humor and plasma clustered by type of biofluid, and appear to correlate reasonably well as evidenced by Spearman's correlation. **(A)** Heatmap, and **(B)** histogram of Spearman's correlation index demonstrates that the correlation, while evident, is not absolute i.e. $0.5 < R^2 < 0.7$.

Figure 3: Principal component (PCA) and differential expression analysis demonstrates differences in proteins identified in **(A)** aqueous humor and **(B)** plasma comparing glaucomatous and non-glaucomatous donors. PCA demonstrates large variance between cataract (non-glaucoma) and glaucomatous donor samples with no discernible separation between the two disease groups. Volcano plots demonstrate that no significant differences were seen comparing cataract (non-glaucoma) and glaucomatous donors for most proteins. Few proteins whose levels were significantly altered are represented in the bar plot.

Figure 4: Elastic net regression analysis (aqueous humor only) demonstrated **(A)** increased protein concentration is observed in aqueous humor of donors with glaucoma, with no specific sex related differences observed, **(B)** changes in some proteins correlate significantly with IOP independent of sex of donor. **(C)** Scatter plots and linear regression of proteins ($R^2 > 0.25$) as a function of IOP indicate NEFL, DLL4, and NFE2L1 to most significantly altered proteins in the aqueous humor.

Figure 5: Heatmap demonstrates correlation of NEFL to top 30 proteins identified in the aqueous humor of glaucomatous donors. Pearson's correlation analysis comparing NEFL with rest of the proteins identified ~30 proteins that either positively (red) or negatively (blue) correlated with NEFL ($|r > 0.4|$). In general, more proteins that negatively correlated with NEFL levels were identified than those that exhibited positive correlation.

Figure 6: Heatmap identifies top 6 proteins either up or down regulated in the aqueous humor of glaucomatous donors as a function of cup-to-disc ratios.

Supplementary figure 1: Outlier and principal component analysis identifying donors to exclude from data set in an unbiased manner.

Supplementary figure 2: Scatter plots of proteins identified in glaucomatous AH as a function of IOP demonstrate large variability between samples.

Supplementary figure 3: (A) Scatter plot of NEFL expression levels as a function of cup-to-disc ratios of glaucomatous donors demonstrate no significant relationship ($R^2 = 0.037$) between the two parameters. **(B)** Cup-to-disc ratios of glaucomatous donors show no significant correlation ($R^2 = 0.008$) with intraocular pressures reported.

References

1.   Tham Y-C, Li X, Wong TY, Quigley HA, Aung T, Cheng C-Y. Global Prevalence of Glaucoma and Projections of Glaucoma Burden through 2040: A Systematic Review and Meta-Analysis. *Ophthalmology* 2014;121:2081-2090.

2.   Allison K, Patel D, Alabi O. Epidemiology of Glaucoma: The Past, Present, and Predictions for the Future. *Cureus* 2020.

3.   Glaucoma. In: Jackson TL (ed), *Moorfields Manual of Ophthalmology.* Edinburgh: Mosby; 2008:274-330.

4.   Freddo TF. A contemporary concept of the blood-aqueous barrier. *Prog Retin Eye Res* 2013;32:181-195.

5.   Freddo TF, Johnson M. Chapter 6 Aqueous Humor Outflow Resistance. *The Eye's Aqueous Humor*, 2008:161-192.

6.   Toris CB, Yablonski ME, Wang Y-L, Camras CB. Aqueous humor dynamics in the aging human eye. *American journal of ophthalmology* 1999;127:407-412.

7.   Toris CB, Koepsell SA, Yablonski ME, Camras CB. Aqueous humor dynamics in ocular hypertensive patients. *Journal of glaucoma* 2002;11:253-258.

8.   Stamer WD, Acott TS. Current understanding of conventional outflow dysfunction in glaucoma. *Curr Opin Ophthalmol* 2012;23:135-143.

9.   Acott TS, Kelley MJ. Extracellular matrix in the trabecular meshwork. *Exp Eye Res* 2008;86:543-561.

10.   Vranka JA, Kelley MJ, Acott TS, Keller KE. Extracellular matrix in the trabecular meshwork: intraocular pressure regulation and dysregulation in glaucoma. *Exp Eye Res* 2015;133:112-125.

11.   Yorio T, Prasanna G, Coca-Prados M. CHAPTER 4 - The Ciliary Body: A Potential Multifaceted Functional Neuroendocrine Unit. In: Yorio T, Clark AF, Wax MB (eds), *Ocular Therapeutics.* London: Academic Press; 2008:69-85.

12.   Ahmad MT, Zhang P, Dufresne C, Ferrucci L, Semba RD. The Human Eye Proteome Project: Updates on an Emerging Proteome. *PROTEOMICS* 2018;18:1700394.

13.   Baek J-H, Lim D, Park KH, et al. Quantitative proteomic analysis of aqueous humor from patients with drusen and reticular pseudodrusen in age-related macular degeneration. *BMC Ophthalmology* 2018;18.

14.   Aghamollaei H, Parvin S, Shahriary A. Review of proteomics approach to eye diseases affecting the anterior segment. *Journal of Proteomics* 2020;225:103881.

15.   Skeie JM, Roybal CN, Mahajan VB. Proteomic Insight into the Molecular Function of the
Vitreous. *PLOS ONE* 2015;10:e0127567.

16.   Kang GY, Bang JY, Choi AJ, et al. Exosomal proteins in the aqueous humor as novel
biomarkers in patients with neovascular age-related macular degeneration. *J Proteome Res*
2014;13:581-595.

17.   Dammeier S, Martus P, Klose F, et al. Combined Targeted Analysis of Metabolites and
Proteins in Tear Fluid With Regard to Clinical Applications. *Translational Vision Science &*
Technology 2018;7:22-22.

18.   De Almeida Borges D, Alborghetti MR, Franco Paes Leme A, et al. Tear proteomic
profile in three distinct ocular surface diseases: keratoconus, pterygium, and dry eye related to
graft-versus-host disease. *Clinical Proteomics* 2020;17.

19.   Agnifili L, Pieragostino D, Mastropasqua A, et al. Molecular biomarkers in primary open-
angle glaucoma: from noninvasive to invasive. *Prog Brain Res* 2015;221:1-32.

20.   Babizhayev MA. Biomarkers and special features of oxidative stress in the anterior
segment of the eye linked to lens cataract and the trabecular meshwork injury in primary open-
angle glaucoma: challenges of dual combination therapy with N-acetylcarnosine lubricant eye
drops and oral formulation of nonhydrolyzed carnosine. *Fundam Clin Pharmacol* 2012;26:86-
117.

21.   Bollinger KE, Crabb JS, Yuan X, Putliwala T, Clark AF, Crabb JW. Quantitative
proteomics: TGFbeta(2) signaling in trabecular meshwork cells. *Invest Ophthalmol Vis Sci*
2011;52:8287-8294.

22.   Burgos-Blasco B, Vidal-Villegas B, Saenz-Frances F, et al. Tear and aqueous humour
cytokine profile in primary open-angle glaucoma. *Acta Ophthalmol* 2020;98:e768-e772.

23.   Castany M, Jordi I, Catala J, et al. Glaucoma patients present increased levels of
diadenosine tetraphosphate, Ap4A, in the aqueous humour. *Experimental Eye Research*
2011;92:221-226.

24.   Csosz E, Deak E, Toth N, Traverso CE, Csutak A, Tozser J. Comparative analysis of
cytokine profiles of glaucomatous tears and aqueous humour reveals potential biomarkers for
trabeculectomy complications. *FEBS Open Bio* 2019;9:1020-1028.

25.   Garweg JG, Zandi S, Pfister IB, Skowronska M, Gerhardt C. Comparison of cytokine
profiles in the aqueous humor of eyes with pseudoexfoliation syndrome and glaucoma. *PLoS*
One 2017;12:e0182571.

26.   Guo T, Guo L, Fan Y, et al. Aqueous humor levels of TGFbeta2 and SFRP1 in different
types of glaucoma. *BMC Ophthalmol* 2019;19:170.

27.   Igarashi N, Honjo M, Asaoka R, et al. Aqueous autotaxin and TGF-betas are promising
diagnostic biomarkers for distinguishing open-angle glaucoma subtypes. *Sci Rep* 2021;11:1408.
28.   Lin JB, Sheybani A, Santeford A, De Maria A, Apte RS. Increased Aqueous Humor
GDF15 Is Associated with Worse Visual Field Loss in Pseudoexfoliative Glaucoma Patients.
Transl Vis Sci Technol 2020;9:16.
29.   Sun C, Zhang H, Jiang J, et al. Angiogenic and inflammatory biomarker levels in
aqueous humor and vitreous of neovascular glaucoma and proliferative diabetic retinopathy. *Int
Ophthalmol* 2020;40:467-475.
30.   Gold L, Ayers D, Bertino J, et al. Aptamer-Based Multiplexed Proteomic Technology for
Biomarker Discovery. *PLoS ONE* 2010;5:e15004.
31.   Belfort R, Jr., Moura NC, Mendes NF. T and B lymphocytes in the aqueous humor of
patients with uveitis. *Arch Ophthalmol* 1982;100:465-467.
32.   Mehan MR, Ostroff R, Wilcox SK, et al. Highly multiplexed proteomic platform for
biomarker discovery, diagnostics, and therapeutics. *Adv Exp Med Biol* 2013;735:283-300.
33.   Gold L, Walker JJ, Wilcox SK, Williams S. Advances in human proteomics at high scale
with the SOMAscan proteomics platform. *N Biotechnol* 2012;29:543-549.
34.   Kauffmann A, Gentleman R, Huber W. arrayQualityMetrics--a bioconductor package for
quality assessment of microarray data. *Bioinformatics* 2009;25:415-416.
35.   Hicks SC, Okrah K, Paulson JN, Quackenbush J, Irizarry RA, Bravo HC. Smooth
quantile normalization. *Biostatistics* 2018;19:185-198.
36.   Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for
RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015;43:e47.
37.   Aleksander SA, Balhoff J, Carbon S, et al. The Gene Ontology knowledgebase in 2023.
GENETICS 2023;224.
38.   Ashburner M, Ball CA, Blake JA, et al. Gene Ontology: tool for the unification of biology.
Nature Genetics 2000;25:25-29.
39.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. *Bioinformatics* 2011;27:1739-1740.
40.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Jill, Tamayo P. The Molecular
Signatures Database Hallmark Gene Set Collection. *Cell Systems* 2015;1:417-425.
41.   Lin JB, Pitts KM, El Helwe H, et al. Neurofilament Light Chain in Aqueous Humor as a
Marker of Neurodegeneration in Glaucoma. *Clinical Ophthalmology* 2023;Volume 17:2209-
2217.

42.   Kaeser SA, Lehallier B, Thinggaard M, et al. A neuronal blood marker is associated with
mortality in old age. *Nature Aging* 2021;1:218-225.

43.   Sasaki Y, Engber TM, Hughes RO, et al. cADPR is a gene dosage-sensitive biomarker
of SARM1 activity in healthy, compromised, and degenerating axons. *Experimental Neurology*
2020;329:113252.

44.   Baka P, Steenken L, Escolano-Lozano F, et al. Studying serum neurofilament light chain
levels as a potential new biomarker for small fiber neuropathy. *Eur J Neurol* 2024;e16192.

45.   Fissolo N, Benkert P, Sastre-Garriga J, et al. Serum biomarker levels predict disability
progression in patients with primary progressive multiple sclerosis. *J Neurol Neurosurg*
Psychiatry 2023.

46.   Frank A, Bendig J, Schnalke N, et al. Serum neurofilament indicates accelerated
neurodegeneration and predicts mortality in late-stage Parkinson's disease. *NPJ Parkinsons Dis*
2024;10:14.

47.   Franthal S, Khalil M, Kern D, et al. Elevated serum neurofilament light chain protein in
patients with essential tremor. *Eur J Neurol* 2024;31:e16143.

48.   Novobilsky R, Bartova P, Licha K, Bar M, Stejskal D, Kusnierova P. Serum
neurofilament light chain levels in patients with cognitive deficits and movement disorders:
comparison of cerebrospinal and serum neurofilament light chain levels with other biomarkers.
Front Hum Neurosci 2023;17:1284416.

49.   Sano T, Masuda Y, Yasuno H, Shinozawa T, Watanabe T. Plasma neurofilament light
chain as a potential biomarker of neurodegeneration in murine brain. *Toxicol Res (Camb)*
2023;12:751-755.

50.   Yang Y, Kim WS, Michaelian JC, et al. Predicting neurodegeneration from sleep related
biofluid changes. *Neurobiol Dis* 2024;190:106369.

51.   Seaman MNJ. Membrane traffic in the secretory pathway. *Cellular and Molecular Life*
Sciences 2008;65:2842-2858.

52.   Vazquez-Sanchez S, Gonzalez-Lozano MA, Walfenzao A, Li KW, Van Weering JRT.
The endosomal protein sorting nexin 4 is a synaptic protein. *Scientific Reports* 2020;10.

53.   Henkel V, Schürmanns L, Brunner M, Hamann A, Osiewacz HD. Role of sorting nexin
PaATG24 in autophagy, aging and development of Podospora anserina. *Mechanisms of Ageing*
and Development 2020;186:111211.

54.   Dada T, Verma S, Gagrani M, et al. Ocular and Systemic Factors Associated with
Glaucoma. *J Curr Glaucoma Pract* 2022;16:179-191.

55.   Zhang T, Xie X, Lu F. Primary open-angle glaucoma: Neuroendocrine–immune
disorder? *Medical Hypotheses* 2014;83:514-515.

56.   Salim S, Shields MB. Glaucoma and Systemic Diseases. *Survey of ophthalmology*
2010;55:64-77.

57.   Pache M, Flammer J. A Sick Eye in a Sick Body? Systemic Findings in Patients with
Primary Open-angle Glaucoma. *Survey of ophthalmology* 2006;51:179-212.

58.   Nucci C, Martucci A, Cesareo M, et al. Links among glaucoma, neurodegenerative, and
vascular diseases of the central nervous system. Elsevier; 2015:49-65.

59.   Määttä M, Heljasvaara R, Pihlajaniemi T, Uusitalo M. Collagen XVIII/endostatin shows a
ubiquitous distribution in human ocular tissues and endostatin-containing fragments accumulate
in ocular fluid samples. *Graefe's Archive for Clinical and Experimental Ophthalmology*
2006;245:74-81.

60.   Haddadin RI, Oh DJ, Kang MH, et al. Thrombospondin-1 (TSP1)-null and TSP2-null
mice exhibit lower intraocular pressures. *Invest Ophthalmol Vis Sci* 2012;53:6708-6717.

61.   Shan S-W, Do C-W, Lam TC, Li H-L, Stamer WD, To C-H. Thrombospondin-1 mediates
Rho-kinase inhibitor-induced increase in outflow-facility. *Journal of Cellular Physiology*
2021;236:8226-8238.

62.   Wirtz MK, Sykes R, Samples J, et al. Identification of Missense Extracellular Matrix
Gene Variants in a Large Glaucoma Pedigree and Investigation of the N700S Thrombospondin-
1 Variant in Normal and Glaucomatous Trabecular Meshwork Cells. *Curr Eye Res* 2022;47:79-
90.

63.   Goyanes AM, Moldobaeva A, Marimoutou M, et al. Functional Impact of Human Genetic
Variants of COL18A1/Endostatin on Pulmonary Endothelium. *American Journal of Respiratory
Cell and Molecular Biology* 2020;62:524-534.

64.   Zhang Z, Liu X, Shen Z, et al. Endostatin in fibrosis and as a potential candidate of anti-
fibrotic therapy. *Drug Delivery* 2021;28:2051-2061.

65.   Young TL, Whisenhunt KN, Jin J, et al. SVEP1 as a Genetic Modifier of TEK-Related
Primary Congenital Glaucoma. *Invest Ophthalmol Vis Sci* 2020;61:6.

66.   Hußmann M, Schulte D, Weischer S, et al. Svep1 is a binding ligand of Tie1 and affects
specific aspects of facial lymphatic development in a Vegfc-independent manner. *eLife*
2023;12:e82969.

67.   Kizhatil K, Ryan M, Marchant JK, Henrich S, John SWM. Schlemm's Canal Is a Unique
Vessel with a Combination of Blood Vascular and Lymphatic Phenotypes that Forms by a Novel
Developmental Process. *PLoS Biology* 2014;12:e1001912.

68. Gharahkhani P, Jorgenson E, Hysi P, et al. Genome-wide meta-analysis identifies 127
open-angle glaucoma loci with consistent effect across ancestries. *Nat Commun* 2021;12:1258.

69. Langhe RP, Gudzenko T, Bachmann M, et al. Cadherin-11 localizes to focal adhesions
and promotes cell-substrate adhesion. *Nat Commun* 2016;7:10909.

70. Schneider DJ, Wu M, Le TT, et al. Cadherin-11 contributes to pulmonary fibrosis:
potential role in TGF-β production and epithelial to mesenchymal transition. *The FASEB Journal*
2012;26:503-512.

71. Passanha FR, Divinagracia ML, Lapointe VLS. Cadherin-11 Regulates Cell Proliferation
via the PDGFRβ-ERK1/2 Signaling Pathway in Human Mesenchymal Stem Cells. *Stem Cells*
2022;40:165-174.

72. Passanha FR, Geuens T, LaPointe VLS. Cadherin-11 Influences Differentiation in
Human Mesenchymal Stem Cells by Regulating the Extracellular Matrix Via the TGFβ1
Pathway. *Stem Cells* 2022;40:669-677.

73. Lodyga M, Cambridge E, Karvonen HM, et al. Cadherin-11–mediated adhesion of
macrophages to myofibroblasts establishes a profibrotic niche of active TGF-β. *Science*
Signaling 2019;12:eaao3469.

74. Wirtz MK, Bradley JMB, Xu H, et al. Proteoglycan expression by human trabecular
meshworks. *Current Eye Research* 1997;16:412-421.

75. Pinto F, Santos-Ferreira L, Pinto M, Gomes C, Reis C. The Extracellular Small Leucine-
Rich Proteoglycan Biglycan Is a Key Player in Gastric Cancer Aggressiveness. *Cancers*
2021;13:1330.

76. Schulz M, Diehl V, Trebicka J, Wygrecka M, Schaefer L. Biglycan: A regulator of
hepatorenal inflammation and autophagy. *Matrix Biol* 2021;100-101:150-161.

77. Poluzzi C, Nastase MV, Zeng-Brouwers J, et al. Biglycan evokes autophagy in
macrophages via a novel CD44/Toll-like receptor 4 signaling axis in ischemia/reperfusion injury.
Kidney Int 2019;95:540-562.

78. Kuroda Y. Vascular cell adhesion molecule-1 as a biochemical marker of left ventricular
mass in the patients with hypertension. *American Journal of Hypertension* 2001;14:868-872.

79. Troncoso MF, Ortiz-Quintero J, Garrido-Moreno V, et al. VCAM-1 as a predictor
biomarker in cardiovascular disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of*
Disease 2021;1867:166170.

80. Papayianni A, Alexopoulos E, Giamalis P, et al. Circulating levels of ICAM-1, VCAM-1,
and MCP-1 are increased in haemodialysis patients: association with inflammation,
dyslipidaemia, and vascular events. *Nephrology Dialysis Transplantation* 2002;17:435-441.

81.   Boardman R, Pang V, Malhi N, et al. Activation of Notch signaling by soluble Dll4
decreases vascular permeability via a cAMP/PKA-dependent pathway. *American Journal of*
Physiology-Heart and Circulatory Physiology 2019;316:H1065-H1075.

82.   Scehnet JS, Jiang W, Kumar SR, et al. Inhibition of Dll4-mediated signaling induces
proliferation of immature vessels and results in poor tissue perfusion. *Blood* 2007;109:4753-
4760.

83.   Lobov IB, Renard RA, Papadopoulos N, et al. Delta-like ligand 4 (Dll4) is induced by
VEGF as a negative regulator of angiogenic sprouting. *Proc Natl Acad Sci U S A*
2007;104:3219-3224.

84.   Camelo S, Raoul W, Lavalette S, et al. Delta-like 4 inhibits choroidal neovascularization
despite opposing effects on vascular endothelium and macrophages. *Angiogenesis*
2012;15:609-622.

85.   Dhamodaran K, Baidouri H, Nartey A, et al. Endogenous expression of Notch pathway
molecules in human trabecular meshwork cells. *Exp Eye Res* 2022;216:108935.

86.   Khalil M, Pirpamer L, Hofer E, et al. Serum neurofilament light levels in normal aging and
their association with morphologic brain changes. *Nature Communications* 2020;11.

87.   Taniguchi T, Shimazawa M, Hara H. Alterations in neurofilament light in optic nerve in
rat kainate and monkey ocular hypertension models. *Brain Research* 2004;1013:241-248.

88.   Markaki I, Bergström S, Tsitsi P, et al. Cerebrospinal Fluid Levels of Kininogen-1
Indicate Early Cognitive Impairment in Parkinson's Disease. *Movement Disorders*
2020;35:2101-2106.

89.   Liu Y, Cao DJ, Sainz IM, Guo YL, Colman RW. The inhibitory effect of HKa in
endothelial cell tube formation is mediated by disrupting the uPA-uPAR complex and inhibiting
its signaling and internalization. *Am J Physiol Cell Physiol* 2008;295:C257-267.

90.   Sharif NA, Katoli P, Scott D, et al. FR-190997, a Nonpeptide Bradykinin B2-Receptor
Partial Agonist, is a Potent and Efficacious Intraocular Pressure Lowering Agent in Ocular
Hypertensive Cynomolgus Monkeys. *Drug Development Research* 2014;75:211-223.

91.   Sharif NA, Li L, Katoli P, et al. Preclinical pharmacology, ocular tolerability and ocular
hypotensive efficacy of a novel non-peptide bradykinin mimetic small molecule. *Experimental*
Eye Research 2014;128:170-180.

92.   Sharif NA, Wang Y, Katoli P, Xu S, Kelly CR, Li L. Human Non-Pigmented Ciliary
Epithelium Bradykinin B2-Receptors: Receptor Localization, Pharmacological Characterization
of Intracellular Ca2+ Mobilization, and Prostaglandin Secretion. *Current Eye Research*
2014;39:378-389.

93.   Sharif NA, Xu SX. Pharmacological Characterization of Bradykinin Receptors Coupled to
Phosphoinositide Turnover in SV40-Immortalized Human Trabecular Meshwork Cells.
Experimental Eye Research 1996;63:631-637.

94.   Webb JG, Shearer TW, Yates PW, Mukhin YV, Crosson CE. Bradykinin enhancement of
PGE2 signalling in bovine trabecular meshwork cells. *Experimental Eye Research* 2003;76:283-
289.

95.   Webb JG, Yang X, Crosson CE. Expression of the kallikrein/kinin system in human
anterior segment. *Experimental Eye Research* 2009;89:126-132.

96.   Vashishtha A, Maina SW, Altman J, et al. Complement System Proteins in the Human
Aqueous Humor and Their Association with Primary Open-Angle Glaucoma. *Journal of*
Personalized Medicine 2023;13:1400.

97.   Kim J-H, Afridi R, Han J, et al. Gamma subunit of complement component 8 is a
neuroinflammation inhibitor. *Brain* 2021;144:528-552.

98.   Patel PA, Lee TJ, Kodeboyina SK, et al. Intra-population differences of apolipoproteins
in the aqueous humor. *Lipids in Health and Disease* 2021;20.

99.   Gharahkhani P, Burdon KP, Fogarty R, et al. Common variants near ABCA1, AFAP1
and GMDS confer risk of primary open-angle glaucoma. *Nature Genetics* 2014;46:1120-1125.

100.   Stewart WC, Sine C, Sutherland S, Stewart JA. Total Cholesterol and High-Density
Lipoprotein Levels as Risk Factors for Increased Intraocular Pressure. *American Journal of*
Ophthalmology 1996;122:575-577.

101.   Stur M, Grabner G, Dorda W, Zehetbauer G. The effect of timolol on the concentrations
of albumin and IgG in the aqueous humor of the human eye. *Am J Ophthalmol* 1983;96:726-
729.

Demographics

Condition	Gender	# of patients	Age	IOP (mmHg)
Cataract	Male	16	66.7 ± 10.2	-n/a-
	Female	19	67.7 ± 12.2	-n/a-
Glaucoma	Male	17	71.2 ± 6.3	19.6 ± 10.7
	Female	17	71.2 ± 9.9	15.4 ± 4.9

(A)

(B)