The landscape of regional missense mutational intolerance quantified from 125,748 exomes

Katherine R. Chao[1,2]*, Lily Wang[1,2,3]*, Ruchit Panchal[1,2], Calwing Liao[4,5,6], Haneen Abderrazzaq[7], Robert Ye[4,5], Patrick Schultz[1,4,5], John Compitello[1,4,5], Riley H. Grant[1], Jack A. Kosmicki[3,4,5], Ben Weisburd[1,2], William Phu[1,2], Michael W. Wilson[1,2], Kristen M. Laricchia[1,2], Julia K. Goodrich[1,2], Daniel Goldstein[1,4,5], Jacqueline I. Goldstein[1,4,5], Christopher Vittal[1,4,5], Timothy Poterba[1,4,5], Samantha Baxter[1], Nicholas A. Watts[1,4], Matthew Solomonson[1,4], gnomAD Consortium, Grace Tiao[1,2], Heidi L. Rehm[1,2], Benjamin M. Neale[1,4], Michael E. Talkowski[1,2,5], Daniel G. MacArthur[1,8,9], Anne O'Donnell-Luria[1,2,10], Konrad J. Karczewski[1,4,5], Predrag Radivojac[7], Mark J. Daly[1,4,11], Kaitlin E. Samocha[1,2,4]

[1]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[2]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
[3]Bioinformatics and Integrative Genomics Program, Harvard Medical School, Boston, MA, USA
[4]Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
[5]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[6]Department of Medicine, Harvard Medical School, Boston, MA, USA
[7]Khoury College of Computer Sciences, Northeastern University, Boston, MA, USA
[8]Centre for Population Genomics, Garvan Institute of Medical Research and UNSW Sydney, Sydney, New South Wales, Australia
[9]Centre for Population Genomics, Murdoch Children's Research Institute, Melbourne, Victoria, Australia
[10]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA
[11]Institute for Molecular Medicine Finland (FIMM), Helsinki, Finland

*Indicates equal contribution

Correspondence should be addressed to K.E.S. (samocha@broadinstitute.org)

Abstract

Missense variants can have a range of functional impacts depending on factors such as the

specific amino acid substitution and location within the gene. To interpret their deleteriousness,

studies have sought to identify regions within genes that are specifically intolerant of missense

variation[1–12]. Here, we leverage the patterns of rare missense variation in 125,748 individuals in

the Genome Aggregation Database (gnomAD)[13] against a null mutational model to identify

transcripts that display regional differences in missense constraint. Missense-depleted regions

are enriched for ClinVar[14] pathogenic variants, *de novo* missense variants from individuals with

neurodevelopmental disorders (NDDs)[15,16], and complex trait heritability. Following ClinGen

calibration recommendations for the ACMG/AMP guidelines, we establish that regions with less

than 20% of their expected missense variation achieve moderate support for pathogenicity. We

create a missense deleteriousness metric (MPC) that incorporates regional constraint and

outperforms other deleteriousness scores at stratifying case and control *de novo* missense

variation, with a strong enrichment in NDDs. These results provide additional tools to aid in

missense variant interpretation.

Main text

Over the last decade, exome and genome sequencing have enabled variant discovery across

hundreds of thousands of individuals[13,17–21]. These large reference databases have provided the

opportunity to study selective forces acting on the human genome and to identify genomic

regions under selective constraint by, for example, identifying regions with fewer variants than

expected based on mutational models[13,18,22–25]. Gene-level metrics of predicted loss-of-function

(pLoF) variant depletion have proven to be valuable in variant classification and identification of

novel disease genes[15,16,26–28]. The functional impact and selective pressures relevant to

missense variation, by contrast, remain challenging to predict, as the effect of a missense

variant is governed by the gene housing the variant, the position of the variant in the gene, and

the specific amino acid substitution caused by the variant. To address this, prior work has

sought to identify regions within coding genes that are specifically intolerant of missense

variation as a way to improve interpretation[1–12]. Here, we expand upon previous work[1] and show

a sub-genic measure of missense intolerance leveraging population-level variation facilitates

variant classification and risk stratification for association studies with *de novo*, rare, and

common variants.

We explored the patterns of rare missense variant presence or absence in 125,748 exomes in

the Genome Aggregation Database (gnomAD) v2.1.1 on GRCh37 to quantify missense

depletion at the sub-genic level. We searched 18,629 canonical protein-coding transcripts for

variability in missense constraint, quantified as the number of rare (allele frequency [AF] <

0.1%) missense variants observed in gnomAD divided by the number expected under neutral

evolution as estimated from previously described mutational models[13](observed/expected [OE]).

For each transcript, we applied a recursive search based on likelihood ratio tests over all

potential rare missense sites looking for breaks that divide the transcript coding sequence

(CDS) into distinct missense constraint regions (MCRs; **Fig. 1a, b**). We discover 5,127

transcripts (28%) harbor regional variability in missense constraint (**Fig. 1c**), i.e., have two or

more MCRs (minimum coding length 49bp, median 461bp; **Supplementary Fig. 1**). We thus

refine the resolution of missense constraint for 42% of coding sites (coding space in the 5,127

transcripts vs. 18,629 total assessed). After recalibrating the missense OE distribution over all

potential sites of missense variants using MCR-wide rather than transcript-wide missense OE

measurements, we discover widespread signatures of negative and neutral selection that are

obscured when quantifying over the unit of whole transcripts (**Fig. 1d**). We find a larger

proportion of the exome lies within strongly constrained sequences (5.6% vs. 1.7% at OE < 0.4;

see **Supplementary Note** for OE threshold selection), and the mode of the distribution shifts

toward an OE indicative of evolutionary neutrality at approximately 1 (40.6% vs. 36.5% at 0.9 <

OE ≤ 1.1).

Fig. 1: 28% of protein-coding genes in the human genome are discovered to harbor regional

variation in population-level missense depletion.

a, An example transcript that has two missense constraint regions (MCRs) with significantly

different levels of population-wide missense depletion, defined as the number of missense

variants observed in gnomAD at rare frequency (AF < 0.1%) divided by the number of rare

missense variants expected under neutral evolution (observed/expected or OE). Lower OE

values correspond to greater variant depletion in the population and suggest stronger constraint.

b, Flow chart describing the process of searching for breakpoints that divide a transcript into

multiple MCRs. Searching for breakpoints is recursive and leverages likelihood ratio tests at a

significance threshold of p = 0.001. **c**, The number of MCRs within the 5,127 transcripts

discovered to harbor regional differences in missense constraint. The other 13,502 transcripts

are deemed to have a single MCR (that is, a constant level of constraint across their entirety)

and are not shown. **d**, The distribution of local missense OE at all coding sites in canonical

transcripts. Local missense OE is defined as the OE calculated over the whole transcript (for

"transcript OE") or over the MCR (for "MCR OE") where the site is located. Transcript OE and
MCR OE are equivalent for transcripts with one MCR.
73
74
Furthermore, we find that constrained MCRs overlap established disease-associated mutational
hotspots, including critical protein domains. One example is in the well-characterized *KCNQ1*, a
voltage-gated potassium channel gene, in which pathogenic variants cause cardiac disorders
such as long QT syndrome. We discover one moderately constrained MCR (missense OE =
0.60) overlapping the highly conserved C-terminus [29] and another (missense OE = 0.66)
encompassing the voltage-sensing and pore domains (**Fig. 2a**). Both the C-terminus of *KCNQ1*
and its voltage-sensing domain are established "hotspot" regions (specific missense-
constrained regions with ACMG/AMP hotspot/functional domain moderate support [PM1] for
pathogenicity)[29–31]. All but two ClinVar pathogenic/likely pathogenic (P/LP) missense variants in
this gene fall within these two missense-constrained MCRs.
85
We also find that missense constraint within MCRs is able to identify regions associated with
severe, early-onset disease. One example is in *BAP1*, which plays a key role in chromatin
modeling by mediating histone deubiquitination. Disease-causing variants in this gene are linked
to cancer or, as recently discovered, Kury-Isidor syndrome[32]. The first highly missense-
constrained MCR (missense OE = 0.33) in *BAP1* encompasses the ubiquitin C-terminal
hydrolase domain connected to Kury-Isidor syndrome[32] (**Fig. 2b**), and all 11 variants reported to
be causal for Kury-Isidor fall within this MCR. The only ClinVar P/LP variants that do not fall
within any missense-constrained MCRs in *BAP1* are associated with cancer phenotypes, which
may be under weaker selection than neurodevelopmental disorders (NDDs).
95

96
Fig. 2: Missense constraint regions (MCRs) and the distribution of ClinVar pathogenic/likely
pathogenic (P/LP) missense variants in two genes associated with early-onset developmental
disorders. Exons are delineated with black outlines and MCRs are delineated by color. MCRs
are colored based on their missense observed/expected (OE) ratio, and MCRs with missense
OEs not significantly different from 1 (p > 0.001) are shaded gray. **a,** *KCNQ1*. Only two of the
210 P/LP missense variants in *KCNQ1* do not fall within either constrained MCR. The first
constrained MCR encompasses the voltage-sensing and pore domains of this gene, and the
other constrained MCR overlaps the C-terminus. Both domains contain previously reported
hotspot regions, with some regions reaching moderate level (PM1) support for pathogenicity[31].
Ion transport protein: domain that contains both the transmembrane voltage-sensing and pore
domains. KCNQ channel: C-terminal cytoplasmic domain that overlaps four helices (A-D). **b,**
BAP1. Variants in this gene can lead to cancer-predisposition syndromes, increased risk of
certain cancers, or the neurodevelopmental disorder Kury-Isidor syndrome[32]. All of the ClinVar
P/LP variants associated with Kury-Isidor fall within the first MCR with a highly depleted
missense OE of 0.33. An additional five variants reported in Kury *et al.*[32] but not ClinVar fall
within either highly constrained MCR in this gene. P/LP variants associated with Kury-Isidor are
colored in black, and all other cancer-associated P/LP variants are colored in gray. UCH:
Ubiquitin carboxyl-terminal hydrolase isozyme L5 domain. ClinVar data are from the October 15,
2023 release.

116
117
Next, we sought to determine if the signatures of selection revealed by MCRs recapitulated
biological and disease relevance of coding sequences. Overall, most transcripts that are

intolerant to pLoF variation (as measured by the loss-of-function observed/expected upper

bound fraction [LOEUF] score[13]) also tend to be intolerant to missense variation. This trend is

markedly more prominent when measuring missense constraint at the sub-genic level vs. the

transcript-level (**Supplementary Note; Supplementary Fig. 2**). We also discovered that 64%

(1697/2659) of genes that are both LOEUF- and MCR missense-constrained do not have

disease associations in OMIM[33], suggesting the existence of many undocumented genes

containing variants of significant consequence for disease (**Supplementary Fig. 3**). In a set of

730 strongly mutationally intolerant genes, defined here as exhibiting both population depletion

of pLoF variants (first three LOEUF deciles) and association with a developmental phenotype

(high-confidence membership in any non-cancer Gene2Phenotype [G2P][34] gene list with

dominant inheritance), we observed strong transcript-wide missense depletion that was even

stronger for genes with multiple MCRs (**Fig. 3a** and **Supplementary Fig. 4**; Wilcoxon $p < 10^{-50}$).

Given that we have greater power to detect missense constraint variability over longer

sequences (**Supplementary Fig. 5**), we controlled for transcript length but still found that

intolerant transcripts are eight times more likely to harbor multiple MCRs ($p < 10^{-50}$). These

strongly intolerant transcripts are highly enriched for severely depleted regions (three times

more likely to have minimum MCR OE < 0.4 after regressing out transcript length, $p < 10^{-18}$),

whereas the most constrained MCRs in not strongly intolerant transcripts are less depleted and

more evenly distributed across the OE spectrum. Finally, we observe a group of genes with

strong overall missense depletion in which we did not detect multiple MCRs (n = 459 with

missense OE < 0.4; **Supplementary Table 1**), suggesting these genes are robustly intolerant to

missense variants across their length. When comparing missense constraint to selection over

longer timescales (measured by evolutionary conservation in placental mammals, phyloP[35]), we

found that genes with more conserved coding sequences also tended to be more overall

depleted of human missense variation (Spearman $\rho = 0.56$, $p < 10^{-50}$). However, a substantial

number of strongly constrained MCRs appear widely unconserved across mammals, potentially

pointing to human-specific negative selection pressures that are obscured at the whole-

transcript level (**Supplementary Fig. 6**).

We next aggregated *de novo* missense variants from 31,058 individuals with a severe

developmental disorder[15] (DD), 15,036 autistic individuals (AUT), and 5,492 siblings not

diagnosed with a DD[16] (**Fig. 3b**). The distribution of *de novo* missense variants across the

missense OE spectrum in unaffected siblings largely mirrored the exome-wide missense OE

distribution. In contrast, *de novo* missense variants in autistic individuals are enriched in

missense-constrained sequences, and this pattern is more striking in individuals with DDs. For

example, relative to unaffected siblings, the rate of *de novo* missense variants in MCRs with OE

< 0.2 is 2-fold higher in autistic individuals ($p < 10^{-23}$) and 6.6-fold higher in individuals with DDs

($p < 10^{-50}$) (**Supplementary Fig. 7;** see **Supplementary Note** for OE threshold selection). This

is consistent with the expectations that a small subset of *de novo* missense variants in

individuals with developmental phenotypes are causal for those traits and that variants causal

for DD are generally more selectively deleterious than those for autism.

Beyond large-effect rare and *de novo* variation in traits under strong negative selection, we

additionally investigated whether our MCR metric, which was calculated using rare variants,

correlates with functional effects of common variants. Prior work found that pLoF-constrained

genes and their flanking 100kb sequences are enriched for SNP heritability across hundreds of

independent traits in the UK Biobank (UKBB) and other large genome-wide association studies

(GWAS) [13]. We partitioned common (AF > 5%) variant heritability of the same 268 independent

traits across MCRs to investigate relative enrichment. To establish a baseline, we computed the

heritability enrichment over all coding sequences comprising MCRs (3-fold). The most

constrained MCRs have the strongest heritability enrichment; the first quintile of MCR missense

OE harbors a 41-fold enrichment (**Fig. 3c**). Coding SNPs in missense-unconstrained MCRs

(e.g., in the two least constrained quintiles of MCR missense OE) harbor no detectable

heritability enrichment relative to the average genome-wide SNP. These findings suggest that:

1) regions depleted of rare missense variation can help prioritize common coding variants

important for complex traits (i.e., improve GWAS fine-mapping variant prioritization), and 2)

there exists a subset of coding sequence with no appreciable heritability enrichment, which rare

variant depletion can help identify.

180
Fig. 3: Regional missense depletion reveals constraint obscured by gene-level measures. **a**,
Left: The distribution of transcript-wide missense observed/expected (OE) across 18,629
transcripts stratified by the combination of two factors: whether the transcript is strongly
mutationally intolerant (within first three LOEUF deciles and association with a developmental
phenotype in Gene2Phenotype [G2P][34]) and whether we detect multiple missense constraint
regions (MCRs). Number of transcripts in each category are: strongly intolerant with multiple
MCRs (n=581; red), strongly intolerant with one MCR (n=149; yellow), not strongly intolerant
with multiple MCRs (n=4,546; dark gray), not strongly intolerant with one MCR (n=13,353; light
gray). X-axis is cut off at 1.6 for visibility. Right: Minimum MCR missense OE using the same
groupings. Minimum MCR missense OE is the same as transcript missense OE for transcripts
with a single MCR. **b**, MCR missense OE at all sites of possible exome-wide missense variants
vs. sites of *de novo* missense variants in controls, autistic individuals (AUT), or individuals with
DD. *De novo* variants from individuals with developmental phenotypes are enriched in more
constrained sequences, with a more pronounced enrichment in DD than autism. **c**, Enrichment
in per-variant heritability explained by common (AF > 5%) protein-coding SNPs stratified by
MCR missense OE quintile, relative to the average SNP genome-wide. Enrichment is estimated
by linkage disequilibrium score regression, accounting for number of SNPs in each quintile, and

is averaged across 268 independent traits in UKBB and other large genome-wide association
studies. Black dashed line at 1 indicates no enrichment. Blue dashed line at 3 indicates average
coding enrichment. Error bars represent 95% confidence intervals.
201
202
We examined the localization of high-quality ClinVar[36] missense variants classified as P/LP
within genes with both unconstrained (missense OE > 0.9) and constrained (missense OE <
0.2) MCRs and found that P/LP variants occur much more frequently in missense constrained
MCRs (odds ratio [OR] = 15.2; $p < 10^{-50}$). We also examined the localization of P/LP and
benign/likely benign (B/LB) variants within MCRs in autosomal dominant disease-associated
genes and found that P/LP variants tend to localize to regions that are more strongly missense-
constrained than the overall transcript (Wilcoxon $p = 3.5 \times 10^{-10}$), while B/LB variants show the
opposite effect and tend to occur in regions with OEs closer to 1 (Wilcoxon $p < 10^{-18}$; **Fig. 4a**).
While more subtle, these same patterns are also significant in autosomal recessive disease-
associated genes (**Supplementary Fig. 8**).
213
To enable use of our missense constraint metric in ACMG/AMP clinical variant classification, we
applied previously established probabilistic frameworks[37] to determine the MCR missense OE
thresholds that met different levels of clinical evidence strengths evaluated under the
hotspot/functional domain (PM1) and benign *in silico* prediction (BP4) criteria codes[30]. MCR
missense OE ≤ 0.37 met supporting (PM1_Supporting) and OE ≤ 0.21 met moderate (PM1)
levels of evidence for pathogenicity (**Fig. 4b**), but no MCR missense OE threshold met any
levels of evidence to support benignity. However, separate calibration specifically in transcripts
with multiple MCRs found that MCR missense OE ≥ 1.56 met moderate and OE ≥ 0.97 met
supporting evidence for BP4, indicating that in transcripts where we are powered to characterize
regional constraint, MCRs with OEs close to one harbor an indication of benignity
(**Supplementary Fig. 9**). Calibration of two additional regional constraint metrics, Constrained
Coding Regions (CCRs[9]) and COntact Set MISsense tolerance (COSMIS[12]), which incorporates
predicted 3D structure information, revealed that these metrics also reach moderate support for
pathogenicity (PM1), and COSMIS only reaches supporting levels for benignity
(**Supplementary Fig. 9; Supplementary Table 2**).
229
230

Fig. 4: ACMG/AMP calibration of missense constraint. **a,** The distribution within genes with autosomal dominant disease associations of transcript-wide missense observed/expected (OE; gray) and missense constraint region (MCR) OE for ClinVar pathogenic/likely pathogenic missense variants (P/LP; red) and benign/likely benign missense variants (B/LB; blue). We filtered to 1,007 transcripts with at least one P/LP and one B/LB missense variant. For the P/LP and B/LB distributions, we annotated each variant with the missense OE across the MCR they fell in and collapsed these values within each transcript by taking the respective medians. **b,** Local posterior probabilities of pathogenicity given MCR missense OE in all transcripts. Gray shading indicates the one-sided 95% confidence interval on the more stringent side. Horizontal lines indicate thresholds required to meet ACMG/AMP evidence levels. From bottom to top: supporting, moderate, strong, very strong. MCR missense OE reaches supporting (OE ≤ 0.37) and moderate (OE ≤ 0.21) level evidence for PM1 (hotspot/functional domain).

We transformed our regional missense constraint measure into a variant-level predictor of missense deleteriousness named MPC (Missense deleteriousness Prediction by Constraint) that additionally incorporates information about amino acid substitution type and local context. The logistic regression-based model integrates regional missense constraint-derived metrics together with BLOSUM[38], Grantham[39], and PolyPhen-2[40] and is trained on ClinVar pathogenic and gnomAD common (AF > 0.1%) variants in 2,987 genes defined as haploinsufficient in Collins *et al*.[41] and 366 genes with DD associations in G2P through non-LoF mechanisms. Higher scores predict greater deleteriousness (**Supplementary Fig. 10, 11**). We assessed the utility of MPC in prioritizing potentially disease-causing variation by evaluating its ability to stratify case and control rare and *de novo* missense variation. Consistent with the regional constraint results, the *de novo* missense variants from DD and AUT cases are enriched for high MPC scores compared to controls (**Supplementary Fig. 12**). We further stratified by presence in 373 genes previously associated with NDD[16] and three bins of MPC scores (< 1.6, 1.6-2.6, ≥ 2.6; see **Supplementary Note** for calibration of these bins), and found a very strong enrichment of *de novo* missense variants in the two most deleterious bins among both the DD (**Fig. 5a**) and

AUT cases (**Fig. 5b**) compared to unaffected individuals. However, while the enrichment in the

373 NDD-associated genes was significant for missense variants with MPC ≥ 2.6 (RR in DD

cases = 22.7, p < 10^{-50}; RR in AUT cases = 6.9, p < 10^{-21}), as well as missense variants with

MPC between 1.6-2.6 (RR in DD cases = 4.5, p < 10^{-35}; RR in AUT cases = 1.9, p = 3.0×10^{-5}), it

was only significant in NDD-unassociated genes for missense variants with MPC ≥ 2.6 (RR in

DD cases = 3.1, p < 10^{-28}; RR in AUT cases = 1.5, p = 5.9×10^{-4}). This suggests that while there

is a sizable reservoir of potentially causal variants in genes yet to be associated with NDDs,

they will be more difficult to find as they must reach stricter deleteriousness criteria. For autism,

we additionally assessed inheritance rates of rare missense variants (AF < 0.1%) from parents

to 13,384 probands and case-control rates for an additional 5,591 cases and 8,597 controls

without *de novo* information. While we did not find substantial enrichment in inheritance rates in

any missense category, we discovered substantial enrichment in the case-control analysis for

variants in the 373 NDD-associated genes with MPC ≥ 2.6 (RR = 1.6, p < 10^{-12}), which we infer

is from *de novo* variants that are not recognizable as such due to lack of parental information.

We extended our assessment of case-control *de novo* stratification for a comparison of our

model against several other missense deleteriousness predictors: AlphaMissense[42], CCRs[9], M-

CAP[43], REVEL[44], PrimateAI-3D[45], MVP[46], Polyphen-2[40], CADD[47,48], mammalian conservation

phyloP[35], and SIFT[49]. For this assessment, we evaluated four additional early-onset

development-related phenotypes: epileptic encephalopathy (EE), orofacial cleft (OFC),

congenital heart disease (CHD), and congenital diaphragmatic hernia (CDH). To compare

across predictors with different score distributions, we used a ranking-based performance

assessment. For each predictor, we ranked the *de novo* missense variants from each case

cohort against those in the 5,492 controls and computed the OR of case vs. control variants in

the top percentiles of these rankings (**Fig. 5c**). At the top 10% of variants, MPC displays the

highest OR for DD (OR = 5.2, Fisher's exact p < 10^{-48}), EE (OR = 3.1, p = 2.2×10^{-7}), AUT (OR =

1.7, p = 8.9×10^{-9}), and OFC (OR = 1.5, p = 0.025), although there is substantial confidence

interval overlap with other predictors. This indicates that MPC effectively ranks high-impact *de*

novo variants in the most deleterious prediction regimes. Of the other predictors,

AlphaMissense also performs consistently well across all phenotypes. In particular, in CHD and

CDH, which have the least *de novo* enrichment across predictors, we observe MPC lagging in

performance, while AlphaMissense is one of the top performers. This may suggest that causal

de novo variants in these phenotypes may occur at a narrow set of sites where 3D structure is

important, which AlphaMissense can more deftly capture through integration of protein structure

prediction. These observations are more or less consistent over a range of thresholds used to

define the ranking top percentiles (**Supplementary Fig. 13**).

Fig. 5: MPC effectively stratifies case and control variation.

a, The difference relative to controls of missense variants stratified by MPC score and

localization to genes associated with neurodevelopmental disorders (NDDs) for **a,** individuals

with DD and **b**, autistic individuals (AUT). Relative difference is calculated as: for *de novo*

variants, the average rate of variants in probands divided by that in sibling controls; for case-

control, the average rate of variants in cases divided by that in controls from case-control data;

for inherited, the average rate in probands of transmitted variants divided by that of

untransmitted variants. Error bars represent 95% confidence intervals calculated from a

binomial test. **c**, The odds ratio of case to control *de novo* missense variants in the top 10% vs.

bottom 90% of respective rankings. *De novo* missense variants from each case cohort are

ranked against those in the 5,492 controls for each predictor. DD: developmental disorders, EE:

epileptic encephalopathy, AUT: autism, OFC: orofacial cleft, CHD: congenital heart disease,

CDH: congenital diaphragmatic hernia. Error bars represent 95% confidence intervals. Only

variants scored by all predictors are included. Points are solid colored if the difference from 1 is

statistically significant (binomial or Fisher exact p < 0.05).

We have developed a method to identify sub-genic regions with differential intolerance to

missense variation at base-level resolution. We demonstrate that coding regions depleted for

missense variation in the general population are enriched for established disease-associated

variation, *de novo* variants from individuals with NDDs, and heritability for 268 complex traits

from the UK Biobank and other large GWAS. Additionally, we have calibrated these constraint

scores to establish that regions with less than 20% of their expected variation can achieve

moderate evidence for association to disease following ACMG/AMP guidelines. Finally, we

incorporated regional missense intolerance information into the missense deleteriousness

metric, MPC, and show that MPC effectively separates potentially risk-carrying variants

identified in various developmental disorder cases from those seen in controls.

At current sample sizes, we are unable to characterize constraint at single amino acid

resolution. Furthermore, because our approach relies on variant presence or absence in a large

reference dataset, many of the constrained regions we find are linked to variants that cause

severe, early-onset disease. However, the true nature of the variation we capture is more

accurately linked to reproductive fitness and the strength of selection acting on heterozygotes[50].

Our methodology specifically searches for linear sub-genic regions in canonical transcripts that

are depleted of missense variants compared to a null mutational model. This means that our

model is unable to find depleted sequences that are clustered specifically in 3D space and is

also currently ignorant of coding sequences not present in the Ensembl canonical transcript.

However, we note that our linear metric achieves similar evidence for both pathogenicity and

benignity as the structural constraint-based COSMIS model[12] (**Supplementary Fig. 9**).

In summary, we identify 28% of canonical transcripts with variable levels of missense constraint

and demonstrate that coding regions specifically depleted of missense variation in the general

population are enriched for disease-associated variation. Additionally, we show that this

depletion of missense variation can be used as moderate evidence when classifying variants

according to ACMG/AMP guidelines and that incorporation of regional missense constraint into

an *in silico* predictor effectively prioritizes a subset of *de novo* missense variation in individuals

with developmental phenotypes for association testing. We have publicly released these data

for use in both research and clinical settings. We anticipate refined resolution of these metrics

as datasets grow, both in size and in ancestral diversity, and with the incorporation of

complementary structural or functional data.

Methods

Transcripts

This study analyzed only canonical, coding transcripts as defined by GENCODE v19/Ensembl
v74. We excluded the same set of transcripts from this analysis that were excluded in the
previous gnomAD v2.1.1 genic constraint estimates[13]. Briefly, we excluded transcripts that had
outlier counts of variants expected under neutrality (zero expected pLoF, missense, or
synonymous variants; too many observed pLoF, missense, or synonymous variants compared
to expectation; or too few observed synonymous variants compared to expectation). In total, this
study analyzed 18,629 transcripts.
359

gnomAD variants

All analyses in this paper were conducted using the 125,748 gnomAD v2.1.1 exomes[13] on
GRCh37. Median coverage was calculated on a random subset of the gnomAD exomes as
described previously[13]. We defined the set of sites with possible missense variants using a
synthetic Hail Table (HT) containing all possible single nucleotide variants in the exome. We
annotated this HT with the Variant Effect Predictor (VEP, version 85) against GENCODE
version 19, and filtered to variants with the consequence "missense_variant" in the canonical,
coding transcripts as defined in *Transcripts*. We then further filtered to variants that fit one of the
following criteria: (1) allele count (AC) > 0 and AF < 0.001, variant QC PASS, and median
coverage > 0 in gnomAD v2.1.1 exomes; or (2) AC = 0, i.e. variants not seen in gnomAD v2.1.1
exomes.
371

372 ClinVar variants

373 We annotated functional consequences for ClinVar[14] (v.20230305) variants using the VEP table
374 described in *gnomAD variants.* Missense ClinVar variants with non-conflicting P, LP, B, LB
375 classification and a review status of at least one star were selected for analysis.
376

Rare and *de novo* variants from developmental cohorts

Case *de novo* mutations for association analyses were obtained from studies of developmental
disorders[15] (DD), autism[16] (AUT), congenital heart disease[51] (CHD), orofacial cleft[52] (OFC),
congenital diaphragmatic hernia[53] (CDH), and epileptic encephalopathy[54] (EE). Control *de novo*
mutations were obtained from neurotypical siblings of the autistic probands[16]. Variants from the
autism study were lifted over from GRCh38 to GRCh37 using the "liftover" function in Hail.
Variant functional consequences were re-annotated using the VEP table described in *gnomAD*
variants. Variants transmitted and not transmitted from parents to autistic probands were
procured from previously published ASC-SSC and SPARK cohorts, and case-control variants
for autism were procured from previously published iPSYCH and Swedish cohorts[16]. Both the
inherited/uninherited and case-control variant sets were filtered to AF < 0.1%.
388

Training, validation, and test datasets

To generate independent training and test sets, we selected 80% (14,894 transcripts) of the
18,629 canonical coding transcripts to comprise the training set and the remaining 20% (3,735
transcripts) to the test set. To ensure the training and test transcripts have similar distributions
of features that may impact constraint estimates, we used stratified randomization to match the
training and test transcripts on s_{het} coefficients (as a measure of selection) and number of
potential missense sites (as a measure of power to detect transcript-wide constraint changes).
The training set was used for MPC model training and MCR model selection, and the test set
was used for MPC model evaluation. No similar hold-outs of data were performed for training of
the mutational model used to compute expected variant counts (see *Modeling of mutation rates*
and expected neutral missense variation).

Modeling of mutation rates and expected neutral missense variation

Expected missense variant counts were determined as described previously[13]. Briefly, we
created a model using the 15,708 gnomAD v2.1.1 genomes that estimated the mutation rate for
each single nucleotide substitution with one base of context (e.g., ACT > AGT) in non-coding
regions of the genome. We then calibrated this mutation rate against the proportion observed of
each context at synonymous sites to adjust for the larger size of the gnomAD v2 exomes,
adjusting for low coverage regions (median coverage < 40x) and methylation levels at CpG sites
using methylation data from the Roadmap Epigenomics Consortium[55]. We created three
separate models (referred to as "plateau" models moving forwards): one for autosomal and
pseudoautosomal sites, one for chromosome X sites, and one for chromosome Y sites. Each of
these models contains mutation rate estimates for each substitution, context, and methylation
level. We then applied the plateau models to the proportion observed of each substitution and
its context, exome coverage, and methylation level. We counted all possible variants in our
synthetic Hail Table (HT) that passed the following criteria: (1) Median coverage > 0; (2) no low-
quality variant observed in gnomAD v2 exomes; (3) no variants above 0.1% AF observed in
gnomAD v2 exomes. We then correlated this proportion observed value with the mutation rate
calculated using the appropriate model above. For low coverage sites (median coverage below
40x), we calculated a scaling factor as described previously[13]: briefly, we computed the total
number of observed synonymous variants in the gnomAD v2 exomes divided by the total
number of possible synonymous variants in the synthetic HT multiplied by the mutation rate
aggregated across all possible substitutions and their contexts and methylation levels. We used
this scaling factor to create a model to adjust the proportion of expected variation for low
coverage sites (coverage model).

Identifying breakpoints within transcripts of regional missense constraint

Observed missense variant counts were calculated using sites from the 125,748 gnomAD
exomes that passed all the following criteria: (1) Allele count (AC) > 0; (2) allele frequency (AF)
< 0.001; (3) variant QC PASS (passed gnomAD variant QC filters, including random forest
filters); (4) median coverage > 0. We filtered the gnomAD v2 exomes Hail Table (HT) to the

sites that matched the above criteria and then annotated the synthetic HT with whether that
variant (chromosome/locus plus reference and alternate alleles) was observed in the gnomAD
exomes. We then aggregated the total number of observed variant counts per locus by
summing the number of observed variants for each possible substitution (reference and
alternate allele) at each locus. Finally, we grouped the synthetic HT by transcript annotation to
sum the total number of observed missense variants per transcript.

As previously described[13], we applied the two models (plateau and coverage) described in
Modeling of mutation rates and expected neutral missense variation to calculate the total
proportion of expected missense variation. Briefly, we summed the mutation rate (mu_agg) for
each substitution, context, and methylation level across the exome. We then applied the
appropriate plateau model (autosomal/pseudoautosomal, chromosome X, chromosome Y) and
adjusted CpG vs. non-CpG sites separately. After applying the appropriate plateau model, we
applied the coverage model to low coverage (median coverage < 40x) sites to create the final
adjusted mutation rate (mu_adj). We then aggregated the raw mutation rate sum (mu_agg) and
adjusted mutation rate (mu_adj) per transcript to get the total mutation rate sum and proportion
of expected missense variation per transcript.

We implemented a minimum number of expected missense variants to prevent finding
breakpoint positions that would create very small (i.e., a handful of base pairs in size) transcript
subsections (see **Supplementary Note**).

We applied a likelihood ratio test to determine whether the missense observed/expected (OE)
ratio was uniform along a transcript or whether a transcript had evidence of distinct sections of
missense constraint. We used the observed and expected missense counts to search for
positions that would divide a transcript into two or more regions with varying levels of missense
depletion. For our analyses, we assume that the observed missense counts should follow a
Poisson distribution around the expected missense counts. We defined our null model as
transcripts not having any evidence of regional variability in missense depletion (where the
expectation, the OE ratio, is consistent across the length of the transcript). Our alternative
model was that transcripts exhibited evidence of distinct sections of missense depletion (OE
ratio calculated per transcript subsection). Because the alternative model should always have a
better fit than the null model, we require a chi square value above a given threshold (p = 0.001)
to establish significance. We used the following formulas to determine the significance of a
breakpoint that would split a transcript into two sections, A and B:

- $p_0 = Pois(obs_A, exp_A * OE) * Pois(obs_B, exp_B * OE)$
- $p_1 = Pois(obs_A, exp_A * OE_A) * Pois(obs_B, exp_B * OE_B)$
- $\chi^2 = 2(ln(p_1) - ln(p_0))$

where OE is the missense observed/expected ratio across the entire transcript, obs_A is the
number of observed missense variants in transcript section A, exp_A is the number of expected
missense variants in transcript section A, OE_A is the OE ratio across transcript section A, obs_B is
the number of observed missense variants in transcript section B, exp_B is the number of

expected missense variants in transcript section A, OE_B is the OE ratio across transcript section
B, and Pois is the Poisson likelihood.
476
477 We used the following formulas to determine the significance of a breakpoint that would split a
478 transcript into three sections, A, B, and C:
479

480 - $p_0 = Pois(obs_A, exp_A * OE) * Pois(obs_B, exp_B * OE) * Pois(obs_C, exp_C * OE)$

481 - $p_1 = Pois(obs_A, exp_A * OE_A) * Pois(obs_B, exp_B * OE_B) * Pois(obs_C, exp_C * OE_C)$

482 - $\chi^2 = 2(ln(p_1) - ln(p_0))$

483
where OE is the missense observed/expected ratio across the entire transcript, obs_A is the
number of observed missense variants in transcript section A, exp_A is the number of expected
missense variants in transcript section A, OE_A is the OE ratio across transcript section A, obs_B is
the number of observed missense variants in transcript section B, exp_B is the number of
expected missense variants in transcript section A, OE_B is the OE ratio across transcript section
B, obs_C is the number of observed missense variants in transcript section C, exp_C is the number
of expected missense variants in section C, and Pois is the Poisson likelihood.
491
492 For the purposes of our analyses, all transcript subsections with more observed variants than
493 expected were capped at an OE of 1, as we were looking for areas of missense depletion and
494 not missense enrichment. We also converted the expected counts for transcript subsections
495 with zero expected variants from 0 to 10^{-9} to prevent nonfinite OE values.
496
To search for a single breakpoint that would divide a transcript into two subsections, we
calculated chi square statistics (as discussed above) to conduct likelihood ratio tests
simultaneously for every eligible position within a transcript. The positions we considered were
positions with a possible missense variant substitution that had at least 16 expected missense
counts in either direction (i.e., both transcript subsections created by dividing the transcript at
this point would have at least 16 expected missense variants). We then aggregated chi square
values across each transcript to find the maximum value per transcript, and we marked any
positions as breakpoints if the chi square calculated at that position was equal to the maximum
chi square value over all sites in the transcript and significant at p = 0.001.
506
Any transcripts that did not have a single significant breakpoint moved forwards into our two
simultaneous breaks search flow. In this search flow, we again calculated chi square statistics to
conduct likelihood ratio tests for every eligible position pair. For every position with a possible
missense, we calculated the chi square statistic of that position paired with each possible
position downstream as long as the two positions created transcript subsections with at least 16
expected missense variants (i.e., all three of the transcript subsections created would have at
least 16 expected missense variants). Because of the large number of pairwise computations,
this step is the most computationally intensive portion of our algorithm. After completing the

single and two simultaneous break search workflows, we merged the results from both search
types.

Our breakpoint search flow is recursive, and the steps are as follows: Search for a single
significant breakpoint dividing a transcript into two subsections. If no single significant
breakpoint was found in the transcript, search for two simultaneous breakpoints. Merge the
results from the single and two simultaneous breakpoint searches. Repeat the steps above,
treating each separate transcript subsection as if it were an independent transcript, until no
more significant breakpoints are found.

Modeling deleteriousness of missense substitution classes with missense constraint

We incorporated two MCR OE-based metrics to measure the increased deleteriousness of
amino acid substitution classes (e.g., Met to Tyr) in functionally important areas of proteins: the
overall OE for each substitution and the second derivative of this OE value per OE bin of
missense constraint (**Supplementary Fig. 14**). To calculate the first metric, the substitution
overall OE, we divided the total number of rare, high quality variants (see *gnomAD variants*)
causing that substitution by the total number of expected variants (see *Modeling of mutation*
rates and expected neutral missense variation). To calculate the second metric, the substitution
OE second derivative, we aggregated the OEs of each substitution by MCR OE bin in 10 bins
from 0 to 1.0+ (i.e., for the 0-0.1 OE bin, we calculated all of the observed substitutions that
occurred within regions with a OE between 0 and 0.1 and divided that number by the total
number of expected substitutions occurring in those regions).

Modeling deleteriousness of individual missense variants

We designed a missense variant deleteriousness predictor to explicitly incorporate information
on amino acid substitution class and position-specific variant effects. A logistic regression model
was first trained to differentiate pathogenic from benign missense variants. The pathogenic
training set consisted of high-quality ClinVar variants (see *ClinVar variants*) labeled as
pathogenic or likely pathogenic in 2,987 likely-haploinsufficient genes, defined as having
probability of haploinsufficiency (pHaplo) ≥ 0.86[41], or in 366 genes with DD associations in G2P
through non-LoF mechanisms. The latter gene set was created by filtering on the G2P DD panel
(accessed October 6, 2023) to select genes where: 1. confidence_category is either definitive or
strong evidence, 2. allelic_requirement was monoallelic, and 3. mutation_consequence included
altered gene product structure or increased gene product level. The benign training set
consisted of high-quality common variants as described in *gnomAD variants*. Variants matching
criteria for both the benign and pathogenic sets were removed from the training data. We
evaluated models with all possible combinations of the following complementary features: amino
acid substitution overall OE and OE second derivative (see *Modeling deleteriousness of*
missense substitution classes); BLOSUM[38] and Grantham[39] scores of amino acid substitution
class severity; the local missense constraint level of a variant (missense OE across the MCR if
applicable, else across the transcript); and PolyPhen-2[40]. We selected BLOSUM, Grantham,

and PolyPhen-2 because of the orthogonal information added on top of our OE-based (and
therefore population allele frequency-dependent) metrics. For each model, only variants with all
relevant annotations were used in training the regression model and the subsequent
calculations to produce deleteriousness scores. The deleteriousness score prediction for any
missense variant i is given as:

$$d_i = -log_{10}(m_i/M)$$
$$m_i = max(0.83, f_i)$$

where d_i is the deleteriousness score prediction, f_i is the number of common missense variants
with a fitted value from the regression that is less than the fitted value for variant i, and M is the
number of common missense variants (equivalent to the number of benign training variants for
the regression). m_i is set to have a minimum value of 0.83 to avoid a mathematical error in the
log when the fitted value for a given variant is less than those of all common variants. Larger
values of d_i indicate stronger predicted-deleteriousness. The best model was chosen to be the
model featurized with all six possible features. The training set for this model consisted of
64,023 benign variants and 12,955 pathogenic variants. This model was then applied to
produce MPC scores for the 68,576,965 possible exome-wide missense variants with all
features, and the distribution of these MPC scores is given in **Supplementary Figs. 10, 11,** and
12.

Comparison of MPC to other predictors

We compared our model to the following missense deleteriousness predictors:
AlphaMissense[42], Constrained Coding Regions (CCRs)[9], MVP[46], M-CAP[43], PrimateAI-3D[45],
REVEL[44], CADD[47,48], PolyPhen-2[40], and SIFT[49]. We annotated the case and control *de novo*
missense variants described in *Rare and de novo variants from developmental cohorts* and
ranked the variants based on their annotated scores. To assess each predictor's ability to
stratify case and control variation, we assessed the proportion of case to control variants among
the variants with the top 10% for each score and compared this number to the overall proportion
of case to control variation using a Fisher exact test.

Data availability

The missense constraint regions (MCRs) are displayed on the gnomAD v2 browser
(https://gnomad.broadinstitute.org) and available for download on the gnomAD website
(https://gnomad.broadinstitute.org/downloads#v2) and in the gnomAD v2 public datasets on
Google, Amazon, and Microsoft clouds. MPC scores for all possible variants in canonical
transcripts is available in the gnomAD v2 public datasets on Google (gs://gcp-public-data--
gnomad/release/2.1.1/regional_missense_constraint/gnomad_v2.1.1_mpc.ht). gnomAD v2
exome, genome, and coverage data and the table of all possible single nucleotide
polymorphisms used to calculate mutational models and search for MCRs are also available in
the gnomAD public buckets and are easily accessed using code in the gnomAD Hail utilities
GitHub repository
(https://github.com/broadinstitute/gnomad_methods/blob/7c0c994883f321492a48962674d5cae

b289df4c7/gnomad/resources/grch37/gnomad.py#L107 and
https://github.com/broadinstitute/gnomad_methods/blob/7c0c994883f321492a48962674d5caeb
289df4c7/gnomad/utils/vep.py#L161).

ClinVar data were downloaded from ClinVar's FTP server
(https://ftp.ncbi.nlm.nih.gov/pub/clinvar/). *De novo* variants were extracted from the
supplemental files of the cited studies.

AlphaMissense scores were downloaded from https://github.com/google-
deepmind/alphamissense. CCRs were downloaded from https://github.com/quinlan-lab/ccrhtml.
MCAP scores were downloaded from http://bejerano.stanford.edu/MCAP/. REVEL scores were
downloaded from https://sites.google.com/site/revelgenomics/. PrimateAI-3D scores were
downloaded from https://primad.basespace.illumina.com/download. MVP scores were
downloaded from
https://figshare.com/articles/dataset/Predicting_pathogenicity_of_missense_variants_by_deep_l
earning/13204118. PolyPhen-2 and SIFT scores were obtained from VEP[56]. CADD scores were
downloaded from the CADD website (https://cadd.gs.washington.edu/download). phyloP scores
were downloaded from the UCSC browser (https://genome.ucsc.edu/cgi-
bin/hgTrackUi?db=hg38&g=cons241way).

Code availability

Code to determine missense constraint regions (MCRs) and calculate MPC is available at
https://github.com/broadinstitute/regional_missense_constraint. Code used to generate the
mutational models is available at https://github.com/broadinstitute/gnomad_lof and
https://github.com/broadinstitute/gnomad-constraint. The Hail library is available at
https://hail.is/.

Acknowledgments

We thank all of the individuals who contributed their data to gnomAD for enabling the research
presented here and in numerous other publications. We also thank members of the gnomAD
Production and Browser Teams for their behind-the-scenes efforts in producing and sharing the
gnomAD dataset, Dan King and the rest of the Hail team for enabling data processing at scale,
and Emma Pierce-Hoffman for her work curating gene lists. This work was supported by the
National Human Genome Research Institute (NHGRI; U24HG011450 to H.L.R. and M.J.D.;
U01HG011755 to A.O.D.L., H.L.R, and M.E.T.; U01HG012022 to P.R., 5T32HG002295-18 to
L.W.), the Eunice Kennedy Shriver National Institute of Child Health and Human Development
(NICHD; F31HD111109 to L.W.), the National Science Foundation (NSF; GRFP#2022339661 to
L.W.) the Simons Foundation (SFARI 1009802 to M.E.T. and K.E.S.), a National Health and
Medical Research Council investigator grant (2009982) to D.G.M., and the Massachusetts
General Hospital Executive Committee on Research, Interim Support Funding (K.E.S.).

Author information

These authors contributed equally: Katherine R. Chao and Lily Wang.

Contributions

K.R.C., L.W., K.E.S., B.M.N, and M.J.D conceived and designed experiments. K.R.C, L.W., and K.E.S. performed primary writing of the manuscript. K.R.C., L.W., R.P., C.L., H.A., and R.Y. performed the analyses and generated figures. P.S. and J.C. were instrumental to developing methods. R.H.G., N.A.W., and M.S. developed visualizations for the web browser. B.W., W.P., M.W.W., K.M.L., J.K.G, K.J.K, and G.T. completed code review for methods. D.G., J.I.G., C.V., and T.P. helped debug runtime compute. J.A.K. provided data and analysis suggestions. S.B. contributed analysis suggestions. H.L.R., B.M.N., M.E.T, D.G.M, A.O.D.L., K.J.K., P.R., M.J.D., and K.E.S. supervised the research. All authors listed under the Genome Aggregation Database Consortium contributed to the generation of the primary data incorporated into the gnomAD resource. All authors reviewed the manuscript.

Ethics declarations

Competing interests

J.A.K is a current employee of Regeneron Genetics Center. T.P. and G.T. are founders of E9 Genomics, Inc.. H.L.R. has received support from Illumina and Microsoft to support rare disease gene discovery and diagnosis. B.M.N. is a member of the scientific advisory board at Deep Genomics and Neumora. M.E.T. has received research support and/or reagents from Illumina, Pacific Biosciences, Microsoft, Oxford Nanopore, and Ionis Therapeutics. D.G.M. is a paid advisor to GSK, Insitro, and Overtone Therapeutics, and receives research funding from Microsoft. A.O.D.L. has consulted for Tome Biosciences and Ono Pharma USA Inc, and is member of the scientific advisory board for Congenica Inc and the Simons Foundation SPARK for Autism study, and received research support from Pacific Biosciences for rare disease diagnosis. K.J.K. is a consultant for Vor Biopharma, Tome Biosciences, and is on the Scientific Advisory Board of Nurture Genomics. M.J.D. is a founder of Maze Therapeutics and Neumora Therapeutics, Inc. (f/k/a RBNC Therapeutics). K.E.S. has received support from Microsoft for work related to rare disease diagnostics. All other authors declare no competing interests.

References

1. Samocha, K. E. *et al.* Regional missense constraint improves variant deleteriousness prediction. *bioRxiv* 148353 (2017) doi:10.1101/148353.
2. Gussow, A. B., Petrovski, S., Wang, Q., Allen, A. S. & Goldstein, D. B. The intolerance to functional genetic variation of protein domains predicts the localization of pathogenic mutations within genes. *Genome Biol.* **17**, 9 (2016).
3. Wiel, L., Venselaar, H., Veltman, J. A., Vriend, G. & Gilissen, C. Aggregation of population-based genetic variation over protein domain homologues and its potential use in genetic diagnostics. *Hum. Mutat.* **38**, 1454–1463 (2017).
4. Sivley, R. M., Dou, X., Meiler, J., Bush, W. S. & Capra, J. A. Comprehensive Analysis of Constraint on the Spatial Distribution of Missense Variants in Human Protein Structures. *Am. J. Hum. Genet.* **102**, 415–426 (2018).
5. Lal, D. *et al.* Gene family information facilitates variant interpretation and identification of disease-associated genes in neurodevelopmental disorders. *Genome Med.* **12**, 28 (2020).
6. Zhang, X. *et al.* Genetic constraint at single amino acid resolution improves missense variant prioritisation and gene discovery. (2022) doi:10.1101/2022.02.16.22271023.
7. Traynelis, J. *et al.* Optimizing genomic medicine in epilepsy through a gene-customized approach to missense variant interpretation. *Genome Res.* **27**, 1715–1729 (2017).
8. Perszyk, R. E., Kristensen, A. S., Lyuboslavsky, P. & Traynelis, S. F. Three-dimensional missense tolerance ratio analysis. *Genome Res.* **31**, 1447–1461 (2021).
9. Havrilla, J. M., Pedersen, B. S., Layer, R. M. & Quinlan, A. R. A map of constrained coding regions in the human genome. *Nat. Genet.* **51**, 88–95 (2019).
10. Silk, M. *et al.* MTR3D: identifying regions within protein tertiary structures under purifying selection. *Nucleic Acids Res.* **49**, W438–W445 (2021).
11. Hicks, M., Bartha, I., di Iulio, J., Venter, J. C. & Telenti, A. Functional characterization of 3D protein structures informed by human genetic diversity. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 8960–8965 (2019).
12. Li, B., Roden, D. M. & Capra, J. A. The 3D mutational constraint on amino acid sites in the human proteome. *Nat. Commun.* **13**, 3273 (2022).
13. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).
14. Landrum, M. J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res.* **48**, (2020).
15. Kaplanis, J. *et al.* Evidence for 28 genetic disorders discovered by combining healthcare and research data. *Nature* **586**, 757–762 (2020).
16. Fu, J. M. *et al.* Rare coding variation provides insight into the genetic architecture and phenotypic context of autism. *Nat. Genet.* **54**, 1320–1331 (2022).
17. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).
18. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**,

285–291 (2016).

19. Taliun, D. *et al.* Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature* **590**, 290–299 (2021).

20. Backman, J. D. *et al.* Exome sequencing and analysis of 454,787 UK Biobank participants. *Nature* **599**, 628–634 (2021).

21. All of Us Research Program Genomics Investigators. Genomic data in the All of Us Research Program. *Nature* (2024) doi:10.1038/s41586-023-06957-x.

22. Petrovski, S. *et al.* The Intolerance of Regulatory Sequence to Genetic Variation Predicts Gene Dosage Sensitivity. *PLoS Genet.* **11**, e1005492 (2015).

23. Samocha, K. E. *et al.* A framework for the interpretation of de novo mutation in human disease. *Nat. Genet.* **46**, 944–950 (2014).

24. Weghorn, D. *et al.* Applicability of the Mutation-Selection Balance Model to Population Genetics of Heterozygous Protein-Truncating Variants in Humans. *Mol. Biol. Evol.* **36**, 1701–1710 (2019).

25. Agarwal, I., Fuller, Z. L., Myers, S. R. & Przeworski, M. Relating pathogenic loss-of-function mutations in humans to their evolutionary fitness costs. *Elife* **12**, (2023).

26. Kosmicki, J. A. *et al.* Refining the role of de novo protein-truncating variants in neurodevelopmental disorders by using population reference samples. *Nat. Genet.* **49**, 504–510 (2017).

27. Bamshad, M. J., Nickerson, D. A. & Chong, J. X. Mendelian Gene Discovery: Fast and Furious with No End in Sight. *Am. J. Hum. Genet.* **105**, 448–455 (2019).

28. Seaby, E. G., Rehm, H. L. & O'Donnell-Luria, A. Strategies to Uplift Novel Mendelian Gene Discovery for Improved Clinical Outcomes. *Front. Genet.* **12**, 674295 (2021).

29. Kapplinger, J. D. *et al.* Enhancing the Predictive Power of Mutations in the C-Terminus of the KCNQ1-Encoded Kv7.1 Voltage-Gated Potassium Channel. *J. Cardiovasc. Transl. Res.* **8**, 187–197 (2015).

30. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet. Med.* **17**, 405–424 (2015).

31. Whiffin, N. *et al.* CardioClassifier: disease- and gene-specific computational decision support for clinical genome interpretation. *Genet. Med.* **20**, 1246–1254 (2018).

32. Küry, S. *et al.* Rare germline heterozygous missense variants in BRCA1-associated protein 1, BAP1, cause a syndromic neurodevelopmental disorder. *Am. J. Hum. Genet.* **109**, 361–372 (2022).

33. Hamosh, A., Amberger, J. S., Bocchini, C., Scott, A. F. & Rasmussen, S. A. Online Mendelian Inheritance in Man (OMIM®): Victor McKusick's magnum opus. *Am. J. Med. Genet. A* **185**, 3259–3265 (2021).

34. Thormann, A. *et al.* Flexible and scalable diagnostic filtering of genomic variants using G2P with Ensembl VEP. *Nat. Commun.* **10**, 2373 (2019).

35. Christmas, M. J. *et al.* Evolutionary constraint and innovation across hundreds of placental mammals. *Science* **380**, eabn3943 (2023).

36. Landrum, M. J. *et al.* ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res.* **46**, D1062–D1067 (2018).

37. Pejaver, V. *et al.* Calibration of computational tools for missense variant pathogenicity

classification and ClinGen recommendations for PP3/BP4 criteria. *Am. J. Hum. Genet.* **109**, 2163–2177 (2022).

38. Henikoff, S. & Henikoff, J. G. Amino acid substitution matrices from protein blocks. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 10915–10919 (1992).

39. Grantham, R. Amino acid difference formula to help explain protein evolution. *Science* **185**, 862–864 (1974).

40. Adzhubei, I. A. *et al.* A method and server for predicting damaging missense mutations. *Nat. Methods* **7**, 248–249 (2010).

41. Collins, R. L. *et al.* A cross-disorder dosage sensitivity map of the human genome. *Cell* **185**, 3041–3055.e25 (2022).

42. Cheng, J. *et al.* Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science* **381**, eadg7492 (2023).

43. Jagadeesh, K. A. *et al.* M-CAP eliminates a majority of variants of uncertain significance in clinical exomes at high sensitivity. *Nat. Genet.* **48**, 1581–1586 (2016).

44. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877–885 (2016).

45. Gao, H. *et al.* The landscape of tolerated genetic variation in humans and primates. *Science* **380**, eabn8153 (2023).

46. Qi, H. *et al.* MVP predicts the pathogenicity of missense variants by deep learning. *Nat. Commun.* **12**, 510 (2021).

47. Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human genetic variants. *Nat. Genet.* **46**, 310–315 (2014).

48. Rentzsch, P., Schubach, M., Shendure, J. & Kircher, M. CADD-Splice-improving genome-wide variant effect prediction using deep learning-derived splice scores. *Genome Med.* **13**, 31 (2021).

49. Ng, P. C. & Henikoff, S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res.* **31**, 3812–3814 (2003).

50. Fuller, Z. L., Berg, J. J., Mostafavi, H., Sella, G. & Przeworski, M. Measuring intolerance to mutation in human genetics. *Nat. Genet.* **51**, 772–776 (2019).

51. Jin, S. C. *et al.* Contribution of rare inherited and de novo variants in 2,871 congenital heart disease probands. *Nat. Genet.* **49**, 1593–1601 (2017).

52. Bishop, M. R. *et al.* Genome-wide Enrichment of De Novo Coding Mutations in Orofacial Cleft Trios. *Am. J. Hum. Genet.* **107**, 124–136 (2020).

53. Qiao, L. *et al.* Rare and de novo variants in 827 congenital diaphragmatic hernia probands implicate LONP1 as candidate risk gene. *Am. J. Hum. Genet.* **108**, 1964–1980 (2021).

54. EuroEPINOMICS-RES Consortium, Epilepsy Phenome/Genome Project & Epi4K Consortium. De novo mutations in synaptic transmission genes including DNM1 cause epileptic encephalopathies. *Am. J. Hum. Genet.* **95**, 360–370 (2014).

55. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–330 (2015).

56. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).