The landscape of regional missense mutational intolerance quantified from 730,947 exomes

Lily Wang[1,2,3]*, Katherine R. Chao[1,2]*, Ruchit Panchal[1,2], Calwing Liao[4,5,6], Haneen Abderrazzaq[7], Robert Ye[4,5], Patrick Schultz[1,4,5], John Compitello[1,4,5], Riley H. Grant[1], Jack A. Kosmicki[3,4,5], Ben Weisburd[1,2], William Phu[1,2], Michael W. Wilson[1,2], Kristen M. Laricchia[1,2], Julia K. Goodrich[1,2], Daniel Goldstein[1,4,5], Jacqueline I. Goldstein[1,4,5], Christopher Vittal[1,4,5], Timothy Poterba[1,4,5], Samantha Baxter[1], Nicholas A. Watts[1,4], Matthew Solomonson[1,4], Genome Aggregation Database Consortium, Grace Tiao[1,2], Heidi L. Rehm[1,2], Benjamin M. Neale[1,4], Michael E. Talkowski[1,2,5], Daniel G. MacArthur[1,8,9], Anne O'Donnell-Luria[1,2,10], Konrad J. Karczewski[1,4,5], Predrag Radivojac[7], Mark J. Daly[1,4,11], Kaitlin E. Samocha[1,2]

[1]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[2]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
[3]Bioinformatics and Integrative Genomics Program, Harvard Medical School, Boston, MA, USA
[4]Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
[5]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[6]Department of Medicine, Harvard Medical School, Boston, MA, USA
[7]Khoury College of Computer Sciences, Northeastern University, Boston, MA, USA
[8]Centre for Population Genomics, Garvan Institute of Medical Research and UNSW Sydney, Sydney, New South Wales, Australia
[9]Centre for Population Genomics, Murdoch Children's Research Institute, Melbourne, Victoria, Australia
[10]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA
[11]Institute for Molecular Medicine Finland (FIMM), Helsinki, Finland

*Indicates equal contribution

Correspondence should be addressed to K.E.S. (samocha@broadinstitute.org)

Abstract

Missense variants can have a range of functional impacts depending on factors such as the

specific amino acid substitution and location within the gene. To interpret their deleteriousness,

studies have sought to identify regions within genes that are specifically intolerant of missense

variation. Here, we leverage the patterns of rare missense variation in 730,947 exome

sequenced individuals in the Genome Aggregation Database (gnomAD v4.1.1) against a null

mutational model to identify transcripts with regional differences in missense constraint.

Missense-depleted regions are enriched for ClinVar pathogenic variants, *de novo* missense

variants from individuals with neurodevelopmental disorders, and complex trait heritability.

Following ClinGen calibration recommendations for the ACMG/AMP variant classification

guidelines, we establish that variants within regions with <36% of their expected missense

variation achieve moderate support for pathogenicity. We integrate this regional constraint

measure into a missense deleteriousness metric (named MPC) that effectively stratifies rare

and *de novo* missense variants in individuals with early-onset developmental conditions from

controls. These results provide additional tools to aid in missense variant interpretation.

Main text

Over the last decade, exome and genome sequencing have enabled variant discovery across hundreds of thousands of individuals[1–6]. As sequencing becomes more routine in both clinical and research settings, it is crucial to improve our ability to interpret the potential impacts of these variants, particularly in understanding disease risk. With the rapidly expanding catalog of observed genomic variation comes an increasingly large number of variants of uncertain significance (VUS) that cannot be conclusively considered harmful (pathogenic) or neutral (benign)[7]. Indeed, well over half of the variants in the ClinVar database[8], which hosts disease-variant associations, are considered VUS, with thousands more added every year[9].

Classification and interpretation of rare variants has been greatly aided by the release of large reference databases, which have provided the opportunity to study selective forces acting on the human genome and to identify genomic regions under selective constraint by, for example, identifying regions with fewer variants than expected based on mutational models[2,3,10–13]. In particular, gene-level metrics of predicted loss-of-function (pLoF) variant depletion have proven to be valuable in variant classification and identification of novel gene-disease relationships[14–18]. The functional impact and selective pressures relevant to missense variation, by contrast, remain challenging to predict, as the effect of a missense variant is governed by the gene housing the variant, the position of the variant in the gene, and the specific amino acid substitution caused by the variant. In fact, 90% of the missense variants in ClinVar are VUS[9,19]. To address this, prior work has sought to identify regions within coding genes that are specifically intolerant of missense variation as a way to improve classification[20–31].

One major application of constraint metrics is in novel gene identification, particularly when searching for a significant burden of *de novo* (newly arising) variants. Both missense deleteriousness metrics and region-specific missense depletion information have been used to improve power to identify genes associated with neurodevelopmental disorders[15,16,32], as both approaches help separate signal from noise by highlighting a subset of missense variants that are more likely to contribute to rare disorders. Beyond enrichment in statistical frameworks, these metrics can also be incorporated into clinical workflows for variant classification. This requires that the metrics are calibrated for use in the published guidelines for evaluating variant pathogenicity from the American College of Medical Genetics and Genomics and the Association for Molecular Pathology (ACMG/AMP)[33]. These ACMG/AMP guidelines provide recommendations for how various lines of evidence (e.g., computational, functional) can be

applied towards determining if a variant is pathogenic or benign as well as the strength of the

evidence (e.g., strong, moderate). For a given type of evidence, such as *in silico* computational

tools[34,35] or functional data from multiplex assays of variant effects[36–38], probabilistic frameworks

can be used to determine the strength of evidence.

Here, we expand upon previous work[20] and develop a sub-genic measure of missense

intolerance leveraging population-level variation. We show that this measure facilitates variant

classification and risk stratification for association studies with *de novo*, rare, and common

variants. Specifically, we identify that 36% of transcripts have evidence of variable missense

constraint, allowing us to refine the missense constraint scores for 50% of all coding bases to

the sub-genic region-level rather than gene-level scores. In missense-depleted coding regions,

we find strong enrichments of disease-associated variants from ClinVar and *de novo* missense

variants identified in individuals with neurodevelopmental disorders (NDDs). For the first time,

we apply a clinical calibration framework to assign a strength of evidence for regional missense

depletion that can be used in clinical classification workflows, demonstrating moderate support

for pathogenicity. Finally, we combine our regional missense constraint results with variant- and

amino acid-level evidence to create the MPC (Missense deleteriousness Prediction by

Constraint) score that effectively stratifies rare variants identified in individuals with NDDs from

those found in controls[39–41].

Results

Identification of missense constraint regions within transcripts

We explored the patterns of rare missense variant presence or absence in 730,947 exomes in

the Genome Aggregation Database (gnomAD) v4.1.1 to quantify missense depletion at the sub-

genic level. We searched 17,841 MANE Select or canonical protein-coding transcripts that

passed quality control (see **Methods, Supplementary Table 1**) for variability in missense

constraint, quantified as the number of rare (allele frequency [AF] < 0.1%) missense variants

observed in gnomAD divided by the number expected under neutral evolution as estimated from

previously described mutational models[3]. For each transcript, we applied a recursive search

based on likelihood ratio tests over all potential rare missense sites looking for breaks that

divide the transcript coding sequence into sequences with statistically significantly different

missense observed/expected (OE) values (see **Supplementary Note** and **Supplementary Fig.**

1), which we refer to as missense constraint regions (MCRs; **Fig. 1a-b**). Through this approach,

we discover 6,361 transcripts (36%) that harbor regional variability in missense constraint (**Fig.**

1c), i.e., have two or more MCRs (minimum coding length 20bp, median 339bp;

Supplementary Figs. 2-3). We thus refine the resolution of missense constraint for 50% of

coding sites (15.7 coding Mb in the 6,361 transcripts vs. 31.2 coding Mb in all 17,841 transcripts

assessed), discovering widespread signatures of negative and neutral selection obscured when

measuring constraint at transcript-resolution (**Fig. 1d**). For example, relative to the transcript-

resolution missense constraint distribution, a larger proportion of the coding exome lies within

strongly constrained sequences (e.g., 3.3% vs. 0.7% at OE < 0.4 for all genes; see

Supplementary Note for OE threshold selection) and the mode shifts toward an OE indicative

of evolutionary neutrality at approximately 1 (49.8% vs. 46.4% at 0.9 < OE ≤ 1.1). Across coding

sequences, the median missense OE is 0.92 and the top 10% most constrained sites have a

missense OE ≤ 0.63 (**Supplementary Table 2**).

Fig. 1: 36% of protein-coding genes in the human genome harbor regional variation in

population-level missense depletion. **a**, An example transcript that has two missense constraint

regions (MCRs) with significantly different levels of population-wide missense depletion, defined

as the number of missense variants observed in gnomAD at rare frequency (AF < 0.1%) divided
by the number of rare missense variants expected under neutral evolution (observed/expected
or OE). Lower OE values correspond to greater variant depletion and suggest stronger selective
constraint. **b**, Flow chart describing the process of searching for breakpoints that divide a
transcript into multiple MCRs. Searching for breakpoints is recursive and leverages likelihood
ratio tests at a significance threshold of p = 0.001. **c**, The number of MCRs identified within the
6,361 transcripts where we discover regional differences in missense constraint. The other
11,480 transcripts are deemed to have a single MCR (i.e., a constant level of constraint across
their entirety) and are not shown. **d**, The distribution of local missense OE at all coding sites in
MANE Select or canonical transcripts. Local missense OE is defined as the OE calculated over
the whole transcript (for "transcript OE") or over the MCR (for "MCR OE") where the site is
located. Transcript OE and MCR OE are equivalent for transcripts with one MCR. The areas
under each of the two curves sum to 1.

We find that missense constraint is able to identify regions associated with severe, early-onset

disease. One example is in *BAP1*, which plays a key role in chromatin modeling by mediating

histone deubiquitination[42–44]. BAP1 has a conserved N-terminal catalytic domain (UCH) and a

predicted C-terminal UCHL5/UCH37-like domain (ULD) likely involved in complexing with other

proteins and nuclear localization[45–47]. Disease-causing variants in this gene are linked to both

cancer[42,48,49] and a recently discovered rare NDD named Küry-Isidor syndrome[44]. We identify

two highly constrained MCRs in this gene with missense OEs of 0.50 (6th percentile) and 0.57

(8th percentile) at the N- and C-termini, closely overlapping the annotated gene domains.

Despite covering only 43% of the gene (944/2,187 bp), these two constrained MCRs capture all

13 missense variants reported to be causal for Küry-Isidor in ClinVar or the initial publication[44]

(binomial p of enrichment in the constrained MCRs = 1.8×10^{-5}, **Fig. 2a**). The only ClinVar

pathogenic/likely pathogenic (P/LP) variants that do not fall within any missense-constrained

MCRs in *BAP1* (n=2) are associated with cancer predisposition diseases and are likely under

weaker selection than early-onset NDDs. Overall, 26/28 (93%) of all ClinVar P/LP variants in

BAP1 reside in constrained MCRs (binomial p = 4.2×10^{-8}).

We also find that missense constraint can identify disease-associated gene regions beyond

annotated protein domains. For example, *TARDBP*, which encodes a regulator of RNA

processing and metabolism, is an extremely pLoF-constrained gene[50] and is classically

associated with amyotrophic lateral sclerosis (ALS) and accompanying frontotemporal dementia

(FTD)[51–54]. All 23 ClinVar P/LP variants in this gene cause a missense change, and all but one

reside in the protein C-terminus, which is predicted to be disordered by AlphaFold[55] and

MobiDB-lite in UniProt[45,56] (**Fig. 2b**). Our missense constraint metric assigns strong constraint

over the entirety of this gene and identifies a highly constrained C-terminal MCR (OE = 0.56, 8th

percentile; amino acids [AAs] 265-414) that aligns precisely with the predicted-disordered C-

terminal region of the gene (AAs 259-414 with AlphaFold predicted Local Distance Difference

Test [pLDDT] < 50) and includes 21 of the 22 ClinVar variants at this end of the gene. Notably,

this particular MCR has the weakest constraint of all five MCRs identified in *TARDBP*, but would

be missed by most 3D structural approaches that remove regions with low pLDDT[30]. Taken

together with the extreme pLoF constraint in this gene yet lack of disease-associated pLoF

variants, this suggests two things. First, it is likely the currently known neurodegenerative

conditions are some of the least severe diseases caused by disruptions of *TARDBP*, and LoF or

other damaging variants outside of the disordered C-terminus are likely to cause unrecognized

severe rare disorders or be incompatible with life. Second, missense variants and constraint can

provide a window into understanding gene functions and disease mechanisms when pLoF-

based approaches are more opaque.

Fig. 2: Missense constraint regions (MCRs) and the distribution of ClinVar pathogenic/likely
pathogenic (P/LP) missense variants in two disease-associated genes. Exons are delineated
with black outlines and MCRs are delineated by color based on their missense
observed/expected (OE) ratio. **a,** *BAP1*. Variants in this gene can lead to cancer-predisposition

syndromes, increased risk of certain cancers, or the neurodevelopmental disorder Küry-Isidor

syndrome[44]. All of the variants associated with Küry-Isidor in ClinVar or the initial publication fall

within the highly constrained N-terminal or C-terminal MCRs (6th and 8th percentiles,

respectively). Küry-Isidor variants are colored in black and all other cancer-associated P/LP

missense variants are colored in gray. UCH: Ubiquitin C-terminal hydrolase domain, ULD:

UCHL5/UCH37-like domain. **b,** *TARDBP*. Missense variants in this gene can cause adult-onset

neurodegenerative disorders. The C-terminal constrained MCR (8th percentile), where the

majority of the ClinVar P/LP variants are located, aligns to a predicted-disordered region of the

protein. pLDDT: predicted Local Distance Difference Test from AlphaFold.

Enrichment of disease-associated variation in missense-depleted regions

Next, we sought to determine if the signatures of selection revealed by MCRs more broadly

recapitulated biological and disease relevance of coding sequences. Here, we filtered to high-

quality, high-coverage sequences (95% of MCRs or 98.3% of coding space; see

Supplementary Note and **Supplementary Fig. 4** for discussion of lower-coverage sequences).

Overall, transcripts that are more intolerant to pLoF variation, as measured by the loss-of-

function observed/expected upper bound fraction (LOEUF) score[3,50], also tend to be more

intolerant to missense variation. This trend is markedly more prominent when measuring

intolerance via the most constrained MCR in a transcript vs. across the whole transcript

(Spearman ρ = 0.58 vs. 0.51; **Supplementary Note** and **Supplementary Fig. 5**). We confirmed

that, as expected, autosomal genes without disease associations in OMIM[57] tended to be less

missense-constrained than genes with autosomal dominant disease inheritance (Welch's t p <

10^{-50} for transcript missense OE and Wilcoxon p < 10^{-50} for minimum MCR missense OE).

However, we found that these non-OMIM genes were also more missense-constrained than

genes with autosomal recessive disease inheritance (Welch's t p = 1.1×10^{-16} for transcript

missense OE and Wilcoxon p = 9.0×10^{-7} for minimum MCR missense OE; **Supplementary Fig.**

6). Taken together with our finding that 62% (1,511/2,430) of genes that are both LOEUF-

constrained and severely missense-depleted (first three deciles of LOEUF and MCR with OE <

0.4) do not have disease associations in OMIM, this highlights a potential search space for

novel gene-disease relationships (**Supplementary Fig. 7**).

We defined a set of 695 strongly pLoF-intolerant genes as those in the first three deciles of

LOEUF or either of two s_{het} measures[58,59] and with monoallelic association with a developmental

phenotype in Gene2Phenotype (G2P)[60]. In these genes, we observed strong transcript-wide

missense depletion that was even stronger for genes with multiple MCRs (**Fig. 3a**; Wilcoxon p <

10^{-50}). As a comparator, we defined 1,580 tolerant genes as those in the last three deciles of

LOEUF and the two s_{het} measures and not associated with a phenotype in OMIM. Intolerant

genes were much more likely to harbor multiple MCRs than tolerant genes, at a 62-fold

increased rate after regressing out transcript coding length (612/695 intolerant genes vs.

103/1,580 tolerant genes with multiple MCRs, logistic regression coefficient $p < 10^{-50}$). Intolerant

genes were also highly enriched for severely depleted regions (2.3-fold more likely to have

minimum MCR OE < 0.4 after regressing out transcript length, logistic regression coefficient $p =$

4.3×10^{-4}), whereas the most constrained MCRs in tolerant genes were overall less depleted and

more evenly distributed across the OE spectrum. We next compared missense constraint to

selection over longer timescales, measured by the phyloP[61] score for conservation across

placental mammals. At the transcript-level, we found that genes with more conserved coding

sequences also tended to be more depleted of human missense variation overall (Spearman ρ

$= 0.50$, $p < 10^{-50}$). However, at the MCR-level, we discovered a substantial number of strongly

constrained MCRs that appear widely unconserved across mammals, suggesting human-

specific negative selection pressures that are obscured at the whole-transcript level

(**Supplementary Fig. 8**, **Supplementary Tables 3-4**).

We next aggregated *de novo* missense variants from 31,058 individuals with a severe

developmental disorder[15] (DD), 15,036 autistic individuals (AUT), and their 5,492 siblings not

diagnosed with a DD[16] (**Fig. 3b**, **Supplementary Fig. 9, Supplementary Tables 5-8**). The

distribution of *de novo* missense variants across the missense OE spectrum in unaffected

siblings largely mirrored the exome-wide missense OE distribution. In contrast, *de novo*

missense variants in autistic individuals are enriched in missense-constrained sequences, with

an even stronger effect observed in individuals with DDs. For example, relative to unaffected

siblings, the rate of *de novo* missense variants in MCRs with OE < 0.6 is 3-fold higher in

individuals with DDs (Poisson two-sided $p < 10^{-50}$) and 1.7-fold higher in autistic individuals

(Poisson two-sided $p < 10^{-50}$; **Supplementary Fig. 9**; see **Supplementary Note** for OE

threshold selection). This is consistent with the expectations that a small subset of *de novo*

missense variants in individuals with developmental phenotypes are causal for those traits and

that variants causal for DD are generally more selectively deleterious than those for autism.

Beyond large-effect rare and *de novo* variation in traits under strong negative selection, we

additionally investigated whether our MCR metric, which was calculated using rare variants,

correlated with functional effects of common variants. Prior work found that pLoF-constrained

genes and their flanking 100kb sequences are enriched for SNP heritability across hundreds of

independent traits in the UK Biobank and other large genome-wide association studies

(GWAS)[3]. We partitioned common (AF > 5%) variant heritability of the same 268 independent

traits across MCRs to investigate relative enrichment. To establish a baseline, we computed the

heritability enrichment relative to the average genome-wide SNP over all coding sequences

comprising MCRs (4.5-fold). The most constrained MCRs have the strongest heritability

enrichment: e.g., the first quintile of MCR missense OE harbors a 20-fold enrichment (**Fig. 3c**,

Supplementary Fig. 10, Supplementary Tables 9-10). Coding SNPs in missense-

unconstrained MCRs (e.g., in the three least constrained quintiles of MCR missense OE) harbor

little to no detectable heritability enrichment. These findings suggest that regions depleted of

rare missense variation can prioritize common coding variants important for complex traits (e.g.,

improve GWAS fine-mapping variant prioritization), and that there exists a subset of coding

sequences with no appreciable heritability enrichment relative to the average genome-wide

SNP, which constraint can help identify.

Fig. 3: Regional missense depletion reveals constraint obscured by gene-level measures. **a**,
Left: The distribution of transcript-wide missense observed/expected (OE) across 17,841
transcripts stratified by the combination of two factors: whether the transcript is strongly
mutationally intolerant or tolerant and whether we detect multiple missense constraint regions
(MCRs). Number of transcripts in each category are: strongly intolerant with multiple MCRs
(n=612; red), strongly intolerant with one MCR (n=83; yellow), tolerant with multiple MCRs
(n=103; dark gray), tolerant with one MCR (n=1,477; light gray). Right: Minimum MCR missense
OE using the same groupings. For transcripts with a single MCR, minimum MCR missense OE
is equivalent to transcript missense OE. **b**, MCR missense OE at all sites of possible exome-
wide missense variants vs. sites of *de novo* missense variants in controls, autistic individuals
(AUT), or individuals with developmental disorders (DDs). *De novo* variants from individuals with
developmental phenotypes are enriched in more constrained sequences, with a more
pronounced enrichment in DD than autism. **c**, Enrichment in per-variant heritability explained by
common (AF > 5%) protein-coding SNPs stratified by MCR missense OE quintile, relative to the
average SNP genome-wide. Enrichment is estimated by linkage disequilibrium score
regression, accounting for number of SNPs in each quintile, and is averaged across 268
independent traits in the UK Biobank and other large genome-wide association studies. Black

259　dashed line at 1 indicates no enrichment. Blue dashed line at 4.5 indicates average coding
260　enrichment. Error bars represent 95% confidence intervals.
261

Evaluating the clinical utility of missense depletion
We then assessed the ability of MCRs to recapitulate clinical disease information by examining
the constraint distribution around rare ClinVar P/LP and benign/likely benign (B/LB) missense
variants. Across genes with autosomal dominant disease inheritance and ≥2 MCRs, we found
that P/LP missense variants tend to occur in regions that are more strongly constrained than the
transcript average (n=529 genes, n=29,928 variants, paired Welch's t p < 10^{-50}), while the
converse was true for B/LB variants (n=615 genes, n=51,705 variants, paired Welch's t p < 10^{-50}
$^{-50}$; **Fig. 4a**, **Supplementary Fig. 11a-b, Supplementary Table 11**). For example, in 337 genes
with both constrained (missense OE < 0.6) and unconstrained (missense OE > 0.9) MCRs,
P/LP variants occur 11-fold more frequently in constrained than unconstrained MCRs, while
B/LB variants occur 9-fold more frequently in unconstrained than constrained MCRs (odds ratio
[OR] = 93, Fisher's exact p < 10^{-50}). These patterns are also significant, though more mild, in
genes associated with autosomal recessive inheritance (**Supplementary Fig. 11c-d,**
Supplementary Table 12). Together, these results demonstrate that our constraint metric is
able to distinguish specific sequences within genes that are enriched for pathogenic or benign
variants.
278

Having confirmed a correlation between constraint and clinical disease information, we then
sought to quantify the utility of our constraint metric as a point of evidence for clinical variant
classification. To do so, we leveraged a probabilistic framework established by ClinGen[34] that
formally identifies score thresholds for computational metrics that match each variant
classification tier (e.g., "strong" or "moderate") under the ACMG/AMP guidelines[33]. We applied
this robust quantitative framework to determine the missense OE thresholds that met different
levels of clinical evidence strengths evaluated under the population data-derived pathogenic
(PP2) and benign (BP1) criteria codes[33]. We did not apply a restriction on gnomAD coverage for
this assessment, as individuals leveraging our constraint metric for clinical interpretation may
not make such considerations, though we do provide coverage information and scores are
coverage-adjusted (see **Supplementary Note**). The resulting thresholds are thus conservative
estimates. We discovered that our constraint metric met moderate (OE ≤ 0.36) and supporting
(OE ≤ 0.59) evidence for pathogenicity, as well as supporting (OE > 0.97) and moderate (OE >
1.23) evidence for benignity (**Fig. 4b, Supplementary Fig. 12**). These correspond respectively

to the 3rd, 9th, 65th, and 99th percentiles of the overall missense OE distribution across coding

sites. Similar assessment of another regional constraint metric, the missense tolerance ratio

(MTR)[62], revealed that MTR also reaches supporting and moderate support for pathogenicity,

but does not meet any level of evidence to support benignity (**Supplementary Fig. 12,**

Supplementary Table 13). These results establish formal thresholds on our missense

constraint metric that may be used within the ACMG/AMP clinical variant classification

guidelines.

Fig. 4: Clinical utility in application of missense constraint. **a,** The distribution of transcript-wide

missense observed/expected (OE; gray) and missense constraint region (MCR) OE for ClinVar

pathogenic/likely pathogenic missense variants (P/LP; red) and benign/likely benign missense

variants (B/LB; blue) in genes with ≥2 MCRs and autosomal dominant disease associations. We

filtered to 501 transcripts with at least one P/LP and one B/LB missense variant. For the P/LP

and B/LB distributions, we annotated each variant with the missense OE across the MCR they

fell in and collapsed these values within each transcript by taking the respective medians.

Asterisks denote $p < 10^{-50}$. **b,** Local posterior probabilities of variant pathogenicity given MCR

missense OE calculated across all transcripts. Gray shading indicates the one-sided 95%

confidence interval on the more stringent side. Horizontal lines indicate thresholds required to

meet ACMG/AMP evidence levels — from bottom to top: supporting, moderate, strong, very

strong. MCR missense OE reaches supporting (OE ≤ 0.59) and moderate (OE ≤ 0.36) level

evidence for the population data-derived pathogenic (PP2) criteria code.

Integrating missense constraint into a variant-level deleteriousness score

We integrated our regional missense constraint measure into a variant-level predictor of

missense deleteriousness named MPC (Missense deleteriousness Prediction by Constraint)

that additionally incorporates information about amino acid substitution type and local context.

The XGBoost-based model integrates regional and gene constraint-derived metrics together

with PolyPhen-2[63] and phyloP[61] and is trained on ClinVar pathogenic variants in 2,987

haploinsufficient genes[64] and 359 genes with non-LoF DD associations in G2P[60] vs. ClinVar

benign variants and gnomAD variants with AF > 0.1% (**Supplementary Tables 14-15,**

Supplementary Fig. 13). Higher scores predict greater deleteriousness (**Supplementary Figs.**

14-15).

We assessed the utility of MPC in prioritizing potentially disease-causing variation by evaluating

its ability to stratify case and control rare and *de novo* missense variation. Consistent with the

regional constraint results, the *de novo* missense variants in DD and AUT are enriched for high

MPC scores compared to the control siblings without DDs (**Supplementary Fig. 16**). We then

calculated burden enrichments in DD and AUT compared to the controls, stratifying by presence

in 373 genes previously associated with NDD[16] and three bins of MPC scores (<2, 2-2.5, ≥2.5;

see **Supplementary Note** for threshold selection; **Fig. 5a-b**, **Supplementary Fig. 17,**

Supplementary Table 16). We found a very strong enrichment of *de novo* variants in NDD-

associated genes for most MPC bins (from most- to least-damaging MPC: rate ratios [RRs] =

5.1, 3.9, 1.2 for AUT; 15.1, 8.0, and 1.4 for DD), consistent with known causal genetic

architectures for these phenotypes. We additionally discovered meaningful enrichments of

predicted-damaging *de novo* variants in genes not yet associated with NDDs. For AUT, this was

only significant in the most severe MPC bin (RR = 1.7, p = 1.3×10^{-5}), suggesting a sizable

reservoir of *de novo* missense variants causal for autism in as-of-yet unassociated genes that

will likely require strict deleteriousness criteria to statistically ascertain. Substantial *de novo*

enrichments were found outside of NDD genes for DD across all missense deleteriousness bins

(RRs = 3.6, 1.4, and 1.1), pointing to considerable latent risk outside of currently associated

genes and even outside of known OMIM genes (RR for MPC ≥2.5 = 2.2, p = 2.0×10^{-9}).

We additionally assessed enrichments of rare missense variants in 5,591 autistic individuals

and 8,597 controls from case-control studies without *de novo* information, as well as parental

transmission for 13,384 autistic individuals[16]. While we did not find meaningful parental

transmission differences for any MPC bin, we did observe substantial enrichment in the case-

control analysis for variants in NDD genes in the two most damaging MPC bins (RRs = 1.4 and

1.2, p = 9.8×10^{-9} and 3.8×10^{-3}, respectively), most likely from *de novo* variants. No significant

case-control enrichment was found in NDD-unassociated genes, pointing to the importance of

de novo information for autism gene discovery.

We extended our assessment of case-control *de novo* stratification for a comparison of our

model against several other *in silico* missense deleteriousness predictors: AlphaMissense[40],

popEVE[41], REVEL[39], Polyphen-2[63], CADD[65,66], phyloP[61], and SIFT[67]. For this assessment, we

evaluated four additional early-onset developmental phenotypes: epileptic encephalopathy

(EE)[68], orofacial cleft (OFC)[69], congenital heart disease (CHD)[70], and congenital diaphragmatic

hernia (CDH)[71]. To compare across predictors with different score distributions, we used a

ranking-based performance assessment. For each predictor, we ranked the *de novo* missense

variants from each case cohort against those in the 5,492 controls without DDs from the autism

study and computed the case-control variant ORs in the top percentiles of these rankings (**Fig.**

5c, Supplementary Table 17). At the top 5% of variants, MPC displays the highest OR for DD

(OR = 9.7, Fisher's exact $p < 10^{-50}$), AUT (OR = 2.4, $p = 1.6 \times 10^{-16}$), CHD (OR = 2.1, $p = 1.4 \times 10^{-8}$

), and CDH (OR = 1.6, $p = 4.8 \times 10^{-3}$), demonstrating that MPC effectively ranks high-impact *de*

novo variants in the most deleterious prediction regimes. Of the other predictors,

AlphaMissense, popEVE, and REVEL tend to perform best. There is generally substantial

overlap in the 95% confidence intervals between the top-performing predictors, indicating that

de novo variant stratification performance tends to be fairly comparable across those models.

The exception is in DD, where MPC has much higher performance than the other predictors.

This is likely driven by the way MPC is designed to predict deleteriousness, which focuses on

strong fitness effects (see **Discussion**). These observations are generally consistent over a

range of thresholds used to define the top percentiles for ranking (**Supplementary Fig. 18**).

Across the phenotype categories assessed, the predictors typically had the highest efficacy in

DD and EE, then AUT and CHD, and finally CDH and OFC, consistent with the expected

contribution of Mendelian *de novo* missense variants to relative risk for each condition. In the

top 1% ranking, CDH and OFC missense enrichments reach substantial suggestive

enrichments, though not Bonferroni significant (CDH OR = 2.8 and $p = 3.8 \times 10^{-3}$, OFC OR = 3.7

and $p = 4.7 \times 10^{-4}$), indicating that while *de novo* missense variants contribute to risk for these

disorders, they do so at a lower rate than for more severe phenotypes.

Fig. 5: MPC effectively stratifies case and control variation. **a**, The difference relative to controls

of missense variants stratified by MPC score and localization to genes associated with

neurodevelopmental disorders (NDDs) for **a,** individuals with a developmental disorder (DD) and

b, autistic individuals (AUT). Relative difference is calculated as: for *de novo* variants, the

average rate of variants in probands divided by that in sibling controls; for case-control, the

average rate of variants in cases divided by that in controls from case-control data; for inherited,

the average rate in probands of transmitted variants divided by that of untransmitted variants.

Error bars represent 95% confidence intervals calculated from a binomial test. **c**, The odds ratio

of case to control *de novo* missense variants in the top 5% vs. bottom 95% of respective

rankings. *De novo* missense variants from each case cohort are ranked against those in the

5,492 controls for each predictor. EE: epileptic encephalopathy, OFC: orofacial cleft, CHD:

congenital heart disease, CDH: congenital diaphragmatic hernia. Error bars represent 95%

confidence intervals. Only variants scored by all predictors are included. In all plots, points are

solid colored if the difference from 1 is statistically significant (binomial or Fisher's exact p <

0.05).

Discussion

We developed a method to identify sub-genic regions with differential intolerance to missense

variation. We demonstrate that coding regions depleted for missense variation in the general

population are enriched for established disease-associated variation, *de novo* variants from

individuals with developmental disorders, and heritability for 268 complex traits from the UK

Biobank and other large GWAS. Additionally, we have calibrated these constraint scores to

establish that regions with less than 36% of their expected missense variation can achieve

moderate evidence towards clinical classification of pathogenicity for missense variants that

occur in these regions, following ACMG/AMP guidelines. Finally, we incorporated regional

missense intolerance information into the missense deleteriousness metric, MPC, and show that

MPC effectively separates potentially risk-carrying variants identified in various developmental

disorder cases from those seen in controls.

Our method of summarizing constraint strength information over contiguous base pair sites into

discretized regions allows for identification of gene segments under distinct selection pressures

that can point to different functions and structures within a gene. This approach can provide

more easily interpretable information about broad gene functions than very fine resolution

metrics at the site- or amino acid-level. For example, we show that MCRs, which do not

incorporate any protein structural information, can align precisely with the locations of protein

domains and functional intrinsically disordered regions as well as the pathogenic disease

variants therein.

Other metrics that have been developed to identify gene segments with distinct missense

intolerance linearly along the genome include subRVIS[21], Constrained Coding Regions

(CCRs)[27], and the constrained segments calculated with MTR[62], among others. SubRVIS is

limited to quantification at the level of exons or domains, lacking dynamic ascertainment of

functional gene regions. The CCR metric is generally more akin to a site-level metric: 60% of

CCR gene segments are 1bp in size due to being the site of a nonsynonymous variant observed

in gnomAD (assigned a score of 0 or unconstrained), and 0.1% of segments are >50bp.

Notably, both subRVIS and CCRs were calculated on datasets much smaller than the dataset

underlying the MCRs presented here (gnomAD v4.1.1): subRVIS used the Exome Sequencing

Project[72] (~112x smaller than gnomAD v4.1.1) and CCRs used gnomAD v2.1.1 (~6x smaller

than v4.1.1). We previously provided a comparison of an earlier version of MCRs calculated

with gnomAD v2.1.1[73] to CCRs. MTR was calculated on the Regeneron Genetics Center Million

Exome (RGC-ME) dataset consisting of 821,979 unrelated samples, which is a comparably

sized dataset to gnomAD v4.1.1 and should therefore afford similar statistical power to our MCR

method. The RGC-ME study released MTR intolerance scores at the amino acid level as well as

at the gene segment level, the latter calculated by aggregating over the former. We find that the

MTR method of segmenting genes into regions is prone to identifying likely spurious

"unconstrained" regions as there are no restrictions placed on the size of unconstrained regions,

only constrained regions. This results in approximately 16% of unconstrained regions falling

under the 10 amino acid minimum applied to constrained regions. Furthermore, the binary

nature of the assignment of amino acids as "constrained" or "unconstrained" for input to

segmentation means that there is no distinction between regions of varying levels of constraint

below the 15th percentile threshold designated "constrained". As a result, consecutive regions

that fall under the "constrained" threshold but have different levels of constraint will be unlikely

to be distinguished from each other by the MTR method unless an "unconstrained region" is

identified that separates them. In contrast, our MCR method delineates gene segments based

on having differing levels of constraint without this binarization, and the size minimum of 16

expected variants is applied to all MCRs for robustness.

We note that constraint, as quantified here by population depletion of observed rare variants, is

essentially a footprint of selection on heterozygous variants. Our methodology is thus most

powerful in sequences undergoing appreciable heterozygous selection, e.g., genes where

severe and early onset diseases are caused by dominant missense variants. However, as

demonstrated in **Supplementary Fig. 11**, disease variants in genes currently known to be only

associated with conditions with recessive inheritance do also associate with detectable

constraint, albeit to a lesser degree than dominant variants. This lends further support to a body

of evidence that few genes are purely recessive: many "recessive-only" genes are likely only

partially recessive, with heterozygotes having measurable phenotypic differences[74–76]. As such,

we show that this constraint method is indeed amenable, to an extent, to characterizing

selective effects even in genes considered to be solely associated with recessive inheritance.

An inherent strength of our regional missense constraint metric is its grounding in population

genetics and independence of, and thus lack of bias from, disease genetics databases and

domain or structural knowledge. Given that constraint measures fitness effects, we expect—and

indeed observe—associations between strong constraint and risk variants in early

developmental disorders which are under strong selective pressure (**Fig. 2a**, **Fig. 3b**), while

genes where germline variants cause adult-onset diseases like hereditary cancers generally

harbor little detectable constraint on missense or pLoF variants. This makes it all the more

notable when we clearly identify highly constrained gene sequences where clinically relevant

missense variants are currently only known to be associated with adult-onset disease, such as

in *TARDBP* (**Fig. 2b**), where the highly constrained C-terminal MCR bounds nearly all clinically

reported variants causing ALS and FTD, diseases that typically begin in post-reproductive life

but can start in earlier adulthood[77–80]. The rest of the gene is even more strongly constrained,

which points to severe fitness effects unexplainable by the known gene-disease associations,

demonstrating the value of this constraint method in revealing novel evidence for gene function.

Additionally, while missense predictive models that rely on 3D structure exclude or have lower

performance in intrinsically disordered regions[81–83], there is no such structure-based

incompatibility for the genetics-derived inference of MCRs. Returning to *TARDBP* as an

example, nearly all of the 23 P/LP missense variants in ClinVar lie in the disordered regions of

this gene. Of the 23, AlphaMissense provides support for pathogenicity for only 7 while

supporting benignity for the remaining 16 (using the developer-recommended score thresholds

of >0.564 and <0.34, respectively[40]; the average score for all 23 is 0.36). Meanwhile MPC,

where missense constraint is a key component, predicts scores >2 for all 23 variants (average

score of 2.76 or 97th percentile). Still, detachment from structural information means that MCR

identification is restricted to a linear search along the genome, and we are thus less powered to

discover 3D-specific constraint clusters such as small binding pockets.

For the MPC variant-level deleteriousness predictor, certain protein domain and structural

information is learned with the PolyPhen-2 model feature, though this has limitations compared

to full 3D structure-based methods like AlphaMissense. We thus evaluated the utility of

combining MPC and AlphaMissense in predicting missense deleteriousness. We re-categorized

the DD, AUT, and control *de novo* missense variants by whether they were predicted to have

high or low deleteriousness scores by MPC and AlphaMissense (≥2.5 and <2 for MPC, ≥0.9985

and <0.761 for AlphaMissense; see **Supplementary Note** for threshold selection) and

calculated case-control enrichments. Variants predicted to be highly deleterious by both MPC

and AlphaMissense have stronger enrichment than when predicting by either metric alone, and

variants where deleteriousness is high in one metric and low in the other also harbor detectable

enrichments (**Supplementary Fig. 19, Supplementary Table 18**). This demonstrates that MPC

and AlphaMissense offer complementary information and joint application can improve

prediction performance. However, we caution against taking the simple maximum of the MPC or

AlphaMissense relative deleteriousness for any variant — this sharply decreases predictive

performance in ranking case vs. control *de novo* variants (**Supplementary Fig. 18,**

Supplementary Table 17), arising from decreased specificity wherein MPC and AlphaMissense

each assign relatively high predicted deleteriousness to subsets of the control *de novo* variants.

Lastly, to guide usage of the MPC model, we note that MPC is best suited to modeling strong

fitness effects: the variants composing the deleterious training set were chosen to be those

likely under strong selection given constraint and disease association evidence, and the

missense constraint feature of the MPC model is, as aforementioned, grounded in heterozygous

selection. This is demonstrated by MPC's excellent performance in distinguishing *de novo*

variants in a severe DD cohort from controls (**Fig. 5c**), which is driven by a combination of

model design and unavoidable training biases as the training set naturally includes known DD

genes that severely disrupt development.

In summary, we identify 36% of MANE Select or canonical transcripts with variable levels of

missense constraint and demonstrate that coding regions specifically depleted of missense

variation in the general population are enriched for disease-associated variation. Additionally,

we show that this depletion of missense variation provides moderate strength evidence when

clinically classifying variants according to ACMG/AMP guidelines and that incorporation of

regional missense constraint into an *in silico* predictor effectively prioritizes a subset of *de novo*

missense variation in individuals with developmental phenotypes for association testing. We

have publicly released these data for use in both research and clinical settings. We anticipate

refined resolution of these metrics as datasets grow, both in size and in ancestral

representation, and with the incorporation of complementary structural or functional data.

Methods

Transcripts

This study followed the same steps of transcript annotation and the Ensembl Variant Effect

Predictor (VEP) table as described in gnomAD v4.1.1[50] and analyzed only coding transcripts as

defined by MANE Select v0.95 and, when not present, the canonical transcript as listed in

GENCODE v39/Ensembl VEP v105. For high quality transcripts, we excluded transcripts that

had outlier variant counts: zero expected or too many observed pLoF, missense, or

synonymous variants; or too few observed synonymous variants. This totaled 17,841

transcripts, 96% from MANE Select and 4% canonical. We also make MCR and MPC

information available for the 1,534 transcripts with outlier counts, but caution that scores may be

less accurate in these sequences.

Genome Aggregation Database (gnomAD) population data

All analyses in this paper were conducted using the 730,947 gnomAD v4.1.1 exomes on

GRCh38[50]. Coverage for gnomAD v4.1.1 was calculated using information from sample

genomic VCF (gVCF) data rather than from read data. As a result, coverage information for

gnomAD v4.1.1 is not as granular due to the reference block structure within gVCFs. To remedy
this, we used allele number percent (%AN) to proxy coverage for all analyses to improve our
ability to capture constraint in lower-coverage sites. Sites with %AN < 20 (0.68% of all possible
missense sites with %AN > 0) were excluded from analysis. "High-coverage" sequences refer to
MCRs with %AN ≥ 90. The gnomAD allele frequencies used in this study are those calculated
across all individuals. LOEUF and transcript missense OE scores are from the gnomAD v4.1.1
gene constraint data[50].
549
ClinVar variants

We annotated functional consequences for ClinVar[8] (v.20250504) variants using the Ensembl
VEP table. For this study, we selected high-quality missense ClinVar variants with non-
conflicting pathogenic (P), likely pathogenic (LP), variant of uncertain significance (VUS), benign
(B), or likely benign (LB) classification and a review status of at least one star. For these high-
quality missense variants, 89.9% are VUS (1,585,522/1,764,005).
556
557   **OMIM data**

558   We used OMIM (v.20230215) to annotate gene-phenotype modes of inheritance (MoI). Genes
559   were considered autosomal dominant if their only MoI noted in OMIM was autosomal dominant,
560   likewise for autosomal recessive. In total, 1,036 genes were considered autosomal dominant,
561   and 2,176 were considered autosomal recessive.
562
Gene2Phenotype (G2P) data

Phenotype associations and mechanistic information for developmental disorder (DD) genes
were taken from the G2P DD panel (v.20251127). We filtered to 810 genes with definitive or
strong evidence and a monoallelic mechanism for DD association. The 359 genes considered to
be non-LoF were those where the noted variant consequence included "altered gene product
structure" or "increased gene product level", and the noted molecular mechanism was not loss-
of-function.
570
Missense Tolerance Ratio (MTR) scores

We obtained MTR scores estimated from the exomes of 983,578 individuals[62]. This method
provides a window-based estimate of observed/expected missense variant counts, similar to the
gnomAD missense OE presented here. Scores are available at the amino acid-level (smoothed
across a window of 31 amino acids), or a segment (region) level calculated by aggregating
consecutive amino acids below a certain "constrained" OE threshold. For the analysis
calibrating metric thresholds for clinical interpretation, we used the amino acid-level scores.
578
Rare and *de novo* variants from developmental disorder (DD) cohorts

Case *de novo* variants for association analyses were obtained from studies of severe DD[15],
autism[16] (AUT), congenital heart disease[70] (CHD), orofacial cleft[69] (OFC), congenital
diaphragmatic hernia[71] (CDH), and epileptic encephalopathy[68] (EE). Control *de novo* mutations
were obtained from siblings of the autistic probands without DDs[16]. Variants were lifted over
from GRCh37 to GRCh38 using the "liftover" function in Hail for the DD, CDH, and EE cohorts.

Variant functional consequences were re-annotated using the Ensembl VEP table. Variants
transmitted and not transmitted from parents to autistic probands were procured from previously
published ASC-SSC and SPARK cohorts, and case-control variants for autism were procured
from previously published iPSYCH and Swedish cohorts[16]. Both the transmitted/untransmitted
and case-control variant sets were filtered to gnomAD AF < 0.1%.

Identifying breakpoints within transcripts of regional missense constraint

We defined the set of sites with possible missense variants using a synthetic Hail Table (HT)
containing all possible single nucleotide variants in the exome. We annotated this HT with the
Ensembl VEP table and filtered it to variants with the consequence "missense_variant" in the
MANE Select or canonical coding transcripts as defined in *Transcripts*. We then further filtered
to (1) variants observed in gnomAD with allele count (AC) > 0, AF < 0.001, %AN ≥ 20, and QC
PASS; or (2) variants unobserved in gnomAD with AC = 0. We then aggregated the total
number of observed variant counts per base pair site by summing the number of observed
variants over each possible substitution (reference and alternate allele) at the locus.

As described in more detail for the gnomAD v4.1.1 release[50], we applied two models (plateau
and coverage) to calculate the proportion of expected missense variation per variant
(chromosome/locus plus reference and alternate alleles). As with the observed missense
counts, we then summed the number of expected variants to get the total expected rare
missense variant count per locus.

We applied a likelihood ratio test to determine whether the missense observed/expected (OE)
ratio was uniform along a transcript or whether a transcript had evidence for discrete
subsections (regions) with distinct missense constraint strengths. We assumed that the
observed missense counts should follow a Poisson distribution around the expected missense
counts. We defined our null model as transcripts not having any evidence of regional variability
in missense depletion (where the OE ratio is consistent across the length of the transcript). Our
alternative model was that transcripts exhibited evidence of distinct subsections of missense
depletion (OE ratio calculated per transcript subsection). Because the alternative model should
always have a better fit than the null model, we require a chi-squared value above a given
threshold (p = 0.001) to establish significance. We used the following formulas to determine the
significance of a breakpoint that would split a transcript into two subsections, A and B:

- $p_0 = Pois(obs_A, exp_A * OE) * Pois(obs_B, exp_B * OE)$
- $p_1 = Pois(obs_A, exp_A * OE_A) * Pois(obs_B, exp_B * OE_B)$
- $\chi^2 = 2(ln(p_1) - ln(p_0))$

where OE is the missense observed/expected ratio across the entire transcript, obs_A is the
number of observed missense variants in transcript subsection A, exp_A is the number of
expected missense variants in transcript subsection A, OE_A is the OE ratio across transcript
subsection A, obs_B is the number of observed missense variants in transcript subsection B, exp_B
is the number of expected missense variants in transcript subsection B, OE_B is the OE ratio
across transcript subsection B, and Pois is the Poisson likelihood.

629
630 Likewise, we used the following formulas to determine the significance of a breakpoint that
631 would split a transcript into three subsections, A, B, and C:

632
633 • $p_0 = Pois(obs_A, exp_A * OE) * Pois(obs_B, exp_B * OE) * Pois(obs_C, exp_C * OE)$
634 • $p_1 = Pois(obs_A, exp_A * OE_A) * Pois(obs_B, exp_B * OE_B) * Pois(obs_C, exp_C * OE_C)$
635 • $\chi^2 = 2(ln(p_1) - ln(p_0))$

636
For the purposes of this breakpoint search, all transcript subsections with more observed
variants than expected were capped at an OE of 1, as we were looking for areas of missense
depletion and not missense enrichment. We also converted the expected counts for transcript
subsections with zero expected variants from 0 to 10^{-9} to prevent nonfinite OE values. We
implemented a minimum number of expected missense variants (16) to prevent identification of
breakpoint positions that would create very small (i.e., a handful of base pairs in size) transcript
subsections that are more likely to be spurious (see **Supplementary Note**).

644
To search for a single breakpoint that would divide a transcript into two subsections, we
calculated chi-squared statistics to conduct likelihood ratio tests simultaneously for every eligible
position within a transcript. The positions we considered were base pair sites with a possible
missense variant substitution that had at least 16 expected missense counts in either direction
(i.e., both transcript subsections created by dividing the transcript at this point would have at
least 16 expected missense variants). Breakpoints were determined by the site with the
maximum chi-squared value that was significant at p = 0.001. Any transcripts that did not have a
single significant breakpoint were searched for two simultaneous breakpoints that would divide
the transcript into three sections. As in the single-breakpoint scenario, we tested all pairs of
eligible base pair sites where the resulting transcript subsections would harbor at least 16
expected missense variants. For transcripts where we identified significant breakpoints, we
repeated this process (single-breakpoint search, then two-breakpoint search) recursively on
each subsection until no further significant breakpoints were found.

658
Clinical calibration of missense constraint regions

We filtered to genes that had at least one pathogenic missense variant in ClinVar and selected
missense variants in those genes with a non-conflicting pathogenic (P), likely pathogenic (LP),
benign (B), or likely benign (LB) classification and a review status of at least one star. Missense
variants with an AF ≥ 1% in gnomAD v4.1.1 were removed, leaving a total of 122,269 ClinVar
variants in 4,334 genes. For this set of genes, we also retrieved QC PASS variants with AF <
1% from gnomAD v4.1.1, which resulted in 3,847,501 gnomAD variants from 4,302 genes.

666
All variants were annotated with MCR missense OE and the amino acid-level missense
tolerance ratio (MTR) metric[62]. Using our previously developed method[34] and the prior
probability of pathogenicity of 4.41%, we computed the local posterior probability curves for all
scores. P/LP and B/LB ClinVar variants were used to estimate the local values of the posterior
probability of pathogenicity, and gnomAD variants were used to smooth out the posterior
estimates. We defined score threshold ranges for each strength of evidence (supporting,

moderate, strong, and very strong) for pathogenicity and benignity based on the ACMG/AMP

criteria (**Supplementary Table 13**). For both metrics, we plotted the local posterior probability

curve estimate along with the one-sided 95% confidence interval, determined using 10,000

bootstrapping iterations.

Modeling deleteriousness of individual missense variants

We designed our missense variant deleteriousness predictor (Missense deleteriousness

Prediction by Constraint or MPC) to explicitly incorporate information on amino acid substitution

class and position-specific variant effects informed by missense constraint. Briefly, we trained a

machine learning model to differentiate "pathogenic" missense variants likely under strong

heterozygous selection from "benign" missense variants likely under neutral to near-neutral

selection. The "pathogenic" training set consisted of high-quality ClinVar variants (see *ClinVar*

variants) labeled as P or LP in 2,987 predicted-haploinsufficient genes (defined as having

probability of haploinsufficiency (pHaplo) ≥ 0.86[64]) or in 359 genes with DD associations in G2P

through non-LoF mechanisms (see *Gene2Phenotype data*). The "benign" training set consisted

of high-quality gnomAD variants AF > 0.1% or ClinVar B/LB variants in the same genes as the

"pathogenic" set (see *gnomAD variants* and *ClinVar variants*). Variants matching criteria for both

the "benign" and "pathogenic" sets were removed from the training data. We further removed

variants with %AN < 20 in the gnomAD v4.1.1 exomes and variants that were not in the 17,841

MANE Select/canonical transcripts.

We evaluated logistic regression and XGBoost (gradient-boosted tree) architectures,

incorporating features of MCR and gene constraint, amino acid substitution class severity,

PolyPhen-2 for orthogonal information from protein homology and structure, and phyloP for

evolutionary selection information across longer timescales. Only variants with all relevant

annotations were retained for training and for the full variant score table. The XGBoost model

was chosen over logistic regression based on performance on a hold-out gene set and then

retrained on all data (20,931 "pathogenic" variants and 93,638 "benign" variants) for the final

model. We applied the XGBoost model across the 70,313,598 possible exome-wide missense

variants in the Ensembl VEP table with all features to obtain a probability of pathogenicity for

each variant. The MPC score for any missense variant *i* is given as:

$d_i = -log_{10}(m_i/M)$

where d_i is the MPC score, m_i is the number of "benign" missense variants in the training set

with a fitted probability of pathogenicity value that is less than the fitted value for variant *i*, and *M*

is the total number of "benign" missense variants in the training set. When m_i is 0, i.e. when the

fitted score for variant *i* is more severe than for all "benign" variants, this introduces a log-zero

error so we set d_i to 6 (the maximum real value for d_i is just over 5 when m_i is 1). Larger values

of d_i indicate stronger predicted-deleteriousness. See **Supplementary Note** and

Supplementary Figs. 13-16 for further detail on these methods and the distribution of MPC

scores.

Comparison of MPC to other predictors

We compared our model to the following *in silico* missense deleteriousness predictors:
AlphaMissense[40], popEVE[41], REVEL[39], CADD[65,66], PolyPhen-2[63], and SIFT[67]. AlphaMissense
scores were downloaded from https://github.com/google-deepmind/alphamissense and mapped
to variants by locus and gene name, as the transcript provided was not always the MANE Select
transcript. popEVE scores were downloaded from https://pop.evemodel.org. REVEL scores
were downloaded from https://sites.google.com/site/revelgenomics/. PolyPhen-2 and SIFT
scores were obtained from VEP[84]. CADD scores were downloaded from the CADD website
(https://cadd.gs.washington.edu/download). phyloP scores were downloaded from the UCSC
browser (https://genome.ucsc.edu/cgi-bin/hgTrackUi?db=hg38&g=cons241way). We annotated
the case and control *de novo* missense variants described in *Rare and de novo variants from*
developmental disorder cohorts and ranked the variants based on their annotated scores. To
assess each predictor's ability to stratify case and control variation, we assessed the proportion
of case to control variants among the variants with the top 1%, 5%, or 10% for each score and
compared to the remaining 99%, 95%, and 90% of variants, respectively, using Fisher's exact
test.

Data availability

The missense constraint regions (MCRs) are displayed on the gnomAD v4 browser
(https://gnomad.broadinstitute.org) and available for download on the gnomAD website
(https://gnomad.broadinstitute.org/data#v4) and in the gnomAD v4 public datasets on Google
and Amazon clouds (gs://gcp-public-data--gnomad/papers/2026-rmc). MPC scores for all
possible variants in MANE Select canonical transcripts are available in the gnomAD v4 public
datasets on Google (gs://gcp-public-data--gnomad/papers/2026-rmc/gnomad_v4.1.1_mpc.ht).
gnomAD v4 exome, genome, and allele number data and the table of all possible single
nucleotide polymorphisms used to calculate mutational models and search for MCRs are also
available in the gnomAD public buckets and are accessible using code in the gnomAD Hail
utilities GitHub repository
(https://github.com/broadinstitute/gnomad_methods/blob/b61aff692041dbf1118773b18104f13d1
31008fa/gnomad/resources/grch38/gnomad.py#L456 and
https://github.com/broadinstitute/gnomad_methods/blob/b61aff692041dbf1118773b18104f13d1
31008fa/gnomad/resources/grch38/reference_data.py#L147).

ClinVar data were downloaded from ClinVar's FTP server
(https://ftp.ncbi.nlm.nih.gov/pub/clinvar/). *De novo* variants were extracted from the
supplementary files of the cited studies.

Code availability

Code to determine missense constraint regions (MCRs) and calculate MPC is available at
https://github.com/broadinstitute/regional_missense_constraint. Code used to generate the
mutational models is available at
https://github.com/broadinstitute/gnomad_methods/blob/main/gnomad/utils/constraint.py and

https://github.com/broadinstitute/gnomad-constraint. Code used to generate figures is available
at https://github.com/broadinstitute/rmc-2026-figures. The Hail library is available at
https://hail.is/.

Acknowledgments

We thank all of the individuals who contributed their data to gnomAD for enabling the research
presented here and in numerous other publications. We also thank members of the gnomAD
Production and Browser Teams for their behind-the-scenes efforts in producing and sharing the
gnomAD dataset, Dan King and the rest of the Hail team for enabling data processing at scale,
and Emma Pierce-Hoffman for her work curating gene lists. This work was supported by the
National Human Genome Research Institute (NHGRI; U24HG011450 to H.L.R. and M.J.D.;
U01HG011755 to A.O.D.L., H.L.R, and M.E.T.; U24HG006834 to H.L.R, U01HG012022 to P.R.;
5T32HG002295 to L.W.; R01HG012867 to K.E.S.), the Eunice Kennedy Shriver National
Institute of Child Health and Human Development (NICHD; F31HD111109 to L.W.), the National
Science Foundation (NSF; GRFP#2022339661 to L.W.; GRFP#2023361442 to H.A.), the
Simons Foundation (SFARI 1009802 to M.E.T. and K.E.S.), a National Health and Medical
Research Council investigator grant (2009982) to D.G.M., and the Massachusetts General
Hospital Executive Committee on Research, Interim Support Funding (K.E.S.).
773

Author information

775 These authors contributed equally: Lily Wang and Katherine R. Chao.

Contributions

L.W., K.R.C., K.E.S., B.M.N, and M.J.D conceived and designed experiments. L.W., K.R.C, and
K.E.S. performed primary writing of the manuscript. L.W., K.R.C., R.P., C.L., H.A., and R.Y.
performed the analyses and generated figures. P.S. and J.C. were instrumental to developing
methods. R.H.G., N.A.W., and M.S. developed visualizations for the web browser. B.W., W.P.,
M.W.W., K.M.L., J.K.G, K.J.K, and G.T. completed code review for methods. D.G., J.I.G., C.V.,
and T.P. helped debug runtime compute. J.A.K. provided data and analysis suggestions. S.B.
contributed analysis suggestions. H.L.R., B.M.N., M.E.T, D.G.M, A.O.D.L., K.J.K., P.R., M.J.D.,
and K.E.S. supervised the research. All authors listed under the Genome Aggregation Database
Consortium contributed to the generation of the primary data incorporated into the gnomAD
resource. All authors reviewed the manuscript.

Ethics declarations

Competing interests/Declaration of interests

J.A.K is a current employee at the Regeneron Genetics Center and a shareholder of Regeneron
Pharmaceuticals. T.P. and G.T. are founders of E9 Genomics, Inc.. H.L.R. has received support
from Illumina and Microsoft to support rare disease gene discovery and diagnosis. B.M.N. is a

member of the scientific advisory board at Deep Genomics and Neumora. M.E.T. has received

research support and/or reagents from Illumina, Pacific Biosciences, Microsoft, Oxford

Nanopore, and Ionis Therapeutics. D.G.M. is a paid advisor to GSK, Insitro, and Overtone

Therapeutics, and receives research funding from Microsoft. A.O.D.L. has consulted for Addition

Therapeutics and received research support from Pacific Biosciences for rare disease

diagnosis. K.J.K. is on the Scientific Advisory Board of Nurture Genomics. M.J.D. is a founder of

Maze Therapeutics and Neumora Therapeutics, Inc. (f/k/a RBNC Therapeutics). K.E.S. has

received research support from Microsoft for work related to rare disease diagnostics. All other

authors declare no competing interests.

References

1. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).
2. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291 (2016).
3. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).
4. Taliun, D. *et al.* Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature* **590**, 290–299 (2021).
5. Backman, J. D. *et al.* Exome sequencing and analysis of 454,787 UK Biobank participants. *Nature* **599**, 628–634 (2021).
6. All of Us Research Program Genomics Investigators. Genomic data in the All of Us Research Program. *Nature* (2024) doi:10.1038/s41586-023-06957-x.
7. McInnes, G. *et al.* Opportunities and challenges for the computational interpretation of rare variation in clinically important genes. *Am. J. Hum. Genet.* **108**, 535–548 (2021).
8. Landrum, M. J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res.* **48**, (2020).
9. McEwen, A. E., Tejura, M., Fayer, S., Starita, L. M. & Fowler, D. M. Multiplexed assays of variant effect for clinical variant interpretation. *Nat Rev Genet* **27**, 137–154 (2026).
10. Petrovski, S. *et al.* The Intolerance of Regulatory Sequence to Genetic Variation Predicts Gene Dosage Sensitivity. *PLoS Genet.* **11**, e1005492 (2015).
11. Samocha, K. E. *et al.* A framework for the interpretation of de novo mutation in human disease. *Nat. Genet.* **46**, 944–950 (2014).
12. Weghorn, D. *et al.* Applicability of the Mutation-Selection Balance Model to Population Genetics of Heterozygous Protein-Truncating Variants in Humans. *Mol. Biol. Evol.* **36**, 1701–1710 (2019).
13. Agarwal, I., Fuller, Z. L., Myers, S. R. & Przeworski, M. Relating pathogenic loss-of-function mutations in humans to their evolutionary fitness costs. *Elife* **12**, (2023).
14. Kosmicki, J. A. *et al.* Refining the role of de novo protein-truncating variants in neurodevelopmental disorders by using population reference samples. *Nat. Genet.* **49**, 504–510 (2017).
15. Kaplanis, J. *et al.* Evidence for 28 genetic disorders discovered by combining healthcare and research data. *Nature* **586**, 757–762 (2020).
16. Fu, J. M. *et al.* Rare coding variation provides insight into the genetic architecture and phenotypic context of autism. *Nat. Genet.* **54**, 1320–1331 (2022).
17. Bamshad, M. J., Nickerson, D. A. & Chong, J. X. Mendelian Gene Discovery: Fast and Furious with No End in Sight. *Am. J. Hum. Genet.* **105**, 448–455 (2019).
18. Seaby, E. G., Rehm, H. L. & O'Donnell-Luria, A. Strategies to Uplift Novel Mendelian Gene Discovery for Improved Clinical Outcomes. *Front. Genet.* **12**, 674295 (2021).
19. Chen, E. *et al.* Rates and Classification of Variants of Uncertain Significance in Hereditary Disease Genetic Testing. *JAMA Netw Open* **6**, e2339571 (2023).
20. Samocha, K. E. *et al.* Regional missense constraint improves variant deleteriousness

prediction. *bioRxiv* 148353 (2017) doi:10.1101/148353.

21. Gussow, A. B., Petrovski, S., Wang, Q., Allen, A. S. & Goldstein, D. B. The intolerance to functional genetic variation of protein domains predicts the localization of pathogenic mutations within genes. *Genome Biol.* **17**, 9 (2016).

22. Wiel, L., Venselaar, H., Veltman, J. A., Vriend, G. & Gilissen, C. Aggregation of population-based genetic variation over protein domain homologues and its potential use in genetic diagnostics. *Hum. Mutat.* **38**, 1454–1463 (2017).

23. Sivley, R. M., Dou, X., Meiler, J., Bush, W. S. & Capra, J. A. Comprehensive Analysis of Constraint on the Spatial Distribution of Missense Variants in Human Protein Structures. *Am. J. Hum. Genet.* **102**, 415–426 (2018).

24. Lal, D. *et al.* Gene family information facilitates variant interpretation and identification of disease-associated genes in neurodevelopmental disorders. *Genome Med.* **12**, 28 (2020).

25. Traynelis, J. *et al.* Optimizing genomic medicine in epilepsy through a gene-customized approach to missense variant interpretation. *Genome Res.* **27**, 1715–1729 (2017).

26. Perszyk, R. E., Kristensen, A. S., Lyuboslavsky, P. & Traynelis, S. F. Three-dimensional missense tolerance ratio analysis. *Genome Res.* **31**, 1447–1461 (2021).

27. Havrilla, J. M., Pedersen, B. S., Layer, R. M. & Quinlan, A. R. A map of constrained coding regions in the human genome. *Nat. Genet.* **51**, 88–95 (2019).

28. Silk, M. *et al.* MTR3D: identifying regions within protein tertiary structures under purifying selection. *Nucleic Acids Res.* **49**, W438–W445 (2021).

29. Hicks, M., Bartha, I., di Iulio, J., Venter, J. C. & Telenti, A. Functional characterization of 3D protein structures informed by human genetic diversity. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 8960–8965 (2019).

30. Li, B., Roden, D. M. & Capra, J. A. The 3D mutational constraint on amino acid sites in the human proteome. *Nat. Commun.* **13**, 3273 (2022).

31. Zhang, X. *et al.* Genetic constraint at single amino acid resolution in protein domains improves missense variant prioritisation and gene discovery. *Genome Med* **16**, 88 (2024).

32. Satterstrom, F. K. *et al.* Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **180**, 568–584.e23 (2020).

33. Richards, S. *et al.* Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet. Med.* **17**, 405–424 (2015).

34. Pejaver, V. *et al.* Calibration of computational tools for missense variant pathogenicity classification and ClinGen recommendations for PP3/BP4 criteria. *Am. J. Hum. Genet.* **109**, 2163–2177 (2022).

35. Bergquist, T. *et al.* Calibration of additional computational tools expands ClinGen recommendation options for variant classification with PP3/BP4 criteria. *Genet Med* **27**, 101402 (2025).

36. Tejura, M. *et al.* A scalable approach to resolving variants of uncertain significance. *bioRxiv* 2026.02.14.705848 (2026) doi:10.64898/2026.02.14.705848.

37. Brnich, S. E. *et al.* Recommendations for application of the functional evidence PS3/BS3 criterion using the ACMG/AMP sequence variant interpretation framework. *Genome Med* **12**, 3 (2019).

38. Zeiberg, D. *et al.* Gene-based calibration of high-throughput functional assays for clinical variant classification. *bioRxiv* (2025) doi:10.1101/2025.04.29.651326.

39. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877–885 (2016).

40. Cheng, J. *et al.* Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science* **381**, eadg7492 (2023).

41. Orenbuch, R. *et al.* Proteome-wide model for human disease genetics. *Nat Genet* **57**, 3165–3174 (2025).

42. Jensen, D. E. *et al.* BAP1: a novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1-mediated cell growth suppression. *Oncogene* **16**, 1097–1112 (1998).

43. Machida, Y. J., Machida, Y., Vashisht, A. A., Wohlschlegel, J. A. & Dutta, A. The deubiquitinating enzyme BAP1 regulates cell growth via interaction with HCF-1. *J Biol Chem* **284**, 34179–34188 (2009).

44. Küry, S. *et al.* Rare germline heterozygous missense variants in BRCA1-associated protein 1, BAP1, cause a syndromic neurodevelopmental disorder. *Am. J. Hum. Genet.* **109**, 361–372 (2022).

45. UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic Acids Res* **53**, D609–D617 (2025).

46. Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic Acids Res* **53**, D444–D456 (2025).

47. Szczepanski, A. P. & Wang, L. Emerging multifaceted roles of BAP1 complexes in biological processes. *Cell Death Discov* **7**, 20 (2021).

48. Murali, R., Wiesner, T. & Scolyer, R. A. Tumours associated with BAP1 mutations. *Pathology* **45**, 116–126 (2013).

49. Louie, B. H. & Kurzrock, R. BAP1: Not just a BRCA1-associated protein. *Cancer Treat Rev* **90**, 102091 (2020).

50. Guez, J. *et al.* Integrating 730,947 exome sequences with clinical literature improves gene discovery. *medRxiv* (2026) doi:10.64898/2026.03.23.26349081.

51. Neumann, M. *et al.* Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* **314**, 130–133 (2006).

52. Arai, T. *et al.* TDP-43 is a component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Biochem Biophys Res Commun* **351**, 602–611 (2006).

53. Gitcho, M. A. *et al.* TDP-43 A315T mutation in familial motor neuron disease. *Ann Neurol* **63**, 535–538 (2008).

54. Sreedharan, J. *et al.* TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science* **319**, 1668–1672 (2008).

55. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

56. Necci, M., Piovesan, D., Dosztányi, Z. & Tosatto, S. C. E. MobiDB-lite: fast and highly specific consensus prediction of intrinsic disorder in proteins. *Bioinformatics* **33**, 1402–1404 (2017).

57. Hamosh, A., Amberger, J. S., Bocchini, C., Scott, A. F. & Rasmussen, S. A. Online

Mendelian Inheritance in Man (OMIM®): Victor McKusick's magnum opus. *Am. J. Med. Genet. A* **185**, 3259–3265 (2021).

58. Zeng, T., Spence, J. P., Mostafavi, H. & Pritchard, J. K. Bayesian estimation of gene constraint from an evolutionary model with gene features. *Nat Genet* **56**, 1632–1643 (2024).

59. Seplyarskiy, V. *et al.* A mutation rate model at the basepair resolution identifies the mutagenic effect of polymerase III transcription. *Nat Genet* **55**, 2235–2242 (2023).

60. Thormann, A. *et al.* Flexible and scalable diagnostic filtering of genomic variants using G2P with Ensembl VEP. *Nat. Commun.* **10**, 2373 (2019).

61. Christmas, M. J. *et al.* Evolutionary constraint and innovation across hundreds of placental mammals. *Science* **380**, eabn3943 (2023).

62. Sun, K. Y. *et al.* A deep catalogue of protein-coding variation in 983,578 individuals. *Nature* **631**, 583–592 (2024).

63. Adzhubei, I. A. *et al.* A method and server for predicting damaging missense mutations. *Nat. Methods* **7**, 248–249 (2010).

64. Collins, R. L. *et al.* A cross-disorder dosage sensitivity map of the human genome. *Cell* **185**, 3041–3055.e25 (2022).

65. Kircher, M. *et al.* A general framework for estimating the relative pathogenicity of human genetic variants. *Nat. Genet.* **46**, 310–315 (2014).

66. Rentzsch, P., Schubach, M., Shendure, J. & Kircher, M. CADD-Splice-improving genome-wide variant effect prediction using deep learning-derived splice scores. *Genome Med.* **13**, 31 (2021).

67. Ng, P. C. & Henikoff, S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res.* **31**, 3812–3814 (2003).

68. EuroEPINOMICS-RES Consortium, Epilepsy Phenome/Genome Project & Epi4K Consortium. De novo mutations in synaptic transmission genes including DNM1 cause epileptic encephalopathies. *Am. J. Hum. Genet.* **95**, 360–370 (2014).

69. Bishop, M. R. *et al.* Genome-wide Enrichment of De Novo Coding Mutations in Orofacial Cleft Trios. *Am. J. Hum. Genet.* **107**, 124–136 (2020).

70. Jin, S. C. *et al.* Contribution of rare inherited and de novo variants in 2,871 congenital heart disease probands. *Nat. Genet.* **49**, 1593–1601 (2017).

71. Qiao, L. *et al.* Rare and de novo variants in 827 congenital diaphragmatic hernia probands implicate LONP1 as candidate risk gene. *Am. J. Hum. Genet.* **108**, 1964–1980 (2021).

72. Fu, W. *et al.* Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants. *Nature* **493**, 216–220 (2013).

73. Chao, K. R. *et al.* The landscape of regional missense mutational intolerance quantified from 125,748 exomes. *bioRxiv* (2024) doi:10.1101/2024.04.11.588920.

74. Cassa, C. A. *et al.* Estimating the selective effects of heterozygous protein-truncating variants from human exome data. *Nat Genet* **49**, 806–810 (2017).

75. Judd, J. *et al.* Allele Frequencies at Recessive Disease Genes are Mainly Determined by Pleiotropic Effects in Heterozygotes. *bioRxiv* (2025) doi:10.64898/2025.12.05.692665.

76. Fridman, H., Khazeeva, G., Levy-Lahad, E., Gilissen, C. & Brunner, H. G. Reproductive and cognitive phenotypes in carriers of recessive pathogenic variants. *Nat Hum Behav* **9**, 1726–1736 (2025).

77. Lombardi, M. *et al.* Variability in Clinical Phenotype in TARDBP Mutations: Amyotrophic Lateral Sclerosis Case Description and Literature Review. *Genes (Basel)* **14**, (2023).

78. Kirby, J. *et al.* Broad clinical phenotypes associated with TAR-DNA binding protein (TARDBP) mutations in amyotrophic lateral sclerosis. *Neurogenetics* **11**, 217–225 (2010).

79. Newell, K. *et al.* Dysregulation of TDP-43 intracellular localization and early onset ALS are associated with a TARDBP S375G variant. *Brain Pathol* **29**, 397–413 (2019).

80. Jih, K.-Y. *et al.* TARDBP variants in Taiwanese ALS patients: Genetic spectrum, clinical features, and founder effects. *J Neurol Sci* **474**, 123531 (2025).

81. Luppino, F., Lenz, S., Chow, C. F. W. & Toth-Petroczy, A. Deep learning tools predict variants in disordered regions with lower sensitivity. *BMC Genomics* **26**, 367 (2025).

82. Zhong, G., Zhao, Y., Zhuang, D., Chung, W. K. & Shen, Y. PreMode predicts mode-of-action of missense variants by deep graph representation learning of protein sequence and structural context. *Nat Commun* **16**, 7189 (2025).

83. Khanna, T., Hanna, G., Sternberg, M. J. E. & David, A. Missense3D-DB web catalogue: an atom-based analysis and repository of 4M human protein-coding genetic variants. *Hum Genet* **140**, 805–812 (2021).

84. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).

Genome Aggregation Database Consortium

Maria Abreu[1], Amina Abubakar[2,3], Rolf Adolfsson[4], Carlos A. Aguilar Salinas[5], Tariq Ahmad[6], Elissa Alarmani[7], Christine M. Albert[8,9], Jessica Alföldi[7,10], Matthieu Allez[11], Irina M. Armean[7,10], Elizabeth G. Atkinson[12,13], Gil Atzmon[14,15], Eric Banks[16], John Barnard[17], Daniel Ben-Isvy[7,18], Emelia J. Benjamin[19,20,21], David Benjamin[16], Louis Bergelson[16], Charles Bernstein[22,23], Douglas Blackwood[24], Michael Boehnke[25], Erwin P. Bottinger[26], Donald W. Bowden[27], Matthew J. Bown[28], Harrison Brand[18,29], Steven Brant[30,31,32], Ted Brookings[16,33], Sam Bryant[10,34], Shawneequa L. Callier[35,36], Hannia Campos[37,38], John C. Chambers[39,40,41], Juliana C. Chan[42], Sinéad Chapman[7,10,43], Daniel I. Chasman[8,44], Lea A. Chen[45], Siwei Chen[7,10], Rex Chisholm[46], Judy Cho[26], Mina K. Chung[47], Wendy K. Chung[48], Kristian Cibulskis[16], Bruce Cohen[49,50], Ryan L. Collins[7,18,51], Kristen M. Connolly[52], Adolfo Correa[53], Aiden Corvin[54], Miguel Covarrubias[16], Nick Craddock[55], Beryl B. Cummings[7,51], Dana Dabelea[56], John Danesh[57], Dawood Darbar[58], Philip W. Darnowsky[7], Stacey Donnelly[59], Richard H. Duerr[60,61,62], Ravindranath Duggirala[63], Josée Dupuis[64,65], Patrick T. Ellinor[7,66], Roberto Elosua[67,68,69], Eleina England[7,70], Jeanette Erdmann[71,72,73], Tõnu Esko[74], Emily Evangelista[7], Yossi Farjoun[75], Diane Fatkin[76,77,78], William Faubion[79], Steven Ferriera[80], Gemma Figtree[81,82,83], Jose Florez[44,84,85], Laurent Francioli[7,10], Andre Franke[86,87], Adam Frankish[88], Jack Fu[7,18,89], Martti Färkkilä[90,91,92], Stacey Gabriel[80], Kiran Garimella[16], Laura D. Gauthier[16], Jeff Gentry[16], Michel Georges[93], Gad Getz[44,94,95], David C. Glahn[96,97], Benjamin Glaser[98], Stephen J. Glatt[99], Fernando S. Goes[100], Clicerio Gonzalez[101], Leif Groop[102,103], Sanna Gudmundsson[7,104,105], Jeremy Guez[7,10], Michael Guo[7], Namrata Gupta[7,80], Andrea Haessly[16], Christopher Haiman[106], Craig L. Hanis[107], James Hanyok[108], Matthew Harms[109,110], Qin He[7], Mikko Hiltunen[111], Matti M. Holi[112], Christina M. Hultman[113,114], Steve Jahl[7,10], Chaim Jalas[115], Thibault Jeandet[16], Mikko Kallela[116], Diane Kaplan[16], Jaakko Kaprio[103], Sekar Kathiresan[18,44,117], Eimear E. Kenny[118], Bong-Jo Kim[119], Young Jin Kim[119], George Kirov[120], Zan Koenig[10,43], Jaspal Kooner[40,121,122], Pragati Kore[12], Seppo Koskinen[123], Harlan M. Krumholz[124,125], Subra Kugathasan[126], Juozas Kupcinskas[127], Nehir E. Kurtas[18], Soo Heon Kwak[128], Markku Laakso[129,130], Nicole Lake[131], Mikael Landén[113,132], Matthew Lebo[133,134,135], Terho Lehtimäki[136,137], Monkol Lek[131], James D. Lewis[138], Cecilia M. Lindgren[139,140,141], Christopher Llanwarne[16], Ruth J.F. Loos[26,142], Edouard Louis[143], Chelsea Lowther[18], Wenhan Lu[7], Steven A. Lubitz[7,66], Ronald C.W. Ma[42,144,145], Dara S. Manoach[146,147], Gregory M. Marcus[148], Jaume Marrugat[149,150], Nicholas A. Marston[151], Daniel M. Marten[7,105], Alicia R. Martin[7,10,43], Steven McCarroll[43,152], Mark I. McCarthy[153,154,155], Jacob L. McCauley[156,157], Dermot McGovern[158], Ruth McPherson[159], Andrew McQuillin[160], James B. Meigs[7,44,161], Olle Melander[162], Andres Metspalu[163], Deborah Meyers[164], Eric V. Minikel[7], Braxton D. Mitchell[165], Paul Moayyedi[166,167,168], Sanghamitra Mohanty[169], Andrés Moreno Estrada[170], Nicola J. Mulder[171,172], " NA, Joshua N. Nadeau[7], Aliya Naheed[173], Andrea Natale[174,175,176], Saman Nazarian[177,178], Charles Newton[179,180], Peter M. Nilsson[181], Sam Novod[16], Michael C. O'Donovan[55], Yukinori Okada[182,183,184], Dost Ongur[44,49], Roel A. Ophoff[185,186], Lorena Orozco[187,188], Willem Ouwehand[189], Michael J. Owen[55], Nick Owen[190], Colin Palmer[191], Nicholette D. Palmer[27], Aarno Palotie[10,43,103], Mara Parellada[192,193,194], Kyong Soo Park[128,195], Carlos Pato[196], Nancy L. Pedersen[113], Tina Pesaran[197], Emma Pierce-Hoffman[7], Sharon Plon[198], Danielle Posthuma[199,200], Ann E. Pulver[201], Aaron Quinlan[202], Dan Rader[177,203], Nazneen Rahman[204], Andreas Reif[205], Alex P. Reiner[206,207], Anne M. Remes[208,209], Stephen Rich[210,211],

John D. Rioux[212,213], Samuli Ripatti[59,103,214], Jason Roberts[215,216], Elise Robinson[18], Dan M. Roden[217,218], Jerome I. Rotter[219], Guy Rouleau[220], Christian T. Ruff[151], Heiko Runz[221], Marc S. Sabatine[151], Nareh Sahakian[16], Andrea Saltzman[7], Nilesh J. Samani[222,223], Alba Sanchis-Juan[18], Akira Sawa[224,225,226], Jeremiah Scharf[7,18,43], Molly Schleicher[7], Heribert Schunkert[227,228], Sebastian Schönherr[229], Eleanor G. Seaby[7,230], Svati H. Shah[231,232], Megan Shand[16], Ted Sharpe[16], Moore B. Shoemaker[233], Tai Shyong[234,235], Edwin K. Silverman[236,237], Moriel Singer-Berk[7], Jurgita Skieceviciene[127], Jonathan T. Smith[16], J. Gustav Smith[238,239,240], Jordan W. Smoller[43,237,241], Hilkka Soininen[242], Harry Sokol[243,244,245], Rachel G. Son[7], Tim Spector[246], David St Clair[247], Sarah Stenton[7], Christine Stevens[7,10,43], Nathan O. Stitziel[248,249,250], Patrick F. Sullivan[113,251], Jaana Suvisaari[252], Yekaterina Tarasova[7], Kent D. Taylor[219], Yik Ying Teo[253,254,255], Kathleen Tibbetts[16], Ming Tsuang[256,257], Tiinamaija Tuomi[103,258,259], Dan Turner[260], Teresa Tusie-Luna[261,262], Rachel Ungar[7], Grace VanNoy[7], Erkki Vartiainen[263], Marquis Vawter[264], Elisabet Vilella[265,266,267], Gordon Wade[16], Mark Walker[16], Qingbo Wang[7,268], Arcturus Wang[7,10,43], James S. Ware[7,269,270], Hugh Watkins[271], Rinse K. Weersma[272], Christopher Whelan[7], Nicola Whiffin[7,273,274], James G. Wilson[275], Lauren Witzgall[7], Ramnik J. Xavier[276,277], Mary T. Yohannes[7], Robert Yolken[278], Xuefang Zhao[7]

[1]University of Miami Miller School of Medicine, Gastroenterology, Miami, USA
[2]Neuroscience Research Group, Department of Clinical Sciences, Kenyan Medical Research Institute, Wellcome Trust, Kilifi, Kenya
[3]Institute for Human Development, Aga Khan University, Nairobi, Kenya
[4]Department of Clinical Sciences, Psychiatry, Umeå University, Sweden
[5]Unidad de Investigacion de Enfermedades Metabolicas, Instituto Nacional de Ciencias Medicas y Nutricion, Mexico City, Mexico
[6]Peninsula College of Medicine and Dentistry, Exeter, UK
[7]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[8]Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA
[9]Division of Cardiovascular Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA
[10]Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
[11]Gastroenterology Department, Hôpital Saint-Louis - APHP, Université Paris Cité, INSERM U1160, Paris, France
[12]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA
[13]Stanley Center for Psychiatric Research, The Broad Institute of MIT and Harvard, Cambridge MA, USA
[14]Department of Biology Faculty of Natural Sciences, University of Haifa, Haifa, Israel
[15]Departments of Medicine and Genetics, Albert Einstein College of Medicine, Bronx, NY, USA
[16]Data Science Platform, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[17]Department of Quantitative Health Sciences, Cleveland Clinic Research, Cleveland, OH, USA
[18]Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
[19]NHLBI and Boston University's Framingham Heart Study, Framingham, MA, USA
[20]Department of Medicine, Boston University Chobanian & Avedisian School of Medicine, Boston, MA, USA
[21]Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

[22]Department of Internal Medicine, Max Rady College of Medicine, University of Manitoba, Winnipeg, Canada

[23]Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, Canada

[24]Anaesthesia and Perioperative Medicine, Division of Surgery and Interventional Science, University College London Hospitals NHS Foundation Trust, University College London, London, UK

[25]Department of Biostatistics and Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA

[26]The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[27]Department of Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC, USA

[28]British Heart Foundation Centre for Research Excellence, NIHR Leicester Biomedical Research Centre, Division of Cardiovascular Sciences, University of Leicester, Leicester, UK

[29]"Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

[30]Department of Medicine, Rutgers Robert Wood Johnson Medical School, Rutgers, The State University of New Jersey, New Brunswick, NJ, USA

[31]Department of Genetics and the Human Genetics Institute of New Jersey, School of Arts and Sciences, Rutgers, The State University of New Jersey, Piscataway, NJ, USA

[32]Meyerhoff Inflammatory Bowel Disease Center, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[33]Fulcrum Genomics, Boulder, CO, USA

[34]Stanley Center for Psychiatric Research, The Broad Intitute of MIT and Harvard, Cambridge MA, USA

[35]Department of Clinical Research and Leadership, George Washington University School of Medicine and Health Sciences, Washington, DC, USA

[36]Center for Research on Genomics and Global Health, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA

[37]Harvard School of Public Health, Boston, MA, USA

[38]Central American Population Center, San Pedro, Costa Rica

[39]Department of Epidemiology and Biostatistics, Imperial College London, London, UK

[40]Department of Cardiology, Ealing Hospital, NHS Trust, Southall, UK

[41]Imperial College, Healthcare NHS Trust Imperial College London, London, UK

[42]Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China

[43]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[44]Department of Medicine, Harvard Medical School, Boston, MA, USA

[45]Department of Medicine, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ, USA

[46]Northwestern University, Evanston, IL, USA

[47]Departments of Cardiovascular, Medicine Cellular and Molecular Medicine Molecular Cardiology, Quantitative Health Sciences, Cleveland Clinic, Cleveland, OH, USA

[48]Department of Pediatrics, Boston Children's Hospital, Harvard Medical School, Boston, MA

USA

[49]McLean Hospital, Belmont, MA, USA

[50]Department of Psychiatry, Harvard Medical School, Boston, MA, USA

[51]Division of Medical Sciences, Harvard Medical School, Boston, MA, USA

[52]Genomics Platform, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[53]Department of Medicine, University of Mississippi Medical Center, Jackson, MI, USA

[54]Neuropsychiatric Genetics Research Group, Dept of Psychiatry and Trinity Translational Medicine Institute, Trinity College Dublin, Dublin, Ireland

[55]Centre for Neuropsychiatric Genetics & Genomics, Cardiff University School of Medicine, Cardiff, Wales

[56]Department of Epidemiology Colorado School of Public Health Aurora, CO, USA

[57]University of Cambridge, Cambridge, England

[58]Department of Medicine and Pharmacology, University of Illinois at Chicago, Chicago, IL, USA

[59]Broad Institute of MIT and Harvard, Cambridge, MA, USA

[60]Department of Medicine, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA

[61]Department of Human Genetics, School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA

[62]Clinical and Translational Science Institute, University of Pittsburgh, Pittsburgh, PA, USA

[63]Department of Life Sciences, College of Arts and Scienecs, Texas A&M University-San Antonio, San Antonio, TX, USA

[64]Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA

[65]Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, QC, Canada

[66]Cardiac Arrhythmia Service and Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA

[67]Cardiovascular Epidemiology and Genetics, Hospital del Mar Medical Research Institute (IMIM), Barcelona, Catalonia, Spain

[68]CIBER CV, Spain

[69]Departament of Medicine, Faculty of Medicine, University of Vic-Central University of Catalonia, Vic Catalonia, Spain

[70]Clalit Genomics Center, Ramat-Gan, Israel

[71]Institute for Cardiogenetics, University of Lübeck, Lübeck, Germany

[72]German Research Centre for Cardiovascular Research, Hamburg/Lübeck/Kiel, Lübeck, Germany

[73]University Heart Center Lübeck, Lübeck, Germany

[74]Estonian Genome Center, Institute of Genomics University of Tartu, Tartu, Estonia

[75]Richards Lab, Lady Davis Institute, Montreal, QC, Canada

[76]Victor Chang Cardiac Research Institute, Darlinghurst, NSW, Australia

[77]Faculty of Medicine and Health, UNSW Sydney, Kensington, NSW, Australia

[78]Cardiology Department, St Vincent's Hospital, Darlinghurst, NSW, Australia

[79]Mayo Clinic, Arizona, USA

[80]Broad Genomics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[81]Cardiovascular Discovery Group, Kolling Institute of Medical Research, University of Sydney, Australia

[82]Department of Cardiology, Royal North Shore Hospital, Australia

[83]Faculty of Medicine and Health, University of Sydney, Australia

[84]Diabetes Unit (Department of Medicine) and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[85]Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[86]Institute of Clinical Molecular Biology, Christian-Albrechts-University of Kiel, Kiel, Germany

[87]University Hospital Schleswig-Holstein, Kiel, Germany

[88]European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, UK

[89]Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

[90]Helsinki University and Helsinki University Hospital Clinic of Gastroenterology, Helsinki, Finland

[91]Helsinki University and Helsinki University Hospital, Helsinki, Finland

[92]Abdominal Center, Helsinki, Finland

[93]Unit of Animal Genomics, GIGA & Faculty of Veterinary Medicine, University of Liège, Liège, Belgium

[94]Bioinformatics Program MGH Cancer Center and Department of Pathology, Boston, MA, USA

[95]Cancer Genome Computational Analysis, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[96]Department of Psychiatry and Behavioral Sciences, Boston Children's Hospitaland Harvard Medical School, Boston, MA, USA

[97]Harvard Medical School Teaching Hospital, Boston, MA, USA

[98]Department of Endocrinology and Metabolism, Hadassah Medical Center and Faculty of Medicine, Hebrew University of Jerusalem, Israel

[99]Department of Psychiatry and Behavioral Sciences, SUNY Upstate Medical University, Syracuse, NY, USA

[100]Department of PSychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[101]Centro de Investigacion en Salud Poblacional, Instituto Nacional de Salud Publica, Mexico

[102]Lund University Sweden, Sweden

[103]Institute for Molecular Medicine Finland, (FIMM) HiLIFE University of Helsinki, Helsinki, Finland

[104]Science for Life Laboratory, Department of Gene Technology, KTH Royal Institute of Technology, Stockholm, Sweden

[105]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA

[106]Center for Genetic Epidemiology, Department of Population and Public Health Sciences, University of Southern California, Los Angeles, CA, USA

[107]Human Genetics Center, University of Texas Health Science Center at Houston, Houston, TX, USA

[108]Daiichi Sankyo, Basking Ridge, NJ, USA

[109]Department of Neurology Columbia University, New York City, NY, USA

[110]Institute of Genomic Medicine, Columbia University, New York City, NY, USA

[111]Institute of Biomedicine, University of Eastern Finland, Kuopio, Finland

[112]Department of Psychiatry, Helsinki University Central Hospital Lapinlahdentie, Helsinki, Finland

[113]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

[114]Icahn School of Medicine at Mount Sinai, New York, NY, USA

[115]Bonei Olam, Center for Rare Jewish Genetic Diseases, Brooklyn, NY, USA

[116]Department of Neurology, Helsinki University, Central Hospital, Helsinki, Finland

[117]Cardiovascular Disease Initiative and Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[118]Institute for Genomic Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[119]Division of Genome Science, Department of Precision Medicine, National Institute of Health, Republic of Korea

[120]MRC Centre for Neuropsychiatric Genetics & Genomics, Cardiff University School of Medicine, Cardiff, Wales

[121]Imperial College Healthcare NHS Trust, London, UK

[122]National Heart and Lung Institute Cardiovascular Sciences, Hammersmith Campus, Imperial College London, London, UK

[123]Department of Health THL-National Institute for Health and Welfare, Helsinki, Finland

[124]Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, New Haven, Connecticut

[125]Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New Haven, Connecticut

[126]Division of Pediatric Gastroenterology, Emory University School of Medicine, Atlanta, GA, USA

[127]Institute for Digestive Research and Department of Gastroenterology, Lithuanian University of Health Sciences, Kaunas, Lithuania

[128]Department of Internal Medicine, Seoul National University Hospital, Seoul, Republic of Korea

[129]The University of Eastern Finland, Institute of Clinical Medicine, Kuopio, Finland

[130]Kuopio University Hospital, Kuopio, Finland

[131]Department of Genetics, Yale School of Medicine, New Haven, CT, USA

[132]Department of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden

[133]Personalized Medicine, Mass General Brigham, Cambridge, MA, USA

[134]Department of Pathology, Brigham and Women's Hospital, Boston, MA, USA

[135]Department of Pathology, Harvard Medical School, Boston, MA, USA

[136]Department of Clinical Chemistry Fimlab Laboratories, Tampere University, Finland

[137]Finnish Cardiovascular Research Center-Tampere Faculty of Medicine and Health Technology, Tampere University, Finland

[138]Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[139]Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK

[140]Wellcome Trust Centre Human Genetics, University of Oxford, Oxford, UK

[141]Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge,

MA, USA

[142]The Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

[143]Department of Gastroenterology, University Hospital CHU of Liège, Liège, Belgium

[144]Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China

[145]Hong Kong Institute of Diabetes and Obesity, The Chinese University of Hong Kong, Hong Kong, China

[146]Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

[147]A. A. Martinos Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

[148]Division of Cardiology, University of California San Francisco, San Francisco, CA, USA

[149]Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain

[150]CIBERCV, Madrid, Spain

[151]TIMI Study Group, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, USA

[152]Department of Genetics, Harvard Medical School, Boston, MA, USA

[153]Oxford Centre for Diabetes, Endocrinology and Metabolism, University of Oxford, Churchill Hospital Old Road Headington, Oxford, OX, LJ, UK

[154]Welcome Centre for Human Genetics, University of Oxford, Oxford, OX, BN, UK

[155]Oxford NIHR Biomedical Research Centre, Oxford University Hospitals, NHS Foundation Trust, John Radcliffe Hospital, Oxford, OX, DU, UK

[156]John P. Hussman Institute for Human Genomics, Leonard M. Miller School of Medicine, University of Miami, Miami, FL, USA

[157]The Dr. John T. Macdonald Foundation Department of Human Genetics, Leonard M. Miller School of Medicine, University of Miami, Miami, FL, USA

[158]F. Widjaja Foundation Inflammatory Bowel and Immunobiology Research Institute Cedars-Sinai Medical Center, Los Angeles, CA, USA

[159]Atherogenomics Laboratory University of Ottawa, Heart Institute, Ottawa, Canada

[160]Division of Psychiatry, University College London, London, UK

[161]Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA

[162]Department of Clinical Sciences University, Hospital Malmo Clinical Research Center, Lund University, Malmö, Sweden

[163]Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Estonia

[164]University of Arizona Health Science, Tuscon, AZ, USA

[165]University of Maryland School of Medicine, Baltimore, MD, USA

[166]The Population Health Research Institute, McMaster University and Hamilton Health Sciences, Hamilton, Ontario, Canada

[167]Division of Gastroenterology, Department of Medicine, McMaster University, Hamilton, Ontario, Canada

[168]Farncombe Family Digestive Health Research Institute, McMaster University, Hamilton, Ontario, Canada

[169]Department of Electrophysiology, Texas Cardiac Arrhythmia Institute, St. David's Medical

Center, Austin, Texas, USA

[170]Aging Research Center, Cinvestav Sede Sur, Center for Research and Advanced Studies of the National Polytechnic Institute, Mexico City, Mexico

[171]Computational Biology Division, Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town, South Africa

[172]Institute of Infectious Disease & Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

[173]Nutrition Research Division, International Centre for Diarrhoeal Disease Research, Bangladesh

[174]Texas Cardiac Arrhythmia Institute, St. David's Medical Center, Austin, TX;

[175]Department of Biomedicine and Prevention, Division of Cardiology, University of Tor Vergata, Rome, Italy

[176]MetroHealth Medical Center, Case Western Reserve University School of Medicine, Cleveland, OH, USA

[177]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[178]Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

[179]Kenya Medical Research Institute-Wellcome Trust Collaborative Programme, Kilifi, Kenya

[180]Dept of Psychiatry, University of Oxford, United Kingdom

[181]Lund University, Dept. Clinical Sciences, Skåne University Hospital, Malmö, Sweden

[182]Department of Genome Informatics, Graduate School of Medicine, the University of Tokyo, Tokyo, Japan

[183]Laboratory for Systems Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan

[184]Premium Research Institute for Human Metaverse Medicine (WPI-PRIMe), Osaka University, Japan

[185]Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles

[186]Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA

[187]Instituto Nacional de Medicina Genómica, (INMEGEN) Mexico City, Mexico

[188]Laboratory of Immunogenomics and Metabolic Diseases, INMEGEN,Mexico City, Mexico

[189]Department of Haematology, University of Cambridge, Cambridgeshire, United Kingdom of Great Britain and Northern Ireland

[190]Applied Biomechanics Department, Swansea University, Singleton Park, Swansea SA2 8PP, UK

[191]Medical Research Institute, Ninewells Hospital and Medical School University of Dundee, Dundee, UK

[192]Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Instituto de Salud Carlos III, Madrid, Spain

[193]Department of Child and Adolescent Psychiatry, Institute of Psychiatry and Mental Health, Madrid, Spain

[194]Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM, Madrid, Spain

[195]Department of Molecular Medicine and Biopharmaceutical Sciences, Graduate School of

Convergence Science and Technology, Seoul National University, Seoul, Republic of Korea

[196]Department of Psychiatry Keck School of Medicine at the University of Southern California, Los Angeles, CA, USA

[197]Ambry Genetics, Aliso Viejo, CA, USA

[198]Department of Pediatrics/Hematology-Oncology, Baylor College of Medicine, Houston, Texas, USA

[199]Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[200]Department of Clinical Genetics, Amsterdam Neuroscience, Vrije Universiteit Medical Center, Amsterdam, The Netherlands

[201]Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[202]Departments of Human Genetics, University of Utah, Salt Lake City, UT, USA

[203]Children's Hospital of Philadelphia, Philadelphia, PA, USA

[204]Division of Genetics and Epidemiology, Institute of Cancer Research, London, UK

[205]Department of Psychiatry, Psychosomatic Medicine and Psychotherapy; University Hospital Frankfurt - Goethe University, Frankfurt am Main, Germany

[206]University of Washington, Seattle, WA, USA

[207]Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[208]Medical Research Center, Oulu University Hospital, Oulu Finland

[209]Research Unit of Clinical Neuroscience Neurology University of Oulu, Oulu, Finland

[210]Department of Genome Sciences, University of Virginia, Charlottesville, VA, USA

[211]Department of Public Health Sciences, University of Virginia, Charlottesville, VA, USA

[212]Research Center Montreal Heart Institute, Montreal, Quebec, Canada

[213]Department of Medicine, Faculty of Medicine Université de Montréal, Québec, Canada

[214]Department of Public Health Faculty of Medicine, University of Helsinki, Helsinki, Finland

[215]Section of Cardiac Electrophysiology, Division of Cardiology, Department of Medicine, Western University, London, Ontario, Canada

[216]Population Health Research Institute, Hamilton Health Sciences, and McMaster University, Hamilton, Ontario, Canada

[217]Department of Medicine, Vanderbilt, University Medical Center, Nashville, TN, USA

[218]Departments of Pharmacology and Biomedical Informatics Vanderbilt, University Medical Center, Nashville, TN, USA

[219]The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA, USA

[220]Department of Neurology and Neurosurgery, Montreal Neurological Institute and Hospital, McGill University Health Center, Montreal, Canada

[221]Translational Sciences, Research & Development, Biogen Inc., Cambridge, MA, USA

[222]Department of Cardiovascular Sciences, University of Leicester, Leicester, UK

[223]NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester, UK

[224]Departments of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[225]Departments of Psychiatry, Johns Hopkins University School of Medicine, Baltimore, MD,

USA

[226]Departments of Biomedical Engineering, Johns Hopkins University School of Medicine, Baltimore, MD, USA

[227]Department of Cardiology, Deutsches Herzzentrum München, Technical University of Munich, DZHK Munich Heart Alliance, Germany

[228]Technische Universität München, Germany

[229]Institute of Genetic Epidemiology, Department of Genetics, Medical University of Innsbruck, 6020 Innsbruck, Austria

[230]Faculty of Medicine, University of Southampton, Southampton, SO16 6YD, UK

[231]Duke Molecular Physiology Institute, Durham, NC

[232]Division of Cardiology, Department of Medicine, Duke University School of Medicine, Durham, NC, USA

[233]Division of Cardiovascular Medicine, Nashville VA Medical Center, Vanderbilt University School of Medicine, Nashville, TN, USA

[234]Division of Endocrinology, National University Hospital, Singapore

[235]NUS Saw Swee Hock School of Public Health, Singapore

[236]Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA, USA

[237]Harvard Medical School, Boston, MA, USA

[238]The Wallenberg Laboratory/Department of Molecular and Clinical Medicine, Institute of Medicine, Gothenburg University

[239]Department of Cardiology, Wallenberg Center for Molecular Medicine and Lund University Diabetes Center, Clinical Sciences, Lund University and Skåne University Hospital, Lund, Sweden

[240]Department of Cardiology, Sahlgrenska University Hospital, Gothenburg, Sweden

[241]Psychiatric and Neurodevelopmental Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[242]Institute of Clinical Medicine Neurology, University of Eastern Finad, Kuopio, Finland

[243]Sorbonne Université, INSERM, Centre de Recherche Saint-Antoine, CRSA, AP-HP, Saint Antoine Hospital, Gastroenterology department, F-75012 Paris, France

[244]INRA, UMR1319 Micalis, Jouy en Josas, France

[245]Paris Center for Microbiome Medicine, (PaCeMM) FHU, Paris, France

[246]Department of Twin Research and Genetic Epidemiology King's College London, London, UK

[247]Institute of Medical Sciences, University of Aberdeen, Aberdeen, Scotland, UK

[248]Department of Medicine, Washington University School of Medicine, Saint Louis, MO, USA

[249]Department of Genetics, Washington University School of Medicine, Saint Louis, MO, USA

[250]The McDonnell Genome Institute at Washington University, Saint Louis, MO, USA

[251]Departments of Genetics and Psychiatry, University of North Carolina, Chapel Hill, NC, USA

[252]National Institute for Health and Welfare, Helsinki, Finland

[253]Saw Swee Hock School of Public Health National University of Singapore, National University Health System, Singapore

[254]Life Sciences Institute, National University of Singapore, Singapore

[255]Department of Statistics and Applied Probability, National University of Singapore, Singapore

[256]Center for Behavioral Genomics, Department of Psychiatry, University of California, San Diego, CA, USA

[257]Institute of Genomic Medicine, University of California San Diego, San Diego, CA, USA

[258]Endocrinology, Abdominal Center, Helsinki University Hospital, Helsinki, Finland

[259]Institute of Genetics, Folkhalsan Research Center, Helsinki, Finland

[260]Juliet Keidan Institute of Pediatric Gastroenterology, Eisenberg R&D Authority, Shaare Zedek Medical Center, The Hebrew University of Jerusalem, Jerusalem, Israel

[261]Instituto de Investigaciones Biomédicas, UNAM, Mexico City, Mexico

[262]Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico

[263]Department of Public Health Faculty of Medicine University of Helsinki, Helsinki, Finland

[264]Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, CA, USA

[265]Hospital Universitari Institut Pere Mata, Reus, Tarragona, Spain

[266]Institut de Recerca IRB CatSud (formerly i-CERCA), Reus, Tarragona, Spain

[267]Centro de investigación biomédica en red- CIBERSAM, Madrid, Spain

[268]Department of Statistical Genetics, Osaka University Graduate School of Medicine, Suita, Japan

[269]National Heart and Lung Institute, Imperial College London, London, UK

[270]MRC Laboratory of Medical Sciences, Imperial College London, London, UK

[271]Radcliffe Department of Medicine, University of Oxford, Oxford, UK

[272]Department of Gastroenterology and Hepatology, University of Groningen and University Medical Center Groningen, Groningen, Netherlands

[273]Big Data Institute, University of Oxford, UK

[274]Wellcome Centre for Human Genetics, University of Oxford, UK

[275]Division of Cardiology, Beth Israel Deaconess Medical Center, Boston, MA USA

[276]Program in Infectious Disease and Microbiome, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[277]Center for Computational and Integrative Biology, Massachusetts General Hospital, Boston, MA, USA

[278]Stanley Division of Developmental Neurovirology, Department of Pediatrics, Johns Hopkins School of Medicine, Baltimore, MD, USA

Authors received funding as follows:

Elizabeth G. Atkinson: R01 HG012869

Emelia J. Benjamin: R01HL092577; American Heart Association AF AHA_18SFRN34110082

Matthew J. Bown: British Heart Foundation awards CS/14/2/30841 and RG/18/10/33842

Steven Brant: National Institutes of Health DK062431

Lea A. Chen: NIH, New York Crohn's Foundation, Crohn's and Colitis Foundation

Richard H. Duerr: U01 DK062420

Ravindranath Duggirala: U01 DK085524 National Institute for Diabetes and Digestive and Kidney Diseases (NIDDK)

Josée Dupuis: National Institute for Diabetes and Digestive and Kidney Diseases (NIDDK) R DK

Roberto Elosua: Agència de Gestió d'Ajuts Universitaris i de Recerca: 2021 SGR 00144

Jeanette Erdmann: VIAgenomics, Leducq network PlaqOmics, Deutsche Forschungsgemeinschaft Cluster of Excellence "Precision Medicine in Chronic Inflammation" (EXC2167);

Martti Färkkilä: State funding for university level health research

Laura D. Gauthier: Intel, Illumina
Benjamin Glaser: 5U01 DK085584
Stephen J. Glatt: U.S. NIMH Grant R MH
Leif Groop: The Academy of Finland and University of Helsinki: Center of Excellence for Complex Disease Genetics (grant number 312063 and 336822), Sigrid Jusélius Foundation; IMI 2 (grant No 115974 and 15881)
Christopher Haiman: U01CA164973
Mikko Hiltunen: Academy of Finland (grant 338182, 353053), Sigrid Jusélius Foundation, the Strategic Neuroscience Funding of the University of Eastern Finland
Chaim Jalas: Bonei Olam
Mikko Kallela: Grants from State funding for university level health research and from Department of Neurology, Helsinki University, Central Hospital; Grant from Maire Taponen Foundation
Jaakko Kaprio: Academy of Finland (grant 352792)
Mikael Landén: Swedish Medical Research Council (2022-00642)
Terho Lehtimäki: Academy of Finland (grant 356405), VTR Pirha
Ruth J.F. Loos: Novo Nordisk Foundation (NNF18CC0034900, NNF20OC0059313); NIH (R01DK110113; R01DK124097)
Ronald C.W. Ma: Research Grants Council of the Hong Kong Special Administrative Region (CU R4012-18), Research Grants; Council Theme-based Research Scheme (T12-402/13N), University Grants Committee Research Grants Matching Scheme
Jaume Marrugat: Agència de Gestió d'Ajuts Universitaris i de Recerca: 2021 SGR 00144
Jacob L. McCauley: National Institute of Diabetes and Digestive and Kidney Disease Grant R01DK104844
Michael C. O'Donovan: Medical Research Council UK: Centre Grant No. MR/L010305/1, Program Grant No. G0800509
Yukinori Okada: JSPS KAKENHI (25H01057), and AMED (JP24km0405217, JP24ek0109594, JP24ek0410113, JP24kk0305022, JP223fa627001, JP223fa627002, JP223fa627010, JP223fa627011, JP22zf0127008, JP24tm0524002, JP24wm0625504, JP24gm1810011), JST Moonshot R&D (JPMJMS2021, JPMJMS2024), Takeda Science Foundation, Ono Pharmaceutical Foundation for Oncology, Immunology, and Neurology, Bioinformatics Initiative of Osaka University Graduate School of Medicine, Institute for Open and Transdisciplinary Research Initiatives, Center for Infectious Disease Education and Research (CiDER), and Center for Advanced Modality and DDS (CAMaD), Osaka University, RIKEN TRIP initiative (AGIS)
Michael J. Owen: Medical Research Council UK: Centre Grant No. MR/L010305/1, Program Grant No. G0800509
Aarno Palotie: the Academy of Finland Center of Excellence for Complex Disease Genetics (grant numbers 312074 and 336824) and Sigrid Jusélius Foundation
John D. Rioux: National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK; DK062432), from the Canadian Institutes of Health (CIHR GPG 102170), from Genome Canada/Génome Québec (GPH-129341), and a Canada Research Chair (#230625)
Samuli Ripatti: the Academy of Finland Center of Excellence for Complex Disease Genetics (grant number) Sigrid Jusélius Foundation

Jerome I. Rotter: Trans-Omics in Precision Medicine (TOPMed) program was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for "NHLBI TOPMed: Multi-Ethnic Study of Atherosclerosis (MESA)" (phs001416.v1.p1) was performed at the Broad Institute of MIT and Harvard (3U54HG003067-13S1). Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who provided biological samples and data for MESA and TOPMed. JSK was supported by the Pulmonary Fibrosis Foundation Scholars Award and grant K23-HL-150301 from the NHLBI. MRA was supported by grant K23-HL-150280, AJP was supported by grant K23-HL-140199, and AM was supported by R01-HL131565 from the NHLBI. EJB was supported by grant K23-AR-075112 from the National Institute of Arthritis and Musculoskeletal and Skin Diseases.The MESA project is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420. Also supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center
Kaitlin E. Samocha: R01HG012867
Jeremiah Scharf: NIH Grants U01 NS40024, K02 NS085048, NS102371
Eleanor G. Seaby: Kerkut Charitable Trust, Foulkes Fellowship, University of Southampton Presidential Scholarship
Edwin K. Silverman: NIH Grants U01 HL089856 and U01 HL089897
J. Gustav Smith: The Swedish Heart-Lung Foundation (2022-0344, 2022-0345), the Swedish Research Council (2021-02273), the European Research Council (ERC-STG-2015-679242), Gothenburg University, Skåne University Hospital, governmental funding of clinical research within the Swedish National Health Service, a generous donation from the Knut and Alice Wallenberg foundation to the Wallenberg Center for Molecular Medicine in Lund, and funding from the Swedish Research Council (Linnaeus grant Dnr 349-2006-237, Strategic Research Area Exodiab Dnr 2009-1039) and Swedish Foundation for Strategic Research (Dnr IRC15-0067) to the Lund University Diabetes Center
Harry Sokol: AgroParisTech, Jouy en Josas, France
Nathan O. Stitziel: National Human Genome Reserach Institute Grant UM1HG008853
Kent D. Taylor: Trans-Omics in Precision Medicine (TOPMed) program was supported by the National Heart, Lung and Blood Institute (NHLBI). WGS for "NHLBI TOPMed: Multi-Ethnic Study of Atherosclerosis (MESA)" (phs001416.v1.p1) was performed at the Broad Institute of MIT and Harvard (3U54HG003067-13S1). Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed

Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I). We gratefully acknowledge the studies and participants who provided biological samples and data for MESA and TOPMed. JSK was supported by the Pulmonary Fibrosis Foundation Scholars Award and grant K23-HL-150301 from the NHLBI. MRA was supported by grant K23-HL-150280, AJP was supported by grant K23-HL-140199, and AM was supported by R01-HL131565 from the NHLBI. EJB was supported by grant K23-AR-075112 from the National Institute of Arthritis and Musculoskeletal and Skin Diseases.The MESA project is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts 75N92020D00001, HHSN268201500003I, N01-HC-95159, 75N92020D00005, N01-HC-95160, 75N92020D00002, N01-HC-95161, 75N92020D00003, N01-HC-95162, 75N92020D00006, N01-HC-95163, 75N92020D00004, N01-HC-95164, 75N92020D00007, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, and UL1-TR-001420. Also supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center Tiinamaija Tuomi: The Academy of Finland and University of Helsinki: Center of Excellence for Complex Disease Genetics (grant number 312072 and 336826), Folkhalsan Research Foundation, Helsinki University Hospital, Ollqvist Foundation, Liv och Halsa foundation; NovoNordisk Foundation
Teresa Tusie-Luna: CONACyT Project 312688
James S. Ware: JSW is supported by Medical Research Council (UK), Sir Jules Thorn Charitable Trust [21JTA], British Heart Foundation [RE/24/130023], and the NIHR Imperial Biomedical Research Centre
Rinse K. Weersma: The Lifelines Biobank initiative has been made possible by subsidy from the Dutch Ministry of Health Welfare and Sport the Dutch Ministry of Economic Affairs the University Medical Centre Groningen (UMCG the Netherlands) the University of Groningen and the Northern Provinces of the Netherlands

Conflicts of interest are as follows:
Mikko Kallela: No related COI
Eimear E. Kenny: EEK has received personal fees from Regeneron Pharmaceuticals, 23&Me, Allelica, and Illumina; has received research funding from Allelica; and serves on the advisory boards for Encompass Biosciences, Foresite Labs, and Galateo Bio
Mikael Landén: M.L. has received lecture honoraria from Lundbeck pharmaceutical.
Ruth J.F. Loos: R.J.F.L has received consultancy and speaker fees from Novo Nordisk and Eli Lilly and Company
Ronald C.W. Ma: No related COI
Edwin K. Silverman: Research grants from GSK and Bayer
James S. Ware: JSW has received research support from Bristol Myers Squibb, has acted as a paid advisor to MyoKardia, Pfizer, Foresite Labs, Health Lumen, Tenaya Therapeutics, and Solid Biosciences, and is a founder with equity in Saturnus Bio.