RNA tertiary structure and conformational dynamics revealed by BASH MaP

Maxim Oleynikov[1], Samie R. Jaffrey[1*]

[1]Department of Pharmacology, Weill Medical College, Cornell University, New York, NY, USA.

*e-mail: srj2003@med.cornell.edu

ABSTRACT

The functional effects of an RNA can arise from complex three-dimensional folds known as tertiary structures. However, predicting the tertiary structure of an RNA and whether an RNA adopts distinct tertiary conformations remains challenging. To address this, we developed BASH MaP, a single-molecule dimethyl sulfate (DMS) footprinting method and DAGGER, a computational pipeline, to identify alternative tertiary structures adopted by different molecules of RNA. BASH MaP utilizes potassium borohydride to reveal the chemical accessibility of the N7 position of guanosine, a key mediator of tertiary structures. We used BASH MaP to identify diverse conformational states and dynamics of RNA G-quadruplexes, an important RNA tertiary motif, in vitro and in cells. BASH MaP and DAGGER analysis of the fluorogenic aptamer Spinach reveals that it adopts alternative tertiary conformations which determine its fluorescence states. BASH MaP thus provides an approach for structural analysis of RNA by revealing previously undetectable tertiary structures.

INTRODUCTION

Chemical probing reagents can reveal extensive information about RNA structure. Typically,

folded RNAs are incubated with chemical probing agents that modify accessible nucleotides.

Atoms that are involved in hydrogen bonding are protected from modification, while unbonded

surface accessible atoms are modified. The chemical probing can also be performed with RNA

that is systematically mutagenized to predict the role of each nucleotide in the overall RNA

structure. In principle, the pattern of modifications can reveal the diverse set of hydrogen-

bonding interactions within an RNA, allowing RNA structures to be predicted.

However, not all modifications that are introduced by a chemical probing agent into RNA can be

detected. For example, one of the most widely used RNA structure-probing reagents, dimethyl

sulfate (DMS), primarily forms three methyl modifications on mRNA: N^1-methyladenosine (m^{1}A),

N^3-methylcytidine (m^{3}C), N^7-methylguanosine (m^{7}G)[1, 2]. Among these, m^{1}A and m^{3}C contain

methyl modifications located on the Watson-Crick face and therefore can be readily detected

due to their ability to impair reverse transcription and thus induce misincorporations in cDNA[3-6].

In contrast, m^{7}G has minimal effects on reverse transcription. Since m^{7}G contains a methyl

modification on the Hoogsteen face, rather than the Watson-Crick face of guanosine, it does not

efficiently perturb reverse transcription. As a result, the accessibility of the N7 position of G in

RNA cannot be readily assessed.

The ability to detect methyl modifications on the Watson-Crick face allows structure-probing

methods to accurately predict RNA secondary structure. RNA secondary structure is mediated

by Watson-Crick base pairs and includes the location of helices and loop regions in an RNA

structure. A's and C's are protected from methylation when they are engaged in Watson-Crick

base pairs, and are readily methylated when they are in single-stranded regions. The resulting

m^{1}A and m^{3}C residues are the primary nucleotides detected in DMS-based RNA structure

mapping methods[7-9].

In contrast to the Watson-Crick face, interactions that occur with the Hoogsteen face have

important roles in mediating RNA tertiary structures. RNA tertiary structure includes long-

distance interactions that coordinate the overall topology of an RNA as well as other structurally

complex non-helical RNA structures. These include G-quadruplexes, which rely on hydrogen

bonding interactions of the N7 position of G (N7G)[10]. G-quadruplexes comprise stacks of two or

more G-tetrads, which are square planar arrangements of four guanine bases that interact

through hydrogen bonds around a central potassium ion[10]. G-quadruplexes are physiologically

important RNA structures associated with translation, splicing, phase separation, and small

molecule binding[11-16]. G-quadruplex folding also appears to be dynamically regulated by

specific helicases[17,18]. However, the inability to probe the accessibility of N7G in RNA has

limited DMS structure mapping experiments from resolving RNA structures such as G-

quadruplexes which depend on N7G interactions.

Here we describe BASH MaP (Borohydride Assisted Structure determination of N7G Hoogsteen

interactions through Misincorporation Profiling), a method for determining RNA tertiary structure

by measuring global patterns of N7G accessibility in RNA. BASH MaP detects DMS-induced

modifications on both the Watson-Crick and Hoogsteen faces by using potassium borohydride

to convert m^7G to an abasic site. The resulting abasic site is detected by a specific signature

misincorporation during reverse transcription. We show that measurement of N7G accessibility

reveals broad insights into RNA tertiary structure. To reveal unexpected conformations states

and dynamics of RNA G-quadruplexes in vitro and in cells, we developed DAGGER

(Deconvolution Assisted by N7G Gaps Enabled by Reduction and depurination). DAGGER is a

computational pipeline that incorporates BASH MaP N7G accessibility data to model RNA

structure and identify alternative conformations. Using BASH MaP and DAGGER, we

demonstrate that the non-fluorescent conformation of the fluorogenic aptamer Spinach is

marked by an alternative tertiary structure that disrupts the folding of Spinach's G-quadruplex.

We show that the tertiary structure information afforded by BASH MaP allows substantially

improved RNA structure prediction, especially for RNAs with G-quadruplexes. Overall, BASH

MaP combines structural information from both the Watson-Crick and Hoogsteen face to reveal

RNA tertiary structures and conformational dynamics.

RESULTS

BASH MaP reveals DMS-induced m^7G sites in RNA

DMS MaP is a popular method for determining RNA structure that primarily reveals the location

of RNA helices. However, many tertiary RNA structures cannot be determined using DMS MaP

because they often rely on hydrogen bonding with the N7 position of G (N7G)[19]. In principle, the

accessibility of N7G can be readily assessed because the N7 position of G is the most

nucleophilic site in RNA and therefore the most reactive atom in RNA for DMS methylation[1].

However, the methyl moiety in m^7G is not on the Watson-Crick face and therefore does not

readily induce misincorporations during reverse transcription[20].

We therefore sought to make m^7G detectable during reverse transcription. Previous studies

showed that m^7G can be selectively converted to an abasic site, which is an efficient inducer of

misincorporations[21]. In this reaction, m^7G is reduced with potassium borohydride[22] and then

converted to an abasic site by heating in an acidic buffer[23, 24] (**Fig. 1a**). Thus, m^7G sites in

DMS-treated RNA can be converted to abasic sites, which readily induces misincorporations

during reverse transcription[22, 23]. Here we refer to BASH MaP as the treatment of RNA with

DMS, conversion of m^7G to abasic sites, and finally reverse transcription to generate

misincorporations at m^1A, m^3C, and m^7G (**Fig. 1b**).

We first characterized the misincorporations induced by m^7G-derived abasic sites. To test this,

we used the 18S ribosomal RNA (rRNA) from HeLa cells, which contains a stoichiometric m^7G

at position 1638[25-27]. We incubated total HeLa RNA, which contains 18S rRNA, with potassium

borohydride (800 mM) for 30 min to 4 h and then subsequently heated the RNA at 45°C in a pH

2.9 buffer for 4 h to induce depurination. The BASH-MaP-treated HeLa RNA was then reverse

transcribed by two commonly used reverse transcriptases in DMS MaP, SuperScript II and

Marathon RT, and the 18S RNA was sequenced.

A plot of misincorporation rate versus reduction duration of DMS treatment showed a ~70%

misincorporation rate at G1638 after 4 h of reduction and depurination (**Fig. 1c**). This high

misincorporation rate is consistent with essentially complete depurination of the m^7G based on

misincorporation rates seen with synthetic RNA oligos containing abasic sites[22]. The

misincorporations were primarily G→T and G deletions when using SuperScript II (**Fig. 1d**).

Similar results were seen with Marathon RT, with fewer deletions (**Supplementary Data Fig.**

1a). Together, these data demonstrate that m^7G can be efficiently converted to an abasic site

and then detected by a misincorporation signature using standard reverse transcription

conditions used for DMS MaP.

We next asked whether BASH MaP detects m^7G generated by DMS treatment of RNA. For

these experiments, we chose the fluorogenic RNA aptamer Spinach[28]. The crystal structure of

Spinach revealed a non-canonical G-quadruplex, G's in helices, as well as G's not involved in

structural interactions[29-31]. The N7G of each G in the G-quadruplexes participates in

Hoogsteen-face hydrogen-bonding interactions with an adjacent G and is thus expected to be

inaccessible to DMS.

We first modified Spinach with DMS (170 mM) for 8 min at 25°C. DMS-modified Spinach was

then split into two fractions and either subjected to BASH MaP or directly reverse transcribed as

in DMS MaP. In BASH MaP, misincorporations were detected at G's, which were

predominantly G→T and G deletions (**Fig. 1d**). The misincorporation signature for these G's

matches the misincorporations observed at the m^7G site in the 18S rRNA, thus demonstrating

that DMS generated m^7G in Spinach.

In contrast, in DMS MaP, misincorporations were rare at G's, and the few misincorporations

were predominantly G→A (**Fig. 1d**). These misincorporations may reflect mutations introduced

during RNA synthesis by T7 RNA polymerase, which are often G→A mutations[32]. Additionally,

the misincorporations may be due to an m^7G tautomeric state which induces G→A

misincorporations at very low rates during reverse transcription[20, 33]. Notably, BASH-MaP-

treated Spinach produced an average G misincorporation rate of 4.6% compared to 0.58% for

DMS MaP and 0.31% for untreated RNA (**Fig. 1e**). Altogether, these results demonstrate that

BASH MaP detects DMS-mediated methylation of N7G.

DMS is typically used at a final concentration between 10 mM and 170 mM, depending on the

level of methylation that is desired[26, 34]. We therefore modified Spinach with various

concentrations of DMS and performed BASH MaP. The misincorporation rates for all four

nucleotides were highly correlated between 85 mM and 170 mM DMS (R^2=0.9928) with a high

Pearson correlation coefficient over a range of DMS concentrations (**Fig. 1f** and

Supplementary Data Fig. 1b). Thus, BASH MaP produces highly reproducible measures of

DMS reactivity over a range of DMS concentrations.

We next asked whether the reduction and depurination steps in BASH MaP impaired detection

of other methylated nucleotides that are formed in single-stranded RNA. In addition to m^1A,

m^3C, and m^7G, DMS also forms m^1G and m^3U to a lesser extent[2, 20]. Of these five modified

nucleotides, only m^7G fails to efficiently induce misincorporations following reverse

transcription[19]. To test whether BASH MaP conditions affect the detection of m^1A, m^3C, or

m^3U, we modified Spinach with DMS (170 mM) for 8 min at 25°C in bicine buffer at pH 8.3,

which increases the formation of m^3U[2, 35]. DMS-modified Spinach was then split into two

fractions and either subjected to BASH MaP or directly reverse transcribed as in DMS MaP.

Notably, the misincorporation rate of A, C, and U bases remained correlated between the two

samples (**Fig. 1g** and **Supplementary Data Fig. 1b**). We could not measure m^1G in BASH

MaP as it is obscured by the high rate of m^7G formation (**Supplementary Data Fig. 1b**). As

expected, A, C, and U bases with high misincorporation rates predominantly mapped to single

stranded regions (**Fig. 1h-j**). Together, these results demonstrate that the reduction and

depurination steps of BASH MaP do not compromise the detection of methylated A, C, and U.

BASH MaP identifies guanosines that form G-quadruplexes

We next wanted to determine the misincorporation rate of G's located in different structural

contexts. Spinach contains G's involved in Hoogsteen interactions involving the N7G, as well

as G•C, G•U, and G•A base pairs. Spinach also displays a complex tertiary fold involving a G

containing a hydrogen-bonded N7 in a mixed tetrad beneath the two G-tetrads[29, 30]

(**Supplementary Data Fig. 1C**).

We performed BASH MaP on Spinach (1 µM) in solution with its cognate ligand DFHBI-1T (5

µM). Although m^7G in BASH MaP can exhibit either misincorporations or deletions, we only

quantified misincorporations. We chose to ignore all deletions because a deletion within a

stretch of two or more consecutive G's cannot be assigned to any specific G. We converted

misincorporation rates for each G into DMS reactivity values through a previously described

normalization scheme[20]. This approach allows DMS reactivities to be binned as either low,

medium, or high. The calculation of DMS reactivity values also enables direct comparison

between different nucleotides which display differences in baseline reactivity to DMS[20]. We

overlayed the DMS reactivity values on a secondary structure model of Spinach which revealed

a complex pattern of N7G reactivity to DMS (**Fig. 2a**).

We first examined G's that are not base-paired and are not hydrogen bonded at the N7 position

in the Spinach crystal structure. A total of three such G's are found in Spinach and an adjacent

stem loop, which is introduced for DNA sequencing[36] (**Supplementary Data Fig. 2a**). For the

following comparisons, we directly used misincorporation rates instead of normalized DMS

reactivities values. Each of these three G's displayed a high misincorporation rate (**Fig. 2b**).

Thus, as expected, unpaired G's with free N7 positions display high N7G reactivity to DMS.

In contrast, most G's in helices exhibited reduced misincorporation rates compared to G's in

single-stranded regions (**Fig. 2b**). Although helical G's do not have hydrogen-bonds at the N7

position, the reduced methylation of helical G's is consistent with low accessibility of the major

groove in duplex RNA[37]. However, we noticed two base-paired G's, G81 and G102, with very

high misincorporation rates (**Fig. 2a-b**). These G's are located at helix termini, a location

previously shown to exhibit enhanced major groove accessibility, and in locations of unusual

base stacking[37]. These results suggest that base-paired G's show moderately low N7 reactivity

to DMS except when located at the end of a helix.

We next asked whether N7-hydrogen bonds block methylation by DMS. We annotated all G's

with N7 positions engaged in a hydrogen bond, which included G's in G-quadruplexes and the

G involved in the mixed tetrad[29, 30]. We found that these G's showed the lowest

misincorporation rates compared to base-paired G's and single-stranded G's (**Fig. 2b**). Notably,

the G's constituting the RNA G-quadruplex of Spinach showed the lowest misincorporation rates

in Spinach (**Fig. 2c**). Thus, G's that are highly resistant to DMS methylation are likely engaged

in hydrogen-bonding interactions involving N7 and may reflect G's within G-quadruplexes.

To further test whether BASH MaP can identify G's in G-quadruplexes, we performed BASH

MaP on a polyUG repeat RNA which was recently shown to adopt an atypical G-quadruplex[38].

We choose a UG repeat length of 15 because we reasoned that this repeat length could

theoretically form four unique G-quadruplexes composed of 12 consecutive GU repeats (**Fig.**

2c). The formation of four unique G-quadruplexes would occur because of "register shifts," a

form of alternative RNA tertiary conformations (**Fig. 2c**). We reasoned that each unique G-

quadruplex register could be differentiated by the reactivity of three G's not engaged in a G-

quadruplex (**Fig. 2c**). Because these three excluded G's should lack N7 hydrogen bonding,

they therefore are expected to display higher DMS-mediated methylation than the G's engaged

in a G-quadruplex in all four registers.

BASH MaP analysis of the 15x polyUG repeat RNA revealed low misincorporation rates for the

nine G's expected to be hydrogen bonded in all four G-quadruplex registers (G26-G42) (**Fig.**

2e). G's that could participate in only one G-quadruplex showed the highest misincorporation

rate (**Fig. 2f**). Interestingly, a plot of the average misincorporation rate for each register

revealed that the propensity to form each register was not equal (**Supplementary Data Fig.**

2b). Instead, the G-quadruplex in register 2, which starts at G22 and ends at G44, displayed

the lowest average misincorporation rate and therefore marks the most abundant register

(**Supplementary Data Fig. 2b**). Overall, these experiments demonstrate that BASH MaP

enables fine resolution of different types of G-quadruplex structures.

Although we focused on the N7 position of G to identify the location of G-quadruplexes, we also

considered the possibility that other positions within G could be useful for G-quadruplex

prediction, particularly the N1 position (N1G). N1G is hydrogen bonded in G-quadruplexes and

in standard Watson-Crick base pairs and is therefore expected to be protected from DMS

methylation. Therefore, N1G reactivity might reveal the location of G-quadruplexes.

Compared to the N7G, we found the reactivity of N1G is between 10 to 100-fold lower, even

under higher pH conditions which promote the formation of m^1G[2, 20] (**Supplementary Data Fig.**

2c). Nevertheless, with sufficient sequencing depth, the presence of m^1G can be detected by

signature G→C and G→U misincorporations[20]. Using this approach, we observed no

discrimination between base-paired G's and G's involved in a G-quadruplex (**Supplementary**

Data Fig. 2c). Thus, the reactivity values of N1G are insufficient to identify G's in a G-

quadruplex. Overall, these data suggest that N7G reactivity measured by BASH MaP uniquely

identifies G's that are involved in a G-quadruplex.

BASH MaP detects G-quadruplexes in cellular mRNAs

The above experiments examined RNA G-quadruplexes in vitro. We next asked whether BASH

MaP could identify G's involved in an RNA G-quadruplex in cells. Methods for mapping G-

quadruplexes in cells are limited. One method involves treating cells with DMS and then

utilizing the property of folded G-quadruplexes to induce reverse transcriptase stops (RT stops)

at their 3' ends only in potassium-rich buffers[18, 39]. Cells are first treated with DMS, which

modifies any N7G's that are not in a G-quadruplex, and then the RNA is harvested and refolded

in potassium or sodium buffers. If the G-quadruplex was folded in the cell, then its N7G's

should be protected from methylation and the G-quadruplex would readily reform when folded in

potassium buffer. If the G-quadruplex was unfolded, then the N7G's would become methylated

which prevents subsequent G-quadruplex refolding. The refolded G-quadruplexes are then

identified as potassium-dependent RT stops and represent G-quadruplexes in cells.

A second method is based on the propensity for terminal G-quadruplex G's to become

chemically acylated in cells by a class of RNA-modifying chemicals called SHAPE reagents[40].

The location of SHAPE reagent adducts in extracted cellular RNA is tested by reverse-

transcription stops in potassium free buffers. Together, these methods predict the 3' end of G-

quadruplexes in cells. However, these G-quadruplex mapping methods only infer the presence

of a G-quadruplex and cannot reveal multiple G-tetrads or the presence of atypical G-

quadruplexes[40].

Despite their low resolution, these methods have suggested that most G-quadruplexes in

mRNAs are globally unfolded in HEK293T and HeLa mammalian cell lines[18, 41]. Although most

G-quadruplexes are unfolded, a small subset of G-quadruplexes appeared to remain folded in

cells. One of these folded G-quadruplexes resides in the 3' UTR of the *AKT2* mRNA[18]

(**Supplementary Data Fig. 2f**). We therefore sought to use BASH MaP to validate the folding

status of the *AKT2* 3' UTR G-quadruplex.

To determine if the 3' UTR of the *AKT2* mRNA contains a G-quadruplex in cells, we treated SH-

SY5Y cells with DMS (170 mM) for 6 min at 37°C[20, 35]. We then performed BASH MaP on the

extracted total RNA. The putative folded G-quadruplex region of *AKT2* was PCR amplified and

sequenced to identify misincorporation rates at G's. Notably, the misincorporations at G's in

AKT2 reflect formation of m^7G since they exhibited the characteristic G→T and G deletion

misincorporation signature of m^7G in BASH MaP (**Supplementary Data Fig. 2e**).

Next, we asked whether G's which were previously identified as the 3' end of the *AKT2* folded

G-quadruplex displayed low DMS reactivity. We converted misincorporation rates into

normalized DMS reactivities and plotted the reactivity profile[20] (**Fig. 2g**). The in-cell G-

quadruplex folding data suggests that the *AKT2* 3' UTR G-quadruplex has multiple

conformations with four unique 3' ends identified by potassium-dependent RT stops (**Fig. 2g**

and **Supplementary Data Fig. 2e**). Within the putative G-quadruplex region, we were able to

distinguish several tracts of G's with very low DMS reactivity, reflecting N7 positions protected

from DMS (**Fig. 2g**). These protected G's mapped specifically to G's previously identified to be

the 3' end of the in-cell folded G-quadruplex (**Fig. 2g** and **Supplementary Data Fig. 2e**).

Together, these results suggest that BASH MaP can identify G's engaged in a G-quadruplex in

cells.

In principle, a G-quadruplex only comprises four tracts of G's. However, *AKT2* contains seven

tracts of three or more G's which show low reactivity. Therefore, the BASH MaP N7G

accessibility data is unclear about which tracts of G's are engaged in the G-quadruplex (**Fig.**

2g). However, as will be described below, specific G-quadruplex G's can be identified using a

method that involves assessing co-occurring methylation events. The *AKT2* mapped G-

quadruplex region contains several tracts of G's with high reactivity to DMS, which reflects

accessible N7Gs and suggests that these G tracts are not engaged in a G-quadruplex (**Fig. 2g**

and **Supplementary Data Fig. 2e**). The location of these highly reactive G tracts suggests that

the *AKT2* G-quadruplex may adopt an unusual topology.

The G-quadruplex core of Spinach is marked by co-occurring G – G misincorporations

In addition to providing information on nucleobase accessibility, DMS MaP can predict

nucleobase-nucleobase interactions due to the phenomenon of "RNA breathing"[7]. During RNA

breathing, a transient local unfolding event can lead to methylation of an otherwise inaccessible

nucleotide. The methylated nucleotide can no longer interact with its cognate nucleotide

partner, which leads to methylation of the nucleotide partner as well. These events produce

pairs of misincorporations that repeatedly co-occur on the same strand of RNA[2, 7, 35, 42]. By

using a statistical analysis of misincorporations co-occurring in individual sequencing reads, with

each read representing a unique RNA molecule [7], Watson-Crick base pairs have been identified

in cells[2]. We therefore wondered whether BASH MaP can identify the specific G's in G-

quadruplexes using a similar statistical analysis of co-occurring misincorporations between G's.

We first asked whether BASH MaP produced enough co-occurring misincorporations needed for

statistical analysis. Statistical analysis of co-occurring misincorporations requires two or more

misincorporations in a sequencing read, which in DMS MaP is limited to misincorporations

induced by m^1A and m^3C due to low rates of m^1G and m^3U formation[7]. Analysis of Spinach

BASH MaP data indicated significantly more misincorporations per RNA molecule than DMS

MaP (**Supplementary Data Fig. 3a**). The percentage of reads with two or more

misincorporations doubled after BASH MaP, and the average number of total misincorporations

per read increased from 1.35 to 2.9 (**Supplementary Data Fig. 3b-c**). These results suggest

that BASH MaP produces data which is ideal for the statistical analysis of co-occurring

misincorporations.

Statistical analysis of co-occurring misincorporations is performed with the program RING

Mapper which identifies pairs of nucleotides which display higher than expected rates of co-

misincorporations[2, 7]. RING Mapper utilizes a likelihood-ratio statistical test called a G-test to

calculate whether a chemical modification event at one location affects the probability of a

chemical modification at a second location. If a chemical modification at a position in an RNA

increases the probability of a chemical modification at a second location, then the two locations

display statistical dependence. In contrast, if a chemical modification at one position has little or

no effect on the probability of modification at a second location, then the two locations display

statistical independence. Nucleotide pairs with a high G-test statistic value indicate high

dependence and are thought to represent pairs of nucleotides which interact in the RNA

structure[2, 7].

We next asked whether G's in BASH-MaP-treated Spinach RNA produced patterns of co-

occurring misincorporations that cannot be seen in DMS MaP. To identify novel patterns of co-

occurring misincorporations in BASH MaP of Spinach, we calculated G-test statistic values with

RING Mapper and converted these to a normalized correlation strength value. We then

compared these values to annotated domains derived from the crystal structure of Spinach[30]

(**Fig. 3c**). A heatmap of normalized correlation strength for all possible combinations of base

pairs showed marked differences between BASH-MaP-treated Spinach and control DMS-MaP-

treated Spinach (**Fig. 3d-e**).

We next asked whether the patterns of co-occurring misincorporations produced by BASH MaP

represent structural interactions in Spinach which are invisible to DMS MaP. DMS-MaP-treated

Spinach displayed strong correlations between A's in the P2 domain and G-quadruplex G

residues but displayed no correlations between G-quadruplex G's (**Fig. 3d**). In contrast, BASH-

MaP treated Spinach produced G – G correlations between G's comprising the G-quadruplex

(**Fig. 3e**). These G – G correlations represent the hydrogen bonding of N7Gs engaged in a G-

quadruplex. We previously relied on measures of N7G reactivity to infer G-quadruplex G's (**Fig.

2**). Instead, the co-occurring misincorporations heatmap suggests that the G – G correlations in

BASH MaP can directly identify the specific G's engaged in a G-quadruplex in an unbiased

manner. Thus, G-quadruplex G's can be discovered using BASH MaP, in contrast to DMS MaP

and produce a specific signature in the RING MaP heatmap.

Since BASH MaP produced many G – G co-occurring misincorporations, we next wondered

whether network analysis could identify discrete groups of structurally linked G's in Spinach

such as G-quadruplex G's. We reasoned that since the G's in the Spinach G-quadruplex form a

hydrogen-bonding network, then the G-quadruplex G's would form a distinct cluster in a network

analysis of G – G co-occurring misincorporations (**Fig. 3c**). To visualize the relationships

between G – G co-occurring misincorporations, we first represented the collection of co-

occurring misincorporations between pairs of G's as a network where vertices represent unique

G's in Spinach and edges denote co-occurring misincorporations[43]. To remove low confidence

co-modified nucleotides, we performed Z-score normalization on the collection of G-test statistic

values. We then filtered the network by Z-scores and included only co-occurring

misincorporations with a Z-score above a certain threshold.

To identify discrete groups of structurally linked G's in Spinach, we performed a detailed visual

analysis of the BASH MaP G – G co-occurring misincorporations network. The BASH MaP

network revealed a substantially higher number of co-occurring misincorporations between G

nucleotides compared to DMS MaP (**Fig. 3f-g**). We implemented a Z-score cutoff of 2.0 which

resulted in the formation of two distinct clusters within the network (**Fig. 3g**). The smaller cluster

was composed of seven G nucleotides, with six of them forming part of the Spinach G-

quadruplex and mixed tetrad (**Fig. 3g**). Notably, three G-quadruplex G's: G50, G51, and G89,

were not represented in the network because they did not display co-occurring

misincorporations with other G's. The lack of co-occurring misincorporations suggests that

methylation of G50, G51, and G89 produces an RNA conformation which retains hydrogen

bonds at the N7G of other G-quadruplex G's. Instead, G97, which is base stacked below G95

of the mixed tetrad, was present in the second, smaller cluster (**Fig. 3g**). These results show

that network analysis of BASH MaP data enables visualization of structurally linked groups of

G's and revealed that G-quadruplex G's form a distinct cluster.

Network analysis of G – G co-occurring misincorporations also revealed a large cluster of G's

which are annotated as base paired (**Fig. 3g**). The network is evident as a diffuse

checkerboard pattern in the RING MaP heatmap for G nucleotides engaged in helical regions

outside of the G-quadruplex core (**Supplementary Data Fig. 3d**). These results reveal that G's

engaged in canonical base pairs contribute to the stability of nearby base-paired G's.

In some cases, base-paired G's displayed moderate levels of co-methylation with other base-

paired G's regardless of if the G's were in the same or different helix (**Supplementary Data Fig.**

3d). The mechanism for these co-occurring methylations is unclear but suggests that

methylation of even a single N7G within Spinach can lead to global destabilization of RNA

structure. Together, these results further show that BASH-MaP-treated RNA produces novel

patterns of co-occurring methylations which correspond to structurally linked G's.

We next asked if G – G co-occurring misincorporations and network analysis could more

confidently identify G-quadruplex G's in *AKT2*. We performed RING MaP analysis of *AKT2*

BASH MaP data and represented the G – G co-occurring misincorporations as a network

(**Supplementary Data Fig. 4a**). The network formed four main clusters, one of which contained

four connected and lowly reactive G's (**Supplementary Data Fig. 4b**). Interestingly, the co-

occurring misincorporation network data indicates an unusual G-quadruplex topology where at

least one conformation of the *AKT2* G-quadruplex involves the first two and last two G tracts of

the putative G-quadruplex region (**Supplementary Data Fig. 4c**). We used the program QGRS

mapper[44] to predict all possible three-tiered G-quadruplex conformations and identified 129

unique G-quadruplex conformers (**Supplementary Data Fig. 4d**). We then calculated the

average misincorporation rate of the G-quadruplex G's for each conformation and ordered the

predicted conformers by average misincorporation rate. Analysis of the ten G-quadruplex

conformers with the lowest average misincorporation rate revealed a common large central loop

element (**Supplementary Data Fig. 4d**). Overall, these results further support the formation of

a G-quadruplex in the *AKT2* 3' UTR in cells and suggests the *AKT2* 3' UTR G-quadruplex may

adopt an unusual topology with a long central loop.

Multiplexed mutagenesis and chemical probing reveal N7G and base stacking

interactions

A second way to identify interactions between nucleobases in RNA is to use the mutate-and-

map method[32, 45] (M2). In the mutate-and-map method, mutagenic PCR is used to create a pool

of RNAs with PCR-derived mutations at every position along the length of the target RNA. DMS

MaP is then performed on the pool of PCR-mutated RNAs (M2 DMS MaP), and the data is

demultiplexed to identify how a change in nucleotide identity at each position along an RNA

affects the global DMS reactivity of that RNA. When a PCR-derived mutation occurs at the

position of a base pairing partner to an A or C, the A or C can no longer base pair and becomes

accessible to DMS. M2 DMS MaP provides a highly comprehensive screen of how each

position in an RNA interacts with other nucleotides in an RNA. However, M2 DMS MaP

primarily detects Watson-Crick base pairs in helical regions in an RNA[34]. We reasoned that

mutate-and-map combined with BASH MaP (M2 BASH MaP) would enable a comprehensive

screen of global N7G interactions, thereby revealing G-quadruplex and other tertiary

interactions of N7G in RNA.

To identify N7G interactions that contribute to RNA structure, we performed M2 BASH MaP and

M2 DMS MaP and compared the resulting heatmaps to identify N7G-specific interactions. We

used mutagenic PCR followed by in vitro transcription to randomly introduce point mutations

within Spinach and the 15x polyUG RNA (**Supplementary Data Fig. 5a**). We then adapted the

M2 DMS MaP method to include the reduction and depurination steps of BASH MaP (M2 BASH

MaP) and created an analysis pipeline to generate the mutate-and-map heatmap.

To create the mutate-and-map heatmap, we only considered sequencing reads which covered

the entire length of the RNA. We then considered only a subset of sequencing reads which

contained a PCR-derived mutation at one location along the RNA. Then, we calculated the

misincorporation rate at all other positions of the RNA from the subset of sequencing reads. We

repeated the previously described data analysis process for each position along the RNA.

Finally, all the calculated misincorporation rate profiles were stacked vertically to create the

mutate-and-map heatmap.

To better visualize changes in nucleotide reactivity due to the presence of a mutation installed

during PCR, we analyzed each column of the heatmap separately by Z-score normalization.

Each row of the heatmap represents the misincorporation rate profile for RNAs with a PCR-

derived mutation at that indicated position. We converted the misincorporation rates in each

column to a Z-score. A Z-score normalization strategy helps to reveal at which location a PCR-

derived mutation increases chemical reactivity the most for a given base in RNA[34]. To restrict

our analysis to increases in chemical reactivity, we plotted only positive Z-scores (**Fig. 4a-b**).

We then compared normalized heatmaps of M2 DMS MaP with M2 BASH MaP.

M2 BASH MaP of Spinach RNA produced a large collection of unique mutate-and-map signals

(**Fig. 4b**). The most notable of these signals mapped specifically to the N7G-interactions of the

G-quadruplex in Spinach. These interactions were absent in the M2 DMS MaP heatmap (**Fig.**

4a). Comparison of mutate-and-map heatmaps suggests that M2 BASH MaP can uniquely

identify N7G interactions in RNA.

In mutate-and-map heatmaps, mutations installed during PCR typically increase the reactivity of

a single position, namely that location's base-paired partner[34]. This led us to ask whether

individual G-tetrads which stack to form a G-quadruplex could be determined from M2 BASH

MaP heatmaps in a manner analogous to base pair identification.

PCR-derived mutations at a G-quadruplex G could potentially increase the reactivity of only G's

in the same tetrad (**Fig. 4c**). However, a zoomed-in plot of the Spinach M2 BASH MaP

heatmap indicated that a mutation installed during PCR in any G-quadruplex G caused

increased N7G methylation in all G-quadruplex tetrad layers (**Fig. 4d**). A similar pattern was

seen with M2 BASH MaP analysis of 15x polyUG RNA (**Supplementary Data Fig. 5b**). These

results suggest that PCR-derived mutations at G bases in the Spinach and polyUG G-

quadruplexes cause large scale increases in N7G accessibility of all G-quadruplex G's. Large

scale increases in N7G accessibility due to PCR-derived mutations may reflect global

destabilization of the G-quadruplex which leads to unfolding. Therefore, M2 BASH MaP

revealed that G-quadruplex folding is easily perturbed by changing any G in the G-quadruplex.

In addition to revealing the G-quadruplex-forming residues, the M2 BASH MaP heatmap

revealed conformational alterations to helices after point mutations. Helices are typically

detected as a diagonal line in the mutate-and-map heatmap, caused by a PCR-induced

mutation at one position, and a DMS-induced increase in reactivity at the complimentary

nucleotide in the helix (**Fig. 4e**). However, helices in M2 BASH MaP appeared as thicker and

jagged diagonal lines as compared to the same helix in M2 DMS MaP (**Fig. 4f**). The additional

signals in the M2 DAGGGER MaP heatmap indicates that PCR-induced mutations in the

Spinach helices cause broader changes in DMS reactivity than to simply the complementary

nucleotide.

To better understand the origin of these differences, we first asked which pairs of nucleotides

displayed novel M2 BASH MaP signals in the P3 stem of Spinach. Enlargement and visual

inspection of the P3 stem in the M2 BASH MaP heatmap revealed that most of the novel signals

corresponded to increased reactivity to DMS for G residues in the P3 stem (**Fig. 4f**).

One explanation for these signals is a transient breathing of the helix leading to increased N7G

accessibility as previously mentioned for G – G co-occurring misincorporations in the RING MaP

heatmap. However, G – G co-occurring misincorporations in the RING MaP heatmap occurred

between G bases in the same and separate helices. In contrast, M2 BASH MaP signals

between pairs of G residues were restricted to the same contiguous helix which suggests a

different mechanism.

We additionally observed vertical lines in the P3 stem for the M2 BASH MaP heatmap (**Fig. 4f**).

PCR-derived mutations at different locations in the P3 stem induced increased reactivity to DMS

for the same G bases to create these vertical lines. PCR-derived mutations within the P3 stem

are expected to create bulges which disrupt helix base stacking[46]. The highly reactive N7G at

helix termini, such as at G81, are also located in regions where base stacking interactions are

disrupted (**Fig. 4g**). The vertical lines in the heatmap suggest that PCR-derived mutations

within helices induce local distortions of base-stacking interactions and consequently increased

N7G reactivity of nearby G's. Therefore, local distortions of base stacking caused by PCR-

derived mutations may explain the novel signals present in the Spinach M2 BASH MaP

heatmap. Together, these data further suggest that N7G reactivity is highly influenced by local

base stacking interactions.

Tertiary folding constraints improve RNA secondary structure modeling

Incorporation of chemical probing data greatly improves RNA secondary structure modeling for

most RNAs[47-49]. The secondary structure of RNA refers to the location of base pairs in RNA and

is typically used to represent RNA structure[19]. The locations of base pairs are usually predicted

by computational algorithms such as mFOLD[50]. To incorporate chemical probing data, chemical

reactivities are commonly converted to free energy folding constraints which are then used by

RNA folding software[47-49].

Notably, computational algorithms are poor at predicting tertiary structures in RNA, such as G-

quadruplexes[19, 51]. The original predicted secondary structure of Spinach[28, 52] was largely

incorrect since it did not anticipate a G-quadruplex at the core of Spinach[31, 32]. Addition of

constraints from DMS MaP data does not improve modeling of G-quadruplexes since DMS MaP

primarily identifies regions of canonical Watson-Crick base pairs[6]. We therefore sought to apply

BASH MaP to improve secondary structure predictions for RNA that contain a G-quadruplex.

To determine if BASH MaP can improve secondary structure prediction, we focused on

Spinach. We first generated a secondary structure model from the Spinach crystal structure

(PDB: 4TS2) as a benchmark for the comparison of future models (**Fig. 5a**). Importantly, we

only considered Watson-Crick interactions and non-complementary base pairs involving only

two bases including G•A, A•A, and U•U base pairs when generating the secondary structure

model (**Fig. 5a**). We included these non-complementary base pairs because secondary

structure modeling algorithms can model these interactions[53]. G-quadruplex, mixed tetrad, and

base-triple interactions are intentionally left out as these tertiary structure elements are not

predictable or able to be modeled by current secondary structure modeling algorithms[53, 54].

We next used a minimum free energy folding approach to generate a secondary structure model

for Spinach[55]. We utilized mFold, a minimum free energy folding algorithm which attempts to

find the base-pairing pattern with the lowest free energy for a given RNA sequence to predict

the secondary structure of Spinach from sequence alone[50] (**Fig. 5b**). The mFold secondary

structure model of Spinach correctly displays stems P1 and most of P3 but lacks stem P2 and

contains two stem loops which are not present in the crystallographic secondary structure (**Fig.**

5a-b). The mFold secondary structure model demonstrates that minimum free energy folding

algorithms fail to accurately model Spinach secondary structure from sequence alone.

We then asked whether incorporation of chemical probing data could improve the secondary

structure prediction for Spinach. The most common method for using chemical probing data to

improve secondary structure modeling is to convert chemical reactivities to free energy folding

constraints which are then used by a minimum free energy folding program[48]. These

constraints reward the algorithms for forming base pairs with RNA bases that display low

chemical reactivity and penalize the formation of base pairs at locations with high chemical

reactivity.

We first used DMS reactivities to generate free energy folding constraints for the folding

program RNAstructure[34, 54]. We then compared the resulting secondary structure model to the

mFold and crystallographic secondary structures[35, 54] (**Fig. 5a-b**). For DMS MaP experiments

we considered chemical reactivities at all four nucleotides to create free energy folding

constraints. For BASH MaP, we only considered chemical reactivities at A, C, and U bases to

create free energy folding constraints as we previously showed the some based-paired G's

displayed high N7G reactivity. If included, folding algorithms would incorrectly force base-paired

G's with high N7G reactivities to be single stranded. Therefore, we ignored all data at G's when

creating free energy folding constraints from BASH MaP data.

Incorporation of chemical reactivity data produced secondary structure models which retained

the correct features of the mFold secondary structure including the P1 and P3 domain (**Fig. 5b**).

However, the secondary structure models continued to lack a properly formed P2 domain and

continued to display two incorrect stem loops, although with less incorrect base pairs than the

mFold model (**Fig. 5b**). Together, these results show that free energy folding constraints from

chemical probing data improve but are insufficient to correctly model the secondary structure of

Spinach.

Next, we asked whether we could generate additional folding constraints from N7G reactivity

data. We noticed that all the previous secondary structure models displayed the same incorrect

stems (**Fig. 5b**). These incorrect stems included G-quadruplex G's that were mis-assigned and

forced into base pairs (**Fig. 5b**). The low N1G reactivity of the G-quadruplex G's in DMS MaP

was incorrectly interpreted by RNAstructure to indicate that G-quadruplex G's were engaged in

a base-pair interaction. In principle, very low N7G reactivity could be used to identify G-

quadruplex G's, which would then be restricted from being assigned as base paired.

We next implemented a two-step approach to confidently annotate G's engaged in a tertiary

interaction (**Fig. 5c**). First, we selected G's in the bottom quartile for N7G reactivity. Of these

G's, we then identified which G's displayed high rates of co-occurring misincorporations with

other lowly reactive G's. Together, these two steps annotate G's engaged in a tertiary

interaction and should thus be excluded from base pair assignments during secondary structure

modeling.

To incorporate N7G reactivity data, we developed a single-molecule RNA secondary structure

modeling pipeline called DAGGER (Deconvolution Assisted by N7G Gaps Enabled by

Reduction and depurination). DAGGER modifies the recently described DaVinci analysis

pipeline to incorporate N7G accessibility data derived from BASH MaP[56] (**Fig. 5c**). In contrast

to minimum free energy folding algorithms like RNAstructure, DAGGER uses a thermodynamic

independent RNA folding algorithm called CONTRAfold[53]. CONTRAfold utilizes a probabilistic

methodology for generating secondary structures through statistical learning on large RNA

secondary structure datasets[53]. A recent benchmark found that CONTRAfold outperformed

thermodynamic folding algorithms in secondary structure modeling accuracy[57].

DAGGER incorporates chemical reactivity data to constrain secondary structure modeling by

CONTRAfold. Like DaVinci, DAGGER uses locations of misincorporations in a sequencing read

to exclude those nucleotides from base-pair assignment by CONTRAfold[56]. First, each

sequencing read is converted to a string of ones and zeros called a bitvector which enables a

simple representation of misincorporation sites within a sequencing read. A zero value

indicates that nucleotide matches the reference sequence whereas a one indicates a

misincorporation site (**Fig. 5c**). DAGGER then folds each sequencing read as a unique

molecule of RNA. Locations with a bitvector value of one are forced to be single stranded

during secondary structure modeling by CONTRAfold[56]. Thus, DAGGER utilizes chemical

reactivity to constrain the secondary structure modeling of individual sequencing reads.

In contrast to chemical reactivity data on the Watson-Crick face, G's engaged in tertiary

interactions display low reactivity and mostly appear as zeros in the DAGGER bitvectors.

Consequently, CONTRAfold misinterprets G's engaged in tertiary interactions as engaged in

secondary structure interactions which leads to inaccurate structure modeling (**Fig. 5a-b**).

To prevent incorrect assignment of G's engaged in tertiary interactions, we created an additional

step in DAGGER to edit each bitvector before folding. We edited each bitvector such that all

G's engaged in tertiary interactions were represented by a one in the bitvector and would

therefore be excluded from base-pair assignment (**Fig. 5c**). We also set all G's not engaged in

tertiary interactions to a bitvector value of zero as we previously determined N7G

misincorporation status would incorrectly force base-paired G's to be single stranded (**Fig. 2b**).

Through N7G-directed editing of bitvector values, we incorporate tertiary-interaction constraints

into the DAGGER pipeline.

Even though G's engaged in tertiary interactions display low rates of misincorporations, in rare

cases, there will be a misincorporation and we wanted a strategy to analyze these events. In

principle, a misincorporation at a G engaged in a tertiary interaction may result from an

alternative conformation of the RNA in which the G is no longer engaged in a tertiary interaction.

Therefore, if a sequencing read contains a misincorporation at a G engaged in a tertiary

interaction, we allow all G's to be considered for base-pair assignment by CONTRAfold. As

such, our integration of N7G reactivity data into DAGGER allows us to model secondary

structures for single RNA molecules in multiple distinct tertiary structures. Together,

identification of G's engaged in tertiary interactions enables N7G reactivity data to be converted

into a folding constraint for secondary structure prediction by DAGGER.

We next asked whether N7G-derived tertiary constraints improved the modeling of Spinach

secondary structure. We implemented these tertiary-folding constraints through our structure

probing deconvolution pipeline DAGGER which is based on the DaVinci pipeline[56] (**Fig. 5c**).

Conventional analysis with DaVinci using either DMS MaP or BASH MaP Spinach datasets

produced consensus secondary structures nearly identical to models obtained via free energy

modeling approaches using RNAstructure (**Fig. 5b**). We then applied the DAGGER pipeline to

the 10,000 most modified sequencing reads in the Spinach BASH MaP dataset. Application of

DAGGER to the BASH MaP Spinach dataset produced a consensus secondary structure that

closely resembled the crystallographic secondary structure (**Fig. 5d**). The resulting secondary

structure model included a properly formed P1, P2, and P3 stem and was devoid of false helical

regions (**Fig. 5d**). We were unable to create tertiary constraints from DMS MaP since this

method does not differentiate between G-quadruplex G's and base-paired G's (**Supplementary**

Data Fig. 2c). These results demonstrate that tertiary-folding constraints from BASH MaP as

implemented through DAGGER enable accurate secondary structure modeling of Spinach.

BASH MaP and DAGGER identifies a misfolded state of Spinach

Spinach has been used as a model RNA for studying RNA G-quadruplex folding[29, 58]. The

fraction of fully folded Spinach transcripts was previously assessed using a fluorescence-based

assay where the fluorescence of a solution of Spinach was compared to a standard solution

containing a known amount of fully folded Spinach. Fluorescence-based assays showed that

~60% of Spinach was in a fluorescent form at 25°C, with the remainder thought to be in a

misfolded conformation[59]. Preventing the misfolded conformation would increase overall

Spinach fluorescence. Critically, it is unclear what, if any, structure is found in the misfolded

form of Spinach. We therefore sought to apply BASH MaP and DAGGER to understand the

conformation of Spinach misfolded states.

First, we sought to identify distinct conformations of Spinach using the RNA structure

deconvolution program DANCE[35]. DANCE uses a Bernoulli mixture model to identify mutually

exclusive patterns of misincorporations on single sequencing reads from DMS MaP

experiments. Sequencing reads that exhibit mutually exclusive patterns of misincorporations

are derived from different conformations of RNA[7]. We first used DANCE with the Spinach M2

DMS MaP dataset, which failed to identify more than one RNA conformation (data not shown).

In contrast, when we used DANCE with the Spinach M2 BASH MaP dataset, we readily

detected two states: State 1 with an abundance of 80.6% and State 2 with an abundance of

19.4% (**Fig. 6a**). To confirm that the two states identified by DANCE were not a result of PCR-

derived mutations, we repeated DANCE deconvolution of a Spinach BASH MaP dataset without

the mutate-and-map step, which therefore uses RNA lacking PCR-derived mutations. DANCE

deconvolution of this dataset also produced two states with abundances of 78.7% and 21.3%

which closely resembles the abundances seen in the DANCE deconvolution of the M2 BASH

MaP dataset (**Supplementary Data Fig. 7a**). Together, these data show that DANCE analysis

of BASH MaP data uniquely identifies multiple conformations of Spinach.

We next asked whether the folding abundances measured by DANCE agreed with

fluorescence-based measurements. The folding abundances identified by DANCE of ~80% are

slightly higher than the ~60% folding abundance previously measured for Spinach[59]. The slight

difference in measured folding abundance may reflect additional conformations in Spinach

which display similar chemical reactivity to the fluorescent-competent form of Spinach but are

nonetheless unable to activate its fluorophore DFHBI-1T. DANCE deconvolution requires at

least 10 nucleotides to display differential reactivities for deconvolution and would therefore be

unable to differentiate between two conformations of Spinach which differ by only a few bases[35].

Evidence for additional conformations within the DANCE identified states is indicated by

intermediate reactivity levels in single stranded regions which should either display low or high

reactivity[60] (**Fig. 6a**). Overall, DANCE deconvolution of Spinach closely approximates

fluorescence-based measurements of Spinach folding.

Next, we asked which regions of Spinach displayed different structure between the two states

revealed by DANCE. Analysis of normalized misincorporation rate differences between State 1

and State 2 revealed large changes in the P2 and G-quadruplex core domains and minimal

differences between stems P1, P3, and a linker region which was added to enable PCR

amplification of Spinach (**Fig. 6b**). The pattern of reactivity differences suggests that the

"misfolded" conformation of Spinach is folded and maintains key helical elements of the folded

form of Spinach but has an alternative conformation of the P2 and G-quadruplex core domains.

Next, we wanted to understand the alternative conformation of the G-quadruplex. We first

compared the reactivity to DMS for each G in both states (**Fig. 6c**). G-quadruplex G's in State 1

displayed strong protections from DMS methylation (**Fig. 6d**). In contrast, G-quadruplex G's in

State 2 were found to display reactivity to DMS like base paired G's (**Fig. 6e**). Stratification of

G-quadruplex G's by either population average, State 1, or State 2 revealed increased N7G

reactivity in State 2 (**Fig. 6f**). Analysis of single stranded A's in the G-quadruplex loops showed

an opposite trend with decreased reactivity to DMS in State 2 (**Fig. 6g**). Together, these results

suggest that the misfolded Spinach adopts an alternative conformation characterized by a

breakdown of N7G hydrogen bonding in the G-quadruplex core.

Next, we asked whether the misfolded Spinach contained alternative interactions of the G-

quadruplex loop residues. The crystal structure of Spinach shows that the loop A's make no

interactions within Spinach[29, 30]. Therefore, Spinach folding, and fluorescence is not expected to

depend on the identity of the loop nucleotides. Since the loop A's showed decreased reactivity

in State 2 and the G-quadruplex G's displayed similar reactivities to base-paired G's, we

reasoned that the G-quadruplex G's may misfold via interactions with loop A's to form non-

canonical G•A base pairs as seen in the P2 domain of Spinach (**Fig. 6e** and **Fig. 6g**).

If the misfolded Spinach contained non-canonical G•A base pairs between loop A's and G-

quadruplex G's then mutation of loop A's to C should increase the stability of these interactions

and therefore increase the fraction of misfolded Spinach transcripts. To increase the stability of

these interactions, we progressively converted loop A's to C's and measured RNA fluorescence

through an in-gel assay[61]. Surprisingly, mutation of two single stranded loop A's to C's resulted

in ~40% of the wildtype fluorescence (**Fig. 6h**). Mutation of three loop A's to C's resulted in

near complete loss of fluorescence (**Fig. 6h**). These results support a misfolded conformation

of Spinach where G-quadruplex G's form non-canonical G•A base pairs with loop A's.

To better understand if the misfolded Spinach contains an alternative conformation in the P2

domain, we next asked whether an orthogonal RNA structure deconvolution technique, our

previously developed DAGGER, produced clusters of RNA structure with alternative base

pairing in the P2 domain. We previously developed DAGGER for modeling the secondary

structure of Spinach through the incorporation of tertiary-folding constraints (see **Fig. 5c-d**).

DAGGER performs deconvolution after secondary structure modeling of individual RNA

molecules[56], which contrasts with DANCE deconvolution which occurs before secondary

structure modeling[35]. To perform RNA structure deconvolution, DAGGER first folds each

sequencing read utilizing misincorporations as folding constraints then represents each

secondary structure as a string of numbers[62]. Then, principal component analysis (PCA) is used

to reduce each string of numbers representing RNA structures down to two unique values and

plots these values as the first two principal components. From PCA plots, clustering methods

such as Kmeans clustering can be used to identify clusters of similar RNA structures[56].

We analyzed the M2 BASH MaP dataset with tertiary constraints as implemented through the

DAGGER pipeline and generated a plot of the first two principal components (**Fig. 6i**).

Interestingly, the DAGGER plot appeared to form two major clusters which agrees with the

number of clusters identified by DANCE. To identify the most representative structure for each

cluster, we performed Kmeans clustering with K equal to two. We then identified the

sequencing read which was closest to the center of Cluster 1 as the most representative

structure of Cluster 1. The same approach was used to identify the most represent structure of

Cluster 2. We then compared the most representative secondary structures of Cluster 1 and

Cluster 2. The key difference between the representative secondary structure of Cluster 1 and

the representative secondary structure of Cluster 2 involved alternative base-pair interactions in

the P2 domain (**Supplementary Data Fig. 6a-b).** The alternative base-pairing pattern of the P2

domain represents a register shift and is expected to prevent proper G-quadruplex folding.

Together, DAGGER analysis supports an alternative base-pairing pattern in the P2 domain of

Spinach's misfolded state.

We next asked whether Broccoli, a variant of Spinach identified through functional SELEX for

better intracellular folding, contained stabilizing mutations in the P2 domain[63]. A previous

alignment of the Broccoli sequence to Spinach suggested the two aptamers contain highly

conserved G-quadruplex domains[63]. However, sequence differences in Spinach which led to

the improved Broccoli aptamer are concentrated in the P2 domain (**Supplementary Data Fig.**

7b). In Spinach, a non-canonical A•A base pair just below the mixed tetrad was present as a

G•C pair in Broccoli. Additionally, another non-canonical U•U base pair in Spinach was present

as an A•U base pair in Broccoli. These sequence differences are expected to enhance folding

of the fluorescent conformation by stabilizing the P2 domain. Furthermore, these mutations

prevent the P2 domain from adopting the alternative Spinach base-pairing pattern seen in the

secondary structure of the DaVinci Cluster 2 by converting a U42•G97 base pair to the unstable

A42•G97 base pair (**Supplementary Data Fig. 7b**). Overall, these data provide a basis for the

improved fluorescence of Broccoli and suggest that the misfolded state of Spinach involves a

specific alternative base-pairing interaction of the P2 domain.

DISCUSSION

Deep sequencing of misincorporations induced by chemical probe-treated RNA is very useful

for determining RNA structures and for identifying different RNA conformational states[8].

Identification of different RNA conformational states relies on converting chemically modified

nucleotides into a misincorporation during cDNA synthesis by a reverse transcriptase. As a

result, only chemical adducts that are "seen" by reverse transcriptases can be detected by

cDNA sequencing. Adducts that occur on the Watson-Crick face perturb base pairing during

nucleotide selection for cDNA synthesis, and therefore result in cDNA misincorporations. In the

case of N7G, however, the methyl adduct is on the Hoogsteen face, rather than the Watson-

Crick face and is thus largely invisible to reverse transcriptases. Importantly, information on the

accessibility of N7G is needed to identify G-quadruplexes and any other tertiary interaction

involving the N7G position. To detect N7G interactions, we developed BASH MaP. BASH MaP

converts the otherwise nearly invisible m^7G into a highly detectable modified nucleotide by

selectively converting it to an abasic site. Using BASH MaP, we reveal the diversity of N7G

interactions in RNA structure and enable precise localization of G-quadruplexes in RNAs in vitro

and in cells. We then applied bioinformatic approaches to identify and measure changes in

N7G interactions for previously intractable RNAs. Thus, BASH MaP expands the structures that

can be detected using DMS-based structure probing methods.

G-quadruplexes have diverse roles in regulating RNA function but have been challenging to

identify and study in vitro and in cells[11-15, 41]. The lack of methods to specifically identify clusters

of the interconnected guanosine residues that constitute a G-quadruplex makes it difficult to

validate and establish the topology of G-quadruplexes in RNA. This limits the ability to establish

the physiologic roles and functions of G-quadruplexes. Current RNA structure mapping

methods cannot detect G-quadruplexes and other complex RNA structures because they

primarily detect the presence of Watson-Crick base pairs. A feature of BASH MaP is its ability

specifically identify G's involved in a G-quadruplex.

G-quadruplex G's can be identified based on their very low reactivity towards DMS, as

measured by a low misincorporation rate. However, G-quadruplex G's are more precisely

identified through co-occurring misincorporations in single reads where individual G's in a G-

quadruplex become methylated with DMS, which disrupts the G-quadruplex, resulting in

coordinated methylation of the otherwise non-reactive G's in the G-quadruplex. These clusters

of reactive G's will occur on single reads, reflecting individual strands of RNA. Notably, the

heatmaps used to visualize BASH MaP analyses of RNA structure produce a new box-like

pattern for G-quadruplexes not previously seen with DMS MaP heatmaps. The statistical co-

occurrence of misincorporations at G's in BASH MaP allows us to create a network of

structurally interconnected G's, which thus identifies G-quadruplexes.

The G's involved in a G-quadruplex can similarly be identified using the mutate-and-map

method, in which mutations are intentionally incorporated into an RNA using mutagenic PCR[32].

In this method, PCR-derived mutation of any G in the G-quadruplex enhances the reactivity of

all other G's in the G-quadruplex, as well as another structural feature that is dependent on G-

quadruplex folding, as seen in Spinach. Overall, the M2 BASH MaP method allows highly

precise identification of specific G's involved in G-quadruplex formation.

BASH MaP provides substantial improvements over previous methods for discovering G-

quadruplex G's in vitro. G-quadruplexes G's have been predominantly identified by performing

chemical or enzymatic probing of an RNA refolded in potassium-containing buffer versus

refolding in lithium- or sodium-containing buffer[65]. Since G-quadruplexes are destabilized by

lithium or sodium ions, refolding an RNA in lithium buffer or sodium buffers should lead to

reactivity changes around G-quadruplexes. Similarly, chemical, or enzymatic reactivity of RNA

synthesized with 7-deaza GTP, which is unable to form G-quadruplexes, can be compared with

the reactivity of RNA synthesized with GTP to identify regions where G-quadruplexes may

form[66]. However, these methods are limited to cell-free RNA and only infer locations of G-

quadruplexes. BASH MaP enables high throughput identification of G-quadruplexes G's in an

RNA of interest by directly detecting N7G accessibility.

BASH MaP also provides substantial improvements over previous methods for identifying G-

quadruplex G's in cells. Chemical probes such as kethoxal[67], which modify the Watson-Crick

face of G, lack the ability to discriminate between G's in helices and G-quadruplexes

(**Supplementary Data Fig. 2c**). Other assays rely on reverse transcriptase stalls at G-

quadruplexes, but these methods only indicate the 3' most guanine and do not definitively

identify a G-quadruplex as the source of the stall[18, 39, 41]. SHAPE reagents were found to

preferentially modify terminal G's in some G-quadruplexes; however, this property does not

have a clear mechanistic basis and does not unambiguously identify G-quadruplexes G's[40].

BASH MaP identifies G-quadruplexes G's in cells with RING MaP co-occurring misincorporation

data by directly revealing G-quadruplex guanosine hydrogen bonding networks.

An important advance in RNA structure mapping was the utilization of multiple modification

events on a single RNA molecule to detect the presence of different conformations of RNA in

solution[7, 35, 56, 60, 64]. The power of deconvolution methods increases with the number of co-

occurring misincorporations on individual reads by identifying additional mutually exclusive

patterns of nucleotide accessibility. Mutually exclusive patterns indicate different structures. In

BASH MaP, m^7G contributes to the misincorporations seen in a read. Since N7G is the most

reactive site to DMS, the detection of m^7G results in substantially more misincorporations per

read than in conventional DMS MaP. As a result, the ability to identify distinct conformational

states of RNA is substantially improved by detecting the m^7G formed in DMS experiments.

We used BASH MaP to understand the distinct conformations of Spinach. Spinach

fluorescence is limited since it exists in both a folded and a non-fluorescent unfolded

conformation[59]. We first developed a computational approach for utilizing BASH MaP data to

create new constraints for RNA folding. Our DAGGER pipeline uses a mixture of chemical

reactivity data and single-molecule co-occurring misincorporation data to identify G's likely to be

engaged in a tertiary interaction. DAGGER then restricts identified G's from forming canonical

base pairs during folding. The incorporation of tertiary folding constraints through DAGGER led

to a remarkably accurate prediction of the Spinach secondary structure (**Fig. 5d**).

Surprisingly, DANCE analysis of Spinach BASH MaP data revealed that the previously

described "misfolded" conformation was instead a highly specific folded conformation. In the

non-fluorescent conformation, the G-quadruplex G's only display a modest increase in reactivity,

and stems P1 and P3 remained intact (**Fig. 6b-c**). Instead, the G-quadruplex G's display N7

reactivity consistent with G's engaged in base-pair interactions (**Fig. 6e**). In addition, the hyper-

reactive G81 undergoes a dramatic decrease in reactivity in the non-fluorescent conformation

which indicates a change in local base stacking. The decrease of G81 reactivity suggests that

G81 is no longer located at the end of a helix in the nonfluorescent conformation (**Fig. 6c**).

Also, DAGGER analysis supports a reorganization the of P2 domain which is likely aided by the

adjacent and flexible J1/2 domain of Spinach (**Fig. 6h**). Together, these results suggest that the

non-fluorescent conformation of Spinach involves the G-quadruplex G's forming an extended P3

domain characterized by base-pairing with loop residues and a reorganization of the P2 domain.

BASH MaP also revealed insights into how Spinach fluorescence is enhanced by sequence

alterations. Notably, Broccoli, an RNA aptamer evolved and selected for higher fluorescence,

shows high sequence similarity to Spinach except for key mutations in the P2 stem and

residues proximal to the G-quadruplex core[63] (**Supplementary Data Fig. 7b**). The mutations of

Broccoli are in locations which do not impact G-quadruplex folding, but instead stabilize the P2

domain, and destabilize the alternative base pairing pattern of the P2 domain (**Supplementary**

Data Fig. 6b and **Supplementary Data Fig. 7b).** In this way, Broccoli enhances the

fluorescence of the Spinach aptamer by destabilizing the non-fluorescent alternative tertiary

conformation[63].

BASH MaP also enables prediction of G-quadruplex conformations in cells. Our analysis

identified a unique G-quadruplex formation in the *AKT2* 3'UTR, characterized by an atypical

topology that includes a significantly elongated internal loop. Indeed, recent studies reported

the in vitro formation of RNA G-quadruplexes with unusually long loop lengths in human

5'UTRs[68]. However, current computational algorithms cannot identify which guanines form the

G-quadruplex in atypical or long-loop G-quadruplexes[69]. BASH MaP simplifies this process by

identifying long-range G – G correlations, thus enabling predictions of precise conformations of

the *AKT2* G-quadruplex.

References

1. Lawley, P. & Brookes, P. Further studies on the alkylation of nucleic acids and their constituent nucleotides. *Biochemical Journal* **89**, 127 (1963).

2. Mustoe, A.M., Lama, N.N., Irving, P.S., Olson, S.W. & Weeks, K.M. RNA base-pairing complexity in living cells visualized by correlated chemical probing. *Proceedings of the National Academy of Sciences* **116**, 24574-24582 (2019).

3. Peattie, D.A. & Gilbert, W. Chemical probes for higher-order structure in RNA. *Proceedings of the National Academy of Sciences* **77**, 4679-4682 (1980).

4. Inoue, T. & Cech, T.R. Secondary structure of the circular form of the Tetrahymena rRNA intervening sequence: a technique for RNA structure analysis using chemical probes and reverse transcriptase. *Proceedings of the National Academy of Sciences* **82**, 648-652 (1985).

5. Lempereur, L., Nicoloso, M., Riehl, N., Ehresmann, C., Ehresmann, B. & Bachellerie, J.-P. Conformation of yeast 18S rRNA. Direct chemical probing of the 5' domain in ribosomal subunits and in deproteinized RNA by reverse transcriptase mapping of dimethyl sulfate-accessible sites. *Nucleic acids research* **13**, 8339-8357 (1985).

6. Wells, S.E., Hughes, J.M., Igel, A.H. & Ares Jr, M. [32] Use of dimethyl sulfate to probe RNA structure in vivo. (2000).

7. Homan, P.J., Favorov, O.V., Lavender, C.A., Kursun, O., Ge, X., Busan, S., Dokholyan, N.V. & Weeks, K.M. Single-molecule correlated chemical probing of RNA. *Proceedings of the National Academy of Sciences* **111**, 13858-13863 (2014).

8. Siegfried, N.A., Busan, S., Rice, G.M., Nelson, J.A. & Weeks, K.M. RNA motif discovery by SHAPE and mutational profiling (SHAPE-MaP). *Nature methods* **11**, 959-965 (2014).

9. Zubradt, M., Gupta, P., Persad, S., Lambowitz, A.M., Weissman, J.S. & Rouskin, S. DMS-MaPseq for genome-wide or targeted RNA structure probing in vivo. *Nature Methods* **14**, 75-82 (2017).

10. Gellert, M., Lipsett, M.N. & Davies, D.R. HELIX FORMATION BY GUANYLIC ACID. *Proceedings of the National Academy of Sciences* **48**, 2013-2018 (1962).

11.   Arora, A., Dutkiewicz, M., Scaria, V., Hariharan, M., Maiti, S. & Kurreck, J. Inhibition of
translation in living eukaryotic cells by an RNA G-quadruplex motif. *Rna* **14**, 1290-1296
(2008).

12.   Marcel, V., Tran, P.L., Sagne, C., Martel-Planche, G., Vaslin, L., Teulade-Fichou, M.-P.,
Hall, J., Mergny, J.-L., Hainaut, P. & Van Dyck, E. G-quadruplex structures in TP53 intron
3: role in alternative splicing and in production of p53 mRNA isoforms. *Carcinogenesis*
32, 271-278 (2011).

13.   Blice-Baum, A.C. & Mihailescu, M.-R. Biophysical characterization of G-quadruplex
forming FMR1 mRNA and of its interactions with different fragile X mental retardation
protein isoforms. *Rna* **20**, 103-114 (2014).

14.   Zhang, Y., Yang, M., Duncan, S., Yang, X., Abdelhamid, M.A.S., Huang, L., Zhang, H.,
Benfey, P.N., Waller, Z.A.E. & Ding, Y. G-quadruplex structures trigger RNA phase
separation. *Nucleic Acids Research* **47**, 11746-11754 (2019).

15.   Asamitsu, S., Yabuki, Y., Matsuo, K., Kawasaki, M., Hirose, Y., Kashiwazaki, G., Chandran,
A., Bando, T., Wang, D.O., Sugiyama, H. & Shioda, N. RNA G-quadruplex organizes stress
granule assembly through DNAPTP6 in neurons. *Science Advances* **9**, eade2035 (2023).

16.   Rodriguez, R., Müller, S., Yeoman, J.A., Trentesaux, C., Riou, J.-F. & Balasubramanian, S.
A novel small molecule that alters shelterin integrity and triggers a DNA-damage
response at telomeres. *Journal of the American Chemical Society* **130**, 15758-15759
(2008).

17.   Creacy, S.D., Routh, E.D., Iwamoto, F., Nagamine, Y., Akman, S.A. & Vaughn, J.P. G4
resolvase 1 binds both DNA and RNA tetramolecular quadruplex with high affinity and is
the major source of tetramolecular quadruplex G4-DNA and G4-RNA resolving activity in
HeLa cell lysates. *Journal of Biological Chemistry* **283**, 34626-34634 (2008).

18.   Guo, J.U. & Bartel, D.P. RNA G-quadruplexes are globally unfolded in eukaryotic cells
and depleted in bacteria. *Science* **353**, aaf5371 (2016).

19.   Vicens, Q. & Kieft, J.S. Thoughts on how to think (and talk) about RNA structure.
Proceedings of the National Academy of Sciences **119**, e2112677119 (2022).

20. Mitchell, D., III, Cotter, J., Saleem, I. & Mustoe, A.M. Mutation signature filtering enables high-fidelity RNA structure probing at all four nucleobases with DMS. *Nucleic Acids Research* **51**, 8744-8757 (2023).

21. Küpfer, P.A., Crey-Desbiolles, C. & Leumann, C.J. Trans-lesion synthesis and RNaseH activity by reverse transcriptases on a true abasic RNA template. *Nucleic Acids Research* **35**, 6846-6853 (2007).

22. Zhang, L.-S., Ju, C.-W., Liu, C., Wei, J., Dai, Q., Chen, L., Ye, C. & He, C. m7G-quant-seq: Quantitative Detection of RNA Internal N7-Methylguanosine. *ACS Chemical Biology* **17**, 3306-3312 (2022).

23. Enroth, C., Poulsen, L.D., Iversen, S., Kirpekar, F., Albrechtsen, A. & Vinther, J. Detection of internal N7-methylguanosine (m7G) RNA modifications by mutational profiling sequencing. *Nucleic Acids Research* **47**, e126-e126 (2019).

24. Zhang, L.-S., Liu, C., Ma, H., Dai, Q., Sun, H.-L., Luo, G., Zhang, Z., Zhang, L., Hu, L. & Dong, X. Transcriptome-wide mapping of internal N7-methylguanosine methylome in mammalian mRNA. *Molecular cell* **74**, 1304-1316. e1308 (2019).

25. Piekna-Przybylska, D., Decatur, W.A. & Fournier, M.J. The 3D rRNA modification maps database: with interactive tools for ribosome analysis. *Nucleic Acids Res* **36**, D178-183 (2008).

26. Choi, Y.C. & Busch, H. Modified nucleotides in T1 RNase oligonucleotides of 18S ribosomal RNA of the Novikoff hepatoma. *Biochemistry* **17**, 2551-2560 (1978).

27. Haag, S., Kretschmer, J. & Bohnsack, M.T. WBSCR22/Merm1 is required for late nuclear pre-ribosomal RNA processing and mediates N7-methylation of G1639 in human 18S rRNA. *Rna* **21**, 180-187 (2015).

28. Paige, J.S., Wu, K.Y. & Jaffrey, S.R. RNA Mimics of Green Fluorescent Protein. *Science* **333**, 642-646 (2011).

29. Huang, H., Suslov, N.B., Li, N.-S., Shelke, S.A., Evans, M.E., Koldobskaya, Y., Rice, P.A. & Piccirilli, J.A. A G-quadruplex–containing RNA activates fluorescence in a GFP-like fluorophore. *Nature Chemical Biology* **10**, 686-691 (2014).

30. Warner, K.D., Chen, M.C., Song, W., Strack, R.L., Thorn, A., Jaffrey, S.R. & Ferré-D'Amaré, A.R. Structural basis for activity of highly efficient RNA mimics of green fluorescent protein. *Nature Structural & Molecular Biology* **21**, 658-663 (2014).

31. Fernandez-Millan, P., Autour, A., Ennifar, E., Westhof, E. & Ryckelynck, M. Crystal structure and fluorescence properties of the iSpinach aptamer in complex with DFHBI. *Rna* **23**, 1788-1795 (2017).

32. Cheng, C.Y., Kladwang, W., Yesselman, J.D. & Das, R. RNA structure inference through chemical mapping after accidental or intentional mutations. *Proceedings of the National Academy of Sciences* **114**, 9876-9881 (2017).

33. Štoček, J.R. & Dračínský, M. Tautomerism of Guanine Analogues. *Biomolecules* **10** (2020).

34. Rouskin, S., Zubradt, M., Washietl, S., Kellis, M. & Weissman, J.S. Genome-wide probing of RNA structure reveals active unfolding of mRNA structures in vivo. *Nature* **505**, 701-705 (2014).

35. Olson, S.W., Turner, A.W., Arney, J.W., Saleem, I., Weidmann, C.A., Margolis, D.M., Weeks, K.M. & Mustoe, A.M. Discovery of a large-scale, cell-state-responsive allosteric switch in the 7SK RNA using DANCE-MaP. *Mol Cell* **82**, 1708-1723.e1710 (2022).

36. Merino, E.J., Wilkinson, K.A., Coughlan, J.L. & Weeks, K.M. RNA Structure Analysis at Single Nucleotide Resolution by Selective 2'-Hydroxyl Acylation and Primer Extension (SHAPE). *Journal of the American Chemical Society* **127**, 4223-4231 (2005).

37. Weeks, K.M. & Crothers, D.M. Major Groove Accessibility of RNA. *Science* **261**, 1574-1577 (1993).

38. Roschdi, S., Yan, J., Nomura, Y., Escobar, C.A., Petersen, R.J., Bingman, C.A., Tonelli, M., Vivek, R., Montemayor, E.J., Wickens, M., Kennedy, S.G. & Butcher, S.E. An atypical RNA quadruplex marks RNAs as vectors for gene silencing. *Nature Structural & Molecular Biology* **29**, 1113-1121 (2022).

39. Kwok, C.K., Marsico, G., Sahakyan, A.B., Chambers, V.S. & Balasubramanian, S. rG4-seq reveals widespread formation of G-quadruplex structures in the human transcriptome. *Nature Methods* **13**, 841-844 (2016).

40. Kwok, C.K., Sahakyan, A.B. & Balasubramanian, S. Structural analysis using SHALiPE to

reveal RNA G-quadruplex formation in human precursor microRNA. *Angewandte Chemie*

128, 9104-9107 (2016).

41. Kharel, P., Fay, M., Manasova, E.V., Anderson, P.J., Kurkin, A.V., Guo, J.U. & Ivanov, P.

Stress promotes RNA G-quadruplex folding in human cells. *Nature Communications* **14**,

205 (2023).

42. Krokhotin, A., Mustoe, A.M., Weeks, K.M. & Dokholyan, N.V. Direct identification of

base-paired RNA nucleotides by correlated chemical probing. *Rna* **23**, 6-13 (2017).

43. Sengupta, A., Rice, G.M. & Weeks, K.M. Single-molecule correlated chemical probing

reveals large-scale structural communication in the ribosome and the mechanism of the

antibiotic spectinomycin in living cells. *PLoS biology* **17**, e3000393 (2019).

44. Kikin, O., D'Antonio, L. & Bagga, P.S. QGRS Mapper: a web-based server for predicting G-

quadruplexes in nucleotide sequences. *Nucleic Acids Research* **34**, W676-W682 (2006).

45. Cordero, P., Kladwang, W., VanLang, C.C. & Das, R. The mutate-and-map protocol for

inferring base pairs in structured RNA. *Methods Mol Biol* **1086**, 53-77 (2014).

46. Davis, A.R., Kirkpatrick, C.C. & Znosko, B.M. Structural characterization of naturally

occurring RNA single mismatches. *Nucleic Acids Res* **39**, 1081-1094 (2011).

47. Deigan, K.E., Li, T.W., Mathews, D.H. & Weeks, K.M. Accurate SHAPE-directed RNA

structure determination. *Proceedings of the National Academy of Sciences* **106**, 97-102

(2009).

48. Mathews, D.H., Disney, M.D., Childs, J.L., Schroeder, S.J., Zuker, M. & Turner, D.H.

Incorporating chemical modification constraints into a dynamic programming algorithm

for prediction of RNA secondary structure. *Proceedings of the National Academy of*

Sciences **101**, 7287-7292 (2004).

49. Cordero, P., Kladwang, W., VanLang, C.C. & Das, R. Quantitative dimethyl sulfate

mapping for automated RNA secondary structure inference. *Biochemistry* **51**, 7037-7039

(2012).

50. Zuker, M. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic*

acids research **31**, 3406-3415 (2003).

51.   Sato, K. & Kato, Y. Prediction of RNA secondary structure including pseudoknots for long
sequences. *Briefings in Bioinformatics* **23** (2021).

52.   Strack, R.L., Song, W. & Jaffrey, S.R. Using Spinach-based sensors for fluorescence
imaging of intracellular metabolites and proteins in living bacteria. *Nat Protoc* **9**, 146-
155 (2014).

53.   Do, C.B., Woods, D.A. & Batzoglou, S. CONTRAfold: RNA secondary structure prediction
without physics-based models. *Bioinformatics* **22**, e90-e98 (2006).

54.   Reuter, J.S. & Mathews, D.H. RNAstructure: software for RNA secondary structure
prediction and analysis. *BMC Bioinformatics* **11**, 129 (2010).

55.   Mathews, D.H., Sabina, J., Zuker, M. & Turner, D.H. Expanded sequence dependence of
thermodynamic parameters improves prediction of RNA secondary structure11Edited
by I. Tinoco. *Journal of Molecular Biology* **288**, 911-940 (1999).

56.   Yang, M., Zhu, P., Cheema, J., Bloomer, R., Mikulski, P., Liu, Q., Zhang, Y., Dean, C. &
Ding, Y. In vivo single-molecule analysis reveals COOLAIR RNA structural diversity.
Nature **609**, 394-399 (2022).

57.   Wayment-Steele, H.K., Kladwang, W., Strom, A.I., Lee, J., Treuille, A., Becka, A., Das, R. &
Eterna, P. RNA secondary structure packages evaluated and improved by high-
throughput experiments. *Nature Methods* **19**, 1234-1242 (2022).

58.   Banco, M.T. & Ferré-D'Amaré, A.R. The emerging structural complexity of G-quadruplex
RNAs. *Rna* **27**, 390-402 (2021).

59.   Strack, R.L., Disney, M.D. & Jaffrey, S.R. A superfolding Spinach2 reveals the dynamic
nature of trinucleotide repeat–containing RNA. *Nature Methods* **10**, 1219-1224 (2013).

60.   Tomezsko, P.J., Corbin, V.D.A., Gupta, P., Swaminathan, H., Glasgow, M., Persad, S.,
Edwards, M.D., McIntosh, L., Papenfuss, A.T., Emery, A., Swanstrom, R., Zang, T., Lan,
T.C.T., Bieniasz, P., Kuritzkes, D.R., Tsibris, A. & Rouskin, S. Determination of RNA
structural diversity and its role in HIV-1 RNA splicing. *Nature* **582**, 438-442 (2020).

61.   Filonov, Grigory S., Kam, Christina W., Song, W. & Jaffrey, Samie R. In-Gel Imaging of
RNA Processing Using Broccoli Reveals Optimal Aptamer Expression Strategies.
Chemistry & Biology **22**, 649-660 (2015).

62.   Thiel, B.C., Beckmann, I.K., Kerpedjiev, P. & Hofacker, I.L. 3D based on 2D: Calculating
helix angles and stacking patterns using forgi 2.0, an RNA Python library centered on
secondary structure elements. *F1000Research* **8** (2019).

63.   Filonov, G.S., Moon, J.D., Svensen, N. & Jaffrey, S.R. Broccoli: Rapid Selection of an RNA
Mimic of Green Fluorescent Protein by Fluorescence-Based Selection and Directed
Evolution. *Journal of the American Chemical Society* **136**, 16299-16308 (2014).

64.   Morandi, E., Manfredonia, I., Simon, L.M., Anselmi, F., van Hemert, M.J., Oliviero, S. &
Incarnato, D. Genome-scale deconvolution of RNA structure ensembles. *Nature*
Methods **18**, 249-252 (2021).

65.   Su, Z., Zhang, Y., Gendron, T.F., Bauer, P.O., Chew, J., Yang, W.-Y., Fostvedt, E., Jansen-
West, K., Belzil, V.V. & Desaro, P. Discovery of a biomarker and lead small molecules to
target r (GGGGCC)-associated defects in c9FTD/ALS. *Neuron* **83**, 1043-1050 (2014).

66.   Weldon, C., Behm-Ansmant, I., Hurley, L.H., Burley, G.A., Branlant, C., Eperon, I.C. &
Dominguez, C. Identification of G-quadruplexes in long functional RNAs using 7-
deazaguanine RNA. *Nat Chem Biol* **13**, 18-20 (2017).

67.   Weng, X., Gong, J., Chen, Y., Wu, T., Wang, F., Yang, S., Yuan, Y., Luo, G., Chen, K., Hu, L.,
Ma, H., Wang, P., Zhang, Q.C., Zhou, X. & He, C. Keth-seq for transcriptome-wide RNA
structure mapping. *Nature Chemical Biology* **16**, 489-492 (2020).

68.   Jodoin, R., Bauer, L., Garant, J.-M., Laaref, A.M., Phaneuf, F. & Perreault, J.-P. The folding
of 5'-UTR human G-quadruplexes possessing a long central loop. *Rna* **20**, 1129-1141
(2014).

69.   Mukundan, V.T. & Phan, A.T. Bulges in G-quadruplexes: broadening the definition of G-
quadruplex-forming sequences. *Journal of the American Chemical Society* **135**, 5017-
5028 (2013).

70.   Busan, S. & Weeks, K.M. Accurate detection of chemical modifications in RNA by
mutational profiling (MaP) with ShapeMapper 2. *Rna* **24**, 143-148 (2018).

1120
1121

Acknowledgements

1123

1124   We thank members of the Jaffrey lab for helpful comments and suggestions. We thank the

1125   Genomics Resources Core Facility at Weill Cornell for performing all DNA sequencing. This

1126   work was supported by NIH grants R35NS111631 to S.R.J, and NIH grant T32 GM141949-03 to

1127   M.D.O.

1128

Author Contributions

1130

1131   M.D.O. conceived and designed the experiments. M.D.O. carried out experiments and M.D.O.

1132   and S.R.J. analyzed data. M.D.O. and S.R.J. wrote the manuscript.

1133

Competing interests

1135

1136   S.R.J. is the co-founder of Lucerna Technologies and has equity in this company. Lucerna has

1137   licensed technology related to Spinach, Broccoli, and other RNA-fluorophore complexes. S.R.J.

1138   is a founder of Chimerna Therapeutics and has equity in this company.

1139
Fig. 1 | Reduction and depuration of DMS-modified RNA enables detection of N7G-DMS
adducts with misincorporations.

(**A**) Strategy for the misincorporation detection of N7G methylation. N7-methylated G is reduced

by potassium borohydride and then heated in acidic conditions to yield an abasic site. The

abasic site then proceeds to induce misincorporations in the cDNA following reverse

transcription.

(**B**) Overall schematic for the BASH MaP experimental workflow. RNA is first treated with

dimethyl sulfate (DMS) which produces the following adducts: m^1A, m^3C, m^7G, and to a lesser

extent m^1G and m^3U. DMS-modified RNA is then subjected to reduction by potassium

borohydride (800 mM) for 4 h at room temperature followed by purification and heating in a pH

2.9 buffer of acetic acid and sodium acetate for 4 h at 45°C. RNA is then purified and subjected

to reverse transcription with enzymes and buffer conditions which promote cDNA

misincorporations at methylated bases.

(**C**) Optimization of reduction duration and efficiency of abasic sites to induce misincorporations.

To determine the optimal reduction duration, we treated total HeLa RNA with potassium

borohydride (800 mM) for various durations of time, performed depurination for 4 h in acid

buffer, and then prepared RT-PCR amplicons surrounding the endogenously methylated base

m^7G1638 in the 18S rRNA. We then quantified the fraction of reads containing a

misincorporations at G1638 divided by the total number of reads to yield a misincorporations

rate at G1638 following amplicon sequencing. Plots of misincorporations rates revealed that 4 h

of borohydride treatment induced an average misincorporations rate of 70% at G1638.

(**D**) Misincorporation signature of abasic sites under reverse transcription conditions for

detection of methylated bases. On the left, quantification of the types of misincorporations at

G1638 as described in **Fig. 1C** for SuperScript II, a reverse transcriptase enzyme commonly

used to detect methylated bases with cDNA misincorporations. On the right, fraction of each

type of misincorporation calculated collectively from all G residues in Spinach following reverse

transcription with SuperScript II. Spinach RNA was either modified with DMS (170 mM) for 8

min at 25°C or treated with an ethanol control. Modified and control RNA was then reduced with

potassium borohydride (800 mM) for 4 h or incubated in water for 4 h. All three Spinach

samples underwent identical heating in acidic buffer conditions before undergoing reverse

transcription. Comparison of types of misincorporations shows that reduction of DMS treated

Spinach RNA produces a misincorporation signature at G residues which mirrors the positive

control G1638 when reverse transcribed with SuperScript II.

(**E**) Reduction of DMS treated Spinach RNA produces novel misincorporation data at G bases.

To determine if Spinach is highly modified by DMS at N7G we utilized the experimental data as

described in **Fig. 1E** with an additional control group in which DMS was omitted but the sample

underwent reduction and depurination. Not shown, all four samples underwent identical heating

in acidic buffer prior to reverse transcription. We then plotted the misincorporation rate of each

G in Spinach for each experimental condition. This misincorporation rate reveals a dramatic

increase in misincorporation rates for G bases in Spinach modified with DMS and reduced with

potassium borohydride.

1181

1182   (**F**) Reproducibility of BASH MaP. Spinach RNA was probed with either 85 mM or 170 mM

1183   DMS for 8 min at 25°C and then reduced and depurinated. The misincorporation rate at each

1184   position in Spinach was compared between the two samples and a linear regression was

1185   performed which showed an R^2 of 0.9928 demonstrating high reproducibility.

1186

(**G**) Effect of reduction and depurination on the detection of m^1A, m^3C, and m^3U. To determine

whether reduction and depurination of DMS-treated RNA impaired the detection of other

methylated bases, we treated Spinach with DMS (170 mM) for 8 min at 25°C using buffer

conditions which promote the methylation of m^1G and m^3U[2, 20]. Then, DMS-treated Spinach was

either directly reverse transcribed (DMS MaP) or subjected to reduction and depurination

(BASH MaP) before reverse transcription. We then compared the misincorporation rate at each

A, C, and U position in Spinach and performed a linear regression. The R^2 of 0.9222

demonstrates that reduction and depurination do not impair the detection of m^1A, m^3C, and m^3U

generated by the modification of RNA by DMS.

1196

(**H-J**) Receiver operator characteristic curves demonstrate that BASH MaP identifies single-

stranded regions of RNA. To determine if BASH MaP could accurately distinguish single-

stranded from base-paired A, C, and U bases, we constructed Receiver Operator Characteristic

(ROC) curves for A, C, and U bases for both BASH MaP and DMS MaP as described in **Fig.**

1G. The larger the area under the curve (AUC), the better a method is at discriminating paired

vs unpaired RNA bases. An AUC = 1.0 demonstrates perfect discrimination ability. Panels **H-J**

demonstrate that BASH MaP accurately discriminates between single-stranded and base-paired

A, C, and U bases.

1205
Fig. 2 | N7G reactivity reveals guanosines involved in tertiary interactions.

(**A**) N7G reactivity overlaid on the secondary structure model of Spinach as derived from the

crystal structure 4TS2[30]. G-quadruplex G's are indicated with a red asterisk. N7G reactivity is

colored on a continuous gradient from grey (low reactivity) to yellow (moderate reactivity) to red

(high reactivity). N7G reactivities represent normalized values derived from misincorporation

rates. Spinach BASH MaP data was obtained as described in **Fig. 1G** and no background

subtraction was utilized.

1213

(**B**) Comparison between structural features in Spinach and observed BASH MaP

misincorporation rate at G bases. To determine which structural features impact N7G

methylation rate we utilized the crystal structure 4TS2, and classified individual G's as either

base-paired, single-stranded, engaged in an RNA G-quadruplex, or engaging in a base triple.

This plot reveals single-stranded G's have relatively high methylation rates whereas RNA G-

quadruplex G's are strongly protected from N7G methylation.

1220

1221   (**C**) Comparison between RNA G-quadruplex G's and all other G's in Spinach reveals RNA G-

1222   quadruplex G's are the most protected G's from N7G methylation in Spinach.

1223

(**D**) Schematic of alternative conformations the 15x polyUG (pUG) repeat RNA can adopt in

solution. The 15x pUG repeat can adopt four distinct RNA G-quadruplex conformations of 12

consecutive GU repeats. Each of these four distinct conformations is indicated as Register 1

through 4. Each register is uniquely defined by a set of three G's not engaged in an RNA G-

quadruplex (colored red). These G's are predicted to be single stranded or base-paired and

thus display much higher reactivity than the G's engaged in the RNA G-quadruplex. To show

how the G-quadruplex is rearranged in each conformation, G42 is labeled with a red asterisk.

1231

(**E**) BASH MaP misincorporations rate plot of pUG G's. 15x pUG RNA was refolded in

potassium buffer (100 mM) and modified with DMS (170 mM) for 6 min at 37°C. G's engaged in

a G-quadruplex for each unique register are indicated at the bottom of the plot. G's predicted to

be engaged in a G-quadruplex in each of the four registers is bolded and bracketed below the

plot. The misincorporation rate plot reveals that G's engaged in a G-quadruplex in all four

registers display a strong protection from DMS methylation. Consequently, G20 and G48,

which are only predicted to be protected in a single register show the highest misincorporation

rates.

1240

1241   (**F**) Quantification of **Fig. 2E** shows the relationship between G inclusion in RNA G-quadruplex

1242   registers and measured misincorporation rate. The plot reveals increased protection from N7G

1243   methylation for G's as the number of G-quadruplex conformations increases.

1244

(**G**) BASH MaP of a G-quadruplex in the 3'UTR of the *AKT2* mRNA. SH-SY5Y cells were

treated with DMS (170 mM) for 6 min at 37°C. Primers were designed to specifically amplify the

putative G-quadruplex region in the *AKT2* mRNA. Misincorporation rates were converted to

normalized reactivity values as described in METHODS. Putative G-quadruplex G's as

identified in rG4-seq[39] are highlighted in red. Red bars below the sequence indicate putative G-

quadruplex G's with strong protections from N7G-DMS methylations. G's previously identified

as engaged in a G-quadruplex in cells are indicated below with a black bar. The normalized

reactivity plot reveals N7G protections from DMS at previously identified in cell G-quadruplex

G's as well as other G-tracts. Together, these data support the formation of a G-quadruplex in

3'UTR of the *AKT2* mRNA in SH-SY5Y cells.

1255
Fig. 3 | BASH MaP reveals networks of co-occurring modifications in the Spinach G-
quadruplex.

(**A-B**) Heatmap of correlation strength between misincorporation that co-occur on the same

sequencing read for Spinach treated with DMS MaP (**A**) or BASH MaP (**B**). Spinach was

treated with DMS (170 mM) for 8 min at 25°C. Each point represents a G-test significance

value as calculated by RING Mapper which was then scaled to a value between zero and one.

Values closer to one appear darker in the correlation heatmap and represent higher G-test

correlation strength.

(**C**) Three-dimensional model of Spinach core ligand binding domain with numbering scheme

used in **Fig. 3A-D** and **Fig. 3F-G**. G-quadruplex and mixed tetrad interactions in Spinach are

indicated with a grey plane. The ligand DFHBI-1T interacts with G52 and sits between the

upper G-quadruplex tetrad and a U-A-U base triple. Structural domains P2, P3, and the core

domain are indicated.

(**D**) Close up of boxed region in **Fig. 3A** shows the pattern of co-occurring misincorporations

surrounding the bases involved in the G-quadruplex of Spinach (marked in red). This heatmap

displays predominantly co-occurring misincorporations between A – A, A – C, and A – G

positions.

(**E**) Close up of boxed region in **Fig. 3B** shows the pattern of co-occurring misincorporations

surrounding the bases involved in the G-quadruplex of Spinach (marked in red). This heatmap

is enriched in co-occurring misincorporations between G – G positions and displays correlations

between G's involved in the G-quadruplex of Spinach.

(**F-G**) Network analysis of G – G correlations in DMS MaP (**F**) and BASH MaP (**G**) of Spinach.

A network was constructed by representing G positions in Spinach as vertices and creating

edges between G positions that display co-occurring misincorporations above a Z-score

threshold of 2.0. BASH MaP of Spinach reveals that G's form two major clusters. The smaller

cluster contains six of nine G's which comprise the Spinach G-quadruplex.

1282

Fig. 4 | Multiplexed probing of single nucleotide mutants identifies N7G and base

stacking interactions.

(**A-E**) Mutate and Map (M2) heatmaps of DMS MaP (**A**) and BASH MaP (**B**) of randomly

mutagenized Spinach RNA. A mutate and map heatmap plots the chemical reactivity profile for

RNAs with a PCR-derived mutation at a specific position along the length of the RNA. When a

position in an RNA is mutated through mutagenic PCR, it is predicted that all interacting

nucleotides will display an increase in chemical reactivity. Each row of the heatmap (Mutation

Position) represents sequencing reads with a PCR-derived mutation at the indicated position

within Spinach. Each column of the heatmap (Mapped Position) represents the

misincorporation rate at the indicated position in Spinach. Visualization of changes in reactivity

to DMS which are induced by point mutations is enabled by performing Z-score normalizations

for each column individually. Only positive Z-scores are plotted to display increases in chemical

reactivity due to PCR-derived mutations. M2 BASH MaP displays unique signals in the mutate-

and-map heatmap in the G-quadruplex (red box) and P3 (grey box) region of Spinach.

(**C**) G-quadruplex topology of Spinach. Spinach contains a two-tiered G-quadruplex with a

mixed tetra stacked below. G-quadruplex numberings and colors are used to denote G-tetrads.

The top G-tetrad is colored blue. The middle G-tetrad is colored black. The bottom mixed

tetrad is colored red. N7G interactions are denoted with a yellow dotted line.

(**D**) Mutate-and-Map has previously been used to identify base-pair interactions in RNA. To

determine whether N7G interactions could be identified by mutate-and-map we examined the

mutate-and-map heatmap generated by M2 BASH MaP of Spinach shown in **Fig. 4B** zoomed in

on the Spinach G-quadruplex core domain. G-quadruplex G's are indicated in red. The plot

reveals that PCR-derived mutations in G-quadruplex G's induce increases in N7G reactivity to

DMS throughout the entire Spinach core domain. This suggests that PCR-derived mutations

within the Spinach G-quadruplex perturb the structure of the entire core domain.

(**E**) Zoom in of the M2 DMS MaP heatmap for the P3 domain of Spinach. P3 stem and loop

nucleotides are indicated below the plot. The base-pairing pattern of the P3 domain is clearly

identified as a diagonal line in the mutate-and-map heatmap. Base pairs are highlighted with

dotted arrows.

(**F**) Zoom in of M2 BASH MaP heatmap for the P3 domain of Spinach. G's which display long

vertical lines in the heatmap and give the P3 region a jagged appearance are highlighted in

blue. The vertical lines indicate that PCR-derived mutations at multiple adjacent nucleotides all

cause increased reactivity to DMS of highlighted G's. This suggests that N7G reactivity to DMS

is sensitive to local disruption of helix stacking.

(**G**) Crystal structure of the Spinach P3 stem. The P3 stem is denoted in grey. G bases which

display signals in the M2 BASH MaP heatmap are highlighted in blue. Hyper-reactive G's at

helix termini such as G81 are colored red.

1320
Fig. 5 | Tertiary folding constraints enable accurate secondary structure modeling of
Spinach.

(**A**) Spinach secondary structure model derived from the crystal structure 4TS2. This secondary

structure model was utilized as a benchmark for comparing structure modeling approaches.

(**B**) Spinach secondary structure models generated by the indicated combination of

experimental data (DMS MaP or BASH MaP) and folding algorithm (mFold, RNAstructure,

DAGGER). To determine whether structure probing data could improve the modeling of

Spinach secondary structure, we assessed the sensitivity and specificity of a variety of

computational approaches. Base pairs which are correctly predicted are indicated by green

bars. Base pairs which are incorrectly predicted are indicated with red dashed lines. A base

pair was determined to be correct if the true base pairing partner was within one base (+/-) from

the indicated pairing partner. For detailed explanation of settings used for each RNA secondary

structure modeling approach see METHODS. Incorporation of experimental data improved

Spinach secondary structure modeling; however, all structures included false helices and lacked

the P2 domain of Spinach.

(**C**) Tertiary-folding constraints derived from N7G-reactivity data are implemented through

modification of the DaVinci data analysis pipeline (DAGGER). To generate tertiary constraints,

G's in the bottom quartile of N7G reactivity are first identified. Then, all pairs of bottom quartile

G's which display significant rates of co-occurring misincorporations with each other are

identified as likely to be engaged in a tertiary interaction. These positions are indicated by

annotating the base as lowercase in the input FASTA file for a modified DaVinci analysis

pipeline. Each sequencing read is first converted to a bitvector where a zero represents no

misincorporation and a one represents a misincorporation. The DaVinci pipeline forces sites of

misincorporations to be single stranded upon subsequent folding. G's identified as likely to be

engaged in a tertiary interaction are forced to be single stranded by editing the bitvectors and

setting the value at each tertiary G to one. Sequencing reads with a misincorporation at any G

identified as tertiary are treated separately because a modification at these positions indicates a

change in tertiary structure. For these bitvectors, tertiary G's are allowed to be considered for

base pairing by ContraFold, the folding engine used in DaVinci. After folding of each unique

sequencing read, RNA secondary structures are converted to forgi vectors which utilize the

Forgi library to encode RNA structure in a string of numbers. Principle component analysis

(PCA) is then performed on the forgi vectors, and the ensemble of RNA structures is visualized

through a plot of the first two principle components. Clustering of related RNA structures is

performed in the PCA reduced dimensional space using techniques such as Kmeans clustering.

Together, this pipeline enables more accurate structure modeling of G-quadruplex containing

RNA.

(**D**) Tertiary constraints and DAGGER analysis of BASH MaP treated Spinach accurately model

Spinach secondary structure. To determine whether tertiary folding constraints could improve

Spinach structure modeling, we implemented the technique as described in **Fig. 5C** and applied

it to Spinach BASH MaP data. Base pairs which are correctly predicted are indicated by green

bars. Base pairs which are incorrectly predicted are indicated with red dashed lines. A base

pair was determined to be correct if the true base pairing partner was within one base (+/-) from

the indicated pairing partner. The resulting secondary structure most closely matches the

crystallographic secondary structure through formation of the P2 domain and absence of false

helices.

1366
Fig. 6 | RNA structure deconvolution reveals G-quadruplex and P2 misfolding in Spinach.

(**A**) RNA structure deconvolution identifies two conformations of Spinach. To identify multiple

conformations of Spinach, we applied the program DANCE which utilizes a Bernoulli mixture

model to identify mutually exclusive patterns of misincorporation in sequencing data. Spinach

was subjected to BASH MaP and the sequencing data was input into the DANCE pipeline.

DANCE identified two conformations denoted as State 1 and State 2 of 80.6% and 19.4%

abundance. Misincorporation rates for each state were converted into reactivity values through

normalization and plotted for comparison (see METHODS).

(**B**) Reactivity differences between DANCE-identified states in Spinach reveal changes in G-

quadruplex core and P2 domains. To identify which regions of Spinach adopted different

structures between the two states identified by DANCE, we plotted the difference between

reactivities of State 1 and State 2. The plot of change in reactivity shows that State 1 and State

2 differ in the core and P2 domains but remain unchanged in stems P1 and P3.

(**C**) Spinach alternative states display differential N7G reactivity at G-quadruplex G's. To

determine whether the alternative states of Spinach display differences in N7G reactivity, we

compared the misincorporation rate of each G for State 1 and State 2. G-quadruplex G's are

indicated below the plot in red. The plot shows that most G's in Spinach display no change in

N7G reactivity to DMS whereas G-quadruplex G's show marked changes in N7G reactivity to

DMS.

(**D-E**) Spinach G-quadruplex G's are differentiated from all other G's in State 1 (**D**) ****p <

0.0001, unpaired t-test with Welch's correction. G-quadruplex G's show no difference in

misincorporation rates for State 2 (**E**) ns p=0.2262, unpaired t-test with Welch's correction.

(**F**) The Spinach G-quadruplex is unfolded in State 2. To determine whether the G-quadruplex

in Spinach was unfolded in State 2 we compared the misincorporation rate of G-quadruplex G's

for the population average and DANCE deconvolved States 1 and 2. The plot shows that G-

quadruplex G's display increased misincorporation rates in State 2 which suggests State 2

consists of an unfolded G-quadruplex. ****p < 0.0001, Tukey's multiple comparison test.

(**G**) Single-stranded loops in the Spinach G-quadruplex show decreased reactivity in State 2.

To determine whether the single-stranded loops in the Spinach G-quadruplex core remain

unpaired, we compared misincorporation rates between State 1 and State 2. The plot shows

that all single-stranded loop residues show reduced reactivity to DMS in State 2 which suggests

these positions display increased base-pairing interactions in State 2.

(**H**) Nucleotide substitutions in Spinach G-quadruplex loop residues reduce Spinach

fluorescence. To determine whether Spinach G-quadruplex loop residues make base-pair

interactions with G-quadruplex G's in the Spinach alternative conformation, we systematically

changed loop residues from A to C which should stabilize any base-pairing interactions with G's.

We quantified Spinach fluorescence through an in-gel fluorescence assay (see METHODS).

The plot shows that conversion of A residues to C residues in the G-quadruplex loop region

induces a progressive loss in Spinach fluorescence.

(**I**) DAGGER clustering with tertiary constraints applied to Spinach BASH MaP data identifies

two clusters with altered base pairing in the P2 domain. To identify alternative base pairing

conformations of the misfolded Spinach, we utilized the orthogonal single molecule analysis

method DaVinci with N7G reactivity data (DAGGER). We incorporated N7G reactivity data to

create tertiary folding constraints before DAGGER folding and clustering (see METHODS).

Dimensional reduction and clustering identified two major clusters denoted Cluster 1 and

Cluster 2. The most representative secondary structure of each cluster is boxed and indicated

by the bit number. The DaVinci clustering plot reveals that the misfolded Spinach displays a

register shift in the P2 domain. Cluster 1, colored green, is consistent with the Spinach

crystallographic secondary structure. Cluster 2 is colored orange.

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427 **METHODS**

1428

1429 **Reagents and equipment**

1430

All buffers and NTPs were purchased from commercial sources. Dimethyl sulfate was

purchased from Sigma Aldrich (77-78-1). Potassium borohydride was purchased from Santa

Cruz Biotechnology (SC-250747). SuperScript II reverse transcriptase was purchased from

Invitrogen. Marathon RT was purchased from Kerafast. DFHBI-1T was purchased from Lucerna

Technologes. N-methyl mesoporphyrin IX (NMM) was purchased from Santa Cruz

Biotechnology. DNAse I was purchased from Sigma Aldrich. DNA constructs and primers were

ordered from Integrated DNA Technologies (IDT) and Twist Bioscience. Spin columns were

purchased from Zymo Research. AMPure XP beads were purchased from Beckman Coulter.

Barcoded sequencing oligos were purchased from New England BioLabs. AmpliScribe T7 High

Yield Transcription Kit were purchased from Biosearch Technologies.

1441

1442 **Cell culture**

1443

1444 HeLa cells were grown in DMEM containing 10% FBS with 100 U ml^{-1} penicillin and 100 μg ml^{-1}

1445 ^{1} streptomycin under standard culturing conditions (37°C with 5.0% CO_2). SH-SY5Y cells were

1446 grown in a 1:1 ratio of DMEM to F12 media containing 10% FBS with 100 U ml^{-1} penicillin and

1447 100 μg ml^{-1} streptomycin under standard culturing conditions (37°C with 5.0% CO_2).

1448

1449 **Reduction and depurination procedure (BASH MaP treatment):**

1450

Reduction and depurination procedures were adapted from a previously published protocol[22].

Between 200 ng and 2 μg of DNAse I treated RNA was brought to 10 μL in water and added to

40 μL of freshy prepared 1M Potassium Borohydride in RNAse free water (final reaction

concentration 800 mM). The reaction was mixed and placed in the dark at room temperature for

various lengths of time. To stop the reaction, 4 volumes of Zymo RNA bindng buffer (200 μL)

was slowly added to avoid foaming. An equal volume of 100% ethanol (250 μL) was then added

to precipitate the RNA. RNA was recovered with RNA clean and concentrator 5 columns and

eluted with 45μL of RNAase free water. Eluted RNA (45 μL) was added to 5μL of 1M sodium

acetate / acetic acid buffer (pH 2.9) and incubated in the dark for 4 hours at 45°C. Reactions

were purified with Oligo clean and concentrator kits following manufacturer's instructions and

eluted in 15µL water.

1462

1463   **HeLa 18s rRNA reduction optimization**

1464

HeLa cells were grown to 80% confluence in 75 mm flasks. Cells were pelleted by centrifugation

at 400 x g for 3 min followed by 1 wash with PBS. Cells were then lysed by the addition of 4 mL

TRIzol reagent. Total RNA was isolated following manufacturer's instructions. 25 µg of total

RNA was subjected to DNAse I treatment and purification with Zymo RNA Clean and

Concentrator 25 columns according to manufacturer's instructions. 1.8 µg of DNAse I treated

HeLa total RNA was input for reduction and depurination and random hexamers were used for

priming the reverse transcription reaction.

1472

1473   **Reverse transcription**

1474

Superscript II: Reverse transcription with SSII was performed as previously described[8, 20].

Briefly, 200 ng of input RNA was mixed with 1µL 50µM random hexamer (HeLa total RNA) or

2µL of 1µM gene specific reverse transcription primer (in vitro transcribed RNAs) and annealed.

Annealed RNA was added to reverse transcription master mix containing 6 mM final

concentration of $MnCl_2$. Reverse transcription reactions were incubated at 23°C for 15 min only

when using random hexamers then 42°C for 3 hours.

1481

Marathon RT: Reverse transcription with Marathon RT was performed with a modified protocol.

First, 200 ng of input RNA was mixed with 2µL10 mM dNTP and 1µL 50µM random hexamer or

2µL of 1µM gene specific RT primer. The volume was brough up to 8µL and the RNA was

annealed to the primer with the following thermocycler settings: 80°C for 1 min, 65°C for 5 min

at which point the RNA was placed on ice for 2 min. Then, an RT master mix was created

containing 4µL of 5x Marathon MaP buffer (250 mM Tris-HCl pH 7.5, 1 M NaCl), 4µLof 100%

glycerol, 1µL of 100 mM DTT, 0.5µL of 20 mM $MnCl_2$, 0.5µL of water. Next, 10µL of RT master

mix was added to 8µL of annealed RNA followed by the addition of 2µL Marathon RT. Reverse

transcription reactions were incubated at 23°C for 15 min only when using random hexamers

then 42°C for 3 hours.

1492

1493 Reverse transcription reactions were inactivated by incubating at 75°C for 10 min. cDNA was

1494 purified with DNA clean and concentrator 5 columns by using 7 volumes of DNA binding buffer

1495 (140 µL) and eluted in 10µL water.

1496

1497 **Sequencing library preparation**

1498

Sequencing libraries were generated via a 2 step PCR. Briefly, 1/5th of purified cDNA (2 µL) was

amplified with step 1 forward and reverse primers. HeLa 18s rRNA samples included a 4-

nucleotide barcode on the forward primer and a 7-nucleotide barcode on the reverse primer

which enabled pooling of purified step 1 PCR products. AKT2 step 1 PCR primers were

designed to amplify a ~200 nucleotide segment of the *AKT2* 3'UTR. In vitro transcribed RNAs

used non-barcoded step 1 forward and reverse primers which were specific to flanking

sequences introduced into the RNA constructs. These flanking sequences are referred to as a

Structure Cassette (SC). Step 1 PCR reactions were assembled on ice by adding to 2µL purified

cDNA: 1.25µL of 10µM forward primer, 1.25µL of 10µM reverse primer, 9µL of water and 12.5µL

of 2x Phusion HF master mix or 2x Phusion GC master mix for AKT2 step 1 PCR. Step 1 PCR

reactions were run with the following thermocycler settings: 98°C for 2 min followed by 8-28

cycles of 98°C for 10 seconds, 65°C for 20 seconds, 72°C for 20 seconds, followed by a final

extension of 2 min. PCR cycle number was optimized for each experiment separately. PCR

reactions were purified with 1.8x AMPure XP beads according to manufactures instructions.

Purified DNA was eluted in 25µL water.

1514

Step 2 PCR was performed with NEB Next Multiplex Oligos for Illumina. Step 2 PCR reactions

were composed of the following: 2-10 ng of purified step 1 DNA brough up to 12µL water, 4µL of

i5 index primer or universal primer, 4µL i7 index primer, and 20µL of 2x Phusion HF or 2x

Phusion GC for AKT2 library prep. Step 2 PCR reactions were run with the following

thermocycler settings: 98°C for 2 min followed by 6-12 cycles of 98°C for 15 seconds, 65°C for

30 seconds, 72°C for 30 seconds, then 72°C for 5 min. PCR cycle number was optimized for

each experiment separately. Step 2 PCR reactions were size selected with AMPure XP beads,

eluted in 25µL of water, and quantified with Qubit 1x dsDNA HS kit. Libraries were pooled and

sequenced on the NovaSeq 6000 PE 2x100, NovaSeq 6000 PE 2x150, MiSeq PE 2x150,

Nextseq2000 P1 600 cycles, or MiSeq Micro 300 cycles.

1525

1526 **RNA construct preparation**

1527

RNA constructs were designed to include flanking stem loops on the 5' and 3' ends of the RNA

that are designed to fold back on themselves as not to interfere with the folding of the RNA of

interest. When present, these flanking sequences are referred to as a structure cassette (SC).

RNAs of interest were ordered from IDT or Twist and then PCR amplified to introduce a T7

promoter as well as the 5' and 3' structure cassette sequences. PCR reactions were purified

with 1.8X AMPure XP beads. RNA was in vitro transcribed with ~200 ng of template dsDNA with

AmpliScribe T7 High Yield Transcription Kit following manufacturer's instructions. Reactions

were purified with RNA Clean and Concentrator 25 kits. RNA quality was assessed on a 10%

denaturing polyacrylamide gel and stained with SYBR gold.

1537

1538   **DMS modification of in vitro transcribed RNA**

1539

In vitro transcribed RNA was probed as previously described[2]. Briefly, 200 ng – 1 µg of RNA

was folded in a buffer designed to promote modification of A and C bases only (200 mM Bicine

pH 7.75 at room temperature, 100 mM KCl) or a buffer which promotes the additional

modification of m^1G and m^3U (200 mM Bicine pH 8.37 at room temperature, 100 mM KCl). RNA

was refolded in buffer by heating to 95°C for 3 min then snap cooling on ice for 1 min. Then

$MgCl_2$ was added to a final concentration of 1-5 mM and the RNA was left at room temperature

for 10 min. Neat DMS was diluted to 1.7M with anhydrous ethanol and added to the folded RNA.

DMS reactions was performed for 10 min at room temperature or 6 min at 37°C. Spinach was

probed at room temperature due to its thermal instability. pUG RNA was probed at 37°C.

Reactions were quenched by the addition of 4 volumes of 20% beta-mercaptoethanol in water

on ice. RNA was purified from the quenched DMS reaction with RNA Clean and Concentrator 5

kit and eluted in 10µL water. RNA was either directly processed or stored at -80°C.

1552

1553   **In cell DMS modification and purification**

1554

DMS modification of SH-SY5Y cells was performed as previously described[35]. Briefly, cells

were grown to 85% confluence in a 10 cm dish. Then, the media was exchanged for a DMS

probing media which consisted of 1:1 DMEM to F12 medium with 10% FBS and 200 mM Bicine

adjusted to pH 8.37 with NaOH at room temperature. Probing media was prewarmed to 37C

and cell media was exchanged with 5.4 mL probing media for 3 min at 37°C. Then 600µL of

prewarmed 1.7 M DMS or prewarmed 600µL ethanol was added to the cells and incubated for 6

min at 37°C. DMS reaction was stopped by the addition of 6 mL 20% 2-mercaptoethanol and

cells were placed on ice. We noticed that DMS probing caused a large portion of adherent cells

to become detached from the plate. We therefore decanted the quenched DMS probing media

off the plate into a 15 mL falcon tube and pelleted the detached cells for 5 min at 4C and 1,000

x g. To isolate the total RNA, 2 mL Trizol reagent was added to the plate for the cells which had

not become detached. Then, the 2 mL Trizol reagent was transferred to solubilize the pelleted

cells which had become detached during DMS probing. Total RNA was then isolated according

to manufacturer's instructions.

1569

1570   **Gene-specific BASH MaP**

1571

Total RNA was treated with DNAse I for 30 min prior to BASH MaP treatment according to

manufacturer's instructions. Three replicates of 1 µg of total RNA was mixed with 1µL 50µM

random hexamers and the volume was adjusted to 11 µL. RNA was annealed by heating to

85°C for 1 min, 65°C for 5 min, then placing on ice for 3 min. Reverse transcription was

performed with SuperScript II as described previously[8]. Reverse transcription reactions were

incubated at 23°C for 15 min and then 42°C for 3 hours. Reactions were inactivated by heating

to 75°C for 10 min. Reverse transcription replicates were pooled and the cDNA was purified with

the Zymo DNA Clean and Concentrator 5 kit and eluted in 10µL water.

1580

1581   **BASH MaP data processing**

1582

1583   Sequencing data was processed using the ShapeMapper pipeline[20, 70].

1584

1585   For HeLa 18s rRNA experiments, sequencing data was aligned to 12 unique 18S amplicons

1586   defined by the unique combinations of 4-nucleotide forward and 7-nucleotide reverse barcode

1587   sequences inserted during step 1 PCR. Sequencing data alignment was performed with

1588   ShapeMapperV2.2 with the following settings: --amplicon --output-parsed --output-aligned-reads

1589   --nproc 15 --output-counted-mutations --min-mutation-separation 0.

1590

Optimal misincorporation separation distance for DMS MaP experiments was identified by

analyzing the misincorporation signatures of m^1G and $m^3psuedoU$ in *E. coli* rRNA as surrogates

for m^1A and m^3C (**Supplementary Data Fig. 8a-b**). *E. coli* 23s rRNA contains two bases

methylated on their Watson-Crick face, m^1G745 and $m^3pseudoU1915$. To analyze whether

reverse transcriptases encode methylated bases as complex misincorporation, as is seen in

SHAPE MaP experiments, we utilized a previously published dataset where *E. coli* 23s rRNA

were reverse transcribed by SSII and Marathon RT[20]. We downloaded the raw data from

GSE225383 and combined all control samples for SSII and Marathon RT. We then aligned the

reads to *E. coli* 23s rRNA sequences with ShapeMapperV2.2 with the following settings: --star-

aligner --random-primer-len 9 --output-aligned-reads. The aligned .sam file was then converted

into a .bam file and imported into integrated genome viewer (IGV). The misincorporation rate

was then recorded for each position relative to the modified bases G745 and U1915. The results

show a clear increase in misincorporation rate for the first position after the modified nucleotide

and this effect is more pronounced for SSII as compared with Martahon RT. This suggests that

methylated bases may produce more complex misincorporation types rather than simple

mismatches, particularly when using SSII. It is important to note that 2 bases beyond the site of

modification, the misincorporation rate falls back to baseline.

1608

We next applied the same analysis procedure to determine the optimal separation distance for

BASH MaP experiments (**Supplementary Data Fig. 8a-b**). We utilized our own HeLa 18s rRNA

dataset where we depurinated m^{7}G1638 into an abasic site and reverse transcribed the rRNA

with SSII and Marathon RT. We similarly converted the aligned .sam file into a .bam file and

utilized IGV to count the misincorporation around G1638. We saw a clear increase in

misincorporation rate after the abasic site for SSII samples but no increase in misincorporation

rate for Marathon RT (**Supplementary Data Fig. 8a-b**). The pattern of misincorporation induced

by abasic sites was like those seen for methylated bases in *E. coli* 23s rRNA. In both cases, an

adduct site does not appear to induce increased misincorporation at distances greater than 2

nucleotides. Together, these data suggest that the optimal misincorporation separation distance

for BASH MaP is 2 nucleotides.

1620

1621   For Spinach, pUG, and AKT2 experiments, sequencing data was aligned to a single fasta

1622   sequence with ShapeMapperV2.2 with the following settings: --amplicon --dms --output-parsed -

1623   -min-mutation-separation 2. Deletions are ambiguously aligned and therefore ignored by

1624   invoking the --dms option.

1625

1626   **Misincorporation signature analysis**

1627

Misincorporation signatures were calculated by aligning the data with ShapeMapperV2.2 with

the following settings: --amplicon --output-counted-mutations --min-mutation-separation 0.

Counts of misincorporation types for each position within an RNA are tabulated by the --output-

counted-mutations field. A minimum misincorporation separation distance of 0 was chosen to

capture the full range of misincorporation types generated in BASH MaP and DMS MaP

experiments.

1634

1635   Receiver operator curve generation

1636

Bases in Spinach were annotated as either base-paired or single stranded based on the

secondary structure provided in **Supplementary Data Fig. 1b**. Misincorporation rates were

derived by probing Spinach in Bicine pH 8.3 buffer which promotes modification of m^3U and

m^1G. Although m^1G reaction is promoted, m^7G formation is favored by an order of magnitude.

Misincorporation rates were calculated in ShapeMapperV2.2 with the follow settings: --amplicon

--dms --min-mutation-separation 2. The crystal structure 4TS2 was utilized and all isolated base

pairs and non-canonical base pairs in which the Watson-Crick face was not hydrogen bonded

were designated as single stranded. This mainly affected annotation of A bases in the J1-2

region. ROC curves were generated with the python script ROC_curve_DAGGER.py and

graphed with PRISM GraphPad.

1647

1648   Correlated misincorporation analysis

1649

ShapeMapper alignment with --output-parsed produces a .mut file which encodes the

misincorporations within a sequencing read as a bitvector where 0 represents no

misincorporation and 1 represents a misincorporation. This .mut file was then input into the

program RingMapper which identifies positions with statistically high rates of co-occurring

misincorporations[7]. The output of RingMapper was processed with the custom python script

ring_pair_to_heatmap.py which filters all negative correlations and creates a square matrix

which was then visualized in Prism GraphPad.

1657

1658   Network analysis of bases with correlated misincorporations

1659

1660   RingMapper was run with default parameters and the output file was filtered to include only

1661   positive correlations between two G nucleotides. A second filter removed all correlations with Z-

1662   scores less than 2.0. Networks of structurally related nucleotides were visualized in Cytoscape

1663   with unique G bases represented as nodes and positive correlations represented as edges.

1664

1665   **AKT2 G-quadruplex conformational analysis**

1666

The sequence GGGUGGGGAGGGGUGGGGUUGGUUCGGGUGGGUGAGGGU,

corresponding to the putative G-rich sequence in the *AKT2* 3'UTR, was input to GQRS Mapper

with the following search parameters: Max length = 45, Min G-Group size = 3, Loop size: from 0

to 36. All predicted G-quadruplex conformations with overlaps were then viewed and saved as

an .html file before undergoing processing with the python script

QGRS_to_average_mutation_rate.py. The output excel file was then filtered by average

misincorporation rate and the ten lowest conformations were plotted in Prism GraphPad where

lowercase 'g' characters indicate G's engaged in a G-quadruplex.

1675

1676   **M2 BASH MaP**

1677

1678   Spinach and polyUG T7 templates were randomly mutagenized by performing 24 rounds of

1679   error-prone PCR as described previously[32]. Mutagenized Spinach and polyUG were then in

1680   vitro transcribed, folded and DMS modified as described prior with minor modifications. $MgCl_2$

1681   was added to a final concentration of 5 mM and Spinach was probed in the presence of 5μM

1682   DFHBI-1T. polyUG RNA was probed in the presence of 2μM NMM.

1683

For M2 Spinach and pUG experiments, sequencing data was aligned to a single fasta sequence

with ShapeMapperV2.2 with the following settings: --amplicon --dms --output-parsed --min-

mutation-separation 2. The output parsed mutation file was then processed with the custom

script mut_to_simple.py which retains only sequencing reads that cover the entire length of the

RNA. Then the output .simple file was used to calculate a unique mutational profile for each

point mutant with the script simple_to_M2_map.py. This script produces a square matrix with

length equal to the length of the RNA and displays how often two mutations co-occur on the

same sequencing read. To better visualize changes in nucleotide reactivity given a mutation at a

certain position, Z-score normalization was performed with the script M2_Map_to_Zscores.py.

The normalized matrix was then visualized in Prism GraphPad as a heatmap.

1694

1695   **Minimum free energy RNA secondary structure modeling**

1696

The baseline secondary structure model of Spinach was generated in mFold with default

parameters. Population average DMS guided RNA secondary structure modeling was

performed as described previously in the DANCE pipeline[35]. Briefly, the python script

DanceMapper.py was run on parsed mutation output files with the following settings for deriving

population average data: --fit --maxc 1. Then, the population average reactivities were

converted to free energy restraints for the RNA folding program RNAstructure through the

foldClusters.py script. When processing BASH MaP data, the additional command --nog was

used with foldClusters.py to ignore the reactivity data for G nucleotides. The resulting RNA

secondary structures were visualized in RNA2D drawer. Mapping of per nucleotide DMS

reactivity onto RNA2D drawer secondary models was performed through the custom python

script color_reactivities_RNA2D_drawer.py. Conformation specific RNA secondary structure

modeling was performed by increasing the maximum allowed conformations from 1 to 5 with --

maxc 5.

1710

1711   **Single molecule probabilistic RNA secondary structure modeling**

1712

1713   A custom pipeline was developed to apply the DaVinci structural analysis method to BASH

1714   MaP[56]. This analysis method utilizes misincorporations to generate a folding constraint for each

1715   sequencing read. Folding constraints are then passed to ContraFold which folds each

1716   sequencing read as a unique RNA molecule. The complete analysis pipeline is as follows:

1717

First, sequencing data is processed through ShapeMapper as described above with --amplicon -

-dms --output-parsed --min-mutation-separation 2 options enabled. The resulting .mut file is

then filtered and converted to a .simple file with mut_to_simple.py. For DMS MaP experiments,

the resulting .simple file is converted into a .bit file by running the simple_to_bit.py python script.

For BASH MaP experiments, the resulting .simple file is instead converted into a .bit file with the

script updated_simple_to_bit_rG42.py. This script requires an input fasta file and sets all upper-

case G positions to be considered for base pairing. In addition, the script constrains all

lowercase letters to be single stranded in the .bit file. If a sequencing read harbors a

misincorporations at the position of one of the lowercase letters, then all lowercase letters in that

sequencing read are instead considered for base pairing. The resulting .bit file is then passed as

an argument to the fold-contrafold-uniq-bits-vectors2.py script. ContraFold was run either with

normal settings or with --noncomplementary enabled. Typically, the fold-contrafold-uniq-bits-

vectors2.py script is stopped after roughly 10,000 RNA molecules have been folded to save on

computation time. The forgi vector encodings of the RNA secondary structures generated by

ContraFold are output in the file forgi-vect-ser2.txt. Dimensional reduction is performed on the

forgi vectors text file with the python script run-pca-on-forgi-vectors.py which reduces the vector

space to 2 dimensions. Kmeans clustering is then performed on the 2-dimensional

representation of RNA structures with the draw-kmeans-clusters.py script. Identification of a

population average structure is done by choosing --num_clusters 1. Multiple conformational

analysis is performed by varying the number of Kmeans clusters. The structure of the most

representative of the Kmeans cluster is retrieved with the python script

fetch_DaVinci_folded_copy.py and visualized with RNA 2D drawer.

1740

1741   **Nucleotide reactivity normalization**

1742

1743   Mutation rates were normalized to reactivity values as previously described with the custom

1744   python script normalize_DAGGER_DANCE.py[20].

1745

1746   **Tertiary folding constraint determination and implementation (DAGGER)**

1747

G's likely to be engaged in tertiary interactions were identified by first selecting G's in the bottom

quartile for misincorporation rate. In situations when reactivity data was available for multiple

conformations, G's in separate conformations were treated as unique G's and all unique G's

were pooled for reactivity normalization. From the bottom quartile of reactive G's, G's which

displayed correlations to other lowly reactive G's as identified by RingMapper were designated

as engaged in a tertiary interaction.

1754

1755   The DaVinci analysis pipeline was modified to utilize tertiary constraints through setting tertiary

1756   G positions to lowercase G's in the fasta input file for generation of .bit files by

1757   updated_simple_to_bit_rG42.py. The modified DaVinci analysis pipeline is reffered to as

1758   DAGGER.

1759

1760   **Crystallographic structures**

1761

1762   The reference Spinach crystal structure is 4TS2[30]. The reference polyUG crystal structure is

1763   7MKT[38].

1764

1765 **In gel fluorescence assay**

1766

Spinach mutants were purchased from Twist and PCR amplified with the Structure cassette T7

forward and reverse primer. Fluorescence was assayed using a previously described in gel

staining assay[61]. The protocol is as follows: 50 ng of RNA was denatured with 2x formamide

RNA loading buffer at 95°C for 3 min and then loaded onto a 10% TBE – Urea PAGE gel. RNA

was run at 280V for 35 min and then the PAGE gel was washed in RNase free water 3 times for

5 minutes each. Then the RNA was refolded and stained inside the gel by incubation with

refolding buffer (10µM DFHBI-1T, 40 mM HEPES pH 7.4, 100 mM KCl, 1 mM $MgCl_2$) for 15-20

minutes with shaking. The gel was imaged then washed in RNase free water 3 times for 5

minutes each. The gel was then imaged in the SYBR gold channel to confirm all DFHBI-1T had

been washed away. The gel was then stained with SYBR gold for 5 min and imaged again.

Fluorescence was normalized to SYBR gold signal.

1778

1779 **Data availability**

1780

1781 Raw and processed sequencing data have been deposited at the GEO under accession number

1782 XXXX. All code used for analysis is available at https://github.com/SamieJaffreyLab.

1783

1784

1785

1786 **Table 1: Step 1 PCR Primers**

RNA sample name	Step 1 forward primer sequence	Step 1 reverse primer sequence
SSII_HeLa_18s_0min_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAAGGCGA</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_0min_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAAGGCGA</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_30min	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATT	GACGTGTGCTCTTCCGATCTNNNNNN<u>CGTACTAG</u>GTGTGTACAAAGGGCAGGGAC

_1	CGTGA	
SSII_HeLa_18s_30min_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>CGTACTAG</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_1hour_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>AGCAGAA</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_1hour_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>AGCAGAA</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_2hour_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TCCTGAGC</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_2hour_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TCCTGAGC</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_3hour_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>GGACTCCT</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_3hour_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>GGACTCCT</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_4hour_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAGGCATG</u>GTGTGTACAAAGGGCAGGGAC
SSII_HeLa_18s_4hour_2	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAGGCATG</u>GTGTGTACAAAGGGCAGGGAC
MaRT_HeLa_18s_4hour_1	ACACGACGCTCTTCCGATCTNNNNN**ATAT**CCCGTTGAACCCCATTCGTGA	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAGGCATG</u>GTGTGTACAAAGGGCAGGGACCATGCCTA
MaRT_HeLa_18s_4ho	ACACGACGCTCTTCCGATCTNNNNN**CGCG**CCCGTTGAACCCCAT	GACGTGTGCTCTTCCGATCTNNNNNN<u>TAGGCATG</u>GTGTGTACAAAGGGCAGGGACCATGCCTA

ur_2	TCGTGA	
Structure Cassette Universal	ACACGACGCTCTTCCGATCTNN NNNGGCTGGCCTTTCGGGCCAA	GACGTGTGCTCTTCCGATCTNNNNNGA ACCGGACCGAAGCCCG
AKT2 3UTR	ACACGACGCTCTTCCGATCTNN NNNAACACCTCTGGGTGTTTGG AGTTTAGC	GACGTGTGCTCTTCCGATCTNNNNNCC GTACAAATATGAAGACGAGGAGAAAGG C

1787

1788

1789 **Table 2: Oligos and primers**

Spinach oligo	**GTATAATACGACTCACTATAGGGCTGGCCTTTCGGGCCAAGGGA CGCGACCGAATGAAATGGTGAAGGACGGGTCCAGCCGGCTGCT TCGGCAGCCGGCTTGTTGAGTAGAGTGTGAGCTCCGTAACTGGT CGCGTCTCGATCCGGTTCGCCGGATCCAAATCGGGCTTCGGTCC GGTTC**
pUG oligo	**GGCTGGCCTTTCGGGCCAAGTGTGTGTGTGTGTGTGTGTGTGTGT GTGTTCGATCCGGTTCGCCGGATCCAAAT**
Structure Cassette T7 Forward	**GTATAATACGACTCACTATAGGGCTGGCCTTTCGGGCCAA**
Structure Cassette T7 Reverse	**GAACCGGACCGAAGCCCGATTTGGATCCGGCGAACCGGAT**
Structure Cassette RT primer	**GAACCGGACCGAAGCCCGA**
Spinach A48C, A90C template	**ACGGGCCAGATATACGCGTAGTTCCTGCTATAATTAGCCTTCCTC ATAAGTTGCACTGCTCCAGGTGATAGTGCGGGAACCTCGATGGT CTTCACACTTTACTTCAGCGTCtggtaggcgtgtacggtgggaggcctatata agcagagctTCTGGCTAACTAGGCTGGCCTTTCGGGCCAAGGGACG CGACCGAATGAAATGGTGAAGGCCGGGTCCAGCCGGCTGCTTC GGCAGCCGGCTTGTTGAGTAGCGTGTGAGCTCCGTAACTGGTCG CGTCTCGATCCGGTTCGCCGGATCCAAAT**

| Spinach A48C, A88C, A90C template | ACGGGCCAGATATACGCGTAGTTCCTGCTATAATTAGCCTTCCTCATAAGTTGCACTGCTCCAGGTGATAGTGCGGGAACCTCGATGGTCTTCACACTTTACTTCAGCGTCtggtaggcgtgtacggtgggaggcctatataagcagagctTCTGGCTAACTAGGCTGGCCTTTCGGGCCAAGGGACGCGACCGAATGAAATGGTGAAGGCCGGGTCCAGCCGGCTGCTTCGGCAGCCGGCTTGTTGAGTCGCGTGTGAGCTCCGTAACTGGTCGCGTCTCGATCCGGTTCGCCGGATCCAAAT |

1790

References

1. Zhang, L.-S., Ju, C.-W., Liu, C., Wei, J., Dai, Q., Chen, L., Ye, C. & He, C. m7G-quant-seq: Quantitative Detection of RNA Internal N7-Methylguanosine. *ACS Chemical Biology* **17**, 3306-3312 (2022).
2. Siegfried, N.A., Busan, S., Rice, G.M., Nelson, J.A. & Weeks, K.M. RNA motif discovery by SHAPE and mutational profiling (SHAPE-MaP). *Nature methods* **11**, 959-965 (2014).
3. Mitchell, D., III, Cotter, J., Saleem, I. & Mustoe, A.M. Mutation signature filtering enables high-fidelity RNA structure probing at all four nucleobases with DMS. *Nucleic Acids Research* **51**, 8744-8757 (2023).
4. Mustoe, A.M., Lama, N.N., Irving, P.S., Olson, S.W. & Weeks, K.M. RNA base-pairing complexity in living cells visualized by correlated chemical probing. *Proceedings of the National Academy of Sciences* **116**, 24574-24582 (2019).
5. Olson, S.W., Turner, A.W., Arney, J.W., Saleem, I., Weidmann, C.A., Margolis, D.M., Weeks, K.M. & Mustoe, A.M. Discovery of a large-scale, cell-state-responsive allosteric switch in the 7SK RNA using DANCE-MaP. *Mol Cell* **82**, 1708-1723.e1710 (2022).
6. Busan, S. & Weeks, K.M. Accurate detection of chemical modifications in RNA by mutational profiling (MaP) with ShapeMapper 2. *Rna* **24**, 143-148 (2018).
7. Homan, P.J., Favorov, O.V., Lavender, C.A., Kursun, O., Ge, X., Busan, S., Dokholyan, N.V. & Weeks, K.M. Single-molecule correlated chemical probing of RNA. *Proceedings of the National Academy of Sciences* **111**, 13858-13863 (2014).
8. Cheng, C.Y., Kladwang, W., Yesselman, J.D. & Das, R. RNA structure inference through chemical mapping after accidental or intentional mutations. *Proceedings of the National Academy of Sciences* **114**, 9876-9881 (2017).

9.   Yang, M., Zhu, P., Cheema, J., Bloomer, R., Mikulski, P., Liu, Q., Zhang, Y., Dean, C. &

Ding, Y. In vivo single-molecule analysis reveals COOLAIR RNA structural diversity.

Nature **609**, 394-399 (2022).

10.   Warner, K.D., Chen, M.C., Song, W., Strack, R.L., Thorn, A., Jaffrey, S.R. & Ferré-

D'Amaré, A.R. Structural basis for activity of highly efficient RNA mimics of green

fluorescent protein. *Nature Structural & Molecular Biology* **21**, 658-663 (2014).

11.   Roschdi, S., Yan, J., Nomura, Y., Escobar, C.A., Petersen, R.J., Bingman, C.A., Tonelli,

M., Vivek, R., Montemayor, E.J., Wickens, M., Kennedy, S.G. & Butcher, S.E. An

atypical RNA quadruplex marks RNAs as vectors for gene silencing. *Nature Structural &*

Molecular Biology **29**, 1113-1121 (2022).

12.   Filonov, Grigory S., Kam, Christina W., Song, W. & Jaffrey, Samie R. In-Gel Imaging of

RNA Processing Using Broccoli Reveals Optimal Aptamer Expression Strategies.

Chemistry & Biology **22**, 649-660 (2015).

1829

1830

1831

1832

1833

1834

1835

1836