Extracellular vesicles adhere to cells predominantly through the interaction of CD151-associated integrin heterodimers and GM1 with laminin

Running head: Mechanisms of extracellular vesicle–cell binding

Tatsuki Isogai[1], Koichiro M. Hirosawa[2], Miki Kanno[3], Ayano Sho[4], Rinshi S. Kasai[5], Naoko Komura[2], Hiromune Ando[1,2,3,4], Keiko Furukawa[6], Yuhsuke Ohmi[6], Koichi Furukawa[6], Yasunari Yokota[7], Kenichi G. N. Suzuki[1,2,3,4,5]

[1]The United Graduate School of Agricultural Science, Gifu University, Gifu 501-1193, Japan

[2]Institute for Glyco-core Research (iGCORE), Gifu University, Gifu 501-1193, Japan

[3]Graduate School of Natural Science and Technology, Gifu University, Gifu 501-1193, Japan

[4]Faculty of Applied Biological Sciences, Gifu University, Gifu 501-1193, Japan

[5]Division of Advanced Bioimaging, National Cancer Center Research Institute (NCCRI), Tokyo 104-0045, Japan

[6]Department of Biomedical Sciences, Chubu University, Kasugai 487-8501, Japan

[7]Department of Information Science, Faculty of Engineering, Gifu University, Gifu 501-1193, Japan

Corresponding author

Kenichi G. N. Suzuki

ORCID: 0000-0002-8436-5757

Institute for Glyco-core Research (iGCORE), Gifu University, Gifu 501-1193, Japan

Tel: 011-81-58-293-3330

E-mail: suzuki.kenichi.b7@f.gifu-u.ac.jp

Summary

Quantitative assessments using single-molecule imaging and super-resolution microscopy revealed that all extracellular vesicle subtypes derived from four distinct tumor cell lines, regardless of size, bind to laminin predominantly via CD151-facilitated integrin heterodimers and GM1, but not as much to fibronectin.

Abstract

Tumor-derived extracellular vesicles (EVs) have attracted significant attention, yet the molecular mechanisms that govern their specific binding to recipient cells remain elusive. Our in vitro study utilizing single-particle tracking demonstrated that integrin heterodimers comprising α6β4 and α6β1 are responsible for the binding of small-EV (sEV) subtypes to laminin. EVs derived from four distinct tumor cell lines, regardless of size, exhibited high binding affinities for laminin but not for fibronectin, although fibronectin receptors are abundant in EVs and have functional roles in EV-secreting cells. Our findings also revealed that the robust binding of integrins in EVs to laminin is preserved by CD151 rather than by talin-1 inside-out signaling and is inhibited by a molecule that associates with CD151 via cholesterol. The sEV–laminin interaction is also induced by GM1. Super-resolution movie observation revealed that sEV integrins bind only to laminin on living recipient cells. Thus, we demonstrated that all EV subtypes bind to laminin predominantly via CD151-facilitated integrin heterodimers and GM1.

Introduction

Extracellular vesicles (EVs) of a variety of sizes (~30-1000 nm) contain a diverse array of cargo,

including microRNAs, metabolites, and proteins, and play a critical role in intercellular

communication (Valadi et al., 2007; Raposo and Stoorvogel, 2013; van Niel et al., 2018). EVs

have attracted extensive attention in a wide range of research fields, and the number of published

papers has increased dramatically (Théry et al., 2018; Witwer et al., 2021). In particular, several

reports have suggested that tumor-derived EVs can induce phenotypic changes in recipient cells,

creating a premetastatic niche that promotes cancer metastasis (Peinado et al., 2012; Ono et al.,

2014). sEV studies might elucidate the underlying mechanism of the 'seed and soil' hypothesis

(Paget, 1989), which posits that cancer cells tend to metastasize and proliferate in organs suitable

for these activities (Hart and Fidler, 1980; Peinado et al., 2017). A previous proteomic study of

sEVs revealed that integrin α6β4 and α6β1 in sEVs from 4175-LuT human breast cancer cells

were associated with lung metastasis, while αVβ5 in small EVs (sEVs) from BxPC3 human

pancreatic adenocarcinoma cells was linked to liver metastasis (Hoshino et al., 2015). In

addition, several studies have shown that blocking integrin in sEVs through anti-integrin

antibodies or integrin ligands inhibits the internalization of integrin by recipient cells (Nazarenko

et al., 2010; Altei et al., 2020). However, no direct evidence that integrin heterodimers in sEVs

bind to the extracellular matrix (ECM) components on recipient cell plasma membranes (PMs)

has been reported. In addition, the binding affinities between EVs and diverse ECM components have yet to be determined via comparative analysis. Furthermore, the detailed molecular-level mechanisms that govern the selective binding of sEVs to recipient cell PMs remain elusive. Elucidating these mechanisms is critical for developing effective strategies to inhibit cancer metastasis (Möller and Lobb, 2020; Marar et al., 2021). Moreover, it is essential to validate the biological functions of sEVs and their roles in physiological and pathological processes (Gurung et al., 2021).

In this study, we aimed to investigate whether integrin heterodimers play a critical role in the binding of sEVs to ECM components on recipient cell PMs. Furthermore, we sought to elucidate the mechanisms of selective binding using our cutting-edge imaging techniques. Since we demonstrated that sEVs can be classified into several subtypes with different tetraspanin marker proteins (CD63, CD81, and CD9) and that each subtype has unique properties (Hirosawa et al., 2024), it was crucial to examine the binding of each sEV subtype to ECM components. To this end, we first established an in vitro assay system that enabled us to quantitatively evaluate the binding of three sizes of EVs (small EVs, sEVs; medium-sized EVs, mEVs; and microvesicles; MVs) to ECM components on glass by single-particle tracking with single-molecule detection sensitivity. Second, we developed a state-of-the-art imaging system that enables the acquisition of super-resolution (dSTORM) movies, as opposed to static images.

These advances enabled us to examine the binding affinities of various EV subtypes to ECM components on living cell PMs.

Results

Characterization of sEVs by single-particle tracking with single-molecule detection sensitivity

To investigate whether integrin subunits in tumor-derived sEVs are essential for selective binding to ECM components, we first characterized sEVs isolated from human prostate cancer (PC3) cells using an ultracentrifugation-based (pellet-down) method (Fig. S1A). These sEVs displayed circular morphology, with a diameter of 83 ± 19 nm (mean ± S.D.) (Fig. S1B), as determined by transmission electron microscopy (TEM) with negative staining. Consistently, the diameter measured by a tunable resistive pulse sensing instrument (qNano) was 69 ± 17 nm (mean ± S.D.) (Fig. S1C).

Recently, we demonstrated that PC3 cell-derived sEVs comprise several distinct subtypes, each harboring different tetraspanin marker proteins, as determined by single-particle imaging analysis (Hirosawa et al., 2024). Approximately 40% of the sEVs contained the CD63, CD81, and CD9 tetraspanin marker proteins, while the remaining 60% of the sEVs contained only CD63. Furthermore, sEVs that contained only CD63 colocalized with caveolae in the

recipient cell PMs, whereas those containing all three tetraspanin proteins did not. These findings

indicate that the sEV subtypes interact differently with recipient cell PM structures (Hirosawa et

al., 2024). Therefore, in this study, to investigate the roles of integrin heterodimers in different

sEV subtypes, we isolated sEVs from PC3 cells that stably expressed CD63-Halo7, CD81-

Halo7, or CD9-Halo7. The tetraspanin-Halo7 in these sEVs was labeled by conjugation of the

SaraFlour650T (SF650T) ligand, henceforth referred to as "sEV-tetraspanin-Halo7-SF650T". We

successfully observed single sEV-tetraspanin-Halo7-SF650T particles using total internal

reflection fluorescence microscopy (TIRFM) with no contamination from free dye molecules or

autofluorescence from nonlabeled sEVs (Fig. S1D). Since the estimated average numbers of

CD63-Halo7, CD81-Halo7, and CD9-Halo7 molecules per sEV particle are between 4.2 and 4.8

(Hirosawa et al., 2024), according to the Poisson distribution, more than 98% of sEV particles

should contain at least one Halo7-tagged tetraspanin protein. To standardize the concentrations of

sEVs derived from different cell lines, we utilized single-particle imaging with single-molecule

detection sensitivity to directly measure the number of sEV-tetraspanin-Halo7-SF650T particles

bound to the antibody-coated glass. After incubating the sEV solution (2×10^{10} particles/ml) on

antibody-coated glass at a dilution factor (df) of 1, 3 or 9, followed by three washes with HBSS,

we obtained TIRFM images of single sEV-CD63-Halo7-SF650T particles (Fig. S1E). The

number of sEVs bound to the glass decreased in accordance with the dilution factor (df), which

allowed us to obtain a calibration curve (Fig. S1F-J). By directly determining the concentration of sEV-tetraspanin-Halo7-SF650T and adjusting the concentration of the sEV solution to 0.8-4 $\times$ 10^{10} particles/ml, we were able to compare the binding numbers of sEVs derived from different cell types to ECM-coated glass.

Knockout of integrin subunits

We employed CRISPR–Cas9 gene editing to knock out the integrin β1, integrin β4, integrin α2 or integrin α6 subunit in PC3 cells to investigate the role of these subunits in the binding of sEVs to ECM components. Western blotting analysis of the sEVs obtained from intact cells revealed the presence of the integrin β1, integrin β4, integrin α2, integrin α6, integrin α3, integrin α5, and integrin αV subunits, as well as greater abundances of the tetraspanin marker proteins CD63, CD81, and CD9 in comparison to those in parental cells (Fig. 1A). We confirmed that the integrin β1, integrin β4, integrin α2, and integrin α6 subunits were indeed eliminated in sEVs derived from integrin-KO cells. Additionally, integrin α2 and α3 were below the detection limit in sEVs from integrin β1-KO cells (Fig. 1A). Moreover, integrin β4 was not detected in sEVs from integrin α6 KO cells, and integrin α6 expression was decreased in sEVs from β4 KO cells (Fig. 1A). Thus, the knockout of integrin subunits in sEVs often results in decreased expression levels of integrin heterodimer counterparts (Fig. 1 B) (Hynes, 2002), while the expression levels

of other integrin subunits remain unaffected.

Integrin β1 in tumor-derived sEVs plays a critical role in binding to laminin and collagen type I on glass

Integrins are responsible for bidirectional signaling, namely, inside-out and outside-in signaling. For instance, intracellular signaling molecules such as talin can activate integrins (Hynes, 2002; Moser et al., 2009; Lu et al., 2022). Although sEVs contain integrin heterodimers of ECM receptors, this does not necessarily imply that the integrin heterodimers in sEVs can bind to the ECM. Hence, to investigate whether integrin subunits in tumor-derived sEVs are responsible for sEV binding to ECM components, we first performed in vitro binding assays. dSTORM images of immunostained fibronectin, laminin, and collagen type I coated on glass showed that all the ECM components were attached to and evenly dispersed on the glass surfaces (Fig. 2A). Furthermore, almost all the cells that secreted sEVs in this study attached to and spread on all three ECM components coated on glass at similar velocities (Fig. 2B and C), indicating that integrin heterodimers that act as receptors for all three ECM components are active on the cell PMs.

Fig. 3A shows TIRFM images of individual sEV-CD63-Halo7-SF650T particles derived from wild-type PC3 cells (PC3-WT) and integrin β1 KO cells (PC3-β1KO) adhered to glass

surfaces coated with fibronectin, laminin, or collagen type I. Quantitative analysis revealed that

the number of all subtypes of sEV-tetraspanin-Halo7-SF650T particles bound to fibronectin per

image area (82 μm × 82 μm) was very small; therefore, no reduction in quantity due to integrin

β1 KO was observed (left in Figs. 3B and S2A and B and Table S1). Conversely, the numbers of

all subtypes of sEV-tetraspanin-Halo7-SF650T particles attached to collagen type I were higher

than those bound to fibronectin and significantly decreased by integrin β1 KO (middle in Figs.

3B and S2A and B and Table S1). Moreover, all subtypes of sEV-tetraspanin-Halo7-SF650T

particles bound to laminin in considerably higher numbers than they bound to fibronectin and

collagen type I, and the number decreased drastically due to integrin β1 KO (right in Figs. 3B

and S2A and B and Table S1). These findings explicitly reveal that integrin β1 in PC3-derived

sEVs is responsible for the binding of sEVs to laminin and collagen type I.

The surface of sEVs derived from myeloma cells and isolated by ultracentrifugation is

covered by fibronectin, which serves as a heparan sulfate-binding ligand. Consequently, the

enrichment of fibronectin in sEVs marginally (by a factor of approximately 1.2) amplifies the

interactions between sEVs and recipient cells (Purushothaman et al., 2016). Therefore, we

performed immunoprecipitation experiments to examine the presence of fibronectin and/or

laminin on the surface of sEVs derived from PC3 cells (Fig. S3A). However, neither fibronectin

nor laminin was detected in the fraction isolated by the immunoprecipitation of sEVs with the

anti-CD63 antibody (Fig. S3B). Furthermore, we did not detect tetraspanin marker proteins

(CD63, CD81, or CD9) in the fraction obtained through the immunoprecipitation of sEVs using

anti-fibronectin antibody (Fig. S3C). These results unequivocally showed that neither laminin

nor fibronectin was present on the surface of PC3 cell-derived sEVs.

Integrin α2 in tumor-derived sEVs is responsible for binding to collagen type I on glass

The in vitro assay revealed that in all the examined sEV subtypes, integrin β1 was critical for

binding to collagen type I and laminin on glass. Accordingly, we focused on the function of

integrin heterodimers containing the β1 subunit. Since the integrin α2β1 heterodimer is a

receptor for collagen type I, we examined the role of the integrin α2 subunit. Fig. 3C shows

TIRFM images of individual particles of sEV-CD63Halo7-SF650T derived from PC3-WT and

PC3-α2KO cells, which attached to ECM components on glass surfaces. Notably, the expression

levels of other integrin subunits in PC3-α2KO cells were comparable to those in PC3-WT cells

(Fig. 1A). Quantitative analysis revealed a significant decrease in the number of all subtypes of

sEV-tetraspanin-Halo7-SF650T particles derived from α2KO cells bound to collagen type I,

while the number that bound to laminin was not significantly reduced (Figs. 3D and S2C and D

and Table S1). These findings, in conjunction with the results of β1KO, unequivocally show the

crucial involvement of integrin α2β1 heterodimers in the binding of sEVs to collagen type I.

Integrins α6 and β4 in tumor-derived sEVs are responsible for binding to laminin on glass

We next investigated the involvement of the counterpart in β1-containing integrin heterodimers in the binding of sEVs to laminin. As integrin α6β1 is a receptor for laminin, we aimed to determine whether the absence of integrin α6 affects the binding of sEVs to laminin on glass. To this end, we performed a western blotting experiment and detected a lack of the integrin β4 subunit in sEVs derived from PC3-α6KO cells. In addition, we observed a considerable decrease in the expression level of the integrin α6 subunit in EVs derived from β4KO cells (Fig. 1A). Therefore, it is plausible that the expression levels of integrins α6 and β4 in sEVs are closely linked. Fig. 3E and G show TIRFM images of single particles of sEV-CD63Halo7-SF650T derived from PC3-WT, PC3-α6KO, and PC3-β4KO cells, which were attached to ECM components on glass. We discovered that knocking out integrin α6 or integrin β4 considerably impaired the binding of all subtypes of sEV-tetraspanin-SF650T to laminin but did not significantly affect the binding to fibronectin or collagen type I (Figs. 3F and H and S2E-H and Table S1). These findings explicitly show that integrin heterodimers α6β1 and α6β4 in sEVs are responsible for binding to laminin on glass.

Thus, single-particle imaging of PC3 cell-derived sEVs on glass revealed that integrin α2β1 in sEVs were critical for binding to collagen type I, while integrin α6β1 and integrin α6β4

in sEVs were responsible for binding to laminin (Figs. 3 and S2 and Table S1). Interestingly, only small amounts of all the PC3 cell-derived sEVs bound to fibronectin on glass despite the presence of fibronectin receptors such as integrin α5β1, αVβ1 and αVβ5 in the PC3-derived sEVs (Fig. 1A and B). Notably, these fibronectin receptors are functional in PC3 cells, as evidenced by their comparable spread on glass surfaces coated with laminin, fibronectin, or collagen type I (Fig. 2C).

GM1, but not other gangliosides, in sEVs specifically binds to laminin

Our results showed that the knockout of integrin α6, integrin β4, or integrin β1 greatly reduced but did not eliminate the binding affinity of PC3-derived sEVs for laminin, suggesting that other molecules may be involved in the binding of sEVs to laminin. Previous reports have shown that the addition of laminin to dorsal root ganglion (DRG) neurons containing Bodipy-GM1 induces clustering of the GM1 probe, suggesting that laminin may bind directly to GM1 on DRG neurons (Ichikawa et al., 2009). Since our dot blot experiments showed that the relative density of GM1 (Fig. 4A) in sEVs derived from PC3 cells was approximately 29.7 ± 8.1 (mean ± S.E., n=3) times greater than that in PC3 cells (Fig. 4B), GM1 might play a role in the binding of sEVs to laminin. To investigate whether GM1 in sEVs binds directly to laminin, we sought to isolate cell lines that abundantly expressed only one type of ganglioside by overexpressing ganglioside

synthase in cells to produce or deplete gangliosides, as reported previously (Yamashiro et al., 1995; Yesmin et al, 2023). However, this strategy did not work in PC3 cells because the amounts of gangliosides produced or depleted by the synthase returned to the original levels during cell culture. Therefore, we applied this method instead to B78 cells, a subclone of B16 melanoma cells that can predominantly express one type of ganglioside (Fig. S4). The number of sEVs that bound to laminin, normalized by the ratio of the expressed amount of integrin α6/CD81 in the sEVs (Fig. 4C), was approximately 3-fold greater for sEVs derived from B78 cells predominantly expressing GM1 than for sEVs derived from B78 cells expressing GM2, GM3, GD2 and GD2/GD3 (Fig. 4A, D and E). Regardless of which gangliosides were expressed, the number of B78 cell-derived sEVs bound to fibronectin was low (Fig. 4F and G). These results suggest that GM1, but not the other examined gangliosides in sEVs, might bind to laminin and that all the gangliosides in sEVs might not bind to fibronectin.

To further examine whether GM1 binds directly to laminin, we investigated whether liposomes (diameter ≈ 100 nm) containing GM1 bound to laminin but not to fibronectin. Since the molar ratio of GM1 to lipids in PC3 cells is known to be 0.0158 mol% (Llorente et al., 2013) and since the relative density of GM1 in the sEVs derived from PC3 was 29.7 ± 8.1 (mean $\pm$ S.E., n=3) times greater than that in PC3 cells (Fig. 4B), the molar ratio of GM1 in the PC3-derived sEVs was estimated to be 0.47 mol%. Since gangliosides are distributed in both the outer

and inner leaflets of liposomal membranes, the density of GM1 in sEVs is similar to that in the

outer leaflet of liposomes containing 0.94 mol% GM1. Consequently, we used liposomes

containing 1.0 mol% GM1 for the binding assay. Fig. 4H and I demonstrate that liposomes

containing GM1 bound extensively to laminin, while those containing 1.0 mol% other

gangliosides (GM2, GM3, GD1a, GD2, and GD3) showed very little binding to laminin.

Conversely, liposomes hardly bound to fibronectin regardless of which gangliosides they

contained (Fig. 4H and J). These results unequivocally establish that GM1 in sEVs directly binds

to laminin but not to fibronectin.

All the sEVs, mEVs and MVs derived from four tumor cell lines exhibited marked binding

affinity for laminin but minimal binding to fibronectin

To investigate whether the high binding affinity for laminin and the low binding affinity for

fibronectin are general features of tumor-derived sEVs, we conducted in vitro binding assays of

sEV-CD63-Halo7-SF650T particles derived from four other types of tumor cells. These sEVs

were purified by ultracentrifugation (200,000×g for 4 h). The diameters of the sEVs derived from

4175-LuT, BxPC3, HeLa and SKBR3 cells, which were assessed via qNano, were 83 ± 16 nm,

97 ± 9 nm, 83 ± 12 nm and 75 ± 8 nm (mean $\pm$ S.D.), respectively (Fig. 5A). Quantitative

analysis of the TIRFM images (Fig. 5B and C) revealed that the numbers of sEV particles

derived from four tumor cell lines (4175-LuT, PC3, BxPC3, and SKBR3) that bound to laminin

on glass were considerably greater than those that bound to fibronectin on glass (the numbers of

sEVs derived from all the cell lines applied to laminin- or fibronectin-coated glass were adjusted

to the same). On the other hand, the numbers of sEVs derived from HeLa cells that bound to

either fibronectin or laminin were very low (Fig. 5C). These results explicitly demonstrated that

tumor-derived sEVs primarily bind to laminin on glass and exhibit low affinity for fibronectin.

The diameter of EVs isolated by ultracentrifugation depends on factors such as the

sample volume, centrifugation tube, centrifugal acceleration, and centrifugal duration (Théry et

al., 2018). Therefore, to investigate whether the stronger binding affinity of EVs for laminin is a

general phenomenon, we quantified the number of larger EVs bound to ECM components coated

on glass. This experiment was necessary because sEVs and MVs are reported to contain distinct

membrane components (Jeppesen et al., 2019). To achieve this goal, we isolated larger EVs by

ultracentrifugation at slower rates. The mEVs were isolated by low-speed centrifugation

(50,000×g for 30 min) and had diameters of 90 ± 12 nm (4175-LuT), 90 ± 18 nm (PC3), $103 \pm$

13 nm (BxPC3), 97 ± 14 nm (HeLa), and 85 ± 16 nm (SKBR3) (Fig. 5A). Our in vitro binding

assay showed that these mEVs share similar characteristics with sEVs (Fig. 5D) (the numbers of

mEVs derived from all the cell lines applied to laminin- or fibronectin-coated glass were

adjusted to the same). Western blotting (Fig. 5F) revealed that fibronectin receptors, such as

integrin α5β1 heterodimers, were expressed in the EVs derived from all tested types of tumor

cells. In addition, the diameters of the MVs purified by lower-speed centrifugation (10,000×g for

30 min) were 142 ± 14 (4175-LuT), 173 ± 28 (PC3), 138 ± 27 (BxPC3), 145 ± 31 (HeLa), and

212 ± 47 (SKBR3) (Fig. 5A). Since CD63 was not highly concentrated in MVs (Fig. 5F), the

MVs were stained with the lipid probe (ExoSparkler exosome membrane labeling kit-Deep Red,

Dojindo), and the numbers of MVs bound to laminin and fibronectin were quantified by TIRFM

(Fig. 5E). Our results indicate that MVs derived from 4175-LuT, PC3, and SKBR3 cells exhibit

greater binding affinity to laminin than to fibronectin (the numbers of MVs derived from all the

cell lines applied to laminin- or fibronectin-coated glass were adjusted to the same). However,

the numbers of MVs derived from BxPC3 and HeLa cells that bound to laminin and fibronectin

were very low (Fig. 5E). Thus, despite the presence of fibronectin receptors such as integrin

α5β1 and αVβ3 in all sizes of EVs (Fig. 5F), these proteins exhibited minimal binding to

fibronectin. Notably, tumor cells spread on fibronectin- and laminin-coated glass at comparable

rates (Fig. 2C), indicating that all receptors for ECM components on the tumor cell PMs were

functional. These results demonstrate that the laminin receptor is markedly active and the

fibronectin receptor activity is low in EVs derived from four tumor cell lines (4175-LuT, PC3,

BxPC3, and SKBR3).

The integrin β1 and α6 subunits in sEVs are responsible for binding to cell plasma

membranes

Although the in vitro assay showed that integrin heterodimers play critical roles in binding to

laminin and collagen type I on glass (Figs. 3 and S2), it remains unknown whether these integrin

heterodimers also mediate the binding of sEVs to the recipient cell PM. To address this question,

we quantitatively determined the number of sEV-CD63-SF650T particles bound to the PM of the

recipient cell by TIRFM. Immunofluorescence imaging revealed that while human fetal lung

fibroblast (MRC-5) PMs were extensively coated with fibronectin, collagen type I, and laminin,

these coatings were poor in human marrow stromal (HS-5) cells (Fig. 6A and B). After 30 min of

incubation, the number of sEV-CD63-SF650T particles derived from PC3-WT cells that bound

to the PM of MRC-5 cells was substantially greater than the number of sEV-CD63-SF650T

particles derived from PC3-β1KO cells (p = 0.046; two-sided Welch's t-test) or PC3-α6KO cells

(p = 0.0082) (Fig. 6C, E, F and H). In contrast, the number of PC3-β1KO cell-derived sEVs that

bound to HS-5 cell PMs was indistinguishable from that of PC3-WT cell-derived sEVs (p =

0.55) (Fig. 6D and G). These results explicitly indicate that integrin β1 and integrin α6 in sEVs

are involved in binding to the MRC-5 cell PM of which surface is densely covered by ECM

components.

Pseudo real-time super-resolution movie observation demonstrated that sEVs bind

predominantly to laminin on the living cell PM

Having demonstrated the binding of integrin α2β1 in PC3 cell-derived sEVs to collagen type I

on glass and the binding of both integrin α6β1 and integrin α6β4 in sEVs to laminin on glass, we

next examined whether these sEVs also bind to ECM components on the living recipient cell

PM. To directly assess this binding, we performed simultaneous single-particle tracking of PC3

cell-derived sEVs-CD63-Halo7-TMR particles and super-resolution microscopy observation of

ECM components on the living MRC-5 cell PM (Fig. 7A-E). Since both sEVs and ECM

components exhibited slow movement on the cell PM and their positions changed throughout the

observation, we sought to more accurately analyze the colocalization between sEVs and ECM

components by acquiring super-resolution "dSTORM movies" of ECM components instead of

the still images. For this purpose, we acquired dSTORM images by the single-molecule

observation of ECM components immunostained with SF650B-conjugated antibodies at high

speed (200 frames/second) for 3504 frames. To obtain one dSTORM image, data acquisition was

performed for 1002 frames and then repeated by shifting the initial frames backward by 6

frames, resulting in a total of 417 dSTORM images (Fig. 7A). By connecting these dSTORM

image sequences, we generated a pseudo-real-time "dSTORM movie". Images of individual sEV

particles were concurrently recorded at 200 frames/second, and images averaged over 6 frames

were connected to create the movie. The pseudo-real-time movies of the ECM components and

sEV particles (33 frames/second) were superimposed (Fig. 7A).

Conventional immunofluorescence microscopy failed to visualize the fibrillary structure

of ECM components on living cells (left-top panels of Fig. 7B-D). In contrast, live-cell

dSTORM movie observation enabled clear visualization of this structure (left-bottom panels of

Fig. 7B-D). We found that sEVs frequently colocalized with the structure of laminin on the cell

PM, as indicated by the yellow arrowheads in Fig. 7D (right) (Video 3), and that the duration of

colocalization occasionally persisted for longer than 4 seconds (Fig. 7E and Video 4). Moreover,

we observed scarcely any colocalization events between the sEVs and the fibrillary structures of

fibronectin (Fig. 7B, right, and Video 1) or collagen type I (Fig. 7C, right, and Video 2). To

quantitatively analyze the colocalization events, we measured the nearest distance from an edge

of the ECM structure to a centroid of the sEV spot and performed this measurement for all pairs

of ECM structures and sEV particles. The data were normalized using an in silico distribution of

randomized sEV spots, and subsequently, we obtained histograms showing the distribution of the

normalized relative frequency of sEVs at each distance from the edge of the ECM structures

(Fig. 7F). Our analysis revealed that sEVs were significantly enriched near the edges of laminin

structures at distances ranging from -100 to 150 nm on the PMs of living MRC-5 cells (Fig. 7G,

bottom). In contrast, we did not observe the presence of sEVs near fibronectin or collagen type I

structures on the MRC-5 cell PM (Fig. 7G, top and middle, respectively). Our results indicate

that PC3 cell-derived sEVs adhere predominantly to laminin on the PMs of recipient cells.

Notably, we did not observe any binding of sEVs to collagen type I on the cell membrane,

although sEVs did bind to collagen I on glass (Figs. 2A-D and S2A-D). This difference may be

attributed to the considerably lower binding of sEVs to collagen type I than to laminin, as

demonstrated in the glass assay (Figs. 3 and S2).

Talin-1 does not facilitate the binding of integrin heterodimers in EVs to laminin

Talin binds to the cytoplasmic tail of integrin β subunits and thereby disrupts the transmembrane

association of integrin heterodimers, induces extended-open conformations, and promotes the

binding of integrin heterodimers to all ECM components (Sun et al., 2019). Talin-1, but not talin-

2, is an important factor for activating integrin β1 in prostate cancer (PCa) cells, which facilitates

bone metastasis (Jin et al., 2015). The phosphorylation of talin-1 at Ser 425 promotes the activity

of integrin β1, accelerating metastasis (Jin et al., 2015). Furthermore, Cdk5 kinase activity is

responsible for talin-1 phosphorylation at Ser 425 (Huang et al., 2009), which subsequently

triggers integrin β1 activation (Jin et al., 2015). Moreover, the present study showed that the

presence of fibronectin receptors in EVs does not correspond to binding with fibronectin.

Therefore, we hypothesized that factors other than the presence of ECM receptors are involved

in regulating the binding of EVs with ECM components. We investigated whether talin-1 and its phosphorylation at Ser 425 play pivotal roles in the activation of integrin heterodimers, which are receptors for ECM components in sEVs.

sEVs derived from talin-1 knockdown (KD) PC3 cells were isolated by ultracentrifugation, and we quantitatively analyzed the numbers of sEV-Halo7-integrinβ1-SF650T and sEV-CD63-Halo7-SF650T particles that bound to ECM components on glass (Fig. 8). The expression level of talin-1 in sEVs was reduced to 40% following talin-1 KD (Fig. 8A, top). Notably, we observed that the binding of talin-1 KD cell-derived sEVs to laminin was similar to that of sEVs derived from intact cells (Fig. 8B, top; 8C; 8E, top; 8F). Furthermore, overexpression (OE) of talin-1 in sEVs (2.3 times greater than the endogenous level; Fig. 8A, bottom) did not alter the number of sEVs bound to laminin (Fig. 8B, bottom; 8D; 8E, bottom; 8G). These results suggest that the amount of talin-1 in sEVs is irrelevant to the binding of sEVs to laminin on glass.

Subsequently, we investigated whether the phosphorylation of talin-1 at Ser 425 promotes the binding of integrin heterodimers in sEVs to ECM components. Western blotting revealed that talin-1 was highly phosphorylated at Ser 425 in PC3 cells, but this phosphorylation was attenuated by treatment with roscovitine, an inhibitor of Cdk5 (Fig. 8H). Moreover, talin-1 at Ser 425 was almost completely dephosphorylated in sEVs (Fig. 8H), suggesting that the

phosphorylation of talin-1 at Ser 425 does not play an important role in the binding of sEVs to

ECM components. Combined with the talin-1 KD and OE experiments, these results strongly

indicate that the presence of talin-1 in sEVs does not increase the binding of integrin

heterodimers to laminin. Moreover, the diminished binding of integrin heterodimers to

fibronectin in EVs might stem from the absence of talin-1 functionality.

Cholesterol attenuates the binding of sEVs to laminin and the PMs of recipient cells

Given that our results demonstrate that talin-1 does not participate in the activation of integrin in

sEVs (Fig. 8), we aimed to identify alternative factors that specifically maintain integrin binding

to laminin even in sEVs where inside-out signaling is lacking. Previous reports have shown that

focal adhesions are highly ordered structures similar to rafts (Gaus et al., 2006) and that some

integrin subunits, such as α6 and β4, undergo palmitoylation and partition into rafts (Gagnoux-

Palacios et al., 2003; Yang et al., 2004). Moreover, a recent study demonstrated that sEV

membranes are richer in rafts than the PMs of parental cells (Yasuda et al., 2022). Therefore, we

hypothesized that rafts in sEV membranes enhance the binding of sEVs to laminin. To test this

possibility, we used TIRFM to observe the binding of sEV-CD63-Halo7-SF650T to laminin on

glass after the depletion or addition of cholesterol. The cholesterol content in PC3-derived sEVs

was reduced to 16% after treatment with MβCD and increased to 186% after treatment with the

MβCD-cholesterol complex. After cholesterol depletion by MβCD or saponin, the number of

sEVs that bound to laminin on glass significantly increased by factors of 1.7 and 1.3,

respectively ($p < 0.01$) (Fig. 9A and B). Conversely, after cholesterol was added to sEVs with the

MβCD-cholesterol complex, the binding affinity was significantly reduced by approximately

20% ($p < 0.01$) (Fig. 9 C). Furthermore, we quantified the binding affinity of cholesterol-

depleted and cholesterol-supplemented sEVs for MRC-5 cell PMs. In agreement with the

findings on glass, cholesterol depletion from sEVs significantly increased the number of sEVs

that bound to the recipient cell PM after 30 min of incubation ($p = 0.039$), whereas the addition

of cholesterol markedly reduced this binding ($p = 0.0059$) (Fig. 9D-F). Thus, these results

explicitly show that, in contrast to our hypothesis, the presence of abundant cholesterol in sEV

membranes impedes the binding of sEVs to both laminin and the recipient cell PM.

CD151 preserves the binding of integrin heterodimers in sEVs to laminin

Because the presence of large amounts of cholesterol inhibits the activity of laminin receptors in

sEVs, we investigated the mechanism by which integrin heterodimers in sEVs lacking inside-out

signaling maintain binding to laminin. CD151 is a ubiquitously expressed tetraspanin protein,

and its large outer loop (Kazarov et al., 2002) specifically interacts with the C-terminal domain

of the extracellular region of the integrin α7, α6, and α3 subunits, selectively strengthening the

binding of integrin heterodimers such as α6β1 and α6β4 to laminin (Lammerding et al., 2003; Winterwood et al., 2006). CD151 has a cholesterol-binding domain (Purushothaman and Thiruvenkatam, 2019), is palmitoylated (Yang et al., 2002), and associates with many other membrane proteins through raft-lipid interactions (Berditchevski, 2001; Charrin et al., 2003; Odintsova et al., 2006). Therefore, we investigated whether the binding activities of integrin heterodimers in sEVs lacking inside-out signaling to laminin are preserved by CD151. After CD151 KD by siRNA (Fig. 9G), there was no change in the levels of integrin α6 or integrin α3 in PC3 cells. However, these cells exhibited a significantly reduced ability to spread on laminin, whereas their ability to spread on fibronectin and collagen type I remained unaltered (Fig. 9H). CD151 KD also decreased the number of sEVs that bound to laminin by 41% (Fig. 9I), although the amounts of integrin α6 and α3 in the sEVs did not change (Fig. 9G). Furthermore, CD151 KD attenuated the marked increase in the laminin-binding activity of sEVs caused by cholesterol depletion (Fig. 9I and J). These results suggest that the presence of cholesterol in sEVs suppresses sEV binding to laminin through a CD151-dependent mechanism. In other words, an unidentified molecule that interacts with CD151 in a cholesterol-dependent manner may inhibit the binding of integrin heterodimers in sEVs to laminin.

Discussion

Our results directly demonstrated that laminin, rather than fibronectin, is the predominant

binding target of tumor-derived sEV subtypes containing CD63, CD81 or CD9, and this binding

is mediated by specific interactions between the integrin α6β1 and integrin α6β4 heterodimers in

sEVs and laminin on both glass (Figs. 3 and S2) and recipient cell PMs (Figs. 6, 7 and 10A).

Furthermore, much greater binding affinities for laminin than for fibronectin were detected for

both mEVs and MVs derived from four different types of tumor cells (Fig. 5C-E), although these

EVs contained receptors for both laminin and fibronectin (Figs. 1A and 5F), and the activities of

all the ECM receptors were comparable in all the examined cells that secreted these EVs (Fig. 2).

Therefore, we surmised that factors other than the presence of these ECM receptors in EVs

contribute significantly to determining their binding activities to ECM components. Our results

showed that inside-out signaling by talin-1 does not increase integrin activity in EVs (Fig. 8);

however, the tetraspanin CD151, which interacts with integrins α6 and α3, considerably

increases the binding of integrin heterodimers in EVs to laminin (Fig. 9G-J).

Although the inhibition of EV binding to recipient cells by antibodies and ligands against

integrins has suggested that integrin subunits in EVs are responsible for the binding of EVs to

recipient cells (Nazarenko et al., 2010; Carney et al., 2017; Zhang et al., 2022), there has been no

direct evidence of the specific binding of integrins to ECM components. The binding affinities of

EVs for fibronectin, laminin, and collagen type I have not been quantitatively compared. In this

study, we established an in vitro experimental system to quantitatively measure the binding

affinities of integrin heterodimers with ECM components, which enabled us to compare the

binding affinities of EVs derived from different types of tumor cells with fibronectin, laminin,

and collagen type I. The strong binding of all the sEVs, mEVs and MVs to laminin via integrin

heterodimers and the weak binding to fibronectin were unexpected because previous studies

reported that the binding of EVs to recipient cells was inhibited by antibodies and inhibitors

against fibronectin receptors (Huang et al., 2016; Altei et al., 2020). However, since our results

did not indicate that tumor-derived EVs have no binding ability to fibronectin at all but rather

that they exhibit lower binding to fibronectin than to laminin, our results are not inconsistent

with previously reported findings. Since the amount of EVs bound to PMs increases with the

amount of EVs added to cells, differences in the numbers of EVs bound to PMs after treatment

with antibodies against integrin subunits of fibronectin receptors may be detected in previous

reports.

Our results also revealed that only GM1, which is approximately 29-fold more enriched

in sEVs compared to sEV-secreting cells, bound to laminin on glass, while all the other examined

gangliosides did not (Fig. 4). Since GM2 has a structure in which N-acetylgalactosamine

(GalNAc) is bound to the C4 position of galactose (Gal) of GM3 and GM1 has an additional Gal

bound to the C3 position of the GalNAc of GM2 (Fig. 4A), these results indicate that the

additional GalNAc and Gal residues are necessary for the effective binding of ganglioside to

laminin. Furthermore, for the binding of ganglioside to laminin, sialic acid linked to the C3

position of the terminal Gal of GM1 should not be important, because GD1a did not bind to

laminin (Fig. 4A, H, and I). Furthermore, none of the examined gangliosides, including GM1,

bound to fibronectin. These results suggest that gangliosides may bind to laminin through

specific glycan–protein interactions, not merely through electrostatic interactions. Since the

binding of the glycan structure of GM1 to laminin does not require complicated regulation by

other membrane molecules, it must constitute a robust binding system that is not prone to loss of

activity unless GM1 is degraded.

Neither the knockdown nor the overexpression of talin-1 in sEVs significantly altered the

binding of sEVs to laminin (Fig. 8). In addition, Ser 425 of talin-1, an activating phosphorylation

site (Jin et al., 2015), was almost completely dephosphorylated, whereas Ser 425 was

phosphorylated in cells that secreted sEVs. Talin-1 is ubiquitously expressed in all cell types in

all tissues, whereas talin-2 is expressed mostly in the brain, heart muscle, and kidney (Monkley

et al., 2001; Thul et al., 2017) and is not expressed in the PC3 cells we examined. These results

explicitly demonstrate that inside-out signaling by integrins is absent in sEVs and that integrins

in sEVs exist in a completely different environment from that in cell PMs. The lack of inside-out

integrin signaling in sEVs seems reasonable because there is no cytoskeletal actin filament

network inside the sEVs, which might be one of the reasons that integrin heterodimers of

fibronectin receptors exhibit relatively low activity in sEVs (Fig. 10B). On the other hand, our

results indicate that α6β1 and α6β4 integrin heterodimers of laminin receptors in sEVs are still

active (Figs. 3, 6, 7 and S2), and a tetraspanin, CD151, preserves the activities of these receptors

(Fig. 9I). CD151 is known to form a complex with integrin early in biosynthesis; this complex is

delivered to tetraspanin-enriched microdomains (TEMs) in the cell PM (Stipp, 2010) and

selectively strengthens integrin-mediated adhesion to laminin (Lammerding et al., 2003). CD151

has a cholesterol-binding domain (Purushothaman and Thiruvenkatam, 2019) and links integrin

heterodimers of laminin receptors with other tetraspanins by cholesterol- and raftophilic lipid-

dependent mechanisms (Berditchevski et al., 2002; Franco et al., 2010). Our results revealed that

cholesterol suppressed the binding of integrin heterodimers in sEVs to laminin in a manner that

was dependent on the presence of CD151 on sEVs (Fig. 9G-J), suggesting that the raft-dependent

interaction of membrane component(s) X with a molecular complex of integrin α6β1, α6β4 and

CD151 may inhibit the binding of integrins in sEV membranes to laminin (Fig. 10C). This

finding is consistent with the notion that the depalmitoylation of CD151 increases the

downstream signaling of integrins in the cell PM (Berditchevski et al., 2002). Since the PC3-

derived sEV membrane possesses a greater abundance of the Lo-like phase than does the cell PM

(Yasuda et al., 2022), the binding of integrins in EVs to laminin would be attenuated by the

membrane component(s) X, which should be identified in future research. Tetraspanin proteins are candidate molecules because CD151 associates with tetraspanin-enriched microdomains (TEMs) in a cholesterol-dependent manner (Berditchevski et al., 2001).

The higher binding affinity of tumor-derived sEVs for laminin than for fibronectin on recipient cell PMs may also influence the response of the recipient cells. For example, embryonic neural crest cells (NCCs) were reported to exhibit both ligand- and receptor-specific integrin regulation (Strachan and Condic, 2004). On laminin-coated glass, NCCs accumulate internalized laminin receptors (integrin α6) but not fibronectin receptors (integrin α5). Furthermore, internalized laminin receptors colocalize with receptor-recycling vesicles and are subsequently recycled to the cell surface, whereas fibronectin receptors do not colocalize with receptor-recycling vesicles. Blocking receptor recycling inhibited NCC motility on laminin. Therefore, these results indicate that internalized laminin receptors can be recycled on laminin, which is essential for rapid cell migration. Recently, our dual-color PALM movies and single-particle observations of sEVs demonstrated that after sEVs were bound to laminin on recipient cells, integrin subunits accumulated underneath the sEVs, which were then internalized via caveolae and phagocytosis. Surprisingly, approximately 20-40% of PC3 cell-derived sEVs accumulate in recycling endosomes (Hirosawa et al., 2024). Therefore, the specific binding of sEVs to laminin on recipient cell PMs may promote the efficient recycling of internalized integrins.

Another example of a specific response in recipient cells after integrin-laminin binding involves signaling pathways that are different from those induced by fibronectin binding. It has been reported that laminin-10/11 (which is the same as laminin 511 used in this study) preferentially activates Rac via p130Cas-CrkII-DOCK180, promoting cell migration, while fibronectin activates Rho via FAK, decreasing cell migration (Gu et al., 2001). Since we recently elucidated that the binding of sEVs to the PMs of recipient cells triggers prompt signal transduction underneath sEVs through the activation of integrin, talin-1, and Src family kinases (Hirosawa et al., 2024), the results of the present study suggest that laminin-specific signaling might occur upon sEV binding. The biological importance of the selective binding of sEVs to laminin but not to fibronectin on the PMs of recipient cells warrants further investigation.

In summary, using single-particle tracking and super-resolution microscopy, we showed definitively that tumor-derived sEVs bound to laminin via integrin $\alpha6\beta1$ and $\alpha6\beta4$ while far fewer EVs attached to fibronectin than to laminin despite the abundant presence of fibronectin receptors in sEVs. This difference may stem from the inability of talin-1 in sEVs to activate the binding of integrin heterodimers. Conversely, CD151 promoted the function of laminin receptors, and a raftophilic molecule X may inhibit laminin receptors on the TEM formed by CD151 and cholesterol in sEVs. GM1 in sEVs directly bound to laminin but other gangliosides did not, while none of the examined gangliosides in sEVs bound to fibronectin. Thus, our

findings demonstrate that sEVs bind to laminin through robust mechanisms that do not require inside-out integrin signaling. The mechanism of EV binding to laminin revealed in this study will also provide useful information for applied research, including clinical applications.

Materials and methods

Materials

The human prostate cancer cell line (PC3), human breast cancer cell line (SKBR3), human pancreas cancer cell line (BxPC-3), human fetal lung fibroblast line (MRC-5), and human marrow stromal cell line (HS-5) were purchased from the American Type Culture Collection (ATCC). The human breast cancer cell line (4175-LuT) was kindly provided by Dr. Joan Massagué and Dr. Hoshino (Hoshino et al., 2015).

All antibodies were obtained from commercial sources. The following primary antibodies were used for western blotting: mouse monoclonal anti-beta actin 15G5A11/E2 (1:5000, Thermo Fisher, Cat# MA1-140), mouse monoclonal anti-CD63 8A12 (1:1000, Cosmo Bio, Cat# SHI-EXO-M02), mouse monoclonal anti-CD81 B11 (1:500, Santa Cruz Biotechnology, Cat# SC166029), rabbit monoclonal anti-CD9 EPR23105-125 (1:1000, Abcam, Cat# ab263019), mouse monoclonal anti-CD151 11G5a (1:250, Abcam, Cat# ab33315), rabbit polyclonal anti-

fibronectin (1:1000, Sigma–Aldrich, Cat# F3648), rabbit monoclonal anti-GAPDH 14C10 (1:2000, Cell Signaling, Cat# 2181S), rabbit monoclonal anti-integrin alpha 2 EPR5788 (1:2500, Abcam, Cat# ab133557), rabbit polyclonal anti-integrin alpha 3 (1:500, Abcam, Cat# ab190731), rabbit polyclonal anti-integrin alpha 5 (1:250, Cell Signaling, Cat# 4705S), rabbit polyclonal anti-integrin alpha 6 (1:500, Cell Signaling, Cat# 3750S), rabbit polyclonal anti-integrin alpha 7 (1:500, Abcam, Cat# ab203254), rabbit polyclonal anti-integrin alpha V (1:500, Abcam, Cat# ab117611), mouse monoclonal anti-CD29 Clone 18 (1:2000, BD Bioscience, Cat# 610467), rabbit polyclonal anti-integrin beta 3 (1:500, Millipore, Cat# AB2984), rabbit polyclonal anti-integrin beta 4 (1:500, Cell Signaling, Cat# 4707S), rabbit polyclonal anti-integrin beta 5 (1:500, Cell Signaling, Cat# 4708S), rabbit polyclonal anti-laminin 1+2 (1:1000, Abcam, Cat# ab7463), mouse monoclonal anti-talin-1 97H6 (1:500, Gene Tex, Cat# GTX38972), rabbit polyclonal anti-phospho-talin (Ser-425), (1:250, ECM Biosciences, Cat# TP4171), and anti-GM3 antibody clone GMR6 (1:1000, Tokyo Chemical Industry, Cat# A2582), along with biotin-conjugated Cholera toxin B subunit (1:500, List Labs, Cat# 112). Additionally, goat anti-mouse IgG-HRP (1:5000, Millipore, Cat# 12-349), donkey anti-rabbit IgG-HRP (1:4000, Cytiva, Cat# NA934), goat anti-rabbit IgG-HRP (1:10000, Sigma–Aldrich, Cat# A0545), goat anti-mouse IgM antibody-HRP (1:10000, Invitrogen, Cat# 31440), and rabbit anti-biotin antibody-HRP clone D5A7 (1:10000, Cell Signaling, Cat# 5571) were used as secondary antibodies. For immunoprecipitation, mouse

monoclonal IgG2a negative control C1.18.4 (Millipore, Cat# MABF1080Z), mouse monoclonal

IgG1-κ negative control MOPC-21 (Millipore, Cat# MABF1081Z), and rabbit polyclonal IgG

negative control (Abcam, Cat# ab37415) were used. For the observation of EVs and ECM

components, mouse monoclonal anti-CD81 JS-81 (BD Pharmingen, Cat# 555675), mouse

monoclonal anti-CD9 M-L13 (BD Pharmingen, Cat# 555370), rabbit polyclonal anti-collagen

type 1 (Abcam, Cat# ab34710), rabbit polyclonal anti-fibronectin (Abcam, Cat# ab23750), rabbit

polyclonal anti-laminin (Sigma–Aldrich, Cat# L9393), goat polyclonal anti-rabbit IgG-

rhodamine (Cappel, Cat# 55666), and goat anti-rabbit IgG (Cappel, Cat# 0212-0081) were used.

We purchased the following purified ECM proteins: fibronectin from Roche (Cat#

11051407001), laminin from Sigma–Aldrich (Cat# L6274), and collagen type I from Sigma–

Aldrich (Cat# C7774). For the experiments involving cholesterol and CD151, we used laminin-

511 (Biolaminin 511 LN, BioLamina). The recombinant laminin (L6274, Sigma–Aldrich) we

used was almost identical to Biolaminin 511 LN because nearly all the laminins in L6274 were

laminin-511 (Wondimu et al., 2006).

Cell culture

Mycoplasma contamination was not detected in any of the cell lines used in this study.

PC3 human prostate cancer cells (ATCC, CRL-1435) were cultured in HAM F12 medium

(Sigma–Aldrich) supplemented with 10% fetal bovine serum (FBS) (Gibco), 100 U/ml

penicillin, and 100 μg/ml streptomycin (Gibco). Human fetal lung fibroblasts (MRC-5; ATCC,

CCL-171) and human uterine cervix cancer cells (HeLa) were cultured in minimum essential

medium (Gibco) supplemented with 10% FBS, 100 U/ml penicillin, and 100 μg/ml streptomycin.

Human marrow stromal cells (HS-5; ATCC, CRL-11882) and human breast cancer cells (4175-

LuT; Hoshino et al., 2015) were cultured in Dulbecco's modified Eagle's medium (Sigma–

Aldrich) supplemented with 10% FBS, 100 U/ml penicillin, and 100 μg/ml streptomycin. Human

breast cancer cells (SKBR3; ATCC, HTB-30) were cultured in McCoy's 5a medium (Gibco)

supplemented with 10% FBS, 100 U/ml penicillin, and 100 μg/ml streptomycin. Human

pancreatic cancer cells (BxPC-3; ATCC, CRL-1687) were cultured in RPMI-1640 medium

(Sigma–Aldrich) supplemented with 10% FBS, 100 U/ml penicillin, and 100 μg/ml

streptomycin.

The mouse melanoma cell line B78 was cultured in Dulbecco's modified Eagle's medium

(Sigma–Aldrich) supplemented with 10% FBS, 100 U/ml penicillin, 100 μg/ml streptomycin and

400 μg/ml G418. B78 cells expressed only GM3 (Fig. S4). pM2T1-1 cDNA transfection induced

the overexpression of GM2/GD2 synthase, resulting in GM3 depletion because this enzyme uses

GM3 as a substrate for synthesizing GM2 (Fig. S4). The transfection of B78 cells with pM1T-9

and pM2T1-1 induced the overexpression of GM1 (Fig. S4), resulting in GM3 and GM2

depletion because these enzymes use GM3 and GM2 as substrates for the synthesis of GM1.

Transfection of B78 cells with pM2T1-1 and subsequently with pD3T31 induced overexpression of GD2, while transfection with pD3T31 and subsequently with pM2T1-1 induced overexpression of both GD2 and GD3 (Fig. S4) (Yesmin et al., 2023).

Flow cytometry of cells expressing gangliosides

The expression of gangliosides on the cell surface was analyzed by flow cytometry with a FACSCalibur (Becton-Dickison) as previously reported (Yamashiro et al., 1995; Yesmin et al., 2023). Intact and glycosyltransferase-transfected B78 cells were stained by incubation with monoclonal antibodies against GM3 (M2590, IgM) [Cosmo Bio], GM2 (10-11, IgM), GD2 (220-51, IgG1), GD3 (R24, IgG3) [10-11 and R24 antibodies kindly provided by P. O. Livingston and L. J. Old at Memorial Sloan-Kettering Cancer Center], GD1a (D-266, IgM), or GD1b (370, IgM); FITC-labeled goat anti-mouse IgM antibodies (Cappel); or goat anti-mouse IgG (H & L chain) antibodies (Cappel). Anti-GD2, anti-GD1a, and anti-GD1b antibodies were generated by K. Furukawa. Alternatively, GM1 on the cells was stained by applying biotinylated cholera toxin subunit B (CTXB; Sigma–Aldrich) and then FITC-labeled avidin (Zymed).

Knockout of integrin subunits

The integrin β1, β4, α2, and α6 subunits were knocked out in PC3 cells via CRISPR–Cas9 gene editing. The single guide RNA (sgRNA) target sequence was selected via the online tool CHOPCHOP (https://chopchop.cbu.uib.no/). The sgRNA target sequences were subsequently incorporated into the multicloning site of pSpCas(BB)-2A-Puro (Addgene) using BbsI-HF (New England Biolabs). The sgRNA target sequences used were TCATCACATCGTGCAGAAGT and ATACAAGCAGGGCCAAATTG for the knockout of integrin β1; GTGCAGGTTCTCGTATCCCT and GCGCTCAGTCAAGGTAAGCG for the knockout of integrin α2; CCGGGTACTTATAACTGGAA and ACACCGCCCAAAGATGTCTC for the knockout of integrin α6; and CTGCGAGATCAACTACTCGG for the knockout of integrin β4. PC3 cells were transfected with plasmids with a 4D-Nucleofector (LONZA). The transfected cells were selected and cloned using 1 μg/ml puromycin (final concentration) to achieve the knockout of integrin β1 or α2. For the knockout of integrin α6 or β4, the transfected cells were sorted by flow cytometry (BD, FACSMelody) using a PE-conjugated anti-integrin α6 antibody (Invitrogen) or a PE-conjugated anti-integrin β4 antibody (BD). The knockout of each integrin subunit was validated by western blotting.

Preparation of cDNA plasmids and transfection

Wild-type PC3 cells and integrin KO PC3 cells were transfected with 1500 ng of cDNA

encoding CD63-Halo7, CD81-Halo7, or CD9-Halo7 using a 4D-nucleofector (LONZA)

according to the manufacturer's recommendations. cDNA plasmids for human CD63

(NM_001780), human CD81 (NM_004356 and pFN21AE5228) and human CD9 (NM_001769

and pFN21AE1546) were purchased from Kazusa DNA Res. Inst. The DNA sequences encoding

CD63, CD81, or CD9 were cloned and inserted into the pEGFP-N1 vector, and the EGFP

sequence was replaced with the Halo7 sequence (Promega). The linker sequence 5'-

ACCGGTGGTGGGCGCGCCTCTGGTGGCGGATCCGGGGGT-3' was incorporated between

CD63, CD81, CD9 and Halo7. Cells stably expressing the molecules of interest were selected

using a flow cytometer (BD, FACSMelody) and were cloned by treatment with 0.2 mg/ml (final

concentration) G418 (Nacalai Tesque). To generate an expression vector for Halo7-integrin β1,

cDNA encoding Halo7-integrin β1 was cloned and inserted into the PiggyBac transposon vector.

Halo7 was inserted into the hybrid domain of human integrin β1 (NM_002211) between residues

Gly101 and Tyr102. The tagging of integrin β1 with Halo7 at this position is known to have no

detrimental effects on integrin activity (Huet-Calderwood et al., 2017). Wild-type or integrin β1

KO PC3 cells were cotransfected with cDNA encoding Halo7 integrin β1 and transposase,

yielding cell lines that exhibited stable expression of Halo7 integrin β1 (Yusa et al., 2009; Nagai

et al., 2021). Cells exhibiting high expression of Halo7-integrin β1 were selected by flow

cytometry.

4175-LuT, PC3, BxPC3, HeLa, and SKBR3 cells were transiently transfected with a plasmid encoding CD63-Halo7 using polyethylenimine (PEI) MAX (Polysciences) to isolate tumor-derived sEVs containing CD63.

To obtain PC3 cells overexpressing human talin-1, PC3 cells were transfected with cDNA encoding human talin-1 (NM_006289) using Lipofectamine 3000 reagent (Invitrogen) according to the manufacturer's recommendation. cDNA encoding talin-1 was cloned and inserted into the pEGFP-N1 vector, after which the EGFP sequence was deleted. Moreover, talin-1 was knocked down by transfecting PC3 cells with talin-1 siRNA (Santa Cruz Biotech. sc-36610). siRNA-A (Santa Cruz Biotech. sc-37007) was used as a negative control. CD151 was knocked down by transfecting PC3 cells with CD151 siRNA (Dharmacon, D-003637-04 and D-003637-13) using Lipofectamine 3000.

Isolation of small extracellular vesicles (sEVs) from cell culture supernatant

Cells that secreted sEVs were cultured either in two 100-mm dishes for observing sEVs on glass or in two 150-mm dishes for western blotting analysis and the observation of sEVs on cell PMs. The cells were grown to approximately 80% confluence ($0.3\text{-}1.6\times10^8$ cells/150-mm dish). These cells were then transfected with plasmids or siRNAs and cultured for 1-2 days. Next, the cell culture medium in the 100-mm or 150-mm dishes was replaced with 10 ml or 30 ml of FBS-free

medium, respectively. After 48 h of incubation, the cell culture supernatant was collected. For

roscovitine treatment, cells were incubated with 20 μM roscovitine (LKT Laboratories) in FBS-

free medium. The collected supernatant was then subjected to centrifugation at 300×g for 10 min

at 4°C to remove cells. Subsequently, the supernatant was further centrifuged at 2,000×g and 4°C

for 10 min to remove sediment and eliminate apoptotic bodies. Then, the supernatant was

centrifuged at 10,000×g and 4°C for 30 min (himac CF16RN, T9A31 angle rotor, 50 ml Falcon

tube) to pellet the microvesicles (MVs). Finally, the supernatant was concentrated by

ultrafiltration using an Amicon Ultra 15 100K (Millipore) or a Centricon Plus 70 100K

(Millipore) filter unit. The collected sEVs were incubated with 50-100 nM (final concentration)

HaloTag TMR ligand (Promega) or HaloTag SaraFluor650T (SF650T) ligand (Goryo Chemical)

for 1 h at 37°C. The sEVs were then pelleted by ultracentrifugation at 200,000×g and 4°C for 4 h

(himac CS100FNX with an S55A2 angle rotor and S308892A microtubes for

ultracentrifugation). Alternatively, medium-sized EVs (mEVs) were pelleted by

ultracentrifugation at 50,000×g and 4°C for 30 min. The resulting pellet was resuspended in

Hank's balanced salt solution (HBSS) for microscopic observation. For western blotting, the

pellet was suspended in RIPA buffer containing protease inhibitor cocktail set III (Millipore).

The ExoSparkler Exosome Membrane Labeling Kit-Deep Red (Dojindo) was also

utilized as a fluorescent probe. Microvesicles (MVs) purified by lower-speed centrifugation

(10,000×g for 30 min) were successfully labeled according to the Exosparkler Technical Manual.

Subsequently, the labeled MVs were purified by ultrafiltration using an Amicon Ultra 0.5 100K

filter unit (Millipore).

Partial cholesterol depletion/addition in sEVs

sEVs were isolated from the cell culture supernatant of two 150-mm dishes using

ultracentrifugation. To deplete cholesterol, the sEVs were incubated with 5 mM methyl-beta-

cyclodextrin (MβCD) (Sigma–Aldrich) overnight at room temperature with gentle shaking or

incubated with 200 μM saponin (Kanto Chemical) for 1 h on ice. These solutions were then

concentrated by ultrafiltration using Amicon Ultra 2 ml 100K. The concentrated sEVs were

incubated with 50-100 nM (final concentration) HaloTag SF650T ligand for 1 h at 37°C. Next,

the sEVs were pelleted by ultracentrifugation at 200,000×g and 4°C for 4 h. The pellet was

resuspended in phosphate-buffered saline (PBS) or RIPA buffer. To add cholesterol, sEVs were

incubated with 3.5 mM MβCD-cholesterol complex (1:1) for 2 h at 37°C, then isolated by the

same method used in cholesterol depletion. The amount of cholesterol in these sEVs was

measured by a LabAssay Cholesterol Kit (Fujifilm Wako Pure Chemical).

Immunoprecipitation of sEVs

The initial step involved treating 0.6 mg of Protein G-conjugated Dynabeads (Invitrogen) with 200 μl of 15 μg/ml anti-CD63 antibody (Cosmo Bio, SHI-EXO-M02) or mouse IgG2a isotype control (Millipore, MABF1080Z) in PBS containing 0.02% Tween 20. Alternatively, 0.6 mg of Protein A-conjugated Dynabeads (Invitrogen) was treated with 200 μl of 15 μg/ml anti-fibronectin antibody (Sigma–Aldrich, F3648) or rabbit IgG isotype control (Abcam, ab172730) in PBS containing 0.02% Tween 20. The mixture was incubated with rotation for 10 min at room temperature, and then the supernatant was removed. The beads were washed once with PBS containing 0.02% Tween 20 and twice with 20 mM sodium phosphate (pH 7.0) and 150 mM NaCl using a magnetic rack. Then, the antibody-conjugated beads were resuspended in 5 mM BS3, 20 mM sodium phosphate (pH 7.0), and 150 mM NaCl for crosslinking between the antibody and the beads. After incubation with rotation for 10 min at room temperature, 12.5 μl of 1 M Tris-HCl (pH 7.5) was added to the solution, which was then incubated with rotation for an additional 15 min at room temperature. Subsequently, the supernatant was removed, and the beads were washed once with PBS containing 0.02% Tween 20 and then twice with PBS using a magnetic rack.

The antibody-conjugated beads were mixed with 300 μl of PC3-derived sEVs (0.1 mg/ml protein concentration) in PBS. After the mixture was incubated with rotation overnight, the supernatant was removed, and the beads were washed three times with PBS. Then, the beads

were resuspended in Laemmle's SDS sample buffer and incubated at 95°C for 5 min. The

supernatant was collected using a magnetic rack and analyzed by western blotting.

Transmission electron microscopy of sEVs after negative staining

A copper grid with a carbon-coated acetylcellulose film (EM Japan) was washed briefly (~1

min). Ten microliters of water was placed on the grid and then blotted to eliminate the excess

liquid. Next, 5 µl of the sEV sample was pipetted onto the grid and incubated for 1 min. The

excess liquid was then removed by blotting. The grid was then briefly (~45 sec) treated with 5 µl

of 2% phosphor tungstic acid (TAAB Laboratory and Microscopy), followed by blotting to

remove excess liquid. Next, the grid was air-dried at room temperature for 3 days in a desiccator

containing silica gel desiccant and subsequently subjected to TEM observation.

sEV images were obtained using a single TEM instrument (JEM-2100F, JEOL) at 200

kV. These images were produced by computing the mean of 3-second acquisitions captured by a

side-mounted CCD camera (Gatan) and processed by image solution software (Gatan Digital

Micrography).

Determination of sEV size by qNano

The diameter of the suspended sEVs in HBSS was analyzed using a tunable resistive pulse

sensing instrument, qNano (Izon Science) (Coumans et al., 2014), according to the

manufacturer's protocol. Izon Control Suite 3.3 (Izon Science) was used for the data analysis.

The measurements were performed with an NP100 pore (particle detection range: 40-320 nm)

equipped with a stretch of 47.00 mm, a voltage of 1.2 V, and a pressure of 0.8 kPa; an NP150

pore with a stretch of 46.85 mm, a voltage of 1.2 V, and a pressure of 1.4 kPa; or an NP200 with

a stretch of 48.00 mm, a voltage of 1.4 V, and a pressure of 1.4 kPa. The samples were calibrated

using 70, 95, and 210 nm polystyrene calibration beads (Izom Science, CPC100).

Western blotting analysis

The quantities of integrin subunits were evaluated in wild-type and integrin-KO PC3 cells, as

well as in sEVs derived from these cells, by western blotting. The cells and sEVs were

suspended in RIPA buffer containing protease inhibitor cocktail set III (Millipore). The protein

concentrations of the samples were determined using a Thermo BCA protein assay kit (Thermo

Fisher Scientific) and adjusted to 1 mg/ml for SDS–PAGE. Then, 5× concentrated Laemmli's

SDS sample buffer was added to the samples, and the mixture was incubated at 95°C for 5 min

in a blocking incubator. Next, 10 µl of this mixture was loaded into the lanes of a precast 4-12%

gradient polyacrylamide gel (4–12% Bolt Bis-Tris Plus Gels, Thermo Fisher Scientific).

Molecular weights were determined using Precision Plus Protein Prestained Standards (Bio-Rad

Laboratories). After electrophoresis, the proteins were transferred onto a 0.45-μm polyvinylidene

difluoride membrane (Millipore). Next, after blocking for 30 min at room temperature and

washing, the membrane was incubated with the primary antibody in Tris-buffered saline (20 mM

Tris and 150 mM NaCl, pH = 7.4) supplemented with 0.1% Tween20 (TBS-T) containing 5%

nonfat milk or Blocking One (Nacalai Tesque) overnight at 4°C. After washing with TBS-T, the

membranes were incubated with horseradish peroxidase (HRP)-conjugated secondary antibody

in TBS-T containing 5% nonfat milk or Blocking One solution at room temperature for 1 h.

After washing with TBS-T, the membranes were treated with either the ECL start reagent (GE

Healthcare) or the ECL select reagent (GE Healthcare) according to the manufacturer's

guidelines. The chemiluminescent images of the membranes were acquired using FUSION-

SOLO.7S (Vilber-Loumat) and analyzed using ImageJ.

Dot blot analysis

Lysates of PC3 cells and PC3 cell-derived sEVs were prepared as described above. The

phospholipid content of these lysates was quantified using a LabAssay Phospholipid (Fujifilm).

Subsequently, 100 μl of these lysates (phospholipid concentration = 6.4 μM) was adsorbed to a

nitrocellulose membrane using a Bio-Dot (Bio-Rad). After blocking with TBS-T containing 3%

BSA for 30 min at room temperature, followed by washing, the membrane was incubated with

biotin-conjugated CTXB for the detection of GM1 (1:500, list labs, Cat# 112) or with the

primary IgM antibody against GM3 clone GMR6 (1:1000, Tokyo Chemical Industry, Cat#

A2582) in TBS-T containing 3% BSA overnight at 4°C. After washing with TBS-T, the

membranes were incubated with the HRP-conjugated primary antibody against biotin (1:10000,

Cell Signaling, Cat# 5571) or the HRP-conjugated secondary antibody against IgM (1:10000,

Invitrogen, Cat# 31440) in TBS-T containing 3% BSA solution at room temperature for 1 h.

After washing with TBS-T, the membranes were incubated with the ECL start reagent (GE

Healthcare). Chemiluminescence images of the membranes were acquired using FUSION-

SOLO.7S (Vilber-Loumat) and analyzed using ImageJ.

Preparation of ganglioside-containing liposomes

The gangliosides GM1 (bovine brain), GD1a (bovine brain) and GD2 (human brain) were

purchased from Avanti Polar Lipids, AdipoGen Life Sciences and Santa Cruz, respectively.

GM2, GM3, and GD3 were prepared by total chemical synthesis (Koikeda et al., 2019;

Takahashi et al., 2020). In an eggplant-shaped flask, 50 μl of 1 mM ganglioside (GM1, GM2,

GM3, GD1a, GD2, or GD3) in methanol, 500 μl of 10 mM DMPC in chloroform, and 0.5 μl of 1

mM Bodipy-SM (Invitrogen) in methanol were mixed in a solution of 99:1:0.01 mol% ratio

DMPC:ganglioside:Bodipy-SM. The solvent was first dried with nitrogen gas and further dried

with a CVE-3000 (EYELA) evaporator with VT-2000 (EYELA). Then, 5 ml of PBS was added

to the flask, which was incubated at room temperature for 1 h. The solution was completely

frozen in liquid nitrogen and then thawed in hot water (45 °C). The freeze–thaw process was

repeated 5 times. Using an extruder (LiposoFAST, AVESTIN), 500 µl of the vesicle suspension

was passed through 100-nm pore-sized polycarbonate membrane filters (10 times). The amount

of sialic acid in the liposomes was determined using a Sialic Acid Assay Kit (MAK314-1KT,

Sigma Aldrich), and the concentration of liposomes containing a ganglioside was adjusted to the

same level for use in microscopic observation.

Immunofluorescence imaging of the extracellular matrix

MRC-5 and HS-5 cells were cultured on glass-bottom dishes (IWAKI) for two days. The cells

were then incubated with 10 µg/ml primary antibodies against fibronectin (Sigma–Aldrich,

F3648), laminin (Sigma–Aldrich, L9393), or collagen type I (Abcam, ab34710) in MEM at 37°C

for 30 min. Next, the cells were washed twice with HBSS and incubated with 12 µg/ml

rhodamine-conjugated anti-rabbit IgG antibody (Cappel, 55666) in MEM at 37°C for 30 min.

After washing twice with HBSS, fresh HBSS was added to the glass-base dishes. Subsequently,

the cells were observed by epi-fluorescence microscopy (ECLIPSE Ts2-FL, Nikon, 60x 1.40 NA

oil objective), which was performed with a CMOS camera (DS-Qi2, Nikon). A 560 nm LED

lamp (Nikon) was used to excite rhodamine. The fluorescence intensities of laminin in the

regions of cells (cell contours were determined using bright-field images) were quantified using

ImageJ. The actual fluorescence intensities of laminin in the region of cells were measured by

subtracting the average fluorescence intensity without the primary antibody from the

fluorescence intensity of laminin stained with both the primary and secondary antibodies.

Determination of the concentrations of sEVs containing fluorescently labeled marker proteins

by total internal reflection fluorescence microscopy (TIRFM)

The concentrations of protein and lipids in the sEV suspension (on the order of nM) were too

low to determine very quantitatively via conventional spectrophotometry. Therefore, we

employed single-fluorescent particle tracking to directly quantify the number of sEVs. In brief,

the glass windows of single-well or triple-well glass-base dishes (IWAKI) were coated with 100

or 50 μl of 10 μg/ml anti-CD63 IgG antibody (8A12, Cosmo Bio), anti-CD81 IgG antibody (JS-

81, BD Biosciences), or anti-CD9 antibody (M-L13, BD Biosciences) in HBSS and then

incubated for 2 h at 37°C. Subsequently, the antibody solution was removed, and the glass

window was coated with either 100 or 50 μl of 50 μg/ml casein (Sigma–Aldrich) in HBSS for 1

h at 37°C. This coating process was essential for mitigating the nonspecific binding of sEVs to

the glass surface during subsequent experiments.

Either 100 or 50 µl of three distinct concentrations of sEVs containing CD63-Halo7,

CD81-Halo7, and CD9-Halo7, all conjugated with SF650T in HBSS, was applied to glass

windows pre-coated with antibodies against CD63, CD81, and CD9, respectively. The samples

were then incubated at 37°C for 1 h. After removing the sEVs and washing twice with HBSS, the

individual fluorescent particles of the sEVs were observed at 37°C with single-molecule

detection sensitivity by total internal reflection fluorescence microscopy (TIRFM) using an

Olympus IX-83 microscope (60x 1.49 NA oil objective) equipped with a high-speed gated image

intensifier (C9016-02MLG; Hamamatsu Photonics) coupled to an sCMOS camera (ORCA-

Flash4.0 V2; Hamamatsu Photonics), as previously described (Komura et al., 2016; Kinoshita et

al., 2017; Morise et al., 2019). SF650T was excited using a 647-nm laser (LuxXPlus647-140,

140 mW, Omicron Laserrange) at an intensity of 0.3 µW/µm^2.

All the acquired sEV movies were analyzed using ImageJ (Fiji). First, a noise reduction

process was carried out, wherein 10 frames of sEV images observed at 30 frames/sec were

averaged. Subsequently, each pixel was replaced by the average of the 3x3 neighborhood pixels

using a smooth filter. Single particles of sEVs whose fluorescence intensity exceeded that of free

SF650T were identified, and the numbers of sEVs were counted. By performing these

observations at three distinct sEV concentrations, we obtained a calibration curve. Thereafter,

sEVs derived from both intact cells and integrin subunit KO cells were prepared at identical

concentrations ($0.8\text{-}4\times10^{10}$ particles/ml).

Determination of the numbers of sEVs bound to ECM components on glass

To prevent nonspecific binding of sEVs to the glass surface, 50 or 100 μl of 50 μg/ml casein

solution in HBSS was added to glass windows in single-well or triple-well glass-base dishes

(IWAKI) and incubated for 1 h at 37°C. Then, the casein solution was replaced with 50 or 100 μl

of 100 μg/ml fibronectin (Roche), 50 μg/ml laminin (Sigma–Aldrich), 20 μg/ml laminin-511

(BioLamina), 100 μg/ml collagen type I (Sigma–Aldrich), or 50 μg/ml casein (Sigma–Aldrich)

for 2 h at 37°C. After removal of the ECM or casein solution, 50 or 100 μl of sEVs containing

CD63/CD81/CD9-Halo7 labeled with SF650T at the same concentration was applied to the glass

and incubated for 1 h at 37°C. After removal of the sEV suspension and two washes with HBSS,

individual fluorescent particles of the sEVs in HBSS were visualized at 37°C with single-

molecule detection sensitivity by TIRFM. The number of sEVs bound to the ECM or casein on

glass was determined using ImageJ (Fiji) as described above. The number of sEVs bound to

ECM-coated glass was estimated by subtracting the number of sEVs bound to casein-coated

glass in a 1024 pixel × 1024 pixel (81.9 μm × 81.9 μm) image.

Determination of the numbers of sEVs bound to cell membranes

sEVs containing CD63-Halo7 labeled with SF650T (sEV-CD63Halo7-SF650T) were incubated

with MRC-5 or HS-5 cells expressing mCherry or GFP in the cytosol at 37°C. sEV-CD63Halo7-

SF650T was visualized with single-molecule detection sensitivity by TIRFM, and mCherry or

GFP was observed by oblique illumination. The number of sEVs on the cell membrane (the cell

contour was determined using binarized mCherry or GFP images) was quantified every 10 min

by ImageJ (Fiji) as described above.

Simultaneous dual-color observation of dSTORM movie of ECM components and single

fluorescent particles of sEVs on living cell plasma membranes

For immunostaining of the ECM components for dSTORM movie observation, 0.67 μl of 5.6

μg/ml SaraFluor650B (SF650B, Goryo Chemical) NHS ester was incubated with 100 μl of 1

mg/ml anti-rabbit IgG antibody (0212-0081, Cappel) in 0.1 M $NaHCO_3$ for 60 min at room

temperature. SF650B-conjugated anti-rabbit IgG was then isolated using a Sephadex G-25

column (GE Healthcare). MRC-5 cells cultured on glass-based dishes for 2 days were

subsequently incubated with 10 μg/ml anti-fibronectin IgG (F3648; Sigma–Aldrich), anti-

laminin IgG (L9393; Sigma–Aldrich) or anti-collagen type I IgG (ab34710; Abcam) for 30 min

at 37°C. After washing twice with HBSS, the cells were incubated with 10 μg/ml SF650B-

conjugated anti-rabbit IgG antibody for 30 min at 37°C. After washing with HBSS 3 times, sEVs

containing CD63-Halo7-TMR were incubated with the cells for another 30 min at 37°C.

Individual fluorescent particles of the sEVs and single molecules of SF650B on the apical

surface of the cells at 512×512×pixel (25.6 μ×25.6 μm) were observed at 5-ms resolution (200

frames/second) for 3504 frames by oblique-angle illumination using an Olympus IX-83 inverted

microscope (100×1.5 NA oil objective) equipped with two high-speed gated image intensifiers

(C9016-02MLG; Hamamatsu Photonics) coupled to two sCMOS cameras (ORCA-Flash4.0 V2;

Hamamatsu Photonics), as described previously (Komura et al., 2016; Kinoshita et al., 2017;

Morise et al., 2019; Kemmoku et al., 2024). TMR and SF650B were excited using a 561-nm

laser (Excelsior-561-100-CDRW, 100 mW, Spectra-Physics) and a 647-nm laser (140 mW,

Omicron Laserrange) at 0.9 and 3.2 $\mu W/\mu m^2$, respectively.

dSTORM image reconstructions with a pixel size of 10 nm were performed using the

ThunderSTORM plugin for ImageJ (Ovesný et al., 2014) installed in the Fiji package

(Schindelin et al., 2012). Gaussian rendering with a localization precision of 24 nm was utilized

in this process.

The dSTORM super-resolution video data were generated using frame information and

the x and y coordinates of all the spots, which were incorporated into the csv data output by

ThunderSTORM. Furthermore, we used the uncertainty that represents the spatial precision of

the spot. A Gaussian distribution was generated for each spot, with the x and y coordinates

serving as the center and the standard deviation (S.D.) being equivalent to 6 times the uncertainty

of the spot. This distribution indicates the existence probability of the spot. The existence

probability distribution for all structures at time t was obtained as a summation of all the

distributions for spots that appeared in [6t+1 6t+1002] frames. The dSTORM super-resolution

video data were generated by repeating this process for time t=0,1,2,… If the value of each time

and coordinate in the dSTORM video data exceeded a threshold value θ, we considered a

structure to exist at the time and coordinate. The optimal threshold value θ was determined for

each dSTORM video data point to minimize the average in-class variance for the two classes

when every value in the dSTORM video data was classified into two classes using the threshold

value θ (Chen et al., 2014; Kemmoku et al., 2024). To synchronize the dSTORM movie of ECM

structures with the single-particle movie of sEVs, the temporal resolution of the single-particle

movie was converted to 33.3 frames/s by averaging 6 frames, and the averaged single-particle

image was merged with the dSTORM image created from 1002 frames, of which the middle

frame was synchronized with the averaged single-particle image (Fig. 7A). Single-particle

tracking in the movie was performed by in-house computer software based on previously

reported methods (Gelles et al., 1988; Kusumi et al., 1993). More details have been described

previously (Suzuki et al., 2007a, 2007b; Kasai et al., 2011; Suzuki et al., 2012; Komura et al.,

2016; Kinoshita et al., 2017; Morise et al., 2019; Fujiwara et al., 2023a, 2023b).

Colocalization analysis between ECM structures in dSTORM movies and single sEV particles

The distances between the centroids of the sEVs and the boundaries of the ECM structures were

quantified using single-particle tracking data and binarized dSTORM movies. To generate

binarized images of the ECM structures in the dSTORM movie, we employed the kernel density

estimation (KDE) method (Kemmoku et al., 2024). KDE is a traditional image segmentation

method that can rapidly determine the criterion for segmenting PALM and dSTORM images, as

reported previously (Chen et al., 2014; Kemmoku et al., 2024). Furthermore, a recent study

demonstrated that KDE is one of the most appropriate methods for segmenting images of clearly

outlined structures, such as protein clusters (Nieves et al., 2023). KDE provides a way to

interpolate object boundaries without bias by using random fluctuations of activated boundary

fluorophores.

The ThunderSTORM datasets were imported into in-house KDE software (MATLAB)

for image segmentation and colocalization analysis. We used not only the localization

coordinates but also the uncertainty, which reflects the spatial detection precision of the spots.

Let x_i, y_i and $u_i, i = 1, 2, \cdots, N$ be the horizontal and vertical localization coordinates and the

uncertainty of spot i, respectively. N represents the total number of spots. Considering the

uncertainty of the location measurement of spot i, the existence probability $p_i(x, y)$ of spot i

is considered to spread to the Gaussian distribution with the center coordinate x_i, y_i and the

standard deviation (S.D.) that is proportional to the uncertainty u_i as follows:

$$p_i(x, y) = \frac{1}{2\pi(Au_i)^2} e^{-\frac{(x-x_i)^2+(y-y_i)^2}{2(Au_i)^2}}$$

in which A is the proportional coefficient of S.D. to the uncertainty u_i. This coefficient was

estimated as $A \approx 6$ by some preliminary experiments. The Gaussian distribution for

constructing each existence probability $p_i(x, y)$ is called the Gaussian kernel. The existence

probability distribution for all spots, i.e., the dSTORM image, is obtained by averaging the

existence probability distribution $p_i(x, y)$ as follows:

$$p(x, y) = \alpha \frac{1}{N} \sum_{i=1}^{N} p_i(x, y) + (1 - \alpha)p_{bg}(x, y)$$

excluding spots having extremely small or large uncertainties u_i because such spots are likely

to contain noise (e.g. electrical shot noise from camera). $p_{bg}(x, y)$ represents the background

noise generated by the probability density distribution $f(v) = \frac{1}{2\pi\sigma_{bg}^2} e^{-\frac{(v-\mu_{bg})^2}{2\sigma_{bg}^2}}$, independent of

x, y. α is a weight determined by the respective occurrence probabilities of the spots and

background noise.

The structures can be considered to exist where the dSTORM image, which represents

the probability of structure existence, is above a certain threshold value θ because the spots

appear uniformly inside the structures with a certain probability. It is necessary to determine the

optimal threshold value θ for each obtained dSTORM image because the optimal value differs

depending on the target molecule and the experimental environment. In general, the histogram of

$p(x, y)$ values for all x, y forms a bimodal shape consisting of both larger values created by

dense spots appearing in structures and smaller values created by sporadic noise. Therefore, the

threshold value θ should be determined to separate these two structures. Otsu's method is a

well-known approach for determining such a threshold value (Otsu, 1979). Let S_{in} and S_{out} be

sets of coordinates (x, y) where $p(x, y) \geq \theta$ and $p(x, y) < \theta$, respectively. The sets S_{in} and

S_{out} indicate the inside and outside of the clusters, respectively. The numbers of elements in sets

S_{in} and S_{out} are represented by N_{in} and N_{out}, respectively. The intraclass variances of the

sets S_{in} and S_{out} are calculated by

$$\sigma^2[S_{in}] = \frac{1}{N_{in}} \sum_{(x,y) \in S_{in}} (p(x, y) - \mu[S_{in}])^2,$$

$$\sigma^2[S_{out}] = \frac{1}{N_{out}} \sum_{(x,y) \in S_{out}} (p(x, y) - \mu[S_{out}])^2.$$

In the above equations, $\mu[S_{in}]$ and $\mu[S_{out}]$ are the respective averages of the set S_{in} and S_{out}

obtained by

$$\mu[S_{in}] = \frac{1}{N_{in}} \sum_{(x,y) \in S_{in}} p(x, y),$$

$$\mu[S_{out}] = \frac{1}{N_{in}} \sum_{(x,y) \in S_{out}} p(x, y).$$

The average of these intraclass variances is

$$\sigma^2[S_{in}, S_{out}] = \frac{N_{in}\sigma^2[S_{in}] + N_{out}\sigma^2[S_{out}]}{N_{in} + N_{out}}.$$

In Otsu's method, the optimal threshold value $\hat{\theta}$ to minimize the average of the intraclass variances $\sigma^2[S_{in}, S_{out}]$ is determined as follows:

$$\hat{\theta} = \arg\min_{\theta} \sigma^2[S_{in}, S_{out}].$$

In this study, the structures were determined using Otsu's method.

Thereafter, the distance from the contour of the determined structures to the center of the sEVs on the PM was measured in all the frames of the superimposed movies of the ECM components and sEVs, and the number densities of pairwise distances at a given distance were plotted. If an sEV particle was within an ECM structure, the measured distance was expressed as a negative value. Furthermore, we simulated the distances between random coordinates and the boundaries of ECM structures in silico and calculated their corresponding relative frequency distributions using a bin width of 50 nm. To normalize the relative frequency of sEVs within each bin, we divided the data by the corresponding relative frequency of random spots. A ratio exceeding 1 indicated a higher incidence of sEVs at the measured distance.

dSTORM imaging of ECM-coated glass

The ECM molecules were coated on triple-well glass-bottom dishes and incubated with 5 μg/ml

anti-fibronectin IgG (F3648; Sigma–Aldrich), anti-laminin IgG (L9393; Sigma–Aldrich), or anti-collagen type I IgG (ab34710; Abcam) for 2 h at room temperature. After washing twice with HBSS, the ECM molecules were incubated with 2.5 μg/ml SF650B-conjugated anti-rabbit IgG antibodies for 1 h at room temperature. After washing with HBSS 5 times, single molecules of SF650B at 512 pixels × 512 pixels (25.6 μm × 25.6 μm) were observed via TIRFM at 5-ms resolution (200 frames/s) for 1008 frames. The dSTORM images were reconstructed from the movie using the ThunderSTORM plugin for ImageJ.

Cell spreading assay

The cells that secreted sEVs were cultured for 3-5 days and subsequently collected using a solution of 0.5% BSA and 5 mM EDTA in PBS. A total of 4×10^5 cells/ml were then evenly dispersed onto an ECM-coated single-well glass-bottom dish. The cells were observed after incubation for 0, 30, 60, or 120 min with an epi-illumination microscope (CKX53, Olympus, 20x 0.40 NA objective) equipped with a camera (WAT-01U2, WATEC). The cell area was quantified using ImageJ software.

Supplemental material

Fig. S1 shows the preparation of sEVs derived from PC3 cells and the determination of their size

and concentration. Fig. S2 shows the number of sEV particles attached to glass coated with ECM components. sEVs containing CD81 or CD9-Halo7 were derived from wild-type, integrin β1, integrin α2, integrin α6, or integrin β4 KO PC3 cells and labeled with SF650T. Fig. S3 shows the results of western blot analysis to determine whether sEVs derived from PC3 cells were covered by fibronectin and/or laminin. Fig. S4 shows the flow cytometry analysis of B78 cells expressing several glycosyltransferases to examine the expression levels of gangliosides. Table S1 presents the numbers of sEVs labeled with tetraspanin-Halo7-TMR attached to glass coated with ECM components. Table S1 is related to Figs. 3 and S2. Videos 1-3 (Video 1: fibronectin, Video 2: collagen type I, Video 3: laminin) show the simultaneous observation of single sEV-CD63Halo7-TMR particles (green) and dSTORM ECM movies (magenta) on an MRC cell. Video 4 shows an enlarged movie of the colocalization of an sEV-CD63Halo7-TMR particle (green) and laminin (dSTORM movie, magenta) on an MRC cell.

Data availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments

We thank Joan Massagué (Memorial Sloan Kettering Cancer Center) and Ayuko Hoshino (University of Tokyo) for kindly providing the human 4175-LuT cells. We also thank Yumi Matsuno for supporting the preparation of extracellular vesicles from tumor cells, Shinobu Kawaguchi for constructing various cDNAs, and Takahiro Fujiwara and Akihiro Kusumi for developing the analysis software for single-molecule imaging. This work was supported in part by Japan Science and Technology Agency (JST) grants from the Core Research for Evolutional Science and Technology (CREST) program in the field of "Extracellular Fine Particles" (JPMJCR18H2) (K.G.N.S., K.M.H., and H.A.), Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS) (21H02424, 18H02401) (K.G.N.S.), a Grant-in-Aid for Challenging Research (Exploratory) from JSPS (20K21387) (K.G.N.S.), a Grant-in-Aid for Core-to-Core Program from JSPS (JSCCA202000007) (K.G.N.S., and H.A.), a Grant-in-Aid for Innovative Areas from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT) (18H04671) (K.G.N.S.), the National Cancer Center Research and Development Fund (2023-A-03) (K.G.N.S.), the Takeda Science Foundation (K.G.N.S.), Uehara Memorial Foundation (K.G.N.S.), the Mizutani Foundation for Glycoscience (K.G.N.S.), the Daiichi Sankyo Foundation of Life Science (K.G.N.S.), the Research Foundation for Opto-Science and Technology (K.G.N.S.), and the Naito Foundation (K.G.N.S.), as well as a Grant-in-Aid from Support for Pioneering Research Initiated by the Next Generation in JST (JST

SPRING) (JPMJSP2125) (T. I.) and a Grant-in-Aid for JSPS Fellows (23KJ1045) (T. I.).

Author contributions

T.I., K.M.H., and K.G.N.S. conceived and formulated the project. T.I., K.M.H., R.S.K. and

K.G.N.S. designed the biological experiments and participated in the discussions. T.I., M.K., and

A.S. performed the experiments on the binding of EVs to ECM components and the data

analysis. Y.M. helped to isolate sEVs from cell culture supernatants. S.K. generated the cDNA

plasmids. Koichi F, Y.O., and Keiko F. generated B78 cells expressing only one type of

ganglioside. H.A. and N.K. synthesized the gangliosides. Y.Y., T.I., K.M.H., and K.G.N.S.

developed the analysis software for super-resolution movies and single-particle tracking. T.I.,

Y.Y. and K.G.N.S. wrote the manuscript, and all the authors discussed the results and

participated in revising the manuscript.

References

Altei, W.F., B.C. Pachane, P.K.D. Santos, L.N.M. Ribeiro, B.H. Sung, A.M. Weaver, and H.S.
Selistre-de-Araújo. 2020. Inhibition of alphavbeta3 integrin impairs adhesion and uptake
of tumor-derived small extracellular vesicles. *Cell Commun. Signal.* 18:158.

Berditchevski, F. 2001. Complexes of tetraspanins with integrins: more than meets the eye. *J.
Cell Sci.* 114:4143–4151.

Berditchevski, F., E. Odintsova, S. Sawada, and E. Gilbert. 2002. Expression of the
palmitoylation-deficient CD151 weakens the association of alpha 3 beta 1 integrin with
the tetraspanin-enriched microdomains and affects integrin-dependent signaling. *J. Biol.
Chem.* 277:36991–37000.

Carney, R.P., S. Hazari, T. Rojalin, A. Knudson, T. Gao, Y. Tang, R. Liu, T. Viitala, M.
Yliperttula, and K.S. Lam. 2017. Targeting tumor-associated exosomes with integrin-
binding peptides. *Adv. Biosyst.* 1:1600038.

Charrin, S., S. Manié, C. Thiele, M. Billard, D. Gerlier, C. Boucheix, and E. Rubinstein. 2003. A
physical and functional link between cholesterol and tetraspanins. *Eur. J. Immunol.*
33:2479–2489.

Chen, K.-C.J., G. Yang, and J. Kovacevic. 2014. Spatial density estimation based segmentation
of super-resolution localization microscopy images. *In* 2014 IEEE International

Conference on Image Processing (ICIP). IEEE, Paris, France. 867–871.

Coumans, F.A.W., E. Van der Pol, A.N. Böing, N. Hajji, G. Sturk, T.G. Van Leeuwen, and R. Nieuwland. 2014. Reproducible extracellular vesicle size and concentration determination with tunable resistive pulse sensing. *J. Extracell. Vesicles* 3:25922.

Franco, M., C. Muratori, S. Corso, E. Tenaglia, A. Bertotti, L. Capparuccia, L. Trusolino, P.M. Comoglio, and L. Tamagnone. 2010. The tetraspanin CD151 is required for Met-dependent signaling and tumor cell growth. *J. Biol. Chem.* 285:38756–38764.

Fujiwara, T.K., S. Takeuchi, Z. Kalay, Y. Nagai, T.A. Tsunoyama, T. Kalkbrenner, K. Iwasawa, K.P. Ritchie, K.G.N. Suzuki, and A. Kusumi. 2023a. Development of ultrafast camera-based single fluorescent-molecule imaging for cell biology. *J. Cell Biol.* 222:e202110160.

Fujiwara, T.K., T.A. Tsunoyama, S. Takeuchi, Z. Kalay, Y. Nagai, T. Kalkbrenner, Y.L. Nemoto, L.H. Chen, A.C.E. Shibata, K. Iwasawa, K.P. Ritchie, K.G.N. Suzuki, and A. Kusumi. 2023b. Ultrafast single-molecule imaging reveals focal adhesion nano-architecture and molecular dynamics. *J. Cell Biol.* 222:e202110162.

Gagnoux-Palacios, L., M. Dans, W. Van't Hof, A. Mariotti, A. Pepe, G. Meneguzzi, M.D. Resh, and F.G. Giancotti. 2003. Compartmentalization of integrin alpha6beta4 signaling in lipid rafts. *J. Cell Biol.* 162:1189–1196.

Gaus, K., S. Le Lay, N. Balasubramanian, and M.A. Schwartz. 2006. Integrin-mediated adhesion

regulates membrane order. *J. Cell Biol.* 174:725–734.

Gelles, J., B.J. Schnapp, and M.P. Sheetz. 1988. Tracking kinesin-driven movements with
nanometre-scale precision. *Nature* 331:450–453.

Gu, J., Y. Sumida, N. Sanzen, and K. Sekiguchi. 2001. Laminin-10/11 and fibronectin
differentially regulate integrin-dependent Rho and Rac activation via p130(Cas)-CrkII-
DOCK180 pathway. *J. Biol. Chem.* 276:27090–27097.

Gurung, S., D. Perocheau, L. Touramanidou, and J. Baruteau. 2021. The exosome journey: from
biogenesis to uptake and intracellular signalling. *Cell Commun. Signal.* 19:47.

Hart, I.R., and I.J. Fidler. 1980. Role of organ selectivity in the determination of metastatic
patterns of B16 melanoma. *Cancer Res.* 40:2281–2287.

Hirosawa, K.M., Y. Sato, E. Yamaguchi, N. Komura, H. Ando, A. Hoshino, Y. Yokota, and K.G.
Suzuki. 2024. Small extracellular vesicles trigger paracrine adhesion signaling that
facilitates their uptake by the target cells.

Hoshino, A., B. Costa-Silva, T.-L. Shen, G. Rodrigues, A. Hashimoto, M.T. Mark, H. Molina, S.
Kohsaka, A. Di Giannatale, S. Ceder, S. Singh, C. Williams, N. Soplop, K. Uryu, L.
Pharmer, T. King, L. Bojmar, A.E. Davies, Y. Ararso, T. Zhang, H. Zhang, J. Hernandez,
J.M. Weiss, V.D. Dumont-Cole, K. Kramer, L.H. Wexler, A. Narendran, G.K. Schwartz,
J.H. Healey, P. Sandstrom, K.J. Labori, E.H. Kure, P.M. Grandgenett, M.A.

Hollingsworth, M. de Sousa, S. Kaur, M. Jain, K. Mallya, S.K. Batra, W.R. Jarnagin, M.S. Brady, O. Fodstad, V. Muller, K. Pantel, A.J. Minn, M.J. Bissell, B.A. Garcia, Y. Kang, V.K. Rajasekhar, C.M. Ghajar, I. Matei, H. Peinado, J. Bromberg, and D. Lyden. 2015. Tumour exosome integrins determine organotropic metastasis. *Nature* 527:329–335.

Huang, C.-C., R. Narayanan, S. Alapati, and S. Ravindran. 2016. Exosomes as biomimetic tools for stem cell differentiation: applications in dental pulp tissue regeneration. *Biomaterials* 111:103–115.

Huang, C., Z. Rajfur, N. Yousefi, Z. Chen, K. Jacobson, and M.H. Ginsberg. 2009. Talin phosphorylation by Cdk5 regulates Smurf1-mediated talin head ubiquitylation and cell migration. *Nat. Cell Biol.* 11:624–630.

Huet-Calderwood, C., F. Rivera-Molina, D.V. Iwamoto, E.B. Kromann, D. Toomre, and D.A. Calderwood. 2017. Novel ecto-tagged integrins reveal their trafficking in live cells. *Nat. Commun.* 8:570.

Hynes, R.O. 2002. Integrins: bidirectional, allosteric signaling machines. *Cell* 110:673–687.

Ichikawa, N., K. Iwabuchi, H. Kurihara, K. Ishii, T. Kobayashi, T. Sasaki, N. Hattori, Y. Mizuno, K. Hozumi, Y. Yamada, and E. Arikawa-Hirasawa. 2009. Binding of laminin-1 to monosialoganglioside GM1 in lipid rafts is crucial for neurite outgrowth. *J. Cell Sci.*

122:289–299.

Jeppesen, D.K., A.M. Fenix, J.L. Franklin, J.N. Higginbotham, Q. Zhang, L.J. Zimmerman, D.C. Liebler, J. Ping, Q. Liu, R. Evans, W.H. Fissell, J.G. Patton, L.H. Rome, D.T. Burnette, and R.J. Coffey. 2019. Reassessment of exosome composition. *Cell* 177:428–445.e18.

Jin, J.K., P.C. Tien, C.J. Cheng, J.H. Song, C. Huang, S.H. Lin, and G.E. Gallick. 2015. Talin1 phosphorylation activates beta1 integrins: a novel mechanism to promote prostate cancer bone metastasis. *Oncogene* 34:1811–1821.

Kasai, R.S., K.G.N. Suzuki, E.R. Prossnitz, I. Koyama-Honda, C. Nakada, T.K. Fujiwara, and A. Kusumi. 2011. Full characterization of GPCR monomer-dimer dynamic equilibrium by single molecule imaging. *J. Cell Biol.* 192:463–480.

Kazarov, A.R., X. Yang, C.S. Stipp, B. Sehgal, and M.E. Hemler. 2002. An extracellular site on tetraspanin CD151 determines alpha 3 and alpha 6 integrin-dependent cellular morphology. *J. Cell Biol.* 158:1299–1309.

Kemmoku, H., K. Takahashi, K. Mukai, T. Mori, K.M. Hirosawa, F. Kiku, Y. Uchida, Y. Kuchitsu, Y. Nishioka, M. Sawa, T. Kishimoto, K. Tanaka, Y. Yokota, H. Arai, K.G.N. Suzuki, and T. Taguchi. 2024. Single-molecule localization microscopy reveals STING clustering at the trans-Golgi network through palmitoylation-dependent accumulation of cholesterol. *Nat. Commun.* 15:220.

Kinoshita, M., K.G.N. Suzuki, N. Matsumori, M. Takada, H. Ano, K. Morigaki, M. Abe, A. Makino, T. Kobayashi, K.M. Hirosawa, T.K. Fujiwara, A. Kusumi, and M. Murata. 2017. Raft-based sphingomyelin interactions revealed by new fluorescent sphingomyelin analogs. *J. Cell Biol.* 216:1183–1204.

Koikeda, M., N. Komura, H.-N. Tanaka, A. Imamura, H. Ishida, M. Kiso, and H. Ando. 2019. Synthesis of ganglioside analogs containing fluorescently labeled GalNAc for single-molecule imaging. *J. Carbohydr. Chem.* 38:509–527.

Komura, N., K.G.N. Suzuki, H. Ando, M. Konishi, M. Koikeda, A. Imamura, R. Chadda, T.K. Fujiwara, H. Tsuboi, R. Sheng, W. Cho, K. Furukawa, K. Furukawa, Y. Yamauchi, H. Ishida, A. Kusumi, and M. Kiso. 2016. Raft-based interactions of gangliosides with a GPI-anchored receptor. *Nat. Chem. Biol.* 12:402–410.

Kusumi, A., Y. Sako, and M. Yamamoto. 1993. Confined lateral diffusion of membrane receptors as studied by single particle tracking (nanovid microscopy). Effects of calcium-induced differentiation in cultured epithelial cells. *Biophys. J.* 65:2021–2040.

Lammerding, J., A.R. Kazarov, H. Huang, R.T. Lee, and M.E. Hemler. 2003. Tetraspanin CD151 regulates alpha6beta1 integrin adhesion strengthening. *Proc. Natl. Acad. Sci. U. S. A.* 100:7616–7621.

Llorente, A., T. Skotland, T. Sylvänne, D. Kauhanen, T. Róg, A. Orłowski, I. Vattulainen, K.

Ekroos, and K. Sandvig. 2013. Molecular lipidomics of exosomes released by PC-3 prostate cancer cells. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1831:1302–1309.

Lu, F., L. Zhu, T. Bromberger, J. Yang, Q. Yang, J. Liu, E.F. Plow, M. Moser, and J. Qin. 2022. Mechanism of integrin activation by talin and its cooperation with kindlin. *Nat. Commun.* 13:2362.

Marar, C., B. Starich, and D. Wirtz. 2021. Extracellular vesicles in immunomodulation and tumor progression. *Nat. Immunol.* 22:560–570.

Möller, A., and R.J. Lobb. 2020. The evolving translational potential of small extracellular vesicles in cancer. *Nat. Rev. Cancer* 20:697–709.

Monkley, S.J., C.A. Pritchard, and D.R. Critchley. 2001. Analysis of the mammalian talin2 gene TLN2. *Biochem. Biophys. Res. Commun.* 286:880–885.

Morise, J., K.G.N. Suzuki, A. Kitagawa, Y. Wakazono, K. Takamiya, T.A. Tsunoyama, Y.L. Nemoto, H. Takematsu, A. Kusumi, and S. Oka. 2019. AMPA receptors in the synapse turnover by monomer diffusion. *Nat. Commun.* 10:5245.

Moser, M., K.R. Legate, R. Zent, and R. Fässler. 2009. The tail of integrins, talin, and kindlins. *Science* 324:895–899.

Nagai, R., A. Sugimachi, Y. Tanimoto, K.G.N. Suzuki, F. Hayashi, D. Weikert, P. Gmeiner, R.S. Kasai, and K. Morigaki. 2021. Functional reconstitution of dopamine D2 receptor into a

supported model membrane in a nanometric confinement. *Adv. Biol.* 5:e2100636.

Nazarenko, I., S. Rana, A. Baumann, J. McAlear, A. Hellwig, M. Trendelenburg, G. Lochnit, K.T. Preissner, and M. Zöller. 2010. Cell surface tetraspanin Tspan8 contributes to molecular pathways of exosome-induced endothelial cell activation. *Cancer Res.* 70:1668–1678.

Nieves, D.J., J.A. Pike, F. Levet, D.J. Williamson, M. Baragilly, S. Oloketuyi, A. De Marco, J. Griffié, D. Sage, E.A.K. Cohen, J.-B. Sibarita, M. Heilemann, and D.M. Owen. 2023. A framework for evaluating the performance of SMLM cluster analysis algorithms. *Nat. Methods* 20:259–267.

Odintsova, E., T.D. Butters, E. Monti, H. Sprong, G. Van Meer, and F. Berditchevski. 2006. Gangliosides play an important role in the organization of CD82-enriched microdomains. *Biochem. J.* 400:315–325.

Ono, M., N. Kosaka, N. Tominaga, Y. Yoshioka, F. Takeshita, R. Takahashi, M. Yoshida, H. Tsuda, K. Tamura, and T. Ochiya. 2014. Exosomes from bone marrow mesenchymal stem cells contain a microRNA that promotes dormancy in metastatic breast cancer cells. *Sci. Signal.* 7:ra63.

Otsu, N. 1979. A threshold selection method from Gray-level histograms. *IEEE Trans. Syst. Man Cybern.* 9:62–66.

Ovesný, M., P. Křížek, J. Borkovec, Z. Svindrych, and G.M. Hagen. 2014. ThunderSTORM: a

comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution

imaging. *Bioinformatics* 30:2389–2390.

Paget, S. 1989. The distribution of secondary growths in cancer of the breast. 1889. *Cancer

Metastasis Rev.* 8:98–101.

Peinado, H., M. Alečković, S. Lavotshkin, I. Matei, B. Costa-Silva, G. Moreno-Bueno, M.

Hergueta-Redondo, C. Williams, G. García-Santos, C. Ghajar, A. Nitadori-Hoshino, C.

Hoffman, K. Badal, B.A. Garcia, M.K. Callahan, J. Yuan, V.R. Martins, J. Skog, R.N.

Kaplan, M.S. Brady, J.D. Wolchok, P.B. Chapman, Y. Kang, J. Bromberg, and D. Lyden.

2012. Melanoma exosomes educate bone marrow progenitor cells toward a pro-

metastatic phenotype through MET. *Nat. Med.* 18:883–891.

Peinado, H., H. Zhang, I.R. Matei, B. Costa-Silva, A. Hoshino, G. Rodrigues, B. Psaila, R.N.

Kaplan, J.F. Bromberg, Y. Kang, M.J. Bissell, T.R. Cox, A.J. Giaccia, J.T. Erler, S.

Hiratsuka, C.M. Ghajar, and D. Lyden. 2017. Pre-metastatic niches: organ-specific homes

for metastases. *Nat. Rev. Cancer* 17:302–317.

Purushothaman, A., S.K. Bandari, J. Liu, J.A. Mobley, E.E. Brown, and R.D. Sanderson. 2016.

Fibronectin on the surface of myeloma cell-derived exosomes mediates exosome-cell

interactions. *J. Biol. Chem.* 291:1652–1663.

Purushothaman, G., and V. Thiruvenkatam. 2019. High yield expression of recombinant CD151

in *E. coli* and a structural insight into cholesterol binding domain. *Mol. Biotechnol.*

61:905–915.

Raposo, G., and W. Stoorvogel. 2013. Extracellular vesicles: exosomes, microvesicles, and

friends. *J. Cell Biol.* 200:373–383.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C.

Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P.

Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image

analysis. *Nat. Methods* 9:676–682.

Stipp, C.S. 2010. Laminin-binding integrins and their tetraspanin partners as potential

antimetastatic targets. *Expert Rev. Mol. Med.* 12:e3.

Strachan, L.R., and M.L. Condic. 2004. Cranial neural crest recycle surface integrins in a

substratum-dependent manner to promote rapid motility. *J. Cell Biol.* 167:545–554.

Sun, Z., M. Costell, and R. Fässler. 2019. Integrin activation by talin, kindlin and mechanical

forces. *Nat. Cell Biol.* 21:25–31.

Suzuki, K.G.N., T.K. Fujiwara, M. Edidin, and A. Kusumi. 2007a. Dynamic recruitment of

phospholipase C gamma at transiently immobilized GPI-anchored receptor clusters

induces IP3-Ca2+ signaling: single-molecule tracking study 2. *J. Cell Biol.* 177:731–742.

Suzuki, K.G.N., T.K. Fujiwara, F. Sanematsu, R. Iino, M. Edidin, and A. Kusumi. 2007b. GPI-anchored receptor clusters transiently recruit Lyn and G alpha for temporary cluster immobilization and Lyn activation: single-molecule tracking study 1. *J. Cell Biol.* 177:717–730.

Suzuki, K.G.N., R.S. Kasai, K.M. Hirosawa, Y.L. Nemoto, M. Ishibashi, Y. Miwa, T.K. Fujiwara, and A. Kusumi. 2012. Transient GPI-anchored protein homodimers are units for raft organization and function. *Nat. Chem. Biol.* 8:774–783.

Takahashi, M., J. Shirasaki, N. Komura, K. Sasaki, H.-N. Tanaka, A. Imamura, H. Ishida, S. Hanashima, M. Murata, and H. Ando. 2020. Efficient diversification of GM3 gangliosides via late-stage sialylation and dynamic glycan structural studies with ^{19}F solid-state NMR. *Org. Biomol. Chem.* 18:2902–2913.

Théry, C., K.W. Witwer, E. Aikawa, M.J. Alcaraz, J.D. Anderson, R. Andriantsitohaina, A. Antoniou, T. Arab, F. Archer, G.K. Atkin-Smith, D.C. Ayre, J.-M. Bach, D. Bachurski, H. Baharvand, L. Balaj, S. Baldacchino, N.N. Bauer, A.A. Baxter, M. Bebawy, C. Beckham, A. Bedina Zavec, A. Benmoussa, A.C. Berardi, P. Bergese, E. Bielska, C. Blenkiron, S. Bobis-Wozowicz, E. Boilard, W. Boireau, A. Bongiovanni, F.E. Borràs, S. Bosch, C.M. Boulanger, X. Breakefield, A.M. Breglio, M.Á. Brennan, D.R. Brigstock, A. Brisson, M.L. Broekman, J.F. Bromberg, P. Bryl-Górecka, S. Buch, A.H. Buck, D. Burger, S.

Busatto, D. Buschmann, B. Bussolati, E.I. Buzás, J.B. Byrd, G. Camussi, D.R. Carter, S. Caruso, L.W. Chamley, Y.-T. Chang, C. Chen, S. Chen, L. Cheng, A.R. Chin, A. Clayton, S.P. Clerici, A. Cocks, E. Cocucci, R.J. Coffey, A. Cordeiro-da-Silva, Y. Couch, F.A. Coumans, B. Coyle, R. Crescitelli, M.F. Criado, C. D'Souza-Schorey, S. Das, A. Datta Chaudhuri, P. de Candia, E.F. De Santana, O. De Wever, H.A. Del Portillo, T. Demaret, S. Deville, A. Devitt, B. Dhondt, D. Di Vizio, L.C. Dieterich, V. Dolo, A.P. Dominguez Rubio, M. Dominici, M.R. Dourado, T.A. Driedonks, F.V. Duarte, H.M. Duncan, R.M. Eichenberger, K. Ekström, S. El Andaloussi, C. Elie-Caille, U. Erdbrügger, J.M. Falcón-Pérez, F. Fatima, J.E. Fish, M. Flores-Bellver, A. Försönits, A. Frelet-Barrand, F. Fricke, G. Fuhrmann, S. Gabrielsson, A. Gámez-Valero, C. Gardiner, K. Gärtner, R. Gaudin, Y.S. Gho, B. Giebel, C. Gilbert, M. Gimona, I. Giusti, D.C. Goberdhan, A. Görgens, S.M. Gorski, D.W. Greening, J.C. Gross, A. Gualerzi, G.N. Gupta, D. Gustafson, A. Handberg, R.A. Haraszti, P. Harrison, H. Hegyesi, A. Hendrix, A.F. Hill, F.H. Hochberg, K.F. Hoffmann, B. Holder, H. Holthofer, B. Hosseinkhani, G. Hu, Y. Huang, V. Huber, S. Hunt, A.G.-E. Ibrahim, T. Ikezu, J.M. Inal, M. Isin, A. Ivanova, H.K. Jackson, S. Jacobsen, S.M. Jay, M. Jayachandran, G. Jenster, L. Jiang, S.M. Johnson, J.C. Jones, A. Jong, T. Jovanovic-Talisman, S. Jung, R. Kalluri, S.-I. Kano, S. Kaur, Y. Kawamura, E.T. Keller, D. Khamari, E. Khomyakova, A. Khvorova, P. Kierulf, K.P. Kim, T. Kislinger, M.

Klingeborn, D.J. Klinke, 2nd, M. Kornek, M.M. Kosanović, Á.F. Kovács, E.-M. Krämer-

Albers, S. Krasemann, M. Krause, I.V. Kurochkin, G.D. Kusuma, S. Kuypers, S.

Laitinen, S.M. Langevin, L.R. Languino, J. Lannigan, C. Lässer, L.C. Laurent, G. Lavieu,

E. Lázaro-Ibáñez, S. Le Lay, M.-S. Lee, Y.X.F. Lee, D.S. Lemos, M. Lenassi, A.

Leszczynska, I.T. Li, K. Liao, S.F. Libregts, E. Ligeti, R. Lim, S.K. Lim, A. Linē, K.

Linnemannstöns, A. Llorente, C.A. Lombard, M.J. Lorenowicz, Á.M. Lörincz, J. Lötvall,

J. Lovett, M.C. Lowry, X. Loyer, Q. Lu, B. Lukomska, T.R. Lunavat, S.L. Maas, H.

Malhi, A. Marcilla, J. Mariani, J. Mariscal, E.S. Martens-Uzunova, L. Martin-Jaular,

M.C. Martinez, V.R. Martins, M. Mathieu, S. Mathivanan, M. Maugeri, L.K. McGinnis,

M.J. McVey, D.G. Meckes, Jr., K.L. Meehan, I. Mertens, V.R. Minciacchi, A. Möller, M.

Møller Jørgensen, A. Morales-Kastresana, J. Morhayim, F. Mullier, M. Muraca, L.

Musante, V. Mussack, D.C. Muth, K.H. Myburgh, T. Najrana, M. Nawaz, I. Nazarenko, P.

Nejsum, C. Neri, T. Neri, R. Nieuwland, L. Nimrichter, J.P. Nolan, E.N. Nolte-'t Hoen, N.

Noren Hooten, L. O'Driscoll, T. O'Grady, A. O'Loghlen, T. Ochiya, M. Olivier, A. Ortiz,

L.A. Ortiz, X. Osteikoetxea, O. Østergaard, M. Ostrowski, J. Park, D.M. Pegtel, H.

Peinado, F. Perut, M.W. Pfaffl, D.G. Phinney, B.C. Pieters, R.C. Pink, D.S. Pisetsky, E.

Pogge von Strandmann, I. Polakovicova, I.K. Poon, B.H. Powell, I. Prada, L. Pulliam, P.

Quesenberry, A. Radeghieri, R.L. Raffai, S. Raimondo, J. Rak, M.I. Ramirez, G. Raposo,

M.S. Rayyan, N. Regev-Rudzki, F.L. Ricklefs, P.D. Robbins, D.D. Roberts, S.C.

Rodrigues, E. Rohde, S. Rome, K.M. Rouschop, A. Rughetti, A.E. Russell, P. Saá, S.

Sahoo, E. Salas-Huenuleo, C. Sánchez, J.A. Saugstad, M.J. Saul, R.M. Schiffelers, R.

Schneider, T.H. Schøyen, A. Scott, E. Shahaj, S. Sharma, O. Shatnyeva, F. Shekari, G.V.

Shelke, A.K. Shetty, K. Shiba, P.R.M. Siljander, A.M. Silva, A. Skowronek, O.L. Snyder,

2nd, R.P. Soares, B.W. Sódar, C. Soekmadji, J. Sotillo, P.D. Stahl, W. Stoorvogel, S.L.

Stott, E.F. Strasser, S. Swift, H. Tahara, M. Tewari, K. Timms, S. Tiwari, R. Tixeira, M.

Tkach, W.S. Toh, R. Tomasini, A.C. Torrecilhas, J.P. Tosar, V. Toxavidis, L. Urbanelli, P.

Vader, B.W. Van Balkom, S.G. Van der Grein, J. Van Deun, M.J. Van Herwijnen, K. Van

Keuren-Jensen, G. Van Niel, M.E. Van Royen, A.J. Van Wijnen, M.H. Vasconcelos, I.J.

Vechetti, Jr., T.D. Veit, L.J. Vella, É. Velot, F.J. Verweij, B. Vestad, J.L. Viñas, T.

Visnovitz, K.V. Vukman, J. Wahlgren, D.C. Watson, M.H. Wauben, A. Weaver, J.P.

Webber, V. Weber, A.M. Wehman, D.J. Weiss, J.A. Welsh, S. Wendt, A.M. Wheelock, Z.

Wiener, L. Witte, J. Wolfram, A. Xagorari, P. Xander, J. Xu, X. Yan, M. Yáñez-Mó, H.

Yin, Y. Yuana, V. Zappulli, J. Zarubova, V. Žėkas, J.-Y. Zhang, Z. Zhao, L. Zheng, A.R.

Zheutlin, A.M. Zickler, P. Zimmermann, A.M. Zivkovic, D. Zocco, and E.K. Zuba-

Surma. 2018. Minimal information for studies of extracellular vesicles 2018

(MISEV2018): a position statement of the International Society for Extracellular Vesicles

and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* 7:1535750.

Thul, P.J., L. Åkesson, M. Wiking, D. Mahdessian, A. Geladaki, H. Ait Blal, T. Alm, A. Asplund, L. Björk, L.M. Breckels, A. Bäckström, F. Danielsson, L. Fagerberg, J. Fall, L. Gatto, C. Gnann, S. Hober, M. Hjelmare, F. Johansson, S. Lee, C. Lindskog, J. Mulder, C.M. Mulvey, P. Nilsson, P. Oksvold, J. Rockberg, R. Schutten, J.M. Schwenk, Å. Sivertsson, E. Sjöstedt, M. Skogs, C. Stadler, D.P. Sullivan, H. Tegel, C. Winsnes, C. Zhang, M. Zwahlen, A. Mardinoglu, F. Pontén, K. von Feilitzen, K.S. Lilley, M. Uhlén, and E. Lundberg. 2017. A subcellular map of the human proteome. *Science* 356:eaal3321.

Valadi, H., K. Ekström, A. Bossios, M. Sjöstrand, J.J. Lee, and J.O. Lötvall. 2007. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat. Cell Biol.* 9:654–659.

Van Niel, G., G. D'Angelo, and G. Raposo. 2018. Shedding light on the cell biology of extracellular vesicles. *Nat. Rev. Mol. Cell. Biol.* 19:213–228.

Winterwood, N.E., A. Varzavand, M.N. Meland, L.K. Ashman, and C.S. Stipp. 2006. A critical role for tetraspanin CD151 in alpha3beta1 and alpha6beta4 integrin-dependent tumor cell functions on laminin-5. *Mol. Biol. Cell*. 17:2707–2721.

Witwer, K.W., D.C. Goberdhan, L. O'Driscoll, C. Théry, J.A. Welsh, C. Blenkiron, E.I. Buzás, D. Di Vizio, U. Erdbrügger, J.M. Falcón-Pérez, Q.-L. Fu, A.F. Hill, M. Lenassi, J. Lötvall, R.

Nieuwland, T. Ochiya, S. Rome, S. Sahoo, and L. Zheng. 2021. Updating MISEV:

evolving the minimal requirements for studies of extracellular vesicles. *J. Extracell.*

Vesicles 10:e12182.

Wondimu, Z., G. Gorfu, T. Kawataki, S. Smirnov, P. Yurchenco, K. Tryggvason, and M.

Patarroyo. 2006. Characterization of commercial laminin preparations from human

placenta in comparison to recombinant laminins 2 (alpha2beta1gamma1), 8

(alpha4beta1gamma1), 10 (alpha5beta1gamma1). *Matrix Biol.* 25:89–93.

Yamashiro, S., M. Haraguchi, K. Furukawa, K. Takamiya, A. Yamamoto, Y. Nagata, K.O. Lloyd,

H. Shiku, and K. Furukawa. 1995. Substrate specificity of beta 1,4-N-

acetylgalactosaminyltransferase in vitro and in cDNA-transfected cells. GM2/GD2

synthase efficiently generates asialo-GM2 in certain cells. *J. Biol. Chem.* 270:6149–6155.

Yang, X., C. Claas, S.-K. Kraeft, L.B. Chen, Z. Wang, J.A. Kreidberg, and M.E. Hemler. 2002.

Palmitoylation of tetraspanin proteins: modulation of CD151 lateral interactions,

subcellular distribution, and integrin-dependent cell morphology. *Mol. Biol. Cell.* 13:767–

781.

Yang, X., O.V. Kovalenko, W. Tang, C. Claas, C.S. Stipp, and M.E. Hemler. 2004.

Palmitoylation supports assembly and function of integrin-tetraspanin complexes. *J. Cell*

Biol. 167:1231–1240.

Yasuda, T., H. Watanabe, K.M. Hirosawa, K.G.N. Suzuki, K. Suga, and S. Hanashima. 2022. Fluorescence spectroscopic analysis of lateral and transbilayer fluidity of exosome membranes. *Langmuir* 38:14695–14703.

Yesmin, F., Furukawa, K., Kambe, M., Ohmi, Y., Bhuiyan, R. H., Hasnat, M. A., Mizutani, M., Tajima, O., Hashimoto, N., Tsuchida, A., Kaneko, K., and Furukawa, K. 2023. Extracellular vesicles released from ganglioside GD2-expressing melanoma cells enhance the malignant properties of GD2-negative melanomas. *Sci. Rep.* 13: 4987.

Yusa, K., R. Rad, J. Takeda, and A. Bradley. 2009. Generation of transgene-free induced pluripotent mouse stem cells by the piggyBac transposon. *Nat. Methods* 6:363–369.

Zhang, D.X., X.T.T. Dang, L.T. Vu, C.M.H. Lim, E.Y.M. Yeo, B.W.S. Lam, S.M. Leong, N. Omar, T.C. Putti, Y.C. Yeh, V. Ma, J.-Y. Luo, W.C. Cho, G. Chen, V.K.M. Lee, A. Grimson, and M.T.N. Le. 2022. alphavbeta1 integrin is enriched in extracellular vesicles of metastatic breast cancer cells: a mechanism mediated by galectin-3. *J. Extracell. Vesicles* 11:e12234.

Author notes

Disclosures: The authors declare no competing interests exist.

Figures

Fig. 1. Western blotting of integrin subunits in PC3 cells and sEVs. (A) sEVs were isolated from

the cell culture supernatant of intact PC3 cells or from cells in which an integrin subunit (β1, β4,

α2, α6) was knocked out via the CRISPR-Cas9 method. (B) The correlation map of integrin

heterodimers and the ECM.

Fig. 2. All the cells that secreted sEVs in this study attached to and spread at similar velocities on

all three ECM components (fibronectin, laminin, and collagen type I) coated on glass. (A)

dSTORM images of the ECM (left-top: fibronectin, left-bottom: collagen type I, right: laminin)

coated on glass. (B) Images of HeLa cells attached to glass coated with ECM components

(fibronectin, collagen type I, laminin) or casein after 0, 30, or 60 min of incubation. (C) The time

course of the area of five tumor cell lines on glass coated with ECM components (fibronectin,

collagen type I, laminin) or casein (n = 8 cells). The data are presented as the mean ± SE.

Fig. 3. Integrin α2β1 in sEVs derived from PC3 cells is responsible for the binding of CD63-

containing sEVs to collagen type I, and integrins α6β1 and α6β4 are responsible for the binding

to laminin. (A, C, E, and G) Single-particle fluorescence images of sEV-CD63Halo7-SF650T on glass coated with fibronectin, collagen type I, and laminin before and after the KO of integrin β1 (A), integrin α2 (C), integrin α6 (E), and integrin β4 (G). (B, D, F, and H) The numbers of sEVs attached to glass coated with these extracellular matrix molecules before and after the KO of integrin β1 (B), integrin α2 (D), integrin α6 (F), and integrin β4 (H). The data are presented as the mean ± SE. * indicates a significant difference (p<0.001) according to Welch's t-test (two-sided).

Fig. 4. GM1 is responsible for the binding of sEVs to laminin. (A) (top) Schematic diagram of

gangliosides and (bottom) chemical structure of GM1. (B) Dot blotting of GM1 and GM3 in PC3

cells and PC3-derived sEVs. (C) Western blot analysis of integrin subunits in B78 cell lines with

abundant expression of one type of ganglioside—GM1, GM2, GM3, GD2 or GD2/GD3—and

sEVs derived from these cells. (D and E) The numbers of sEVs attached to glass coated with

laminin (D) and the numbers normalized to the ratio of integrin $\alpha6$/CD81 in the sEVs (E). (F and

G) The numbers of sEVs attached to glass coated with fibronectin (F) and the numbers

normalized to the ratio of integrin $\alpha5$/CD81 in the sEVs (G). (H) Single-particle fluorescence

images of DMPC-liposomes containing GM1, GM2, GM3, GD1a, GD2, or GD3 on glass coated

with fibronectin or laminin. (I and J) The numbers of liposomes attached to glass coated with

laminin (I) or fibronectin (J). The data are presented as the mean $\pm$ SE. * indicates a significant

difference ($p<0.001$) according to Welch's t-test (two-sided).

Fig. 5. Tumor-derived sEVs bind much more strongly to laminin than to fibronectin. (A) The

diameters of sEVs, mEVs, and MVs derived from the 4175-LuT, PC3, BxPC3, HeLa, and

SKBR3 cell lines were measured by qNano. (B) Single-particle fluorescence images of sEV-CD63Halo7-SF650T particles derived from the 4175-LuT, PC3, BxPC3, HeLa, and SKBR3 cell lines on glass coated with fibronectin or laminin. (C-E) The numbers of EV-CD63Halo7-SF650T particles (sEVs (C), mEVs (D), and MVs (E)) attached to glass coated with ECM components. The data are presented as the mean ± SE. (F) Western blotting of integrin subunits in sEVs, mEVs, and MVs derived from 4175-LuT, PC3, BxPC3, HeLa, and SKBR3 cells.

Fig. 6. Integrins β1 and α6 in sEVs and the extracellular matrix (ECM) are responsible for the

binding of sEVs to cell plasma membranes. (A) Immunofluorescence images of fibronectin,

collagen type I, and laminin in human normal embryonic lung fibroblast (MRC-5) cells and

human bone marrow stromal (HS-5) cells. (B) The fluorescence intensities of laminin on cells in

1000 μm^2. The data are presented as the mean ± SE. *p<0.01 by Welch's t-test (two-sided). (C)

Fluorescence images of MRC-5 cells expressing mCherry and sEV-CD63Halo7-SF650T

particles (arrowhead) attached to cell membranes after 10, 20, and 30 min of incubation. sEVs

isolated from the cell culture supernatant of intact PC3 cells (top panels) and from the cell

culture supernatant of integrin β1 KO PC3 cells (bottom panels) bound to MRC-5 cells. (D)

Fluorescence images of HS-5 cells and sEV-CD63Halo7-SF650T particles (arrowhead) attached

to cell membranes after 30 min of incubation. sEVs were isolated from intact PC3 cells (top

panels) and integrin β1 KO PC3 cells (bottom panels). (E) Fluorescence images of MRC-5 cells

and the attached sEV-CD63Halo7-SF650T particles derived from intact PC3 cells (top panels)

and from integrin α6 KO PC3 cells (bottom panels) after 30 min of incubation. (F-H) Time

course of the numbers of intact sEVs and integrin β1-KO sEVs attached to the MRC-5 cell

membrane (*n* = 8 images) (F) and to the HS-5 cell membrane (*n* = 15 images) (G) per 1000 μm^2.

(H) Time course of the numbers of intact sEVs and integrin α6 KO sEVs attached to the MRC-5

cell membrane per 1000 mm^2 (*n* = 12 images). The data are presented as the mean ± SE.

Fig. 7. sEVs bind to laminin on the plasma membranes of living MRC-5 cells, as revealed by

pseudo-real-time super-resolution movie observation. (A) Schematic diagram for generating

merged movies of dSTORM images of the ECM structures and single-particle images of sEVs.

Data acquisition was performed by observing single fluorescent molecules of SF650B at 200

frames/s (3,504 frames), and dSTORM images were reconstructed using the data acquired every

5.0 s (=1,002 frames). The first dSTORM image was reconstructed using frames 1-1,002 and the

process was then repeated by shifting the initial frames backward by 6 frames; thus, 417

dSTORM images were obtained. These dSTORM still images were connected to construct the

"dSTORM movie". The single-particle movies of sEVs were subjected to a rolling-average for 6

frames and synchronously merged with the dSTORM movie. (B-D) Conventional

immunofluorescence images (top-left) and dSTORM images (bottom-left) of the ECM structures

(B: fibronectin, C: collagen type I, D: laminin) on the cells. dSTORM images of the ECM

structures (magenta) and single-particle images of sEV-CD63Halo7-TMR particles (green) were

merged (right). sEVs localized near (<100 nm) the boundary of ECM structures and sEVs

localized alone are indicated by yellow and white arrowheads, respectively. (E) The image

sequence (every 0.3 sec) of laminin obtained by dSTORM (magenta) and a single sEV-

CD63Halo7-TMR particle (green) on the living MRC-5 cell membrane. sEVs colocalized with

laminin, as indicated by the yellow arrowhead. (F) Colocalization analysis method. The

normalized relative frequency was defined as the ratio of the average value of the spatial pair

correlation function of the actual image to that of randomly distributed spots generated by a

computer. Zero on the x-axis indicates the contour of the ECM structures in the dSTORM images

determined by the kernel density estimation (KDE) method. When sEVs are enriched near the

ECM structures, the normalized relative frequency is greater than 1. (G) Probability density

analysis of the sEVs and ECM structures. The colored areas indicate regions within the ECM

structures. The sEV-CD63Halo7-TMR particles localized near the contour of laminin ($n = 20$

cells).

Fig. 8. Talin-1 in sEVs does not regulate the binding affinity of integrins for laminin. (A)

Western blot analysis of PC3 cell-derived sEVs after talin-1 KD by siRNA or overexpression of

talin-1. (B, C, and D) The fluorescence images (B) and the numbers of PC3-sEVs attached to

glass coated with laminin before and after (C) talin-1 KD or (D) overexpression of talin-1.

Halo7-integrin β1 in sEVs was labeled with SF650T. (E, F, and G) The fluorescence images (E)

and the numbers of CD63-labeled sEVs attached to glass coated with laminin before and after

(F) talin-1 KD and (G) overexpression of talin-1. (H) Western blot analysis of the

phosphorylation of Ser425 on talin-1 in PC3 cells and sEVs. Roscovitine: an inhibitor of CDK5

that phosphorylates Ser425 of talin-1.

Fig. 9. CD151 and cholesterol regulate the binding affinity of sEVs for laminin. (A)

Fluorescence images of sEV-CD63Halo7-SF650T particles bound to laminin (LN) on glass before and after cholesterol depletion by MβCD and the numbers of attached sEVs per image (82 μm × 82 μm). (B and C) The numbers of sEV-CD63-Halo7-SF650T particles attached to glass coated with laminin before and after treatment with saponin (B) and the addition of cholesterol by MβCD-cholesterol complex (C). (D) Fluorescence images of an MRC-5-GFP cell and sEV-CD63Halo7-SF650T particles on the MRC-5 cell after 30 min of incubation. (E and F) Time course of the number of sEV-CD63-Halo7-SF650T particles per 1000 μm^2 attached to the MRC-5 cell membrane before and after treatment with MβCD ($n = 16$ cells) (E) or the MβCD-cholesterol complex ($n = 8$ cells) (F). (G) Western blot analysis of CD151 and integrin subunits in PC3 cells and sEVs after CD151 KD. The amount of cell proteins loaded in one lane was 2.5 times greater than that of the sEVs in the other lane. (H) Images of wild-type (WT) PC3 cells and CD151-knockdown cells on glass coated with ECM components (fibronectin (FN), laminin (LN), or collagen type I (COL1)) after 2 h of incubation. The areas of the cells were quantified. (I) Single-particle fluorescence images of sEV-CD63Halo7-SF650T particles bound to laminin (LN) on glass before and after CD151 was knocked down and cholesterol was depleted by MβCD (left). The number of attached sEVs increased (right). (J) The binding affinity ratio of cholesterol-depleted sEVs to intact sEVs was compared with that of CD151 KD sEVs. The data are presented as the mean ± SE. * indicates a significant difference (p<0.01) according to

Welch's t-test (two-sided).

Fig. 10. Schematic model showing the binding of EVs to the recipient cell PM. (A) Integrin heterodimers in EVs, α6β1/α6β4 and α2β1, can bind to laminin and collagen type I, respectively, but integrin heterodimers in EVs, such as α5β1, bind only weakly to fibronectin. (B) Because Ser 425 of talin-1 is hardly phosphorylated in EVs, talin-1 is inactive and is not involved in promoting the ECM-binding activity of integrin heterodimers. (C) CD151 facilitates the binding of integrins α6β1 and α6β4 in sEVs to laminin. CD151 also links these integrins with tetraspanin-enriched membrane microdomains (TEMs) in a cholesterol-dependent manner. A molecule X in TEMs attenuates the binding of integrin heterodimers in EVs to laminin.