Androgenic shifts and external cues mediate parental care versus

infanticide in mimic poison frogs

Amaris R. Lewis[1†], Billie C. Goolsby[1†], Bryan H. Juarez[1], Madison P. Lacey[1], Lauren A. O'Connell[1]*

1 Department of Biology, Stanford University, Stanford, CA 94305, USA

Word count: Abstract: 200, Main text: 6,582

Key words: parental care, cannibalism, testosterone, glucocorticoid, kin recognition,

Ranitomeya imitator, Dendrobatidae

*To whom correspondence should be addressed:

Lauren A. O'Connell

Department of Biology

Stanford University

371 Jane Stanford Way

Stanford, CA 94305

loconnel@stanford.edu

†These authors contributed equally to this manuscript.

Abstract

Infanticide is widespread across the animal kingdom. However, the physiological mechanisms underlying infanticide versus care or neglect are relatively unexplored. Here, we identified salient environmental and physiological antecedents of care and infanticide in the mimic poison frog (*Ranitomeya imitator*), a biparental and monogamous amphibian in which female parents feed their tadpoles with unfertilized eggs. Specifically, we explored potential environmental cues by evaluating changes in the frequency of food provisioning and tadpole mortality after cross-fostering tadpoles within and between morphs and displacing tadpoles within the terraria of their parents. After analyzing trophic egg deposition and observing tadpole mortality in the latter condition, we concluded that changes to offspring location reduce care and increase infanticide. Following this observation, we investigated whether steroid hormones relate to infanticide in an unfamiliar, resource-limited setting. Infanticidal behavior towards fertilized eggs and hatchlings included cannibalism and was associated with lower testosterone concentrations upon displacement to the new environment, while corticosterone concentrations did not correlate with infanticide at any sampling event. Ratios of testosterone to corticosterone were overall lower in infanticidal individuals. Overall, these findings support earlier investigations of environmental cues for offspring-directed behavior in poison frogs, while offering a novel perspective on temporal endocrine correlates of infanticide.

Highlights

- Offspring location drives parental decisions of care vs. infanticide.
- In novel territories, adults cannibalize conspecific, unrelated young.
- Lower circulating T in novel territory is associated with infanticide.
- Infanticidal adults exhibit lower T/CORT ratios.

1. Introduction

Animal parents exhibit remarkable behavioral flexibility, performing parental care (e.g. feeding, protecting, and transporting young), neglect, and infanticide, regardless of kinship (Hrdy, 1979; Royle et al. 2012; Lukas and Huchard, 2014; Lukas and Huchard, 2019). As care often represents a significant energy investment, parental decision-making is sensitive to evolutionary trade-offs dependent on ecological and physiological constraints (Clutton-Brock, 1991; Alonso-Alvarez and Velando, 2012). Moreover, parental behavior is often informed by environmental cues relevant to discriminating between related and unrelated offspring (Waldman, 1988). Therefore, parents provide a valuable framework for studying flexible social decision-making at multiple levels. While behavioral ecologists have reported numerous adaptive rationales for infanticide across diverse taxa, including mammals, fish, amphibians, insects, and birds (Hrdy, 1979; Ebensperger and Blumstein, 2007; Bose, 2022), efforts to link this behavior to its hormonal substrates is limited, especially beyond mammals. By connecting endocrine states to the performance of parental care versus infanticide, we can clarify how evolutionary innovations in physiology enable rapid responses to social challenges.

Steroid hormones present a compelling avenue to investigate the physiological basis of infanticide due to their well-established role in the proximate facilitation of social behaviors, especially those relating to aggression and parental care. For example, acutely elevated glucocorticoids promote social aggression in mammals and non-mammals alike (Summers et al., 2005; Haller, 2014), but are also positively associated with parental care in many vertebrates (Bales et al., 2006; O'Connor et al., 2009; Crossin et al., 2012). There are also reports of a positive association between infanticide and postnatal testosterone in rodents, although this relationship appears inconsistent, possibly owing to differences in genetic backgrounds and prenatal exposures (Elwood and Stolzenberg, 2020). More generally, across the animal kingdom, increases in male aggressive behavior sometimes co-occur with increases in circulating androgens during periods of social instability, such as intraspecific competition for

territory or mates (Wingfield et al., 1990; Hirschenhauser and Oliveira, 2006; Rodriguez et al.,

2022). Accordingly, testosterone is generally hypothesized to be lower in adult males during

parental care, although research suggests a more complicated relationship, as the trade-off

between sexual and paternal behavior is variably detrimental to lifetime fitness across ecological

contexts (Wingfield et al., 2001; Hirschenhauser and Oliveira, 2006). There also exists emerging

support for a link between testosterone, parental care, and aggression in female vertebrates,

including mammals, birds, lizards, and fishes (Rosvall, 2020). Additionally, proneness to social

aggression may be better understood in terms of hormonal interactions, as proposed by the

dual hormone hypothesis (Mehta and Prasad, 2015), where higher testosterone/cortisol ratios

are generally associated with dominance attainment, risk-taking, and social aggression in some

mammals including humans (Sapolsky, 2005; Terburg et al., 2009; Goll et al., 2023), but this

trend has yet to be directly explored in other vertebrates.

Hormonal transitions between care and infanticide are unpredictable across species.

Although some work on this topic has emerged in mammalian models (Elwood and Stolzenberg,

2020), the lack of comparable investigations in other taxa impedes the development of a

comparative, predictive framework for parental decision-making. Poison frogs demonstrate

extraordinary diversity in care strategies, including unimaternal, uniparental, and biparental

care, in addition to a range of parental behaviors, namely tadpole transport, maternal egg

provisioning, and cannibalism (Weygoldt, 1987; Summers and Tumulty, 2014). Therefore, these

amphibians present an ideal system to study how changes in physiology resulting from

environmental challenges influence parental decisions. Previous work has shown that, in poison

frogs, offspring location, territorial status, and recent reproductive activity can be determining

factors for whether adults cannibalize young or not (Stynoski, 2009; Ringler et al., 2016; Ringler

et al., 2017; Spring et al., 2019). Specifically, work by Ringler et al., found that in the brilliant-

thighed poison frog, *Allobates femoralis,* territorial status defines parental care or infanticidal

decisions in males, whereas females perform care dependent on exact location of offspring

within their territory (2016, 2017). These findings have established a framework for investigating

environmental cues for infanticide in other poison frogs, but the concurrent hormonal states

underlying this behavior remain unexplored.

We used mimic poison frogs (*Ranitomeya imitator*) to better understand how hormonal

shifts in response to environmental changes may elicit changes in parental behavior. Mimic

poison frogs are biparental and sexually monogamous, forming pair bonds that last several

months and performing sex-specific parental roles to cooperate in the care of their eggs and

tadpoles (Brown et al., 2008). Males of this species transport tadpoles after egg hatching, and

females provision trophic eggs to their tadpoles. By studying a monogamous, biparental

amphibian, we contribute to a comparative framework for infanticide relating to the evolution of

mating systems and parental care. Based on prior work in poison frogs (Stynoski, 2009; Ringler

et al., 2016; Ringler et al., 2017), we hypothesized that care versus infanticide decisions in this

species depend more on tadpole location than direct recognition cues. We performed kin

recognition assays including cross-fostering and displacement within parents' terraria, then

identified differences in the frequencies of egg provisioning and tadpole mortality. Following the

identification of cues for care versus infanticide, we sought to uncover hormonal correlates of

infanticidal behavior by moving individual adults to a novel environment with an unrelated

fertilized egg or hatchling and measuring water-borne testosterone and corticosterone

concentrations. We reasoned that testosterone and corticosterone should be reflective of

perceived social competition and stress, respectively, and that individuals would react to these

challenges by eliminating unrelated young.

2. Methods

2.1 Animal husbandry

All *R. imitator* eggs and tadpoles used in this study were captive-bred in our poison frog

colony. Adults were purchased from Indoor Ecosystems between 2019-2021 (Whitehouse,

Ohio) or Ruffing's Ranitomeya between 2021-2023 (Tiffin, Ohio, USA). One adult male and one

adult female were housed together in a 30 x 30 x 45 cm terraria containing sphagnum moss

substrate, driftwood, live plants, egg deposition sites, and film canisters filled with water for

tadpole deposition. Terraria were automatically misted ten times daily and frogs were fed live

Drosophila fruit flies dusted with Repashy Calcium Plus (Oceanside, CA, USA) three times per

week and with springtail isopods once a week. Tadpoles used in this study were observed in the

home or foster terraria as described below (all with the same dimensions and substrates). All

procedures in this study were approved by the Stanford University Animal Care and Use

Committee (Protocol #34242).

2.2 Offspring recognition assays

Ranitomeya imitator pairs used for this study were required to have previously

successfully raised a tadpole within 31 days of the experiment. Locations of tadpole deposition

were marked on the outside of the terrarium with the date and time to ensure consistency in

placing tadpole canisters in the same location throughout the duration of the study. Tadpoles

were randomly placed into three experimental conditions (**Fig. 1A**): same-morph cross-foster (n

= 16 across 6 terraria), different-morph cross-foster (n = 12 across 5 terraria), or intra-tank

displacement (n = 18 across 7 terraria). Control tadpoles (n=26 across 11 terraria) served as a

measure of parental investment under normal conditions. Whenever a tadpole was moved to

another terrarium, it was replaced by another tadpole such that the number of tadpoles in that

terrarium was consistent between assays.

Tadpole canisters were observed daily for 14 days in each trial, specifically noting

tadpole condition and trophic egg deposits as a proxy for parental investment. Tadpole stage,

size, and age were randomized, with all tadpoles being less developed than Gosner stage 32

(Gosner, 1960) to avoid counting feeding patterns that resulted from changes in investments as

tadpoles neared metamorphosis. Every tadpole underwent one trial as a positive control, such

that all tadpoles in the experimental groups were also observed in the control group. We did this

in order to measure parental effort per tadpole under normal conditions. Sampling each

terrarium several times for positive controls enabled us to account for the number of tadpoles in

that terrarium (one to four) and to capture individual variations in care by the focal pair.

In cases where tadpoles were exchanged between morphs, tadpoles were assigned to

the "different-morph cross-foster" category, while replacements that were consistent with the

morph of the previous tadpole were assigned to the "same-morph cross-foster" category. We

chose to manipulate morphs because morphs are conspecific color variations tightly linked to

geographical areas where coloration divergence occurs. For our study, we focused on two

morphs, *Yurimaguas* (striped) and *Rapidos* (banded), previously established to have genetic

diversity (Twomey et al., 2015). Tadpoles were always transported in their respective canisters

to retain any tadpole-specific olfactory cues already present in the water and to minimize stress.

Tadpoles displaced within their parents' terraria were moved at a minimum of 5 cm. Offspring

remained in their original canisters and were moved either to a laterally opposite corner of the

terrarium or elsewhere along the terrarium edge. Tadpoles were always moved to a location not

previously occupied by a canister.

2.3 "Takeover" behavior assays

To identify changes in physiology in infanticidal versus non-infanticidal parents, we

simulated a territory takeover in an assay inspired by Ringler et al. (2017). We selected paired

adults that raised at least one offspring in their terrarium within 31 days of the takeover trial as a

proxy for parental experience and reproductive status. Frogs from the same reproductive pair

were put into separate takeover trials at the same time to mitigate any risk of mate absence

influencing behavior directed towards offspring. The takeover terrarium had the same

dimensions, with elements of the environment (vegetation, replacement of dead leaves and

substrate, fresh water) disrupted and replaced to introduce unfamiliarity. Rearrangements and

cleaning occurred between trials. Takeover terraria always housed at least two tadpole

canisters, one upright and filled with water and the second empty and sideways, to remove bias

for water availability and shelter, respectively. Frogs were fed the same diet on the same

schedule in their takeover tanks as their home tanks with a dish of springtails available *ad*

libitum. While baseline hormone collection of the adult occurred, we placed a single, unrelated

fertilized late-stage egg to hatchling (Gosner stage 19-22) on a leaf on the terrarium floor. The

leaf was positioned below a Wyze Cam v3 to record behavior using previously described

methods (Goolsby et al., 2023). Frogs (n=12 females, 12 males) were introduced to the

terrarium and recorded for a maximum of seven days. Infanticidal behavior was characterized

as either consumption of the fertilized egg or hatchling (**Video S1**) or as repetitive physical

disturbances to the egg, where the snout repeatedly dug at the jelly-like casing in what we

interpreted as an attempt at egg cannibalism impeded by the encasement (**Video S2**).

2.4 Steroid hormone collection and processing

Frog hormones were collected immediately before frogs were displaced to the new tank

("baseline"), 24 ± 3 hours later ("move"), and following tadpole transport or infanticidal

behaviors, or on Day 7 if no such behavior occurred ("final"), with hormone collections occurring

at the same time of day unless an infanticidal behavior happened earlier or later in the day (**Fig.**

2A-B). Three subjects demonstrated infanticide on the day after displacement, so hormone

measurements counted for both "move" and "final" sampling events. Baseline collection always

occurred between 11:30 AM and 2:30 PM to avoid confounding effects of circadian physiology

on hormone concentrations. We ultimately analyzed data from 65 sampling events across 23

individuals.

As shown in previous studies with poison frogs and amphibians (Gabor et al., 2013;

Baugh et al., 2018, Baugh and Gray-Gailliard, 2021; Rodríguez et al., 2022), water-borne

testosterone and corticosterone were collected as a non-invasive measurement that reflects

circulating levels of both hormones. Frogs were individually moved to a petri dish containing 40

mL of distilled water treated with reverse osmosis (Josh's Frogs RO R/x, Osowo, MI, USA). The

collection lasted for one hour before either transferring the frog to the takeover terrarium (on

Day 0), 24 hours after takeover (on Day 1), or returning the frog to the original terrarium

(dependent on behavior; if none observed, Day 7). Immediately following 60 minutes, frogs were

measured for length (snout-vent length) using a digital caliper (Shahe Measuring Tools,

Amazon) along with mass, using a Maxus precision pocket scale (sensitivity 200 x 0.01 g,

Amazon). Individual mass was measured at each hormone sampling event, then averaged to

account for variations related to body fluctuations.

In sample processing, steroid hormones were pre-extracted using Sep-Pak C18

cartridges (Waters, Milford, MA, USA). The cartridges were conditioned using 2 mL of 100%

ethanol, followed by 2 mL of Milli-Q water treated with reverse osmosis. Water samples were

pushed through the column at a rate of approx. 10mL/minute, and then eluted with 4 mL of

100% ethanol. Ethanolic extracts were stored in glass vials at 4☐ until nitrogen evaporation.

One day prior to hormone quantification, the 4 mL of eluted hormone samples were divided into

2 mL new glass tubes for separate corticosterone and testosterone analyses from the same

frog. The 2 mL tubes were placed in a 37☐ water bath, and evaporated with gentle flow N_2 gas.

After evaporation, hormone samples were resuspended in 250 µl of assay buffer specific to the

corresponding ENZO kit, and stored at 4☐ overnight until analysis the following day.

2.5 Steroid hormone quantification via enzyme-linked immunosorbent assays (ELISA)

After overnight incubation at 4☐, samples were warmed to room temperature, and

corticosterone and testosterone concentrations were determined using commercially available

ELISA kits (ENZO Life Sciences, Farmingdale, NY; Corticosterone: cat. no. ADI-900-097,

antibody: donkey anti-sheep IgG, sensitivity: 27.0 pg/mL; Testosterone: cat. no. ADI-900-065,

antibody: goat anti-mouse IgG, sensitivity: 5.67 pg/mL) according to the manufacturer's

instructions. We chose to analyze corticosterone rather than cortisol concentrations because

corticosterone is suspected to be the main adrenocorticotropic hormone (ACTH)-responsive

glucocorticoid for *R. imitator* (Cockrem, 2013; Westrick et al., 2023). Samples were vortexed

and 100 µL of the samples were added in duplicate in individual wells of microtiter plates.

Following manufacturer instructions, plates were read at 405 nm, with correction between 570

and 590 nm, using a microplate reader (Synergy H1, BioTek Instruments, Winooski, VT, USA).

Hormone concentrations were calculated using a four-parameter logistic curve in the software

Gen5 (version 3.05, BioTek Instruments, Winooski, VT, USA). Samples with duplicate

measurements that yielded a coefficient of variation (CV) > 30% were set aside for imputation.

Before removing these duplicates, intra-assay CV values for testosterone and corticosterone

were 9.7% and 13.9%. After removal, intra-assay CV values were 9.1% and 9.0%. Inter-assay

CV values were 13.7% and 11.8%.

2.6 Statistics

We cleaned and analyzed all data using RStudio (R version 4.2.3, R Core Team,

Boston, MA). We implemented all statistical analyses using a generalized linear mixed model

(GLMM) framework. For each analysis, we used Akaike Information Criterion corrected for small

sample sizes (AICc; Burnham and Anderson, 2002) to choose between alternative model fits.

We evaluated standard model diagnostics and also tested for outliers and quantile deviations

using the '*DHARMa'* package (version 0.4.6, Hartig, 2022). All models in this manuscript passed

all diagnostic tests. To evaluate the significance of main effects, we performed a Type III

analysis of variance with Satterthwaite's method. We evaluated pairwise comparisons and

accounted for multiple comparisons using the Tukey method in the *'emmeans'* package (version

1.9.0, Lenth, 2023). We accounted for effect sizes in multi-termed modeling using the

'*effectsize'* package (version 0.8.6, Ben-Shachar et al., 2020) and with pairwise effect size

comparisons using '*emmeans*'. Figures were created using the package '*ggplot2*' (version 3.4.4,

Wickham, 2016) and assembled in Adobe Illustrator (Adobe Illustrator 2023).

2.6.1 Offspring manipulations

We analyzed the number of eggs deposited for each tadpole among the four conditions

(control, same-morph cross-foster, different-morph cross-foster, intra-tank displacement; **Fig.**

1A). To test whether egg deposition changes due to cross-fostering and intra-tank

displacement, we used a GLMM with a zero inflation component implemented using the

'*glmmTMB*' package (version 1.1.8, Brooks et al., 2017). We chose a Poisson distributed model

based on best fit, with eggs fed as a dependent variable, experiment type and number of

siblings within a terrarium as independent variables, and focal terrarium as a random effect.

2.6.2 Takeover trials

To determine how testosterone, corticosterone, and testosterone:corticosterone ratios

change relative to cannibalizing behavior, we implemented a GLMM using the '*lmerTest*'

package (3.1-3, Kuznetsova et al., 2017). We set the outcome ("infanticide" vs "non-

infanticide"), sampling event ("baseline" vs. "move" vs. "final"), sex, and their interactions as

fixed effects and individual frog identity, average individual mass, volume of collection water,

and body length (snout-vent length) as random effects. We estimated average individual mass

as the average body mass across each sampling event and this was necessary to account for

intra-individual differences in feeding. Since mass was not collected for a single frog, we used

mean imputation to estimate the single missing value as the averaged individual mass across

the entire study. To improve interpretation of regression coefficients, we removed insignificant

interaction terms using stepwise backward regression.

We ran separate models for testosterone, corticosterone, and ratios of testosterone to

corticosterone. We selected appropriate data transformations (natural log, square root, inverse,

no transformation) for each dependent variable using AICc. Using AICc and diagnostic criteria,

we selected log transformations for corticosterone and ratio modeling, and inverse

transformations for testosterone modeling.

Since corticosterone, testosterone, and ratio values of samples exceeding the CV

threshold of 30% are unreliable, we treated these values as missing data. We estimated these

missing data using regression imputation (Ellington et al., 2015; Nakagawa and Freckleton,

2011) and the methods described above. Briefly, regression imputation estimates missing data

as predictions obtained from a regression model fit using the dataset without missing data.

Thus, all available information from complete and incomplete samples is used to predict missing

data while simultaneously preserving all correlations found in the original dataset and without

introducing new correlations. Out of 65 measurements, we imputed only 6 (9.2%) of our

corticosterone values, 2 (3.1%) of our testosterone values, and 8 (12.3%) of our ratio values. All

of our final models used for imputation and inference passed all diagnostic tests.

3. Results

3.1 Offspring location as an indirect recognition cue

To identify which offspring features *R. imitator* parents use when caring for young, we

subjected tadpoles to either cross-fostering or displacement within their parents' terraria and

observed egg deposition and tadpole mortality for 14 days. We hypothesized that in addition to

offspring manipulations, like cross-fostering, the number of siblings present may also influence

how many eggs can be provisioned. We manipulated offspring identity (same-morph cross-

foster, different-morph cross-foster) and offspring location (intratank displacement, **Fig. 1A**). We

found that offspring manipulation type (χ^2 = 17.74, p < 0.0005; **Table S1**) was an overall better

predictor of eggs provisioned than the number of siblings present in a tank at a time (CI: -0.51 –

0.22, χ^2 = 0.62, p = 0.43; **Table S1**). Surprisingly, even within positive controls, sibling number

does not correlate with number of eggs fed (CI: -0.46 – 0.47, χ^2 = 6.00E-04, p = 0.98; **Table**

S1).

Figure 1. Disrupting offspring location impedes egg provisioning and promotes infanticide. (A) Control

tadpoles were the offspring of the parents in that terrarium and were not moved from their original locations. Cross-

fostered tadpoles were exchanged between terraria, such that they were unrelated to the parents in the new

terrarium. They were specifically moved to the same location previously occupied by the offspring they replaced. The

different-morph cross-foster manipulation followed the same procedure as same-morph cross-fosters, with the

additional condition that tadpoles were exchanged between terraria housing different morphs (striped, banded),

discerned by parents' coloration patterns. Finally, intra-tank displacement tadpoles were related to the parents in the

terrarium but were displaced by a minimum of five centimeters to a different location in the terrarium. **(B)** Tadpoles

displaced within their parents' tanks received significantly less feeding in the form of trophic eggs from parents (Tukey

adjustment for a family of 4 estimates: p < 0.0008). Significance: 0 < *** < 0.001 < ** < 0.01 < * < 0.05 < ns < 0.1. "ns"

= not significant. **(C)** Only tadpoles displaced within their parents' tanks (n=4 out of 18) were subjected to infanticide

with no deposition of a second tadpole.

We next asked whether offspring displacement influenced parental provisioning of young

and found a strong effect on the number of eggs provisioned (CI: -2.43 – -0.37, z = -2.66, p <

0.01, **Table 1**; model diagnostics may be found in **Fig S2** with details for analysis of deviance in

Table S1). Specifically, over 14 days, displaced tadpoles received significantly fewer trophic

eggs compared to positive controls (**Fig. 1B**, p_{adj} = 0.0008, $\square$ = 1.81). Across our experiment,

control tadpoles were fed the most, at an average of 1.77 eggs every two weeks, whereas same

morph and different morph cross fosters were fed less than positive controls, being fed 1.13 and

0.58 eggs on average, respectively. Displaced tadpoles were the least cared for, being fed an

average of 0.27 eggs within two weeks. In other words, positive control and cross fosters

averaged between 1-2 eggs being fed whereas tadpoles that were spatially moved, even with

their biological parents, were fed 0-1 eggs over 2 weeks. Tadpoles cross-fostered within and

outside of their morph did not experience a significant reduction in feeding compared to positive

controls (**Fig. 1B**, p_{adj} = 0.71-0.15, $\square$ = -0.910 - -0.409, respectively; **Table S2**). Tadpoles

displaced within their parents' terraria were subjected to a variety of infant-directed behaviors,

including "care" as exemplified by egg feeding, "non-infanticide" as exemplified by the absence

of observed feeding, and "infanticide" as shown by physical injury to the tadpole without

deposition of a second tadpole, which was only observed in the displacement manipulation (**Fig.**

1C).

Table 1. Final model fit of eggs fed with offspring manipulations. SD = Standard Deviation. SE = Standard

Error, z = standardized Z value, Pr() = P value.

Random Effects	Variance	SD		
Tank	0.23	0.48		
Fixed Effects	Estimate	SE	z	Pr (>|z|)
Intercept	0.40	0.64	0.63	0.53
Different Morph	-0.50	0.50	-1.01	0.31
Control	0.41	0.31	1.32	0.19
Displacement	-1.40	0.53	-2.66	**7.80E-03**

| | Siblings | -0.15 | 0.19 | -0.78 | 0.43 |

3.2 Infanticide in new territory

After concluding that offspring location is an important factor in parental decisions of care versus infanticide, we performed a "takeover" behavioral assay to test for hormonal correlates of infanticide. We analyzed corticosterone concentration, testosterone concentration, and ratios of testosterone/corticosterone with respect to outcome ("infanticide" versus "non-infanticide"), sampling event ("baseline", "move", and "final"), and interactions between outcome and event themselves **(Fig. 2A-B)**.

Figure 2. Takeover setup and behavioral outcomes. (A) Individual adult frogs were displaced from their home terraria to novel terraria along with a single, unrelated, fertilized egg or hatchling placed on a leaf. Trials lasted for a maximum of seven days. Hormones were collected via water-borne hormone sampling on the day of the move (Day 0), one day afterward (Day 1), and upon observation of infanticidal behavior, or on Day 7 if no such behavior occurred. **(B)** Behavioral outcomes towards eggs or recently hatched tadpoles as observed on motion-trigger Wyze v3 cameras included "infanticide" in the form of cannibalism or physical disturbance to the egg, or "non-infanticide" where these behaviors were not

observed, including one observation of transport by a female. **(C)** Females exhibited both outcomes (non-

infanticide: n = 6, infanticide: n = 6), as did males (non-infanticide: n = 8, infanticide: n = 4).

Of the twelve trials performed with females, six of the twelve females fell within the

"infanticide" outcome, while the other five were categorized as ignoring the conspecific young.

One individual demonstrated parental care, performing tadpole transport (**Video S3**). As we only

observed transport by one individual, we did not have enough statistical power to analyze this

result distinct from other non-infanticidal females, and therefore we placed them in the "non-

infanticide" category. We further rationalized this choice with the observation that concentrations

of circulating androgens and glucocorticoids do not significantly differ between females that do

or do not perform transport in other poison frogs (Fischer and O'Connell, 2020). Of the twelve

trials performed with males, eight showed no infanticidal behavior towards conspecific young

while four showed infanticidal behavior (**Fig. 2C**). Latency to infanticide decisions were variable,

with the behavior occurring on a range of days, from the day after movement to the new

terrarium to the full seven days (**Fig. S1**).

3.3 Infanticide and corticosterone

Log corticosterone (corticosterone hereafter) concentrations did not vary significantly

between infanticidal versus non-infanticidal adults (**Table 2**, η_p^2 = 0.01, t = -0.47, p = 0.65; **Fig.**

3A, Table S7) nor did they vary significantly between males and females (**Table 2,** η_p^2 = 0.17, t

= 1.99, p = 0.06; **Fig. 3A, Table S7**). On average, corticosterone levels increased during

takeover of novel territory (η_p^2 = 0.17, t = 2.45, p = 0.02, **Table 2**). Model diagnostics may be

found in **Fig S3** with analysis of deviance details in **Table S6-7.** In pairwise analyses across

sampling events, corticosterone levels significantly decreased from "move" to "final" sampling

days (Cohen's *d* = 0.83, t = 2.46, p_{adj} = 0.047; **Table S3**). No other pairwise comparisons

between behavioral outcomes and sampling events were significant (**Table S3**). In summary,

corticosterone levels changed the most between 24 hours into a takeover and subsequent

sampling periods; this change was consistent between males and females, as well as between

infanticidal and non-infanticidal adults.

Figure 3. Hormone concentrations by event and behavioral outcome. (A) Corticosterone and **(B)**

testosterone concentrations by sampling events and behavioral outcomes. **(C)** Testosterone

concentrations of infanticidal individuals 24 hours after being moved into novel territory by day of

infanticide. There is a moderate, but not significant, correlation between testosterone and day of

infanticide (Pearson's product-moment correlation r = 0.48, p = 0.18). **(D)** Natural log of

testosterone:corticosterone concentration ratios by sampling events and behavioral outcomes. The "X" in

each box plot represents the average. The horizontal lines represent the medians of each group. All

results are shown in raw units. Signif. codes: 0 < *** < 0.001 < ** < 0.01 < * < 0.05 < . < 0.1 < ns < 1.

Table 2. Final corticosterone model fit of log(corticosterone) by sampling event, behavioral outcome, and

sex. Frog identity, frog length, and average mass are random effects. SD = Standard Deviation. SE =

Standard Error, t = t-statistic, Pr() = P value.

Random Effects	Variance	*SD*			
Body length	0.02	0.13			
Frog identity	0.03	0.17			
Average mass across trial	0.01	0.12			
Residual	0.08	0.29			
Fixed Effects	Estimate	*SE*	df	*t*	Pr (>|t|)
Intercept	6.25	0.12	31.14	52.52	**<2.00E-16**
Outcome	-0.06	0.13	18.56	-0.47	0.65
Move	0.22	0.09	40.02	2.45	**0.02**
Final	-0.03	0.09	43.24	-0.28	0.78
Sex	0.25	0.13	19.86	1.99	0.06

3.4 Infanticide and testosterone

Across all samples, the inverse transformation of testosterone (testosterone hereafter)

concentration varied significantly both with sampling events (η_p^2 = 0.28, p=0.01, F=5.02, **Fig.**

3B, Table S7). Testosterone concentrations also changed between behavioral outcomes (η^2 =

0.27, p=0.02, F = 6.43, **Fig. 3B, Table S7**), but were not sex-specific (η_p^2 = 0.02, p=0.49,

F=0.50, **Fig. 3B, Table S7**). Infanticidal individuals, across their trial, averaged about ⅔ of the

testosterone levels (67.94 pg/mL) of non-infanticidal individuals (99.44 pg/mL). In other words,

testosterone decreased, on average, during potentially stressful events (baseline compared to

move), and during infanticidal behaviors. Most notably, testosterone levels significantly

decreased 24 hours after takeover onset ($t = 3.45$, $p < 0.01$), but the extent of this decrease was

significantly dependent on whether or not the individual performed infanticide ($\eta_p^2 = 0.24$, $t = -$

2.45, $p = 0.02$, **Table 3**). Model diagnostics may be found in **Fig S4** with full analysis of

deviance details in **Table S9.**

Table 3. Final testosterone model fit of 1/(testosterone) by sampling event, behavioral outcome, and sex. Frog body length, identity, and water collection volume are random effects. SD = Standard Deviation. SE = Standard Error, t = t-statistic, Pr (>|t|) = P value.

Random Effects	Variance	*SD*			
Body length	6.44E-06	2.53E-03			
Frog Identity	2.03E-05	4.50E-03			
Water volume	2.74E-05	5.24E-03			
Residual	8.24E-06	2.87E-03			
Fixed Effects	Estimate	*SE*	df	*t*	**Pr (>|t|)**
Intercept	1.60E-02	2.58E-03	29.17	6.18	**9.6E-07**
Outcome	-3.96E-03	2.58E-03	27.36	-1.54	0.14
Move	6.03E-03	1.75E-03	23.50	3.45	**2.16E-03**
Final	3.31E-03	1.88E-03	22.83	1.76	0.09
Sex	-1.64E-03	2.32E-03	19.64	-0.71	0.49
Move : Outcome	-5.06E-03	2.07E-03	19.40	-2.45	**0.02**
Final : Outcome	-6.14E-04	2.32E-03	23.03	-0.27	0.79

In pairwise comparisons, we examined testosterone levels relating to interactions between sampling events and outcomes. We found that, within infanticidal adults, levels of testosterone between baseline hormone sampling and moving to a new territory were significantly different (Cohen's d = -2.08, t = -3.21, p = 0.03, Tukey's method for comparing family of 6 estimates, **Fig. 3B; Table S4**). Similarly, baseline testosterone levels in adults who did not perform infanticide were significantly greater than testosterone concentrations in infanticidal individuals moving into a new territory (Cohen's d = 3.48, t ratio = 3.57, p = 0.01, Tukey's method for comparing family of 6 estimates, **Fig. 3B; Table S4**). Finally, infanticidal and non-infanticidal individuals have different testosterone levels when moved into a new territory (average of 50.62 pg/mL and 88.26 pg/mL, respectively; Cohen's d = 3.15, t ratio = 3.32, p = 0.026, Tukey's method for comparing family of 6 estimates, **Fig. 3B; Table S4**). Within infanticidal individuals, we found a moderately positive (r = 0.48), albeit statistically insignificant, correlation between testosterone levels 24 hours into the territory takeover assay and latency to infanticide (p = 0.19; **Fig 3C**), which further suggests a relationship between lower testosterone and infanticide.

3.5 Infanticide and testosterone:corticosterone ratios

Finally, we examined whether the (natural) log ratios of testosterone to corticosterone (ratios hereafter) changed across the takeover sampling events or between infanticidal and non-infanticidal individuals. The ratio decreased from baseline to move (η_p^2 = 0.38, t = -5.04, p < 1.22E-05, **Table 4**) and between outcomes (η_p^2 = 0.24, t = 2.31, p < 0.03, **Table 4**). Similar to corticosterone and testosterone, ratios were not sex-specific in their changes (η_p^2 = 0.07, t = -1.22, p < 0.24, **Table 4**). Model diagnostics may be found in **Fig S5** with analysis of deviance details in **Table S11.** In pairwise comparisons, we examined testosterone:corticosterone ratios relating to interactions between sampling events and outcomes. We found that, regardless of behavioral outcome, ratio levels changed from baseline to 24 hours into the territory takeover

significantly changed (Cohen's *d* = 1.56, t ratio = 4.91, p < 0.001, **Fig. 3D**, **Table S5**). Likewise,

ratio levels of infanticidal individuals moving to a new territory were significantly different from

baseline ratios and final ratios of individuals that did not perform infanticide (Cohen's *d* = -2.87

and -2.25, t ratio = -4.31 and -3.31, p < 0.01 and 0.05, respectively, **Fig 3D, Table S5**).

Table 4. Final ratio model fit of log(ratio) by sampling event, behavioral outcome, and sex. Frog body

length and identity are random effects.

Random Effects	Variance	*SD*			
Body length	0.06	0.24			
Frog Identity	0.14	0.36			
Residual	0.12	0.35			
Fixed Effects	Estimate	*SE*	df	*t*	Pr (>|t|)
Intercept	-1.95	0.18	27.36	-10.80	**2.29E-11**
Outcome	0.45	0.20	16.81	2.30	**0.03**
Move	-0.54	0.11	38.90	-5.04	**1.10E-05**
Final	-0.22	0.12	42.23	-1.88	0.07
Sex	-0.24	0.20	20.67	-1.20	0.24

4. Discussion

In this investigation, we first delineated tadpole-based features for parental care versus

infanticide, identifying location as a salient kin recognition cue. Then, we documented adult

infanticidal behavior towards eggs and tadpoles for the first time in *Ranitomeya* frogs. By

analyzing concentrations of water-borne steroid hormones corticosterone and testosterone, we

demonstrated that lower testosterone concentrations and testosterone:corticosterone ratios

precede infanticide.

 4.1 Parental decision-making in familiar and unfamiliar territories

We observed that displaced tadpoles experienced significant decreases in trophic egg

deposition compared to control tadpoles, unlike tadpoles cross-fostered within or between

morphs **(Fig. 1B)**. Similar to other poison frogs (Stynoski, 2009; Ringler et al., 2016; Ringler et

al., 2017), *R. imitator* parents likely discriminate between young on the basis of location. For

example, in the brilliant-thighed poison frog (*Allobates femoralis*), males care for any clutch in

their home terraria but cannibalize clutches upon establishing new territories, and females

perform care based on the exact location of offspring (Ringler et al., 2016; Ringler et al., 2017).

Offspring location therefore appears to be a critical cue driving parental care across poison

frogs despite diversity in care systems and ecological constraints. Parental care triggered by

cues of relatedness has also been observed beyond amphibians (Klug and Bonsall, 2007).

Critical decisions between care or cannibalism are especially abundant in anamniotes and

squamates (Ray and Maruska, 2023). For example, spawning appears to impede cannibalism in

parenting African *Neolamprologus caudopunctatus* cichlids, resulting in the care of foreign

young (Cunha-Saraiva et al., 2018), whereas South American Midas cichlids (*Amphilophus*

citrinellus) require olfactory recognition of offspring and will adopt unrelated fry with the same

odor (McKaye and Barlow, 1976). Similarly, brooding children's pythons (*Antaresia children*) will

provide care when eggs are cross-fostered or even replaced with stones (Brashears and

DeNardo, 2012).

A proportion of displaced tadpoles were found dead shortly after being moved to a new

location in their home terrarium, which we interpreted to be caused by infanticide (**Fig. 1C**).

These offspring did not encounter any other tadpoles, making intra-sibling aggression unlikely,

despite being well documented in *Ranitomeya* (Brown et al., 2009; Schulte and Mayer, 2017;

McKinney et al., 2022). Offspring cannibalism as a behavioral consequence of intraspecific

competition has been previously suggested in other poison frogs (Summers, 1989), where

adults might commit infanticide of unrecognized offspring to prevent misdirected care, to

promote survival of their own offspring by parasitizing occupied deposition sites, or to prevent

offspring cannibalism by unrelated tadpoles. Other species of *Ranitomeya* have been found to

parasitize the care of other parents, another indicator that brood parasitism is a weapon

between conspecific competitors (Poelman and Dicke, 2007; Brown et al., 2009, respectively).

To understand why tadpoles were killed rather than eaten by adults, we theorize that dead

tadpoles may yield a precious protein source for cannibalistic *R. imitator* tadpoles that adults

may deposit. Protein can otherwise be difficult to obtain in low resource environments and may

minimize future costs of females having to provision unfertilized egg meals to nutritionally needy

tadpoles (Yoshioka et al., 2016). Thus, we hypothesize that infanticide by *R. imitator* adults

evolved for parents to maximize their individual reproductive success and opportunities while

minimizing those of potential competitors, following evaluation of their environments.

We quantified the behavioral responses of reproductive adults to unrelated, conspecific

young in an unfamiliar environment and found that both males and females cannibalized the

stimulus. This marks the first documented report of adults cannibalizing young in the

Ranitomeya genus and supports previous reports of non-filial cannibalism in other poison frogs

(Townsend et al., 1984; Summers, 1989; Ringler et al., 2017; Spring et al., 2019; Dugas et al.,

2023). We considered multiple adaptive rationales for non-filial cannibalism in *R. imitator.* First,

cannibalism could be a behavioral decision relevant to a trade-off between acquiring caloric

content (which risks eating offspring) and providing care (which risks caring for unrelated

young), an idea which has been tested extensively in fish (Bose, 2022). Fertilized conspecific

eggs constitute part of other frog diets (Beard, 2007). Furthermore, feeding behavior appears

mechanistically linked to parental status across taxa including mammals, birds, and fish

(O'Rourke and Renn, 2015; Fischer and O'Connell, 2017). Thus, we reasoned that some frogs

might be more likely to cannibalize the stimulus egg or hatchling based on differences in how

recently they parented, although all subjects produced a tadpole within the previous 30 days. To

account for various degrees of hunger and potential effects of parental status, frogs were

supplied with food *ad libitum*, so it is unlikely that the frogs in the present study cannibalize to

compensate for nutritional deficits. Therefore, it appears more likely that cannibalism occurred

as a response to intraspecific competition rather than as a means of acquiring food.

In our experimental paradigm, we disrupted pair bonds; therefore, cannibalism of

conspecific, unrelated young might occur to procure access to new mates. Sexually selected

infanticide is well-documented in mammals, birds, and insects (Palombit, 2015), and some

evidence exists for this rationale exists in other poison frogs (Summers, 1989; Spring et al.,

2019). For instance, *A. femoralis* females are less likely to cannibalize unrelated young in the

presence of a male with whom they produced a clutch (Spring et al., 2019). While *R. imitator* are

usually monogamous, mate poaching is nonetheless a proposed rationale for male aggression

in this species (Brown et al., 2009). Another area for future exploration is clarifying the

importance of territoriality, as *A. femoralis* males demonstrate a nearly universal switch to clutch

cannibalism in new territories, while caring for those in their home environment (Ringler et al.,

2017). As infanticide was still performed by a minority of frogs in the takeover assay, it is worth

clarifying the importance of territoriality relative to other potential decision-making cues in this

species, namely nutritional need, mate access, parental experience, and offspring size or age.

4.2 Hormonal correlates of infanticide

To clarify whether physiological stress correlates with infanticide, we analyzed

cannibalistic behavior and sampling events in relation to corticosterone levels. Parental

glucocorticoid responses to social instability have mostly been studied in mammals, which

presents a logistical challenge for comparison to amphibians, as glucocorticoids are also

involved in lactation (Engh et al., 2006). Still, we expected to find that cannibalistic adults would

exhibit greater corticosterone concentrations upon displacement to new territory, as

glucocorticoids increase when encountering stressful and challenging environments (Sapolsky

et al., 2000). Additionally, we sought to test the idea that adults which underwent this

physiological stress would be more likely to perform infanticide. Instead, corticosterone

concentrations were not significantly associated with infanticide at any sampling event, but

significantly changed over the course of the trial regardless of behavioral outcomes (**Fig. 3A**).

Elevated glucocorticoids have been proposed to promote both avoidant and approaching

behaviors (Terburg et al., 2009), which may relate to decisions to cannibalize and ignore

unrelated offspring, respectively. It is also possible that the stress of displacement masks any

corticosterone-specific effects associated with cannibalism. Based on our observations, we

conclude that although physiological stress responses in *R. imitator* do change with time in a

novel territory, the behavioral reactions to this stress likely depend more so on non-

corticosteroid pathways, potentially involving other steroids or neuromodulators.

Previous investigations have shown conflicting results regarding the relationship

between testosterone levels and infanticide (Rosenberg and Sherman, 1975; Martínez et al.,

2015). However, circulating androgens are classically associated with territorial aggression in

mammals, birds, and recently, other poison frogs (Wingfield et al., 1990; Duque-Wilckens et al.,

2019; Rodriguez et al., 2022). Therefore, we expected to find that frogs which performed

infanticide upon displacement to a new territory would exhibit greater levels of testosterone.

Contrary to what we expected, circulating testosterone appears to be negatively associated with

infanticide in this species (**Fig. 3B**); in fact, latency to cannibalism showed a positive

relationship to testosterone levels the day after territory takeover begins (**Fig. 3C**). Our finding

aligns with investigations in some fish, where plasma 11-ketotestosterone concentrations were

lower in cannibals and comparable to those typical of the parental phase (Takegaki et al., 2023).

Interestingly, low testosterone is also associated with parental care in male poison frogs

(Townsend and Moger, 1987); therefore, it seems plausible that decreased androgens can be

associated with both care and infanticide in amphibian parents.

Moreover, local aromatization of testosterone to estradiol has been linked to aggression

in several mammals, fish, and birds (Trainor et al., 2006; Huffman et al., 2013). The conversion

of testosterone to estradiol by aromatase is a highly conserved process in vertebrates, including

amphibians (Coumailleau et al., 2015), and the co-option of estrogens rather than androgens to

facilitate aggression can potentially avoid the costs of elevated testosterone, which can be

especially detrimental in parents (Wingifeld et al., 2001). Investigations in mice have suggested

that estrogen receptor α expressing (*ESR1+)* neurons are necessary and sufficient for

infanticide (Hashikawa et al., 2017; Liu et al., 2022; Mei et al., 2023). Within birds, estrogens are

thought to increase aggression more broadly, as male aromatase activity is positively

associated with aggressive pecking responses, aggressive song production, and chasing

behavior (Schlinger and Callard, 1989; 1990; Soma et al., 2000). However, estrogen signaling

has yet to be adequately explored in any form of amphibian aggression.

Similar to testosterone, testosterone:corticosterone ratios changed significantly from

baseline to move in the overall sample, where lower ratios were associated with cannibalism.

Therefore, testosterone:corticosterone ratios may be similarly reflective of environmental

perturbations and the behavioral niches that poison frogs occupy to cope with a perceived social

challenge. However, the interpretation of our result is impeded by the dearth of exploration of

androgen:glucocorticoid ratios in amphibians. Our result may hint at a currently uncharted

avenue for investigating the importance of steroidal ratios in infanticidal behavior, or it might

only reflect the association of cannibalism with low testosterone. Because testosterone and

cortisol/corticosterone canonically regulate each other's biological functions as terminals of the

HPG and HPA axes, respectively, interactions between the two may inform behavioral

outcomes, as has been suggested previously for even non-mammals, such as fish (Agues-

Barbosa et al., 2022). However, the trend observed here is opposite of that reported in the

largely mammalian literature, where higher ratios are proposed to represent proneness to social

aggression in humans (Terburg et al., 2009). Reasons for this distinction may include those

previously listed with respect to testosterone and corticosterone individually, and the lack of

studies in this field which address non-mammalian taxa or aggression directed towards young.

A promising future avenue for dissecting the physiology of parental care versus

infanticide in poison frogs lies in the concerted analysis of steroid hormones and neural activity.

Oxytocin and vasopressin are well-known for their roles in mammalian parenting (Feldman and

Bakermans-Kranenberg, 2017), but their homologs do not appear to mediate parental care in *R.*

imitator (Schulte and Summers, 2017). Emerging research on the neural underpinnings of

parental behavior have implicated galanin and *Urocortin-3* in infanticide within cichlids and mice

respectively (Butler et al., 2020; Autry et al., 2021), while galanin is also associated with

parental care in both cichlids and mice, in addition to poison frogs (Wu et al. 2014; Fischer et

al., 2019; Butler et al., 2020). This apparent overlap in the neural circuitry responsible for

seemingly antagonistic parental behaviors merits greater consideration of peripheral mediators.

An emerging area of research centers the principle that steroid hormones can alter sensory

representations and information coding by directly acting on neural circuits (Remage-Healey,

2014). Therefore, biological underpinnings of parental behavior in this species might be more

aptly investigated in terms of interactions between steroids and neuropeptides, rather than

studying either in isolation. This seems especially plausible for infanticide, a behavior which

appears to be guided by a striking interplay between physiological and environmental inputs.

5. Conclusions

In this study, we aimed to clarify environmental and hormonal cues for parental care and

infanticide in mimic poison frogs. Our results suggest that, consistent with other poison frogs,

care and infanticide in this species are antagonistically linked on the basis of a simple external

cue of offspring location. Although infanticide has been observed in other poison frogs, this is

the only such report in a monogamous or biparental amphibian, which indicates that infanticide

by both sexes occurs regardless of mating or parenting systems in this clade. We also showed

that mimic poison frogs perform infanticide in both familair and unfamiliar territories, wherein

they targeted related and unrelated young, respectively. Based on the ecological history of *R.*

imitator, we posit that infanticide in this species serves to prevent misdirected care and

eliminate intraspecific competition for limited resources. Moreover, our hormonal analyses

indicate that low concentrations of circulating androgens can precede infanticide following social

and environmental perturbations, which contributes a unique perspective to the broader

aggression literature. Overall, these findings offer fresh insights into how adults adjust their

behavior and physiology to make life-or-death decisions towards offspring in the face of social

instability. In the future, a concerted analysis of endocrine and neural activities compared

between individuals performing infanticide, neglect, and care is a promising next direction to

uncover evolutionary innovations in physiology underlying offspring-directed behaviors.

Funding

This work was supported by a McKnight Pecot Fellowship to ARL and grants from the National

Institutes of Health (DP2HD102042) and the New York Stem Cell Foundation to LAO. ARL was

additionally supported by the Stanford Bio-X Undergraduate Research Fellowship and grants

from the Office of the Vice Provost for Undergraduate Education. BCG was supported by a

HHMI Gilliam Fellowship (GT15685) and a National Institutes of Health Cellular Molecular

Biology Training Grant (T32GM007276). BHJ was supported by funding from the New York

Stem Cell Foundation. LAO is a New York Stem Cell Foundation–Robertson Investigator.

CRediT authorship contribution statement

Amaris R. Lewis: Conceptualization, Methodology, Formal Analysis, Investigation, Data

Curation, Writing - Original Draft, Writing - Review and Editing, Visualization, Project

Administration, Funding Acquisition. **Billie C. Goolsby:** Conceptualization, Methodology,

Formal Analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review and

Editing, Visualization, Project Administration. **Bryan H. Juarez:** Methodology, Formal Analysis,

Resources, Writing - Original Draft, Writing - Review and Editing, Visualization, Supervision.

Madison P. Lacey: Investigation, Resources, Writing - Review and Editing. **Lauren A.**

O'Connell: Resources, Writing- Review & Editing, Funding Acquisition, Supervision**.**

Acknowledgments

This research was conducted at Stanford University, which is located on the ancestral and

unceded land of the Muwekma Ohlone tribe. We thank Dr. Camilo Rodríguez Lopez and Dr.

Ricardo Cossio for feedback on this manuscript. We are additionally grateful to CRL for

guidance on steroid hormone collection and processing. Finally, we thank the members of the

Laboratory for Organismal Biology for providing routine frog care and continuous support.

Data availability

Data and code used in this study can be found on DataDryad at

https://doi.org/10.5061/dryad.1c59zw43m. The link is currently private but can be accessed for

peer review at https://datadryad.org/stash/share/6hImFvb9nW-

bq0gXHuDj0G230tCxjFfd3LolYfr9I0M.

References

Agues-Barbosa, T., Andrade, P. V. C., Silva, P. F., Moura, C. de A., Galvão, N. L.,

664   Freire, F. A. M., & Luchiari, A. C. (2022). Variation in nest building, aggression, learning,

665   and steroid hormone levels in Betta splendens. *General and Comparative*

666   *Endocrinology*, *323–324*, 114044. https://doi.org/10.1016/j.ygcen.2022.114044

667   Alonso-Alvarez, C., & Velando, A. (2012). Benefits and costs of parental care. *The evolution of*

668   *parental care*, 40-61. https://doi.org/10.1093/acprof:oso/9780199692576.003.0003

669   Autry, A. E., Wu, Z., Kapoor, V., Kohl, J., Bambah-Mukku, D., Rubinstein, N. D., Marin-

670   Rodriguez, B., Carta, I., Sedwick, V., Tang, M. and Dulac, C. (2021). Urocortin-3

neurons in the mouse perifornical area promote infant-directed neglect and aggression.

Elife, *10*, e64680. https://doi.org/10.7554/eLife.64680

Bales, K. L., Kramer, K. M., Lewis-Reese, A. D., & Carter, C. S. (2006). Effects of stress on

parental care are sexually dimorphic in prairie voles. *Physiology & behavior*, *87*(2), 424-

429. https://doi.org/10.1016/j.physbeh.2005.11.002

Baugh, A. T., Bastien, B., Still, M. B., & Stowell, N. (2018). Validation of water-borne steroid

hormones in a tropical frog (Physalaemus pustulosus). *General and Comparative*

Endocrinology, *261*, 67-80. https://doi.org/10.1016/j.ygcen.2018.01.025.

Baugh, A. T., & Gray-Gaillard, S. L. (2021). Excreted testosterone and male sexual proceptivity:

a hormone validation and proof-of-concept experiment in túngara frogs. *General and*

Comparative Endocrinology, *300*, 113638. https://doi.org/10.1016/j.ygcen.2020.113638

Beard, K. (2007). Diet of the invasive frog, *Eleutherodactylus coqui*, in Hawaii. *Copeia, 2007*(2),

281-291. https://doi.org/10.1643/0045-8511(2007)7[281:DOTIFE]2.0.CO;2

Ben-Shachar, M. S., Lüdecke, D., & Makowski, D. (2020). effectsize: Estimation of effect size

indices and standardized parameters. *Journal of Open Source Software*, 5(56), 2815.

https://doi.org/10.21105/joss.02815

Bose, A. P. (2022). Parent–offspring cannibalism throughout the animal kingdom: a review of

adaptive hypotheses. *Biological Reviews*, *97*(5), 1868-1885.

https://doi.org/10.1111/brv.12868

Brashears, J., & DeNardo, D. F. (2012). Do Brooding Pythons Recognize their Clutches?

Investigating External Cues for Offspring Recognition in the Children's Python, *Antaresia*

childreni . *Ethology*, *118*(8), 793–798. https://doi.org/10.1111/j.1439-0310.2012.02070.x

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R*

Journal, *9*(2), 378. https://doi.org/10.32614/RJ-2017-066

Brown, J., Twomey, E., Morales, V., & Summers, K. (2008). Phytotelm size in relation to

parental care and mating strategies in two species of Peruvian poison frogs. *Behaviour*,

145(9), 1139-1165. https://doi.org/10.1163/156853908785387647

Brown, J. L., Morales, V., & Summers, K. (2009). Tactical reproductive parasitism via larval

cannibalism in Peruvian poison frogs. *Biology Letters*, *5*(2), 148-151.

https://doi.org/10.1098/rsbl.2008.0591

Burnham, K. P., & Anderson, D. R. (Eds.). (2004). *Model Selection and Multimodel Inference*.

Springer New York. https://doi.org/10.1007/b97636

Butler, J. M., Herath, E. M., Rimal, A., Whitlow, S. M., & Maruska, K. P. (2020). Galanin neuron

activation in feeding, parental care, and infanticide in a mouthbrooding African cichlid

fish. *Hormones and Behavior*, *126*, 104870. https://doi.org/10.1016/j.yhbeh.2020.104870

Clutton-Brock, T. H. (1991). *The evolution of parental care* (Vol. 64). Princeton University Press.

10.1093/acprof:oso/9780199692576.001.0001

Cockrem, J. F. (2013). Individual variation in glucocorticoid stress responses in animals.

General and Comparative Endocrinology, 181, 45-58.

https://doi.org/10.1016/j.ygcen.2012.11.025

Coumailleau, P., Pellegrini, E., Adrio, F., Diotel, N., Cano-Nicolau, J., Nasri, A., ... & Kah, O.

(2015). Aromatase, estrogen receptors and brain development in fish and amphibians.

Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms, 1849(2), 152-162.

https://doi.org/10.1016/j.bbagrm.2014.07.002

Crossin, G. T., Trathan, P. N., Phillips, R. A., Gorman, K. B., Dawson, A., Sakamoto, K. Q., &

Williams, T. D. (2012). Corticosterone predicts foraging behavior and parental care in

macaroni penguins. *The American Naturalist*, *180*(1), E31-E41.

https://doi.org/10.1086/666001

Cunha-Saraiva, F., Balshine, S., Wagner, R. H., & Schaedelin, F. C. (2018). From cannibal to

caregiver: tracking the transition in a cichlid fish. *Animal Behaviour*, 139, 9–17.

https://doi.org/10.1016/j.anbehav.2018.03.003

Dugas, M. B., Brooks, O. L., Saporito, R. A., & Cossio, R. (2023). Adult poison frogs can

capture and consume aquatic tadpoles. *Evolutionary Ecology*, 1-7.

https://doi.org/10.1007/s10682-023-10257-1

Ebensperger, L. A., & Blumstein, D. T. (2007). Nonparental infanticide. *Rodent societies: an*

ecological and evolutionary perspective, 267-279.

Ellington, E. H., Bastille☐Rousseau, G., Austin, C., Landolt, K. N., Pond, B. A., Rees, E. E.,

Robar, N., & Murray, D. L. (2015). Using multiple imputation to estimate missing data in

meta☐regression. *Methods in Ecology and Evolution*, 6(2), 153–163.

https://doi.org/10.1111/2041-210X.12322

Elwood, R. W., & Stolzenberg, D. S. (2020). Flipping the parental switch: from killing to caring in

male   mammals.   *Animal   Behaviour*,   *165*,   133-142.

https://doi.org/10.1016/j.anbehav.2020.05.001

Engh, A. L., Beehner, J. C., Bergman, T. J., Whitten, P. L., Hoffmeier, R. R., Seyfarth, R.

M., & Cheney, D. L. (2006). Female hierarchy instability, male immigration and

infanticide increase glucocorticoid levels in female chacma baboons. *Animal Behaviour*,

71(5), 1227–1237. https://doi.org/10.1016/j.anbehav.2005.11.009

Feldman, R., & Bakermans-Kranenburg, M. J. (2017). Oxytocin: a parenting hormone. *Current*

Opinion in Psychology, *15*, 13-18. https://doi.org/10.1016/j.copsyc.2017.02.011

Fischer, E. K., & O'Connell, L. A. (2017). Modification of feeding circuits in the evolution of social

behavior.   *Journal   of   Experimental   Biology*,   *220*(1),   92-102.

https://doi.org/10.1242/jeb.143859

Fischer, E. K., Roland, A. B., Moskowitz, N. A., Tapia, E. E., Summers, K., Coloma, L. A., &

O'Connell, L. A. (2019). The neural basis of tadpole transport in poison frogs.

Proceedings of the Royal Society B, 286(1907), 20191084.

https://doi.org/10.1098/rspb.2019.1084

Fischer, E. K., & O'Connell, L. A. (2020). Hormonal and neural correlates of care in active

versus observing poison frog parents. *Hormones and Behavior, 120*, 104696.

https://doi.org/10.1016/j.yhbeh.2020.104696

Gabor, C. R., Bosch, J., Fries, J. N., & Davis, D. R. (2013). A non-invasive water-borne

hormone assay for amphibians. *Amphibia-Reptilia, 34*(2), 151-162.

https://doi.org/10.1163/15685381-00002877

Goll, Y., Bordes, C., Weissman, Y. A., Shnitzer, I., Beukeboom, R., Ilany, A., Koren, L., &

Geffen, E. (2023). The interaction between cortisol and testosterone predicts leadership

within rock hyrax social groups. *Scientific Reports, 13*(1), 14857.

https://doi.org/10.1038/s41598-023-41958-w

Goolsby, B. C., Fischer, M. T., Pareja-Mejia, D., Lewis, A. R., Raboisson, G., & O'Connell, L. A.

(2023). Home security cameras as a tool for behavior observations and science equity.

bioRxiv. https://doi.org/10.1101/2023.04.17.537238

Gosner, K. L. (1960). A Simplified Table for Staging Anuran Embryos and Larvae with Notes on

Identification. *Herpetologica, 16*(3), 183–190.

Haller, J. (2014). The glucocorticoid/aggression relationship in animals and humans: an analysis

sensitive to behavioral characteristics, glucocorticoid secretion patterns, and neural

mechanisms. *Neuroscience of aggression*, 73-109.

https://doi.org/10.1007/7854_2014_284

Hartig F (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.6.

Hashikawa, K., Hashikawa, Y., Tremblay, R., Zhang, J., Feng, J. E., Sabol, A., Piper, W.T., Lee,

H., Rudy, B., & Lin, D. (2017). Esr1+ cells in the ventromedial hypothalamus control

female aggression. *Nature neuroscience*, 20(11), 1580-1590. https://doi.org/10.1038/nn.4644

Hirschenhauser, K., & Oliveira, R. F. (2006). Social modulation of androgens in male vertebrates: meta-analyses of the challenge hypothesis. *Animal Behaviour*, *71*(2), 265-277. https://doi.org/10.1016/j.anbehav.2005.04.014

Hrdy, S. B. (1979). Infanticide among animals: a review, classification, and examination of the implications for the reproductive strategies of females. *Ethology and Sociobiology*, *1*(1), 13-40. https://doi.org/10.1016/0162-3095(79)90004-9

Huffman, L. S., O'Connell, L. A., & Hofmann, H. A. (2013). Aromatase regulates aggression in the African cichlid fish Astatotilapia burtoni. *Physiology & Behavior*, 112–113, 77–83. https://doi.org/10.1016/j.physbeh.2013.02.004

Klug, H., & Bonsall, M. B. (2007). When to Care for, Abandon, or Eat Your Offspring: The Evolution of Parental Care and Filial Cannibalism. *The American Naturalist*, 170(6), 886–901. https://doi.org/10.1086/522936

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13). https://doi.org/10.18637/jss.v082.i13

Lenth R (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.9.0. https://doi.org/10.1080/00031305.1980.10483031

Liu, M., Kim, D. W., Zeng, H., & Anderson, D. J. (2022). Make war not love: the neural substrate underlying a state-dependent switch in female social behavior. *Neuron*, 110(5), 841-856. https://doi.org/10.1016/j.neuron.2021.12.002

Lukas, D., & Huchard, E. (2014). The evolution of infanticide by males in mammalian societies. *Science*, 346(6211), 841-844. https://doi.org/10.1126/science.1257226

Lukas, D., & Huchard, E. (2019). The evolution of infanticide by females in mammals.

Philosophical Transactions of the Royal Society B, 374(1780), 20180075.

https://doi.org/10.1098/rstb.2018.0075

Martínez, A., Ramos, G., Martínez-Torres, M., Nicolás, L., Carmona, A., Cárdenas, M., &

Luis, J. (2015). Paternal behavior in the Mongolian gerbil (Meriones unguiculatus):

Estrogenic and androgenic regulation. *Hormones and Behavior*, *71*, 91–95.

https://doi.org/10.1016/j.yhbeh.2015.04.009

McKaye, K. R., & Barlow, G. W. (1976). Chemical Recognition of Young by the Midas Cichlid,

Cichlasoma citrinellum. *Copeia*, 1976(2), 276. https://doi.org/10.2307/1443947

McKinney, J. E., Ludington, S. C., Butler, J. M., & O'Connell, L. A. (2022). Proopiomelanocortin

(POMC) is a negative regulator of tadpole aggression through opioid receptor signaling.

bioRxiv, 2022-11. https://doi.org/10.1101/2022.11.28.518266

Mehta, P. H., & Prasad, S. (2015). The dual-hormone hypothesis: a brief review and future

research agenda. *Current Opinion in Behavioral Sciences*, *3*, 163–168.

https://doi.org/10.1016/j.cobeha.2015.04.008

Mei, L., Yan, R., Yin, L., Sullivan, R. M., & Lin, D. (2023). Antagonistic circuits mediating

infanticide and maternal care in female mice. *Nature*, 1-11.

https://doi.org/10.1038/s41586-023-06147-9

Nakagawa, S., & Freckleton, R. P. (2011). Model averaging, missing data and multiple

imputation: a case study for behavioural ecology. *Behavioral Ecology and Sociobiology*,

65(1), 103–116. https://doi.org/10.1007/s00265-010-1044-7

O'Connor, C. M., Gilmour, K. M., Arlinghaus, R., Van Der Kraak, G., & Cooke, S. J. (2009).

Stress and parental care in a wild teleost fish: insights from exogenous

supraphysiological cortisol implants. *Physiological and Biochemical Zoology*, *82*(6), 709-

719. https://doi.org/10.1086/605914

O'Rourke, C. F., & Renn, S. C. (2015). Integrating adaptive trade-offs between parental care

and feeding regulation. *Current opinion in behavioral sciences*, 6, 160-167.

https://doi.org/10.1016/j.cobeha.2015.11.010

Palombit, R. A. (2015). Infanticide as Sexual Conflict: Coevolution of Male Strategies

and Female Counterstrategies. *Cold Spring Harbor Perspectives in Biology*, *7*(6),

a017640. https://doi.org/10.1101/cshperspect.a017640

Poelman, E. H., & Dicke, M. (2007). Offering offspring as food to cannibals: oviposition

strategies of Amazonian poison frogs (Dendrobates ventrimaculatus). *Evolutionary*

Ecology, *21*(2), 215–227. https://doi.org/10.1007/s10682-006-9000-8

Ray, E. J., & Maruska, K. P. (2023). Sensory Mechanisms of Parent-Offspring Recognition in

Fishes, Amphibians, and Reptiles. *Integrative And Comparative Biology*, 63(6), 1168–

1181. https://doi.org/10.1093/icb/icad104

Remage-Healey, L. (2014). Frank Beach Award Winner: Steroids as neuromodulators of

brain circuits and behavior. *Hormones and Behavior*, *66*(3), 552–560.

https://doi.org/10.1016/j.yhbeh.2014.07.014

Ringler, E., Pašukonis, A., Ringler, M., & Huber, L. (2016). Sex-specific offspring discrimination

reflects respective risks and costs of misdirected care in a poison frog. *Animal*

Behaviour, *114*, 173-179. https://doi.org/10.1016/j.anbehav.2016.02.008

Ringler, E., Barbara Beck, K., Weinlein, S., Huber, L., & Ringler, M. (2017). Adopt, ignore, or

kill? Male poison frogs adjust parental decisions according to their territorial status.

Scientific Reports, *7*(1), 43544. https://doi.org/10.1038/srep43544

Rodríguez, C., Fusani, L., Raboisson, G., Hödl, W., Ringler, E., & Canoine, V. (2022). Androgen

responsiveness to simulated territorial intrusions in Allobates femoralis males: Evidence

supporting the challenge hypothesis in a territorial frog. *General and Comparative*

Endocrinology, *326*, 114046. https://doi.org/10.1016/j.ygcen.2022.114046

Rosenberg, K. M., & Sherman, G. F. (1975). The role of testosterone in the organization,

maintenance and activation of pup-killing behavior in the male rat. *Hormones and*

Behavior, *6*(2), 173–179. https://doi.org/10.1016/0018-506X(75)90033-1

Rosvall, K. A., Bentz, A. B., & George, E. M. (2020). How research on female vertebrates

contributes to an expanded challenge hypothesis. *Hormones and Behavior*, 123,

104565. https://doi.org/10.1016/j.yhbeh.2019.104565

Royle, N. J., Smiseth, P. T., & Kölliker, M. (Eds.). (2012). *The evolution of parental care*. Oxford

University Press. https://doi.org/10.1093/acprof:oso/9780199692576.001.0001

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). How do glucocorticoids influence stress

responses? Integrating permissive, suppressive, stimulatory, and preparative actions.

Endocrine Reviews, *21*(1), 55-89. https://doi.org/10.1210/edrv.21.1.0389

Sapolsky, R. M. (2005). The influence of social hierarchy on primate health. *Science*,

308(5722), 648-652. https://doi.org/10.1126/science.1106477

Schlinger, B. A., & Callard, G. v. (1989). Estrogen Receptors in Quail Brain: A Functional

Relationship to Aromatase and Aggressiveness1. *Biology of Reproduction*, *40*(2), 268–

275. https://doi.org/10.1095/biolreprod40.2.268

Schlinger, B. A., & Callard, G. v. (1990). Aromatization mediates aggressive behavior in

quail. *General and Comparative Endocrinology*, *79*(1), 39–53.

https://doi.org/10.1016/0016-6480(90)90086-2

Schulte, L. M., & Mayer, M. (2017). Poison frog tadpoles seek parental transportation to escape

their cannibalistic siblings. *Journal of Zoology*, *303*(2), 83-89.

https://doi.org/10.1111/jzo.12472

Schulte, L. M., & Summers, K. (2017). Searching for hormonal facilitators: Are vasotocin and

mesotocin involved in parental care behaviors in poison frogs?. *Physiology & Behavior*,

174, 74-82. https://doi.org/10.1016/j.physbeh.2017.03.005

Soma, K. K., Tramontin, A. D., & Wingfield, J. C. (2000). Oestrogen regulates male

aggression in the non–breeding season. *Proceedings of the Royal Society of London.*

Series   B: Biological   Sciences,   *267*(1448),   1089–1096.

https://doi.org/10.1098/rspb.2000.1113

Spring, S., Lehner, M., Huber, L., & Ringler, E. (2019). Oviposition and father presence reduce

clutch cannibalism by female poison frogs. *Frontiers in zoology*, *16*, 1-10.

https://doi.org/10.1186/s12983-019-0304-2

Stynoski, J. L. (2009). Discrimination of offspring by indirect recognition in an egg-feeding

dendrobatid frog, *Oophaga pumilio. Animal Behaviour*, *78*(6), 1351-1356.

https://doi.org/10.1016/j.anbehav.2009.09.002

Summers, C. H., Watt, M. J., Ling, T. L., Forster, G. L., Carpenter, R. E., Korzan, W. J., Lukkes,

J.L., & Øverli, Ø. (2005). Glucocorticoid interaction with aggression in non-mammalian

vertebrates: reciprocal action. *European Journal of Pharmacology*, *526*(1-3), 21-35.

https://doi.org/10.1016/j.ejphar.2005.09.059

Summers, K. (1989). Sexual selection and intra-female competition in the green poison-dart

frog, Dendrobates auratus. *Animal Behaviour*, *37*, 797–805.

https://doi.org/10.1016/0003-3472(89)90064-X

Summers, K., & Tumulty, J. (2014). Parental Care, Sexual Selection, and Mating Systems in

Neotropical Poison Frogs. In *Sexual Selection* (pp. 289–320). Elsevier.

https://doi.org/10.1016/B978-0-12-416028-6.00011-6

Takegaki, T., Nakatake, Y., Matsumoto, Y., Suga, K., & Amiya, N. (2023). Early filial

cannibalism in fish revisited: endocrinological constraint, costs of parental care, and

mating possibility. *The American Naturalist*, 201(6), 841-850.

https://doi.org/10.1086/724284

Terburg, D., Morgan, B., & van Honk, J. (2009). The testosterone–cortisol ratio: A hormonal

896   marker for proneness to social aggression. *International Journal of Law and Psychiatry*,

897   *32*(4), 216–223. https://doi.org/10.1016/j.ijlp.2009.04.008

898   Townsend, D. S., & Moger, W. H. (1987). Plasma androgen levels during male parental care in

899   a tropical frog (Eleutherodactylus). *Hormones and Behavior*, *21*(1), 93-99.

900   https://doi.org/10.1016/0018-506X(87)90034-1

901   Townsend, D. S., Stewart, M. M., & Pough, F. H. (1984). Male parental care and its adaptive

902   significance in a neotropical frog. *Animal Behaviour*, *32*(2), 421-431.

903   https://doi.org/10.1016/0018-506X(87)90034-1

904   Trainor, B. C., Kyomen, H. H., & Marler, C. A. (2006). Estrogenic encounters: how interactions

905   between aromatase and the environment modulate aggression. *Frontiers in*

906   *Neuroendocrinology*, *27*(2), 170-179. https://doi.org/10.1016/j.yfrne.2005.11.001

907   Tsuneoka, Y., Tokita, K., Yoshihara, C., Amano, T., Esposito, G., Huang, A. J., Yu, L.M.,

908   Odaka, Y., Shinozuka, K., McHugh, T.J., & Kuroda, K. O. (2015). Distinct preoptic☐BST

909   nuclei dissociate paternal and infanticidal behavior in mice. *The EMBO Journal*, *34*(21),

910   2652-2670. https://doi.org/10.15252/embj.201591942

911   Twomey, E., Mayer, M., & Summers, K. (2015). Intraspecific Call Variation in the Mimic Poison

912   Frog   *Ranitomeya*   *imitator*.   *Herpetologica*,   *71*(4),   252–259.

913   https://doi.org/10.1655/HERPETOLOGICA-D-15-00004

914   Weygoldt, P. (1987). Evolution of parental care in dart poison frogs (Amphibia: Anura:

915   Dendrobatidae). *Journal of Zoological Systematics and Evolutionary Research*, *25*(1),

916   51–67. https://doi.org/10.1111/j.1439-0469.1987.tb00913.x

917   Westrick, S. E., Paitz, R. T., & Fischer, E. K. (2023). Why not both? A case study measuring

918   cortisol and corticosterone in poison frogs. *bioRxiv*, 2023-06.

919   https://doi.org/10.1101/2023.06.19.545597

Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

ISBN 978-3-319-24277-4. https://doi.org/10.1007/978-3-319-24277-4

Wingfield, J. C., Hegner, R. E., Dufty Jr, A. M., & Ball, G. F. (1990). The "challenge hypothesis":

theoretical implications for patterns of testosterone secretion, mating systems, and

breeding strategies. *The American Naturalist*, *136*(6), 829-846.

https://doi.org/10.1086/285134

Wingfield, J. C., Lynn, S. E., & Soma, K. K. (2001). Avoiding the 'costs' of testosterone:

ecological bases of hormone-behavior interactions. *Brain Behavior and Evolution*, 57(5),

239-251. https://doi.org/10.1159/000047243

Waldman, B. (1988). The ecology of kin recognition. *Annual review of ecology and systematics*,

19(1), 543-571. 10.1146/annurev.es.19.110188.002551

Wu, Z., Autry, A. E., Bergan, J. F., Watabe-Uchida, M., & Dulac, C. G. (2014). Galanin neurons

in the medial preoptic area govern parental behaviour. *Nature*, *509*(7500), 325-330.

https://doi.org/10.1038/nature13307

Yoshioka, M., Meeks, C., & Summers, K. (2016). Evidence for begging as an honest signal of

offspring need in the biparental mimic poison frog. *Animal Behaviour*, 113, 1–11.

https://doi.org/10.1016/j.anbehav.2015.12.024