1   **Infanticide is driven by unfamiliarity with offspring location and associated**

2   **with androgenic shifts in mimic poison frogs**

3   Amaris R. Lewis[1†], Billie C. Goolsby[1†], Bryan H. Juarez[1], Madison P. Lacey[1], Lauren A. O'Connell[1]*

4   [1]Department of Biology, Stanford University, Stanford, CA 94305, USA

*To whom correspondence should be addressed:

Lauren A. O'Connell

Department of Biology

Stanford University

371 Jane Stanford Way

Stanford, CA 94305

loconnel@stanford.edu

†These authors contributed equally to this manuscript.

Abstract

Infanticide is widespread across the animal kingdom, but the physiological drivers of

infanticide versus care or neglect are relatively unexplored. Here, we identified salient

environmental and physiological antecedents of infanticide in the mimic poison frog

(*Ranitomeya imitator*), a biparental amphibian in which female parents feed their tadpoles

unfertilized eggs. Specifically, we explored potential environmental cues influencing infant-

directed behavior by evaluating changes in the frequency of food provisioning and tadpole

mortality after either cross-fostering tadpoles between family units or displacing tadpoles within

the terraria of their parents. We found that changes in offspring location reduce care and increase

infanticide. Specifically, parents fed their displaced offspring less and, in some instances, tadpole

mortality increased. We also investigated whether care and infanticide were related to changes in

steroid hormone concentrations in an unfamiliar setting. Infanticide of fertilized eggs and

hatchlings in the new territory included cannibalism and was associated with lower testosterone

concentrations, but not with changes in corticosterone. Overall, our results support earlier

findings that familiarity with offspring location drives parental investment in poison frogs, while

indicating an association between low androgen levels and infanticidal behavior in an amphibian.

Highlights

● Offspring location drives parental decisions of care vs. infanticide.

● In novel territories, adults cannibalize conspecific, unrelated young.

● Lower circulating testosterone in novel territory is associated with infanticide.

1. Introduction

Animal parents exhibit remarkable variation in behavior toward conspecific young, performing parental care (e.g. feeding, protecting, and transporting young), neglect, and infanticide, regardless of kinship (Hrdy, 1979; Royle et al. 2012; Lukas and Huchard, 2014; Lukas and Huchard, 2019). Care requires significant energy investment, rendering parental decision-making sensitive to evolutionary trade-offs dependent on environmental and physiological constraints (Clutton-Brock, 1991; Alonso-Alvarez and Velando, 2012). Thus, parental behavior is often informed by cues that help discriminate between related and unrelated offspring, including indirect identifiers such as location (Stynoski, 2009; Huang and Pike, 2011; Ringler et al., 2016, 2017) and more direct sensory signals like acoustics and smell (Neff and Sherman, 2005; Vergne et al., 2011). While behavioral ecologists have reported numerous adaptive rationales for infanticide across diverse animal groups (Hrdy, 1979; Ebensperger and Blumstein, 2007; Bose, 2022), evidence linking infanticide to hormones is limited outside of mammals. Understanding the physiological mechanisms and evolution of parental responses to offspring requires connecting infanticide to physiological states in a broad range of animal species.

Steroid hormones present a compelling avenue to investigate the physiological basis of infanticide. Infanticide may be considered a form of social aggression, which is often studied in the context of elevated androgens, like testosterone, and elevated glucocorticoids, like corticosterone, especially in males (Summers et al., 2005; Haller, 2014; Wingfield et al., 1990; Hirschenhauser and Oliveira, 2006; Rodríguez et al., 2022). Changes in steroid hormone concentrations that mediate socially aggressive responses can persist even after the behavior occurs (Rodríguez et al., 2022). Within female vertebrates, previous authors have proposed a link

between testosterone, parental care, and aggression in mammals, birds, lizards, and fishes

(Rosvall, 2020), but also see Bentz et al. (2019), where the relationship between testosterone and

aggression in females is less clear. Investigations in both mammalian and non-mammalian taxa

indicate that social aggression may be facilitated by the acute elevation of peripheral

glucocorticoids (Summers et al., 2005; Haller, 2014). Still, the lack of studies clarifying the

physiology of infanticide in non-mammalian taxa prevents the development of a general,

comparative, and predictive framework for decision-making toward young.

Poison frogs display an extraordinary diversity of parental care strategies, including

tadpole transport, maternal egg provisioning, and cannibalism (Weygoldt, 1987; Summers and

Tumulty, 2014). Previous work in poison frogs has suggested that offspring location, territorial

status, and recent reproductive activity can influence decisions to perform care or infanticide

(Stynoski, 2009; Ringler et al., 2016, 2017; Spring et al., 2019). For example, territorial status

determines the performance of tadpole transport or cannibalism in brilliant-thighed poison frog

(*Allobates femoralis*) males, while females transport tadpoles based on recent reproductive

activity and the exact location of offspring within their territory (Ringler et al., 2016, 2017).

These findings have established a framework for investigating environmental cues for infanticide

in other poison frogs, but the hormonal correlates of this behavior remain unexplored.

We used mimic poison frogs (*Ranitomeya imitator*) to better understand how external

cues elicit hormonal shifts and infanticide. Mimic poison frogs are biparental and sexually

monogamous, forming pair bonds that last several months and performing sex-specific parental

roles to cooperate in the care of their eggs and tadpoles (Brown et al., 2008). Males of this

species transport tadpoles after egg hatching and call females to tadpole sites, whereas females

provision trophic eggs to their tadpoles. Additionally, *R. imitator* differ in coloration and

patterning based on geographic distribution, and it is unclear whether morph identity may also

play a role in offspring recognition.

We used two morphs of *R. imitator (Yurimaguas* and *Rapidos)* to identify key

components of offspring recognition, then analyzed adult hormone concentrations during an

environmental challenge. We began by identifying salient recognition cues so that we might later

expose frogs to a challenge in which they were unlikely to misidentify unrelated young as their

own. Based on prior work in poison frogs (Stynoski, 2009; Ringler et al., 2016; Ringler et al.,

2017), we hypothesized that offspring recognition depends more on tadpole location than direct

recognition cues, including morph-specific features. We performed kin recognition assays,

namely cross-fostering within morphs, cross-fostering between morphs, and displacement within

parents' terraria. After measuring egg provisioning and tadpole mortality as proxies for parental

investment, we concluded that offspring location is a critical indirect recognition cue for this

species.

We then leveraged this finding to conduct simulated territory takeover trials, where adults

were individually moved to an unfamiliar environment and simultaneously exposed to unrelated,

conspecific offspring. Our hypothesis is that changes in the concentrations of circulating

testosterone and corticosterone correlate with infanticidal behavior, which would be reflected by

changes in hormone concentrations from baseline to immediately after infanticide. However, it is

important to consider that correlations between hormone levels and behaviors do not indicate

whether hormone levels drive behavior, whether behavior influences hormone levels, or if

another factor is responsible for both Nonetheless, we can glean information on the potentially

bi-directional relationship between hormones and behavior using this paradigm. Based on earlier

findings in other vertebrates, we anticipated that all adults would exhibit elevated testosterone

and corticosterone during this challenge. However, we also predicted that infanticidal adults

would have higher testosterone and corticosterone concentrations compared to non-infanticidal

adults. We reasoned that testosterone and corticosterone should be reflective of perceived social

competition and stress, respectively, and that individuals would react to these challenges by

eliminating unrelated young.

2. Methods

2.1 Animal husbandry

All *R. imitator* eggs and tadpoles in this study were captive-bred in our colony. Adults

were purchased from Indoor Ecosystems between 2019-2021 (Whitehouse, Ohio) or Ruffing's

Ranitomeya between 2021-2023 (Tiffin, Ohio, USA). Breeding pairs were housed in 30 x 30 x

45 cm terraria containing sphagnum moss substrate, driftwood, live plants, egg deposition sites,

and film canisters filled with water for tadpole deposition. Terraria were automatically misted ten

times daily and frogs were fed live *Drosophila* fruit flies dusted with Repashy Calcium Plus

(Oceanside, CA, USA) three times per week and springtails once a week. All procedures in this

study were approved by the Stanford University Animal Care and Use Committee (Protocol

#34242).

2.2 Offspring recognition assays

Tadpoles used in this study were observed in home or foster terraria as described (all with

the same dimensions and substrates). *Ranitomeya imitator* pairs used for this study were required

to have previously successfully raised a tadpole within 31 days of the experiment. Locations of

tadpole deposition were marked on the outside of the terrarium with the date and time to ensure

consistency in placing tadpole canisters in the same location throughout the duration of the

study. Tadpoles were randomly placed into three experimental conditions (**Fig. 1A**): same-morph

cross-foster (n = 16 across 6 terraria), different-morph cross-foster (n = 12 across 5 terraria), or

intra-tank displacement (n = 18 across 7 terraria). Control tadpoles (n = 26 across 11 terraria)

served as a measure of parental investment under normal conditions.

We observed tadpoles in each trial for 14 days and measured trophic egg deposits as a

proxy for parental investment. We randomized treatments across tadpoles, using individuals

below Gosner stage 32 (Gosner, 1960) to avoid counting feeding patterns that resulted from

changes in investments as tadpoles neared metamorphosis. We collected data for every tadpole

under controlled conditions before randomly applying treatments. Sampling multiple (one to

four) tadpoles per terrarium allowed us to capture focal pair variation in parental care.

Tadpoles were always transported in their respective canisters to retain any tadpole-

specific olfactory cues already present in the water and to minimize stress. Tadpoles displaced

within their parents' terraria ("intra-tank displacement") were moved at a minimum of 5 cm and

maximum of 30 cm, depending on availability on the terrarium floor. We purposely selected a

minimum distance greater than 2 cm, as this was the minimum distance for indirect offspring

discrimination identified in other poison frogs (Stynoski, 2009). Tadpoles were moved either to a

laterally opposite corner of the terrarium or elsewhere along the terrarium edge. When

undergoing this manipulation, tadpoles were always moved to a location not previously occupied

by a canister.

2.3 "Takeover" behavioral assays

We designed an assay to identify hormonal changes in infanticidal versus non-

infanticidal parents during a simulated territory takeover based on Ringler et al. (2017). We

selected paired adults that raised at least one offspring in their terrarium within 31 days of the

takeover trial as a proxy for parental experience and reproductive status. Frogs from the same

reproductive pair were put into separate takeover trials at the same time to mitigate any risk of

mate absence influencing behavior directed toward offspring. The takeover terrarium had the

same dimensions, with elements of the environment (vegetation, replacement of dead leaves and

substrate, fresh water) disrupted and replaced to introduce unfamiliarity. Rearrangements and

cleaning occurred between trials. Takeover terraria always housed at least two tadpole canisters,

one upright and filled with water and the second empty and sideways, to remove bias for water

availability and shelter, respectively. At the beginning of each trial, frogs were provided with a

dish of springtails available *ad libitum* for the length of the trial and were provided with fruit

flies on the same schedule as their home tanks.

While baseline hormone collection of the adult occurred, we placed a single, unrelated

fertilized late-stage egg to hatchling (Gosner stage 19-22) on a leaf on the terrarium floor. The

leaf was positioned below a Wyze Cam v3 to record behavior using previously described

methods (Goolsby et al., 2023). Frogs (n = 12 females, 12 males) were introduced to the

terrarium and recorded for a maximum of seven days. Infanticidal behavior was characterized as

either consumption of the fertilized egg or hatchling (**Supplementary Video 1**) or as repetitive

physical disturbances to the egg, where the snout repeatedly dug at the jelly-like casing in what

we interpreted as an attempt at egg cannibalism impeded by the encasement (**Supplementary**

Video 2).

2.4 Steroid hormone collection and processing

We collected hormones immediately before displacing frogs to the new tank ("baseline"),

24 ± 3 hours later ("move"), and following infanticide or tadpole transport, or after seven days if

no such behavior occurred ("final"), with hormone collections occurring at approximately the

same time of day or immediately following infanticide (**Fig. 2A-B**). Three subjects demonstrated

infanticide on the day after displacement, so hormone measurements counted for both "move"

and "final" sampling events. Baseline collection always occurred between 11:30 AM and 2:30

PM to avoid confounding effects of circadian physiology on hormone concentrations. We

ultimately analyzed data from 59 samples across 22 individuals.

As shown in previous studies with poison frogs and other amphibians (Gabor et al., 2013;

Baugh et al., 2018, Baugh and Gray-Gailliard, 2021; Rodríguez et al., 2022; Love et al., 2023),

water-borne testosterone and corticosterone were collected as a non-invasive measurement that

reflects circulating levels of both hormones. Frogs were individually moved to a petri dish

containing 40 mL of distilled water treated with reverse osmosis conditioner to prevent osmotic

stress (Josh's Frogs RO R/x, Osowo, MI, USA). After 60 minutes, we measured body length

(snout-vent length) using a digital caliper (Shahe Measuring Tools, Amazon) and body mass

using a Maxus precision pocket scale (sensitivity 200 x 0.01 g, Amazon). We also measured

mass at each hormone sampling event, then averaged to account for variations related to body

fluctuations.

We pre-extracted steroid hormones using Sep-Pak C18 cartridges (Waters, Milford, MA,

USA). The cartridges were conditioned using 2 mL of 100% ethanol, followed by 2 mL of Milli-

Q water treated with reverse osmosis conditioner. Water samples were pushed through the

column at a rate of approx. 10mL/minute, and then eluted with 4 mL of 100% ethanol. Ethanolic

extracts were stored in glass vials at 4 □ until nitrogen evaporation. One day prior to hormone

quantification, the 4 mL of eluted hormone samples were divided into 2 mL new glass tubes for

separate corticosterone and testosterone analyses from the same frog. The 2 mL tubes were

placed in a 37 □ water bath and evaporated with gentle flow N_2 gas. After evaporation, hormone

samples were resuspended in 250 μl of assay buffer specific to the corresponding ENZO kit and

stored at 4 □ overnight until analysis the following day.

2.5 Steroid hormone quantification via enzyme-linked immunosorbent assays (ELISA)

After overnight incubation at 4 □, samples were warmed to room temperature, and

corticosterone and testosterone concentrations were determined using commercially available

ELISA kits (ENZO Life Sciences, Farmingdale, NY; Corticosterone: cat. no. ADI-900-097,

antibody: donkey anti-sheep IgG, sensitivity: 27.0 pg/mL; Testosterone: cat. no. ADI-900-065,

antibody: goat anti-mouse IgG, sensitivity: 5.67 pg/mL) according to the manufacturer's

instructions. We chose to analyze corticosterone rather than cortisol concentrations because

corticosterone is suspected to be the main adrenocorticotropic hormone (ACTH)-responsive

glucocorticoid for *R. imitator* (Cockrem, 2013; Westrick et al., 2023). Samples were vortexed

and 100 µL of the samples were added in duplicate in individual wells of microtiter plates.

Following manufacturer instructions, plates were read at 405 nm, with correction between 570

and 590 nm, using a microplate reader (Synergy H1, BioTek Instruments, Winooski, VT, USA).

Hormone concentrations were calculated using a four-parameter logistic curve in the software

Gen5 (version 3.05, BioTek Instruments, Winooski, VT, USA). Samples with duplicate

measurements that yielded a coefficient of variation (CV) > 30% were excluded from later

analyses (corticosterone: 7 of 59 samples, 11.9%; testosterone: 2 of 59 samples, 3.4%). Before

removing these duplicates, intra-assay CV values for corticosterone and testosterone were 14.0%

and 10.0%. After removal, intra-assay CV values were 8.4% and 9.4%. Inter-assay CV values

were 11.8% and 13.7%.

2.6 Statistics

We cleaned and analyzed all data using RStudio (R version 4.2.3, R Core Team, Boston,

MA). We implemented all statistical analyses using a generalized linear mixed model (GLMM)

framework. For each analysis, we used the Akaike Information Criterion corrected for small

sample sizes (AICc; Burnham and Anderson, 2004) to choose between alternative model fits. We

evaluated standard model diagnostics and tested for outliers and quantile deviations using the

'*DHARMa'* package (version 0.4.6, Hartig, 2022). All models in this manuscript passed all

diagnostic tests. To evaluate the significance of main effects, we performed a Type III analysis

of variance with Satterthwaite's method. We evaluated pairwise comparisons and accounted for

multiple comparisons using the Tukey method in the *'emmeans'* package (version 1.9.0, Lenth,

2023). We estimated effect sizes in multi-termed modeling using the '*effectsize'* package

(version 0.8.6, Ben-Shachar et al., 2020) and with pairwise effect size comparisons using

'*emmeans'*. Figures were created using the package '*ggplot2*' (version 3.4.4, Wickham, 2016)

and assembled in Adobe Illustrator (Adobe Illustrator 2023).

2.6.1 Offspring manipulations

We analyzed the number of eggs deposited for each tadpole among the four experimental conditions (**Fig. 1A**). To test whether egg deposition changes due to cross-fostering and intra-tank displacement, we used a GLMM with a zero-inflation component implemented using the '*glmmTMB*' package (version 1.1.8, Brooks et al., 2017). We chose a Poisson distribution based on best fit, with eggs fed as a dependent variable, experiment type and number of siblings within a terrarium as independent variables, and focal terrarium as a random effect.

2.6.2 Takeover trials

To determine how testosterone, corticosterone, and testosterone:corticosterone ratios change relative to infanticidal behavior, we implemented a GLMM using the '*lmerTest*' package (3.1-3, Kuznetsova et al., 2017). We set the outcome ("infanticide" vs "non-infanticide"), sampling event ("baseline" vs. "move" vs. "final"), sex, and their interactions as fixed effects and individual frog identity, average individual mass, volume of collection water, and body length (snout-vent length) as random effects. We selected this method rather than linearly normalizing by mass, body length, or water collection volume following correlation analyses (**Figs. S1-2**). We estimated average individual mass as the average body mass across sampling events as this was necessary to account for intra-individual differences in feeding. Since mass was not collected for one individual frog, we used mean imputation to estimate the single missing value as the averaged individual mass across the entire study. To improve interpretation of regression coefficients, we removed insignificant interaction terms using stepwise backward regression.

We ran separate models for testosterone and corticosterone. We were additionally

interested in ratios of testosterone to corticosterone, results for which are available in the

supplement (**Figs. S3-4**; **Tables S1-3**). We selected appropriate data transformations (natural log,

square root, inverse, no transformation) for modeling each dependent variable using AICc. Using

AICc and diagnostic criteria, we selected a log transformation for corticosterone and an inverse

transformation for testosterone. Ratios were modeled using the untransformed testosterone and

corticosterone values to compute an initial ratio value, which was then log-transformed. Because

corticosterone, testosterone, and ratio values of samples exceeding the CV threshold of 30% are

deemed unreliable, we treated these values as missing data. All of our final models passed all

diagnostic tests.

3. Results

3.1 Offspring location as an indirect recognition cue

To identify which offspring features parents use to recognize and care for their own

young, we either cross-fostered or displaced tadpoles within their parents' terraria and observed

egg deposition and tadpole mortality for 14 days. We hypothesized that in addition to offspring

manipulations, the number of siblings present may also influence how many eggs can be

provisioned. We manipulated offspring identity (same-morph cross-foster, different-morph

cross-foster) and offspring location (intra-tank displacement, **Fig. 1A**). We found that offspring

manipulation type ($\chi^2 = 17.74$, p < 0.0005; **Table S4**) was an overall better predictor of eggs

provisioned than the number of siblings present in a tank at a time (CI: -0.51 – 0.22, $\chi^2 = 0.62$, p

= 0.43; **Table S4**).

Figure 1. Disrupting offspring location impedes egg provisioning and promotes infanticide. (A) Control tadpoles were the offspring of the parents in that terrarium and were not moved from their original locations. Cross-fostered tadpoles were exchanged between terraria, such that they were unrelated to the parents in the new terrarium. They were specifically moved to the same location previously occupied by the offspring they replaced. The different-morph cross-foster manipulation followed the same procedure as same-morph cross-fosters, with the additional condition that tadpoles were exchanged between terraria housing different morphs (striped, banded), discerned by parents' coloration patterns. Finally, intra-tank displacement tadpoles were related to the parents in the terrarium but were displaced by a minimum of five centimeters to a different location in the terrarium. **(B)** Tadpoles displaced within their parents' tanks received significantly less feeding in the form of trophic eggs from parents (Tukey adjustment for a family of 4 estimates: $p < 0.0008$). Significance: $0 < *** < 0.001 < ** < 0.01 < * < 0.05 <$ ns. "ns" = not significant. **(C)** Only tadpoles displaced within their parents' tanks (n = 4 out of 18) were subjected to infanticide with no deposition of a second tadpole.

We next asked whether offspring displacement influenced parental provisioning of young and found a strong effect on the number of eggs provisioned (CI: -2.43 – -0.37, z = -2.66, p < 0.01, **Table 1**; model diagnostics may be found in **Fig. S5** with details for analysis of deviance in **Table S4.** Specifically, over 14 days, displaced tadpoles received significantly fewer trophic eggs compared to positive controls (**Fig. 1B**, $p_{adj} = 0.0008$, $\square = 1.81$). Across our experiment, control tadpoles were fed the most, at an average of 1.77 eggs every two weeks, whereas same-

morph and different-morph cross-fosters were fed less than positive controls, being fed 1.13 and

0.58 eggs on average, respectively. Displaced tadpoles were the least cared for, being fed an

average of 0.27 eggs within two weeks. Tadpoles cross-fostered within and outside of their

morph did not experience a significant reduction in feeding compared to positive controls (**Fig.**

1B, p_{adj} = 0.16, 0.55; □ = -0.409, -0.910; **Table S5**). Tadpoles displaced within their parents'

terraria were subjected to a variety of infant-directed behaviors, including "care" as exemplified

by egg feeding, "non-infanticide" as exemplified by the absence of observed feeding, and

"infanticide" as shown by physical injury to the tadpole without deposition of a second tadpole,

which was only observed in the intra-tank displacement manipulation (**Fig. 1C**).

Table 1. Final model fit of eggs fed with offspring manipulations. SD = Standard Deviation. SE = Standard Error, z

= standardized Z value, Pr() = P value.

Random Effects	Variance	SD		
Tank	0.23	0.48		
Fixed Effects	Estimate	SE	z	Pr (>|z|)
Intercept	0.40	0.64	0.63	0.53
Different Morph	-0.50	0.50	-1.01	0.31
Control	0.41	0.31	1.32	0.19
Displacement	-1.40	0.53	-2.66	**7.80E-03**
Siblings	-0.15	0.19	-0.78	0.43

3.2 Infanticide in new territory

After concluding that offspring location is an important factor in parental decisions of

care versus infanticide, we performed a "takeover" behavioral assay to identify hormonal

correlates of infanticide. We analyzed adult corticosterone and testosterone concentrations with

respect to outcome ("infanticide" versus "non-infanticide"), sampling event ("baseline", "move",

and "final"), and interactions between outcome and event themselves (**Fig. 2A-B**).

Figure 2. Takeover setup and behavioral outcomes. **(A)** Individual adult frogs were displaced from their home

terraria to unfamiliar terraria along with a single, unrelated, fertilized egg or hatchling placed on a leaf. Trials lasted

for a maximum of seven days. Hormones were collected via water-borne hormone sampling on the day of the move

("baseline"), ~24 hours afterward ("move"), and upon observation of infanticidal behavior, or on Day 7 if no such

behavior occurred ("final"). **(B)** Behavioral outcomes toward eggs or recently hatched tadpoles as observed on

motion-trigger Wyze v3 cameras included "infanticide" in the form of cannibalism or physical disturbance to the

egg, "transport" when tadpole transport was observed, or "non-infanticide" where these behaviors were not

observed. **(C)** Females exhibited all outcomes (non-infanticide: n = 5, infanticide: n = 6, transport: n = 1), unlike

males (non-infanticide: n = 8, infanticide: n = 4).

Of the twelve trials performed with females, six of the twelve females fell within the

"infanticide" outcome, while the other five were categorized as ignoring the conspecific young

("non-infanticide"). One individual demonstrated parental care, performing tadpole transport

(**Supplementary Video 3**). As we only observed transport by one individual, we did not have

enough statistical power to analyze this result distinct from other non-infanticidal females, and

therefore we excluded it from downstream analyses. Of the twelve trials performed with males,

eight showed no infanticidal behavior toward conspecific young while four performed

infanticide (**Fig. 2C**). Latency to infanticide decisions were variable, with the behavior occurring

on a range of days, from the day after movement to the new terrarium to the full seven days (**Fig.**

S6). In addition to interactions with offspring, we measured frog mass at each sampling event.

The median frog mass decreased from baseline to final sampling (**Fig. S7**).

3.3 Infanticide and corticosterone

Log corticosterone (corticosterone hereafter) concentrations did not vary significantly

between infanticidal versus non-infanticidal adults (**Table 2**, $\eta_p^2 = 0.01$, t = -0.59, p = 0.56; **Fig.**

4A, Table S6) nor did they vary significantly between males and females (**Table 2,** $\eta_p^2 = 0.17$, t

= 2.00, p = 0.06; **Fig. 4A, Table S6**). On average, corticosterone concentrations increased

approximately 54 pg/mL during takeover of novel territory, although this increase was not

significant ($\eta_p^2 = 0.12$, t = 1.69, p = 0.10, **Table 2**). Model diagnostics may be found in **Fig. S8**

with analysis of deviance details in **Table S6**. No pairwise comparisons between combinations of

behavioral outcomes, sampling events, and sex were significant (**Table S7**).

Figure 3. Hormone concentrations by event and behavioral outcome. (A) Corticosterone and **(B)** testosterone

concentrations by sampling events and behavioral outcomes. The "X" in each box represents the average. The

horizontal lines in each box plot represent the medians of each group. All results are shown in raw units.

376   **Table 2.** Final corticosterone model fit of log(corticosterone) by sampling event, behavioral outcome, and sex. Frog

377   identity, frog length, and average mass are random effects. SD = Standard Deviation. SE = Standard Error, t = t-

378 statistic, Pr (>|t|) = P value.

Random Effects	Variance	SD			
Body length	0.03	0.18			
Average mass across trial	0.05	0.22			
Residual	0.09	0.30			
Fixed Effects	Estimate	*SE*	df	*t*	Pr (>|t|)
Intercept	6.25	0.14	30.92	44.04	**<2.00E-16**
Outcome	-0.08	0.14	23.15	-0.59	0.56
Move	0.19	0.11	27.63	1.69	0.10
Final	-0.04	0.11	36.94	-0.32	0.76
Sex	0.30	0.15	20.18	2.00	0.06

3.4 Infanticide and testosterone

Across all samples, the inverse transformation of testosterone (testosterone hereafter)

concentration varied significantly with sampling events (η_p^2 = 0.31, p=0.03, F=4.12, **Fig. 3B,**

Table S8). Testosterone concentrations also changed between behavioral outcomes (η^2 = 0.28,

p=0.02, F = 6.40, **Fig. 3B, Table S8**), but were not sex-specific (η_p^2 = 0.04, p=0.40, F=0.75, **Fig.**

3B, Table S8). Infanticidal individuals, across their trial, averaged about two-thirds of the

testosterone concentrations (67.94 pg/mL) of non-infanticidal individuals (97.81 pg/mL). In

other words, testosterone decreased, on average, during potentially stressful events (baseline

compared to move), and during infanticidal behaviors. Most notably, testosterone concentrations

significantly decreased 24 hours after takeover onset (t = 3.30, p < 0.005), but the extent of this

decrease was significantly dependent on whether the individual eventually performed infanticide

(η_p^2 = 0.35, t = -2.60, p = 0.02, **Table 3**). Model diagnostics may be found in **Fig. S9** with full

analysis of deviance details in **Table S8.**

Table 3. Final testosterone model fit of 1/(testosterone) by sampling event, behavioral outcome, and sex. Frog body length, identity, and water collection volume are random effects. SD = Standard Deviation. SE = Standard Error, t = t-statistic, Pr (>|t|) = P value.

Random Effects	Variance	SD			
Body length	8.05E-06	2.83E-03			
Frog Identity	2.24E-05	4.74E-03			
Water volume	2.57E-05	5.07E-03			
Residual	8.38E-06	2.89E-03			
Fixed Effects	Estimate	SE	df	t	**Pr (>|t|)**
Intercept	1.65E-02	2.72E-03	21.57	6.08	**4.42E-06**
Outcome	-4.49E-03	2.82E-03	24.95	-1.59	0.12
Move	5.97E-03	1.81E-03	17.23	3.30	**4.18E-03**
Final	3.37E-03	1.91E-03	15.43	1.77	0.10
Sex	-2.20E-03	2.54E-03	18.16	-0.87	0.40
Move : Outcome	-5.90E-03	2.27E-03	12.95	-2.61	**0.02**
Final : Outcome	-2.10E-04	2.49E-03	17.26	0.08	0.94

In pairwise comparisons, we examined testosterone concentrations in relation to

interactions between sampling events and outcomes. We found that baseline testosterone

concentrations in adults who did not perform infanticide were significantly greater than

testosterone concentrations in infanticidal individuals moving into a new territory (Cohen's *d* = -

3.62, t ratio = 3.38, p = 0.03). Finally, infanticidal and non-infanticidal individuals have different

testosterone concentrations when moved into a new territory (average of 50.62 pg/mL and 90.12

pg/mL, respectively; Cohen's d = 3.59, t ratio = 3.38, p = 0.03). Within infanticidal individuals,

we found a moderately positive (r = 0.48), albeit statistically insignificant, correlation between

testosterone concentrations 24 hours into the territory takeover assay and latency to infanticide (p

= 0.19; **Fig. S10**).

4. Discussion

In this investigation, we first delineated tadpole-based features that influence parental

decisions to perform care or infanticide, identifying location as a salient indirect kin recognition

cue. Then, we documented adult behavior toward eggs and hatchlings in unfamiliar territories,

recording infanticide for the first time in *Ranitomeya* frogs. By analyzing concentrations of

water-borne corticosterone and testosterone, we demonstrated that lower testosterone

concentrations precede infanticide in this species.

4.1 Parental decision-making in familiar and unfamiliar territories

We observed that displaced tadpoles experienced significant decreases in trophic egg

deposition compared to control tadpoles, unlike tadpoles cross-fostered within or between

morphs **(Fig. 1B)**. Like other poison frogs (Stynoski, 2009; Ringler et al., 2016; Ringler et al.,

2017), *R. imitator* parents likely discriminate between young based on location. Offspring

location therefore appears to be a critical cue driving parental care across poison frogs despite

diversity in care systems and ecological constraints. Critical decisions between care or

cannibalism depending on internal physiological states are especially abundant in anamniotes

and squamates (Ray and Maruska, 2023). For example, spawning appears to impede cannibalism

in parenting African *Neolamprologus caudopunctatus* cichlids, resulting in the care of foreign

young (Cunha-Saraiva et al., 2018). Similarly, brooding children's pythons (*Antaresia children*)

will provide care when eggs are cross-fostered or even replaced with stones (Brashears and

DeNardo, 2012).

A proportion of displaced tadpoles were found dead shortly after being moved to a new

location in their home terrarium, which we interpreted to be caused by infanticide (**Fig. 1C**).

These offspring did not encounter any other tadpoles, making intra-sibling aggression unlikely,

despite being well-documented in *Ranitomeya* (Brown et al., 2009; Schulte and Mayer, 2017;

McKinney et al., 2022). Offspring cannibalism as a behavioral consequence of intraspecific

competition has been previously suggested in other poison frogs (Summers, 1989), where adults

might commit infanticide of unrecognized offspring to prevent misdirected care, to promote

survival of their own offspring by parasitizing occupied deposition sites, or to prevent offspring

cannibalism by unrelated tadpoles. Other species of *Ranitomeya* have been found to parasitize

the care of other parents, indicating brood parasitism as a weapon between conspecific

competitors (Poelman and Dicke, 2007; Brown et al., 2009, respectively). To understand why

tadpoles were killed rather than eaten by adults, we theorize that dead tadpoles may yield a

precious protein source for cannibalistic *R. imitator* tadpoles that adults may deposit. Protein can

otherwise be difficult to obtain in low-resource environments and may minimize future costs of

females having to provision unfertilized egg meals to nutritionally needy tadpoles (Yoshioka et

al., 2016). Thus, we hypothesize that infanticide by *R. imitator* adults evolved for parents to

maximize their individual reproductive success and opportunities while minimizing those of

potential competitors, following evaluation of their environments.

Following the identification of offspring location as an indirect offspring recognition cue,

we analyzed the behavioral responses of reproductive adults to unrelated, conspecific young in a

novel terrarium, where adults presumably would not mistake young as their own. We found that

both males and females cannibalized the stimulus, marking the first documented report of adults

cannibalizing young in the *Ranitomeya* genus and supporting previous reports of non-filial

cannibalism in other poison frogs (Townsend et al., 1984; Summers, 1989; Ringler et al., 2017;

Spring et al., 2019; Dugas et al., 2023). We considered multiple adaptive rationales for non-filial

cannibalism in *R. imitator*. First, cannibalism could be a behavioral decision relevant to a trade-

off between acquiring caloric content (which risks eating offspring) and providing care (which

risks caring for unrelated young), an idea which has been tested extensively in fish (Bose, 2022).

Fertilized conspecific eggs constitute part of other frog diets (Beard, 2007). Furthermore, feeding

behavior appears mechanistically linked to parental status across taxa including mammals, birds,

and fish (O'Rourke and Renn, 2015; Fischer and O'Connell, 2017). Thus, we reasoned that some

frogs might be more likely to cannibalize the stimulus egg or hatchling based on differences in

how recently they parented, although all subjects produced a tadpole within the previous 31 days.

To account for various degrees of hunger and potential feeding-related effects of parental status,

frogs were supplied with food *ad libitum*, so it is unlikely that the frogs in the present study

cannibalize to compensate for nutritional deficits. Therefore, it appears more likely that

cannibalism occurred as a response to stress or intraspecific competition rather than as a means

of acquiring food. However, it may be worthwhile for future investigations to clarify the

importance of environmental factors not directly addressed here, including mate access, recent

reproductive activity, and territoriality.

 4.2 Hormonal correlates of infanticide

To clarify whether steroid hormones correlate with infanticide, we analyzed infanticidal behavior and sampling events in relation to corticosterone and testosterone concentrations. We expected to find that adults which performed infanticide would exhibit greater corticosterone concentrations upon displacement to new territory. Instead, corticosterone concentrations were not significantly associated with infanticide at any sampling event (**Fig. 3A**). We were also surprised to find no significant differences in corticosterone between sampling events. However, it is possible that sampling at a time point sooner after displacement would have better reflected the physiological changes associated with an acutely stressful environmental change. Interestingly, glucocorticoids have been proposed to promote both avoidant and approaching behaviors (Terburg et al., 2009), which may relate to decisions to cannibalize and ignore unrelated offspring, respectively. Based on our observations, we conclude that behavioral reactions to unrelated young in this species likely depend more on non-corticosteroid pathways, potentially involving other steroids or neuromodulators.

Circulating androgens are classically associated with territorial aggression in mammals, birds, and recently, other poison frogs (Wingfield et al., 1990; Duque-Wilckens et al., 2019; Rodríguez et al., 2022). Therefore, we expected to find that frogs that performed infanticide upon displacement to a new territory would exhibit greater levels of testosterone. Contrary to what we expected, circulating testosterone appears to be negatively associated with infanticide in this species (**Fig. 3B**). Our finding aligns with investigations in some fish, where plasma 11-ketotestosterone concentrations were lower in cannibals but comparable to those typical of parents (Takegaki et al., 2023). Interestingly, low testosterone is also associated with parental care in male poison frogs (Townsend and Moger, 1987); therefore, it seems plausible that

decreased androgens can be associated with both care and infanticide in amphibians. Potential

mechanisms for such a relationship may include the local aromatization of testosterone to

estradiol, which has been linked to aggression in several mammals, fish, and birds (Trainor et al.,

2006; Huffman et al., 2013) and is a conserved process in amphibians (Coumailleau et al., 2015).

The co-option of estrogens rather than androgens to facilitate aggression can potentially avoid

the costs of elevated testosterone, which can be especially detrimental in parents (Wingfield et

al., 2001).

Finally, while our work necessitates hormone collections *after* a behavioral outcome, this

does not suggest that hormone collections cause behavioral outputs. Rather, it is equally possible

that behavioral outputs may drive variation in hormone levels, which has been richly

documented across contexts where animals encounter opportunities such as social ascent,

resources, or sexual opportunities (Nelson, 2009; Maruska and Fernald, 2010). We hope that

future work can functionally delineate the relationship between androgens and infanticidal

behavior in animals broadly.

5. Conclusions

In this study, we aimed to clarify environmental and hormonal cues for infanticide in

mimic poison frogs. Our results suggest that, consistent with other poison frogs, care and

infanticide in this species are antagonistically linked on the basis of a simple external cue:

offspring location. We showed that mimic poison frogs perform infanticide in both familiar and

unfamiliar territories, wherein they targeted related and unrelated young, respectively. Based on

the ecological history of *R. imitator*, we posit that infanticide in this species serves to prevent

misdirected care and eliminate intraspecific competition. Although infanticide has been observed

in other poison frogs, to our knowledge, this is the first such report in a monogamous or

biparental amphibian, indicating that infanticide by both sexes can occur regardless of mating or

parenting systems in this clade. Moreover, our hormonal analyses indicate that low

concentrations of circulating androgens can precede infanticide following social and

environmental perturbations, which contributes a unique perspective to the broader aggression

literature. Overall, these findings offer fresh insights into how adults adjust their behavior and

physiology to make life-or-death decisions toward offspring in the face of social instability. In

the future, a concerted analysis of endocrine and neural activities compared between individuals

performing infanticide, neglect, and care is a promising next direction to uncover evolutionary

innovations in physiology underlying offspring-directed behaviors.

Funding

This work was supported by a McKnight Pecot Fellowship to ARL and grants from the National

Institutes of Health (DP2HD102042) and the New York Stem Cell Foundation to LAO. ARL

was additionally supported by the Stanford Bio-X Undergraduate Research Fellowship and

grants from the Office of the Vice Provost for Undergraduate Education. BCG was supported by

a HHMI Gilliam Fellowship (GT15685) and a National Institutes of Health Cellular Molecular

Biology Training Grant (T32GM007276). BHJ was supported by funding from the New York

Stem Cell Foundation. LAO is a New York Stem Cell Foundation–Robertson Investigator.

CRediT authorship contribution statement

Amaris R. Lewis: Conceptualization, Methodology, Formal Analysis, Investigation, Data

Curation, Writing - Original Draft, Writing - Review and Editing, Visualization, Project

Administration, Funding Acquisition. **Billie C. Goolsby:** Conceptualization, Methodology,

Formal Analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review and

Editing, Visualization, Project Administration. **Bryan H. Juarez:** Methodology, Formal

Analysis, Resources, Writing - Original Draft, Writing - Review and Editing, Visualization,

Supervision. **Madison P. Lacey**: Investigation, Resources, Writing - Review and Editing.

Lauren A. O'Connell: Resources, Writing- Review & Editing, Funding Acquisition,

Supervision**.**

Acknowledgments

This research was conducted at Stanford University, which is located on the ancestral and

unceded land of the Muwekma Ohlone tribe. We thank Dr. Camilo Rodríguez Lopez (CRL) and

Dr. Ricardo Cossio for feedback on this manuscript. We are additionally grateful to CRL for

guidance on steroid hormone collection and processing. We also thank the members of the

Laboratory for Organismal Biology for providing routine frog care and continuous support.

Finally, we kindly thank the two reviewers for their thoughtful comments which improved the

quality of this manuscript.

References

Alonso-Alvarez, C., & Velando, A. (2012). Benefits and costs of parental care. *The evolution of parental care*, 40-61. https://doi.org/10.1093/acprof:oso/9780199692576.003.0003

Baugh, A. T., Bastien, B., Still, M. B., & Stowell, N. (2018). Validation of water-borne steroid hormones in a tropical frog (Physalaemus pustulosus). *General and Comparative Endocrinology*, *261*, 67-80. https://doi.org/10.1016/j.ygcen.2018.01.025.

Baugh, A. T., & Gray-Gaillard, S. L. (2021). Excreted testosterone and male sexual proceptivity: a hormone validation and proof-of-concept experiment in túngara frogs. *General and Comparative Endocrinology*, *300*, 113638. https://doi.org/10.1016/j.ygcen.2020.113638

Beard, K. (2007). Diet of the invasive frog, *Eleutherodactylus coqui*, in Hawaii. *Copeia, 2007*(2), 281-291. https://doi.org/10.1643/0045-8511(2007)7[281:DOTIFE]2.0.CO;2

Ben-Shachar, M. S., Lüdecke, D., & Makowski, D. (2020). effectsize: Estimation of effect size indices and standardized parameters. *Journal of Open Source Software*, 5(56), 2815. https://doi.org/10.21105/joss.02815

Bentz, A. B., Philippi, K. J., & Rosvall, K. A. (2019). Evaluating seasonal patterns of female aggression: Case study in a cavity☐nesting bird with intense female–female competition. *Ethology*, *125*(8), 555–564. https://doi.org/10.1111/eth.12881

Bose, A. P. (2022). Parent–offspring cannibalism throughout the animal kingdom: a review of adaptive hypotheses. *Biological Reviews*, *97*(5), 1868-1885. https://doi.org/10.1111/brv.12868

Brashears, J., & DeNardo, D. F. (2012). Do Brooding Pythons Recognize their Clutches? Investigating External Cues for Offspring Recognition in the Children's Python,

Antaresia childreni . *Ethology*, *118*(8), 793–798. https://doi.org/10.1111/j.1439-0310.2012.02070.x

Brooks, M. E., Kristensen, K., Van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378. https://doi.org/10.32614/RJ-2017-066

Brown, J., Twomey, E., Morales, V., & Summers, K. (2008). Phytotelm size in relation to parental care and mating strategies in two species of Peruvian poison frogs. *Behaviour*, *145*(9), 1139-1165. https://doi.org/10.1163/156853908785387647

Brown, J. L., Morales, V., & Summers, K. (2009). Tactical reproductive parasitism via larval cannibalism in Peruvian poison frogs. *Biology Letters*, *5*(2), 148-151. https://doi.org/10.1098/rsbl.2008.0591

Burnham, K. P., & Anderson, D. R. (Eds.). (2004). *Model Selection and Multimodel Inference*. Springer New York. https://doi.org/10.1007/b97636

Clutton-Brock, T. H. (1991). *The evolution of parental care* (Vol. 64). Princeton University Press. 10.1093/acprof:oso/9780199692576.001.0001

Cockrem, J. F. (2013). Individual variation in glucocorticoid stress responses in animals. *General and Comparative Endocrinology*, 181, 45-58. https://doi.org/10.1016/j.ygcen.2012.11.025

Coumailleau, P., Pellegrini, E., Adrio, F., Diotel, N., Cano-Nicolau, J., Nasri, A., ... & Kah, O. (2015). Aromatase, estrogen receptors and brain development in fish and amphibians. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1849(2), 152-162. https://doi.org/10.1016/j.bbagrm.2014.07.002

Cunha-Saraiva, F., Balshine, S., Wagner, R. H., & Schaedelin, F. C. (2018). From cannibal to

caregiver: tracking the transition in a cichlid fish. *Animal Behaviour*, 139, 9–17.

https://doi.org/10.1016/j.anbehav.2018.03.003

Dugas, M. B., Brooks, O. L., Saporito, R. A., & Cossio, R. (2023). Adult poison frogs can

capture and consume aquatic tadpoles. *Evolutionary Ecology*, 1-7.

https://doi.org/10.1007/s10682-023-10257-1

Duque-Wilckens, N., Trainor, B. C., & Marler, C. A. (2019). Aggression and territoriality.

Encyclopedia of animal behavior, 2, 539-546.

Ebensperger, L. A., & Blumstein, D. T. (2007). Nonparental infanticide. *Rodent societies: an*

ecological and evolutionary perspective, 267-279.

Fischer, E. K., & O'Connell, L. A. (2017). Modification of feeding circuits in the evolution of

social behavior. *Journal of Experimental Biology*, *220*(1), 92-102.

https://doi.org/10.1242/jeb.143859

Gabor, C. R., Bosch, J., Fries, J. N., & Davis, D. R. (2013). A non-invasive water-borne

hormone assay for amphibians. *Amphibia-Reptilia*, *34*(2), 151-162.

https://doi.org/10.1163/15685381-00002877

Goolsby, B. C., Fischer, M. T., Pareja-Mejia, D., Lewis, A. R., Raboisson, G., & O'Connell, L.

A. (2023). Home security cameras as a tool for behavior observations and science equity.

bioRxiv. https://doi.org/10.1101/2023.04.17.537238

Gosner, K. L. (1960). A Simplified Table for Staging Anuran Embryos and Larvae with Notes

on Identification. *Herpetologica*, *16*(3), 183–190.

Haller, J. (2014). The glucocorticoid/aggression relationship in animals and humans: an analysis

sensitive to behavioral characteristics, glucocorticoid secretion patterns, and neural

mechanisms. *Neuroscience of aggression*, 73-109.

https://doi.org/10.1007/7854_2014_284

Hartig, F. (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)

Regression Models. R package version 0.4.6.

Hirschenhauser, K., & Oliveira, R. F. (2006). Social modulation of androgens in male

vertebrates: meta-analyses of the challenge hypothesis. *Animal Behaviour*, *71*(2), 265-

277. https://doi.org/10.1016/j.anbehav.2005.04.014

Hrdy, S. B. (1979). Infanticide among animals: a review, classification, and examination of the

implications for the reproductive strategies of females. *Ethology and Sociobiology*, *1*(1),

13-40. https://doi.org/10.1016/0162-3095(79)90004-9

Huang, W. S., & Pike, D. A. (2011). Does maternal care evolve through egg recognition or

directed territoriality?. *Journal of Evolutionary Biology*, 24(9), 1984-1991. https://

10.1111/j.1420-9101.2011.02332.x

Huffman, L. S., O'Connell, L. A., & Hofmann, H. A. (2013). Aromatase regulates aggression in

the African cichlid fish Astatotilapia burtoni. *Physiology & Behavior*, 112–113, 77–83.

https://doi.org/10.1016/j.physbeh.2013.02.004

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13).

https://doi.org/10.18637/jss.v082.i13

Lenth, R (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.9.0. https://doi.org/10.1080/00031305.1980.10483031

Love, N., Preininger, D., & Fuxjager, M. J. (2023). Social regulation of androgenic hormones and gestural display behavior in a tropical frog. Hormones and Behavior, 155, 105425. https://doi.org/10.1016/j.yhbeh.2023.105425

Lukas, D., & Huchard, E. (2014). The evolution of infanticide by males in mammalian societies. *Science*, 346(6211), 841-844. https://doi.org/10.1126/science.1257226

Lukas, D., & Huchard, E. (2019). The evolution of infanticide by females in mammals. *Philosophical Transactions of the Royal Society B*, 374(1780), 20180075. https://doi.org/10.1098/rstb.2018.0075

Maruska, K. P., & Fernald, R. D. (2010). Behavioral and physiological plasticity: rapid changes during social ascent in an African cichlid fish. *Hormones and behavior*, *58*(2), 230–240. https://doi.org/10.1016/j.yhbeh.2010.03.011

McKinney, J. E., Ludington, S. C., Butler, J. M., & O'Connell, L. A. (2022). Proopiomelanocortin (POMC) is a negative regulator of tadpole aggression through opioid receptor signaling. *bioRxiv*, 2022-11. https://doi.org/10.1101/2022.11.28.518266

Neff, B. D., & Sherman, P. W. (2005). In vitro fertilization reveals offspring recognition via self□referencing in a fish with paternal care and cuckoldry. *Ethology*, 111(4), 425-438. https://doi.org/10.1111/j.1439-0310.2005.01075.x

Nelson, R. J. (2009). Hormones and behavior: basic concepts. *Encyclopedia of animal behavior*, 97-105.

O'Rourke, C. F., & Renn, S. C. (2015). Integrating adaptive trade-offs between parental care and feeding regulation. *Current opinion in behavioral sciences*, 6, 160-167. https://doi.org/10.1016/j.cobeha.2015.11.010

Poelman, E. H., & Dicke, M. (2007). Offering offspring as food to cannibals: oviposition

strategies of Amazonian poison frogs (Dendrobates ventrimaculatus). *Evolutionary*

Ecology, *21*(2), 215–227. https://doi.org/10.1007/s10682-006-9000-8

Ray, E. J., & Maruska, K. P. (2023). Sensory Mechanisms of Parent-Offspring Recognition in

Fishes, Amphibians, and Reptiles. *Integrative And Comparative Biology*, 63(6), 1168–

1181. https://doi.org/10.1093/icb/icad104

Ringler, E., Pašukonis, A., Ringler, M., & Huber, L. (2016). Sex-specific offspring

discrimination reflects respective risks and costs of misdirected care in a poison frog.

Animal Behaviour, *114*, 173-179. https://doi.org/10.1016/j.anbehav.2016.02.008

Ringler, E., Barbara Beck, K., Weinlein, S., Huber, L., & Ringler, M. (2017). Adopt, ignore, or

kill? Male poison frogs adjust parental decisions according to their territorial status.

Scientific Reports, *7*(1), 43544. https://doi.org/10.1038/srep43544

Rodríguez, C., Fusani, L., Raboisson, G., Hödl, W., Ringler, E., & Canoine, V. (2022).

Androgen responsiveness to simulated territorial intrusions in Allobates femoralis males:

Evidence supporting the challenge hypothesis in a territorial frog. *General and*

Comparative Endocrinology, *326*, 114046. https://doi.org/10.1016/j.ygcen.2022.114046

Rosvall, K. A., Bentz, A. B., & George, E. M. (2020). How research on female vertebrates

contributes to an expanded challenge hypothesis. *Hormones and Behavior*, 123, 104565.

https://doi.org/10.1016/j.yhbeh.2019.104565

Royle, N. J., Smiseth, P. T., & Kölliker, M. (Eds.). (2012). *The evolution of parental care*.

Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199692576.001.0001

Schulte, L. M., & Mayer, M. (2017). Poison frog tadpoles seek parental transportation to escape

their cannibalistic siblings. *Journal of Zoology*, *303*(2), 83-89.

https://doi.org/10.1111/jzo.12472

Spring, S., Lehner, M., Huber, L., & Ringler, E. (2019). Oviposition and father presence reduce

clutch cannibalism by female poison frogs. *Frontiers in zoology*, *16*, 1-10.

https://doi.org/10.1186/s12983-019-0304-2

Stynoski, J. L. (2009). Discrimination of offspring by indirect recognition in an egg-feeding

dendrobatid frog, *Oophaga pumilio*. *Animal Behaviour*, *78*(6), 1351-1356.

https://doi.org/10.1016/j.anbehav.2009.09.002

Summers, C. H., Watt, M. J., Ling, T. L., Forster, G. L., Carpenter, R. E., Korzan, W. J., Lukkes,

J.L., & Øverli, Ø. (2005). Glucocorticoid interaction with aggression in non-mammalian

vertebrates: reciprocal action. *European Journal of Pharmacology*, *526*(1-3), 21-35.

https://doi.org/10.1016/j.ejphar.2005.09.059

Summers, K. (1989). Sexual selection and intra-female competition in the green poison-dart frog,

Dendrobates auratus. *Animal Behaviour*, *37*, 797–805. https://doi.org/10.1016/0003-

3472(89)90064-X

Summers, K., & Tumulty, J. (2014). Parental Care, Sexual Selection, and Mating Systems in

Neotropical Poison Frogs. In *Sexual Selection* (pp. 289–320). Elsevier.

https://doi.org/10.1016/B978-0-12-416028-6.00011-6

Takegaki, T., Nakatake, Y., Matsumoto, Y., Suga, K., & Amiya, N. (2023). Early filial

cannibalism in fish revisited: endocrinological constraint, costs of parental care, and

mating possibility. *The American Naturalist*, 201(6), 841-850.

https://doi.org/10.1086/724284

Terburg, D., Morgan, B., & van Honk, J. (2009). The testosterone–cortisol ratio: A hormonal

marker for proneness to social aggression. *International Journal of Law and Psychiatry*,

32(4), 216–223. https://doi.org/10.1016/j.ijlp.2009.04.008

Townsend, D. S., & Moger, W. H. (1987). Plasma androgen levels during male parental care in a

tropical frog (Eleutherodactylus). *Hormones and Behavior*, *21*(1), 93-99.

https://doi.org/10.1016/0018-506X(87)90034-1

Townsend, D. S., Stewart, M. M., & Pough, F. H. (1984). Male parental care and its adaptive

significance in a neotropical frog. *Animal Behaviour*, *32*(2), 421-431.

https://doi.org/10.1016/0018-506X(87)90034-1

Trainor, B. C., Kyomen, H. H., & Marler, C. A. (2006). Estrogenic encounters: how interactions

between aromatase and the environment modulate aggression. *Frontiers in*

Neuroendocrinology, *27*(2), 170-179. https://doi.org/10.1016/j.yfrne.2005.11.001.

Vergne, A. L., Aubin, T., Taylor, P., & Mathevon, N. (2011). Acoustic signals of baby black

caimans. *Zoology*, 114(6), 313-320. https://10.1016/j.zool.2011.07.003

Weygoldt, P. (1987). Evolution of parental care in dart poison frogs (Amphibia: Anura:

Dendrobatidae). *Journal of Zoological Systematics and Evolutionary Research*, *25*(1),

51–67. https://doi.org/10.1111/j.1439-0469.1987.tb00913.x

Westrick, S. E., Paitz, R. T., & Fischer, E. K. (2023). Why not both? A case study measuring

cortisol and corticosterone in poison frogs. *bioRxiv*, 2023-06.

https://doi.org/10.1101/2023.06.19.545597

Wickham, H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

ISBN 978-3-319-24277-4. https://doi.org/10.1007/978-3-319-24277-4.

Wingfield, J. C., Hegner, R. E., Dufty Jr, A. M., & Ball, G. F. (1990). The "challenge
hypothesis": theoretical implications for patterns of testosterone secretion, mating
systems, and breeding strategies. *The American Naturalist*, *136*(6), 829-846.
https://doi.org/10.1086/285134
Wingfield, J. C., Lynn, S. E., & Soma, K. K. (2001). Avoiding the 'costs' of testosterone:
ecological bases of hormone-behavior interactions. *Brain Behavior and Evolution*, 57(5),
239-251. https://doi.org/10.1159/000047243
Yoshioka, M., Meeks, C., & Summers, K. (2016). Evidence for begging as an honest signal of
offspring need in the biparental mimic poison frog. *Animal Behaviour*, 113, 1–11.
https://doi.org/10.1016/j.anbehav.2015.12.024