1   **The neonicotinoid acetamiprid is highly toxic to wild non-target insects**
2

3   Jan Erik Sedlmeier[1*], Ingo Grass[3], Prasanth Bendalam[1], Birgit Höglinger[2], Frank Walker[2],
4   Daniel Gerhard[4], Hans-Peter Piepho[5], Carsten A. Brühl[6] and Georg Petschenka[1*]
5
[1]Applied Entomology, University of Hohenheim, Stuttgart, Germany
[2]Analytical Chemistry, University of Hohenheim, Stuttgart, Germany
[3]Ecology of Tropical Agricultural Systems, University of Hohenheim, Stuttgart, Germany
[4]School of Mathematics and Statistics, University of Canterbury, Christchurch, New Zealand
[5]Biostatistics, University of Hohenheim, Stuttgart, Germany
[6]Community Ecology & Ecotoxicology, University of Kaiserslautern-Landau, Landau,
Germany
13
[*]**Corresponding authors:**
Jan Erik Sedlmeier
Department of Applied Entomology, Institute of Phytomedicine, University of Hohenheim,
Otto-Sander-Str. 5, 70599 Stuttgart; jan.sedlmeier@uni-hohenheim.de
Georg Petschenka
Department of Applied Entomology, Institute of Phytomedicine, University of Hohenheim,
Otto-Sander-Str. 5, 70599 Stuttgart; georg.petschenka@uni-hohenheim.de
21

Abstract

Although pesticides are seen as one of the main drivers of insect decline, there are still only
few studies that assess their effects on non-target species under field conditions. Here we
investigated the effects of the neonicotinoid insecticide Mospilan®SG (active ingredient
acetamiprid) on plant bugs (Heteroptera: Miridae), a dominant group of European grassland
insect communities. Abundance of three focal mirid species was reduced by up to 78% two
days after field exposure at concentrations expected at field margins, with mortality varying
considerably among species. Follow-up feeding assays with insecticide-treated host plants in
the greenhouse and controlled dose-response assays in the laboratory confirmed the strong
negative effects on non-target species. Strikingly, the neonicotinoid was nearly 10,000 times
more toxic to plant bugs than to honeybees. In addition, male bugs were 20 times more
sensitive than females in two of the three tested species. Thus, continuous exposure to
neonicotinoids in the field may reduce the plant bug populations and promote a shift towards
more insecticide-tolerant species, altering community composition. We suggest that sex-
specific sensitivity should be considered in risk assessment and conclude that the real risk to
non-target insects is currently greatly underestimated.
38
39

1. Main

We are facing an unprecedented decline in biodiversity, which is particularly evident for insects
in the past years, as many studies have shown worldwide declines in insect biomass,
abundance, and richness (Dirzo et al. 2014, Hallmann et al. 2017, IPBES 2019, Cardoso et
al. 2020, Wagner et al. 2021). The intensive use of agrochemicals is considered to be a major
driver of insect decline (Brühl and Zaller 2019, Wagner et al. 2021). Non-target insects can be
exposed to insecticides in both agricultural and non-agricultural areas. For terrestrial insects,
direct contact and oral uptake are likely the main routes of exposure. Direct contact can occur

through overspray or contact with plant or soil surfaces containing residues. Oral uptake can occur through consumption of contaminated food, e.g. plants or other invertebrates (EFSA 2015). In terrestrial non-target habitats (e.g. field margins, neighbouring meadows, hedges), such exposure is mainly caused by overspray and spray drift (Schmitz et al. 2014). Spray drift is caused by wind during insecticide application and can contaminate habitats up to 100 m away from the field (van de Zande et al. 2015). Consequently, areas close to the field such as field margins, are particularly affected, since the first metre of a field margin is exposed to both overspray and spray drift. Estimates of pesticide deposition here range from 30% to 58% of the field rate (Schmitz et al. 2014, van de Zande et al. 2015). At the same time, semi-natural habitats such as field margins make up a significant proportion of insect habitats in modern agro-ecosystems (Hahn et al. 2014). It is therefore all the more alarming that there is increasing evidence that modern agro-ecosystems, as well as non-agricultural landscapes and even insects in nature preserves, are highly contaminated with pesticides (Brühl et al. 2021, 2024, Sabzevari and Hofman 2022).

While pesticide exposure of non-target insects seems quite obvious, the effect is still poorly understood. Although modern insecticides are sometimes claimed to be less toxic to important beneficial insects such as the Western honeybee *Apis mellifera* than to the target pest (e.g. based on the German bee risk levels B1-B4), sensitivity to insecticides can vary even between closely related species and is generally difficult to predict (Sánchez-Bayo 2012, Cabrera et al. 2017). The current EU insecticide registration protocol requires sensitivity testing for a limited number of non-target insects. These are typically the honeybee, parasitoid wasps, predatory mites, and individual representatives of large beetle families such as the Coccinellidae, Carabidae or Staphylinidae (Main et al. 2018). Furthermore, the vast majority of available ecotoxicological studies involve either aquatic invertebrates or terrestrial beneficial insects such as pollinators or predators (Teder and Knapp 2019, Rumschlag et al. 2020, Siviter and Muth 2020). Herbivorous insects, which account for about 50% of the total insect richness, have been largely neglected (Pisa et al. 2015). Another shortcoming in the ecotoxicology literature is the underrepresentation of community-level studies, i.e. evaluating a range of species with different sensitivities and effects on changes in predator prey relationships. The study of community-level effects in realistic exposure scenarios is an important cornerstone for assessing the true ecological consequences of insecticide contamination. Consistent community structure effects of insecticide contamination have already been shown for aquatic systems (Rumschlag et al. 2020). However, similar studies are lacking for terrestrial systems.

In this study, we address this research gap on insecticide exposure of non-target herbivorous insects, focusing on three main aspects: (1) realistic exposure scenarios, (2) community-level effects, and (3) differential sensitivity between closely related species and between sexes of the same species. We chose plant bugs (Heteroptera: Miridae) as a model group because they are a highly diverse family within the Heteroptera and a common component of non-target insect communities in agroecosystems (Wachmann et al. 2004). Plant bugs are highly abundant in meadows, but can also be found in grassy field margins, as many species are associated with Poaceae (Denys and Tscharntke 2002, Wachmann et al. 2004). These species feed on the panicles at the top of grasses, making them even more vulnerable to insecticide deposition than ground-dwelling insects (Kühne et al. 2002). However, this group of insects has been overlooked in previous studies on non-target effects of insecticides and is not considered in the risk assessment procedure for plant protection products. We argue that plant bugs are not only an ecologically important group to study, as true bugs make up a

significant proportion of many bird diets (Exernová et al. 2003), but also provide an opportunity
to investigate community-level effects and potential differences in sensitivity between species.

We conducted field, greenhouse, and laboratory experiments to mimic insecticide exposure
in different scenarios. We used the neurotoxic neonicotinoid insecticide Mospilan®SG
(Cheminova Deutschland GmbH & Co. KG, Stade, Germany) with the active ingredient
acetamiprid in all experiments. Mospilan®SG is applied by spraying and is used in field crops
such as rapeseed and potato, in orchards, in wine growing and in floriculture, in particular
against chewing sucking pests. The active substance acetamiprid is both a contact and a
systemic insecticide, as it can also be absorbed by plants and distributed throughout their
tissues. Among other neonicotinoids, acetamiprid is used worldwide, but in the European
Union it is the only neonicotinoid still registered for open-field use. In 2022, sales of the active
ingredient in Germany alone amounted to almost 28 tonnes (BVL 2023).

In the field experiment, we sprayed three concentrations of Mospilan®SG in semi-open field
plots with wild plant bug populations to simulate in-field and off-field exposure scenarios (Fig.
1A). The field plots were located in an extensively managed meadow in southern Germany.
We selected treatments based on deposition estimates as follows: '100%' = maximum field
rate; '30%' = exposure estimate for 1 m field distance (overspray + spray drift); '0.15%' =
estimate for 20 m field distance (spray drift) (Fig. 1B). Plant bugs were sampled at 2, 9, 16, 24
and 30 days after application using a sweep net. We evaluated the total number of individuals
of mirid populations within the field plots and conducted species-level analyses focusing on
the three most abundant species: *Stenotus binotatus* (Fabricius, 1794), *Leptopterna dolabrata*
(Linnaeus, 1758), and *Megaloceroea recticornis* (Geoffroy, 1785). These three species
overlap in phenology (May – August) and host plants (Poaceae), and also differ in size (on
average 6.6 mm *S. b.*, 8.3 mm *L. d.*, and 9.1 mm *M. r.*): We also conducted pesticide analyses
of vegetation samples from the plots to monitor insecticide exposure and metabolism in plants
over time.
We further assessed insecticide susceptibility in a greenhouse experiment using host plants
(Poaceae from the same meadow) that were sprayed with Mospilan®SG two days earlier to
simulate an indirect exposure, where non-target insects enter a habitat after insecticide
deposition. In this experiment, we recorded the survival of the three focal species after four
days of exposure (i.e., after 6 days of plant exposure). We also analysed the insecticide
exposure of the host plants two days after application. Finally, we performed dose-response
assays in the laboratory, where we applied different concentrations of Mospilan®SG dorsally
to individual plant bugs (see Methods) and recorded mortality after 24 hours. This approach
allowed us to investigate intrinsic differences in contact sensitivity between species and
between conspecific males and females without confounding factors, such as potential
behavioural avoidance of exposure.

Figure 1: Overview of the field experiment and concept of treatment choice. (A) Field
enclosures are separated by 2 m. Coloration indicates Mospilan®SG – treatment based on
contamination estimates. Drawings created with AUTODESK AutoCAD 2023 and Fusion 360.
(B) Treatment choice (schematic drawing) is based on literature estimates: 100% = maximum
field rate, 30% = contamination estimate for 1 m field distance (overspray + spray drift, Schmitz
et al. 2013), 0.15% = contamination estimate for 20 m field distance (spray drift, JKI 2020).
Graph in panel B created with BioRender.com.

2. Results

2.1. Field experiment

Neonicotinoid application reduced mirid abundance by 83% (100% treatment) and 82% (30%
treatment) after two days compared to the control (100%: p<0.0001; 30%: p<0.0001; Tukey-
test) (Fig. 2A). Nine days after exposure, the abundance was reduced by 70% (100%
treatment) and 57% (30% treatment) compared to the control (100%: p<0.0001; 30%:
p<0.001; Tukey-test). This pattern was less pronounced in later samples, while the total plant
bug population declined in all treatments, including the control, presumably due to plant bug
phenology. No difference in mean counts was observed between the control and 0.15%
treatments and between the 30% and 100% treatments at any sampling event.

Focusing on the three most abundant species, the mean counts for *S. binotatus*, *L. dolabrata*
and *M. recticornis* in the 30% and 100% treatment were significantly reduced compared to the
control two days after insecticide application (*S. b.* 100%: p<0.0001, *S. b.* 30%: p<0.0001; *L.*
d. 100%: p=0.0021; *L. d.* 30%: p<0.001; *M. r.* 100%: p=0.031; M. r. 30%: p=0.024) (Fig. 2B).
This pattern partially shifted with later samplings, however, in line with the known phenologies
the populations drastically declined in later samplings, including the control (Fig. 3).

Days after Application	Contrast	Estimate	Standard Error	p-Value
2	100% - Control	-1.77	0.20	<0.0001
	30% - Control	-1.73	0.20	<0.0001
9	100% - Control	-1.25	0.21	<0.0001
	30% - Control	-0.85	0.20	=0.0001

Species	Contrast	Estimate	Standard Error	p-Value
S. binotatus	100% - Control	-1.85	0.36	<0.0001
	30% - Control	-1.83	0.37	<0.0001
L. dolabrata	100% - Control	-1.70	0.48	=0.0021
	30% - Control	-2.45	0.64	=0.0007
M. recticornis	100% - Control	-1.10	0.40	=0.0313
	30% - Control	-1.18	0.42	=0.0240

Figure 2: Exposure to Mospilan®SG reduces plant bug abundance and causes a species-specific response. A: Mean number of imagines and nymphs two- and nine days after insecticide application (± standard error). B: Mean number of imagines two days after insecticide application separated by species (± standard error). *Stenotus binotatus* (left), *Leptopterna dolabrata* (center) and *Megaloceroea recticornis* (right). Means within the same sampling day (A) or the same species (B) not sharing the same letter are significantly different (Tukey-test at the 5% level of significance). Results for significant contrasts are provided in the adjoining tables. Numbers above bars indicate percentage difference to control (rounded numbers); n = 5 for all treatments and species (panel A + B). Insect images © Gerhard Strauss, Biberach.

Days After Application

Figure 3: Population development in the field trial for *Stenotus binotatus* (left), *Leptopterna dolabrata* (center) and *Megaloceroea recticornis* (right). Curves are fitted with the Kernel method (local fit: linear). Mean number of imagines per treatment (± standard error); n = 5 for all treatments and samplings. Insect images © Gerhard Strauss, Biberach.

Pesticide analysis of vegetation samples from the enclosures taken after 7-, 17- and 30 days revealed residues of acetamiprid and its metabolite acetamiprid-N-desmethyl in the 100% and the 30% treatments (Table 1). No residues were found in the 0.15% treatment or in the control at any sampling time. Further information on pesticide analysis is provided in supplementary file 1.

Table 1: Mean concentrations of acetamiprid and the metabolite acetamiprid-N-desmethyl (± SD) per gram dry mass of plant tissue sampled inside the 100% and 30% field enclosures (LC-MS) after application of Mospilan®SG in the field experiment. Numbers are rounded, n = 5 for each treatment and substance.

	Treatment	Mean Amount Day 7 [ng/g]	Mean Amount Day 17 [ng/g]	Mean Amount Day 30 [ng/g]	Mean Amount Day 60 [ng/g]
acetamiprid	100%	865 (± 628)	118 (±66)	219 (±135)	0
acetamiprid-N-desmethyl	100%	497 (±175)	119 (±52)	81 (±47)	0

acetamiprid	30%	255 (±111)	46 (±34)	1 (±3)	0
acetamiprid-N-desmethyl	30%	157 (±16)	59 (±51)	0	0

2.2. Greenhouse experiment

Survival of the three plant bug species four days after being placed on plants sprayed 48 h before with 30% of the maximum field application rate of Mospilan®SG was reduced in all species. Specifically, survival was reduced by 88% for *S. binotatus*, 39% for *L. dolabrata*, and 35% for *M. recticornis* compared to the respective controls (Fig. 4). In none of the nine replicates did any individuals of *S. binotatus* survive the insecticide treatment.

Figure 4: Survival of mirids after four days of exposure to plants treated with 30%-Mospilan®SG. Plants were sprayed with insecticide two days prior to plant bug release. Survival rates at day four are given for *Stenotus binotatus*, *Leptopterna dolabrata* and *Megaloceroea recticornis*. Numbers within bars indicate difference between control survival and treatment survival; n = 9. Insect images © Gerhard Strauss, Biberach.

Acetamiprid and the metabolite acetamiprid-N-desmethyl were found in all samples collected from treated plants two days after application (i.e. the time of plant bug release). The mean concentration was 267 (±78) ng/g plant material for acetamiprid and 62 (±33) ng/g for acetamiprid-N-desmethyl (dry weight; n = 9; rounded numbers). Further information on pesticide analysis is provided in supplementary file 1.

2.3. Dose-Response assays

All three species showed high sensitivity to the Mospilan®SG treatment in the dose-response assays in the laboratory (Fig. 4). The observed LD_{50} values correspond to only 12.8% of the maximum field rate for *L. dolabrata*, 9.4% for *M. recticornis*, and 1.0% for *S. binotatus*. In

addition, species-specific differences were observed, with *S. binotatus* being significantly more sensitive to the insecticide than the other two species based on LD_{50} comparisons (*L. dolabrata* – *S. binotatus*: p<0.01, t=3.33; *M. recticornis* – *S. binotatus*: p=0.02, t=2.33).

Species	LD50 of Mospilan®SG (± SE) [ng/µl]			N
	species	male	female	
L. dolabrata	53.17[a] (± 14.75)	6.8[a] (± 2.7)	150.4[b] (± 22.5)	8
M. recticornis	39.04[a] (± 14.99)	5.5[a] (± 2.1)	114.5[b] (± 31.2)	6
S. binotatus	4.12[b] (± 1.41)	—	—	6

Figure 4: Mospilan®SG dose-response curves for all species (A) and separated by sex for *Leptopterna dolabrata* (B), *Megaloceroea recticornis* (C) and *Stenotus binotatus* (D). Data points represent mean values per concentration. Derived LD_{50} values are presented in the table below. Separate LD_{50} values for male and female specimens were generated only for those species in which the sensitivity to the insecticide was significantly different between the sexes based on the fitted generalised linear mixed model (GLMM). LD_{50} values of species

not sharing a common letter are significantly different. LD_{50} values were compared using the
compParm function of the 'drc' package in R. LD_{50} values of males and females of the same
species not sharing a common letter are significantly different. Abbreviations: SE = standard
error; N = number of biological replicates (i.e., petri dishes) per concentration. Total number
of specimens tested: 245 for *S. binotatus*, 332 for *L. dolabrata*, and 224 for *M. recticornis*. Note
that mean values for the lowest Mospilan®SG treatment for males and females of *M. r.* (C)
consist of only two data points instead of three. Insect images © Gerhard Strauss, Biberach.
239
240
241 Besides a very high overall toxicity of Mospilan®SG, we observed significant differences
242 between males and females of two species. Based on LD_{50} values, males of *L. dolabrata* and
243 *M. recticornis* were >20 times more susceptible than females (*L. d.* t=6.33, p<0.0001; *M. r.*
244 t=3.48, p=0.0005). There was no difference between sexes for *S. binotatus* based on the fitted
245 GLMM prior to the dose-response analysis.
246
247
248

3. Discussion

Insecticides are a prime suspect in insect declines, but there is a lack of empirical evidence of
real-world effects on non-target species outside the cropping area. To understand the impact
of terrestrial insecticide contamination on non-target insect communities, we conducted a
series of field, greenhouse, and laboratory experiments to investigate community-, species-
and sex-specific responses in the model insect group of mirid bugs. Mirid bug species in the
field experiment were highly sensitive to neonicotinoid exposure. We found species-specific
responses across different exposure scenarios and drastic differences in sensitivity between
males and female of two species tested in the laboratory.

3.1. Mirid communities are highly sensitive to insecticide contamination

Consistent across experiments, our study shows a high susceptibility of mirids to insecticide
exposure. In the field experiment, we aimed to simulate a scenario of direct drift contamination
with specimens present in the respective off-field environment during spray application. A
drastic reduction in mirid abundance was observed in the scenario representing exposure
within one metre of the field (30% of maximum field rate).
A second scenario of indirect drift contamination for the same field distance was simulated in
the greenhouse experiment, as it is also possible that insects enter an area only after drift
contamination occurred (i.e., after insecticides were applied). Residues of the compound that
may still be present on surfaces (soil or plant material) hours or days later can still be absorbed
through an insect's cuticle, e.g. via the tarsi (Yu 2015), or ingested during feeding. A special
case, however, is that of systemic insecticides, where the active ingredient can also be taken
up by host plants of non-target insects, providing a route for oral contamination of piercing
sucking herbivorous insects (e.g., mirids) for presumably even longer periods of time. In our
experiment, even two days after host plants were treated with 30% of the maximum field rate
of Mospilan®SG, the survival rates of mirids placed on these plants were significantly reduced.
274
These results clearly demonstrate a detrimental effect of insecticide contamination on non-
target mirid populations located close to arable fields. Further, a generally high sensitivity to
neonicotinoid insecticides of this ecologically important insect group that occupies an
important position in the trophic pyramid, is revealed (Exernová et al. 2003). This was further

supported by the insecticide dose-response assays. Here, LD_{50} values for three mirid species
were at concentrations as low as 1% of the original field rate. To put this difference into context,
these values can be compared to those of common model organisms such as the Western
honeybee *A. mellifera*. However, LD_{50} values from dose-response assays are typically given
as a concentration of active ingredient per microliter or per individual. To establish
comparability, the values we obtained using a commercial insecticide formulation can be
adjusted to reflect the 20% active ingredient acetamiprid content. This corresponds to 10.63
ng/bug for *L. dolabrata*, 7.81 ng/bug for *M. recticornis* and 0.82 ng/bug for *S. binotatus.* For
honeybees an LD_{50} of ~ 8090 ng active ingredient per bee is reported as contact exposure
value (European Commission 2004). Accordingly, the sensitivity of the tested mirid species is
~ 760 times (*L. dolabrata*) to ~ 9866 times (*S. binotatus*) higher.
However, the toxicity of a pure active ingredient may differ from that of a product formulation
(Nagy et al. 2020). Therefore, such comparisons should be made with caution. Nevertheless,
this huge difference in LD_{50} values shows that mirids, and probably other non-target insects,
are very sensitive to neonicotinoid insecticides and consequently to environmental
contamination at only a fraction of the applied field rates. This group of insects must therefore
be given greater consideration in risk assessment (see 3.4 below).
296
3.2. Species-specific sensitivity
The three mirid species that were tested in this study can be considered as model species
representing non-target phytophagous insects. In this context, these species are of particular
interest because they share a similar ecology (e.g., feeding style, phenology and host plants,
Wachmann et al. 2004) and yet showed markedly different insecticide susceptibility under the
experimental contamination scenarios. Differences in species response were already
suggested by the field experiment and further confirmed by the greenhouse experiment with
treated plants and the dose-response assays in the laboratory. As differences in species
response were not only observed in the field experiment, it is unlikely that they are related to
feeding preferences or the spatial position of the bugs on the plants during spraying, but rather
follow from species-specific physiological traits.
Physiological factors that have been proposed to influence sensitivity include, for example,
differences in the activity and abundance of detoxification enzymes, or simply body size
(Müller 2018). Notably, the smallest (and lightest) species *S. binotatus* was always the most
sensitive in our greenhouse- and dose-response assays, suggesting that body size or mass
is an important factor. Goulson (2013) compared available LD_{50} values from neonicotinoid
studies (topical and oral) and stated that much of the variation in sensitivity between several
insect taxa could be explained by differences in body size/weight. For example, the LD_{50} for
the same neonicotinoid of the highly sensitive *Nilaparvata lugens* (body weight ~1 mg) is
similar to the one of the least sensitive *Leptinotarsa decemlineata* (body weight ~ 130 mg)
expressed as ng active ingredient per mg body weight (0.82 and 0.67). Considering the
species we tested, although *L. dolabrata* has more than twice the body mass of *M. recticornis*,
the LD_{50} values are almost identical when expressed as concentration of active ingredient per
body mass (L. d.: 0.53, M. r.: 0.66; see supplementary file 2). At the same time, there is a
more than 4-fold difference to the LD_{50} of *S. binotatus* (0.13). Therefore, body weight alone
cannot explain the difference in sensitivity between these species.
While the physiological mechanisms involved remain to be investigated, differences in species
susceptibility could permanently alter local community structure under a constant or repeated
contamination scenario. Changes in invertebrate community composition have been
associated with pesticide contamination in the past. However, this has mainly been studied in

freshwater communities (Beketov et al. 2013, Rumschlag et al. 2020), whereas the effects on terrestrial communities are less well understood. A recent meta-analysis showed that pesticide contamination reduces soil invertebrate diversity (Beaumelle et al. 2023). Main et al. (2020) have also linked neonicotinoid soil contamination to reduced species richness of wild bees in field margins. More research is needed to understand the long-term consequences of pesticide exposure on terrestrial non-target invertebrate communities on a landscape scale. Particular attention should be paid to univoltine species with short phenologies that may overlap with common pesticide application regimes. Populations of such species, such as the mirids in this study, should be particularly limited to cope with pesticide contamination should it occur during their short and only reproductive phase throughout the year.

3.3. Sex-specific sensitivity

In addition to differences in sensitivity at the species level, we found that males of *L. dolabrata* and *M. recticornis* were significantly more sensitive to the insecticide than the corresponding females. Sex-specific insecticide sensitivity is a pattern reported for many insects, with males often being more susceptible than females (Huntzinger et al. 2008, Andreazza et al. 2020). However, there are also contrary reports in other species (Navarro-Roldán et al. 2017).

Potential factors involved in sex-specific susceptibility partly overlap with those likely driving the observed species-specific differences. The most likely factor is again body mass, especially in sexually dimorphic species such as *L. dolabrata* and *M. recticornis*. In both species, males had a lower mean fresh body mass than females, with a 2.9-fold difference in *L. dolabrata* and a 2.2-fold difference in *M. recticornis* (n = 10 per species and sex). This is not correlated with the difference in LD_{50} values between males and females, which is 22-fold for *L. dolabrata* and 21-fold for *M. recticornis*. However, such relationships may not be necessarily linear.

Another source for differential susceptibility is at the enzymatic level. Frohlich (1988) showed that the activity of detoxification enzymes decreased with age in males of the solitary bee *Megachile rotunda*, but not in females. An age-dependent change in enzyme activity was also observed by Guirgus and Brindley (1975) when they compared 1-day-old males of *Megachile pacifica* with 4-day-old males, both of which had been previously treated with a fungicide. Thus, age also appears to play a critical role in susceptibility of insects to insecticides.

The higher control mortality observed in males in our dose-response study (which was accounted for in the models) would suggest an older initial age compared to females in the same cohort. However, to our knowledge, there is no evidence of a sex-specific time lag in the emergence of adult specimens of these species, and according to our observations, males and females appeared simultaneously in the field. However, for *L. dolabrata*, Wachmann et al. (2004) suggest that females live longer than males. A lower vitality of male specimens due to a shorter life cycle could therefore explain a higher control mortality and further higher susceptibility to stressors such as an insecticide. Regardless of the underlying mechanisms, the observed differences could lead to a shift in sex ratios, in this case towards females. This could ultimately lead to a reduction in abundance and, in the long term, a shift in species composition.

3.4. Environmental risk assessment

To assess the risk of insecticide contamination to a non-target species, it is essential to determine the true vulnerability of that species in its environment. Vulnerability involves both physiological sensitivity and exposure to an insecticide (de Lange et al. 2012). We have shown

that mirids, as a non-target group, are physiologically highly sensitive to insecticides.
Furthermore, the species tested here are potentially found in highly exposed areas of the off-
field environment, i.e. in the upper vegetation of field margins (Kühne et al. 2002, Wachmann
et al. 2004). We also demonstrated that exposure to a systemic insecticide extends beyond
the initial contamination, as plant bugs placed on previously sprayed plants showed a drastic
reduction in survival. Considering that in the field experiment we were able to detect
acetamiprid in host plant tissue up to 30 days after application, the threat to non-target plant
bugs is clear. It is therefore all the more surprising that the current risk assessment does not
consider this group or other herbivorous insects in the non-target insect community. In
addition, field margins less than 3 m wide are not currently considered to be terrestrial non-
target habitats for pesticide application (Kühne et al. 2000). Consequently, this habitat and the
insects living in it are not protected from pesticide contamination (Hahn et al. 2014).
Roß-Nickoll et al. (2004) argue that the test procedures used in the EU risk assessment are
insufficient because the test species used do not adequately reflect the non-target insect
community. As a result, the disruption to these communities caused by pesticide
contamination is not taken into account. This discrepancy is further illustrated by the drastic
difference in LD_{50} values between the honeybee model and the mirid species we tested which
suggests an up to 10,000x higher toxicity for mirids compared to the honeybee. There is also
the problem that risk assessment is currently only done at the species level, and that sex-
specific sensitivity is not considered. However, the up to 22-fold difference in LD50 values
between sexes that we found could scale to detrimental long-term effects on community
composition. This calls for an increase of the so-called uncertainty factors used to cover
different species and sexes from currently 10 to more than 1000. Consequently, and in light
of the renewal of the authorisation of acetamiprid until 2033 (European Commission 2018),
we argue that the current European risk assessment system urgently needs to be adapted
accordingly to adequately address the environmental risk of pesticide contamination.

4. Methods

4.1. Statistical Analysis and Graphical Representation

Statistical analysis was conducted using R (version 4.3.0, R Core Team 2023) and R-Studio
(version 2023.6.1.524, Posit team 2023). Therefore, only specific packages are cited in the
following sections.
All graphs showing bar or stacked bar charts were generated using JMP®Pro (version 17.2.0,
SAS Institute Inc. 2023). Dose-response curves were generated with R using the package
'drc' (Ritz et al. 2015) and edited with Inkscape (version 1.3.2, Inkscape Project 2020).

4.2. Field Experiment

4.2.1. Construction

In May 2022, we constructed 20 field plots (2 x 4 m) on an extensively managed meadow
(48°42'31''N 9°12'48''E) in the botanical garden of the University of Hohenheim, Stuttgart,
Germany. In every corner of each plot, a 1.4 m long wooden pole was vertically inserted 40
cm into the soil. An 85 cm high insect gauze (mesh size: 0.27 x 0.79 mm, *Howitec Netting bv*,
AH Joure, the Netherlands) was vertically stretched around these poles. About 10 cm of the
gauze was inserted into the soil to form a barrier, while 75 cm remained aboveground fixed to
the poles. Thereby, the top of the field plot remained uncovered. These semi-open field plots
(henceforth, enclosures) were additionally braced with tent cords that were attached at every

corner. On an area of 340 m², we set up 20 enclosures in seven rows with each being
separated by the neighbor by 2 m on all sides. The field experiment was laid out as a
randomized block design with four treatments: three different concentrations of insecticide and
a control. Each enclosure was defined as an individual block with every treatment being
replicated five times (Fig. 1). The choice of blocks was made with respect to the environmental
conditions on the site. Due to surrounding trees, the duration of solar radiation was longest in
the northwestern area and shortest in the southeastern area. In addition, the test site was
located on a south-facing slope. This resulted in a soil moisture gradient towards the south
slope. The following blocks were assigned accordingly: A = upper slope + high solar radiation,
B = middle slope + high solar radiation, C = middle slope + medium solar radiation, D = lower
slope + medium solar radiation, E = lower slope + low solar radiation (see supplementary file
3).
433
4.2.2. Insecticide application
We used the nerve-acting neonicotinoid insecticide Mospilan®SG (FMC Agricultural
Solutions, Cheminova Deutschland GmbH & Co. KG, PO Box 2047, Stade, Germany), with
the active ingredient acetamiprid for all experiments described in this study.
The following concentrations were applied in the field experiment by spraying: (1) typical field
application rate of 250 g/ha in 600 l of water (100%), (2) 30% of the field application rate, (3)
0.15% of the field application rate. These values are based on estimates for overspray- and
drift contamination with respect to the field distance (Schmitz et al. 2014, JKI 2020). The
underlying concept is illustrated in figure 1A. In the control, only water was applied. The
application rate was scaled down to the field plot area of eight square meters (Tab. 2). The
amount of 0.48 l containing the respective insecticide concentration was applied equally to
each field plot using the *GLORIA prima 3* hand sprayer (GLORIA Haus- & Gartengeräte
GmbH, Witten, Germany).
447
448 **Table 2:** Application rates of Mospilan®SG of the field experiment.

Treatment	Application Rate	Application Rate in Field Plot	Stock Solution
100%	250 g/ha in 600 l water	200 mg/8 m² in 0.48 l water	1.25 g in 3 l water
30%	75 g/ha in 600 l water	60 mg/8 m² in 0.48 l water	0.375 g in 3 l water
0.15%	0.375 g/ha in 600 l water	0.3 mg/8 m² in 0.48 l water	3.125 mg in 5 l water

449
The hand compression sprayer was calibrated before use and the volume output was
measured. The mean deviation of liquid output was around 1%. Details of the calibration
procedure are given in supplementary file 4.
Application started with the control (water only) followed by the insecticide treatments in
increasing concentration. Between treatments, the hand sprayer was completely emptied and
re-calibrated. To control for uneven application due to a change in volumetric flow rate
(because of decreasing pressure during spraying), the field plots were divided into four equal
quarters. Each quarter was assigned a number from one to four. Spraying would always start
in the quarter numbered one and end in the quarter numbered four. The order of numbers was
randomized within plots of the same treatment.
During application inside the plots, the spray gun was opened with the nozzle pointing
downwards towards the soil while moving in a standardized pattern of vertical and horizontal
trajectories at a height of 80 cm above the ground. This was done for 21 seconds per quarter
(1/4 of 84 s being the total application time per plot) and then repeated for the other quarters

of the same field plot. To avoid contamination of surrounding plots, each plot was enclosed by
a 2 m high plastic sheet during application.

4.2.3. Exposure analysis

Seven days after insecticide application, vegetation samples were collected from inside the
field plots to quantify the amounts of the active ingredient acetamiprid and the metabolite
acetamiprid-N-desmethyl by LC-MS analysis. This was done to demonstrate the potential
presence of the insecticide inside the enclosures with respect to the different treatments even
days after application as well as its metabolization within plant the tissue. The full analytical
procedure and results are given in supplementary file 1.

4.2.4. Insect sampling and identification

Insects in the upper vegetation of each field plot were regularly sampled with a sweep net
(Flexi-Kescher Dreieck; Bioform, Nürnberg, Germany). Sampling took place 2, 9, 16, 24 and
30 days after insecticide application. While walking parallel to the plot, the net was swept four
times forwards and backwards across the upper part of the vegetation. After the fourth sweep,
the net was immediately closed by hand. The entire contents were then transferred into a
three-litre zip-lock plastic bag (Profissimo, dm-drogerie markt GmbH + Co. KG, Karlsruhe,
Germany). The plastic bags were stored at -20 °C for at least 24 hours to kill all arthropods
inside.
For evaluation, individual bags were stored at 8 °C for 30 min to carefully unfreeze the
specimens. The entire contents of the plastic bag were then transferred to a plastic tray and
the plant bugs (imagines and nymphs) were separated with tweezers. Nymphs were counted
and transferred to a 2 ml screw-capped microtube (SARSTED AG & Co. KG, Numbrecht,
Germany) filled with 70% ethanol for preservation. Imagines were sorted by morphospecies,
and the corresponding individuals were counted. For each morphospecies, one male and one
female specimen (if available) were separated for preparation. The specimens were mounted
on paper cards using wallpaper glue. In the case of male specimens, the aedeagi were
removed beforehand and glued next to the respective specimen. Remaining specimens of
each morphospecies were preserved in 70% ethanol as described above. All morphospecies
were identified to species level using the Hackston (2023) Miridae key and the Corisa software
(Strauss and Simon 2021).

4.2.5. Statistical evaluation

A generalised linear mixed model (GLMM) was fitted for the combined individual counts of
nymphal and adult plant bugs using the 'lme4' package (version 1.1-33, Bates et al. 2015)
with '*count*' as the explanatory variable, '*treatment*' as a fixed effect, '*sampling*' as a co-factor,
and '*block*' and '*plot*' as random effects assuming a Poisson distribution. An interaction
between treatment and sampling was included. Model fit was assessed by residual analysis,
considering the R^2 value and the Akaike Information Criterion (AIC), and by checking for
possible over-dispersion. Significant factors and interactions within the model were identified
using a post-model ANOVA. Those were '*treatment*', '*sampling*' and the interaction between
both factors. Pairwise post hoc multiple comparisons by sampling were then executed using
the Tukey-test based on estimated marginal means using the emmeans package (Lenth 2023)
considering the *treatment-sampling* interaction.
In a second step, a separate GLMM was fitted including only the count data of *S. binotatus*,
L. dolabrata and *M. recticornis* and only the data of the first sampling (i.e., two days after
insecticide application). In this GLMM, '*count*' was the explanatory variable with '*treatment*' as

a fixed effect, '*species*' as a co-factor, an interaction between '*treatment*' and '*species*' and
'*block*' and '*plot*' as random effects, assuming a Poisson distribution. Model fit was assessed
as described above. Significant factors and interactions within the model were identified using
a post-model ANOVA. Pairwise post hoc multiple comparisons by species were then executed
using the Tukey-test based on estimated marginal means considering the treatment-species
interaction.
Based on the control sampling data of the three species, the populations were stable in the
early samples, but declined dramatically in later samples as expected based on the known
phenologies. In order to compare acute species-specific treatment response, we therefore did
not include data beyond the first sampling.

The complete R-code is provided in supplementary file 5.

4.3. Greenhouse Experiment
523
4.3.1. Host plant preparation
Eighteen plots of 30 cm diameter with live plants, consisting mainly of grasses, which are the
host plants of the three mirid species, were collected from a meadow with a shovel and
transferred to 12-liter plant pots. These were previously filled halfway with a 50:50 mixture of
sand and humus. Only sections of the meadow with a high proportion of grasses were selected
to provide sufficient host plants for the plant bugs. The pots were placed in an outdoor test
facility for one month and watered as needed until the grasses had developed seed heads.
531
4.3.2. Insecticide application and experimental procedure
The pots were placed in an 8 m^2 rectangular area, surrounded by 2 m high vertical plastic
sheathing. Spray application was carried out using the same protocol and equipment as
described for the field experiment. Two treatments were applied: a solution containing 30%
field rate concentration of Mospilan®SG and a control containing only water. For each
treatment application, nine randomly selected plant pots were placed within the application
area, all 30 cm apart. The control was applied first. The pots were then placed in the outdoor
test facility. A total of four blocks were formed, with two blocks for each treatment, one
containing five pots and the other one containing four. Treatment and control blocks with the
same number of replicates were placed opposite to each other (Fig. 5). In the center of each
pot, a 2 m wooden stick was vertically inserted. At the top of the stick, a 30 cm diameter
herbacaeus ring was fixed horizontally. Fine insect gauze was then placed over the
construction from the top of the stick down to the plant pot, forming a closed tube around the
host plants.
546
547
548
549

Figure 5: Schematic representation of the experimental setup. "M"=Mospilan®SG
treatment, "C"=Control treatment. Treatment and control blocks with the same number of
replicates are placed opposite to each other.

Two days after application, plant bugs were caught using a butterfly sweep net (Flexi-Kescher
Dreieck; Bioform, Nürnberg, Germany) in the same extensively managed meadow from which
the plant samples were collected. Specimens were immediately transferred to 30 cm x 30 cm
AERARIUM mesh cages (Bioform, Nürnberg, Germany), which were then placed in an 8 °C
room. After 15 min of settling, the cages were opened and specimens of *S. binotatus*, *L.
dolabrata* and *M. recticornis* were sorted by species (identified by naked eye) and transferred
to petri dishes. For each replicate (= one plant pot), three petri dishes were prepared, each
containing six specimens of either species (males and females mixed). After assuring that all
bugs were vital, 18 plant bugs (6 per species) were then released on top of the plants inside
the insect gauze. At the same time, plant samples for HPLC analysis were collected from each
Mospilan®SG treated plot to demonstrate that the active ingredient acetamiprid was present
at the time of plant bug release. Each morning the plants were sprayed with a pressurized
water bottle to simulate morning dew. After four days (six days after insecticide application)
the experiment was terminated, and alive specimens were counted. Specimens from each
replicate were stored separately in 2 ml micro tubes with screw caps (SARSTEDT AG & Co.
KG, Germany) filled with 70% ethanol. Following the assay, species identity was again verified
using a binocular and the Hackston (2023) Miridae key and the Corisa software (Strauss and
Simon 2021).

4.3.3. Exposure analysis
Two days after insecticide application, just before the plant bugs were released into the cages,
vegetation samples were collected from each replicate of the insecticide treatment group.
Unlike the field experiment, the control group was not sampled as the control pots were
sprayed first and then removed before insecticide application. LC-MS analysis was used to
quantify the amounts of the active ingredient acetamiprid and the metabolite acetamiprid-N-
desmethyl. The full analytical procedure and results are provided in supplementary file 1.

4.3.4. Statistical evaluation

A linear mixed regression model, was fitted using the 'lme4' package (Bates et al. 2015) with '*alive*'-specimens as the response variable, '*treatment*' as fixed effect and '*species*' as a co-factor, and an interaction between '*treatment*' and '*species*'. The random factor '*block*' was first included but did not explain any variance of the data and was therefore excluded from the model. The random factor '*pot*', providing a unique identifier for each pot, was included to account for the fact that '*species*' was nested within '*treatment*', allowing an individual baseline morality for each replicate. Hence, if the interaction was significant, it was not due to a replicate effect, allowing to explore species differences in survival only. Significance of the interaction between '*treatment*' and '*species*' was confirmed by a post model ANOVA.

Fitting a follow-up model with 'species' as the explanatory variable was not possible because there were no individuals of *S. binotatus* that survived the insecticide treatment in any of the nine replicates, making it impossible to estimate a standard error. Accordingly, we report the difference in treatment effect as the percentage difference in survival rates between the control and insecticide treatments ('insecticide effect'). This considers the baseline mortality of each species observed in the control.

The complete R-code is provided in supplementary file 5.

4.4. Dose-Response Assays

4.4.1. Insecticide preparation

A stock solution of 4.16 mg/ml Mospilan®SG, dissolved in tap water, was prepared corresponding to 10 times the highest recommended field rate of 0.416mg/ml. Stock and treatment solutions were stored in 1.5 ml ND8 amber glass screw-neck vials (LABSOLUTE – TH GEYER) in the dark at 4°C. Stock solutions were renewed weekly. Prior to each dose-response assay, a dilution series was prepared based on the stock solution on the day of application (Tab. 3).

Table 3: Treatments applied in the dose-response assay and respective Mospilan®SG concentrations. The 0.14% treatment was not applied to *Leptopterna dolabrata* and *Megaloceroea recticornis* due to lower sensitivity.

Treatment	Concentration [ng/µl]
Control	0.00
100%	416.00
33.30%	138.67
11.10%	46.22
3.70%	15.41
1.24%	5.14
0.41%	1.71
0.14%	0.57

Here, 0.5% TWEEN (20 was added to the solvent to break the surface tension and ensure easy application to the bug's body surface. Each treatment concentration was prepared from

617 the stock solution by pipetting volumes from higher to lower concentration treatment tubes
618 prefilled with pure solvent (see dilution protocol supplementary file 6).
619

4.4.2. Insect sampling

Plant bugs were caught in the meadows of the Hohenheim Gardens (extensively managed
grassland) using a butterfly sweep net (Bioform, Nürnberg, Germany). Specimens were
immediately transferred to 30x30 cm insect mesh cages (Bioform, Nürnberg, Germany), which
were then placed in an 8 °C room. After 15 min of settling, the cages were opened and
specimens of *S. binotatus*, *L. dolabrata* and *M. recticornis* were sorted by species (identified
by naked e
ye) and transferred to petri dishes. Two petri dishes were prepared per species and treatment
with six to four specimens each (depending on availability of specimens). Where possible,
male and female specimens were equally distributed. Only vital specimens that were actively
climbing the net were included. Petri dishes were taken to the laboratory (20°C) where the
specimens were allowed to settle again for 20 min. Prior to treatment application, all
specimens were again checked for vitality and replaced if necessary.
633

4.4.3. Topical application and evaluation

Several identical assays were run in sequence to allow for a possible cohort effect (*S.*
binotatus + *M. recticornis*: three assays; *L. dolabrata*: four assays). Each assay included two
physical replicates per concentration, each consisting of four to six specimens (2 male + 2
female / 3 male + 3 female) based on availability of specimens in the Hohenheim Gardens.
The specimens of a physical replicate were first placed in a petri dish and later transferred to
rearing boxes (see below). Accordingly, each concentration consisted of six- (*S. b.* + *M. r.*) or
eight replicates (*L. d.*) nested within three or four cohorts (Tab. 4). Note that for *M.* recticornis
the number of replicates in the lowest tested Mospilan®SG concentration (0.41%) is four.
The number of treatments for *S. binotatus* was increased for the second and third cohort
adding the 0.14% treatment due to higher sensitivity.
645

646 **Table 4:** Number of cohorts, treatments and individuals tested in the dose-response assays.
647 Eight treatments: 100% - 0.14% + control; seven treatments: 100% - 0.41% + control. Note
648 that the number of treatments for *S. binotatus* was increased for the second and third cohort.
649

Species	Cohort	Number of treatments	Individuals tested per treatment
S. binotatus	1	8	4
	2	8	6
	3	8	6
L. dolabrata	1	7	6
	2	7	6
	3	7	6
	4	7	6
M. recticornis	1	7	4
	2	7	6
	3	7	6

Bugs inside petri dishes were anaesthetized with CO2, then 1 µl of insecticide solution (and
pure solvent as a control) was applied dorsally to each specimen. Application was conducted
using the Hamilton repeating dispenser with a 50 µl Hamilton syringe (Hamilton Bonaduz AG,
Switzerland). Specimens of each replicate were then transferred to 500 ml plastic cups. The
cups contained a water supply (Eppendorf tubes filled with water and cotton wool), a bundle
of grasses from the original habitat with the top 10-15 cm of the plants cut off (stems and
heads) placed in an Eppendorf tube filled with water and cotton wool, and a paper tissue
covering the bottom of the cup. The lid was provided with several air holes. In addition, water
was sprayed into the cup to cover the grasses with droplets providing sufficient humidity and
an additional source of drinking water. Cups were stored under standard laboratory conditions.
After 24 h, mortality rates for all treatments were recorded for each replicate and sex. The
experiment was then terminated and dead and surviving specimens from each replicate were
stored separately in 2 ml micro tubes with screw caps (SARSTEDT AG & Co. KG, Germany)
filled with 70% ethanol. Following the assay, species identity and sex of all specimens were
again verified using a binocular and the Hackston (2023) Miridae key and the Corisa software
(Strauss and Simon 2021).
681
4.4.4. Dose-response analysis

Data from each assay were first analyzed with a generalized linear mixed model using the
'lme4' package (Bates et al. 2015). '*Mortality*' was included as the explanatory variable,
'*concentration*' as fixed effect, '*sex*' as co-factor and '*replicate*' and '*cohort*' as random effects.
Additionally, an interaction between '*concentration*' and '*sex*' was included. Model fit was
assessed by residual analysis, considering the R2- and AIC value, and by checking for
possible over-dispersion. Including the random effect '*replicate*' allowed to control for the fact
that '*sex*' was nested within '*concentration*', allowing an individual baseline morality for each
replicate. If the interaction was significant, it was not due to replicate effect, allowing to
generate LD_{50} values for male and female specimens of the same species separately.
A post-model ANOVA was performed confirming that '*concentration*' was a significant
predictor in all assays. A significant interaction between '*concentration*' and '*sex*' was found
for *L. dolabrata* and *M. recticonis*.
Log-logistic dose-response models were then fitted for a combined species analysis and for
each species while modelling male and female response separately using the 'drc' package
(Ritz et al. 2015). In the case of control mortality, the lower asymptote of the fitted regression
curve was not set to '0' to generate unbiased parameter estimates. The upper asymptote was
set to '1'. The best model fit was selected based on visual inspection of the regression line
and evaluation of the confidence intervals of the model parameters, considering the AIC
estimator. Dose-response curves were generated using the plot() operation. LD_{50} values were
obtained using the ED() function, including a control for possible over-dispersion of the logistic
regression model. Differences between LD_{50} values of individual curves within the same model
were analyzed using the compParm() function executing approximate t-tests with α = 0.05 and
including a correction for possible over-dispersion of the underlying model.
The complete R-code for the linear and log-logistic models is provided in supplementary file
5.
708
5. References

Andreazza F, Haddi K, Nörnberg SD, Guedes RNC, Nava DE, Oliveira EE (2020) Sex-
dependent locomotion and physiological responses shape the insecticidal

susceptibility of parasitoid wasps. Environmental Pollution 264: 114605.
https://doi.org/10.1016/j.envpol.2020.114605

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using
lme4. Journal of Statistical Software 67: 1–48. https://doi.org/10.18637/jss.v067.i01

Beaumelle L, Tison L, Eisenhauer N, Hines J, Malladi S, Pelosi C, Thouvenot L, Phillips
HRP (2023) Pesticide effects on soil fauna communities—A meta-analysis. Journal of
Applied Ecology 60: 1239–1253. https://doi.org/10.1111/1365-2664.14437

Beketov MA, Kefford BJ, Schäfer RB, Liess M (2013) Pesticides reduce regional biodiversity
of stream invertebrates. Proceedings of the National Academy of Sciences 110:
11039–11043. https://doi.org/10.1073/pnas.1305618110

Brühl CA, Zaller JG (2019) Biodiversity Decline as a Consequence of an Inappropriate
Environmental Risk Assessment of Pesticides. Frontiers in Environmental Science 7.
Available from: https://www.frontiersin.org/articles/10.3389/fenvs.2019.00177
(January 27, 2023).

Brühl CA, Engelhard N, Bakanov N, Wolfram J, Hertoge K, Zaller JG (2024) Widespread
contamination of soils and vegetation with current use pesticide residues along
altitudinal gradients in a European Alpine valley. Communications Earth &
Environment 5: 1–9. https://doi.org/10.1038/s43247-024-01220-1

Brühl CA, Bakanov N, Köthe S, Eichler L, Sorg M, Hörren T, Mühlethaler R, Meinel G,
Lehmann GUC (2021) Direct pesticide exposure of insects in nature conservation
areas in Germany. Scientific Reports 11: 24144. https://doi.org/10.1038/s41598-021-
03366-w

BVL (2023) BVL - Berichte über Inlandsabsatz und Export von Pflanzenschutzmitteln.
Inlandsabsatz und Ausfuhr von Pflanzenschutzmitteln und deren Wirkstoffen.
Available from:
https://www.bvl.bund.de/DE/Arbeitsbereiche/04_Pflanzenschutzmittel/01_Aufgaben/0
2_ZulassungPSM/03_PSMInlandsabsatzAusfuhr/psm_PSMInlandsabsatzAusfuhr_n
ode.html (February 22, 2024).

Cabrera P, Cormier D, Lucas É (2017) Differential sensitivity of an invasive and an
indigenous ladybeetle to two reduced-risk insecticides. Journal of Applied
Entomology 141: 690–701. https://doi.org/10.1111/jen.12391

Cardoso P, Barton PS, Birkhofer K, Chichorro F, Deacon C, Fartmann T, Fukushima CS,
Gaigher R, Habel JC, Hallmann CA, Hill MJ, Hochkirch A, Kwak ML, Mammola S, Ari
Noriega J, Orfinger AB, Pedraza F, Pryke JS, Roque FO, Settele J, Simaika JP,
Stork NE, Suhling F, Vorster C, Samways MJ (2020) Scientists' warning to humanity
on insect extinctions. Biological Conservation 242: 108426.
https://doi.org/10.1016/j.biocon.2020.108426

Denys C, Tscharntke T (2002) Plant-insect communities and predator-prey ratios in field
margin strips, adjacent crop fields, and fallows. Oecologia 130: 315–324.
https://doi.org/10.1007/s004420100796

Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B (2014) Defaunation in the
Anthropocene. Science 345: 401–406. https://doi.org/10.1126/science.1251817

EFSA (2015) Scientific Opinion addressing the state of the science on risk assessment of
plant protection products for non-target arthropods. EFSA Journal 13: 3996.
https://doi.org/10.2903/j.efsa.2015.3996

European Commission (2004) Review report for the active substance acetamiprid. Finalised
in the Standing Committee on the Food Chain and Animal Health at its meeting on 29
June 2004 in view of the inclusion of acetamiprid in Annex I of Council Directive
91/414/EEC. SANCO/1392/2001.

European Commission (2018) Renewing the approval of the active substance acetamiprid in
accordance with Regulation (EC) No 1107/2009 of the European Parliament and of
the Council concerning the placing of plant protection products on the market, and
amending the Annex to Commission Implementing Regulation (EU) No 540/2011.
Official Journal of the European Commission.
http://data.europa.eu/eli/reg_impl/2018/113/oj

Exernová A, Štys P, Krištín A, Volf O, Pudil M (2003) Birds as predators of true bugs
(Heteroptera) in different habitats. Biologica, Bratislava 58: 253–264.

Frohlich D (1988) Characterization of Esterases and Glutathione *S*-Transferases in a
Solitary Bee, Megachile rotundata (Fab.): Variability, Developmental
Multiplicity, Substrate Specificity, and Inhibition. Utah State University Available from:
https://digitalcommons.usu.edu/etd/8282.

Goulson D (2013) REVIEW: An overview of the environmental risks posed by neonicotinoid
insecticides. Journal of Applied Ecology 50: 977–987. https://doi.org/10.1111/1365-
2664.12111

Guirguis GN, Brindley WA (1975) Carbaryl penetration into and metabolism by alfalfa
leafcutting bees, Megachile. Journal of Agricultural and Food Chemistry 23: 274–279.
https://doi.org/10.1021/jf60198a056

Hackston M (2023) Family Miridae - Key to subfamilies and tribes occuring in Britain.
translated and adapted from Wagner E & Weber HH (1964): Héteroptères Miridae.
Faune de France 67. Fédération Française des Sociétés de Sciences Naturelles,
Paris. Available at: https://sites.google.com/view/mikes-insect-keys/mikes-insect-
keys. Available from: https://sites.google.com/view/mikes-insect-keys/mikes-insect-
keys.

Hahn M, Lenhardt PP, Brühl CA (2014) Characterization of field margins in intensified agro-
ecosystems—why narrow margins should matter in terrestrial pesticide risk
assessment and management. Integrated Environmental Assessment and
Management 10: 456–462. https://doi.org/10.1002/ieam.1535

Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W, Müller A,
Sumser H, Hörren T, Goulson D, Kroon H de (2017) More than 75 percent decline
over 27 years in total flying insect biomass in protected areas. PLOS ONE 12:
e0185809. https://doi.org/10.1371/journal.pone.0185809

Huntzinger CI, James RR, Bosch J, Kemp WP (2008) Fungicide Tests on Adult Alfalfa
Leafcutting Bees (Hymenoptera: Megachilidae). Journal of Economic Entomology
101: 1088–1094. https://doi.org/10.1093/jee/101.4.1088

Inkscape Project (2020) URL: https://inkscape.org.

IPBES (2019) Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. S. Díaz, J. Settele, E. S. Brondízio, H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.). IPBES secretariat, Bonn, Germany: 56. https://doi.org/10.5281/zenodo.3553579

JKI (2020) Tabelle der Abdrifteckwerte für Einfachanwendungen - Stand: 18. Mai 2020. Julius Kühn-Institut (JKI), Quedlinburg, Germany.

Kühne S, Freier B, Kaul P, Jüttersonke B, Schenke D, Forster R, Baier B, Moll E (2002) Auswirkung der Abdrift von Insektiziden in einem Saumbiotop. Agrarökologie, Berne, Hannover, 121 pp.

Kühne S, Enzian S, Jüttersonke B, Freier B, Forster R, Rothert H (2000) Beschaffenheit und Funktion von Saumstrukturen in der Bundesrepublik Deutschland und ihre Berücksichtigung im Zulassungsverfahren im Hinblick auf die Schonung von Nichtzielarthropoden. Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft Berlin und Braunschweig 378: 1–128. https://doi.org/10.5073/20210706-100746

de Lange HJ, Lahr J, Brouwer J h. d., Faber J h. (2012) Review of available evidence regarding the vulnerability of off-crop non-target arthropod communities in comparison to in-crop non-target arthropod communities. EFSA Supporting Publications 9: 348E. https://doi.org/10.2903/sp.efsa.2012.EN-348

Lenth R (2023) emmeans: Estimated Marginal Means, aka Least-Squares Means. Available from: https://CRAN.R-project.org/package=emmeans.

Main AR, Webb EB, Goyne KW, Mengel D (2018) Neonicotinoid insecticides negatively affect performance measures of non-target terrestrial arthropods: a meta-analysis. Ecological Applications: A Publication of the Ecological Society of America 28: 1232–1244. https://doi.org/10.1002/eap.1723

Main AR, Webb EB, Goyne KW, Mengel D (2020) Reduced species richness of native bees in field margins associated with neonicotinoid concentrations in non-target soils. Agriculture, Ecosystems & Environment 287: 106693. https://doi.org/10.1016/j.agee.2019.106693

Müller C (2018) Impacts of sublethal insecticide exposure on insects — Facts and knowledge gaps. Basic and Applied Ecology 30: 1–10. https://doi.org/10.1016/j.baae.2018.05.001

Nagy K, Duca RC, Lovas S, Creta M, Scheepers PTJ, Godderis L, Ádám B (2020) Systematic review of comparative studies assessing the toxicity of pesticide active ingredients and their product formulations. Environmental Research 181: 108926. https://doi.org/10.1016/j.envres.2019.108926

Navarro-Roldán MA, Avilla J, Bosch D, Valls J, Gemeno C (2017) Comparative Effect of Three Neurotoxic Insecticides With Different Modes of Action on Adult Males and Females of Three Tortricid Moth Pests. Journal of Economic Entomology 110: 1740–1749. https://doi.org/10.1093/jee/tox113

Pisa LW, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Downs CA, Goulson D, Kreutzweiser DP, Krupke C, Liess M, McField M, Morrissey CA, Noome DA, Settele J, Simon-Delso N, Stark JD, Van der Sluijs JP, Van Dyck H, Wiemers M (2015) Effects of neonicotinoids and fipronil on non-target invertebrates. Environmental Science and Pollution Research 22: 68–102. https://doi.org/10.1007/s11356-014-3471-x

Posit team (2023) Integrated Development Environment for R. Posit Software, PBC, Boston, MA. http://www.posit.co/. Available from: http://www.posit.co/.

R Core Team (2023) R: A Language and Environment for Statistical Computing. Foundation for Statistical Computing, Vienna, Austria. Available from: https://www.R-project.org/.

Ritz C, Baty F, Streibig JC, Gerhard D (2015) Dose-Response Analysis Using R. PLOS ONE 10: e0146021. https://doi.org/10.1371/journal.pone.0146021

Roß-Nickoll M, Fürste A, Mause R, Ottermanns R, Theißen B, Toschki A, Ratte H-T (2004) Die Arthropodenfauna von Nichtzielflächen und die Konsequenzen für die Bewertung der Auswirkungen von Pflanzenschutzmitteln auf den terrestrischen Bereich des Naturhaushaltes. Umweltbundesamt, Berlin. Forschungsbericht Available from: https://www.umweltbundesamt.de/publikationen/arthropodenfauna-von-nichtzielflaechen-konsequenzen (February 1, 2024).

Rumschlag SL, Mahon MB, Hoverman JT, Raffel TR, Carrick HJ, Hudson PJ, Rohr JR (2020) Consistent effects of pesticides on community structure and ecosystem function in freshwater systems. Nature Communications 11: 6333. https://doi.org/10.1038/s41467-020-20192-2

Sabzevari S, Hofman J (2022) A worldwide review of currently used pesticides' monitoring in agricultural soils. Science of The Total Environment 812: 152344. https://doi.org/10.1016/j.scitotenv.2021.152344

Sánchez-Bayo F (2012) Insecticides Mode of Action in Relation to Their Toxicity to Non-Target Organisms. Journal of Environmental & Analytical Toxicology S4. https://doi.org/10.4172/2161-0525.S4-002

SAS Institute Inc. (2023) JMP®Pro. Cary, NC, 1989–2023.

Schmitz J, Hahn M, Brühl CA (2014) Agrochemicals in field margins – An experimental field study to assess the impacts of pesticides and fertilizers on a natural plant community. Agriculture, Ecosystems & Environment 193: 60–69. https://doi.org/10.1016/j.agee.2014.04.025

Siviter H, Muth F (2020) Do novel insecticides pose a threat to beneficial insects? Proceedings of the Royal Society B: Biological Sciences 287: 20201265. https://doi.org/10.1098/rspb.2020.1265

Strauss G, Simon H (2021) Miridae. Available from: https://www.corisa.de/index.html (January 16, 2024).

Teder T, Knapp M (2019) Sublethal effects enhance detrimental impact of insecticides on non-target organisms: A quantitative synthesis in parasitoids. Chemosphere 214: 371–378. https://doi.org/10.1016/j.chemosphere.2018.09.132

Wachmann E, Melber A, Deckert J (2004) Wanzen 2. Goecke & Evers, Keltern.

Wagner DL, Grames EM, Forister ML, Berenbaum MR, Stopak D (2021) Insect decline in
the Anthropocene: Death by a thousand cuts. Proceedings of the National Academy
of Sciences 118: e2023989118. https://doi.org/10.1073/pnas.2023989118

Yu SJ (2015) The Toxicology and Biochemistry of Insecticides. 2nd ed. CRC Press, New
York, 380 pp.

van de Zande J, Rautmann D, Holterman H, Huijsmans JFM (2015) Joined spray drift curves
for boom sprayers in The Netherlands and Germany. Plant Research International,
part of Wageningen UR.

Acknowledgements

We would like to thank Sarah Rißmann for her constant support throughout this project and
Nina Estler for her help with the dose-response assays. Further, we would like to thank Denise
Kuhn, Anja Betz, Robert Bischoff and Prayan Pokharel for their support and critical feedback.
Special thanks go to Manfred Konert for his help in setting up and maintaining the field
experiment. In addition, we would like to thank the team at the Botanical Garden Hohenheim,
led by Dr Helmut Dalitz, for their support in carrying out the field trial. We are also grateful to
Jürgen Deckert, who advised us on the subject of soft bug handling, and to Gerhard Strauss
for the permission to use insect images from the Corisa software. Finally, we would like to
thank the Graduate Academy of the University of Hohenheim, which made this project possible
by providing a scholarship as part of the Baden-Württemberg Graduate Support Programme.

904
Contributions

CA.B., I.G. and G.P. supervised the project. JE.S., CA.B., I.G. and G.P. drafted the
experiments. JE.S. and P.B. conducted the field experiment. JE.S. performed the greenhouse
experiment. JE.S. performed the dose-response assays. JE.S. performed the analysis of the
field- and greenhouse experiments. JE.S., D.G and HP.P. performed the analysis of the dose-
response assays. B.H., F.W. and JE.S. conducted the pesticide analysis. JE.S. wrote the
original draft of the manuscript. G.P., CA.B., I.G., HP.P. edited the manuscript.