INSTRUCTION ALTERS THE INFLUENCE OF ALLOCENTRIC LANDMARKS IN A REACH TASK

Lina Musa[a,b], Xiaogang Yan[a], J. Douglas Crawford[a,b,c]

[a] *Centre for Vision Research and Vision: Science to Applications (VISTA) Program, York University, ON, Toronto, Canada*

[b] *Department of Psychology, York University, ON, Toronto, Canada*

[c] *Departments of Biology and Kinesiology & Health Sciences, York University, ON, Toronto, Canada*

Abstract

Allocentric landmarks have an implicit influence on aiming movements, but it is not clear how an explicit

instruction (to aim relative to a landmark) influences reach accuracy and precision. Here, 12 participants

performed a task with two instruction conditions (*egocentric* vs. *allocentric*), but with similar sensory and

motor conditions. Participants fixated gaze near the centre of a display aligned with their right shoulder

while a target stimulus briefly appeared alongside a visual landmark in one visual field. After a brief

mask/memory delay the landmark then re-appeared at a different location (same or opposite visual field),

creating an ego/allocentric conflict. In the *egocentric* condition, participants were instructed to ignore the

landmark and point towards the remembered location of the target. In the *allocentric* condition,

participants were instructed to remember initial target location relative to the landmark and then reach

relative to the shifted landmark (same/opposite visual field). To equalize motor execution between tasks,

participants were instructed to anti-point (point to the visual field opposite to the remembered target) on

50% of the egocentric trials. Participants were more accurate, precise, and quicker to react in the

allocentric condition, especially when pointing to the opposite field. We also observed a visual field effect,

where performance was worse overall in the right visual field. These results suggest that when egocentric

and allocentric cues conflict, explicit use of the visual landmark provides better reach performance than

reliance on noisy egocentric signals. Such instructions might aid rehabilitation when the egocentric system

is compromised by disease or injury.

Keywords: **Spatial Representation, Allocentric, Egocentric, Memory-guided reaching**

Highlights:

• 12 participants reached to remembered targets in the presence of a visual landmark

• Participants were instructed to ignore, or point relative to, the landmark

• The landmark instruction improved reaction time, precision, and accuracy

• These effects were stronger when pointing was cued toward the opposite visual field

• Knowledge of these rules might be used to enhance performance or in rehabilitation

1. Introduction

Humans use specific spatial reference frames to retain remembered visual information for a goal-directed

action (Soechting & Flanders, 1992; Crawford et al., 2011). The visual system is thought to utilize two

types of spatial reference frames, observer-centered, egocentric reference frames versus world-fixed,

allocentric reference frames, often anchored relative to reliable landmarks (Byrne et al., 2010; Howard

& Templeton, 1996; Vogeley & Fink, 2003). In ordinary circumstances, the brain integrates information

from these two reference frames in Bayesian manner (Byrne & Crawford, 2010; Fiehler et al., 2014).

However, certain circumstances may require one to ignore surrounding landmarks (i.e., when they are

unstable or irrelevant to the task) or focus strongly on surrounding landmarks (such as a workspace fixed

to a moving based). Various experiments have tapped into these mechanisms by instructing participants

to use one reference frame over the other (Bryne & Crawford, 2010; Lemay et al., 2004). The question

thus arises, which of these of these instructions leads to better performance?

Various studies have explored the implicit influence of visual landmarks on goal-directed actions such

as saccades, reaches, or pointing towards a seen or remembered target. Previous findings have shown

that reach targets can be remembered reasonably well in the absence of visual landmarks (Lemay &

Stelmach, 2005; McIntyre et al., 1997; Vindras & Viviani, 1998) with certain stereotypical errors such

as gaze-centered overshoots (Bock, 1986; Henriques et al., 1998). However, the addition of a visual

landmark can influence reaching, reducing both constant and variable errors (Byrne et al., 2010;

Krigolson & Heath, 2004; Lemay et al. 2004; Redon & Hay, 2005). The stabilizing influence of a

landmark was particularly prominent in task that involved remapping the reach target to the opposite

visual hemifield, where one would expect egocentric signals to be less stable (Byrne et al., 2010).

Conversely, the landmark had less stabilizing influence on behavior when they were shifted and rotated

relative to the reach goal (Thaler & Todd 2009). Finally, the addition of visual landmarks can negate the

accumulation of reach errors after prolonged memory delays in the dark (Chen et al., 2011).

Normally, egocentric and allocentric these cues agree, but they can also conflict, either in tasks that

introduce egocentric noise or when the visual environment is unstable (Byrne & Crawford 2010; Byrne

et al. 2010, Chen et al. 2011). The latter has been replicated experimentally in cue-conflict tasks where

the landmark is surreptitiously shifted relative to egocentric coordinates during a memory delay (Byrne

& Crawford 2010). In this situation, ego / allocentric cues appear to be integrated optimally, based on

their relative reliability (Byrne & Crawford, 2010). Usually, more weight is placed on egocentric

coordinates, such that the movement shifts approximately 1/3 in the direction of the landmark shift

(Byrne & Crawford 2010; Fiehler et al. 2014; Li et al. 2017). However, the specific weighting depends

on task details. For example, Byrne & Crawford (2010) found that participants rely more on a landmark

when it is perceived to be stable, or when gaze position was less stable. Further, in simulated naturalistic

settings, landmarks had more influence when they were task-relevant, when more than one landmark was

shifted in the same direction, and when the landmark was closer to the target (Fiehler et al., 2014). Thus,

in the absence of explicit instructions, the visual system uses implicit algorithms to determining how to

weight ego / allocentric cues. Recent physiological studies suggest that this implicit integration may

occur in frontal cortex (Bharmauria et al. 2020, 2021).

Alternatively, people can be instructed to ignore or put full weight on a visual landmark, as in common

driving instructions like 'ignore the first stop sign and turn right at the second'. Likewise, experimental

participants can be instructed to either ignore a landmark or reach at a to a fixed egocentric location

relative to the landmark. Such instructions were used in neuropsychological and neuroimaging

experiment that suggesting involvement of the ventral visual stream in allocentric representations and

dorsal stream in egocentric transformations (Goodale et al., 2004; Schenk, 2006; Chen et al. 2014, 2018).

However, the influence of these instructions on behavior is less clear. For example, in their control tasks

Byrne & Crawford (2010) found no overall difference in performance when participants were explicitly

instructed to use egocentric or allocentric cues, but there was no cue-conflict in these tasks and the visual

stimuli were different. Thus, it remains unclear what affect the instruction itself (to prioritize egocentric

vs. allocentric cues) has on behavior, particularly when these two cues conflict.

Here, we tested the influence of instruction (to use or not use a landmark) in a cue-conflict, memory-guided

reach task where the visual stimuli were equal and balanced across tasks,. In this way we could isolate and

directly compare how top-down reliance on allocentric vs. egocentric cues affects reach behavior. It has

been suggested that landmark-centered encoding is a less noisy process and more stable over a time delay

(Byrne et al., 2010; Chen et al., 2011). Thus, we predicted that instruction to attend to and use the

landmark for spatial coding would increase weighting on the more stable code, and therefore improve

performance, especially when egocentric noise is higher (Byrne & Crawford 2010, Chen et al. 2011). To

simulate the latter case, we included an 'anti-reach' condition where participants had to reach to the

opposite visual field as the viewed target. We found that 1) reaching was less variable and more accurate in

the allocentric instruction tasks than egocentric instruction tasks (especially in the right visual field) and

2) the beneficial effect of allocentric encoding is more pronounced when participants were required to to

respond in the visual field opposite to stimulus encoding.

90   **2. Material and Methods**

2.1 Participants

13 individuals (7 males and 7 females; ages 20 – 33) provided informed consent to participate in this study.

All participants were right-handed with no neuromuscular or uncorrected visual deficits, based on self-

report. Data from 1 participant was excluded from further analysis because they did not meet the inclusion

criteria described below, leaving 12 participants' data for analysis. This met the sample size required for

sufficient power (see §1.5. *Sample Size Analysis*). All participants were naïve to the purpose of the

experiment and were given monetary compensation for their time. The experimental procedures were

approved by the York Human Participants Review Subcommittee.

2.2 Apparatus

The experiment was conducted in darkness and participants wore a black glove to avoid any ambient reflections off the right hand. Participants sat behind a table, on a chair with an adjustable height (Fig. 1). The head was stabilized on a personalized bite bar made of a dental impression compound (Kerr Corporation, Orange, CA). Participants' right hand was positioned on a button box on the table-top, directly in front of them. The button box was used to control the pace of the experiment and was the designated start position for reaching. A customized ring with a 3x3 array of Infrared-emitting diodes (IREDS) was attached to participants' right index finger and their 3D-position was continuously recorded by two OptoTrack 3020 tracking systems (Northern Digital, Waterloo, Ontario, Canada). Gaze direction was monitored from only the right eye, through the EyeLink II infrared eye-tracking system, (SR Research, Mississauga, Ontario, Canada), that was mounted to the bite bar stand. The stimulus display (described below) was presented 50 cm ahead of the eye and 152 mm to the right / 91 mm bellow the centre of the bite bar stand, approximately aligned with the right shoulder bone (acromion). The height of the chair was adjusted so that the acromion was aligned with the centre of the display panel. This arrangement ensured that stimuli were centred in the right arms mechanical range and within easy reach. During reaches participants' head was facing forward, but gaze was fixated the centre of this display, such that shoulder and central visual coordinates aligned. Audio instructions were delivered from a speaker. Two 40-watt desk lamps, placed on either side of the desk, were turned on every 10 trials (2 min), to eliminate potential dark adaptation. The experimental set-up is shown in Fig. 1.

Fig. 1: Experimental set-up. From left to right and top to bottom: OptoTrack 3020 tracking systems (Northern Digital, Waterloo, Ontario, Canada), on either side of the room (second one not shown in the figure) were used to track the finger motion. Eyelink II cameras (SR Research, Mississauga, Ontario, Canada) were dismounted from the headset and installed on the bite-bar stand. The left camera is occluded in this view of the set-up. Home button box (yellow in the figure), on the desk immediately in front of participants was used to control the pace of the experiment. Participants' height was adjusted relative to the fixed bite-bar height using a metallic screw on a rigid chair with an adjustable height. Two 40W dark adaptation desk lamps illuminated the dark room during breaks and every 3 trials. A custom made 307 mm x 161 mm wooden panel, fitted with light emitting diodes (LEDs) was used to display stimuli. Audio instructions were played from two desktop speakers, one of the speakers is not shown in the figure. The finger pointing device was a customized ring with a 3x3 array of Infrared-emitting diodes (IREDS) that continuously relayed signal to the OptoTrack 3020 tracking systems.

2.3 Calibration

Before each experiment, participants completed a set of calibration procedures. Eye position calibration was

done through sequential fixation of a 5 light emitting diodes (LEDs) including the centre position and the

four corners of the display panel (See Fig. 2A). They then completed two OptoTrak calibration sessions.

Finger-tip position was calibrated by having participants point with a pre-calibrated 'cross' of IREDs on a

rigid body fixed to the right index finger. Further calibration was done by having participants successively

point to the 4 corner positions in the LED display. IRED position data in the OptoTrak intrinsic coordinate

system was compared offline with known positions of the calibration dot positions to create a linear mapping

between IRED positions and the screen coordinates. This procedure allowed for the conversion of recorded

finger-tip position into screen and then visual coordinates, as described below.

2.4 Visual Stimuli and Basic Task

The experimental paradigm was based on a simpler paradigm used previously in an fMRI experiment (Chen

et al., 2014). The task involved touching the remembered location of a transient visual target (that always

appeared simultaneously with a visual landmark) as accurately as possible. Fig. 2A shows the visual/touch

display. LEDs (1.15° light emitting diodes; LEDs) were fixed to a wooden panel (307 mm x 161 mm).

Target, landmark and fixation LEDs were placed at 1 cm (~1.15 visual degrees visual angle) intervals along

this panel. Seven white gaze fixation LEDs were placed at centre and 1.15, 2.29, 3.43 degrees to the left and

right. 18 green reach target LEDs were placed in the periphery, starting at a 4.57 degree distance from the

centre. 6 red landmark LEDs were positioned 7.96, 9.09 or 11.31 degrees, to the right or left of the centre.

40 LEDs used as a visual mask were placed 1.15 degrees above and below the target/landmark stimuli, in-

between the target/landmark LED positions.

The same stimulus paradigm was used in all experiments, with LED positions and task instructions

randomized as described below. The sequence of stimulus events is shown in Fig. 2B. Participants started

seated comfortably with their hand placed on the button box near their chest. At the beginning of each trial,

they received an audio instruction from the speakers telling them how to reach (see next section for details).

Participants were then required to visually fixate one of the 6 non-centre white LEDs. After 2 seconds a

green target LED was displayed to the right or the left of the centre. This was accompanied simultaneously

by presentation of one of 6 red landmark LEDs relative to the target, on the same side of the centre as the

target. After 2 seconds, both the target and landmark stimuli disappeared and the gaze fixation point moved

to centre, allowing 300ms for a saccade to that position. This was done so that the fixation LED could not

be used itself as a reliable allocentric landmark (Chen et al. 2014). The array of white 'mask' LEDs was then

illuminated for 150ms to negate visual afterimages and focus memory resources (Medendorp et al. 2003).

After the mask, the centre fixation LED reappeared, requiring continued central fixation for the rest of the

trial. A landmark then reappeared for 2 s, but always at a different location than it was initially viewed,

creating a conflict between the location of the target in egocentric coordinates versus allocentric coordinates.

At this point participants were instructed to reach via the speakers, and then were given 4 seconds to touch

the stimulus panel with the right index finger as accurately as possible, based on the instruction they had

received at the start of the trial. They then returned their hand to starting position and pressed the button

when they were ready for another trial.

Fig. 2: Experimental Stimuli and Paradigm. **(A)** Stimuli. The stimuli were displayed in a horizontal array. Left side and center of the array are shown, while the right side is the same as the left side (mirror image. The central fixation, off-centre fixation locations and eye calibration points were displayed by white LEDs. The green LED target was randomly displayed in of the 18 (9 L, 9 R) LEDs. The position of the first target LED was 3 circles (4.57°) from screen centre. The red landmark simultaneously appeared on the same side of the screen as the target, displayed by one of the 3 LEDs in the middle of the 9 target LEDs (half red, half green circles in the figure), while the second red landmark randomly appeared in one of the remaining landmark positions. **(B)** Paradigm. The order of a typical trial is shown in the figure. Each trial began with an audio instruction, where participants were instructed to remember the spatial location of the target, or the position of the target relative to the landmark. The response audio depended on task; *EGO* trials were followed by "Target", instructing participants to point toward the target, or "Opposite", instructing participants to point to the mirror opposite side, while all *ALLO* trials were simply followed by "Reach", instructing participants to point to the remembered target position relative to the second landmark.

2.5 Task Instruction and Conditions

As noted above, the visual paradigms used in this task were always the same for different task conditions,

what differed between conditions was the task instruction delivered via the speakers at the start of the trial. In

the *Allocentric (ALLO) Condition,* participants were instructed to "*Reach relative to cue*", i.e., they were

required to remember the position of the green target, relative to the red landmark. Since the landmark was

shifted to a new position at this time, egocentric coordinates provided no useful cues for this task. In this case,

the instruction to reach at the end was simply '*reach*'. In contrast, in the *Egocentric (EGO) Condition*,

participants heard "*Reach to target"* at the beginning of the task. In this condition, participants were instructed

to ignore the landmark (which in this condition provides invalid information). Henceforth, we refer to these

as the *EGO Instruction* and the *ALLO Instruction* conditions.

To further challenge the system, in half the trials participants were required to perform 'anti-reaches' to the

visual field opposite to the field where they saw the target (Gail & Andersen, 2006 Cappadocia et al. 2018).

In the *ALLO Condition*, this was done by shifting the landmark to the opposite hemifield. In the *EGO*

condition, an additional instruction was provided. When participates heard '*target*', they reached toward the

original location of the target ('*PRO-Task*'). When participants heard '*opposite*' they were expected to reach

toward the mirror opposite position, relative to screen midline ('*ANTI Task'*). To be consistent with the

literature, we then divided our data into *EGO PRO/ANTI -* or *ALLO PRO/ANTI -reach* trials (based on the

final reach direction relative to initial target). Finally, in both *PRO* and *ANTI* trials, stimuli were arranged so

EGO/ALLO goals covered the same range.

Experimental Design

Before each experiment, participants engaged in a practice session to ensure that they understood the

instructions and were able to perform each element of the task correctly. This consisted of 3 practice blocks

of 10 trials. The first two practice blocks were made up of each *EGO/ALLO Instruction* condition separately

and the last practice block was randomly interleaved. After they were successful and clear on the instructions,

they proceeded to the actual experiment. Actual experiments were divided into 3 blocks of 72 trials.

Participants had a 5 min rest period between blocks and four 2 min breaks within each block (with room

lights fully illuminated). Individual trials lasted 12.05 s in both instruction conditions, but the experiment

was self-paced, subsequent trials were initiated through a button press. On average, participants took 21.65

mins to complete an entire block. The order of instructions, target locations, and landmark locations was

pseudorandomized beforehand so that they were unpredictable for the participants but provided an equal

dataset with distributed stimuli for each experimental condition.

188   2.6 Data Analysis

All data obtained from OptoTrak and EyeLink was analysed offline using a custom software written in

MatLab R2019a (Math Works). A program was written to generate a mapping between the Optotrak

coordinates and the position of the fingers in screen coordinates of the fingertip. This was done by utilizing

the data exported from the screen calibration session and known screen coordinates of the 4 calibration points

on the screen corners. This mapping was then utilized to obtain reach endpoints in screen coordinates for

analysis. Finally, target and touch positions in screen coordinates were converted into visual angle (relative

to the central fixation point) using geometric measures obtained from the laboratory set-up.

196 *2.7.1 Exclusion Criteria*

Movement kinematics were inspected to ascertain that participants followed instructions and a program was

written to automatically average eye fixation, from the period of the second landmark to the go signal. Trials

were excluded if eye variation was greater than 2 degree in the horizontal direction for more than 20% of

time. Trials with brief saccades to remembered target location that then returned were not excluded, because

this does not appear to influence pointing accuracy (Van Pelt & Medendorp 2007). However, trials were

participants looked at the landmark were excluded. Finally, trials that included anticipatory reaches (reaches

initiated before the audio instructions) were excluded from further analysis.

Overall, 12 participants who had at least 175 viable trials (81.02%) were included in the study. Participants

completed on average 72.90 ± 20.58 *EGO Instruction* trials, on average 37.80 ± 8.98 of trials were trials in

which the target was viewed, and the movement was executed in the same visual field while 36.70 ± 10.60

were the movement was executed in the opposite visual field. On average, participants also completed 70.60

± 18.53 *ALLO Instruction* trials, 32.70 ± 7.81 of trials were on the same visual field and 35.70 ± 12.99 of

trials were in the opposite visual field.

2.7.2 Kinematic Analysis

Using a customized MatLab program, movement start was marked at 20% maximum resultant velocity

(Vmax), while movement end was marked at 8% of Vmax. Reach endpoint was found by averaging 30% of

points at 8% Vmax and at 1.15 times the minimum finger to screen distance (distance in the z direction). The

screen relative coordinates of reach endpoints in in the horizontal (x) and vertical (y) dimension were used

to compute the variable error, a measure of the distance of reach endpoints from the mean final position. R

4.1.2 (Core Team, 2021) was used to create 95% confidence ellipse of the scatter of reach endpoints. The

area of the ellipse was found by first computing eigenvalues ($\sigma 1$, $\sigma 2$) of the covariance matrix of reach

endpoints. The eigenvalues were then used to derive half the lengths of the of the semi-major (principal axis)

and semi-minor axes (orthogonal to the principal axis) of the 95% confidence ellipse and used to compute

the ellipse area (formula [1] shown below). The ellipse areas were then used to compare the *EGO* and *ALLO*

Instruction data. Mean ellipses for each target were constructed by averaging the covariance matrices of the

corresponding ellipses of individual participants.

Ellipse area $= \pi *$ sqrt(5.991$*$ $\sigma 1$) $*$ sqrt(5.991$*$ $\sigma 2$)   [1]

The overshoot error in horizontal reach endpoints were used to analyse response accuracy. The horizontal

overshoot error in reaching response at each target location was found by subtracting the difference between

the reach endpoint to screen midpoint distance, and the expected target location to screen midpoint distance.

The expected target location was the same as the initial location for *EGO/PRO* reaches, mirror opposite to

the target location for *EGO/ANTI* reaches, but based on the shifted landmark for all allocentric reaches. The

mean overshoot error was found by averaging the magnitude and direction of the reaching error (positive to

the right, negative to the left), separately for each instruction condition (*EGO/ALLO*), task condition

(*PRO/ANTI*), and visual field of response. Ellipse areas, reaction times and movement times were averaged

in a similar manner. The reaction time was the period between go signal and movement start, while the

movement time was the period between the movement start and movement end.

2.7 Sample Size Analysis

Sample size was calculated based on the simulation approach described by Green & MacLeod, (2016), using

the simr R package (Core Team, 2021), which calculates the power for generalized linear mixed models

from the lme4 R package (Core Team, 2021). The power calculations were based on Monte Carlo

simulations. Reaching variance was obtained from a pilot study of 2 individuals and 72 observations for each

(*EGO/ALLO*) instruction condition. A mixed effects model was fitted on pilot data, to obtain an estimated

effect size of 0.267 for reaching variance. Using the obtained effect size and the simulation package, the

sample size was gradually increased to achieve a power of 85%. The sample size required to achieve this

power was 12.

2.8 Statistical Analysis

Linear mixed effects models were used to analyse differences in ellipse areas, overshoot errors, reaction

times and movement times, using the lme4 R package (Core Team, 2021). The reaching responses relative

to the expected target positions were fit with quadratic mixed effects models, using lme4 R package (Core

Team, 2021), to determine differences in reaching behaviour due to the task conditions. Interaction contrasts

were performed using the eemeans R package (Core Team, 2021).

For statistical analysis we grouped our data into 4 task conditions, as follows:

1. EGO Instruction / PRO Task

2. EGO Instruction / ANTI Task

3. ALLO Instruction / PRO Task

4. ALLO Instruction / ANTI Task

In addition, based on the findings of Byrne et al. (2010), we also sorted these data into hemifields (*LEFT /*

RIGHT), based on the final direction of the instructed reach goals.

3. Results

Our experiment was designed to test if an explicit instruction to reach relative to an allocentric visual cue (as

opposed to ignoring this cue and relying on egocentric coordinates) would alter reach performance in the

presence of otherwise identical stimuli. For example, the top panel (A) of Figure 3 represents the case where

both the target and landmark were presented in the right visual field, and after the target disappeared the

landmark shifted to the left visual field. The bottom two panels of Figure 3 (B) each display 6 representative

hand trajectories (black lines) and gaze trajectories (grey lines) associated with the stimuli in the top panel.

Both panels show *ANTI-Task* trials but in (B1) the participant was given the egocentric 'opposite' instruction

whereas in the other case (B2) the participant was instructed to point relative to the (shifted) landmark. As

one can see, the participant was able to perform both tasks, although with subtle differences. In total, 12 (of

13) participants completed a sufficient number of these trials in all versions of our task (including the easier

PRO-task versions) for statistical analysis of their data. An analysis focusing on comparisons of precision,

accuracy, and reaction time between conditions will follow.

Fig. 3: Typical Eye and Finger Trajectories. **(A)** Spatial location of example stimuli (see graphic key for details). **(B1)** *EGO / ANTI condition*, where the participant was instructed to touch mirror opposite spatial location of the remembered target position. The same color conventions as A are used, except the unfilled green circle shows the ideal goal position, the black lines show the 2D finger trajectory for 6 example trials, and the grey lines show the corresponding 2D gaze locations during central fixation, where gaze was required to remain within +/- 2° of the fixation point 80% of the time. **(B2)** *ALLO / ANTI condition*, where the landmark appeared in the opposite side of the fixation (same graphic conventions as B1).

3.1 Reach Accuracy and precision: General Observations

Fig. 4 provides a spatial overview of our pointing results in all four conditions, plotted in visual coordinates.

The top portion of each panel shows the labeled 2-D target positions (black dots), corresponding reach data

points (medium circles), and their 95% confidence ellipses (large ovals), using the same color code for each

f(light shade near centre, darker toward periphery). The lower parts of each panel show the corresponding

1-D probability densities for each target position along the horizontal dimension (the dimension in which

targets varied). The leftward panels (A & B) exhibit the plots for a single participant, while the rightward

panels (C & D) portray the mean ellipses and probability densities across participants. The plots pertaining

to the *EGO Instruction* (blue) are showcased in blue on the top panels (A & C), whereas those concerning

the *ALLO Instruction* (red) are presented in red on the bottom panels (B & D).

Some qualitative observations are that the distributions of error are that 1) the distribution of error was

relatively large, as one might expect in a memory guided task, 2) the vertical distributions tended to be larger

than the horizontal distribution, and 3), the distributions did not overlap, but rather shifted with target

location, and 4) the *ALLO Instruction* distributions seem slightly larger in the *PRO-Task* conditions. Fig 4

right column, panels (E & F), illustrate the average ellipses for the *ANTI -task* data using the same

conventions as the middle column of Fig. 4. Here, the *EGO* distributions were larger still, and thus the

difference between the *EGO* and *ALLO* distributions seem clearer, at least in the left visual field, where the

ALLO distributions are visibly smaller than the *EGO* distributions. Each of these observations will be

quantified in more detail below.

Fig. 4: Reach Endpoint Variability. Panels for a representative participant **(A)** & **(B):** 95% confidence ellipses (2D reach endpoint variability) are shown at the top of each panel and horizontal reach endpoints probability densities (1D reach endpoint variability) are shown at the bottom of each panel. The small, filled circles are reach individual endpoints relative to the central fixation. Ellipses, probability density curves and reach endpoints are all color coded by initial target spatial location. The color gradient, from light to dark, represents targets that are central to peripheral. Panels **(C)**, **(D)**, **(E)** & **(F):** Mean ellipses, mean horizontal probability densities and mean reach endpoints of 12 participants. Panels **(A)** & **(C):** *PRO-Task, EGO Instruction* (top) in blue and panels **(B)** & **(D):** *PRO-Task, ALLO Instruction* (bottom) in red. **(E):** *ANTI-Task, EGO Instruction* (top) in blue and panels. **(F):** *ANTI-Task, ALLO Instruction* (top) in red and panels. The target positions are referenced using black dots in the ellipse figures (the exact distances are labelled on the plot) and as axis ticks on the x-axis in the 1-D density plot, both target position labels are color-coded using the same color gradient.

3.2 Quantification of Reaching Variance

Fig. 5 quantifies illustrates the distributions of the ellipse fit areas (averaged within each participant) for the

left and right visual field targets, with *EGO Instruction* data depicted in blue and *ALLO* in red. The *PRO*

Task data are depicted in the top panel (A), while the *ANTI task* data are depicted in the bottom panel (B).

As one can see, the general trend is for smaller ellipses fits (i.e., higher precision) in *ALLO instruction* data.

To quantify these data, we performed a statistical analysis of the instruction influence (*EGO/ALLO*), task

(*PRO/ANTI*) and visual field (*LEFT/RIGHT*) ellipse fits using a generalized linear mixed model. The

analysis included the interaction of spatial instructions with both task and visual field of response. No

significant three-way interaction effect was found, leading to the fitting of a simpler model (Table 1).

The outcome of this analysis revealed a significant effect of spatial instruction, with ellipse areas in the *EGO*

Instruction condition (mean $29.87 \pm$ SD $4.66^{\circ 2}$) being significantly larger i.e., less precise, than the *ALLO*

Instruction condition ($20.32 \pm 3.77^{\circ 2}$) (p = 2.0 e -15). The *ALLO Instruction* led to improved precision

consistently across visual fields of response, especially when participants had to point opposite to the visual

field of response. Post-hoc interaction contrasts, averaged across visual fields, showed that participants were

significantly less precise in the *EGO* than the *ALLO Instruction* condition, especially when they had to point

opposite to the visual field of response (p = 1.0 e -5).

Plot Legend

Fig. 5: Reach Endpoint Ellipse Areas. Violin Plots of ellipse areas averaged over right/left visual field targets, and for each participant. The legend on top explains the conventions used to generate these plots. Each plot (A/B) summarises the observed trends in the left visual field (left side) and right visual field (right) for each task/instruction condition. **(A)** Mean areas of 95% confidence ellipses for the *PRO-Task* data. **(B)** Mean areas of 95% confidence ellipses for the *ANTI Task* data. *EGO* and *ALLO Instruction* conditions are shown in the figure as blue and red, respectively. Significant differences are shown by an asterisk, $*p < 0.05$, $**p < 0.01$.

Table 1

Fixed Effects of Multi-level Model using Ellipse Areas as the Criterion

Predictor	B	B 95% CI [LL, UL]	t value	p
(Intercept)	16.67**	[9.70, 23.61]	4.77	8.5 e -6**
Task (*PRO*)	1.55	[-6.62, 9.71]	0.377	0.71
Instruction (*EGO*)	-5.65 **	[-6.37, -4.92]	-11.57	2.0 e -15**
Visual Field (RIGHT)	4.60	[-3.56, 12.75]	1.12	0.26
Task (*PRO*) x Instruction (*EGO*)	-1.65**	[-2.37, -0.92]	-4.44	1.0 e -5**
Visual Field (RIGHT) x Instruction (*EGO*)	2.21	[-9.33, 13.74]	0.38	0.71

Note. B represents standardized regression weights. *LL* and *UL* indicate the lower and upper limits of the confidence interval, respectively. Model Adjusted R^2 = 0.145* , 95% CI[0.00,0.24]
* indicates p < 0.05. ** indicates p < .01.

3.2 Reaching Accuracy

As noted above, the horizontal positions of the response distributions tended to vary with corresponding desired target locations (Fig. 4), suggesting that participants did not simply point to the left or right visual field. To quantify the accuracy of these responses, the absolute horizontal values of participants' end points (averaged across trials for each target within each participant) were regressed relative to the corresponding ideal target locations (Fig. 6). A general observation is that the distribution of the *ALLO Instruction* data (red) appears to be more compact and follows the line of unity (ideal accuracy) more closely that the *EGO Instruction* data (blue).

To select the most appropriate model to quantify these data, we employed a k-fold (leave-one-out) cross-

validation, comparing the fit of three models: a linear regression model, a quadratic regression model centered

at the first target, and a model with nominal predictors. The quadratic model exhibited a significantly better

fit, as indicated by a lower root-mean-squared-error (RMSE) and a higher R-squared value. These fits are

shown as curved lines, color coded lines in Fig. 6.

Table 2 summarizes the relative contributions (β) of the model parameters, indicating those which were

significant (*), and their 95% confidence intervals (CI). Overall, this fit yielded a significant ($p < 0.01$) R^2 of

0.534 [0.48, 0.57]. The relationship between participants' reach endpoints and the targets demonstrated a

significant negative quadratic trend ($P = 0.02$), suggesting a decrease in slope for more peripheral targets,

possibly due to decreased perceptual acuity of peripheral targets (see 'Overshoot Errors' below). This effect

was significantly more pronounced in the *EGO Instruction* Condition ($p = 4.0$ e-3). In the *EGO instruction*

condition, participants exhibited a significant decrease in the slope between reach endpoints and target

location, indicating more accuracy in the *ALLO Instruction* condition. Reaching trajectory was not

significantly influenced by the task (*PRO/ANTI*), but a significant visual field effect was observed ($P = 8.9$ e-

3), where participants exhibited a more negative quadratic trend and an increase in the tangential linear slope

at the 4 degrees target. This suggests that participants' reaches in the right visual field were relatively accurate

for more central targets but demonstrated poor reaching performance for peripheral targets.

Fig. 6 Scatterplots of the Horizontal component of Reach Endpoints vs Expected Target Positions. See graphic key (upper right) for details. The scatter plots of participants' horizontal reach endpoints in the right visual field (right panel) and left visual field (left panel) were fitted with a quadratic line (solid coloured line). **(A)** In the *PRO Task* data. **(B)** In the *ANTI-Task* data. The *EGO* and *ALLO Instruction* conditions are shown in blue and red, respectively. The dashed colored lines are the LOESS lines. Black dashed line in figures is the line of unity. Shaded grey area shows the standard error of the estimate.

Table 2

Fixed Effects of Multi-level Quadratic Model using Reach Endpoints as the Criterion

Predictor	B	B 95% CI [LL, UL]	t value	p
(Intercept)	1.34*	[0.04, 2.66]	2.02	0.04
Targets	1.27**	[0.88, 1.66]	6.41	2.7 e-10
Targets^2	-0.04*	[-0.06, -0.01]	-2.43	0.02
Targets x *PRO task*	0.02	[-0.11, 0.15]	0.034	0.737
Targets^2 x *PRO task*	0.00	[-0.01, 0.02]	0.03	0.97
Targets x *EGO Instruction*	-0.20**	[-0.33, -0.06]	-2.88	4.0 e-3
Targets^2 x *EGO Instruction*	0.02*	[0.00, 0.03]	2.17	0.03
Targets x *RIGHT* Visual Field	0.29**	[0.07, 0.51]	2.62	8.9 e-3
Targets^2 x *RIGHT* Visual Field	-0.05*	[-0.08, -0.01]	.00	0.01

Note. B represents standardized regression weights. *LL* and *UL* indicate the lower and upper limits of the confidence interval, respectively. Model Adjusted R^2 = 0.53 * , 95% CI[0.48,0.57]
* indicates p < 0.05. ** indicates p < .01.

3.4 Influence of Landmark Proximity and Shift Distance on Reach Endpoints

As described above, reach precision and accuracy were influenced by the EGO / ALLO Instruction, but was there an influence of specific stimulus parameters. Specifically, the initial distance between the landmark and target (1-3°), and the amplitude of the landmark shift (1-3° for the *PRO-Task*, 9-14° for the *ANTI-task*). If participants were able to suppress landmark influence after the *EGO-Instruction*, one would expect these parameters to have no effect, whereas they might have an influence after the *ALLO-Instruction*. To test this, two mixed-effects models were employed separately on *EGO-Instruction* and *ALLO-Instruction* conditions to assess the impact of landmark proximity and shift on accuracy (OE) and precision (ellipse areas), as well as their interaction with instruction (*EGO/ALLO*) and visual field of response (Table 3). To treat the PRO and ANTI-task data on a similar footing, we used separate predictors for these tasks.

In summary, target-landmark distance and landmark shift amplitude had no significant influence on either reach precision or accuracy in the *EGO-Instruction* condition. In the *ALLO Condition,* target-landmark distance had a significant influence on both precision (p = 7.9 e-10) and accuracy (p = 4.1 e-3), when separate shift amplitudes are included in the model. Specifically, performance was improved at the largest (3°) distance from the target (or its mirror opposite). Finally, an interaction with Task was also observed (p = 0.03) , where the *ANTI Task* showed a stronger association between the degree of landmark shift and precision, compared to the *PRO Task*.

Table 3

A: Fixed Effects of Multi-level Model using Ellipse Areas in ALLO Instruction conditions as the Criterion

Predictor	B	B 95% CI [LL, UL]	t value	p
(Intercept)	5.79**	[5.49, 6.08]	39.33	< 2.0 e -16**
Landmark Shift	1.11	[-0.10, 1.44]	1.07	0.29
Target-Landmark Distance	-3.88**	[-4.40, -3.69]	-6.23	7.9 e -10**
Landmark Shift x Task (*PRO*)	2.29*	[0.19, 0.36]	2.19	0.03*
Target-Landmark Distance x Task (*PRO*)	0.68	[0.30, 1.12]	1.07	0.29

B: Fixed Effects of Multi-level Model using Absolute Overshoot Error in ALLO Instruction conditions as the Criterion

Predictor	B	B 95% CI [LL, UL]	t value	p
(Intercept)	1.25**	[1.75, 2.25]	6.41	< 2.7 e -10**
Landmark Shift	-0.05	[-0.08, 0.03]	-1.22	0.20
Target-Landmark Distance	-1.05 **	[-3.04, -0.69]	-2.88	4.1 e -3**
Landmark Shift x Task (*PRO*)	0.08	[-0.19, 0.36]	0.60	0.54
Target-Landmark Distance x Task (*PRO*)	0.10	[-0.32, 1.10]	0.35	0.74

Note. B represents standardized regression weights. *LL* and *UL* indicate the lower and upper limits of the confidence interval, respectively. A: Model Adjusted R^2 = 0.18* , 95% CI[0.15,0.21]. B: Model Adjusted R^2 = 0.09 * , 95% CI[0.03,0.15]
* indicates p < 0.05. ** indicates p < .01.

3.3 Gaze-Centred Overshoot

One potential source of error is the phenomenon of gaze-centered overshoot, sometimes called 'retinal magnification' (Bock, 1986; Enright, 1995; Henriques et al., 1998). In this case, the overshoot is expected to be 'outward' relative to the central fixation point. Fig. 7 shows the data distributions of overshoot errors (across participant means) for both the *EGO-Instruction* (blue) and *ALLO-Instruction* data, separated into LEFT and RIGHT visual fields. Overall, the data showed the expected trends, i.e., leftward overshoots in the left visual field and rightward overshoots in the right visual field, but the magnitudes were smaller for the *ALLO-Instruction*, especially in the *ANTI-Task* (Fig. 8 B).

We analyzed the influence of instruction, task, and visual field on gaze-centered overshoots using a generalized linear mixed model (Table 4). This analysis revealed a significant effect of *EGO/ALLO Instruction* ($p = 8.9$ e-9) and interaction effects which were analysed using post-hoc t-tests. The effect of *EGO/ALLO* instructions was conditional, depending on the visual field of response and whether participants pointed to the same or opposite side of the target. In the *PRO task*, the *ALLO-Instruction* only produced significantly smaller overshoot errors the left visual field ($p = 0.03$), whereas it produced significantly smaller errors in both the left ($p = 3.1$ e-4) and right ($p = 2.7$ e-4) visual fields in the *ANTI-Task*. The second interaction indicated a significantly higher overshot error in the *PRO-Task* compared to the *ANTI-Task*, but only in the right visual field. Notably, this effect was solely driven by overshoot reaching errors in the *ALLO Instruction* condition, as the absolute overshoot error was less in the *EGO Instruction* condition when participants had to pro-point rather than anti-point. Overall, the main finding here was that the *ALLO Instruction* appeared to suppress gaze-centered overshoot errors in these tasks.

Fig. 7: Mean Overshoot Errors for *EGO Instruction* (blue) and *ALLO Instruction* (red) Idata. (same graphic conventions as Fig. 5). To obtain each data point, the horizontal distance between reach endpoints and expected target location (relative to fixation) was averaged separately for each goal position and then across goals within the left and right visual fields respectively. **(A)** *PRO Task* data. **(B)** *ANTI-Task* data. Significant differences are shown by an asterisk, '*' p < 0.0, '**' p < 0.001.

Table 4

Fixed Effects of Multi-level Model using Absolute Overshoot Error as the Criterion

Predictor	B	B 95% CI [LL, UL]	t value	p
(Intercept)	2.00**	[1.75, 2.25]	15.69	< 2.0 e -16**
Task (*PRO*)	-0.17	[-0.44, 0.10]	-1.23	0.21
Instruction (*EGO*)	-1.05 **	[-1.40, -0.69]	-5.82	8.9 e -9**
Visual Field (RIGHT)	0.08	[-0.19, 0.36]	0.60	0.54
Task (*PRO*) x Instruction (*EGO*)	-0.71**	[-1.10, -0.32]	-3.60	3.5e -4**
Visual Field (RIGHT) x Task x Instruction (*EGO*)	0.19*	[0.09, 0.49]	1.67	0.04*

Note. B represents standardized regression weights. *LL* and *UL* indicate the lower and upper limits of the confidence interval, respectively. Model Adjusted R^2 = 0.33*, 95% CI [0.25,0.41]
* indicates p < 0.05. ** indicates p < .01.

3.4 Reaction Time

Reaction times were assessed from the moment of the GO signal until movement onset. In general, the *ALLO*

Instuction data showed reduced reaction times relative to the *EGO Instruction* data. This advantage was

modest in the *PRO-Task* data (Fig. 8 A) but much stronger in the *ANTI-Task* data (Fig. 8 B), where *EGO*

Instruction reaction times appeared to be elevated relative to *PRO*-data.

This was assessed for the 2 (*EGO/ALLO*) conditions x 2 (*PRO/ANTI*) Tasks and 2 (*LEFT/RIGHT*) visual

fields using a generalized linear mixed model. The simplified model in Table 4 was fitted as there was no

significant three-way interaction effect. A significant effect of *EGO/ALLO Instruction* was observed (P = 4.9

e-4). Participants exhibited significantly slower reaction times in the *EGO Instruction* condition, with a

relative delay of 242.75 ms, 95% CI [109.30, 376.21]. Additionally, a significant interaction effect was

identified (p = 9.8 e-3), which was analyzed using a post-hoc interaction contrast: when averaging over the

visual field of response, the impact of *EGO/ALLO Instructions* conditions depended on the Task requirements

(i.e., whether participants had to point to the same or opposite side from the target). *ALLO Instruction* resulted

in faster reaction times, although the effect was only significant when participants had to point in the opposite

visual field (p = 0.03), with a reduction of 245.90 ms, 95% CI [136.9, 355.0]. A similar analysis showed no

significant difference in movement duration.

Fig. 8: Mean Reaction Times (same graphic conventions as Fig. 5). The reaction times were averaged for targets on the right and left visual fields of response (right and left side of the figure, respectively). **(A)** In the *PRO-Task*. **(B)** In the *ANTI Task* data. *EGO* and ALLO *Instruction* conditions are shown as blue and red in the figure, respectively. Significant differences are shown by an asterisk, '*' $p < 0.01$, '**' $p < 0.001$

Table 5

Fixed Effects of Multi-level Model using Reaction Time as the Criterion

Predictor	*B*	*B* 95% CI [LL, UL]	*t value*	*p*
(Intercept)	1001.82**	[907.45, 1096.19]	21.09	< 2.0 e -16**
Task (*PRO*)	14.30	[-94.67, 123.26]	0.26	0.79
Instruction (*EGO*)	-242.75 **	[-376.21, -109.30]	-3.61	4.9 e -4**
Visual Field (RIGHT)	18.77	[-127.73, 90.20]	0.34	0.73
Task (*PRO*) x Instruction (*EGO*)	-204.65	[-358.75, -50.55]	-2.64	9.8 e -3**
Visual Field (RIGHT) x Instruction (*EGO*)	-6.32	[-147.78, 160.42]	-0.08	0.94

Note. B represents standardized regression weights. *LL* and *UL* indicate the lower and upper limits of the confidence interval, respectively. Model Adjusted R^2 = 0.22* , 95% CI[0.18,0.26]

* indicates p < 0.05. ** indicates p < .01.

4. Discussion

Previous studies have reported the sensory influence of a visual landmark on reaching behavior (e.g.

references), or the weighting of egocentric and allocentric cues (e.g. references), but here we provide a

detailed behavioral analysis of the specific influence of *instructions* on performance in an otherwise

stimulus-matched, cue-conflict task. Specifically, we measured how the instruction to use (or ignore)

landmark-centered coordinates influences the accuracy, precision, and timing of memory-guided reaches,

either to the same or opposite visual hemifield as the target. Our results show that participants were

generally more accurate, precise, and quicker to react in the allocentric instruction condition, especially

when reaching to the visual field opposite to the target. We also observed a left / right visual field effect,

where performance was worse overall in the right visual field. We will interpret each of these findings

and other details below, in terms of previous literature, potential physiological mechanisms, and their

practical implications.

4.1 Egocentric vs. Allocentric Aiming: Natural vs. Experimental Conditions

Before discussing our data, we are obliged to consider how our experimental design relates to natural

reaching behavior (Fooken et al. 2023). First, the normal visual feedback of the goal and hand was removed

throughout the memory delay and reach. Reaches based on initial sensory conditions are thought to rely on

internal models of the eye-head-hand system (Blohm & Crawford, 2007; Blohm et al. 2009). In general, this

provides accurate behavior (Vercher et al., 1994), but with certain errors variable and systematic errors

(Henriques et al., 1998; Medendorp & Crawford, 2002; Van Pelt & Medendorp, 2007). It is noteworthy that

even without an instruction, the presence of a visual landmark tends to dampen these errors (Krigolson &

Heath, 2004; Lemay et al. 2004; Obhi & Goodale, 2005; Schütz et al., 2013), especially as the memory delay

increases (Chen et al., 2011).

Second, egocentric and allocentric cues normally complement each other, whereas we introduced a conflict

by surreptitiously shifting the landmark during the memory delay. It has been argued that this simulates

situations where visual landmarks are unstable (Byrne & Crawford 2010) or egocentric signals are unreliable

(Byrne et al. 2010, Chen et al. 2011). In the absence of explicit instruction, healthy individuals tend to

optimally weigh egocentric vs. allocentric cues based on their relative reliability and perceived stability

(Byrne & Crawford, 2010). This weighting is remarkably consistent (~1/3 allocentric, ~2/3 egocentric) and

generalizes well to naturalistic situations, although the exact ego / allocentric ratio is modulated by various

factors such as distance and similarity between the target and landmark (Fiehler et al., 2014; Klinghammer

et al., 2017). Although we instructed our participants to shift this weighting completely toward one cue or

the other, we cannot assume that the other cue had no implicit influence.

Finally, unlike most natural reaches (which employ the implicit processes above) we explicitly required our

participants to follow two different rules (egocentric vs. allocentric) based on a verbal instruction and a set of

color-coded cues. Specifically, the allocentric instruction required participants to both attend to the landmark

and remember its spatial relationship to the target. This required the integration of bottom-up perception, top-

down cognition, and environmental factors (Caduff & Timpf 2008). This additional degree of cognitive

processing could interfere with reaching performance. Thus, one cannot assume either that the allocentric

instruction would improve performance, or conversely, that the landmark would have no influence in the

Egocentric condition.

4.2 The EGO Instruction Condition as a Control: General Observations

Our *EGO Instruction* condition was designed as a control for the ALLO condition, based on several

assumptions confirmed in our results. First, as expected from previous studies (Goodale & Milner, 1992; Hu

et al., 1999; McIntyre et al., 1997; Westwood & Heath, 2003) participants were able to perform the task, i.e.,

reach positions correlated with the required goal positions, but with considerable variable and systematic

errors. In particular, the EGO Instruction data showed the expected gaze-centered overshoots (Henriques &

Crawford, 2000; Henriques et al., 1998; Van Pelt & Medendorp, 2007), which likely originate in the

comparison between hand and target position signals used to compute the reach vector in visual coordinates

(Dessing et al., 2009).

In the *ANTI-task* version of the *EGO condition*, participants had to suppress the normal pro-reach, and then

calculate an opposite reach goal in visual coordinates (Everling & Munoz, 2000; Cappadocia et al., 2017).

As predicted, this appeared to produce additional egocentric noise in the system, increasing both systematic

and variable errors increased in the *ANTI* version of the *EGO* condition.

Finally, neither target-landmark distance or landmark shift amplitude had a significant influence on

performance in the *EGO Instruction* task. This result is consistent with active suppression of landmark

information (although we cannot quantify this without a 'no instruction' control). Overall, the *EGO-*

instruction produced the expected results, providing a control for comparison with the ALLO-Instruction.

We will focus on the difference between these conditions below.

4.3 Influence of the Landmark Instruction: Precision and Accuracy

Our main result was that the instruction to reach relative to the landmark generally enhanced both accuracy

and precision relative to egocentric instruction. In particular, there was a systematic reduction of the gaze-

centered overshoot effect that was observed in our egocentric data and many previous studies (Henriques &

Crawford, 2000; Henriques et al., 1998; Van Pelt & Medendorp, 2007). We also found significant

improvements in variable error. Likely this advantage was due to the greater attentional weighting on

landmark cues, which tend to improve performance (Lemay et al. 2004), especially after a delay (Goodale

& Milner, 1992; Hu et al., 1999; McIntyre et al., 1997; Hay & Redon, 2006; Krigolson & Heath, 2004;

Milner & Goodale, 2006; Chen et al. 2011).

Although we equalized the stimuli in conditions, our instructions may have interacted with stimulus

perception in some way. In contrast previous studies (Fiehler et al., 2014; Aagten-Murphy and Bays, 2019)

landmark influences on performance increased with distance from the target. This is likely because target-

landmark difference used here was small (1.15-3.44°) compared to those previous studies, and relative to the

distance of stimuli from the fovea 4.57-15.07. The beneficial influence of our nearest landmarks might have

been negated by sensory confusion within response fields during initial perception (Ransom-Hogg &

Spillmann, 1980) or more likely, confusion between their representations at the time of response. It is also

possible that our choice of color cues interacted with the instruction in some fashion (Nakshian, 1964;

Nathans, 1999). But overall, these factors do not negate or explain the benefits of instruction, especially in

light of the known advantages of landmark-centered coding for spatial memory (Chen et al. 2011; Lemay et

al., 2004).

Finally, we cannot find previous literature that focused on the influence of ego / allocentric *instruction* on

behavior, but other studies have reported conficting behavior data from tasks that employed such

instructions. For example, Byrne & Crawford (2010) reported that their control egocentric and allocentric

reach tasks did not show significantly different variable error. However, there was no cue conflict in their

control tasks, and stimuli were not held constant in these tasks (i.e., landmark present in the landmark task

but not in the egocentric task). Our results also differed from those of Thaler & Tod (2009), who found

reaching precision to be lowest in their allocentric task. But again there was no cue-conflict in their paradigm,

and the task involved hand alignment, which is thought to employ different mechanisms than reach (Goodale

et al., 1994; Monaco et al. 2024).

4.4 Results Specific to Anti-Pointing: Accuracy and Timing

Our secondary finding was that the advantage of the allocentric instruction was more pronounced for reaches

to the opposite visual field than the target, in terms of accuracy, precision and reaction time. As expected,

these effects appear to be driven by decreased performance (including larger gaze-centered overshoots) in the

EGO/ANTI condition relative to the *EGO/PRO* condition, whereas the *ALLO* condition showed similar

performance for both the *ANTI* and *PRO* task versions. This is likely because anti-reaching requires specific

cognitive transformations and neural mechanisms (Fischer & Weber, 1992; Fernandez-Ruiz et al. 2007; Gail

et al., 2009), whereas the mechanisms for reaching relative to the landmark (Chen et al., 2014). However, anti-

movements either require the reversal of a motor vector plan. It is noteworthy that anti-movements fall within

the class of 'non-standard' rule-based spatial transformations that pervade much of our modern life (Sergio et

al., 2009), so these behaviors might benefit the most from the presence and use of allocentric landmarks.

4.5 Visual Field Dependence

Our tertiary observation was that the benefit of the allocentric instruction for anti-reaching was asymmetric

across the visual hemifields. Whereas the egocentric anti-pointing task produced symmetric gaze-centered

errors, the allocentric instruction only mitigated these errors for goals / landmarks that shifted from the right to

left visual field. Byrne et al. (2010) found that allocentric landmarks had more influence on pointing when

saccades caused remembered targets to remap from the right to left hemifield. This might be related somehow

to a tendency of our participants to be right-hand and right eye dominant. Also, various studies have suggested

that the right cortical hemisphere (corresponding to the left visual) plays a stronger role in allocentric

processing than the left (Faillenot et al. 1999; Galati et al. 2000; Zaehle et al., 2007; Fink et al., 2003, Weiss et

al., 2006).

4.6 Possible Physiological Mechanisms

Most sensorimotor neuroscience studies only consider the egocentric transformation from sensory to motor

coordinates, but to interpret the current results we must consider both 1) the integration of sensory input to

construct two different spatial rules, and 2) the implementation of these rules through separate allocentric

and egocentric transformations.

For the first 'instruction' component, one might safely assume that the auditory instruction is processed

through auditory cortex and higher-level language areas (Wernicke 1874; Friederici et al, 2000; Humphries

et al., 2001; Vandenberghe et al., 2002), whereas initial color and spatial information from the visual stimuli

are processed by well-known pathways in occipital cortex (DeYoe & Van Essen, 1988; Livingstone &

Hubel, 1988; Zeki, 1993). The key question is, how does the former impose a rule on how the latter is

processed? One likely candidate area is prefrontal cortex. For example, lateral prefrontal cortex receives

complex multisensory inputs and is thought to play roles in imposing rules on sensory inputs (Miller &

Cohen, 2001), including both inhibiting incorrect responses (DeSouza et al., 2003) and selecting correct

responses (Rowe et al., 2000). It is also likely that the parietofrontal attention network is involved in directing

attention to the landmark (Buschman & Miller, 2007; Bisley & Goldberg, 2003). Consistent with these

speculations, parietofrontal cortex is thought to exert recurrent influence on early visual areas during reach

(Cappadocia et al. 2017; Blohm et al. 2019) and appears to alter visual processing in these areas (Velji-

Ibrahim et al. 2018; Monaco et al. 2024).

Once such rules are enacted, it is thought that egocentric and allocentric transformations for reach are

processed through separate pathways. Classic neuropsychology experiments suggest that the dorsal visual

stream (via parietal cortex) handles egocentric visual transformations (Goodale & Milner 1992; Carey et al.,

2006; Schenk, 2006). Numerous studies have examined the role of parietal cortex in action; too many to

review here (e.g., Buneo & Andersen, 2006; Crawford et al. 2011; Gallivan & Culham, 2015; Vesia &

Crawford, 2012). However, it is pertinent that damage to the dorsal stream alters the gaze-centred errors

reported here (Khan et al. 2005a, 2005b), and that the visual field effects observed here could be attributed

to interactions between hand and hemifield lateralization in parietal cortex (Perenin & Vighetto, 1988;

Rossetti et al., 2003; Medendorp et al. 2005). Parietal cortex also plays a role in coding rule-based egocentric

transformations, such as the anti-reach task (Gail & Andersen, 2006; Cappadocia et al. 2017).

In contrast, the ventral visual stream (via temporal cortex) is thought to handles allocentric transformations

(Goodale & Milner 1992; Carey et al., 2006; Schenk, 2006). This distinction is supported by neuroimaging

experiments that showed spatial tuning for egocentric targets in the dorsal stream and allocentrically defined

reach targets in the ventral stream and (Chen et al. 2014). Ultimately the latter must be integrated into the

motor system for egocentric control of action, and this appears to happen in both parietal and frontal cortex

(Chen et al. 2018). Consistent with this, frontal cortex visual responses integrate target and landmark location

(Schütz et al. 2023), whereas their memory and motor responses for gaze are able to integrate allocentric

and egocentric coordinates in a cue conflict task similar to that used in reach studies (Bharmauria et al. 2020,

2021).

5. Conclusions

Whereas a number of studies have reported the influence of allocentric landmarks on reach, and some studies

employed explicit instructions to reach in allocentric coordinates, this study examined the interaction of these

two factors: whether an explicit instruction to attend to and use a landmark to encode reach direction

provides additional behavioral benefits. We found that reach performance (accuracy, precision, reaction

time) was generally enhanced, especially in the more difficult task of reaching toward a location defined in

the visual hemifield opposite to the target. This has the practical implication that explicit instructions to use

visual landmark cues may enhance performance, especially in high-demand 'non-standard' spatial tasks

(Sergio et al. 2009; Dalecki et al., 2019), and for individuals with degraded egocentric transformations due

to age, developmental disorders, or brain damage (e.g., Tippett et al., 2007; Niechwiej-Szwedo et al. 2011;

Goodale et al., 1994; Khan et al., 2005).

6. References

Aagten-Murphy, D., & Bays, P. M. (2019). Independent working memory resources for egocentric and allocentric spatial information. *PLoS Computational Biology*, 1–20.

Bharmauria, V., Sajad, A., Schütz, A., Yan, X., Wang, H., & Crawford, J. D. (2020). Predictive role of supplementary eye fields for egocentric/allocentric integration. Journal of Vision, 20(11), 977-977.

Bharmauria, V., Schütz, A., Khoozani, P. A., Yan, X., Wang, H., Bremmer, F., & Crawford, J. D. (2021). Gaze-Related Activity in Primate Frontal Cortex Predicts and Mitigates Spatial Uncertainty. bioRxiv, 2021-06.

Bisley, J. W., & Goldberg, M. E. (2003). Neuronal activity in the lateral intraparietal area and spatial attention. Science, 299(5603), 81-86.

Blohm, G., & Crawford, J. D. (2007). Computations for geometrically accurate visually guided reaching in 3-D space. *Journal of Vision*, *7*(5), 4-4.

Blohm, G., Keith, G. P., & Crawford, J. D. (2009). Decoding the cortical transformations for visually guided reaching in 3D space. Cerebral Cortex, 19(6), 1372-1393.

Bock, O. (1986). Contribution of retinal versus extraretinal signals towards visual localization in goal-directed movements. Experimental brain research, 64(3), 476-482.

Buneo, C. A., & Andersen, R. A. (2006). The posterior parietal cortex: sensorimotor interface for the planning and online control of visually guided movements. Neuropsychologia, 44(13), 2594-2606.

Buschman, T. J., & Miller, E. K. (2007). Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. science, 315(5820), 1860-1862.

Byrne, P. A., & Crawford, J. D. (2010). Cue reliability and a landmark stability heuristic determine relative weighting between egocentric and allocentric visual information in memory-guided reach. *Journal of Neurophysiology*, *103*(6), 3054–3069.

Byrne, P. A., Cappadocia, D. C., & Crawford, J. D. (2010). Interactions between gaze-centered and allocentric representations of reach target location in the presence of spatial updating. *Vision Research*, *50*(24), 2661–2670.

Caduff, D., & Timpf, S. (2008). On the assessment of landmark salience for human navigation. *Cognitive processing*, *9*(4), 249-267.

Cappadocia, D. C., Monaco, S., Chen, Y., Blohm, G., & Crawford, J. D. (2017). Temporal evolution of target representation, movement direction planning, and reach execution in occipital–parietal–frontal cortex: an fMRI study. Cerebral Cortex, 27(11), 5242-5260.

Carey, D. P., Dijkerman, H. C., Murphy, K. J., Goodale, M. A., & Milner, A. D. (2006). Pointing to places and spaces in a patient with visual form agnosia. *Neuropsychologia*, *44*(9), 1584-

Chen, Y., Byrne, P., & Crawford, J. D. (2011). Time course of allocentric decay, egocentric decay, and allocentric-to-egocentric conversion in memory-guided reach. *Neuropsychologia*, *49*(1), 49-60.

Chen, Y., Monaco, S., Byrne, P. A., Yan, X., Henriques, D. Y. P., & Crawford, J. D. (2014). Allocentric versus Egocentric Representation of Remembered Reach Targets in Human Cortex. *Journal of Neuroscience*, *34*(37), 12515–12526.

Chen, Y., Monaco, S., & Crawford, J. D. (2018). Neural substrates for allocentric-to-egocentric conversion of remembered reach targets in humans. European Journal of Neuroscience, 47(8), 901-917.

Core Team, R. (2021). R: A language and environment for statistical computing [computer software]. Vienna, Austria: R Foundation for Statistical Computing.

Crawford, J. D., Henriques, D. Y., & Medendorp, W. P. (2011). Three-dimensional transformations for goal-directed action. Annual review of neuroscience, 34, 309-331.

Dalecki, M., Gorbet, D. J., Macpherson, A., & Sergio, L. E. (2019). Sport experience is correlated with complex motor skill recovery in youth following concussion. European journal of sport science, 19(9), 1257-1266.

DeSouza, J. F., Menon, R. S., & Everling, S. (2003). Preparatory set associated with pro-saccades and anti-saccades in humans investigated with event-related FMRI. Journal of Neurophysiology, 89(2), 1016-1023.

Dessing, J. C., Oostwoud Wijdenes, L., Peper, C. E., & Beek, P. J. (2009). Visuomotor transformation for interception: catching while fixating. Experimental brain research, 196, 511-527.

DeYoe, E. A., & Van Essen, D. C. (1988). Concurrent processing streams in monkey visual cortex. Trends in neurosciences, 11(5), 219-226.

Enright, J. T. (1995). The non-visual impact of eye orientation on eye—hand coordination. Vision research, 35(11), 1611-1618.

Everling, S., & Munoz, D. P. (2000). Neuronal correlates for preparatory set associated with pro-saccades and anti-saccades in the primate frontal eye field. Journal of Neuroscience, 20(1), 387-400.

Faillenot, I., Decety, J., & Jeannerod, M. (1999). Human brain activity related to the perception of spatial features of objects. *Neuroimage*, *10*(2), 114-124.

Fernandez-Ruiz, J., Goltz, H. C., DeSouza, J. F., Vilis, T., & Crawford, J. D. (2007). Human parietal "reach region" primarily encodes intrinsic visual direction, not extrinsic movement direction, in a

visual–motor dissociation task. Cerebral Cortex, 17(10), 2283-2292.

Fiehler, K., Wolf, C., Klinghammer, M., & Blohm, G. (2014). Integration of egocentric and allocentric information during memory-guided reaching to images of a natural environment. *Frontiers in Human Neuroscience*, *8*, 636.

Fink, G. R., Marshall, J. C., Weiss, P. H., Stephan, T., Grefkes, C., Shah, N. J., ... & Dieterich, M. (2003). Performing allocentric visuospatial judgments with induced distortion of the egocentric reference frame: an fMRI study with clinical implications. Neuroimage, 20(3), 1505-1517.

Fischer, B., & Weber, H. (1992). Characteristics of "anti" saccades in man. Experimental Brain Research, 89, 415-424.

Fooken, J., Baltaretu, B. R., Barany, D. A., Diaz, G., Semrau, J. A., Singh, T., & Crawford, J. D. (2023). Perceptual-Cognitive Integration for Goal-Directed Action in Naturalistic Environments. Journal of Neuroscience, 43(45), 7511-7522.

Friederici, A. D., Meyer, M., & Von Cramon, D. Y. (2000). Auditory language comprehension: an event-related fMRI study on the processing of syntactic and lexical information. Brain and language, 74(2), 289-300.

Gail, A., & Andersen, R. A. (2006). Neural dynamics in monkey parietal reach region reflect context-specific sensorimotor transformations. Journal of Neuroscience, 26(37), 9376-9384.

Gail, A., Klaes, C., & Westendorff, S. (2009). Implementation of spatial transformation rules for goal-directed reaching via gain modulation in monkey parietal and premotor cortex. Journal of Neuroscience, 29(30), 9490-9499.

Galati, G., Lobel, E., Vallar, G., Berthoz, A., Pizzamiglio, L., & Le Bihan, D. (2000). The neural basis of egocentric and allocentric coding of space in humans: a functional magnetic resonance study. *Experimental brain research*, *133*(2), 156-164.

Gallivan, J. P., & Culham, J. C. (2015). Neural coding within human brain areas involved in actions. Current opinion in neurobiology, 33, 141-149.

Goodale, M. A., & Milner, A. D. (1992). Separate visual pathways for perception and action. *Trends in neurosciences*, *15*(1), 20-25.

Goodale, M. A., Jakobson, L. S., Milner, A. D., Perrett, D. I., Benson, P. J., & Hietanen, J. K. (1994). The nature and limits of orientation and pattern processing supporting visuomotor control in a visual form agnosic. Journal of Cognitive Neuroscience, 6(1), 46-56.

Goodale, M. A., Westwood, D. A., & Milner, A. D. (2004). Two distinct modes of control for object-directed action. *Progress in brain research*, *144*, 131-144.

Green, P., & MacLeod, C. J. (2016). SIMR: an R package for power analysis of generalized linear mixed models by simulation. *Methods in Ecology and Evolution*, *7*(4), 493-498.

Hay, L., & Redon, C. (2006). Response delay and spatial representation in pointing movements. *Neuroscience letters*, *408*(3), 194-198.

Henriques, D. Y. P., & Crawford, J. D. (2000). Direction-dependent distortions of retinocentric space in the visuomotor transformation for pointing. *Experimental brain research*, *132*(2), 179-194.

Henriques, D. Y., Klier, E. M., Smith, M. A., Lowy, D., & Crawford, J. D. (1998). Gaze-centered remapping of remembered visual space in an open-loop pointing task. *Journal of Neuroscience*, *18*(4), 1583-1594.

Howard, I. P., & Templeton, W. B. (1966). Human spatial orientation.

Hu, Y., Eagleson, R., & Goodale, M. A. (1999). The effects of delay on the kinematics of grasping. *Experimental Brain Research*, *126*(1), 109-116.

Humphries, C., Willard, K., Buchsbaum, B., & Hickok, G. (2001). Role of anterior temporal cortex in auditory sentence comprehension: an fMRI study. Neuroreport, 12(8), 1749-1752.

Khan, A. Z., Pisella, L., Vighetto, A., Cotton, F., Luauté, J., Boisson, D., ... & Rossetti, Y. (2005). Optic ataxia errors depend on remapped, not viewed, target location. Nature neuroscience, 8(4), 418-420.

Khan, A. Z., Pisella, L., Rossetti, Y., Vighetto, A., & Crawford, J. D. (2005). Impairment of gaze-centered updating of reach targets in bilateral parietal–occipital damaged patients. *Cerebral Cortex*, *15*(10), 1547-1560.

Klinghammer, M., Blohm, G., & Fiehler, K. (2017). Scene configuration and object reliability affect the use of allocentric information for memory-guided reaching. Frontiers in Neuroscience, 11, 229248.

Krigolson, O., & Heath, M. (2004). Background visual cues and memory-guided reaching. *Human movement science*, *23*(6), 861-877.

Lemay, M., Bertram, C. P., & Stelmach, G. E. (2004). Pointing to an allocentric and egocentric remembered target. *Motor control*, *8*(1), 16-32.

Lemay, M., & Stelmach, G. E. (2005). Multiple frames of reference for pointing to a remembered target. Experimental Brain Research, 164, 301-310.

Li, J., Sajad, A., Marino, R., Yan, X., Sun, S., Wang, H., & Crawford, J. D. (2017). Effect of allocentric landmarks on primate gaze behavior in a cue conflict task. Journal of vision, 17(5), 20-20.

Livingstone, M., & Hubel, D. (1988). Segregation of form, color, movement, and depth: anatomy, physiology, and perception. Science, 240(4853), 740-749.

McIntyre, J., Stratta, F., & Lacquaniti, F. (1997). Viewer-centered frame of reference for pointing to memorized targets in three-dimensional space. *Journal of neurophysiology*, *78*(3), 1601-1618.

Medendorp, P. W., & Crawford, D. J. (2002). Visuospatial updating of reaching targets in near and far space. *Neuroreport*, *13*(5), 633-636.

Medendorp, W. P., Goltz, H. C., Crawford, J. D., & Vilis, T. (2005). Integration of target and effector information in human posterior parietal cortex for the planning of action. Journal of Neurophysiology, 93(2), 954-962.

Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. Annual review of neuroscience, 24(1), 167-202.

Milner, D., & Goodale, M. (2006). *The visual brain in action* (Vol. 27). OUP Oxford.

Monaco, S., Menghi, N., & Crawford, J. D. (2024). Action-specific feature processing in the human cortex: An fMRI study. Neuropsychologia, 194, 108773.

Nakshian, J. S. (1964). The effects of red and green surroundings on behavior. *The Journal of General Psychology*, *70*(1), 143-161.

Nathans, J. (1999). The evolution and physiology of human color vision: insights from molecular genetic studies of visual pigments. *Neuron*, *24*(2), 299-312.

Niechwiej-Szwedo, E., Goltz, H. C., Chandrakumar, M., Hirji, Z., Crawford, J. D., & Wong, A. M. (2011). Effects of anisometropic amblyopia on visuomotor behavior, part 2: visually guided reaching. Investigative ophthalmology & visual science, 52(2), 795-803.

Obhi, S. S., & Goodale, M. A. (2005). The effects of landmarks on the performance of delayed and real-time pointing movements. Experimental Brain Research, 167, 335-344.

Perenin, M. T., & Vighetto, A. (1988). Optic ataxia: A specific disruption in visuomotor mechanisms: I. Different aspects of the deficit in reaching for objects. *Brain*, *111*(3), 643-674.

Ransom-Hogg, A., & Spillmann, L. (1980). Perceptive field size in fovea and periphery of the light-and dark-adapted retina. Vision Research, 20(3), 221-228.

Redon, C., & Hay, L. (2005). Role of visual context and oculomotor conditions in pointing accuracy. *Neuroreport*, *16*(18), 2065-2067.

Rossetti, Y., Pisella, L., & Vighetto, A. (2003). Optic ataxia revisited. *Experimental Brain Research*, *153*(2), 171-179.

Rowe, J. B., Toni, I., Josephs, O., Frackowiak, R. S., & Passingham, R. E. (2000). The prefrontal cortex: response selection or maintenance within working memory?. Science, 288(5471), 1656-1660.

Sergio, L. E., Gorbet, D. J., Tippett, W. J., Yan, X., & Neagu, B. (2009). Cortical mechanisms of vision for complex action. Cortical mechanisms of vision.

Schenk, T. (2006). An allocentric rather than perceptual deficit in patient DF. *Nature neuroscience*, *9*(11), 1369-1370.

Soechting, J. F., & Flanders, M. (1992). Moving in three-dimensional space: frames of reference, vectors, and coordinate systems. *Annual review of neuroscience*, *15*(1), 167-191.

Schütz, I., Henriques, D. Y. P., & Fiehler, K. (2013). Gaze-centered spatial updating in delayed reaching even in the presence of landmarks. Vision Research, 87, 46-52.

Schütz, A., Bharmauria, V., Yan, X., Wang, H., Bremmer, F., & Crawford, J. D. (2023). Integration of landmark and saccade target signals in macaque frontal cortex visual responses. Communications Biology, 6(1), 938.

Thaler, L., & Todd, J. T. (2009). The use of head/eye-centered, hand-centered and allocentric representations for visually guided hand movements and perceptual judgments. *Neuropsychologia*, *47*(5), 1227-1244.

Tippett, W. J., Krajewski, A., & Sergio, L. E. (2007). Visuomotor integration is compromised in Alzheimer's disease patients reaching for remembered targets. European neurology, 58(1), 1-11.

Van Pelt, S., & Medendorp, W. P. (2007). Gaze-centered updating of remembered visual space during active whole-body translations. Journal of neurophysiology, 97(2), 1209-1220.

Vandenberghe, R., Nobre, A. C., & Price, C. J. (2002). The response of left temporal cortex to sentences. Journal of cognitive neuroscience, 14(4), 550-560.

Velji-Ibrahim, J., Crawford, J. D., Cattaneo, L., & Monaco, S. (2018). Action intention modulates the representation of object features in early visual cortex. bioRxiv, 480574.

Vercher, J. L., Magenes, G., Prablanc, C., & Gauthier, G. M. (1994). Eye-head-hand coordination in pointing at visual targets: spatial and temporal analysis. *Experimental brain research*, *99*(3), 507-523.

Vesia, M., & Crawford, J. D. (2012). Specialization of reach function in human posterior parietal cortex. Experimental brain research, 221, 1-18.

Vindras, P., & Viviani, P. (1998). Frames of reference and control parameters in visuomanual pointing. *Journal of Experimental Psychology: Human Perception and Performance*, *24*(2), 569.

Vogeley, K., & Fink, G. R. (2003). Neural correlates of the first-person-perspective. *Trends in cognitive sciences*, *7*(1), 38-42.

Weiss, P. H., Rahbari, N. N., Lux, S., Pietrzyk, U., Noth, J., & Fink, G. R. (2006). Processing the spatial configuration of complex actions involves right posterior parietal cortex: an fMRI study with clinical implications. Human brain mapping, 27(12), 1004-1014.

Wernicke, C., Cohen, R. S., & Wartofsky, M. W. (1874). Boston studies in the philosophy of science. The symptom complex of aphasia: A psychological study on an anatomical basis, D. Reichel, Dordrecht, 1969, 34-97.

Westwood, D. A., Heath, M., & Roy, E. A. (2003). No evidence for accurate visuomotor memory: Systematic and variable error in memory-guided reaching. Journal of motor behavior, 35(2), 127-133.

Zaehle, T., Jordan, K., Wüstenberg, T., Baudewig, J., Dechent, P., & Mast, F. W. (2007). The neural basis of the egocentric and allocentric spatial frame of reference. *Brain research*, *1137*, 92-103.

Zeki, S. (1993). A vision of the brain. Blackwell scientific publications.