Adolescent THC impacts on mPFC dopamine-mediated cognitive processes in male and female rats

Martinez*, Maricela X.[1], Alizo Vera, Vanessa[1], Ruiz, Christina M.[1], Floresco, Stan B.[2], Mahler*, Stephen V.[1]

Affiliations:

[1]Department of Neurobiology and Behavior, University of California, Irvine. 1203 McGaugh Hall. Irvine, CA 92697

Ph: (949) 824-6128

[2]Department of Psychology, University of British Columbia, Vancouver, British Columbia, V6T 1Z4, Canada

Corresponding authors*: maricexm@uci.edu, mahlers@uci.edu

Funding: Funding was provided by NIH grants P50 DA044118, R01 MH132680, R01 DA055849, U01 DA053826 to SVM,

NSF GRFP DGE-1839285 to MXM, and NSERC (RGPIN-2018-04295) and CIHR (PJT-162444) grants to SBF. The authors

announce no conflicts of interest.

Abstract

Rationale: Adolescent cannabis use is linked to later-life changes in cognition, learning, and memory. Rodent experimental studies suggest Δ^9-tetrahydrocannabinol (THC) influences development of circuits underlying these processes, especially in the prefrontal cortex, which matures during adolescence.

Objective: We determined how 14 daily THC injections (5mg/kg) during adolescence persistently impacts medial prefrontal cortex (mPFC) dopamine-dependent cognition.

Methods: In adult Long Evans rats treated as adolescents with THC (AdoTHC), we quantify performance on two mPFC dopamine-dependent reward-based tasks—strategy set shifting and probabilistic discounting. We also determined how acute dopamine augmentation with amphetamine (0, 0.25, 0.5 mg/kg), or specific chemogenetic stimulation of ventral tegmental area (VTA) dopamine neurons and their projections to mPFC impacts probabilistic discounting.

Results: AdoTHC sex-dependently impacts acquisition of cue-guided instrumental reward seeking, but has minimal effects on set-shifting or probabilistic discounting in either sex. When we challenged dopamine circuits acutely with amphetamine during probabilistic discounting, we found reduced discounting of improbable reward options, with AdoTHC rats being more sensitive to these effects than controls. In contrast, neither acute chemogenetic stimulation of VTA dopamine neurons nor pathway-specific chemogenetic stimulation of their projection to mPFC impacted probabilistic discounting in control rats, although stimulation of this cortical dopamine projection slightly disrupted choices in AdoTHC rats.

Conclusions: These studies confirm a marked specificity in the cognitive processes impacted by AdoTHC exposure. They also suggest that some persistent AdoTHC effects may alter amphetamine-induced cognitive changes in a manner independent of VTA dopamine projections to mPFC, or via alterations of non-VTA dopamine neurons.

Key words: THC, cognition, dopamine, ventral tegmental area, medial prefrontal cortex, chemogenetics

Introduction

Cannabis is one of the most widely used drugs among adolescents, and its availability is increasing around the world. Human studies show that early exposure to cannabis, and especially its main psychoactive constituent Δ^9-tetrahydrocannabinol (THC), is associated with later-life cognitive impairments, and increased risk for psychiatric disorders including schizophrenia and addiction (Curran et al. 2016; Ehrenreich et al. 1999; Jenni et al. 2017; Malone et al. 2010; Murray et al. 2022; Rubino and Parolaro 2016; Schneider 2008; Volkow et al. 2016). However, in humans it is difficult to dissect whether THC exposure causes these associations, or whether early cannabis use and long-term deficits both result instead from other underlying comorbidities or risk factors. Rodent models are thus essential for establishing casual effects of THC on the developing adolescent brain.

Neurodevelopmental disruptions persisting long after adolescent cannabis use are plausible because adolescence is a dynamic critical period for structural and functional brain remodeling, especially in late-developing structures like the prefrontal cortex (PFC) (Andersen 2003; Casey et al. 2000). Some of this age-dependent plasticity seems to involve the endocannabinoid system, with dynamic changes in cannabinoid receptors (CBRs) and endocannabinoids (ECB) occurring across adolescence (Bara et al. 2021; Ellgren et al. 2008; Heng et al. 2011; Lee et al. 2016; Simone et al. 2022). Might THC, which also acts via CBRs, disrupt this age-dependent ECB signaling system and thus leave long-lasting consequences on the brain? If so, the adolescent-developing PFC (Peters et al. 2022; Scheyer et al. 2023; Spear 2000), and its dopaminergic inputs from ventral tegmental area (VTA), which are actively innervating during this period (Hoops and Flores 2017; Manitt et al. 2011; Reynolds et al. 2018), are a likely candidate for cognition-relevant neurodevelopmental insults caused by adolescent THC (Molla and Tseng 2020; Renard et al. 2017a; Renard et al. 2017b).

The adult PFC is crucial for purposeful, goal-directed behaviors driven by the ability to flexibly converge our internal states, like reward motivation, with outside external information about contexts, cues, and rules (Miller 2000; Miller and Cohen 2001; Ott and Nieder 2019). Executive functions like working memory, attention, rule shifting, and decision-making require PFC-dependent cognitive control. Rodent studies show adolescent cannabinoid drug exposure can cause persistent deficits in working memory (De Melo et al. 2005; Kirschmann et al. 2017; O'Shea et al. 2004; Quinn et al. 2008; Schneider and Koch 2003), social cognition (O'Shea et al. 2004; Renard et al. 2017a; Renard et al. 2017b; Zamberletti et al. 2014), and cognitive flexibility (Egerton et al. 2005; Gomes et al. 2015; Hill et al. 2006; Jacobs-Brichford et al. 2019; Szkudlarek et al. 2019) that may depend upon PFC.

Furthermore, dopamine in PFC plays a major role in decision making, working memory, cognitive flexibility, and goal-directed behaviors (Floresco 2013; Goldman-Rakic 1995; Goto et al. 2007; Seamans and Yang 2004). PFC dopamine dysfunction has also been implicated in schizophrenia symptoms (Davis et al. 1991; Howes and Kapur 2009), and there is a clear link between recent cannabis use and onset of psychosis in humans (Andreasson et al. 1987; D'Souza et al. 2004; Hambrecht and Hafner 2000), though it is not clear that this link is causal in nature (Fergusson et al. 2005; Henquet et al. 2005; Sewell et al. 2009). Since VTA dopamine neuron axons actively infiltrate mPFC during adolescence and are thus subject to disruption by THC exposure (Behan et al. 2012; Renard et al. 2017b), and since THC impacts dopamine signaling at several key nodes in reward and salience circuits (Bloomfield et al. 2016; Bossong et al. 2015; Chen et al. 1990; French et al. 1997; Gardner 2002; Gessa et al. 1998; Malone and Taylor 1999; Pistis et al. 2002; Renard et al. 2017b; Sakurai-Yamashita et al. 1989; Wu and French 2000), it is logical that adolescent THC might exert some of its neurodevelopmental disruptions in cognition-relevant non-VTA dopaminergic inputs to mPFC.

We therefore explored this possibility in a series of experiments characterizing effects of a well-characterized, translationally relevant adolescent THC exposure protocol in rats (Ruiz et al. 2021a; Ruiz et al. 2021b; Torrens et al. 2022; Torrens et al. 2020) upon adulthood mPFC dopamine-dependent cognition.

Materials and Methods

Subjects

Long Evans rats (n = 29 males, 25 females) were used for set shifting, probabilistic discounting and amphetamine experiments, and transgenic TH:Cre (n = 19 males, 9 females) and wildtype littermates (n= 14 males, 4 females) were used for chemogenetic experiments. Rats were pair or triple-housed in same-sex groups at weaning (postnatal day; PD21), in a temperature, humidity, and pathogen-controlled colony under a 12:12hr reverse light/dark cycle. Water was provided *ad libitum* and food was restricted to ~85% of free-feeding weight during behavioral testing, starting at ~PD70. Experiments were approved by University of California Irvine's Institutional Animal Care and Use Committee and were conducted in accordance with the NIH Guide for the Care and Use of Laboratory Animals.

Drugs

THC was provided in ethanol by the NIDA Drug Supply Program. For injection, THC was prepared fresh each day; ethanol is evaporated under N_2, and THC is reconstituted in 5% Tween-80 in saline, with heat and sonication, to 2 ml/kg for intraperitoneal (IP) injection. D-amphetamine hemisulfate salt (amphetamine) was attained from Sigma and mixed in saline at 0.5 mg/ml for injection. Clozapine-N-oxide (CNO) was obtained from the NIDA Drug Supply Program, stored at 4°C in

powder aliquots with desiccant, protected from light. For systemic injection, CNO (5 mg/kg) was dissolved daily for IP

injection in 5% dimethyl sulfoxide (DMSO; Sigma Aldrich) in 0.9% saline. For microinjections, CNO (1mM; 0.5 µl/side over

60s) was dissolved in artificial cerebrospinal fluid (Fisher) with 0.5% DMSO, stored in aliquots at -20°C, and thawed/vortexed

just before use.

Adolescent THC and Washout

All rats received daily IP injections of THC (5 mg/kg) or vehicle (VEH; 5% Tween-80 in saline) from PD30-43, followed by a

washout period of 21+ days, allowing full THC clearance in both sexes (Lee et al. 2022) prior to behavioral testing in

adulthood (**Fig. 1a**).

Experimental Design

Following adolescent THC/VEH treatment, 48 adult (PD 70+) rats (n = 29 males, 21 females) underwent strategy set-shifting

training, followed by training on probabilistic discounting until a group displayed stable levels of choice for 3 consecutive

days, determined with previously established criteria (St. Onge and Floresco 2009). Thereafter they underwent 3

counterbalanced amphetamine challenge tests, each (0, 0.25, 0.5 mg/kg IP) delivered 5 min before behavioral testing.

Following the challenge rats were retrained for at least 2 days before receiving their next challenge.

For chemogenetic experiments, another adolescent THC/VEH-treated group (n = 33 males, 13 females; n = 28 TH:Cre+,18

wildtype (WT)) underwent stereotaxic VTA virus injection of a Cre-dependent hM3Dq vector at ~PD65, and intra-mPFC

bilateral cannulae implantation at least 45 days later, at least 8 days prior to the first microinjection. Following recovery from

surgery, they were trained on the probabilistic discounting task to stability, then subjected to a series of counterbalanced

tests, again with re-stabilization training occurring between them. First, rats received 4 counterbalanced IP injections of the

DREADD agonist CNO (5 mg/kg) or VEH, 30 min prior to behavioral testing. Both CNO and VEH were administered twice

on separate days, and data was combined between both tests for analysis. Next, they underwent 4 additional

counterbalanced tests of probabilistic discounting, each held 5 min after intra-mPFC microinjections of CNO or VEH. Results

from intra-mPFC VEH and CNO tests were again averaged to increase reliability of findings. For choice data, the raw

number of risky choices in each block made on both VEH or CNO tests were summed and divided by the total number of

choices made in those respective blocks on the two VEH/CNO tests, thereby factoring out trial omissions. Win-stay/lose

shift values were combined in a similar manner, dividing the sum total of "stays" or "shifts" by the sum total of "wins" and

"losses" on the two respective tests. Latency and omission values were averaged across the tests.

Behavioral Methods

Operant Boxes

Training and testing took place in Med-Associates rat operant conditioning chambers (30.5 x 24 x 21 cm; St Albans, VT) within sound-attenuating boxes, equipped with two retractable levers with white lights above them, and white house light.

Operant Pretraining

Methods for both strategy set shifting and reward probabilistic discounting tasks closely followed prior reports (Floresco et al. 2008; St. Onge and Floresco 2009). Rats were first homecage-habituated to highly palatable, 45 mg banana-flavored reward pellets (Bio-Serv catalogue #: F0059), then given 2 days of magazine training in the operant boxes, where they received 38 pellets at variable intervals over a 60 min session. They were next sequentially trained to press each of two levers to receive a pellet over 2-5 days. On each lever training day, a lever extended into the chamber (side counterbalanced across rats), and one pellet was delivered on a fixed ratio 1 (FR1) schedule. Once they reliably pressed 50+ times in 30 min on a lever, they were transitioned to learning to press the other lever on the subsequent day, and training continued until meeting this criterion. Thereafter, they entered the next phase of the task, in which each lever was periodically extended into the chamber throughout a 30 min session. Levers extended in a pseudorandom order such that there were 45 left-lever trials and 45 right-lever trials, but no more than two consecutive trials on which the same lever was extended. Each lever extension was accompanied by illumination of a house light signaling the start of a trial (90 trials/session), and trials occurred every 20s. If the rat pressed the lever within 10s of the start of a trial, the lever was retracted and a pellet was delivered, and the house light remained on for 4s. If the rat failed to press in 10s, the lever retracted and the house light extinguished until the next trial, and the trial was considered an omission. Importantly, cue lights present above each lever were never illuminated during this training phase. Rats were trained in this manner for at least 5 days, or until they omitted less than 5 trials per session. Since individual rats frequently display idiosyncratic lever position biases that can influence interpretation of subsequent behavior (Brady and Floresco 2015), and to minimize such impacts of stochastic side-preference on subsequent tests, lever side preference was next assessed using a published protocol (Brady and Floresco 2015; Floresco et al. 2008).

Visual Cue Discrimination Training

Next, rats were trained to respond on only one of the two levers extended on each trial—whichever was signaled as the correct response by illumination of a cue light just above it (**Fig. 1b**). Sessions began with both levers retracted and the house light off. Every 20s, one of the two stimulus cue lights was illuminated in a randomized order, 3s later both levers extended, and the house light turned on. If the rat pressed the lever that had a cue light illuminated above it within 10s of

lever insertion, it received a pellet, the levers retracted, the stimulus light was extinguished and the house light remained

illuminated for 4s. If the rat did not choose a lever in 10s, the trial ended and levers retracted until the next trial, and the

trial was scored as an omission. This training proceeded for at least 30 trials, ending when rats either made 10 consecutive

correct responses, or after 150 trials had elapsed. If rats did not achieve 10 consecutive correct responses, they received

a second identical training session on the following day.

Strategy Set Shifting Test

After learning to follow a cue light to respond for reward in the prior training phase, rats then underwent a single 40 min

session on which the response rule was suddenly shifted; a procedure analogous to the Wisconsin Card Sorting Task used

in humans to aid diagnosis of PFC dysfunction (Owen et al. 1991; Pantelis et al. 1999). On this day, the non-preferred lever

(determined in side-bias training described above) became the correct response, and pressing it during trials yielded a pellet

and commencement of the next inter-trial interval (**Fig. 1h**). Light cues above one of the levers were presented just before

and during trials as in the prior visual cue discrimination training, but now their location was irrelevant to the receipt of

reward. Instead, rats needed to recognize that this old rule (follow the cue light) no longer worked, and that pressing of the

previously less-preferred lever, regardless of cue light position, was now the correct strategy for obtaining reward. Trials

continued until rats performed 10 consecutive correct responses, or after 150 trials had occurred. Errors during set-shift

were categorized into two subtypes as in (Floresco et al. 2008): perseverative errors, where rats responded on the incorrect

lever when the previously-relevant visual cue was illuminated above it, and never-reinforced (or non-perseverative) errors,

where rats pressed the incorrect lever despite when the visual cue was illuminated above the correct lever.

Probabilistic Discounting

In this task rats chose between a lever that always delivers a small (1 pellet) "certain" reward, or another that delivers a

large (4 pellet) "risky" reward, delivered at various probabilities throughout the 52.5 min session (**Fig. 2a**). The probability

of receiving 4 pellets upon pressing the risky lever decreases in each of five sequential choice blocks during each session

(100%, 50%, 25%, 12.5%, 6.25%). Each block began with 8 forced trials (35s apart), in which the houselight was illuminated

and 3s later, one of the two levers extended one at a time for 10s (4 trials for each lever, randomized in pairs). This permitted

rats to sample the probability of reward receipt upon pressing of the risky lever in the upcoming certain/risky lever choice

phase of the block. During the subsequent choice phase of each block, both levers were extended, and after rats pressed

one of them, both levers retracted. Certain lever presses always yielded 1 tasty banana pellet, and risky lever choices

delivered 4 or 0 banana pellets, at a probability that varied across blocks. Failure to respond on a lever within 10s on any

trial resulted in lever(s) retraction and extinguishing of the house light, and the trial was scored as an omission. Animals

were trained on this task for 21 days, at which point choice behavior had stabilized across groups. For rats tested with systemic amphetamine during probabilistic discounting, training on this task commenced following strategy set shifting training and testing described above. Rats in chemogenetic experiments did not undergo visual cue discrimination or set shifting tests. Instead, these rats received pellet habituation, magazine training, initial FR1 training, and retractable lever training prior to training on the probabilistic discounting task.

For rats that received acute amphetamine challenges (0, 0.25, 0.5 mg/kg IP) on risky decision making, 3 counterbalanced tests were conducted on separate days, with re-training between tests to reestablish stable baseline responding. When rats in chemogenetic experiments achieved stable baseline responding, they similarly received 4 counterbalanced tests, 2 with CNO and 2 with VEH (data from both tests combined for analysis as described above), all injected IP 30 min prior to probabilistic discounting testing, with retraining between tests. After completing these systemic CNO/VEH tests, rats were implanted with mPFC cannulae, allowed to recover, and re-stabilized on discounting performance for 2+ days. They were then tested on the discounting task in 4 additional counterbalanced tests with intra-mPFC microinjections of CNO (1mM/0.5 µl; 2 tests) and VEH (2 tests).

Chemogenetic Methods

Virus Surgery

Rats were anesthetized with ketamine (56.5 mg/kg) and xylazine (8.7 mg/kg) and given meloxicam (1.0 mg/kg) for pain prophylaxis. An AAV2 vector containing a Cre-dependent, mCherry-tagged hM3Dq excitatory DREADD (hSyn-DIO-hM3Dq-mCherry; titer: 6 x 10^{12} vg/ml; Addgene catalog #: 44361) was injected bilaterally into VTA (relative to bregma (mm): AP: -5.5, ML: ±0.8, DV: -8.1; 0.75 µL/hemisphere) using a Picospritzer and glass micropipette (Martinez et al. 2023). Injections occurred steadily over 1 min, and the pipette was left in place for 5 min after injection to limit spread. Both TH:Cre and WT rats were injected with the active hM3Dq DREADD virus. Colocalization of mCherry expression to TH+ VTA neurons was verified in each TH:Cre rat, and lack of mCherry expression was confirmed in each WT rat. At least 3 weeks elapsed between virus injections and the first CNO administration. In a second surgery held following behavioral training and systemic CNO tests (to maximize accuracy of placement) guide cannulae (22 ga, 2MM, Plastics One) were implanted bilaterally in the mPFC (relative to bregma (mm): AP: 2.7, ML: ±1, DV: -3.1) to allow intra-mPFC CNO injection upon VTA dopamine neuron axon terminals, and occluded with steel stylets between tests.

Histological Validation

Following behavioral testing, chemogenetic rats were perfused with chilled 0.9% saline and 4% paraformaldehyde, brains

were cryoprotected in 20% sucrose-azide, and they were sectioned coronally at 40 μm. VTA virus expression was amplified

with mCherry immunohistochemistry, and expression verified to be in dopamine neurons via co-staining of tyrosine

hydroxylase. VTA sections were then blocked in 3% normal donkey serum PBST, tissue was incubated overnight at room

temperature in rabbit anti-DSred (Clontech; 1:2500) and mouse anti-TH (Immunostar; 1:2000). After washing, sections were

incubated in dark at room temperature for 4 hours in AlexaFluor-donkey anti-Rabbit 594 and donkey anti-Mouse 488

(Thermo Fisher Scientific; 1:500). Sections were mounted and cover slipped (Fluoromount; Thermo FisherScientific),

mCherry/TH expression was imaged at 10X on a Leica DM4000 epifluorescent scope, and expression or lack thereof in

VTA was verified in each animal. To verify cannula placement within mPFC, sections were nissl stained with cresyl violet,

and cannula tracks were mapped using a rat brain atlas (Paxinos and Watson 2006).

Data Analysis

Effects of adolescent treatment (AdoTX: THC or VEH) and Sex (M or F) on learning and cognition employed 2 x 2 ANOVA

and Sidak posthoc tests. Effects of amphetamine (0 x 0.25 x 0.5mg/kg) on probabilistic discounting were tested with

repeated measures ANOVA, and interactions of these acute treatements with AdoTX and Sex were examined by adding

these between subjects variables to multivariate General Linear Model ANOVAs. Simple-main effects analyses conducted

after observing statiically significant interactions included one-way ANOVA models using the appropriate error term from

the overal multifactor analyses. Effects of CNO versus VEH were treated as within-subjects variable, while Genotype

(TH:Cre x WT), AdoTX, and sex were treated as between subjects variables in multivariate General Linear Model ANOVAs.

Prior to testing, rats were verified to display stable patterns of risky choice by examining discounting performance over at

least two consecutive prior training days. Rats of both sexes were tested in chemogenetic experiments, but sample sizes

for each sex were insuffecient to allow formal analysis of this variable. Six rats (n = 2 males, 4 females; n = 1 VEH, 5 THC)

were excluded from set shifting analyses for failure to meet training critera for instrumental or visual cue rule performance,

and nine rats (n = 9 females; n = 2 VEH, 7 THC) tested with amphetamine were excluded for failing to achieve stable

performance on the probabilistic discounting task. Nine rats (n = 3 males, 6 females; n = 4 VEH, 5 THC) were excluded

from the DREADD experiments for failure to stabilize on the probabilistic discounting, death, or inability to confirm virus

expression.

Results

Effects of Adolescent THC on Strategy Set Shifting in Each Sex

Initial Instrumental Training: AdoTHC did not affect acquisition of instrumental food seeking behavior during initial training

(no main effect of AdoTX: $Fs_{(1, 45)} < 0.396$, $ps > 0.532$), and acquisition was similar in both sexes (No main effect of sex:

$Fs_{(1, 45)} < 3.604$, $ps > 0.064$; no Sex x AdoTX interaction: ($Fs_{(1, 45)} < 1.122$, $ps > 0.295$).

Visual Cue Discrimination: Rats were next trained to press whichever lever that had a cue light illuminated above it. AdoTHC

exposed rats were more likely to learn the visual cue rule in one session rather than two, relative to AdoVEH rats (X^2: $p =$

0.039). This was driven by AdoTHC females, as evidenced by these animals requiring fewer trials and making fewer errors

to reach criterion performance of 10 consecutive correct choices (**Fig. 1c-g**, AdoTX x Sex interaction; trials to criterion: $F_{(1,44)}$

$= 5.308$, $p = 0.026$; errors to criterion: $F_{(1,44)} = 5.032$, $p = 0.03$; Šídák's posthoc for trials $p = 0.022$, errors $p = 0.024$). In

addition, relative to controls, AdoTHC females were slower to respond ($F_{(1, 44)} = 4.546$, $p = 0.039$) but also made fewer

omissions: ($F_{(1, 44)} = 8.460$, $p = 0.006$). In contrast, no differences were observed on these measures between AdoTHC

males and controls (trials to criterion: $p = 0.873$; errors: $p = 0.892$; latency: $p = 0.684$, omissions: $p = 0.719$). More generally,

we also saw some sex-differences on performance measures irrespective of treatment, as females were slower to respond

(Sex: $F_{(1,44)} = 13.02$, $p < 0.001$) and omitted more trials ($F_{(1,44)} = 5.602$, $p = 0.022$) compared to males.

Strategy Set Shifting: Following training on the initial visual cue discrimination, rats were then trained to use an egocentric

spatial rule (always press the left/right lever regardless of cue location), and tested in a single-session (Fig. 1i-m; Brady and

Floresco 2015). AdoTHC had few effects on the ability to learn this new response rule with no change in either sex seen

on the number of trials to reach criterion on the new rule (No main effect of AdoTX: $F_{(1,44)} = 0.416$, $p = 0.522$; or Sex: $F_{(1,44)}$

$= 0.011$, $p = 0.916$; or interaction: $F_{(1,44)} = 0.796$, $p = 0.377$). Total errors to criterion were also unaffected (AdoTX: $F_{(1,44)} =$

0.971, $p = 0.33$; Sex: $F_{(1,44)} = 0.323$, $p = 0.573$; AdoTX x Sex interaction: $F_{(1,44)} = 0.192$, $p = 0.664$), as were errors of either

a perseverative or never-reinforced subtype (AdoTX: $F_{(1,44)} = 1.556$, $p = 0.219$; Sex: $F_{(1,44)} = 0.134$, $p = 0.716$; AdoTX x Sex:

$F_{(1,44)} = 0.583 \times 10^3$, $p = 0.981$). Additionally, we saw no effect of AdoTX on response latency or on omissions (AdoTX: Fs

$_{(1,44)} < 2.620$, $ps > 0.113$; AdoTX x Sex: Fs $_{(1,44)} < 0.006$, $ps > 0.937$). Again, we saw that females took longer to respond

(Sex: $F_{(1,44)} = 7.523$, $p = 0.009$) and omitted more trials ($F_{(1,44)} = 6.201$, $p = 0.017$) compared to males.

Adolescent THC History Improves Learning in Females

Adolescent THC History Does Not Affect Cognitive Flexibility

Fig. 1 Adolescent THC history selectively impacts adulthood learning and cognition a) Experimental timeline. **b)** Schematic of the visual cue discrimination task. **c)** AdoTHC rats (green wedges) were more likely than AdoVEH rats (black wedges) to acquire the visual cue discrimination task to criterion in only one training session (unfilled wedges), rather than requiring 2 sessions to acquire (filled wedges). Both sexes showed similar patterns. **d, e)** AdoTHC females took fewer trials to meet criterion, and made fewer errors when learning visual cue discrimination than AdoVeh females, no such effects were seen in males. **f, g)** AdoTHC females took longer to respond, and made fewer errors than AdoVEH females during cue discrimination training, without effects in males. **h)** Schematic of the subsequently tested strategy set-shifting task. **i)** AdoTHC did not alter the number of trials to learn the new rule to criterion in either sex. Likewise, AdoTHC did not alter in either sex **j)** the number of errors, **k)** the types of errors, **l)** latency to respond, or **m)** omitted trials during set shifting training. Individual rats shown as grey dots in each graph: AdoVEH (n = 14 males, 11 females), AdoTHC (n = 13 males, 10 females). X^2 $p^* < 0.05$ and repeated measure two-way ANOVA, Sidak post hoc: $p^* < 0.05$. Data presented as mean + SEM.

Effects of Adolescent THC on Probabilistic Discounting in Each Sex:

Acquisition of Probabilistic Discounting: The same rats were next trained on a probabilistic discounting task for 21 days. All rats had acquired stable performance by the last three days of training (no block X day interaction: $F_{(5.498,241.89)}$ = 1.695, p = 0.130). No effect of AdoTX ($F_{(1, 44)}$ = 0.817, p = 0.37), Sex ($F_{(1,44)}$ = 0.116, p = 0.735), or AdoTX x Sex interactions ($F_{(1, 44)}$ = 0.585, p = 0.448) were found, suggesting that rats of both sexes and AdoTX histories acquired the discounting task at similar rates.

Fig. 2 AdoTHC history does not affect baseline probabilistic discounting a) Schematic of the probabilistic discounting task. Sessions consisted of 5 training blocks, in which 2 levers are presented, and rats must choose either a small reward lever that always delivers one pellet, or a large reward-delivering lever which becomes increasingly unlikely to deliver any reward as the session progresses. **b)** Data is shown for males and females, indicating that AdoTHC rats did not differ from their AdoVEH counterparts in probabilistic discounting, with both shifting from nearly exclusively preferring the large reward lever, but appropriately shifting away from it as it became less likely to deliver reward. AdoVEH (n = 14 males, 11 females), AdoTHC (n = 15 males, 8 females). Data represented as average within each probability block from three consecutive days of stable performance. Mean $\pm$ SEM.

Stable Probabilistic Discounting: Following training, all rats exhibited stable and comparable performance on the task by the last three days, with decreasing choice of the high-reward lever as delivery of this reward became increasingly unlikely or "risky" (main effect of Block: $F_{(4,176)}$ = 75.01, p < 0.001). No effect of AdoTX ($F_{(1,44)}$ = 0.854, p = 0.361), Sex ($F_{(1,44)}$ = 0.103, p = 0.75), or AdoTX x Sex interactions ($F_{(1, 44)}$ = 0.616, p = 0.437) were found, suggesting that rats of both sexes and AdoTX histories displayed comparable levels of risky choice across blocks (**Fig. 2b**). Likewise, neither AdoTX nor Sex affected win stay or lose shift choice strategies (no main effect of AdoTX: $Fs_{(1,44)}$ < 1.642, ps > 0.207; no main effect of Sex $Fs_{(1,44)}$ < 0.549, ps > 0.463; no AdoTX x Sex $Fs_{(1,44)}$ < 1.51, ps > 0.226; data not shown).

With respect to other performance measures, AdoTX did not affect choice latencies (no AdoTx: $F_{(1, 44)}$ = 0.161, p = 0.69; no AdoTX X Sex interaction: $F_{(1, 44)}$ = 0.1.294, p = 0.262). Additionally, we found that AdoVEH rats omitted more on the "riskier" probability blocks compared to AdoTHC animals (AdoTX x Sex x Block: $F_{(4, 176)}$ = 3.949, p = 0.004, AdoTX x Block: $F_{(4, 176)}$ = 4.391, p = 0.002; no AdoTx: $F_{(1, 44)}$ = 0.939, p = 0.338), an effect that was particularly apparent in females (AdoTX x Block: $F_{(4, 68)}$ = 3.188, p = 0.018; data not shown). Overall, we saw that females took longer to respond than males (Sex: $F_{(1,44)}$ =

40.03, $p < 0.0001$) and omitted more trials ($F_{(1,44)} = 63.83$, $p < 0.0001$), especially in low-probability training blocks at the end of the session (Sex x Block: Latency: $F_{(4,176)} = 4.572$, $p = 0.0015$; Omissions: $F_{(4,176)} = 26.57$, $p < 0.0001$).

Fig. 3 Amphetamine-induced 'risky' responding is potentiated after adolescent THC. a) Data from saline (black line/bar), low dose (0.25mg/kg; pink line/bar) and high dose (0.5mg/kg; red line/bar) amphetamine tests in each sex are shown. When performance patterns were further interrogated, we found that high dose of AMPH in Males increased inflexibility across both AdoTX groups, however in **b)** females, the high dose of AMPH in AdoVEH rats reduced risky responding at high probability blocks not seen in AdoTHC females. **c)** AMPH did not alter win-stay,but reduced lose-shift in AdoTX males. **d)** In AdoVEH females, AMPH decreased win-stay and increased lose-shift, while in AdoTHC AMPH had no effect on win-stay but decreased lose-shift. AMPH = amphetamine. AdoVEH ($n = 14$ males, 10 females), AdoTHC ($n = 15$ males, 6 females). Repeated measure three-way ANOVA; Sidak post hoc: $p* < 0.05$, $p** < 0.01$. Data represented as mean + SEM, individual animals shown as grey dots.

Effects of Acute Amphetamine on Probabilistic Discounting: As previously reported (St. Onge et al. 2010), amphetamine increased choice of the large/risky option in a dose dependent manner when analyzed across both groups and sexes (**Fig. 3a**, amphetamine x Block interaction: $F_{(8,328)} = 10.02$, $p < 0.0001$; main effect of amphetamine: $F_{(2,82)} = 9.88$, $p < 0.0001$). However, the effects of different doses of amphetamine varied as a function of AdoTX treatment and sex. Analysis of the choice data also revealed significant amphetamine x AdoTX ($F_{(2,82)} = 3.33$, $p = 0.041$) and amphetamine x AdoTX x Sex interactions ($F_{(2,82)} = 3.20$, $p = 0.045$). Partitioning this latter interaction by AdoTX group revealed that for control rats, amphetamine exerted different effects on choice in males vs females (amphetamine x Sex: $F_{(2,44)} = 3.73$, $p = 0.032$). Post-hoc comparisons showed that, in males, amphetamine increased risky choice following treatment with the 0.5 mg/kg dose ($p < 0.016$) but not the lower, 0.25 mg/kg dose ($p = 0.19$). In contrast, the 0.5 mg/kg dose had more deleterious effects on choice in females, reducing risky choice in the high probability blocks and increasing it in the lower ones, while the 0.25 mg/kg dose induced a modest increase in risky choice in the latter block. This yielded an overall lack of effect of amphetamine in control females (**Fig. 3b**, amphetamine: $F_{(2,18)} = 0.568$, $p = 0.57$). On the other hand, amphetamine induced a more reliable and pronounced increases in risky choice in both males and females AdoTHC animals, with analysis of

these data producing significant main effects of amphetamine treatment (**Fig. 3a,b**, $F_{(2,38)}$ = 12.07, $p < 0.0001$) in the absence of an interaction with the sex factor ($F_{(2,38)}$ = 1.56, $p = 0.22$). When collapsed across sex, post hoc comparisons showed that both the 0.25 and 0.5 mg/kg dose increase risky choice (both ps < 0.05), although inspection of Fig. 3b indicates that the effect of the 0.25 mg/kg dose was driven primarily by females. From these data, we conclude that AdoTHC treatment makes rats more sensitive to the ability of amphetamine to increase risky choice, and this effect appears to be more prominent in females.

Subsequent analyses examined how amphetamine alters sensitivity to recent rewarded or non-rewarded choices by comparing win-stay and lose shift ratios. Analysis of the win-stay data yielded a significant amphetamine x AdoTX x Sex interaction (**Fig. 3c,d**, $F_{(2,82)}$ = 4.34, $p = 0.016$). This was driven by the fact that in control rats, the 0.5 mg/kg dose resulted in lower win-stay values in females vs males ($p < 0.01$), although neither group showed significant changes in these values relative to saline (both Fs < 2.5, both ps > 0.10). Win-stay behavior was unaltered in AdoTHC rats (all Fs < 2.4, all p > 0.10). In contrast, amphetamine had more pronounced effects on sensitivity to reward omissions, as indexed by changes in lose shift behavior. The analyses here revealed a significant amphetamine x AdoTX interaction ($F_{(2,38)}$ = 4.84, $p = 0.013$) and a three-way interaction with the sex factor ($F_{(2,38)}$ = 5.15, $p = 0.007$). In controls, amphetamine reduced lose shift behavior in males, but actually increased it in females (**Fig. 3c,d**, amphetamine x Sex: $F_{(2,44)}$ = 3.86, $p = 0.028$, whereas in AdoTHC rats, these treatments uniformly reduced lose-shift behavior across sexes (amphetamine: $F_{(2,38)}$ = 8.87, $p < 0.001$; amphetamine x Sex: $F_{(2,38)}$ = 2.61, $p = 0.08$). From these data, we conclude that AdoTHC treatment makes rats more sensitive to the ability of amphetamine to increase risky choice and reduce sensitivity to losses, and this effect appears to be more prominent in females.

With respect to other performance measures, amphetamine increased choice latency and number of omissions across all probability blocks (amphetamine: $Fs_{(2,82)}$ < 15.19, ps < 0.001, no amphetamine x Block interaction: $Fs_{(8,328)}$ < 0.98, ps > 0.454). Analysis of the latency data revealed a significant amphetamine x AdoTX x Block ($F_{(8,328)}$ = 2.27, $p = 0.023$) and amphetamine x AdoTX x Sex x Block interaction ($F_{(2,82)}$ = 2.67, $p = 0.008$). Partitioning this latter interaction by AdoTX group revealed that in AdoVEH rats, females took longer to respond than males across the session (Sex x Block: $F_{(4,88)}$ = 3.02, $p = 0.022$, Sex: $F_{(1,22)}$ = 44.13, $p < 0.001$). In AdoTHC rat, females took longer to respond compared to males (Sex: $F_{(1,19)}$ = 21.44, $p < 0.001$; no Sex X Block: $F_{(4,76)}$ = 2.41, $p = 0.057$). Further analysis of the omission data revealed that overall females omitted more on trials compared to males (Sex x Block: $F_{(4,164)}$ = 12.93, $p < 0.001$; Sex: $Fs_{(1,41)}$ < 71.39, $p < 0.001$).

Chemogenetic Dopamine Neuron Stimulation During Probabilistic Discounting

The above experiments showed that AdoTHC enhances visual cue learning in females but had few other effects on cognitive flexibility or probabilistic discounting under basal conditions. However, when treated with the monoamine-enhancing drug amphetamine, we found evidence for stronger enhancement of "risky" responding in AdoTHC rats, relative to AdoVEH controls. We therefore next asked whether this effect relates to changes in the functions of VTA dopamine neurons in particular, by using Gq-coupled DREADDs to acutely stimulate VTA dopamine neurons or VTA dopamine neuron projections to mPFC in rats with both AdoTX histories.

Initial Training: Rats in this experiment did not undergo strategy set shifting training prior to probabilistic discounting training, so we confirmed that AdoTX again did not affect initial acquisition of instrumental food seeking behavior during initial training (no main effect of AdoTX: $Fs_{(1, 46)} < 0.78$, $ps > 0.381$). We also confirmed that genotype (Geno: TH:Cre+ or WT littermate) did not alter instrumental training ($Fs_{(1,46)} < 2.55$, $ps > 0.117$; No AdoTX x Geno interaction: $Fs_{(1,46)} < 0.629$, $ps > 0.432$).

Probabilistic Discounting Training: Likewise, all included rats (TH:Cre and WT) successfully learned the discounting task (Main effect of Block: $F_{(4,184)} = 129.26$, $p < 0.001$, no main effects of, or interactions involving AdoTX or Geno: $Fs_{(1,46)} < 0.884$, $ps > 0.352$). Additionally, AdoTX nor genotype affected latency, omissions, or win-stay/lose-shift choice strategies (no main effects of, or interactions involving AdoTX or Geno: $Fs_{(1,46)} < 3.34$, $ps > 0.074$).

Impact of Acutely Activating VTA Dopamine Neurons on Probabilistic Discounting

Despite the robust changes in locomotion and reward seeking that is seen following chemogenetic VTA dopamine neuron manipulations in TH:Cre rats (Boekhoudt et al. 2016; Boekhoudt et al. 2018; Halbout et al. 2019; Lawson et al. 2023; Mahler et al. 2019; Runegaard et al. 2018), we found few effects of VTA dopamine neuron stimulation on the probabilistic discounting task. First, we looked at VTA dopamine neuron stimulation in only TH:Cre rats. All rats displayed normal discounting profiles (TH:Cre+: main effect of Block: $F_{(4, 84)} = 105.934$, $p < 0.001$), and as in the prior experiment, no effect of AdoTX alone was seen on discounting (AdoTX x Block: $F_{(4, 84)} = 0.908$, $p = 0.463$). Moreover, when we tested the effects of systemic CNO to activate DA neurons, we did not observe any effects of this treatment in either control or AdoTX groups (**Fig. 4e**, no main effect of, or interactions involving AdoTX or CNO: $Fs < 2.27$, $ps > 0.068$). Additionally, we did not see any effects of CNO, AdoTX, nor interactions therein on choice latency, omissions, or win-stay/lose-shift choice strategy ($Fs_{(1,21)} < 3.00$, $ps > 0.098$; data not shown). These findings are inconsistent with our hypothesis that increased excitability of mesolimbic dopamine neurons would be sufficient to recapitulate the ability of amphetamine to increase risky/perseverative responding in either control or in AdoTHC-experienced rats.

Fig. 4 Stimulation of VTA dopamine neurons, or VTA dopamine projections to mPFC does not affect probabilistic discounting a) Bilateral Cre-dependent hM3Dq DREADD AAV injections were made in VTA of TH:Cre rats, and of wildtype (WT) littermates. **b)** Example hM3Dq DREADD expression (red) is localized to tyrosine hydroxylase+ (TH; green) neurons within VTA (yellow=merge). Scale bar, 300 µm. **c)** For pathway-specific stimulation of VTA dopamine projections to mPFC, Cre-dependent hM3Dq DREADDs were injected into VTA as in Experiment 1, and cannulae targeting mPFC allowed CNO microinjection (1mM, 0.5 µl) upon DREADD-expressing dopamine neuron axons in this pathway. **d)** Cannula placements of each rat in pathway stimulation Experiment 2 is shown. **e)** Neither VTA dopamine neuron stimulation induced by systemic CNO in TH:Cre rats, nor **f)** stimulation of the VTA dopamine projection to mPFC induced by mPFC CNO microinjections in TH:Cre rats robustly altered probabilistic discounting in AdoTHC or AdoVEH rats. Likewise, lower panels of **e,f)** indicate that CNO did not have robust effects on WT rats without DREADDs. Data is represented as average % choice of the large reward lever in each probability block across the two CNO, and two VEH tests conducted in each rat, mean±SEM, $p* < 0.05$. TH:Cre+: VTA dopamine stimulation; AdoVEH ($n = 8M, 3F$) AdoTHC ($n = 8M, 4F$); VTA dopamine to mPFC: AdoVEH ($n = 10M, 4F$) AdoTHC ($n = 9M, 5F$). Wildtype: VTA dopamine stimulation; AdoVEH ($n = 5M, 3F$) AdoTHC ($n = 5M$); VTA dopamine to mPFC: AdoVEH ($n = 9M, 4F$) AdoTHC ($n = 8M$).

Impact of Selectively Activating VTA Dopamine Projections to mPFC

We next asked whether selectively stimulating VTA dopamine neuron projections to mPFC would recapitulate the

potentiation of amphetamine effects in AdoTHC rats. We did so by locally applying CNO upon DREADD-expressing axons

of VTA dopamine neurons in mPFC. We have shown that this pathway-specific stimulation approach is capable of

potentiating both axonal dopamine release and motivated reward seeking (Halbout et al. 2019; Mahler and Aston-Jones

2012; Mahler et al. 2019). We found few effects of this manipulation on probabilistic discounting. On the probabilistic

discounting task, "risky" responding across the session varied across CNO and AdoTX (**Fig. 4**; AdoTX x CNO x Block

interaction: $F_{(4,104)} = 2.991$, $p = 0.022$; no main effect of AdoTX: $F_{(1,26)} = 0.153$, $p = 0.699$; no main effect of CNO: $F_{(1,26)} =$

0.000 p = 0.997), in AdoVEH animals there were no measurable effects of or interactions with CNO (all Fs < 1.80, ps >

0.143). However, in AdoTHC rats CNO modestly reduced risky responding at the 100% probability block followed by an

increase at the 50% probability block (CNO x Block: $F_{(4,52)}$ = 2.99, p = 0.027). Rats performed similarly on choice latency,

omissions, and win-stay/lose-shift regardless of AdoTX, CNO, or interactions therein with block (All Fs < 1.996, ps > 0.101).

Minimal DREADD-independent Effects of CNO

Neither systemic nor intra-mPFC CNO had major behavioral effects in WT rats lacking DREADD expression (**Fig. 4e,f**).

Systemic CNO in WT rats seemed to promote a more risk-prone phenotype, based on increased preference for the risky

lever across all blocks (Main effect of CNO in WT rats: $F_{(1, 11)}$ = 9.424, p = 0.011; no interactions of CNO, AdoTX, or Block:

$Fs_{(4, 44)}$ < 2.04, ps > 0.105), but did not affect response latency, omissions, or win-stay/lose-shift choice strategies ($Fs_{(1,11)}$ <

2.962, ps > 0.113). Intra-mPFC CNO in WT rats altered risk responding such that "risky" choice was decreased in AdoTHC

rats, whereas it was increased in AdoVEH animals (AdoTX x CNO: $F_{(1,16)}$ = 5.404, p = 0.034). These effects were driven

by changes in reward sensitivity, as CNO differentially affected win-stay behavior in the two groups ($F_{(1,160}$ = 5.781, p =

0.029; no main effect of CNO $F_{(1,16)}$ = 0.726, p = 0.407). Thus, CNO in AdoTHC WT rats were less likely to follow a risky win

with another risky choice, while CNO had the opposite effects in in AdoVEH WT (data not shown). Additionally, we saw that

AdoTHC animals had lower risky responding in the higher probability blocks, that then increased in the latter probability

blocks (AdoTX x Block: $F_{(4,64)}$ = 2.524, p = 0.049). We did not see any effect of AdoTX or CNO on lose-shift strategies (CNO

x AdoTX: $F_{(1,16)}$ = 0.667, p = 0.426; no main effect of CNO $F_{(1,16)}$ = 1.712, p = 0.209) or response latency and omissions

($Fs_{(1,16)}$ < 3.742, ps > 0.071; data not shown).

Discussion

Here we show that administration of a well-characterized, human-relevant dose of THC (Ruiz et al. 2021a; Torrens et al.

2022; Torrens et al. 2020) during adolescence has subtle effects on behavioral tests of instrumental learning, without

measurably altering performance on mPFC dopamine-dependent strategy set shifting or risk/reward decision making

assessed with a probabilistic discounting task. However, when monoamine signaling was acutely enhanced with systemic

amphetamine, an underlying effect of AdoTHC history was revealed. Relative to AdoVEH controls, amphetamine in AdoTHC

rats were more sensitive to the ability of lower doses of amphetamine to increase perseverative responding for a large

reward option when this choice was unlikely to result in reward. However, this potentiation of amphetamine effects in

AdoTHC rats was not recapitulated by more specific chemogenetic stimulation of VTA dopamine neurons, or of their

projections to mPFC in particular. This could suggest non-VTA dopaminergic mechanisms underlying potentiation of this

cognitive effect of amphetamine in AdoTHC rats, suggesting potential impacts of THC on adolescent development of other systems upon which amphetamine acts. Alternatively, the enhanced effect of amphetamine reported here may be driven by alterations at the level of the dopamine terminal rather than changes in dopamine cell excitability, as dopamine receptor antagonism is able to block this amphetamine-induced risky responding (St. Onge and Floresco 2009). Further, we thoroughly characterize sex differences in decision-making across these behavioral tasks, some of which mediate the persistent impacts of AdoTHC on behavior. Results open new directions for investigating long-term impacts of AdoTHC on non-VTA dopaminergic modulation of cognition, and may inform associational studies of the long-term impacts of adolescent cannabis use in humans.

Adolescent THC Exposure Enhances Initial Discrimination Learning

AdoTHC rats learned a visually cued instrumental response rule quicker than AdoVEH-treated controls, and this effect was most robust in females. This finding adds to the growing literature on potentially pro-learning/cognitive effects of AdoTHC (Bilkei-Gorzo et al. 2017; Hernandez et al. 2021; Sarne 2019; Stringfield and Torregrossa 2021b). This said, previous studies using other adolescent cannabinoid exposure models have not found analogous increases in initial rule discrimination learning (Freels et al. 2024; Gomes et al. 2015; Hernandez et al. 2021), though this might be due to the fact that few studies included female subjects, and the THC administration protocols employed were quite different.

We found no clear effects of AdoTHC across our set shifting performance metrics, including trials to criterion, number of errors made, or error types on the set shift day. These findings are consistent with others that found no major AdoTHC-induced changes in cognitive flexibility, as quantified in with different set shifting tasks, including the intra/extra dimensional "digging" task and an operant-based task similar to the one used here (Gomes et al. 2015; Hernandez et al. 2021; Poulia et al. 2021). However, in a study utilizing the same strategy set-shifting task as used here, females exposed to self-administered cannabis extract vapor during adolescence took longer to learn the new rule, and made more errors on the set-shift day, though this deficit was not seen after experimenter-administered cannabis vapor (Freels et al. 2024). It is presently unclear whether differences in patterns of results reflect differences in route of THC administration or dose, the specific timing of THC exposure during adolescence, or other experimental details. As such, this remains an important topic for future research intended to probe how adolescent cannabinoid exposure may alter executive functioning.

Though untested here, it is possible that AdoTHC altered other related cognitive processes such as transitioning between tasks, mental sets, or rule structures, which depend upon more lateral PFC subregions such as orbitofrontal cortex (OFC; Birrell and Brown 2000; Floresco et al. 2008; McAlonan and Brown 2003). For example, adolescent pubertal administration of the potent CBR agonist WIN 55,212 alters OFC-dependent reversal learning in and attentional set-shifting task (Gomes

et al. 2015), though adolescent vaporized cannabis or cannabis smoke did not impact this same type of reversal learning (Freels et al. 2024; Hernandez et al. 2021). Again, discrepancies between studies may reflect differences in effects of cannabinoid drugs, doses, exposure timing, and washout period; further underscoring the need for a consistent, rationally-designed AdoTHC exposure model in the field—we argue that the model used here is the best-characterized to date in the field (Halbout et al. 2023; Lee et al. 2022; Lee et al. 2024; Lin et al. 2023; Ruiz et al. 2021a; Torrens et al. 2022; Torrens et al. 2020). This said, the possibility that OFC is even more sensitive to disruption by AdoTHC than mPFC should be directly tested in future studies.

Adolescent THC Does Not Alter Basal Probabilistic Discounting

In the mPFC-dependent probabilistic discounting task (St. Onge and Floresco 2010), rats choose between a small reward that is always delivered when chosen (1 palatable banana pellet), and a larger reward that becomes increasingly unlikely to be delivered over the course of the ~1h session (providing either 4 or 0 pellets). Efficient performance on this task demands evaluation of both risk and opportunity, and is dependent on intact functioning of both the mPFC and mesocortical and mesoaccumbens dopamine transmission (Jenni et al. 2017; St. Onge et al. 2012; St. Onge et al. 2010; Stopper et al. 2013). Impaired PFC activity results in deficits in adjusting choice in response to changes in reward probabilities, loss assessment, and a diminished ability to appropriately compare and favor larger rewards even when the probability of receiving them is higher (Bercovici et al. 2023; St. Onge and Floresco 2010). We found no major impacts of AdoTHC history on acquisition of, or stable performance on this task, as measured by choices of the 'risky' lever, choice strategies following rewarded vs unrewarded trials, latency to decide, and decisions to omit trials. One prior study (Jacobs-Brichford et al. 2019) found that administration of WIN 55,212 in both sexes during adolescence elevated preference for the 'risky' lever at lower reward probabilities, which we did not see in either sex as a result of our AdoTHC exposure model. Though several other differences exist between this study and the present one, we note that several prior reports have also shown more severe lasting effects of synthetic CBR agonists versus THC when administered in adolescents (Higuera-Matas et al. 2015; Renard et al. 2014; Stringfield and Torregrossa 2021a).

Adolescent THC History Potentiates Amphetamine Effects on Probabilistic Discounting

Dopamine markedly influences mPFC-dependent cognition (Floresco and Magyar 2006; Goldman-Rakic 1995; Goto et al. 2007; Seamans and Yang 2004), including in the probabilistic discounting task employed here (Floresco and Whelan 2009; Islas-Preciado et al. 2020; Jenni et al. 2017; St. Onge et al. 2012; St. Onge et al. 2010; St. Onge and Floresco 2009). In the present study we replicated prior findings that pharmacologically challenging monoamine systems with amphetamine increased perseveration of responding for a large reward in both males and females (Islas-Preciado et al. 2020; St. Onge

and Floresco 2009). This change was accompanied by a reduction in lose-shift behaviors following 'risky' losses, indicating

that enhancing DA levels attenuates the impact that non-rewarded actions exert over subsequent choice. Interestingly, we

saw that amphetamine led to an overall increase in choice latency, and omitted trials. This may have been driven in part by

increased psychomotor activity induced by amphetamine that may have displaced animals from the levers, thus delaying

choice.

Moreover, we found that this effect of amphetamine was more potent in AdoTHC, relative to AdoVEH rats, especially in

females. At the highest dose of amphetamine, AdoTHC rats chose the 'risky' lever more when probabilities of that reward

were low. These effects were also markedly sex-dependent. In males, amphetamine reduced overall lose-shift behavior. In

females, the highest dose of amphetamine induced a disruptive reduction in win-stay behavior only in AdoVEH females,

whereas it reduced lose-shift behaviors only in AdoTHC females, contributing to the heightened increased 'risky' responding

observed in the AdoTHC groups. These results suggest that although probabilistic discounting under basal conditions is not

altered by AdoTHC, underlying differences were nonetheless revealed upon acute activation of monoaminergic signaling.

Amphetamine blocks and reverses the transporter for dopamine, norepinephrine, serotonin, and changes in one or more of

these systems could be responsible for the potentiated response to amphetamine we see in this task in AdoTHC rats

(especially females).

Effects of Chemogenetic VTA Dopamine Neuron, and mPFC Dopamine Projection Stimulation on Probabilistic

Discounting

In our next experiments, we sought to determine whether, as predicted, changes in VTA dopamine neuron signaling are

responsible for the potentiated effect of amphetamine in AdoTHC rats. We therefore prepared transgenic TH:Cre rats with

excitatory DREADDs, allowing stimulation of either all VTA dopamine neurons, or of dopamine neuron projections to mPFC

in particular. Surprisingly, no major effects of chemogenetically stimulating all VTA dopamine neurons were found on

probabilistic discounting. In this regard, it bears mentioning that pharmacological stimulation of dopamine D2 or D1 receptors

in the nucleus accumbens (a main target of the VTA dopamine projection) either had no effect on risky choice or led to more

optimal decision making, respectively- an effect distinct from that induced by amphetamine (Stopper et al. 2013).

Furthermore, when we stimulated VTA dopamine neuron projections to mPFC, this did not markedly affect probabilistic

discounting in AdoVEH rats, which is similar to a prior finding examining manipulations of VTA dopamine projections to PFC

(Verharen et al. 2018). On the other hand, chemogenetic stimulation of mPFC dopamine terminals in AdoTHC rats

attenuated risky responding during the 100% probability block, but then reduced it in the subsequent 50% block. This effect

loosely resembles that induced by intra-PFC D_2 receptor stimulation on probabilistic discounting, which flattened the

discounting curve (St. Onge et al. 2011). This is consistent with the possibility that AdoTHC exposure causes a slight

enhancement in mesocortical dopamine D_2 receptor signaling—a possibility that should be directly investigated. Moreover,

no other differential effects of DREADD stimulation were seen in AdoTHC versus AdoVEH groups. This result is unlikely to

be due to a lack of efficacy of the chemogenetic manipulations employed, since our prior work and that of others' shows

that such stimulation in TH:Cre rats has pronounced effects on locomotion, reward intake, and motivation (Boekhoudt et al.

2016; Boekhoudt et al. 2018; Halbout et al. 2019; Lawson et al. 2023; Mahler et al. 2019; Runegaard et al. 2018). This may

imply that alterations of VTA and non-VTA dopamine caused by amphetamine might simply be qualitatively distinct from

manipulations of VTA dopamine neuron activity and dopamine release, as were conducted here chemogenetically (Mahler

et al. 2019; Mahler et al. 2014). In this regard, performance of this probabilistic discounting task is associated with

fluctuations in mesocortical and mesoaccumbens dopamine that relate to changes in the amount of rewards received and

adjustments in risky choice biases (St. Onge et al, 2012). Amphetamine would cause robust increases in tonic dopamine

levels, leading to a more static dopamine signal that impedes flexible adjustments in choice biases, whereas chemogenetic

increases in dopamine neuron excitability would be a comparatively more subtle manipulation that could still permit

variations in dopamine signaling over a session. Alternatively, amphetamine-induced changes in risk taking seen in AdoTHC

rats may not solely depend upon VTA dopamine neurons themselves, but may instead depend upon other neural systems

targeted by amphetamine, such as nigrostriatal dopamine projections to the dorsomedial striatum (Schumacher et al. 2021),

other non-mPFC targets of VTA dopamine neurons, or other monoamine systems affected by amphetamine. Regardless,

this intriguing finding refines our understanding of the neural substrates of probability-based decision making and

underscores the need for further exploration to identify the specific mechanisms involved.

Sex Differences

We conducted these studies in female as well as male rats, and found several consistent sex differences, both overall and

in interaction with other experimental variables. Consistent with existing literature (Gargiulo et al. 2022; Islas-Preciado et al.

2020), we saw sex differences in response characteristics during operant set-shifting as well as probabilistic discounting,

with females showing longer deliberation times (response latency), and more omitted trials than males. Strikingly, these

specific sex differences consistently surfaced in the probabilistic discounting task across various experiments and cohorts,

underscoring the robustness of the response latency and trial omission effects in females. Some of these sex differences

may derive from females attaining satiety on the palatable reward quicker than males, but since both omissions and longer

latencies at least partially emerged in females early in sessions during otherwise apparently vigorous reward seeking, this

finding may also reflect different strategies taken by females relative to males (Chen et al. 2021; Orsini and Setlow 2017).

For example, female rats have previously been shown to be more risk-averse than males on average (Orsini et al. 2016),

and may exhibit prolonged response times and higher omission rates due to aversion to the risk of losing a reward, as

demonstrated by a heightened sensitivity to loss (van den Bos et al. 2012). Alternatively, extended choice latencies and

omissions may signify females' tendency to take more time in learning about the probability distribution of the outcomes.

This aligns with both rodent and human findings, indicating that females take longer to develop a preference for the more

advantageous option when learning about probability distributions of reward versus punishment (van den Bos et al. 2013;

van den Bos et al. 2012). Both possibilities warrant further exploration in future.

Limitations

The present report has a number of limitations that should be considered. A single dose of THC (5mg/kg IP) was

administered to adolescents daily from postnatal days 30 to 43, and persistent effects of THC are known to be dose-

dependent (Amal et al. 2010; Freeman-Striegel et al. 2023; Irimia et al. 2015), and dependent upon the specific

developmental stage at which it is experienced (Cha et al. 2006; Gorey et al. 2019; Mokrysz et al. 2016; Murray et al. 2022;

Schramm-Sapyta et al. 2007; Torrens et al. 2022; Torrens et al. 2020). Also relevant to dose, higher blood and brain levels

of the active THC metabolite 11-OH-THC are seen in female relative to male adolescent and adult rats and mice (Baglot et

al. 2021; Craft et al. 2019; Ruiz et al. 2021a; Ruiz et al. 2021b; Torrens et al. 2022; Torrens et al. 2020; Tseng et al. 2004;

Wiley and Burston 2014), and this sex difference is also seen in humans after cannabis use (Arkell et al. 2022; Matheson

et al. 2020; Sholler et al. 2021). This metabolic sex difference may at least in part account for the more robust persistent

effects of adolescent THC in females than males seen here, and in numerous prior reports (Cha et al. 2007; Le et al. 2022;

Rubino and Parolaro 2011; Rubino et al. 2009; Ruiz et al. 2021a; Ruiz et al. 2021b; Tseng et al. 2004). Another limitation

of this report is the lack of data on complementary cognitive tasks, which are dependent upon other cortical regions like

OFC that are notably sensitive to adolescent cannabinoid exposure (Egerton et al. 2005; Gomes et al. 2015; Klugmann et

al. 2011), and which are mediated by ECB-dependent processes that may be particularly sensitive to AdoTHC's long-lasting

impacts.

Summary

In sum, this paper provides novel information about the impact of AdoTHC on the developing brain of males, and especially

of females. Results point to long-lasting changes that influence cue learning, and the ability of acute amphetamine to alter

risky decision making. A key finding is the pervasive nature of sex-differences—both in behavioral strategies employed by

rats in these tasks, and in the severity of long-lasting consequences of AdoTHC.

References

Amal H, Fridman-Rozevich L, Senn R, Strelnikov A, Gafni M, Keren O, Sarne Y (2010) Long-term consequences of a single treatment of mice with an ultra-low dose of Delta9-tetrahydrocannabinol (THC). Behav Brain Res 206: 245-53.

Andersen SL (2003) Trajectories of brain development: point of vulnerability or window of opportunity? Neurosci Biobehav Rev 27: 3-18.

Andreasson S, Allebeck P, Engstrom A, Rydberg U (1987) Cannabis and schizophrenia. A longitudinal study of Swedish conscripts. Lancet 2: 1483-6.

Arkell TR, Kevin RC, Vinckenbosch F, Lintzeris N, Theunissen E, Ramaekers JG, McGregor IS (2022) Sex differences in acute cannabis effects revisited: Results from two randomized, controlled trials. Addict Biol 27: e13125.

Baglot SL, Hume C, Petrie GN, Aukema RJ, Lightfoot SHM, Grace LM, Zhou R, Parker L, Rho JM, Borgland SL, McLaughlin RJ, Brechenmacher L, Hill MN (2021) Pharmacokinetics and central accumulation of delta-9-tetrahydrocannabinol (THC) and its bioactive metabolites are influenced by route of administration and sex in rats. Sci Rep 11: 23990.

Bara A, Ferland JN, Rompala G, Szutorisz H, Hurd YL (2021) Cannabis and synaptic reprogramming of the developing brain. Nat Rev Neurosci 22: 423-438.

Behan AT, Hryniewiecka M, O'Tuathaigh CM, Kinsella A, Cannon M, Karayiorgou M, Gogos JA, Waddington JL, Cotter DR (2012) Chronic adolescent exposure to delta-9-tetrahydrocannabinol in COMT mutant mice: impact on indices of dopaminergic, endocannabinoid and GABAergic pathways. Neuropsychopharmacology 37: 1773-83.

Bercovici DA, Princz-Lebel O, Schumacher JD, Lo VM, Floresco SB (2023) Temporal Dynamics Underlying Prelimbic Prefrontal Cortical Regulation of Action Selection and Outcome Evaluation during Risk/Reward Decision-Making. J Neurosci 43: 1238-1255.

Bilkei-Gorzo A, Albayram O, Draffehn A, Michel K, Piyanova A, Oppenheimer H, Dvir-Ginzberg M, Racz I, Ulas T, Imbeault S, Bab I, Schultze JL, Zimmer A (2017) A chronic low dose of Delta(9)-tetrahydrocannabinol (THC) restores cognitive function in old mice. Nat Med 23: 782-787.

Birrell JM, Brown VJ (2000) Medial frontal cortex mediates perceptual attentional set shifting in the rat. J Neurosci 20: 4320-4.

Bloomfield MAP, Ashok AH, Volkow ND, Howes OD (2016) The effects of Δ9-tetrahydrocannabinol on the dopamine system. Nature 539: 369-377.

Boekhoudt L, Omrani A, Luijendijk MC, Wolterink-Donselaar IG, Wijbrans EC, van der Plasse G, Adan RA (2016) Chemogenetic activation of dopamine neurons in the ventral tegmental area, but not substantia nigra, induces hyperactivity in rats. Eur Neuropsychopharmacol 26: 1784-1793.

Boekhoudt L, Wijbrans EC, Man JHK, Luijendijk MCM, de Jong JW, van der Plasse G, Vanderschuren LJMJ, Adan RAH (2018) Enhancing excitability of dopamine neurons promotes motivational behaviour through increased action initiation. European Neuropsychopharmacology 28: 171-184.

Bossong MG, Mehta MA, van Berckel BNM, Howes OD, Kahn RS, Stokes PRA (2015) Further human evidence for striatal dopamine release induced by administration of a†9-tetrahydrocannabinol (THC): selectivity to limbic striatum. Psychopharmacology 232: 2723-2729.

Brady AM, Floresco SB (2015) Operant procedures for assessing behavioral flexibility in rats. J Vis Exp: e52387.

Casey BJ, Giedd JN, Thomas KM (2000) Structural and functional brain development and its relation to cognitive development. Biol Psychol 54: 241-57.

Cha YM, Jones KH, Kuhn CM, Wilson WA, Swartzwelder HS (2007) Sex differences in the effects of delta9-tetrahydrocannabinol on spatial learning in adolescent and adult rats. Behav Pharmacol 18: 563-9.

Cha YM, White AM, Kuhn CM, Wilson WA, Swartzwelder HS (2006) Differential effects of delta9-THC on learning in adolescent and adult rats. Pharmacol Biochem Behav 83: 448-55.

Chen CS, Ebitz RB, Bindas SR, Redish AD, Hayden BY, Grissom NM (2021) Divergent Strategies for Learning in Males and Females. Curr Biol 31: 39-50 e4.

Chen JP, Paredes W, Li J, Smith D, Lowinson J, Gardner EL (1990) Delta-9-Tetrahydrocannabinol Produces Naloxone-Blockable Enhancement of Presynaptic Basal Dopamine Efflux in Nucleus-Accumbens of Conscious, Freely-Moving Rats as Measured by Intracerebral Microdialysis. Psychopharmacology 102: 156-162.

Craft RM, Britch SC, Buzitis NW, Clowers BH (2019) Age-related differences in Delta(9)-tetrahydrocannabinol-induced antinociception in female and male rats. Exp Clin Psychopharmacol 27: 338-347.

Curran HV, Freeman TP, Mokrysz C, Lewis DA, Morgan CJ, Parsons LH (2016) Keep off the grass? Cannabis, cognition and addiction. Nat Rev Neurosci 17: 293-306.

D'Souza DC, Perry E, MacDougall L, Ammerman Y, Cooper T, Wu YT, Braley G, Gueorguieva R, Krystal JH (2004) The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. Neuropsychopharmacology 29: 1558-72.

Davis KL, Kahn RS, Ko G, Davidson M (1991) Dopamine in schizophrenia: a review and reconceptualization. Am J Psychiatry 148: 1474-86.

De Melo LCS, Cruz AP, Valentim SJR, Marinho AR, Mendonça JB, Nakamura-Palacios EM (2005) Δ9-THC administered into the medial prefrontal cortex disrupts the spatial working memory. Psychopharmacology 183: 54-64.

Egerton A, Brett RR, Pratt JA (2005) Acute delta9-tetrahydrocannabinol-induced deficits in reversal learning: neural correlates of affective inflexibility. Neuropsychopharmacology 30: 1895-905.

Ehrenreich H, Rinn T, Kunert HJ, Moeller MR, Poser W, Schilling L, Gigerenzer G, Hoehe MR (1999) Specific attentional dysfunction in adults following early start of cannabis use. Psychopharmacology (Berl) 142: 295-301.

Ellgren M, Artmann A, Tkalych O, Gupta A, Hansen HS, Hansen SH, Devi LA, Hurd YL (2008) Dynamic changes of the endogenous cannabinoid and opioid mesocorticolimbic systems during adolescence: THC effects. Eur Neuropsychopharmacol 18: 826-34.

Fergusson DM, Horwood LJ, Ridder EM (2005) Tests of causal linkages between cannabis use and psychotic symptoms. Addiction 100: 354-366.

Floresco SB (2013) Prefrontal dopamine and behavioral flexibility: shifting from an "inverted-U" toward a family of functions. Front Neurosci 7: 62.

Floresco SB, Block AE, Tse MT (2008) Inactivation of the medial prefrontal cortex of the rat impairs strategy set-shifting, but not reversal learning, using a novel, automated procedure. Behav Brain Res 190: 85-96.

Floresco SB, Magyar O (2006) Mesocortical dopamine modulation of executive functions: beyond working memory. Psychopharmacology (Berl) 188: 567-85.

Floresco SB, Whelan JM (2009) Perturbations in different forms of cost/benefit decision making induced by repeated amphetamine exposure. Psychopharmacology (Berl) 205: 189-201.

Freels TG, Westbrook SR, Zamberletti E, Kuyat JR, Wright HR, Malena AN, Melville MW, Brown AM, Glodosky NC, Ginder DE, Klappenbach CM, Delevich KM, Rubino T, McLaughlin RJ (2024) Sex Differences in Response-Contingent Cannabis Vapor Administration During Adolescence Mediate Enduring Effects on Behavioral Flexibility and Prefrontal Microglia Activation in Rats. Cannabis Cannabinoid Res.

Freeman-Striegel L, Hamilton J, Kannappan R, Bell T, Robison L, Thanos PK (2023) Chronic Δ9-tetrahydrocannabinol treatment has dose-dependent effects on open field exploratory behavior and [3H] SR141716A receptor binding in the rat brain. Life Sciences: 121825.

French ED, Dillon K, Wu X (1997) Cannabinoids excite dopamine neurons in the ventral tegmentum and substantia nigra. Neuroreport 8: 649-52.

Gardner EL (2002) Addictive potential of cannabinoids: the underlying neurobiology. Chem Phys Lipids 121: 267-90.

Gargiulo AT, Hu J, Ravaglia IC, Hawks A, Li X, Sweasy K, Grafe L (2022) Sex differences in cognitive flexibility are driven by the estrous cycle and stress-dependent. Front Behav Neurosci 16: 958301.

Gessa GL, Melis M, Muntoni AL, Diana M (1998) Cannabinoids activate mesolimbic dopamine neurons by an action on cannabinoid CB1 receptors. Eur J Pharmacol 341: 39-44.

Goldman-Rakic PS (1995) Cellular basis of working memory. Neuron 14: 477-85.

Gomes FV, Guimaraes FS, Grace AA (2015) Effects of Pubertal Cannabinoid Administration on Attentional Set-Shifting and Dopaminergic Hyper-Responsivity in a Developmental Disruption Model of Schizophrenia. International Journal of Neuropsychopharmacology 18.

Gorey C, Kuhns L, Smaragdi E, Kroon E, Cousijn J (2019) Age-related differences in the impact of cannabis use on the brain and cognition: a systematic review. Eur Arch Psychiatry Clin Neurosci 269: 37-58.

Goto Y, Otani S, Grace AA (2007) The Yin and Yang of dopamine release: a new perspective. Neuropharmacology 53: 583-7.

Halbout B, Hutson C, Hua L, Inshishian V, Mahler SV, Ostlund SB (2023) Long-term effects of THC exposure on reward learning and motivated behavior in adolescent and adult male rats. Psychopharmacology (Berl) 240: 1151-1167.

Halbout B, Marshall AT, Azimi A, Liljeholm M, Mahler SV, Wassum KM, Ostlund SB (2019) Mesolimbic dopamine projections mediate cue-motivated reward seeking but not reward retrieval in rats. Elife 8: e43551.

Hambrecht M, Hafner H (2000) Cannabis, vulnerability, and the onset of schizophrenia: an epidemiological perspective. Aust N Z J Psychiatry 34: 468-75.

Heng L, Beverley JA, Steiner H, Tseng KY (2011) Differential developmental trajectories for CB1 cannabinoid receptor expression in limbic/associative and sensorimotor cortical areas. Synapse 65: 278-86.

Henquet C, Murray R, Linszen D, van Os J (2005) The environment and schizophrenia: the role of cannabis use. Schizophr Bull 31: 608-12.

Hernandez CM, Orsini CA, Blaes SL, Bizon JL, Febo M, Bruijnzeel AW, Setlow B (2021) Effects of repeated adolescent exposure to cannabis smoke on cognitive outcomes in adulthood. J Psychopharmacol 35: 848-863.

Higuera-Matas A, Ucha M, Ambrosio E (2015) Long-term consequences of perinatal and adolescent cannabinoid exposure on neural and psychological processes. Neurosci Biobehav Rev 55: 119-46.

Hill MN, Froese LM, Morrish AC, Sun JC, Floresco SB (2006) Alterations in behavioral flexibility by cannabinoid CB1 receptor agonists and antagonists. Psychopharmacology (Berl) 187: 245-59.

Hoops D, Flores C (2017) Making Dopamine Connections in Adolescence. Trends Neurosci 40: 709-719.

Howes OD, Kapur S (2009) The dopamine hypothesis of schizophrenia: version III--the final common pathway. Schizophr Bull 35: 549-62.

675 Irimia C, Polis IY, Stouffer D, Parsons LH (2015) Persistent effects of chronic Delta9-THC exposure on motor impulsivity
676   in rats. Psychopharmacology (Berl) 232: 3033-43.
677 Islas-Preciado D, Wainwright SR, Sniegocki J, Lieblich SE, Yagi S, Floresco SB, Galea LAM (2020) Risk-based decision
678   making in rats: Modulation by sex and amphetamine. Horm Behav 125: 104815.
679 Jacobs-Brichford E, Manson KF, Roitman JD (2019) Effects of chronic cannabinoid exposure during adolescence on
680   reward preference and mPFC activation in adulthood. Physiol Behav 199: 395-404.
681 Jenni NL, Larkin JD, Floresco SB (2017) Prefrontal Dopamine D(1) and D(2) Receptors Regulate Dissociable Aspects of
682   Decision Making via Distinct Ventral Striatal and Amygdalar Circuits. J Neurosci 37: 6200-6213.
683 Kirschmann EK, Pollock MW, Nagarajan V, Torregrossa MM (2017) Effects of Adolescent Cannabinoid Self-
684   Administration in Rats on Addiction-Related Behaviors and Working Memory. Neuropsychopharmacology 42:
685   989-1000.
686 Klugmann M, Goepfrich A, Friemel CM, Schneider M (2011) AAV-Mediated Overexpression of the CB1 Receptor in the
687   mPFC of Adult Rats Alters Cognitive Flexibility, Social Behavior, and Emotional Reactivity. Front Behav Neurosci
688   5: 37.
689 Lawson KA, Ruiz CM, Mahler SV (2023) A head-to-head comparison of two DREADD agonists for suppressing operant
690   behavior in rats via VTA dopamine neuron inhibition. Psychopharmacology 240: 2101-2110.
691 Le AA, Quintanilla J, Amani M, Piomelli D, Lynch G, Gall CM (2022) Persistent sexually dimorphic effects of adolescent
692   THC exposure on hippocampal synaptic plasticity and episodic memory in rodents. Neurobiol Dis 162: 105565.
693 Lee H-L, Jung K-M, Fotio Y, Squire E, Palese F, Lin L, Torrens A, Ahmed F, Tagne AM, Ramirez J (2022) Frequent low-
694   dose Δ9-tetrahydrocannabinol in adolescence disrupts microglia homeostasis and disables responses to
695   microbial infection and social stress in young adulthood. Biological psychiatry 92: 845-860.
696 Lee HL, Squire E, Fotio Y, Mabou Tagne A, Lee J, Yoon JJ, Hong Y, Kim LH, Jung KM, Piomelli D (2024) Frequent low-
697   impact exposure to THC during adolescence causes persistent sexually dimorphic alterations in the response to
698   viral infection in mice. Pharmacol Res 199: 107049.
699 Lee TT, Hill MN, Lee FS (2016) Developmental regulation of fear learning and anxiety behavior by endocannabinoids.
700   Genes Brain Behav 15: 108-24.
701 Lin L, Jung KM, Lee HL, Le J, Colleluori G, Wood C, Palese F, Squire E, Ramirez J, Su S, Torrens A, Fotio Y, Tang L, Yu
702   C, Yang Q, Huang L, DiPatrizio N, Jang C, Cinti S, Piomelli D (2023) Adolescent exposure to low-dose THC
703   disrupts energy balance and adipose organ homeostasis in adulthood. Cell Metab 35: 1227-1241 e7.
704 Mahler SV, Aston-Jones GS (2012) Fos activation of selective afferents to ventral tegmental area during cue-induced
705   reinstatement of cocaine seeking in rats. J Neurosci 32: 13309-26.
706 Mahler SV, Brodnik ZD, Cox BM, Buchta WC, Bentzley BS, Quintanilla J, Cope ZA, Lin EC, Riedy MD, Scofield MD,
707   Messinger J, Ruiz CM, Riegel AC, Espana RA, Aston-Jones G (2019) Chemogenetic Manipulations of Ventral
708   Tegmental Area Dopamine Neurons Reveal Multifaceted Roles in Cocaine Abuse. J Neurosci 39: 503-518.
709 Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kaufling J, Wilson SP, Deisseroth K, Woodward JJ,
710   Aston-Jones G (2014) Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine
711   seeking. Nature Neuroscience 17: 577-U136.
712 Malone DT, Hill MN, Rubino T (2010) Adolescent cannabis use and psychosis: epidemiology and neurodevelopmental
713   models. Br J Pharmacol 160: 511-22.
714 Malone DT, Taylor DA (1999) Modulation by fluoxetine of striatal dopamine release following Delta9-tetrahydrocannabinol:
715   a microdialysis study in conscious rats. Br J Pharmacol 128: 21-6.
716 Manitt C, Mimee A, Eng C, Pokinko M, Stroh T, Cooper HM, Kolb B, Flores C (2011) The Netrin Receptor DCC Is
717   Required in the Pubertal Organization of Mesocortical Dopamine Circuitry. Journal of Neuroscience 31: 8381-
718   8394.
719 Martinez MX, Farrell MR, Mahler SV (2023) Pathway-Specific Chemogenetic Manipulation by Applying Ligand to Axonally
720   Expressed DREADDs Vectorology for Optogenetics and Chemogenetics. Springer, pp 207-220
721 Matheson J, Sproule B, Di Ciano P, Fares A, Le Foll B, Mann RE, Brands B (2020) Sex differences in the acute effects of
722   smoked cannabis: evidence from a human laboratory study of young adults. Psychopharmacology (Berl) 237:
723   305-316.
724 McAlonan K, Brown VJ (2003) Orbital prefrontal cortex mediates reversal learning and not attentional set shifting in the
725   rat. Behav Brain Res 146: 97-103.
726 Miller EK (2000) The prefontral cortex and cognitive control. Nature reviews neuroscience 1: 59-65.
727 Miller EK, Cohen JD (2001) An integrative theory of prefrontal cortex function. Annu Rev Neurosci 24: 167-202.
728 Mokrysz C, Freeman TP, Korkki S, Griffiths K, Curran HV (2016) Are adolescents more vulnerable to the harmful effects
729   of cannabis than adults? A placebo-controlled study in human males. Translational psychiatry 6: e961-e961.
730 Molla HM, Tseng KY (2020) Neural substrates underlying the negative impact of cannabinoid exposure during
731   adolescence. Pharmacol Biochem Behav 195: 172965.
732 Murray CH, Huang ZY, Lee RY, de Wit H (2022) Adolescents are more sensitive than adults to acute behavioral and
733   cognitive effects of THC. Neuropsychopharmacology 47: 1331-1338.
734 O'Shea M, Singh ME, McGregor IS, Mallet PE (2004) Chronic cannabinoid exposure produces lasting memory impairment
735   and increased anxiety in adolescent but not adult rats. J Psychopharmacol 18: 502-8.

Orsini CA, Setlow B (2017) Sex differences in animal models of decision making. J Neurosci Res 95: 260-269.
Orsini CA, Willis ML, Gilbert RJ, Bizon JL, Setlow B (2016) Sex differences in a rat model of risky decision making. Behav
Neurosci 130: 50-61.
Ott T, Nieder A (2019) Dopamine and Cognitive Control in Prefrontal Cortex. Trends Cogn Sci 23: 213-234.
Owen AM, Roberts AC, Polkey CE, Sahakian BJ, Robbins TW (1991) Extra-dimensional versus intra-dimensional set
shifting performance following frontal lobe excisions, temporal lobe excisions or amygdalo-hippocampectomy in
man. Neuropsychologia 29: 993-1006.
Pantelis C, Barber FZ, Barnes TR, Nelson HE, Owen AM, Robbins TW (1999) Comparison of set-shifting ability in
patients with chronic schizophrenia and frontal lobe damage. Schizophr Res 37: 251-70.
Paxinos G, Watson C (2006) The rat brain in stereotaxic coordinates: hard cover edition. Elsevier
Peters KZ, Zlebnik NE, Cheer JF (2022) Cannabis exposure during adolescence: A uniquely sensitive period for
neurobiological effects. Int Rev Neurobiol 161: 95-120.
Pistis M, Ferraro L, Pira L, Flore G, Tanganelli S, Gessa GL, Devoto P (2002) Δ9-tetrahydrocannabinol decreases
extracellular GABA and increases extracellular glutamate and dopamine levels in the rat prefrontal cortex:: an in
vivo microdialysis study. Brain Research 948: 155-158.
Poulia N, Delis F, Brakatselos C, Polissidis A, Koutmani Y, Kokras N, Dalla C, Politis PK, Antoniou K (2021) Detrimental
effects of adolescent escalating low-dose Δ9-tetrahydrocannabinol leads to a specific bio-behavioural profile in
adult male rats. British Journal of Pharmacology 178: 1722-1736.
Quinn HR, Matsumoto I, Callaghan PD, Long LE, Arnold JC, Gunasekaran N, Thompson MR, Dawson B, Mallet PE,
Kashem MA, Matsuda-Matsumoto H, Iwazaki T, McGregor IS (2008) Adolescent rats find repeated Δ9-THC less
aversive than adult rats but display greater residual cognitive deficits and changes in hippocampal protein
expression following exposure. Neuropsychopharmacology 33: 1113-1126.
Renard J, Krebs MO, Le Pen G, Jay TM (2014) Long-term consequences of adolescent cannabinoid exposure in adult
psychopathology. Front Neurosci 8: 361.
Renard J, Rosen LG, Loureiro M, De Oliveira C, Schmid S, Rushlow WJ, Laviolette SR (2017a) Adolescent Cannabinoid
Exposure Induces a Persistent Sub-Cortical Hyper-Dopaminergic State and Associated Molecular Adaptations in
the Prefrontal Cortex. Cereb Cortex 27: 1297-1310.
Renard J, Szkudlarek HJ, Kramar CP, Jobson CEL, Moura K, Rushlow WJ, Laviolette SR (2017b) Adolescent THC
Exposure Causes Enduring Prefrontal Cortical Disruption of GABAergic Inhibition and Dysregulation of Sub-
Cortical Dopamine Function. Sci Rep 7: 11420.
Reynolds LM, Pokinko M, Torres-Berrio A, Cuesta S, Lambert LC, Del Cid Pellitero E, Wodzinski M, Manitt C, Krimpenfort
P, Kolb B, Flores C (2018) DCC Receptors Drive Prefrontal Cortex Maturation by Determining Dopamine Axon
Targeting in Adolescence. Biol Psychiatry 83: 181-192.
Rubino T, Parolaro D (2011) Sexually dimorphic effects of cannabinoid compounds on emotion and cognition. Front
Behav Neurosci 5: 64.
Rubino T, Parolaro D (2016) The Impact of Exposure to Cannabinoids in Adolescence: Insights From Animal Models. Biol
Psychiatry 79: 578-85.
Rubino T, Realini N, Braida D, Guidi S, Capurro V, Vigano D, Guidali C, Pinter M, Sala M, Bartesaghi R, Parolaro D
(2009) Changes in hippocampal morphology and neuroplasticity induced by adolescent THC treatment are
associated with cognitive impairment in adulthood. Hippocampus 19: 763-72.
Ruiz CM, Torrens A, Castillo E, Perrone CR, Cevallos J, Inshishian VC, Harder EV, Justeson DN, Huestis MA, Swarup V
(2021a) Pharmacokinetic, behavioral, and brain activity effects of Δ9-tetrahydrocannabinol in adolescent male
and female rats. Neuropsychopharmacology 46: 959-969.
Ruiz CM, Torrens A, Lallai V, Castillo E, Manca L, Martinez MX, Justeson DN, Fowler CD, Piomelli D, Mahler SV (2021b)
Pharmacokinetic and pharmacodynamic properties of aerosolized ("vaped") THC in adolescent male and female
rats. Psychopharmacology (Berl) 238: 3595-3605.
Runegaard AH, Sorensen AT, Fitzpatrick CM, Jorgensen SH, Petersen AV, Hansen NW, Weikop P, Andreasen JT,
Mikkelsen JD, Perrier JF, Woldbye D, Rickhag M, Wortwein G, Gether U (2018) Locomotor- and Reward-
Enhancing Effects of Cocaine Are Differentially Regulated by Chemogenetic Stimulation of Gi-Signaling in
Dopaminergic Neurons. eNeuro 5.
Sakurai-Yamashita Y, Kataoka Y, Fujiwara M, Mine K, Ueki S (1989) Δ9-Tetrahydrocannabinol facilitates striatal
dopaminergic transmission. Pharmacology Biochemistry and Behavior 33: 397-400.
Sarne Y (2019) Beneficial and deleterious effects of cannabinoids in the brain: the case of ultra-low dose THC. Am J Drug
Alcohol Abuse 45: 551-562.
Scheyer AF, Laviolette SR, Pelissier AL, Manzoni OJJ (2023) Cannabis in Adolescence: Lasting Cognitive Alterations and
Underlying Mechanisms. Cannabis Cannabinoid Res 8: 12-23.
Schneider M (2008) Puberty as a highly vulnerable developmental period for the consequences of cannabis exposure.
Addiction Biology 13: 253-263.
Schneider M, Koch M (2003) Chronic pubertal, but not adult chronic cannabinoid treatment impairs sensorimotor gating,
recognition memory, and the performance in a progressive ratio task in adult rats. Neuropsychopharmacology 28:
1760-9.

Schramm-Sapyta NL, Cha YM, Chaudhry S, Wilson WA, Swartzwelder HS, Kuhn CM (2007) Differential anxiogenic, aversive, and locomotor effects of THC in adolescent and adult rats. Psychopharmacology (Berl) 191: 867-77.

Schumacher JD, van Holstein M, Bagrodia V, Le Bouder HB, Floresco SB (2021) Dorsomedial striatal contributions to different forms of risk/reward decision making. Neurobiol Learn Mem 178: 107369.

Seamans JK, Yang CR (2004) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. Prog Neurobiol 74: 1-58.

Sewell RA, Ranganathan M, D'Souza DC (2009) Cannabinoids and psychosis. Int Rev Psychiatry 21: 152-62.

Sholler DJ, Strickland JC, Spindle TR, Weerts EM, Vandrey R (2021) Sex differences in the acute effects of oral and vaporized cannabis among healthy adults. Addiction biology 26: e12968.

Simone JJ, Green MR, McCormick CM (2022) Endocannabinoid system contributions to sex-specific adolescent neurodevelopment. Prog Neuropsychopharmacol Biol Psychiatry 113: 110438.

Spear LP (2000) The adolescent brain and age-related behavioral manifestations. Neurosci Biobehav Rev 24: 417-63.

St. Onge JR, Abhari H, Floresco SB (2011) Dissociable contributions by prefrontal D1 and D2 receptors to risk-based decision making. Journal of Neuroscience 31: 8625-8633.

St. Onge JR, Ahn S, Phillips AG, Floresco SB (2012) Dynamic fluctuations in dopamine efflux in the prefrontal cortex and nucleus accumbens during risk-based decision making. Journal of Neuroscience 32: 16880-16891.

St. Onge JR, Chiu YC, Floresco SB (2010) Differential effects of dopaminergic manipulations on risky choice. Psychopharmacology (Berl) 211: 209-21.

St. Onge JR, Floresco SB (2009) Dopaminergic modulation of risk-based decision making. Neuropsychopharmacology 34: 681-97.

St. Onge JR, Floresco SB (2010) Prefrontal cortical contribution to risk-based decision making. Cereb Cortex 20: 1816-28.

Stopper CM, Khayambashi S, Floresco SB (2013) Receptor-specific modulation of risk-based decision making by nucleus accumbens dopamine. Neuropsychopharmacology 38: 715-28.

Stringfield SJ, Torregrossa MM (2021a) Disentangling the lasting effects of adolescent cannabinoid exposure. Prog Neuropsychopharmacol Biol Psychiatry 104: 110067.

Stringfield SJ, Torregrossa MM (2021b) Intravenous self-administration of delta-9-THC in adolescent rats produces long-lasting alterations in behavior and receptor protein expression. Psychopharmacology (Berl) 238: 305-319.

Szkudlarek HJ, Desai SJ, Renard J, Pereira B, Norris C, Jobson CEL, Rajakumar N, Allman BL, Laviolette SR (2019) Delta-9-Tetrahydrocannabinol and Cannabidiol produce dissociable effects on prefrontal cortical executive function and regulation of affective behaviors. Neuropsychopharmacology 44: 817-825.

Torrens A, Roy P, Lin L, Vu C, Grimes D, Inshishian VC, Montesinos JS, Ahmed F, Mahler SV, Huestis MA (2022) Comparative pharmacokinetics of Δ9-tetrahydrocannabinol in adolescent and adult male and female rats. Cannabis and Cannabinoid Research 7: 814-826.

Torrens A, Vozella V, Huff H, McNeil B, Ahmed F, Ghidini A, Mahler SV, Huestis MA, Das A, Piomelli D (2020) Comparative Pharmacokinetics of Δ9-Tetrahydrocannabinol in Adolescent and Adult Male Mice. Journal of Pharmacology and Experimental Therapeutics 374: 151-160.

Tseng AH, Harding JW, Craft RM (2004) Pharmacokinetic factors in sex differences in Δ9-tetrahydrocannabinol-induced behavioral effects in rats. Behavioural Brain Research 154: 77-83.

van den Bos R, Homberg J, de Visser L (2013) A critical review of sex differences in decision-making tasks: focus on the Iowa Gambling Task. Behav Brain Res 238: 95-108.

van den Bos R, Jolles J, van der Knaap L, Baars A, de Visser L (2012) Male and female Wistar rats differ in decision-making performance in a rodent version of the Iowa Gambling Task. Behav Brain Res 234: 375-9.

Verharen JPH, de Jong JW, Roelofs TJM, Huffels CFM, van Zessen R, Luijendijk MCM, Hamelink R, Willuhn I, den Ouden HEM, van der Plasse G, Adan RAH, Vanderschuren L (2018) A neuronal mechanism underlying decision-making deficits during hyperdopaminergic states. Nat Commun 9: 731.

Volkow ND, Swanson JM, Evins AE, DeLisi LE, Meier MH, Gonzalez R, Bloomfield MA, Curran HV, Baler R (2016) Effects of Cannabis Use on Human Behavior, Including Cognition, Motivation, and Psychosis: A Review. JAMA Psychiatry 73: 292-7.

Wiley JL, Burston JJ (2014) Sex differences in Δ9-tetrahydrocannabinol metabolism and in vivo pharmacology following acute and repeated dosing in adolescent rats. Neuroscience Letters 576: 51-55.

Wu X, French ED (2000) Effects of chronic delta9-tetrahydrocannabinol on rat midbrain dopamine neurons: an electrophysiological assessment. Neuropharmacology 39: 391-8.

Zamberletti E, Beggiato S, Steardo L, Jr., Prini P, Antonelli T, Ferraro L, Rubino T, Parolaro D (2014) Alterations of prefrontal cortex GABAergic transmission in the complex psychotic-like phenotype induced by adolescent delta-9-tetrahydrocannabinol exposure in rats. Neurobiol Dis 63: 35-47.

Figure Legends

Fig. 1 Adolescent THC history selectively impacts adulthood learning and cognition a) Experimental timeline. **b)** Schematic of the visual cue discrimination task. **c)** AdoTHC rats (green wedges) were more likely than AdoVEH rats (black wedges) to acquire the visual cue discrimination task to criterion in only one training session (unfilled wedges), rather than requiring 2 sessions to acquire (filled wedges). Both sexes showed similar patterns. **d, e)** AdoTHC females took fewer trials to meet criterion, and made fewer errors when learning visual cue discrimination than AdoVeh females, no such effects were seen in males. **f, g)** AdoTHC females took longer to respond, and made fewer errors than AdoVEH females during cue discrimination training, without effects in males. **h)** Schematic of the subsequently tested strategy set-shifting task. **i)** AdoTHC did not alter the number of trials to learn the new rule to criterion in either sex. Likewise, AdoTHC did not alter in either sex **j)** the number of errors, **k)** the types of errors, **l)** latency to respond, or **m)** omitted trials during set shifting training. Individual rats shown as grey dots in each graph: AdoVEH (n = 14 males, 11 females), AdoTHC (n = 13 males, 10 females). X^2 p^* < 0.05 and repeated measure two-way ANOVA, Sidak post hoc: p^* < 0.05. Data presented as mean + SEM.

Fig. 2 AdoTHC history does not affect baseline probabilistic discounting a) Schematic of the probabilistic discounting task. Sessions consisted of 5 training blocks, in which 2 levers are presented, and rats must choose either a small reward lever that always delivers one pellet, or a large reward-delivering lever which becomes increasingly unlikely to deliver any reward as the session progresses. **b)** Data is shown for males and females, indicating that AdoTHC rats did not differ from their AdoVEH counterparts in probabilistic discounting, with both shifting from nearly exclusively preferring the large reward lever, but appropriately shifting away from it as it became less likely to deliver reward. AdoVEH (n = 14 males, 11 females), AdoTHC (n = 15 males, 8 females). Data represented as average within each probability block from three consecutive days of stable performance. Mean $\pm$ SEM.

Fig. 3 Amphetamine-induced 'risky' responding is potentiated after adolescent THC. a) Data from saline (black line/bar), low dose (0.25mg/kg; pink line/bar) and high dose (0.5mg/kg; red line/bar) amphetamine tests in each sex are shown. When performance patterns were further interrogated, we found that high dose of AMPH in Males increased inflexibility across both AdoTX groups, however in **b)** females, the high dose of AMPH in AdoVEH rats reduced risky responding at high probability blocks not seen in AdoTHC females. **c)** AMPH did not alter win-stay, but reduced lose-shift in AdoTX males. **d)** In AdoVEH females, AMPH decreased win-stay and increased lose-shift, while in AdoTHC AMPH had no effect on win-stay but decreased lose-shift. AMPH = amphetamine. AdoVEH (n = 14 males, 10 females), AdoTHC (n = 15 males, 6 females). Repeated measure three-way ANOVA; Sidak post hoc: p^* < 0.05, p^{**} < 0.01. Data represented as mean + SEM, individual animals shown as grey dots.

Fig. 4 Stimulation of VTA dopamine neurons, or VTA dopamine projections to mPFC does not affect probabilistic discounting a) Bilateral Cre-dependent hM3Dq DREADD AAV injections were made in VTA of TH:Cre rats, and of wildtype (WT) littermates. **b)** Example hM3Dq DREADD expression (red) is localized to tyrosine hydroxylase+ (TH; green) neurons within VTA (yellow=merge). Scale bar, 300 µm. **c)** For pathway-specific stimulation of VTA dopamine projections to mPFC, Cre-dependent hM3Dq DREADDs were injected into VTA as in Experiment 1, and cannulae targeting mPFC allowed CNO microinjection (1mM, 0.5 µl) upon DREADD-expressing dopamine neuron axons in this pathway. **d)** Cannula placements of each rat in pathway stimulation Experiment 2 is shown. **e)** Neither VTA dopamine neuron stimulation induced by systemic CNO in TH:Cre rats, nor **f)** stimulation of the VTA dopamine projection to mPFC induced by mPFC CNO microinjections in TH:Cre rats robustly altered probabilistic discounting in AdoTHC or AdoVEH rats. Likewise, lower panels of **e,f)** indicate that CNO did not have robust effects on WT rats without DREADDs. Data is represented as average % choice of the large reward lever in each probability block across the two CNO, and two VEH tests conducted in each rat, mean$\pm$SEM, p^* < 0.05. TH:Cre+: VTA dopamine stimulation; AdoVEH (n = 8M, 3F) AdoTHC (n = 8M, 4F); VTA dopamine to mPFC: AdoVEH (n = 10M, 4F) AdoTHC (n = 9M, 5F). Wildtype: VTA dopamine stimulation; AdoVEH (n = 5M, 3F) AdoTHC (n = 5M); VTA dopamine to mPFC: AdoVEH (n = 9M, 4F) AdoTHC (n = 8M).

Fig. 1

a.

THC or VEH Injections	Washout Period	Operant Set-Shifting	Probabilistic Discounting + Amphetamine Challenge
Posnatal Day: 30 43	44 69	70+	~85+

Adolescent THC History Improves Learning in Females

b. Visual Cue Discrimination

Respond Under Light, Regardless of Side

c. Days to Criterion

Adolescent
VEH
THC *

Males
2 Days 14.3%
7.7%
35.7%
42.3%
1 Day

Females
2 Days 22.7%
27.3%
50%
1 Day

d. Trials to Criterion
*
Males   Females

e. Errors Prior to Criterion
*
Adolescent
VEH
THC
Males   Females

f. Response Latency (sec)
*
Males   Females

g. Omissions
*
Males   Females

Adolescent THC History Does Not Affect Cognitive Flexibility

h. Shift to Response Discrimination

Respond Based on Side Regardless of Light

i. Trials to Criterion
Adolescent
VEH
THC
Males   Females

j. Errors Prior to Criterion
Males   Females

k. Error Breakdown
Perseverative   Never Reinforced
Adolescent
VEH
THC
# of Errors
Males   Females   Males   Females

l. Response Latency (sec)
Males   Females

m. Omissions
Males   Females

Fig. 2

a. Probabilistic Discounting

b.

Fig. 3

Fig. 4

Experiment 1
Ventral Tegmental Area Dopamine Neuron Stimulation

Experiment 2
VTA Dopamine to Medial Prefrontal Cortex Pathway Stimulation

a. Stimulating all VTA DA Neurons

b. VTA^DA DREADD Localization and hM3Dq Validation

c. Pathway Specific Stimulation

d. Microinjector Placement in mPFC

e. Adolescent VEH | Adolescent THC

f. Adolescent VEH | Adolescent THC