Environmental rather than spatial factors determine trait-based community structure: a case study with zooplankton

Hiromichi Suzuki[1*], Hidetaka Ichiyanagi[2], Jamie M. Kass[1], Jotaro Urabe[1]

1, Graduate School of Life Sciences, Tohoku University, Sendai, Miyagi 980-8578, Japan

2. Water Resources Environmental Center, Chiyoda, Tokyo 102-0083, Japan

*…. Corresponding author: hiromichi.suzuki.t2@dc.tohoku.ac.jp

Open research statement

All the data used in this study was obtained from disclosed sources including the Database of Dams in Japan (http://mudam.nilim.go.jp/home) for the location and watershed areas; the Japan Dam Foundation (http://damnet.or.jp/) for the dam specification data; and the River Environmental Database (Red: http://www.nilim.go.jp/lab/fbg/ksnkankyo/index.html) for water chemistry, fish, and zooplankton. Data obtained from local reports of the national reservoir management office, Japan, can be provided by the authors upon request. All analyses were conducted in the R programming language and all functions used in the present study originate from existing packages.

Keywords: dispersal limitation, functional traits, niche filtering, zooplankton

Abstract

Ecological community structure, which has traditionally been described in terms of taxonomic

units, is driven by a combination of dispersal and environmental filters. Traits have recently been

recognized as alternative units for quantifying community parameters, but they may have

important differences with taxonomic units. For example, as taxon-based community structures

are determined by the identities of individual species, they may be more limited by species'

dispersal constraints, whereas trait-based community structures may be more regulated by

environmental conditions because functional traits should have stronger linkages to habitat types

rather than their specific locations. This suggests that the relative importance of dispersal and

environmental filters may vary depending on the units employed to define community structure,

yet how the contributions of these filters compare remains unclear. Zooplankton in lakes and

reservoirs are an ideal system for examining the relative importance of these filters as they have

moderate dispersal ability and inhabit communities in isolated habitats. In this study, we

examined zooplankton assemblages in 87 artificial reservoirs throughout the Japanese

archipelago to test the hypothesis that environmental filters play a more dominant role in

determining trait-based community structures than taxon-based structures. To describe trait-

based communities, we categorized zooplankton taxa into 12 functional groups using

ecologically meaningful traits, including body size, feeding habits and mode, and population

growth rate. We then examined the effects of both spatial configuration (reflecting dispersal

filters) and environmental variables such as reservoir size and depth, trophic conditions, and fish

assemblages for taxon- and trait-based community structures. Although variation in the taxon-

based structure was explained equally well by spatial and environmental variables, variation in

the trait-based structure explained by environmental variables was nearly twice that of spatial

variables. These results support the idea that environmental filters play a more central role in

determining trait-based community structures, and show that the relative importance of spatial

and environmental filters changes with the way we define community structure.

Introduction

The structure of an ecological community, defined as a biological assemblage in a discrete

habitat, is determined by a combination of filters, including those for species dispersal and the

environment (Bilton et al., 2001; Palmer et al., 1996; Cottenie et al., 2001). Success of dispersal

into a habitat patch is a function of the migratory abilities of the propagules, the distance from

other habitats the species has already colonized, and the presence of geographic barriers between

the habitats (Lockwood et al., 2005; Heino et al., 2015). For a species that has successfully

passed the dispersal filter, its colonization success then depends on whether or not the local

environmental conditions are a subset of its realized niche. These conditions can be defined by

physical and chemical variables, such as habitat complexity, temperature, and nutrient supply

(Schuler et al., 2017; Pearman et al., 2008), but also by biological variables, such as food

resources, predators, and competitors (Dodson et al., 2000; Louette & De Meester, 2007; Hessen

et al., 2006). Although these two filters are conceptually well understood, their relative

importance in determining local community structure remains less so, and this is further

complicated by the fact that community structure can be defined in different ways.

Community structure is commonly described using two different units: taxon-based ones that

describe the community in terms of species composition, and trait-based ones that use

ecologically meaningful functional traits, such as body size and feeding habits. Taxon-based

units are essential for understanding the number and diversity of species in a community

(Hutchinson, 1959; Tilman, 1982; Sax & Gaines, 2003), while trait-based units can help us to

understand how communities are maintained under particular environments (Lindeman, 1942;

Simberloff & Dayan, 1991; Cadotte et al., 2011; Louca et al., 2016; Pomerleau et al., 2015).

Furthermore, the factors most important for regulating community structure may differ

depending on how the structure itself is quantified. On one hand, similarity in community

composition should decrease with increasing habitat distance (Nekola & White, 1999) as species

have different dispersal capacities over limited spatial scales that lead to different species pools

among habitat patches. Thus, dispersal filters should play a substantial role in determining taxon-

based community structure. On the other hand, the niche space of a habitat is structured by local

physical, chemical, and biological conditions that may be similar among habitats regardless of

distance. In the case that different habitats share available niche space but differ in species

composition, these species are likely to be similar in functional traits (Messier et al., 2010). This

suggests that, regardless of distance from other habitats, environmental filters should play a

critical role in determining trait-based community structure. Therefore, it is hypothesized that

environmental filters play a more important role in regulating trait-based community structures,

while dispersal filters have a greater effect on taxon-based community structures. In partial

support of this hypothesis, several studies have demonstrated the importance of environmental

filters in determining the functional structure of animal communities (e.g., Loewen et al., 2019;

Vogt et al. 2013). However, no studies have yet compared the relative importance of both filters

in regulating different community structures, perhaps because this would necessitate the difficult

task of comparing a large number of isolated communities at different distances.

A good model system for testing this hypothesis is freshwater zooplankton, which live in

discrete aquatic habitats surrounded by land (i.e., lakes, ponds, and reservoirs) with distinct

communities that can be easily defined (Hortal et al., 2014). These lentic habitats vary

environmentally depending on their size, depth, and watershed characteristics (Arbuckle &

Downing, 2001). Further, as lentic habitats are scattered across landscapes and zooplankton are

easily collected via water samples, comparing communities established in different

environmental conditions and locations is relatively straightforward. In addition, zooplankton

represent various taxa, including rotifers, cladocerans, and copepods, and differ in body size,

feeding habits, morphology, and behavior. Thus, zooplankton community structure can be

described using both taxon-based and trait-based units.

In general, zooplankton have dormant stages that can tolerate harsh conditions, such as

drought and low temperatures. These tolerances allow them to migrate and disperse propagules

to distant habitats, for example, by drifting on floods (Bozelli et al., 2015), riding the wind

(Pinceel et al., 2016), or hitchhiking on waterfowl (Green & Figuerola, 2005; Hessen et al.,

2019), although this ability appears to vary widely among taxa (Jenkins & Underwood, 1998;

Gray & Arnot, 2011). As such, zooplankton are ideal organisms to examine the relative

importance of environmental and dispersal filters in determining taxon- and trait-based

community structure.

In the present study, we analyzed the zooplankton communities in 87 reservoirs that span the

wide latitudinal range of the Japanese archipelago. First, we examined the taxon-based and trait-

based structures of the zooplankton communities in these reservoirs, then analyzed the dominant

factors regulating these structures. In this analysis, we considered geographic locations of

reservoirs at different scales as spatial factors, and dam specifications, water chemistry, and fish

assemblages as environmental factors. Using the results of these analyses, we tested the

hypothesis that the environmental filter plays a dominant role in regulating the trait-based

community structure, while the dispersal filter has a greater effect on the taxon-based community

structure (Fig. 1).

Materials and Methods

We examined zooplankton communities in 87 reservoirs located in a wide geographical area of

the Japanese archipelago, from 26.48194° to 44.11527° north latitude, from 127.94944° to

143.38778° east longitude, and from 28 to 1123 m in elevation (Fig. 2). Zooplankton data have

been routinely collected in these reservoirs by the National Census on River and Dam

Environment, which has been conducted by the Ministry of Land, Infrastructure, Transport and

Tourism (MLIT) of Japan every five years since 1990. In the program, surveys were carried out

for one of the five years, with samples being taken 4-12 times per survey year. In this study, we

analyzed the zooplankton data collected in each reservoir at seasonal (four times) and monthly

(twelve times) scales for one year each during the 2001-2005 period (3rd round of sampling) and

the 2006-2010 period (4th round of sampling). In 52 of the 87 reservoirs, zooplankton were

collected in both the 3rd and 4th sampling rounds, but in the remaining reservoirs, zooplankton

were collected in only one of these single rounds. Since the 3rd and 4th sampling rounds were five

years apart and we used year-specific environmental data, we treated each sampling round

independently, even for the same reservoirs. Thus, we obtained a total of 139 datasets of the

zooplankton community.

In each reservoir, the zooplankton samples were collected at a site near the dam or at the

deepest point by vertical tows of conical plankton nets with 100-µm mesh from the bottom or

mid-depth to the surface. Collected zooplankton were preserved in Lugol's solution or 4%

formalin solution and examined under a microscope for identification. Specimens that could not

be identified to the species level were classified to the genus level, except for calanoid copepods,

which were often classified to the family or order level due to uncertainty in identification even

for genera. As the volume of water passing through the plankton nets was different among

reservoirs, we did not consider abundance data for zooplankton taxa, and instead reduced counts

to binary presence/absence data.

Taxa and trait units

For each reservoir, we assigned a score of 1 (presence) to the genus or species that occurred in at

least one sample for that year, and a 0 (absence) when no samples were available within a year.

We then estimated the frequency of occurrence for each species or genus across all the reservoirs

and years and created a taxonomic list consisting of species or genera that occurred in more than

15% of the 139 zooplankton datasets. We used these binary datasets as taxon-based community

structures.

To describe the zooplankton community using trait-based units, we first classified all

zooplankton taxa based on body size as follows: large: > 1 mm, medium: 0.5-1 mm, and small: <

0.5 mm in total length excluding the apical spine and external appendage. Although these body-

size categories are arbitrary, body size is known to be closely related to predator vulnerability

(Zaret, 1980; Lampert & Sommer, 2007) and physiological rates, including metabolic (Ikeda 1985;

Lampert 1984) and feeding rates (Peters & Downing 1984). Accordingly, it is often considered a

"master trait" (Litchman & Klausmeier 2008; Litchman et al. 2013). We then divided each of the

body-size groups further into several distinct groups according to trophic level (carnivore,

omnivore, herbivore, detriti-herbivore, and detritivore), feeding mode (filter feeders, raptorial

feeders, and browsers) and population growth rate. Here, "detriti-herbivore" refers to the diet of

animals that feed on both suspended algae and sedimented detritus. Trophic level was classified

based on previous knowledge (Benzie, 2005; Barnett et al., 2007; Branstator, 2005; Rizo et al.,

2017; Kuczyńska-Kippen et al., 2020; Dussart & Defaye, 2001; Koste & Shiel, 1991; Duggan,

2001; Gilbert, 2022). Feeding mode, or the method of resource acquisition, was classified

according to Sommer & Stibor (2002), Branstator (2005) and Barnett et al. (2007) for cladocerans;

Dussart & Defaye (2001) for copepods; and Hillbricht-Ilkowska (1983) and Gilbert (2022) for

rotifers. For some rotifer taxa, no published information regarding feeding mode was available; in

that case, feeding mode was determined based on the morphology of the trophis (i.e., the grinding

apparatus) according to Oh et al. (2017) and Palazzo et al. (2021). Groups with different maximum

population growth rates were classified according to Allan (1976). Thus, using body size, feeding

habits, feeding mode and population growth rate, we classified the zooplankton taxa into 12

functional groups (Table 1, see Table S2 for all taxa). For each of the 139 zooplankton community

records, we assigned a score of 1 (presence) to the functional group in which at least one taxon

occurred, and a 0 (absence) when this group was not represented. The functional groups present in

each community were then used to describe the trait-based community structure.

<u>Spatial and environmental variables</u>

The elevation, latitude, and longitude of the reservoirs were obtained from the Database of Dams

in Japan (<u>http://mudam.nilim.go.jp/home</u>). In addition, data specifying dam characteristics,

including watershed area (Watershed), effective storage volume (Volume), dam height (Height),

dam width (Width), and years elapsed since the reservoir first impounded its full capacity after

dam construction (Age) were obtained from the database published by the Japan Dam

Foundation (http://damnet.or.jp/Dambinran/binran/) and the periodic reports issued by each

reservoir management office. In some reservoirs, a vertical water aeration-based circulation

facility (VWCF) was installed to reduce algal production by moving them to deeper depths. As

algae are important food resources for many zooplankton, such facilities may affect community

structures. We therefore included the presence or absence of a VWCF as an environmental factor.

Water retention time of the reservoir was estimated by dividing the effective water storage

volume by the annual inflow rate in that year, which was obtained from the Database of Dams in

Japan. For reservoirs with no data available from the database, we obtained the annual inflow

data directly from the periodic reports issued by the reservoir management office.

Water chemistry data relating to total phosphorus (TP), total nitrogen (TN), and chlorophyll *a*

(Chl-*a*) measured monthly in the reservoir were obtained from the River Environmental

Database (RED http://www.nilim.go.jp/lab/fbg/ksnkankyo/index.html) established by MLIT. For

the analyses described below, we used the mean annual values of these variables for the year in

which the zooplankton were collected. From the RED, for each site we also obtained

presence/absence records of fish species (total 84) distributed in each reservoir during the period

of the 3rd or 4th round of collection (depending on the record).

<u>Statistical analyses</u>

As above, we analyzed a total of 139 zooplankton communities obtained from 87 reservoirs. All

statistical analyses were performed using R version 4.0.4 (R core team 2021). Spatial

configurations among the reservoirs were modeled using distance-based Moran's eigenvector

maps (dbMEM) based on the geographic coordinates of the reservoirs using the function

"*dbmem*" in the package *adespatial* (Dray et al., 2022). Moran's eigenvalues were estimated

from the geographic distance matrix (latitude and longitude) and used as variables representing

the degree of spatial autocorrelation at different scales between study sites, but also the degree of

dispersal limitation of zooplankton communities. The significance of each Moran's eigenvalue

was tested using the function "*moran.randtest*" in the R package *adespatial*. We labeled the

eigenvalues in order from the largest to the smallest spatial scale, using the notation MEM 1,

MEM 2, and so on.

We summarized the fish assemblages in the reservoirs using a principal coordination

analysis (PCoA) based on the Sorensen dissimilarity matrix using the function *"cmdscale"* in the

R package *stats* (R Core Team, 2021). Before running the analysis, we removed fish species that

occurred in less than 15% of the 139 species lists to minimize the effect of rare species on the

characterization of the fish assemblage. This criterion was arbitrary, but sensitivity tests for 5%

and 10% showed that results were not largely changed. For zooplankton analyses, we used the

scores of the first three axes of the PCoA as fish assemblage data (list the percentages of variance

explained for each axis). In addition, to characterize each axis we calculated species scores for

each axis using the function "*add.spec.score*" in the R package *BiodiversityR* (Kindt & Coe,

2005).

To determine the ability of the environmental and spatial filters to explain each of the taxon-

and trait-based community structures, we performed a partial distance-based redundancy analysis

(partial-dbRDA). This analysis is an extension of multiple regression for community data that has

been often used to compare between different candidate explanatory variable-sets (Legendre &

Legendre, 2012). For environmental filter variables, we used the variables listed in Table 2 and

fish assemblage PCs 1–3. For spatial filter variables, we used MEM 1–12 and elevation. All

variables besides the PCoA scores and Moran's eigenvalues were prior to analysis. We calculated

variance inflation factors among these explanatory variables using function "*cor*" in the R package

stats, and "*solve*" and "*diag*" in the R package base (R core team, 2021). These were less than 10,

indicating negligible variable collinearity (O'Brien 2007). To perform patial-dbRDA, we first

estimated the Sorensen dissimilarity index of the zooplankton composition between the reservoirs

using the function "*capscale*" in the R package *vegan* (Oksanen et al., 2020). To distinguish the

effects of environmental and spatial filter variables, one model was constructed in the partial-

dbRDA to focus on each variable and adjust for the other. For these models, we tested the statistical

significance of the focal variables using permutation tests (n = 1999) with the function "*anova.cca*"

in the R package *vegan*. If the focal variables were significant, we performed variation partitioning

using the function "*varpart*" in R the package *vegan*. This procedure helped us to examine the

relative importance of the different filter variables in determining the taxon- and trait-based

community structures.

As the above analysis included multiple data records that originated from repeat surveys

within the same reservoirs, which might bias our results in favor of conditions at sites with multiple

survey records, we performed a randomization test to determine if and to what extent our results

changed if repeat surveys were excluded. For the reservoirs with records from both the 3rd and

4th round surveys (n=52), we first randomly selected one of these two survey records from each

reservoir. Together with the data from reservoirs with only one survey (3rd or 4th round; n=25),

we created a rarefied dataset of individual records from each of the 87 reservoirs. We then

performed the same analysis as above using this rarefied dataset and estimated the difference in

the percentage of variance explained by spatial and environmental variables. We repeated this

procedure 1000 times. With this analysis, we tested the robustness of our results using all data

from all reservoirs.

Finally, we examined the effects of each individual explanatory variable on community

structure using partial-dbRDA to determine the relative importance of these variables in

determining the community structure. In this analysis, the target variable in each partial-dbRDA

model was treated as the objective variable and the other variables as "conditions". We tested the

statistical significance of each explanatory variable using the permutation tests as described above.

Results

<u>Environmental conditions of reservoirs</u>

The reservoirs varied widely in effective storage volume (Arakawa: 1,250,000 m^3 to Sameura:

289,000,000 m^3), dam height (Ikeda: 24 m to Miyagase: 156 m), watershed area (Arakawa: 7.4

km^2 to Maruyama: 2409 km^2), and age (Ishibuchi: 54 years to Origawa: 2 years [from full

capacity when zooplankton were first sampled]) (Table 2). Of the 87 reservoirs sampled, 13 had

at least one VWCF. Nutrient status varied among the reservoirs from oligotrophic to eutrophic as

TP, TN, and chlorophyll-*a* concentrations ranged from 0.003 to 0.09 mg L^{-1} (median 0.012 mg L^{-1}

$^{-1}$), 0.1 to 1.511 mg/L (0.361 mg L^{-1}), and 0.004 to 35 μg L^{-1} (4.113 mg L^{-1}), respectively.

<u>Spatial configurations of reservoirs</u>

We obtained 12 statistically significant Moran's eigenvalues for the reservoirs, which we refer to

as MEM 1 through MEM 12 (Table S1; Fig S1). Some of these eigenvalues are visualized as

examples in Figure 3, and maps of all 12 significant Moran's eigenvalues are shown in Figure

S1. The number of spatial groups ranged from 2 (the largest scale) to 20 (the smallest scale) (Fig.

3).

<u>Zooplankton</u>

In a total of 139 lists of zooplankton community composition obtained from one- or two-year

surveys for the 87 reservoirs, we identified 131 taxa, including 33 cladocerans, 29 copepods, and

69 rotifers (Table S2). The reservoir with the highest number of taxa per year was Origawa with

52, while the one with the lowest was Omachi with 5, and the overall average among reservoirs

per year was 21 taxa. Of these, 38 taxa were recorded in more than 15% of the zooplankton lists

(Table S2).

We then assigned the 131 taxa to the 12 functional groups based on body size, feeding

habits, feeding mode, and maximum growth rate (Table S2). The reservoir with the highest

number of functional groups per year was Hitokura with 11, while that with the lowest was

Maruyama with 4, and the average 8.2 groups per reservoir per year. The Sorensen index ranged

from 0 to 1 with a mean of 0.559 for the taxon-based community and from 0 to 0.828 with a

mean of 0.202 for the trait-based community (Fig. S2). Because the number of units describing

community structure was smaller for the trait-based community (12 functional groups) than for

the taxon-based community (38 taxa), the coefficient of variation (CV) of the Sorensen

dissimilarity index was smaller for the taxon-based communities (30.3%) than for the trait-based

community (59.5%).

<u>Fish assemblages</u>

The PCoA for fish communities explained a total of 32.0% of the variation in community structure

among reservoirs with the first three axes (Tables 3 and S3). For PC1, warm-water fishes, such as

Pseudogobio ecocinus ecosinus (pike gudgeon), *Micropterus salmoides salmoides* (largemouth

bass)*, Zacco platypus* (pale chub), and *Rhinogobius* sp. (goby) were positively related, whereas

cold-water fishes, such as *Salvelinus* sp. (char) and *Oncorhynchus mykiss* (rainbow trout) were

negatively related. *M. salmoides salmoides* is piscivorous whereas *P. ecocinus ecosinus* and

Rhinogobius sp. are omnivorous and prey on benthic animals. Therefore, we refer to PC1 as the

warm-water euryphagous fish assemblage (WEF). For PC2, *Leuciscus hakonensis / ezoensis*

(Japanese dace*), Hemibarbus labeo barbus (barbel steed)*, *Opsariichthys uncirostris uncirostris*

(three-lips), *Squalidus* sp. (gudgeon), and *Zazzo platypus* showed relatively large positive

eigenvectors (Table 3; Table S3). Since all these fishes were cyprinids, we refer to PC2 as the

cyprinid fish assemblage (CYF). For PC3, *Hypomesus nipponensis* (pond smelt) was positively

associated with the highest loading (Table S3). This fish species is planktivorous and has been

artificially stocked in various lakes and reservoirs for commercial and sport fishing. Therefore, we

refer to PC3 as the smelt-dominated fish assemblage (SDF).

<u>Relative importance of environmental and spatial filters</u>

Partial-dbRDA showed that both environmental and spatial filter variables significantly explained

the variation in both the taxon-based and trait-based structures of zooplankton communities (Table

S4). The variation partitioning analysis showed that about one-third of the total variation in each

community structure was explained by these variables, with different relative proportions for the

different variable types. Of the 29.0% total variance explained for the taxon-based structure, the

unique proportions for environmental (10.9%, 18.8% with covariance) and spatial (10.2%, 18.1%

with covariance) variables were similar (Fig. 4; Table S5). On the other hand, of the 39.2% total

variance that was explained for the trait-based structure, the unique proportion for environmental

variables (19.9%, 27% with covariance) was 1.63 times higher than that for the spatial variables

(12.2%, 19.3% with covariance) (Fig. 4; Table S5). We obtained the same results for the rarefied

datasets created for the randomization test: the difference in the total variance explained between

environmental and spatial variables was always larger for trait-based community than taxon-based

community in 99.9% of the 1000 rarefied datasets (Fig. S3).

<u>Effects of individual variables on the taxon-based community</u>

The partial-dbRDA showed that while dam characteristics (watershed area, reservoir water volume

and dam height) and fish communities [WEF (PC 1) and SDF (PC 3)] had significant effects on

taxon-based community structure (Table 4), lake trophic conditions (TP, TN, and Chl-*a*) did not.

The highest percentages of contribution to the variation (adjusted R^2) were watershed area and

dam height, followed by water volume and fish communities WEF and SDF. Specifically, rotifer

species such as *Ascomorpha* spp., *Asplanchna priodonta*, and *Ploesoma truncatum* were positively

associated with reservoir water volume and SDF, while the calanoid copepods and *Daphnia*

galeata were positively associated with dam height and WEF (Table S6). In contrast, *Cyclops*

vicinus, *Alona* spp. and *Asplanchna priodonta* were negatively associated with WEF (Table S6).

The presence of VWCF had no significant effect on the variation in taxon-based community

structure among reservoirs.

Among the spatial variables, elevation and large- and small-scale MEMs significantly

affected the taxon-based community structure (Table 4). Although the percentage contributions of

these variables to the total variation were relatively small (1.4% to 2.2%), the number of significant

variables affecting taxon-based community structure was greater for spatial factors than for

environmental ones. Among the spatial variables, the cladocera *Daphnia dentifera* was positively

associated with elevation and MEM 10, while the copepod *C. vicinus* was associated negatively

with elevation and MEM 4 (Table S7). Similarly, different rotifer species were associated with

different MEM variables. Some cladocerans, such as *Bosminopsis deitersi*, *Daphnia galeata* and

D. dentifera, showed high eigenvector values for small-scale MEMs (Table S7). In contrast, some

rotifers such as *Ascomorpha ovalis, Cephalodella* sp. and *Collotheca* spp. showed high eigenvector

values for larger-scale MEMs (MEM 2 to 5).

<u>Effects of individual variables on the trait-based community</u>

For the trait-based community structure, the environmental variables for fish assemblages WEF

and CYF, as well as reservoir age, watershed area, water volume, and dam height had significant

effects (Table 5). The highest percentages of contribution to variation (adjusted R^2) were WEF,

CYF, and reservoir age (Table 5). Similar to the taxon-based community, TP, TN, Chl-*a*, and the

presence of VWCF had no significant effect on this community structure. Among functional

groups, large filter-feeding herbivores and omnivores were positively associated with WEF and

negatively associated with watershed area (Table S8). In addition, water volume had positive

associations with large raptorial-feeding carnivores and medium and small filter-feeding

herbivores, yet negative ones with small filter-feeding detriti-herbivores and small raptorial-

feeding herbivores. Large filter feeding omnivores and small filter-feeding herbivores were

associated with reservoir age (Table S8). Large filter-feeding omnivores and small raptorial-

feeding herbivores were positively associated with CYF (Table S8).

Also similar to the taxon-based structure, spatial variables for elevation and large- and small-

scale MEMs significantly affected the trait-based community structure (Table 5). Among these

variables, MEM 10 explained the most variation in community structure, followed by MEM 9 and

MEM 1 (Table 5). Among the functional groups, large filter-feeding herbivores were positively

associated with elevation and different scales of spatial configuration (MEM 1, 3, 4, 9, and 10).

Similarly, large filter-feeding omnivores were positively associated with MEM 1, 9, and 10. In

addition, small raptorial omnivores were positively associated with MEM 3 and 4, but negatively

associated with MEM 1. Large browsing omnivores were negatively associated with MEM 4

(Table S9).

Discussion

Our results show that when zooplankton communities were defined by traits, they were almost

twice as well explained by environmental filters than by dispersal ones, whereas the contributions

of these filters were indistinguishable for communities defined by taxonomy. These results support

our hypothesis that environmental filters play a dominant role in determining functional

community structure and that environmental and dispersal filters play relatively equal roles for

taxonomic structure. Our results confirm that the relative importance of environmental and

dispersal filters in determining community structure can vary with the descriptive unit of the

community.

Among the variables related to the environmental filter, ordination axes characterizing fish

communities had significant effects on the trait-based zooplankton communities. PC1,

representing warm-water euryphagous fish assemblages (WEF) positively represented by

largemouth bass and omnivorous cyprinids, significantly influenced both the taxon-based and trait-

based community structures. Although these fishes often prey on zooplankton (Misawa, 2007),

they are not purely planktivorous. As invasions of largemouth bass have resulted in decreased fish

richness in Japanese lakes (Tsunoda & Mitsuo, 2012), it is likely that reservoirs with high values

of WEF had fewer species of planktivorous fish, making predation pressure on zooplankton more

moderate. In support of this, large zooplankton taxa that outcompete smaller zooplankton yet are

favored by planktivorous fish (Brooks & Dodson, 1965; Hall et al. 1976; DeMott 1989; Urabe

1990) occurred in the trait-based community in such reservoirs. These larger zooplankton included

large filter-feeding herbivores such as *Daphnia* and large omnivores such as calanoid copepods.

The trait-based structure was also significantly affected by PC2, representing the cyprinid fish

assemblage (CYF). Most of these cyprinid fish species are omnivorous but are known to

aggressively feed on zooplankton (Sunaga, 1970; Urabe & Maruyama, 1986). Therefore,

reservoirs with such fish communities are expected to select for zooplankton functional groups

resistant to fish predation. Indeed, in reservoirs with high CYF scores, functional groups were

dominant that can escape predation by quick swimming, such as the large filter-feeding omnivores.

Calanoid copepods, the main components of this group, can sense vibrations from fish movement

to expertly escape predation (Nassal et al., 1998). These results indicate that the top-down effects

of fish strongly influence trait-based community structure.

Fish assemblages categorized as PC1 (WEF) and PC3 (SDF) also influenced the taxon-based

community structure. However, the contribution of fish assemblages was much smaller for the

taxon-based community than for the trait-based community. The classical size-efficiency

hypothesis proposed that the size structure, rather than the species composition, of the

zooplankton community is regulated by predators at higher trophic levels such as fish (Brooks

and Dodson 1965). Since then, many studies have supported the adequacy of this hypothesis

(e.g., Dodson 1974; Zaret 1980; Lynch 1979; Hülsmann 2006; Ye et al. 2013). It follows that

zooplankton communities structured by size class (i.e., traits) should be more affected by

predators than those structured by taxonomy, but no study had yet demonstrated this.

Importantly, our results clearly show that this is true for the zooplankton communities we

studied. It should be noted that although the pond smelt *Hypomesus nipponiensis* is a plankton

feeder (Yoshioka et al. 1994; Kawabata, 2002; Oh et al., 2019), the partial-dbRDA showed that

the copepod *Cyclops vicinus* was positively related to SDF, while other *Cyclopidae* species were

negatively related to SDF, although both were classified in the functional group of medium

raptorial-feeding omnivores. These results suggest that the effects of fish predation are not

necessarily the same among different species even within the same functional groups.

Among the local environmental variables, watershed area, water volume, and dam height

also explained substantial portions of the variation in both community structures, but the effect

of these variables differed among taxa and functional groups. For example, groups with large

body sizes, such as large filter-feeding herbivores and omnivores, tended to occur in reservoirs

with small watershed areas and those that were deep. As zooplankton can escape fish predation

via vertical migration in deep reservoirs (Lampert, 1989; Gliwicz, 1986), they can likely harbor

some functional groups with large body sizes. However, at the taxonomic level, although the

large species *Daphnia galeata* was abundant in reservoirs with high dams (i.e. deep reservoir),

other large zooplankton taxa, such as calanoid copepods, were rarely found in reservoirs with

high dams. The results demonstrate clearly that although functional groups may have tight

connections to certain environmental conditions, this may not be true for all species in that

group.

According to Alfonso et al. (2010), species richness and abundance of copepods and

cladocerans tended to increase up to 50 years after lake formation. In the present study, however,

the trait-based community structure, rather than the taxon-based structure, was related with

reservoir age. This suggests that functional diversity may have a stronger relationship with

reservoir age than species richness.

Zooplankton can disperse passively by wind, flooding, and hitchhiking on mobile animals

such as waterfowl (Vanschoenwinkel et al., 2008). Thus, it is likely that community structure is

limited by the dispersal ability of zooplankton species. Indeed, the present study showed that the

taxon-based community structure of zooplankton was substantially influenced by the spatial

configuration of reservoirs at both smaller (MEM 9 to 11) and larger (MEM 2 to 5) scales.

However, the effects of spatial variables varied among taxa. For example, some cladoceran

species, such as *Bosminopsis deitersi*, *Daphnia galeata* and *D. dentifera*, showed high

eigenvector values for one or more smaller spatial configurations, suggesting that they have

limited dispersal abilities, or alternatively that they can easily disperse to neighboring reservoirs

regardless of environmental conditions. On the other hand, some rotifers, such as *Ascomorpha*

ovalis, *Cephalodella* sp. and *Collotheca* spp., showed high eigenvector values for the larger-scale

spatial configurations (MEM 2 to 5), suggesting that they can disperse to more distant reservoirs.

Although we lack direct knowledge about the dispersal abilities of these species, our results

suggest that dispersal ability is highly variable among zooplankton taxa.

In a similar analysis, Padial et al. (2014) reported that environmental variables were more

important than spatial variables in explaining variation in zooplankton communities defined by

taxonomy among dozens of habitats in a floodplain. However, the present study that covers the

Japanese archipelago is much greater in spatial scale and landscape diversity than a single

floodplain, so our results are likely more reflective of general trends in zooplankton

communities. In addition, our analysis may have underestimated the potential impact of dispersal

filters by excluding less abundant species from the data. Nevertheless, the present study showed

that dispersal and niche factors were equally important for zooplankton taxon-based community

structures. Thus, the spatial configuration of habitats may be more important in determining the

taxonomic structure of the zooplankton community.

Although they were less important compared to environmental variables, some Moran's

eigenvalues affected the trait-based community structure. For example, among the functional

groups, large filter-feeding herbivores was strongly associated with both smaller (MEM 9 and

10) and larger spatial scales (MEM 1, 3, and 4), indicating that its distribution pattern is spatially

complex, probably due to the inclusion of multiple species with different dispersal abilities in

this group. However, some functional groups were associated with particular spatial scales: large

and medium raptorial carnivores dispersed mainly at smaller spatial scales, whereas small and

large filter-feeding omnivores dispersed mainly at larger spatial scales. Regardless of the fact that

we did not consider dispersal ability when categorizing zooplankton taxa into functional groups,

these results suggest that dispersal ability did indeed differ somewhat at a functional group level.

Although the rates of variation we were able to explain for zooplankton community

structure using our variable sets were within previously described ranges for this system, our

study does have some potential shortfalls. For example, Cottenie (2005) showed that multivariate

analyses examining the effects of environmental and spatial variables explained 30 to 60% of the

variation in animal and plant community structure in 95% of studies at the time. However, our

analysis could not explain more than 50% of the total variation in taxon- and trait-based

community structure. As sampling frequency varied among reservoirs, we defined our

community structures based on annual averages. But as zooplankton community structure varies

seasonally (Burns & Mitchell, 1980; Yoshida et al., 2001; Perera et al., 2021), the inclusion of

seasonality may increase the resolution of the analysis when examining the relative importance

between the effects of environmental and spatial variables on community structure. Another

potential shortfall of our study is that we analyzed zooplankton communities using binary data,

possibly causing us to miss some important factors affecting zooplankton structure. For example,

we investigated the effects of vertical water circulation by aeration (VWCF) on the zooplankton

community, which was installed in some reservoirs to reduce algal production. We initially

suspected that VWCF may have affected some zooplankton species by physically disturbing

them through water currents or movement, or by changing the composition and abundance of

phytoplankton. However, we detected no significant effects of the VWCF on taxon- and trait-

based zooplankton community structures, nor did we detect significant effects of TN, TP and

Chl-*a* on these structures. However, it is often reported that taxon-based zooplankton community

structures are related to the trophic status of the studied lakes through bottom-up effects (Pace,

1986; Yoshida et al., 2003; Perga et al., 2010). Lemmens et al. (2018) showed that in a shallow

lake complex in Belgium, zooplankton biomass was mainly regulated by bottom-up effects,

while zooplankton composition in terms of taxa was mainly determined by top-down effects

from fish. Thus, the inclusion of continuous data may allow us to examine more deeply the

effects of VWCR and trophic conditions on zooplankton community structure, and thus increase

the relative importance of environmental variables in determining structures defined by

taxonomy and function.

<u>Conclusion</u>

Our extensive study of zooplankton in reservoirs showed that the relative importance of

environmental and spatial filters in determining community structure varied depending on the

unit used to describe the structure. Trait-based community structure was more influenced by

environmental filters than by dispersal filters, but the taxonomic structure was equally regulated

by both kinds of filters. These results suggest that even if the taxonomic structure of local

zooplankton communities differs due to taxon-specific differences in range and dispersal ability,

the communities tend to comprise functionally similar groups if the environmental conditions of

local habitats are similar. However, because these results were obtained from binary data without

considering the abundance of individual zooplankton taxa, the present study may have

underestimated the effect of environmental filters in determining both taxon- and trait-based

community structure compared to dispersal filters. Therefore, future studies seeking to tackle

these important questions should examine whether the present finding is also supported when the

abundance of composite organisms is considered.

Acknowledgments

We thank the members of the Aquatic Ecology Laboratory at the Graduate School of Life Sciences,

Tohoku University, for their help and comments. This study was financially supported by the Japan

Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (KAKENHI)

20H03315 and 23H02548, the Environment Research and Technology Development Fund

(JPMEERF20214003) of the Environmental Restoration and Conservation Agency provided by

the Ministry of the Environment of Japan, and a Dam-research fund by the Water Resources

Environmental Center, Japan.

Author Contributions

J. U. and H. S. planned and designed the study, H. S., J. U. and H. I. complied data, H. S. and J. U.

performed analyses, H. S. and J. U. wrote the draft with J. M. K. providing major edits, and all

authors contributed to the final manuscript.

Competing interests

The authors declare no competing interests.

References

Alfonso, G., G. Belmonte, F. Marrone, and L. Naselli-Flores. 2010. "Does lake age affect

zooplankton diversity in Mediterranean lakes and reservoirs? A case study from southern

Italy." *Hydrobiologia* 653: 149-164.

Allan, J. D. 1976. "Life history patterns in zooplankton." *The American Naturalist* 110 (971):

165-180.

Arbuckle, K. E., and J.A. Downing. 2001. "The influence of watershed land use on lake N : P in

a predominantly agricultural landscape." *Limnology and Oceanography* 46 (4): 970-975.

https://doi.org/10.4319/lo.2001.46.4.0970.

Barnett, A. J., K Finlay, and B. E. Beisner. 2007. "Functional diversity of crustacean

zooplankton communities: towards a trait-based classification." *Freshwater Biology* 52

(5): 796-813. https://doi.org/10.1111/j.1365-2427.2007.01733.x.

Benzie, J. A. H. 2005. "Guides to the Identification of the Microinvertebrates of the Continental

Waters of the World: Cladocera: the Genus Daphnia (including Daphniopsis)." Leiden:

Backhuys Publishers. Ghent: Kenboi Productions

Bilton, D. T., J. R. Freeland, and B. Okamura. 2001. "Dispersal in freshwater

invertebrates." *Annual Review of Ecology and Systematics* 32: 159-181.

https://doi.org/10.1146/annurev.ecolsys.32.081501.114016.

Bozelli, R. L., S. M. Thomaz, A. A. Padial, P. M. Lopes, and L. M. Bini. 2015. "Floods decrease

zooplankton beta diversity and environmental heterogeneity in an Amazonian floodplain

system." *Hydrobiologia* 753 (1): 233-241. https://doi.org/10.1007/s10750-015-2209-1.

Branstrator, D. K. 2005. "Contrasting life histories of the predatory cladocerans Leptodora

kindtii and Bythotrephes longimanus." *Journal of Plankton Research* 27 (6): 569-585.

https://doi.org/10.1093/plankt/fbi033.

Brooks, J. L., and S. I. Dodson. 1965. "Predation body size and composition of

plankton." *Science* 150 (3692): 28-&. https://doi.org/10.1126/science.150.3692.28.

Burns, C. W., and S. F. Mitchell. 1980. "Seasonal Succession and vertical-distribution of

zooplankton in lake hayes and lake Johnson." *New Zealand Journal of Marine and*

Freshwater Research 14 (2): 189-204. https://doi.org/10.1080/00288330.1980.9515860.

Cadotte, M. W., K. Carscadden, and N. Mirotchnick. 2011. "Beyond species: functional diversity

and the maintenance of ecological processes and services." *Journal of Applied*

Ecology 48 (5): 1079-1087. https://doi.org/10.1111/j.1365-2664.2011.02048.x.

Cottenie, K. 2005. "Integrating environmental and spatial processes in ecological community

dynamics." *Ecology Letters* 8 (11): 1175-1182. https://doi.org/10.1111/j.1461-

0248.2005.00820.x.

Cottenie, K., N. Nuytten, E. Michels, and L. De Meester. 2001. "Zooplankton community

structure and environmental conditions in a set of interconnected

ponds." *Hydrobiologia* 442 (1-3): 339-350. https://doi.org/10.1023/A:1017505619088.

DeMott, W. R. 1989. "The role of competition in zooplankton succession." In *Plankton ecology:*

succession in plankton communities, 195-252. Springer.

Dodson, S. I. 1974. "Adaptive change in plankton morphology in response to size-selective

predation: A new hypothesis of cyclomorphosis." *Limnology and Oceanography* 19 (5):

721-729.

Dodson, S. I., S. E. Arnott, and K. L. Cottingham. 2000. "The relationship in lake communities

between primary productivity and species richness." *Ecology* 81 (10): 2662-2679.

https://doi.org/10.1890/0012-9658(2000)081[2662:TRILCB]2.0.CO;2.

Dray S., Bauman D., Blanchet G., Borcard D., Clappe S., Guénard G., Jombart T., Larocque G.,

Legendre P., Madi N., Wagner H. H. (2022). *adespatial: Multivariate Multiscale Spatial*

Analysis. R package version 0.3-20, https://CRAN.R-project.org/package=adespatial.

Duggan, I. C. 2001. "The ecology of periphytic rotifers." Rotifera IX: Proceedings of the IXth

International Rotifer Symposium, held in Khon Kaen, Thailand, 16–23 January 2000.

Dussart, B. H., and D. Defaye. 2001. "İntroduction to the Copepoda. 2th edition." Leiden :

Backhuys Publisher

Gilbert, J. J. 2022. "Food niches of planktonic rotifers: Diversification and

implications." *Limnology and Oceanography* 67 (10): 2218-2251.

https://doi.org/10.1002/lno.12199.

Gliwicz, M. Z. 1986. "Predation and the evolution of vertical migration in

zpooplankton." *Nature* 320 (6064): 746-748. https://doi.org/10.1038/320746a0.

Gray, D. K., and S. E. Arnott. 2011. "Does dispersal limitation impact the recovery of

zooplankton communities damaged by a regional stressor?" *Ecological Applications* 21

(4): 1241-1256. https://doi.org/10.1890/10-0364.1.

Green, A. J., and J. Figuerola. 2005. "Recent advances in the study of long-distance dispersal of

aquatic invertebrates via birds." *Diversity and Distributions* 11 (2): 149-156.

https://doi.org/10.1111/j.1366-9516.2005.00147.x.

Hall, D. J., S. T. Threlkeld, C. W. Burns, and P. H. Crowley. 1976. "The size-efficiency

hypothesis and the size structure of zooplankton communities." *Annual Review of*

Ecology and Systematics 7 (1): 177-208.

Heino, J., A. S. Melo, T. Siqueira, J. Soininen, S. Valanko, and L. M. Bini. 2015.

"Metacommunity organisation, spatial extent and dispersal in aquatic systems: patterns,

processes and prospects." *Freshwater Biology* 60 (5): 845-869.

https://doi.org/10.1111/fwb.12533.

Hessen, D. O., B. A. Faafeng, V. H. Smith, V. Bakkestuen, and B. Walsdeng. 2006. "Extrinsic

and intrinsic controls of zooplankton diversity in lakes." *Ecology* 87 (2): 433-443.

https://doi.org/10.1890/05-0352.

Hessen, D. O., T. C. Jensen, and B. Walseng. 2019. "Zooplankton Diversity and Dispersal by

Birds; Insights From Different Geographical Scales." *Frontiers in Ecology and*

Evolution 7. https://doi.org/10.3389/fevo.2019.00074.

Hillbricht-Ilkowska, A. 1983. "Response of planktonic rotifers to the eutrophication process and

to the autumnal shift of blooms in Lake Biwa, Japan. I. Changes in abundance and

composition of rotifers." *Japanese Journal of Limnology (rikusuigaku Zasshi)* 44 (2): 93-

106.

Hortal, J., J. C. Nabout, J. Calatayud, F. M. Carneiro, A. Padial, A. M. C. Santos, T. Siqueira, F.

Bokma, L. M. Bini, and M. Ventura. 2014. "Perspectives on the use of lakes and ponds as

model systems for macroecological research." *Journal of Limnology* 73: 46-60.

https://doi.org/10.4081/jlimnol.2014.887.

Hutchinson, G. E. 1959. "Homage to Santa-Rosalia or why are there so many kinds of

animals." *American Naturalist* 93 (870): 145-159. https://doi.org/10.1086/282070.

Hülsmann, S., K. Rinke, and W. M. Mooij. 2005. "A quantitative test of the size efficiency

hypothesis by means of a physiologically structured model." *Oikos* 110 (1): 43-54.

Ikeda, T. 1985. "Metabolic rates of epipelagic marine zooplankton as a function of body-mass

and temperature." *Marine Biology* 85 (1): 1-11. https://doi.org/10.1007/BF00396409.

Jenkins, D. G., and M. O. Underwood. 1998. "Zooplankton may not disperse readily in wind,

rain, or waterfowl." International Rotifer Symposium (8 ; Collegeville, Minn. 1997-06-

22).

Kawabata, K., T. Narita, M. Nagoshi, and M. Nishino. 2002. "Stomach contents of the

landlocked dwarf ayu in Lake Biwa, Japan." *Limnology* 3: 0135-0142.

Kindt, R., and R. Coe. 2005. "Tree diversity analysis: a manual and software for common

statistical methods for ecological and biodiversity studies." *World Agroforestry Centre*.

Koste W. & R. J. Shiel. 1991. "Rotifera from Australian inland waters. VII. Notomattidae

(Rotifera: Monogonata)." *Transactions of the Royal Society of South Australist* 115 (3):

111-159.

Kuczynska-Kippen, N., M. Spoljar, C. Zhang, and M. Pronin. 2020. "Zooplankton functional

traits as a tool to assess latitudinal variation in the northern-southern temperate European

regions during spring and autumn seasons." *Ecological*

Indicators 117. https://doi.org/10.1016/j.ecolind.2020.106629.

Lampert, W. 1984. "The measurement of respiration." *A manual on methods for the assessment*

of secondary productivity in fresh waters 17: 413-468.

Lampert, W. 1989. "The adaptive significance of diel vertical migration of

zooplankton." *Functional Ecology* 3 (1): 21-27. https://doi.org/10.2307/2389671.

Lampert, W., and U. Sommer. 2007. *Limnoecology: the ecology of lakes and streams*. New

York: Oxford university press.

Legendre, P., and L. Legendre. 2012." Numerical ecology third English edition." Elsevier

Lemmens, P, S. A. J. Declerck, K. Tuytens, M. Vanderstukken, and L. De Meester. 2018.

"Bottom-Up Effects on Biomass Versus Top-Down Effects on Identity: A Multiple-Lake

Fish Community Manipulation Experiment." *Ecosystems* 21 (1): 166-177.

https://doi.org/10.1007/s10021-017-0144-x.

Lindeman, R. L. 1942. "The trophic-dynamic aspect of ecology." *Ecology* 23 (4): 399-418.

https://doi.org/10.2307/1930126.

Litchman, E., and C. A. Klausmeier. 2008. "Trait-Based Community Ecology of

Phytoplankton." *Annual Review of Ecology Evolution and Systematics* 39: 615-639.

https://doi.org/10.1146/annurev.ecolsys.39.110707.173549.

Litchman, E., M. D. Ohman, and T. Kiorboe. 2013. "Trait-based approaches to zooplankton

communities." *Journal of Plankton Research* 35 (3): 473-484.

https://doi.org/10.1093/plankt/fbt019.

Lockwood, J. L., P. Cassey, and T. Blackburn. 2005. "The role of propagule pressure in

explaining species invasions." *Trends in Ecology & Evolution* 20 (5): 223-228.

https://doi.org/10.1016/j.tree.2005.02.004.

Loewen, C. J. G., A. L. Strecker, G. L. Larson, A. Vogel, J. M. Fischer, and R. D. Vinebrooke.

2019. "Macroecological drivers of zooplankton communities across the mountains of

western North America." *Ecography* 42 (4): 791-803.

https://doi.org/10.1111/ecog.03817.

Louca, S., L. W. Parfrey, and M. Doebeli. 2016. "Decoupling function and taxonomy in the

global ocean microbiome." *Science* 353 (6305): 1272-1277.

https://doi.org/10.1126/science.aaf4507.

Louette, G., and L. De Meester. 2007. "Predation and priority effects in experimental

zooplankton communities." *Oikos* 116 (3): 419-426. https://doi.org/10.1111/j.2006.0030-

1299.15381.x.

Lynch, M. 1979. "Predation, competition, and zooplankton community structure: an

experimental study 1, 2." *Limnology and Oceanography* 24 (2): 253-272.

Messier, J., B. J. McGill, and M. J Lechowicz. 2010. "How do traits vary across ecological

scales? A case for trait-based ecology." *Ecology letters* 13 (7): 838-848.

Misawa, K., T. Yoneda, S. Inoue, M. Tanigawa, H. Konagaya, and A. Kimura. 2007.

"Interspecific competition for food resources between Dolly Varden (Salvelinus malma)

and rainbow trout (Oncorhynchus mykiss) in the Satsunai River reservoir, upper Tokachi

River System, Hokkaido." *Japanese Journal of Ichthyology* 54 (1): 1-13. (in Japanese)

Nassal, B., W. Burghard, and G. Maier. 1998. "Predation by juvenile roach on the calanoid

copepod Eudiaptomus gracilis and the cyclopoid copepod Cyclops vicinus: a laboratory

investigation with mixed and single prey." *Aquatic Ecology* 32 (4): 335-340.

https://doi.org/10.1023/A:1009995432725. https://doi.org/10.1023/A:1009995432725.

Nekola, J. C., and P. S. White. 1999. "The distance decay of similarity in biogeography and

ecology." *Journal of Biogeography* 26 (4): 867-878. https://doi.org/10.1046/j.1365-

2699.1999.00305.x.

O'Brien, R. M. 2007. "A caution regarding rules of thumb for variance inflation factors." *Quality*

& Quantity 41 (5): 673-690. https://doi.org/10.1007/s11135-006-9018-6.

Oh, H. J., H. G. Jeong, G. S. Nam, Y. Oda, W. Dai, E. H. Lee, D. Kong, S. J. Hwang, and K. H.

Chang. 2017. "Comparison of taxon-based and trophi-based response patterns of rotifer

community to water quality: applicability of the rotifer functional group as an indicator of

water quality." *Animal Cells and Systems* 21 (2): 133-140.

https://doi.org/10.1080/19768354.2017.1292952.

Oh, H. J., Y. Oda, J. Y. Ha, T. Nagata, T. Hanazato, Y. Miyabara, M. Sakamoto, and K. H.

Chang. 2019. "Responses of daphnids and other zooplankton populations to massive fish

kill in Lake Suwa." *Ecological Research* 34 (6): 856-863. https://doi.org/10.1111/1440-

1703.12054.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B.

O'hara, G. L. Simpson, and P. Solymos. 2020. vegan: community ecology package (2.5-7)

https://CRAN.R-project.org/package=vegan

Pace, M. L. 1986. "An empirical-analysis of zooplankton zommunity size structure across lake

trophic gradients." *Limnology and Oceanography* 31 (1): 45-55.

https://doi.org/10.4319/lo.1986.31.1.0045.

Padial, A. A., F. Ceschin, S. A. J. Declerck, L. De Meester, C. C. Bonecker, F. A. Lansac-Toha,

L. Rodrigues, L. C. Rodrigues, S. Train, L. F. M. Velho, and L. M. Bini. 2014. "Dispersal

Ability Determines the Role of Environmental, Spatial and Temporal Drivers of

Metacommunity Structure." *Plos One* 9

(10). https://doi.org/10.1371/journal.pone.0111227.

Palazzo, F., F. F. Bomfim, J. D. Dias, N. R. Simões, F. A. Lansac-Tôha, and C. C. Bonecker.

2021. "Temporal dynamics of rotifers' feeding guilds shaped by chlorophyll-a, nitrate,

and environmental heterogeneity in subtropical floodplain lakes." *International Review of*

Hydrobiology 106 (2): 95-105.

Palmer, M. A., J. D. Allan, and C. A. Butman. 1996. "Dispersal as a regional process affecting

the local dynamics of marine and stream benthic invertebrates." *Trends in Ecology &*

Evolution 11 (8): 322-326. https://doi.org/10.1016/0169-5347(96)10038-0.

Pearman, P. B., A. Guisan, O. Broennimann, and C. F. Randin. 2008. "Niche dynamics in space

and time." *Trends in Ecology & Evolution* 23 (3): 149-158.

https://doi.org/10.1016/j.tree.2007.11.005.

Perera, I. U., N. Maruoka, W. Makino, and J. Urabe. 2021. "Temporal β-diversity of zooplankton

at various time scales in a small mountain lake." *Limnology* 22 (1): 89-99.

710   https://doi.org/10.1007/s10201-020-00637-9. https://doi.org/10.1007/s10201-020-00637-

711   9.

712 Perga, M. E., M. Desmet, D. Enters, and J. L. Reyss. 2010. "A century of bottom-up- and top-

713   down-driven changes on a lake planktonic food web: A paleoecological and paleoisotopic

714   study of Lake Annecy, France." *Limnology and Oceanography* 55 (2): 803-816.

715   https://doi.org/10.4319/lo.2009.55.2.0803.

716 Peters, R. H., and J. A. Downing. 1984. "Empirical analysis of zooplankton filtering and feeding

717   rates 1." *Limnology and Oceanography* 29 (4): 763-784.

718 Pinceel, T., L. Brendonck, and B. Vanschoenwinkel. 2016. "Propagule size and shape may

719   promote local wind dispersal in freshwater zooplankton-a wind tunnel

720   experiment." *Limnology and Oceanography* 61 (1): 122-131.

721   https://doi.org/10.1002/lno.10201.

722 Pomerleau, C., A. R. Sastri, and B. E. Beisner. 2015. "Evaluation of functional trait diversity for

723   marine zooplankton communities in the Northeast subarctic Pacific Ocean." *Journal of*

724   *Plankton Research* 37 (4): 712-726. https://doi.org/10.1093/plankt/fbv045.

R Core Team, A, and R Core Team. 2021. R: A language and environment for statistical

computing. *R Foundation for Statistical Computing*, Vienna, Austria. 2012.

Rizo, E. Z. C., Y. Gu, R. D. S. Papa, H. J. Dumont, and B. P. Han. 2017. "Identifying functional

groups and ecological roles of tropical and subtropical freshwater Cladocera in

Asia." *Hydrobiologia* 799: 83-99.

Sax, D. F., and S. D. Gaines. 2003. "Species diversity: from global decreases to local

increases." *Trends in Ecology & Evolution* 18 (11): 561-566.

https://doi.org/10.1016/S0169-5347(03)00224-6.

Schuler, M. S., J. M. Chase, and T. M. Knight. 2017. "Habitat size modulates the influence of

heterogeneity on species richness patterns in a model zooplankton

community." *Ecology* 98 (6): 1651-1659. https://doi.org/10.1002/ecy.1833.

Simberloff, D., and T. Dayan. 1991. "The guild concept and the structure of ecological

communities." *Annual Review of Ecology and Systematics* 22: 115-143.

https://doi.org/10.1146/annurev.ecolsys.22.1.115.

Sommer, U., and H. Stibor. 2002. "Copepoda–Cladocera–Tunicata: the role of three major

mesozooplankton groups in pelagic food webs." *Ecological Research* 17: 161-174.

Sunaga, T. 1970. "Relation between food consumption and food selectivity in fishes I. food

selectivity of juvenile Hasu (Opsaiichthys uncirostris Teminck et Schlegel), in lake

Biwa." *Japanese journal of ecology* 20 (4): 129-137.

https://doi.org/10.18960/seitai.20.4_129.

Tilman, David. 1982. *Resource competition and community structure*. Vol. 17: Princeton:

Princeton university press.

Tsunoda, H., and Y. Mitsuo. 2012. "Multiple effects of exotic largemouth bass (*Micropterus*

salmoides) and environmental factors on species richness and composition of pond-

dwelling fishes." *Aquatic Living Resources* 25 (2): 163-171.

https://doi.org/10.1051/alr/2012013.

Urabe, J., and T. Maruyama. 1986. "Prey Selectivity of Two Cyprinid Fishes in Ogochi

Reservoir." *Nippon suisan gakkaishi* 52 (12): 2045-2054.

https://doi.org/10.2331/suisan.52.2045.

Urabe, J. 1990. "Stable horizontal variation in the zooplankton community structure of a

reservoir maintained by predation and competition." *Limnology and oceanography* 35

(8): 1703-1717.

Vanschoenwinkel, B., S. Gielen, H. Vandewaerde, M. Seaman, and L. Brendonck. 2008.

"Relative importance of different dispersal vectors for small aquatic invertebrates in a

rock pool metacommunity." *Ecography* 31 (5): 567-577. https://doi.org/10.1111/j.0906-

7590.2008.05442.x.

Vogt, R. J., P. R. Peres-Neto, and B. E. Beisner. 2013. "Using functional traits to investigate the

determinants of crustacean zooplankton community structure." *Oikos* 122 (12): 1700-

1709. https://doi.org/10.1111/j.1600-0706.2013.00039.x.

Ye, L., C. Y. Chang, C. García-Comas, G. C. Gong, and C. Hsieh. 2013. "Increasing

zooplankton size diversity enhances the strength of top-down control on phytoplankton

through diet niche partitioning." *Journal of Animal Ecology* 82 (5): 1052-1061.

Yoshida, T., J. Urabe, and J. J. Elser. 2003. "Assessment of 'top-down' and 'bottom-up' forces as

determinants of rotifer distribution among lakes in Ontario, Canada." *Ecological*

Research 18 (6): 639-650. https://doi.org/10.1111/j.1440-1703.2003.00596.x.

Yoshida, T., M. Kagami, T. B. Gurung, and J. Urabe. 2001. "Seasonal succession of zooplankton

in the north basin of Lake Biwa." *Aquatic Ecology* 35 (1): 19-29.

https://doi.org/10.1023/A:1011498202050.

Yoshioka, T., E. Wada, and H. Hayashi. 1994. "A stable isotope study on seasonal food web

dynamics in a eutrophic lake." *Ecology* 75 (3): 835-846.

Zaret, Thomas M. 1980. "Predation and freshwater communities." London: Yale University

Press

Table 1. Functional groups used for description of the trait-based community.

Functional category	Body size	Feeding mode	Food habit	Population growth rate
Large raptorial-feeding carnivore	Large	Raptorial	Carnivore	Middle
Large filter-feeding herbivore	Large	Filter	Herbivore	Middle
Large browsing omnivore	Large	Browser	Omnivore	Low
Large filter-feeding omnivore	Large	Filter	Omnivore	Low
Medium filter-feeding herbivore	Medium	Filter	Herbivore	Middle
Medium raptorial-feeding omnivore	Medium	Raptorial	Omnivore	Low
Medium raptorial-feeding carnivore	Medium	Raptorial	Carnivore	High
Small filter-feeding herbivore	Small	Filter	Herbivore	Middle
Small filter-feeding detriti-herbivore	Small	Filter	Detritivore / Herbivore	Middle
Small raptorial-feeding herbivore	Small	Raptorial	Herbivore	High
Small raptorial-feeding omnivore	Small	Raptorial	Omnivore	High
Small filter-feeding detritivore	Small	Filter	Detritivore	High

785   **Table 2.** Summary of environmental variables showing maximum, minimum, mean, median, and

786   standard deviation (SD).

Variables	Min	Max	Mean	Median	SD
Nutrient condition					
TP (mg/l)	0.003	0.09	0.017	0.012	0.149
TN (mg/l)	0.100	1.511	0.449	0.361	0.279
Chl *a* (mg/m^3)	0.004	35.000	6.351	4.113	6.266
Dam specifications					
Watershed (km^2)	7.4	2409	342.5	225.5	409.5
Volume (1000m^3)	1250	289000	55960.1	42000	52427
Height (m)	24	156	89.06	86	31.3
Width (m)	127.6	1480	340.2	322	173
VWCF	0	1	-	-	-
Turnover (/year)	1.106	922.140	29.025	7.546	92.127
Age (years)	2	54	25.91	26	13

794   **Table 3.** Results of the principal coordination analysis showing the eigenvalue, viability, and

795   cumulative variability of the first three coordinates, with the top 5 fish taxa having the highest

796   absolute eigenvector value in these coordination axes. The direction of the eigenvector is shown

797   in parentheses.

Axis	PC1	PC2	PC3
Eigenvalue	33.04	15.86	11.32
Cumulative variability (%)	17.58	26.02	32.04
Rank			
1	*Zacco platypus* (+)	*Zacco platypus* (+)	*Hypomesus nipponiensis* (+)
2	*Salvelinus* sp. (-)	*Leuciscus hakonensis / ezoensis* (+)	Oncorhynchus masou masou *(+)*
3	*Pseudogobio ecocinus ecocinus* (+)	*Hemibarbus labeo barbus* (+)	*Phoxinus* sp. (+)
4	*Rhinogobius* sp. (+)	*Squalidus* sp. (+)	*Misgurmus anguillicaudatus* (+)
5	*Micropterus salmoides salmoides* (+)	*Opsariichthys uncirostris uncirostris* (+)	*Cyprinus carpio* (+)
Name of fish assemblage	*Warm-water euryphagous fish assemblage* (WEF)	*Cyprinid fish assemblage* (CYF)	*Smelt-dominated fish assemblage* (SDF)

802　**Table 4** Results of partial-dbRDA for the taxon-based community showing adjusted coefficients

803　of determination for environmental and spatial variables with significant probability. For

804　significant variables, the direction of the effects is shown in parentheses. Bold type indicates the

805　variables that were significant.

Environment variables	adjusted R^2	F	p	Spatial variables	adjusted R^2	F	p
TP	0.011	1.510	0.067	**Elevation (+)**	**0.020**	**2.300**	**0.004**
TN	0.006	1.041	0.395	MEM1	0.007	1.100	0.324
Chl *a*	0.004	0.843	0.65	**MEM2 (+)**	**0.020**	**2.308**	**0.003**
WEF (PC1) (+)	**0.017**	**2.085**	**0.004**	**MEM3 (+)**	**0.0201**	**2.402**	**0.002**
CYF (PC2)	0.005	0.955	0.515	**MEM4 (+)**	**0.015**	**1.852**	**0.012**
SDF (PC3) (-)	**0.013**	**1.661**	**0.034**	**MEM5 (+)**	**0.016**	**1.941**	**0.014**
Watershed (+)	**0.036**	**3.755**	**< 0.001**	MEM6	0.010	1.378	0.123
Volume (+)	**0.025**	**2.766**	**< 0.001**	MEM7	0.010	1.407	0.106
Height (-)	**0.029**	**3.166**	**< 0.001**	MEM8	0.010	1.405	0.109
Width	0.003	0.757	0.766	**MEM9 (+)**	**0.022**	**2.483**	**< 0.001**
VWCF	0.005	0.986	0.468	**MEM10 (-)**	**0.014**	**1.742**	**0.025**
Turnover rate	0.004	0.809	0.638	**MEM11 (+)**	**0.015**	**1.825**	**0.02**
Age	0.007	1.135	0.289	MEM12	0.003	0.732	0.777

810 **Table 5** Results of partial-dbRDA for the trait-based community showing adjusted coefficients of

811 determination for environmental and spatial variables with significant probability. For significant

812 variables, the direction of the effects is shown in parentheses. Bold type indicates the variables that

813 were significant.

Environment variables	adjusted R^2	F	p	Spatial variables	adjusted R^2	F	p
TP	<0.001	0.298	0.964	**Elevation (+)**	**0.030**	**2.988**	**0.003**
TN	0.004	0.645	0.743	**MEM1 (-)**	**0.039**	**3.730**	**< 0.001**
Chl *a*	0.011	1.309	0.218	MEM2	0.018	1.940	0.066
WEF (PC1) (+)	**0.069**	**6.367**	**< 0.001**	**MEM3 (+)**	**0.020**	**2.073**	**0.050**
CYF (PC2) (-)	**0.037**	**3.565**	**0.002**	**MEM4 (+)**	**0.028**	**2.767**	**0.006**
SDF (PC3)	0.009	1.062	0.407	MEM5	0.006	0884	0.503
Watershed (+)	**0.061**	**5.703**	**< 0.001**	MEM6	0.007	0.898	0.500
Volume (+)	**0.059**	**5.504**	**< 0.001**	MEM7	0.016	1.721	0.099
Height (-)	**0.021**	**2.158**	**0.035**	MEM8	0.009	1.101	0.343
Width	0.007	0.954	0.454	**MEM9 (+)**	**0.043**	**4.132**	**0.001**
VWCF	0.007	0.964	0.463	**MEM10 (-)**	**0.051**	**4.841**	**< 0.001**
Turnover	0.008	1.012	0.316	MEM11	0.014	1.506	0.146
Age (+)	**0.018**	**1.934**	**0.046**	MEM12	0.002	0.448	0.865

FIGURE LEGENDS

Figure 1. Conceptual diagram showing the workflow of our analysis. We quantified the relative

importance of dispersal and niche filters by examining the relationship between taxon- or trait-

based community structures and environmental or spatial factors.

Figure 2. Map showing locations of artificial reservoirs in the Japanese archipelago from which

we sampled zooplankton in this study.

Figure 3. Maps showing examples of Moran's eigenvalues that we calculated for reservoir

locations and used as spatial variables for dispersal filters in our analysis. The largest spatial

scale is represented by MEM 1 (a) and scales become smaller with increasing numbers: MEM 4

represents a medium scale (b), while MEM 11 represents a small scale (c). The white and black

circles represent positive and negative eigenvalues, respectively, and the size of the circle

represents the magnitude of the absolute value.

Figure 4. Venn diagrams showing the contributions of environmental variables, spatial variables,

and residuals to the total variation in the taxon-based (a) and trait-based (b) zooplankton

community structures.

Figure 1.

Figure 2

848 **Figure 3**

860 **Figure 4**

(a)

(b)

Residuals = 0.710

Residuals = 0.608

861