# Temporally controlled nervous system-to-gut signaling bidirectionally regulates longevity in *C. elegans*

Lingxiu Xu[1,#], Chengxuan Han[1,#], Lei Chun[1,#, *], X.Z. Shawn Xu[2,*], and Jianfeng Liu[1,*]

[1]College of Life Science and Technology, Key Laboratory of Molecular Biophysics of MOE, Huazhong University of Science and Technology, Wuhan, Hubei, 430074, China

[2]Life Sciences Institute and Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI 48109, U.S.A.

[#]Co-first authors

*Correspondence: Lei Chun (lchun@hust.edu.cn)

X.Z. Shawn Xu (shawnxu@umich.edu)

Jianfeng Liu (jfliu@mail.hust.edu.cn)

**Abstract**

The nervous system modulates aging by secreting signaling molecules to cell-nonautonomously regulate the physiological state of distal tissues such as the gut. However, the underlying mechanisms are not well understood. Here, using *C. elegans* as a model, we identified two distinct neuroendocrine signaling circuits through which motor neurons signal the gut in early life to shorten lifespan but in mid-late life to extend lifespan. Both circuits employ the same neurotransmitter acetylcholine (ACh), while recruiting two different gut ACh receptors ACR-6 and GAR-3 to regulate the transcription factor DAF-16 and HSF-1 in early and mid-late life, respectively. Strikingly, the gut expression of ACR-6 is restricted to early life, whereas that of GAR-3 is confined to mid-late life, providing a potential mechanism for the temporal control of the two circuits. These results identify a novel mechanism that empowers the nervous system to bidirectionally regulate longevity by differentially signaling the gut at different life stages.

**Introduction**

Aging is a complex physiological process characterized by a progressive

functional decline in tissues and organs, leading to increased incidents in age-

related diseases and probability of death[1,2]. Such an aging process occurs body

wide in multiple tissues/organs in a coordinated manner, though different

tissues/organs may age at different rates[3-8]. Recent work has uncovered an

increasingly important role for the nervous system in orchestrating such a

coordinated aging process through tissue-tissue communications[5,9-14]. The

nervous system mediates tissue-tissue communications by secreting various

signaling molecules such as small neurotransmitters and neuropeptides, which

modulate the physiological state of distal tissues. For example, in *C. elegans*,

the nervous system can detect and process internal stress signals from

mitochondria and external sensory cues, such as temperature and odors, and

then communicates with the distal tissue gut to regulate aging in a cell-

nonautonomous manner[14-21]. How the nervous system signals distal tissues to

regulate aging remains a question under intensive investigation.

The nervous system modulates aging in a complex fashion. Research in

the past decade has uncovered both negative and positive impact of the

nervous system on aging. The notion that the output of neural cells, including

both neurons and glial cells, plays an important role in aging has gained

consensus[19,22,23], though conflicting results have been reported as to whether

enhanced neural output is beneficial or detrimental to longevity. For example,

several studies have demonstrated that neuronal hyper-excitability shortens

lifespan in organisms ranging from *C. elegans* to humans[24-26]. Specifically, the

transcription factor REST suppresses neural gene expression in humans and

mice, and overexpression of the *C. elegans* REST orthologue *spr-4* suppresses

neuronal output and extends lifespan[24].  On the other hand, several studies

support the opposing view that enhanced neuronal output extends lifespan. For

example, genetic and pharmacological potentiation of motor neuron activity and

synaptic transmission at neuromuscular junctions (NMJs) slow down motor

aging and extend lifespan in *C. elegans*[27-29].  These seemingly conflicting

results reflect the complexity of the role of the nervous system in aging

modulation. One possibility is that the nervous system can both negatively and

positively regulate aging, depending on the context; however, whether and how

the nervous system does so is not well understood.

In this study, we sought to address these questions in *C. elegans*, a

widely used genetic model organism for the study of aging. Using the *C.*

*elegans* motor nervous system as a model, we developed a strategy to

temporally manipulate the motor neuron output at different life stages. We

identified two distinct neuroendocrine signaling circuits through which neuronal

output inhibits longevity in early life but promotes it in mid-late life.  Both circuits

require the neurotransmitter acetylcholine (ACh) as a signaling molecule that

signals the gut via two different gut ACh receptors, ACR-6 and GAR-3, to

regulate two different transcription factors, DAF-16 and HSF-1, at early and

mid-late life stages, leading to shortened and extended lifespan, respectively.

Strikingly, the expression of the ACh receptors ACR-6 and GAR-3 in the gut

undergoes a temporal switch in early and mid-late life, providing a potential

mechanism to temporally control the activity of the two circuits.  Our results

96 define the molecular and circuit mechanisms by which the nervous system

97 bidirectionally regulates longevity by differentially signaling the gut at different

98 life stages, highlighting an exquisite role of the nervous system in aging

99 modulation.

## Results

**Cholinergic motor neurons differentially regulate lifespan at different stages of worm life**

To ascertain whether the nervous system promotes or inhibits longevity, we focused on the motor nervous system using *C. elegans* as a model. Previous work has uncovered a role of *C. elegans* cholinergic but not GABAergic motor neurons in the ventral nerve cord in lifespan regulation[28,30]. To further explore the role of cholinergic motor neurons in lifespan control, we ablated the output of these motor neurons by expressing tetanus toxin (TeTx) as a transgene in these neurons using the *acr-2* promoter. TeTx cleaves the SNARE protein synaptobrevin, thereby blocking the release of signaling molecules such as neurotransmitters and neuropeptides from the neurons of interest[31]. Surprisingly, such intervention led to a complex effect on the lifespan curve by initially promoting worm's survival followed by inhibiting it at later stages (Figure 1A). Specifically, the 25% lifespan of these worms was prolonged, while their 75% and maximal lifespan were slightly shortened, leading to a mean lifespan slightly increased or unchanged compared to that of wild-type worms. This suggests that inhibiting cholinergic motor neurons may extend lifespan at early stages but shorten it at late stages. To test this model, we performed the converse experiment by promoting the output of cholinergic motor neurons. This was achieved by expressing in these neurons a transgene encoding syntaxin(T254I), a gain-of-function form of *Drosophila* syntaxin, another SNARE protein required for neuronal exocytosis[32]. Expression of this syntaxin mutant enhances synaptic release without causing vesicle depletion, thereby promoting

neuronal output[32], an approach that has previously been successfully applied to

manipulate neuronal output in lifespan assays[20]. This transgene simply

shortened lifespan without causing a pleotropic effect (Figure 1B). This

suggests that enhancing the output from cholinergic motor neurons in early life

may cause an irreversible detrimental effect that overwhelms its beneficial effect

in mid-late life.

To further characterize the temporal effect of cholinergic motor neurons

on lifespan, we sought to vary the output of cholinergic motor neurons in a

temporally controlled manner.  To do so, we employed an auxin-induced protein

degradation (AID) system to express syntaxin(T254I) fused with a degron tag as

a transgene in cholinergic motor neurons[33]. Transgenic worms were then

crossed with a line stably expressing TIR1 in these neurons. TIR1 is the plant

ubiquitin ligase, an essential component of the AID system[33]. Expression of

syntaxin(T254I) can be suppressed by auxin treatment and induced by

subsequently removing auxin. In doing so, we were able to potentiate the output

of cholinergic motor neurons in a temporally controlled manner by expressing

syntaxin(T254I) at different stages of worm life (Figure 1C).  As a control, we

first shut down the expression of *syntaxin(T254I)* transgene throughout the

entire worm life by treating worms with auxin at all life stages, and found that

this had no notable effect on lifespan (Figure 1D).  Conversely, enhancing the

output of cholinergic neurons throughout the entire worm life (no auxin

treatment) shortened lifespan, consistent with the view that the detrimental

effect resulting from enhanced motor neuron output in early life overwrites its

beneficial effect at later stages (Figure 1E). Indeed, enhancing the output of

cholinergic motor neurons during the larval stage only (no auxin from egg to L4)

shortened lifespan, confirming the detrimental effect of motor neuron output on

lifespan in early life (Figure 1F). Notably, this detrimental effect began to fade

when motor neuron output was instead enhanced during the adult stage (auxin

from egg to adult stage), and eventually worms became long-lived when motor

neuron output was enhanced after Day 5 adulthood (Figure 1G-1L). These data

reveal a dichotomous role of cholinergic motor neurons in lifespan regulation,

shortening adult lifespan in early life but extending it in mid-late life.

**ACh mediates the lifespan-shortening effect of cholinergic motor neurons**

**in early life**

How do cholinergic motor neurons differentially control lifespan at

different life stages? First, we focused on their lifespan-shortening effect.

Cholinergic motor neurons release both small neurotransmitters and

neuropeptides[34-37]. To determine which type of signaling molecules might be

released from motor neurons to shorten lifespan in early life, we first tested *unc-*

*31* mutant worms, which are defective in secreting neuropeptides[38-40]. Loss of

*unc-31* exhibited no defect on the ability of cholinergic motor neurons to shorten

lifespan, indicating that neuropeptides may not play an important role in this

process (Figure 2A and 2B). This led us to test small neurotransmitters.

Cholinergic motor neurons release ACh, and indeed, loss of *unc-17* gene, which

encodes the sole *C. elegans* vesicular ACh transporter[41], eliminated the

lifespan-shortening effect of these motor neurons (Figure 2A and 2C).  In *unc-*

*17* mutant worms, no ACh is expected to be released from cholinergic motor

neurons. As a control, we also tested mutants deficient in other types of small

neurotransmitters, including glutamate (*eat-4*), GABA (*unc-25*), serotonin (*tph-1*),

dopamine (*cat-2*), tyramine (*tdc-1*), and octopamine (*tbh-1*), but detected no

effect (Figure S1A-S1F). These results uncover a key role of the

neurotransmitter ACh in mediating the lifespan-shortening effect of cholinergic

motor neurons in early life.

**The nicotinic AChR ACR-6 acts in the intestine to mediate the lifespan-**

**shortening effect of cholinergic motor neurons in early life**

ACh is expected to act through ACh receptors (AChRs).  To identify the

AChRs that mediate the lifespan-shortening effect of cholinergic motor neurons,

we examined all of the 37 *C. elegans* AChR genes, including nicotinic AChRs

(nAChRs), muscarinic AChRs (mAChRs), and ACh-gated Cl⁻ channels[35]. By

screening strains with each of these 37 AChRs knocked down by RNAi, we

identified three candidate genes: *acr-6* (nAChR), *acr-8* (nAChR) and *gar-2*

(mAChR) (Figure S2). To validate these candidates, we examined their mutants.

Mutation in *acr-6* blocked the ability of cholinergic motor neurons to shorten

lifespan, while mutations in the other two candidates did not (Figure 2D and

Figure S3). This identifies a key role for the nAChR ACR-6 in mediating the

lifespan-shortening effect of cholinergic motor neurons in early life.

We then asked in which tissue ACR-6 regulates lifespan. Transgenic

expression of wild-type *acr-6* gene in the intestine (Figure 2E), but not in

neurons (Figure 2F), rescued the *acr-6* mutant phenotype, indicating that ACR-

6 acts in the intestine. This suggests that ACh released from cholinergic motor

neurons may shorten lifespan by acting on its cognate receptor ACR-6 in the

intestine, revealing a nervous system-to-gut signaling axis.

**Cholinergic motor neurons require the transcription factor DAF-16 in the**

**intestine to regulate lifespan in early life**

Aging pathways tend to converge on a subset of transcription factors[12]. We

went on to search for a transcription factor that functions in the intestine to

mediate the lifespan-shortening effect of cholinergic motor neurons in early life.

RNAi of *daf-16,* which encodes a FOXO transcription factor that is a key

regulator of lifespan, did not further shorten the lifespan of worms expressing

*syntaxin(T254I)* transgene (Figure 3A and 3B); as a control, RNAi of other

transcription factor such as *hsf-1*/HSF1, *skn-1*/Nrf2 *and pha-4*/FOXA did (Figure

3C-3E). This suggests that *daf-16* may act in the same pathway as cholinergic

motor neurons. DAF-16 regulates lifespan by controlling the expression of its

target genes. As expected, the expression level of *sod-3* and *mtl-1*, two

commonly characterized DAF-16 target genes, was upregulated in transgenic

worms deficient in releasing ACh from cholinergic motor neurons (Figure 3F),

consistent with the notion that DAF-16 acts downstream of cholinergic motor

neurons.

DAF-16 is known to regulate lifespan primarily in the intestine[4]. To

determine whether DAF-16 also functions in the intestine, we performed tissue

specific RNAi experiments in *sid-1* mutant background to restrict RNAi to

specific tissues. dsRNA against *daf-16* was expressed as a transgene

specifically in the intestine. RNAi of *daf-16* in the intestine abolished the ability

of cholinergic motor neurons to regulate lifespan at early life stage (Figure 3G

and 3H). Thus, similar to ACR-6, DAF-16 also appears to function in the

intestine, suggesting that the intestine is an important target tissue to which

cholinergic motor neurons signals to regulate lifespan. It also suggests that

cholinergic motor neurons shorten lifespan at early life stage by regulating DAF-

16 activity in the intestine.

Collectively, the data presented above suggests a model in which

cholinergic motor neurons regulate lifespan in early life through a

neuroendocrine signaling circuit (Figure 6D, below). In this circuit, cholinergic

motor neurons signal the intestine via the neurotransmitter ACh that acts

through its cognate receptor ACR-6 to regulate DAF-16 in the intestine, leading

to a shortened lifespan (Figure 6D).

**ACh mediates the lifespan-extending effect of cholinergic motor neurons**

**in mid-late life**

Having characterized the mechanisms by which cholinergic motor

neurons signal the intestine to shorten lifespan in early life, we then set out to

explore how these motor neurons extend lifespan in mid-late life. Given that

ACh is the only neurotransmitter known to be released by these motor neurons

and its role in mediating motor neuron-to-gut signaling in early life, we asked

whether ACh also plays a role in mediating the lifespan-extending effect of

these motor neurons in mid-late life. Again, we resorted to the AID system to

specifically enhance the output of cholinergic motor neurons in mid-late life by

temporally restricting the expression of *syntaxin(T254I)* transgene at this stage.

We found that cholinergic motor neurons lost the ability to extend lifespan at

mid-late life stage in *unc-17* mutant worms that are deficient in ACh release,

revealing an essential role of ACh in mediating the lifespan-extending effect of

these motor neurons in mid-late life and suggesting it as a key signaling

molecule in this process (Figure 4A and 4B). Thus, the same neurotransmitter

ACh appears to mediate both the lifespan-shortening and extending effects of

cholinergic motor neurons.

**The muscarinic AChR GAR-3 acts in the intestine to mediate the lifespan-**

**extending effect of cholinergic motor neurons in mid-late life**

We then asked how the same neurotransmitter ACh secreted from the

same type of neurons mediates two opposite effects on lifespan. To address

this question, we re-screened all the 37 AChR genes (Figure S4). Two

candidate genes emerged from the screen: *gar-2* and *gar-3*, which encode two

different mAChRs. RNAi of *gar-2* or *gar-3* completely suppressed the ability of

cholinergic motor neurons to extend lifespan in mid-late life (Figure S5A-S5C).

*gar-2* and *gar-3* mutant worms exhibited a similar phenotype (Figure S5D, S5E

and Figure 4C), suggesting that GAR-2 and GAR-3 may be the receptors for

ACh that mediate the lifespan-extending effect of cholinergic motor neurons.

To determine in which tissue GAR-2 and GAR-3 act to regulate lifespan,

we specifically knocked down *gar-2* and *gar-3* genes in the intestine and

neurons, as both genes are known to be expressed in these two tissues[42,43].

RNAi of *gar-3* in the intestine (Figure 4D and 4E), but not in neurons (Figure 4D

and 4F), abolished the ability of cholinergic motor neurons to extend lifespan at

mid-late life stage. Thus, GAR-3 may function in the intestine to regulate

lifespan.  Surprisingly, RNAi of *gar-2* in the intestine or in neurons had no

notable effect on the ability of cholinergic motor neurons to extend lifespan in

mid-late life (Figure S5F-S5H), indicating that GAR-2 acts in a non-intestine,

non-neuronal tissue to regulate lifespan. Given our focus on neuron-gut

signaling, we decided to focus on GAR-3 for further characterizations.  These

results identify GAR-3 as a candidate AChR that acts in the intestine to mediate

the effect of the neurotransmitter ACh and cholinergic motor neurons on

lifespan extension in mid-late life.

**Cholinergic motor neurons require the transcription factor HSF-1 in the**

**intestine to regulate lifespan in mid-late life**

In light of our observation that the transcription factor DAF-16 acts in the

intestine to regulate lifespan in early life, we wondered if DAF-16 also regulates

lifespan in mid-late life.  Surprisingly, RNAi of *daf-16* and other transcription

factors such as *skn-1* and *pha-4* failed to suppress the lifespan-extending effect

of cholinergic motor neurons (Figure S6). By contrast, RNAi of *hsf-1* completely

abolished the ability of these motor neurons to extend lifespan in mid-late life

(Figure 5A and 5B), suggesting that HSF-1, rather than DAF-16, is required by

cholinergic motor neurons to regulate lifespan at mid-late life stage. Additional

support came from the observation that the expression of *hsp-16.2* and *pat-10*,

two commonly characterized HSF-1 target genes, were upregulated in long-

lived worms in which the output of cholinergic motor neurons was enhanced at

mid-late life stage (Figure 5C). HSF-1 is known to regulate lifespan primarily in

the intestine and neurons[44,45]. To identify in which tissue HSF-1 functions, we

inactivated *hsf-1* specifically in the intestine or neurons by RNAi (Figure 5D-5F).

RNAi of *hsf-1* in the intestine but not in neurons abolished the ability of

cholinergic motor neurons to extend lifespan at mid-late life stage. Thus, HSF-1

appears to function in the intestine to regulate lifespan, suggesting that

cholinergic motor neurons extend lifespan in mid-late life through HSF-1 in the

intestine.

In summary, this set of data suggests a model in which cholinergic motor

neurons signal the distal tissue intestine in mid-late life to extend lifespan

through a distinct neuroendocrine circuit (Figure 6D). Similar to the lifespan-

shortening circuit described above, this circuit is also formed by cholinergic

motor neurons and the intestine, and employs the same neurotransmitter ACh;

however, it recruits a different AChR GAR-3 to regulate a different type of

transcription factor HSF-1 in the intestine, leading to the opposite outcome in

lifespan (Figure 6D).

**Temporal control of ACR-6 and GAR-3 expression in the intestine at**

**different life stages**

The identification of two distinct neuroendocrine circuits provides a

potential explanation to the opposing effects of cholinergic motor neurons on

lifespan at different life stages. However, it prompts an important question as to

how these two circuits are temporally controlled at different stages of worm life.

One possibility is that the expression of some components in the two circuits

are temporally regulated at different life stages. To test this idea, we sought to

examine the expression of the two AChRs ACR-6 and GAR-3 in the intestine.

Using CRISPR/Cas9 genome editing, we inserted a mCherry tag at the C-

terminal end of the *acr-6* and *gar-3* gene to label the endogenous ACR-6 and

GAR-3 protein.  At the larval stage, ACR-6::mCherry fluorescence was detected

in the intestine, while GAR-3::mCherry was not (Figure 6A). Remarkably,

beginning at the adult stage, ACR-6::mCherry expression started to decrease

and became undetectable at Day 2 adulthood  (Figure 6B). On the other hand,

GAR-3::mCherry began to show expression at the adult stage and became

prominent at Day 2-4 adulthood (Figure 6A and 6B). We were unable to

examine GAR-3::mCherry expression at Day 5 adulthood and beyond, due to

the interference by increasingly bright gut auto-fluorescence even in the *glo-*

*4(ok623)* background with reduced gut auto-fluorescence[46]**.** Nevertheless,

these results unveil a clear temporal switch in the intestinal expression of ACR-

6 and GAR-3 at different life stages (Figure 6C). Though additional mechanisms

may also contribute, such a dichotomy in the intestinal expression of ACR-6 and

GAR-3 at larval and adult stages provides a potential mechanism that underlies

the temporal control of the two neuroendocrine circuits, leading two opposing

outcomes in lifespan.

**Discussion**

Research in the past decade has revealed an increasingly important role

of the nervous system in orchestrating a body-wide aging process by signaling

distal tissues, particularly the gut[3,4,11,12,14-19,30]. How the nervous system signals

the gut to regulate aging, however, is not well understood, and fwork in this area

has yielded some seemingly inconsistent data. This points to a model in which

nervous system-gut signaling can both positively and negatively regulate aging,

highlighting a complex role of the nervous system-gut signaling axis in

aging[17,20,30]. Nervous system-gut communications may differentially regulate

aging through spatial mechanisms. For example, different neurons may play

distinct roles in aging by positively and/or negatively regulating longevity via

secreting distinct neurotransmitters and neuropeptides[17,18,20,47-49]. By sharp

contrast, whether and how the nervous system temporally regulates aging is

poorly understood. Here, using the *C. elegans* motor nervous system as a

model, we identified a temporal mechanism by which the nervous system both

positively and negatively regulates lifespan by temporally controlling nervous

system-to-gut signaling at different life stages. Specifically, we identified two

distinct neuroendocrine signaling circuits by which cholinergic motor neurons

act to shorten lifespan in early life, whereas extending lifespan in mid-late life.

The two circuits employ the same neurotransmitter ACh, but recruit two different

AChRs ACR-6 and GAR-3 that acts in the intestine to regulate two different

transcription factors DAF-16 and HSF-1 to shorten and extend lifespan at early

and mid-late life stage, respectively.  To the best of our knowledge, this

represents the first such example illustrating how the nervous system may

differentially regulate lifespan by temporally controlling nervous system-to-gut

signaling.

One interesting observation is that the expression level of ACR-6 and

GAR-3 in the intestine features a temporal switch at the larval and adult stages.

The appearance followed by disappearance of ACR-6 expression in the

intestine at the larval and adult stages nicely correlate with the temporal window

under which cholinergic motor neurons shorten lifespan. This provides a

potential mechanism by which cholinergic motor neurons shorten lifespan in

early life. Remarkably, the temporal expression pattern of GAR-3 in the intestine

follows a reciprocal pattern: it is not expressed at the larval stage and only

appears at the adult stage, providing a potential mechanism for the lifespan-

extending effect exerted by cholinergic motor neurons in mid-late life. It should

be pointed out that the intestinal expression of GAR-3 emerges at Day 1

adulthood, earlier than the time point at which the beneficial role of cholinergic

motor neurons began to take effect. Several factors might have contributed to

this delay. For example, the lifespan-shortening signaling in the intestine might

not have been completely shut off during early adulthood, particularly

considering that the role of the transcription factor DAF-16 might be long-lasting.

This might antagonize the effect of the lifespan-extending signaling initiated by

GAR-3 in early adulthood.  Additionally, though GAR-3 expression can be

detected in early adulthood, it might take some time for it to fully engage other

signaling components, which might cause a delay for the lifespan-extending

signaling to take effect. Future studies are needed to elucidate the mechanism

that directs the temporal switch of ACR-6 and GAR-3 expression in the intestine.

Though the temporally-controlled dichotomous expression pattern of

ACR-6 and GAR-3 in the intestine provides a potential mechanism underlying

the opposing effects of cholinergic motor neurons on longevity, we do not

exclude the possibility that additional mechanisms may apply. For example, the

expression pattern of other components in the two neuroendocrine circuits may

undergo a similar temporal shift; similarly, their function may also be subjected

to temporal control. Our work provides an entry point to understand how

temporal mechanisms may contribute to the complex roles of the nervous

system as well as nervous system-gut communications in aging.

## Acknowledgments

We thank Wenxin Jia, Jiyong Meng, Yang Fu, Jinghan Shi and Guanghui Hu for technical assistance. Some strains were obtained from the CGC. This work was supported by the NSFC (32171146 to L.C. and 31130028, 31225011 and 31420103909 to J.L.) and funds from the University of Michigan.

## Declaration of interests

The authors declare that they have no competing interests.

## Author contributions

L.X., C.H. and L.C. conducted the experiments and analyzed the data. L.X., C.H., L.C., X.Z.S.X., and J.L. interpreted the data. L.C., X.Z.S.Z., and J.L. wrote the article with assistance from other authors.

## Data availability

The datasets generated and analyzed during this study are either included within the manuscript or are available from the authors upon request.

## MATERIALS AND METHODS

### Strains and genetics

Wild-type: N2. *Ex263a[Pacr-2::Tetx::sl2::yfp]. Is4a[Pacr-2::TIR1::sl2::yfp].*

*Is4a; Ex329a[Pacr-2::syntaxin(T254I)-degron::sl2::mcherry]. Ex294a[Pacr-*

*2::syntaxin(T254I)::sl2::mcherry]. unc-31(e169). unc-31; Ex294a. cat-2(e1112).*

*cat-2; Ex294a. tbh-1(n3247). tbh-1; Ex294a. tph-1(mg280). tph-1; Ex294a. eat-*

*4(ky5). eat-4; Ex294a. unc-25(e156). unc-25; Ex294a. tdc-1(n3419). tdc-1;*

*Ex294a. unc-17(e245). unc-17; Ex294a. acr-6(ok3117). acr-6; Ex294a. acr-*

*8(ok1240). acr-8; Ex294a. gar-2(ok520). gar-2; Ex294a. acr-6; Ex321a[Pges-*

*1::acr-6::sl2::yfp]. acr-6; Ex294a; Ex321a. acr-6; Ex324a[Prgef-1::acr-6::sl2::yfp].*

*acr-6; Ex294a; Ex324a. sid-1(qt9). sid-1; Ex263a. sid-1; Ex380a[Pvha-6::daf-16*

*(RNAi)+Pvha-6::sl2::mcherry]. sid-1; Ex263a; Ex380a. sid-1; Ex371a[Prgef-*

*1::daf-16 (RNAi)+Prgef-1::sl2::mcherry]. sid-1; Ex263a; Ex371a. Is4a;*

*Is5a[Pacr-2::syntaxin(T254I)-degron::sl2::mcherry]. unc-17; Is4a. unc-17; Is4a;*

*Is5a. gar-3(gk337); Is4a. gar-3; Is4a; Ex329a. sid-1; Is4a. sid-1; Is4a; Is5a.*

*sid-1; Is4a; Ex359a[Pges-1::gar-3(RNAi)+Pges-1::sl2::cfp]. sid-1; Is4a; Is5a;*

*Ex364a[Pges-1::gar-3(RNAi)+Pges-1::sl2::cfp]. sid-1; Is4a; Ex395a[Prgef-*

*1::gar-3(RNAi)+Prgef-1::sl2::cfp]. sid-1; Is4a; Is5a; Ex398a[Prgef-1::gar-*

*3(RNAi)+Prgef-1::sl2::cfp]. gar-2; Is4a. gar-2; Is4a; Ex329a. sid-1; Is4a;*

*Ex357a[Pges-1::gar-2(RNAi)+Pges-1::sl2::cfp]. sid-1; Is4a; Is5a; Ex363a[Pges-*

*1::gar-2(RNAi)+Pges-1::sl2::cfp]. sid-1; Is4a; Ex390a[Prgef-1::gar-*

*2(RNAi)+Prgef-1::sl2::cfp]. sid-1; Is4a; Is5a; Ex392a[Prgef-1::gar-*

*2(RNAi)+Prgef-1::sl2::cfp]. sid-1; Is4a; Ex373a[Pges-1:hsf-1(RNAi)+Pges-*

*1::sl2:cfp]. sid-1; Is4a; Is5a; Ex374a[Pges-1:hsf-1(RNAi)+Pges-1::sl2:cfp]. sid-*

*1; Is4a; Ex384a[Prgef-1::hsf-1(RNAi)+Prgef-1::sl2::cfp]. sid-1; Is4a; Is5a;*

*Ex385a[Prgef-1::hsf-1(RNAi)+Prgef-1::sl2::cfp]. acr-6(xu129a[acr-6::mCherry]).*

*gar-3(xu130a[gar-3::mCherry]).*

Mutant strains were outcrossed at least four times before use. The *acr-6*,

*acr-8*, *acr-15*, *acr-16*, *acr-19*, *acr-21*, *acr-24*, *gar-1*, *acc-1*, *acc-2*, *acc-4*, *deg-3*,

*lev-8*, *pbo-5*, *eat-2* and *unc-63* RNA interference (RNAi) clones were generated

in the laboratory, while other RNAi clones were from the Ahringer library and

were confirmed by sequencing.

**Worm synchronization**

Strains were grown at 20°C for at least three generations before lifespan

determination. Twenty Day 2 adult worms were transferred to fresh 60 mm

nematode growth medium (NGM) plates. The adult worms were removed from

the plates after laying eggs for 4 hours. The plates were placed at 20°C for two

days. L4 larva were used to start the normal lifespan or RNAi assays.

**Lifespan assay**

Lifespan assays were performed on 60 mm NGM plates at 20°C as

previously described [15]. ~120 worms of each genotype were used for lifespan

assays and transferred every other day to fresh NGM plates. Survival rate was

scored every 1 – 2 days. Worms were censored if they crawled off the plate,

bagged, or exhibited protruding vulva. In all cases, the first day of adulthood

was scored as day 1. Lifespan data was analyzed with GraphPad Prism 8

(GraphPad Software, Inc.) and IBM SPSS Statistics 21 (IBM, Inc.). Log-rank

(Kaplan-Meier) was used to calculate *P* values. All lifespan assays were

repeated at least twice. 5-Fluoro-2'-deoxyuridine (FUDR) was included in

assays involving TeTx transgene worms, *unc-31* and *unc-17* mutant worms,

which show a defect in egg laying.

**RNA interference (RNAi) assay**

RNAi was performed using the RNAi-compatible OP50 bacterial strain

OP50(xu363) and HT115 as previously described [50]. RNAi plates included

carbenicillin (25 µg/mL) and isopropyl-β-D-thiogalactoside (0.5 mM).

OP50(xu363) or HT115 bacteria with vector or RNAi plasmid were seeded on

RNAi plates 2 days before experiment.

**Auxin treatment**

Auxin treatment was performed by transferring worms to bacteria-

seeded NGM plates containing 0.5 mM auxin as previously described [33]. Auxin

was dissolved in cholesterol solution (25g/L) and KP buffer (2.5g/L) to avoid

additional ethanol. Cholesterol solution and KP buffer with auxin were added to

NGM agar that has cooled down to about 60°C before pouring the plates.

**Microscopy**

Different stage worms were anesthetized with 50 mM sodium azide

(Sigma) on 2% agarose pads. Confocal images were acquired by FLUOVIEW

FV3000 series of confocal laser scanning microscopes (Olympus) equipped

with a 100X objective as Z-stacks and were processed using ImageJ (NIH).

Maximum intensity Z-projection images were shown to illustrate the

endogenous ACR-6 and GAR-3 protein expression pattern. To quantify the

ACR-6 and GAR-3 protein expression level in the intestine, maximum intensity

Z-projection were used.

**CRISPR genome editing**

ACR-6::mCherry and GAR-3::mCherry are both knockin alleles made by

injecting RNP mixtures as described previously [51]. In brief, repair templates

were amplified using unmodified primers with 70 bp homology arms against the

target genes from linker-mCherry plasmid. The RNP mixture was assembled at

37°C for 15 min and contained Cas9 protein, tracrRNA, and crRNA (all from

IDT). crRNA corresponding DNA target sequences are 5'-

ATTCACGGATTGATAGCAAT-3' (*acr-6*), and 5'-

ATTCACGGATTGATAGCAAT-3' (*gar-3*). The double-strand donor templates

were melted right before adding them to the assembled RNPs [52]. Microinjection

quality was scored using a co-injection marker, PRF4::*rol-6 (su1006)* plasmid.

In general, 50 P0 were injected, singled, and maintained at 25°C for 3 days.

Two plates with the highest number of F1 rollers (usually more than 20 rollers

for a good injection) were selected, and 48 F1 rollers from each plate were

picked and singled for further genotyping [51]. The positive insertions were

confirmed by Sanger sequencing. Using this CRISPR-RNP technology, the

coding sequence of mCherry and a 27 bp linker sequence were inserted

immediately after the last amino acid codon of the *acr-6* gene on chromosome I

and of *gar-3* gene on chromosome V, respectively.

## References

1    Sawhney, R., Sehl, M. & Naeim, A. Physiologic aspects of aging: impact on
cancer management and decision making, part I. *Cancer J* **11**, 449-460,
doi:10.1097/00130404-200511000-00004 (2005).

2    Sehl, M., Sawhney, R. & Naeim, A. Physiologic aspects of aging: impact on
cancer management and decision making, part II. *Cancer J* **11**, 461-473,
doi:10.1097/00130404-200511000-00005 (2005).

3    Apfeld, J. & Kenyon, C. Cell nonautonomy of C. elegans daf-2 function in the
regulation of diapause and life span. *Cell* **95**, 199-210, doi:10.1016/s0092-
8674(00)81751-1 (1998).

4    Libina, N., Berman, J. R. & Kenyon, C. Tissue-specific activities of C. elegans
DAF-16 in the regulation of lifespan. *Cell* **115**, 489-502, doi:10.1016/s0092-
8674(03)00889-4 (2003).

5    Alcedo, J. & Kenyon, C. Regulation of C. elegans longevity by specific
gustatory and olfactory neurons. *Neuron* **41**, 45-55, doi:10.1016/s0896-
6273(03)00816-x (2004).

6    Murphy, C. T., Lee, S. J. & Kenyon, C. Tissue entrainment by feedback
regulation of insulin gene expression in the endoderm of Caenorhabditis elegans.
*Proc Natl Acad Sci U S A* **104**, 19046-19050, doi:10.1073/pnas.0709613104
(2007).

7    Medkour, Y., Svistkova, V. & Titorenko, V. I. Cell-Nonautonomous
Mechanisms Underlying Cellular and Organismal Aging. *Int Rev Cell Mol Biol*
**321**, 259-297, doi:10.1016/bs.ircmb.2015.09.003 (2016).

8    Savini, M. *et al.* Lysosome lipid signalling from the periphery to neurons
regulates longevity. *Nat Cell Biol* **24**, 906-916, doi:10.1038/s41556-022-00926-
8 (2022).

9    Allen, E., Ren, J., Zhang, Y. & Alcedo, J. Sensory systems: their impact on C.
elegans survival. *Neuroscience* **296**, 15-25,
doi:10.1016/j.neuroscience.2014.06.054 (2015).

10    Apfeld, J. & Kenyon, C. Regulation of lifespan by sensory perception in
Caenorhabditis elegans. *Nature* **402**, 804-809, doi:10.1038/45544 (1999).

11    Jeong, D. E., Artan, M., Seo, K. & Lee, S. J. Regulation of lifespan by
chemosensory and thermosensory systems: findings in invertebrates and their
implications in mammalian aging. *Front Genet* **3**, 218,
doi:10.3389/fgene.2012.00218 (2012).

12    Kenyon, C. J. The genetics of ageing. *Nature* **464**, 504-512,
doi:10.1038/nature08980 (2010).

13  Wolkow, C. A., Kimura, K. D., Lee, M. S. & Ruvkun, G. Regulation of C.
elegans life-span by insulinlike signaling in the nervous system. *Science* **290**,
147-150, doi:10.1126/science.290.5489.147 (2000).

14  Artan, M. *et al.* Food-derived sensory cues modulate longevity via distinct
neuroendocrine insulin-like peptides. *Genes & development* **30**, 1047-1057,
doi:10.1101/gad.279448.116 (2016).

15  Xiao, R. *et al.* A genetic program promotes C. elegans longevity at cold
temperatures via a thermosensitive TRP channel. *Cell* **152**, 806-817,
doi:10.1016/j.cell.2013.01.020 (2013).

16  Zhang, B. *et al.* Environmental Temperature Differentially Modulates C. elegans
Longevity through a Thermosensitive TRP Channel. *Cell reports* **11**, 1414-1424,
doi:10.1016/j.celrep.2015.04.066 (2015).

17  Zhang, B., Jun, H., Wu, J., Liu, J. & Xu, X. Z. S. Olfactory perception of food
abundance regulates dietary restriction-mediated longevity via a brain-to-gut
signal. *Nat Aging* **1**, 255-268, doi:10.1038/s43587-021-00039-1 (2021).

18  Burkewitz, K. *et al.* Neuronal CRTC-1 governs systemic mitochondrial
metabolism and lifespan via a catecholamine signal. *Cell* **160**, 842-855,
doi:10.1016/j.cell.2015.02.004 (2015).

19  Lee, S. J. & Kenyon, C. Regulation of the longevity response to temperature by
thermosensory neurons in Caenorhabditis elegans. *Curr Biol* **19**, 715-722,
doi:10.1016/j.cub.2009.03.041 (2009).

20  Zhang, B. *et al.* Brain-gut communications via distinct neuroendocrine signals
bidirectionally regulate longevity in C. elegans. *Genes & development* **32**, 258-
270, doi:10.1101/gad.309625.117 (2018).

21  Mutlu, A. S., Gao, S. M., Zhang, H. & Wang, M. C. Olfactory specificity
regulates lipid metabolism through neuroendocrine signaling in Caenorhabditis
elegans. *Nat Commun* **11**, 1450, doi:10.1038/s41467-020-15296-8 (2020).

22  Yin, J. A. *et al.* Genetic variation in glia-neuron signalling modulates ageing
rate. *Nature* **551**, 198-203, doi:10.1038/nature24463 (2017).

23  Frakes, A. E. *et al.* Four glial cells regulate ER stress resistance and longevity
via neuropeptide signaling in C. elegans. *Science* **367**, 436-440,
doi:10.1126/science.aaz6896 (2020).

24  Zullo, J. M. *et al.* Regulation of lifespan by neural excitation and REST. *Nature*
**574**, 359-364, doi:10.1038/s41586-019-1647-8 (2019).

25  Xia, Y. *et al.* Neuronal C/EBPbeta/AEP pathway shortens life span via selective
GABAnergic neuronal degeneration by FOXO repression. *Sci Adv* **8**, eabj8658,
doi:10.1126/sciadv.abj8658 (2022).

26    Wong, C. O. *et al.* Regulation of longevity by depolarization-induced activation
598            of PLC-beta-IP(3)R signaling in neurons. *Proc Natl Acad Sci U S A* **118**,
599            doi:10.1073/pnas.2004253118 (2021).

600    27    Liu, J. *et al.* Functional aging in the nervous system contributes to age-
601            dependent motor activity decline in C. elegans. *Cell metabolism* **18**, 392-402,
602            doi:10.1016/j.cmet.2013.08.007 (2013).

603    28    Zhou, B. *et al.* Mitochondrial Permeability Uncouples Elevated Autophagy and
604            Lifespan Extension. *Cell* **177**, 299-314 e216, doi:10.1016/j.cell.2019.02.013
605            (2019).

606    29    Ching, T. T. *et al.* Short-term enhancement of motor neuron synaptic exocytosis
607            during early aging extends lifespan in Caenorhabditis elegans. *Experimental*
608            *biology and medicine* **245**, 1552-1559, doi:10.1177/1535370220950639 (2020).

609    30    Chun, L. *et al.* Metabotropic GABA signalling modulates longevity in C.
610            elegans. *Nat Commun* **6**, 8828, doi:10.1038/ncomms9828 (2015).

611    31    Link, E. *et al.* Tetanus toxin action: inhibition of neurotransmitter release linked
612            to synaptobrevin proteolysis. *Biochem Biophys Res Commun* **189**, 1017-1023,
613            doi:10.1016/0006-291x(92)92305-h (1992).

614    32    Lagow, R. D. *et al.* Modification of a hydrophobic layer by a point mutation in
615            syntaxin 1A regulates the rate of synaptic vesicle fusion. *PLoS Biol* **5**, e72,
616            doi:10.1371/journal.pbio.0050072 (2007).

617    33    Zhang, L., Ward, J. D., Cheng, Z. & Dernburg, A. F. The auxin-inducible
618            degradation (AID) system enables versatile conditional protein depletion in C.
619            elegans. *Development* **142**, 4374-4384, doi:10.1242/dev.129635 (2015).

620    34    Li, C. & Kim, K. Neuropeptides. *WormBook*, 1-36,
621            doi:10.1895/wormbook.1.142.1 (2008).

622    35    Rand, J. B. Acetylcholine. *WormBook*, 1-21, doi:10.1895/wormbook.1.131.1
623            (2007).

624    36    Jorgensen, E. M. Gaba. *WormBook*, 1-13, doi:10.1895/wormbook.1.14.1 (2005).

625    37    Chase, D. Biogenic amine neurotransmitters in C. elegans. *WormBook*,
626            doi:10.1895/wormbook.1.132.1 (2007).

627    38    Speese, S. *et al.* UNC-31 (CAPS) is required for dense-core vesicle but not
628            synaptic vesicle exocytosis in Caenorhabditis elegans. *J Neurosci* **27**, 6150-
629            6162, doi:10.1523/JNEUROSCI.1466-07.2007 (2007).

630    39    Lin, X. G. *et al.* UNC-31/CAPS docks and primes dense core vesicles in C.
631            elegans neurons. *Biochem Biophys Res Commun* **397**, 526-531,
632            doi:10.1016/j.bbrc.2010.05.148 (2010).

40    Berwin, B., Floor, E. & Martin, T. F. J. CAPS (Mammalian UNC-31) Protein
Localizes to Membranes Involved in Dense-Core Vesicle Exocytosis. *Neuron*
**21**, 137-145, doi:10.1016/s0896-6273(00)80521-8 (1998).

41    Alfonso, A., Grundahl, K., Duerr, J. S., Han, H. P. & Rand, J. B. The
Caenorhabditis elegans unc-17 gene: a putative vesicular acetylcholine
transporter. *Science* **261**, 617-619, doi:10.1126/science.8342028 (1993).

42    Yemini, E. *et al.* NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal
Identification in C. elegans. *Cell* **184**, 272-288 e211,
doi:10.1016/j.cell.2020.12.012 (2021).

43    Chan, J. P. *et al.* Extrasynaptic muscarinic acetylcholine receptors on neuronal
cell bodies regulate presynaptic function in Caenorhabditis elegans. *J Neurosci*
**33**, 14146-14159, doi:10.1523/JNEUROSCI.1359-13.2013 (2013).

44    Douglas, Peter M. *et al.* Heterotypic Signals from Neural HSF-1 Separate
Thermotolerance from Longevity. *Cell reports* **12**, 1196-1204,
doi:10.1016/j.celrep.2015.07.026 (2015).

45    Hsu, A. L., Murphy, C. T. & Kenyon, C. Regulation of aging and age-related
disease by DAF-16 and heat-shock factor. *Science* **300**, 1142-1145,
doi:10.1126/science.1083701 (2003).

46    Hermann, G. J. *et al.* Genetic Analysis of Lysosomal Trafficking
inCaenorhabditis elegans. *Molecular Biology of the Cell* **16**, 3273-3288,
doi:10.1091/mbc.e05-01-0060 (2005).

47    Higuchi-Sanabria, R. *et al.* Divergent Nodes of Non-autonomous UPR(ER)
Signaling through Serotonergic and Dopaminergic Neurons. *Cell reports* **33**,
108489, doi:10.1016/j.celrep.2020.108489 (2020).

48    Zhang, Q. *et al.* The Mitochondrial Unfolded Protein Response Is Mediated
Cell-Non-autonomously by Retromer-Dependent Wnt Signaling. *Cell* **174**, 870-
883 e817, doi:10.1016/j.cell.2018.06.029 (2018).

49    Ozbey, N. P. *et al.* Tyramine Acts Downstream of Neuronal XBP-1s to
Coordinate Inter-tissue UPR(ER) Activation and Behavior in C. elegans. *Dev*
*Cell* **55**, 754-770 e756, doi:10.1016/j.devcel.2020.10.024 (2020).

50    Xiao, R. *et al.* RNAi Interrogation of Dietary Modulation of Development,
Metabolism, Behavior, and Aging in C. elegans. *Cell reports* **11**, 1123-1133,
doi:10.1016/j.celrep.2015.04.024 (2015).

51    Ghanta, K. S., Ishidate, T. & Mello, C. C. Microinjection for precision genome
editing in Caenorhabditis elegans. *STAR Protoc* **2**, 100748,
doi:10.1016/j.xpro.2021.100748 (2021).

52  Ghanta, K. S. & Mello, C. C. Melting dsDNA Donor Molecules Greatly
Improves Precision Genome Editing in Caenorhabditis elegans. *Genetics* **216**,
643-650, doi:10.1534/genetics.120.303564 (2020).

**Figure legends**

**Figure 1. Cholinergic motor neurons differentially regulate lifespan at different stages of worm life.**

(A) Ablating the output of cholinergic motor neurons promotes survival at early life stage, whereas inhibits survival at later stages. Tetanus toxin (TeTx) was expressed as a transgene in cholinergic motor neurons using *acr-2* promoter to block exocytosis from these neurons.

(B) Promoting the output of cholinergic motor neurons shortens lifespan. The gain-of-function form of *Drosophila* syntaxin(T254I) was expressed as a transgene in cholinergic motor neurons using *acr-2* promotor to potentiate exocytosis from these neurons.

(C) Schematic describing the assay. The gain-of-function form of *Drosophila* syntaxin(T254I) fused with a degron tag was expressed as a transgene in cholinergic motor neurons to promote exocytosis from these neurons. Transgenic worms were then crossed with a line stably expressing TIR1 in these neurons. Expression of Syntaxin(T254I) was suppressed by 0.5 mM auxin treatment and induced by subsequently removing auxin.

(D) *Syntaxin(T254I)* transgene has slight effect on lifespan when suppressing its expression throughout the entire worm life.

(E) *Syntaxin(T254I)* transgene shortens lifespan when inducing its expression throughout the entire worm life.

(F) *Syntaxin(T254I)* transgene shortens lifespan when inducing its expression from egg to L4 larval stage.

(G-I) *Syntaxin(T254I)* transgene modestly affects lifespan when inducing its

expression from L4 larval stage (G), Day 3 (H) or Day 5 (I) adulthood.

(J-K) *Syntaxin(T254I)* transgene extends lifespan when inducing its expression

from Day 7 (J), Day 9 (K) or Day 11 (L) adulthood. See Table S1 for detailed

statistical analysis of lifespan data. Logrank (Kaplan-Meier) was used to

calculate p values.

**Figure 2. ACh and the nicotinic AChR ACR-6 mediate the lifespan-**

**shortening effect of cholinergic motor neurons in early life.**

(A) *Syntaxin(T254I)* transgene shortens lifespan.

(B) Loss of *unc-31* exhibits no defect on the ability of *syntaxin(T254I)* transgene

to shorten lifespan.

(C) Loss of *unc-17* blocks the ability of *syntaxin(T254I)* transgene to shorten

lifespan.

(D-F) Loss of *acr-6* blocks the ability of *syntaxin(T254I)* transgene to shorten

lifespan (D), a phenotype rescued by transgenic expression of wild-type *acr-6*

gene in the intestine (E), but not in neurons (F). The *ges-1* and *rgef-1* promoters

were used to drive *acr-6* gene expression in the intestine and neurons,

respectively. See Table S1 for detailed statistical analysis of lifespan data.

Logrank (Kaplan-Meier) was used to calculate p values.

**Figure 3. The FOXO transcription factor DAF-16 is required in the intestine**

**for cholinergic motor neurons to regulate lifespan in early life.**

(A) *Syntaxin(T254I)* transgene shortens lifespan.

(B-E) RNAi of *daf-16* does not further shorten the lifespan of *syntaxin(T254I)*

transgenic worms, while RNAi of *hsf-1* (C), *skn-1* (D) and *pha-4* (E) does.

(F) qPCR analysis of DAF-16 target genes. qPCR reactions were run in

triplicates for each genotype. Each experiment was repeated three times. Error

bars represent s.e.m. ****$p < 0.0001$ (ANOVA with Bonferroni's test).

(G-H) RNAi of *daf-16* in the intestine (H) abolishes the ability of TeTx transgene

to regulate lifespan (G). dsRNA against *daf-16* was expressed as a transgene

specifically in the intestine using *vha-6* promoter. These experiments were

carried out in a *sid-1* mutant background where systemic effect of RNAi is

absent. See Table S1 for detailed statistical analysis of lifespan data. Log-rank

(Kaplan-Meier) was used to calculate p values.

**Figure 4. ACh and the intestinal muscarinic AChR GAR-3 mediate the**

**lifespan-extending effect of cholinergic motor neurons in mid-late life.**

(A) Enhancing the output of cholinergic motor neurons in mid-late life extends

lifespan. The *syntaxin(T254I)* transgene was induced from Day 7 adulthood as

described in Figure 1C.

(B-C) Loss of *unc-17* (B) or *gar-3* (C) blocks the ability of cholinergic motor

neurons to extend lifespan in mid-late life.

(D-F) RNAi of *gar-3* in the intestine (E), but not in neurons (F), abolishes the

ability of cholinergic motor neurons to extend lifespan in mid-late life (D). dsRNA

against *gar-3* was expressed as a transgene specifically in the intestine and

neurons using *ges-1* and *rgef-1* promoter, respectively. These experiments

were carried out in a *sid-1* mutant background where systemic effect of RNAi is

absent. See Table S1 for detailed statistical analysis of lifespan data. Log-rank

(Kaplan-Meier) was used to calculate p values.

**Figure 5. The transcription factor HSF-1 is required in the intestine for**

**cholinergic motor neurons to regulate lifespan in mid-late life.**

(A-B) Enhancing the output of cholinergic motor neurons in mid-late life extends

lifespan (A). This effect was fully suppressed by RNAi of *hsf-1* (B). The

*syntaxin(T254I)* transgene was induced from Day 7 adulthood as described in

Figure 1C.

(C) qPCR analysis of HSF-1 target genes. qPCR reactions were run in

triplicates for each genotype. Each experiment was repeated three times. Error

bars represent s.e.m. ****p< 0.0001 (ANOVA with Bonferroni's test).

(D-F) RNAi of *hsf-1* in the intestine (E), but not in neurons (F), abolishes the

ability of cholinergic motor neurons to extend lifespan in mid-late life (D). dsRNA

against *hsf-1* was expressed as a transgene specifically in the intestine and

neurons using *ges-1* and *rgef-1* promoter, respectively. These experiments

were carried out in a *sid-1* mutant background where systemic effect of RNAi is

absent. See Table S1 for detailed statistical analysis of lifespan data. Logrank

(Kaplan-Meier) was used to calculate p values.

**Figure 6. Temporal control of ACR-6 and GAR-3 expression in the**

**intestine at different life stages.**

(A-B) The expression pattern of endogenous ACR-6 and GAR-3 proteins at L3

Larva stage (A) and Day 2 adult stage (B). Arrowheads point to the area of the

intestine. Scale bars, 20 μm.

(C) Quantification curves summarizing the data in (A) and (B). Error bars

represent s.e.m. n ≥ 10.

(D) A schematic model illustrating the two cholinergic motor neurons-to-gut

signaling circuits that act in early life to shorten lifespan and in mid-late life to

extend lifespan.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

**A**

*acr-6::mcherry* larva (L3)

DIC | mCherry

*gar-3::mcherry* larva (L3)

DIC | mCherry

**B**

*acr-6::mcherry* adult (Day 2)

DIC | mCherry

*gar-3::mcherry* adult (Day 2)

DIC | mCherry

**C**

Fluorescence intensity vs Age

Legend: *gar-3::mcherry*, *acr-6::mcherry*

**D**

Motor Neurons → ACh

early life: ACR-6 → DAF-16 ⊣ Longevity (Intestine)

mid-late life: GAR-3 → HSF-1 → Longevity (Intestine)