Chemogenetic stimulation of phrenic motor output and diaphragm activity

Ethan S. Benevides[1,2], Prajwal P. Thakre[1,2,3], Sabhya Rana[1,2,3], Michael D. Sunshine[1,2,3], Victoria N. Jensen[1,2,3], Karim Oweiss[3,4] and David D. Fuller[1,2,3]

[1] Department of Physical Therapy, University of Florida, Gainesville, FL, 32601
[2] Breathing Research and Therapeutics Center, University of Florida, Gainesville, FL, 32601
[3] McKnight Brain Institute, University of Florida, Gainesville, FL, 32601
[4] Department of Electrical and Computer Engineering, University of Florida, Gainesville, FL, 32601

Running Title:	Chemogenetic stimulation of phrenic motor output and diaphragm activity
Article Type:	Original Research
Number of Figures:	9
Number of Tables:	6

Corresponding Author: David Fuller, Ph.D
Professor
Department of Physical Therapy
University of Florida
PO Box 100514, UFHSC
Gainesville, FL, 32610-0154
Phone (352) 273-6634
dfuller@phhp.ufl.edu

ORCIDs:
Ethan S. Benevides: 0000-0002-2472-0606
Thakre P. Prajwal: 0000-0001-5440-7023
Sabhya Rana: 0000-0002-1303-6614
Michael D. Sunshine: 0000-0002-7133-4185
Victoria N. Jensen: 0000-0002-6374-6323
Karim Oweiss: 0000-0002-1175-7258
David D. Fuller: 0000-0001-5013-5972

Author Contributions: E.S.B.: design of the experiment, data acquisition, analysis, and interpretation; initial draft of the manuscript. T.P.P: design of the study, data acquisition, analysis, and interpretation; manuscript editing. S.R.: design of the study, acquisition of the data, analysis, and interpretation; review of manuscript draft. M.D.S: data analysis and interpretation; review of manuscript draft. V.N.J.: acquisition; review of manuscript draft. K.O.: provided ChAT-Cre rats; review of manuscript draft. D.D.F.: conception and design of the study, interpretation of the data, revisions, and final approval of the manuscript. All authors read and approved the submitted version.

Authors' Disclosure Statement: Authors declare no competing financial interests.

Funding Information: Support for this work was provided by the National Institutes of Health: 5R01HD052682-14 (DDF), and a Graduate School Funding Award from the University of Florida (ESB).

Data and Code Availability: The datasets along with MATLAB and R code generated during the current study are available from the corresponding author upon reasonable request.

Animal Studies: All procedures described in this manuscript involving rats, mice, and tissue were approved by the University of Florida Institutional Animal Care and Use Committee (protocol #202107438) and in strict accordance with the US National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals.

Abbreviations: DREADD (designer receptors exclusively activated by designer drugs); EMG (electromyography), AUC (area under the curve); J60 (JHU37160); AAV (adeno-associated virus); VT (tidal volume); ChAT (choline acetyltransferase); Cre (Cre recombinase)
Keywords: DREADD, Chemogenetics, Phrenic, Diaphragm EMG, Plethysmography, Adeno-associated virus, Respiratory neural control.

ABSTRACT

Impaired diaphragm activation contributes to morbidity and mortality in many neurodegenerative diseases

and neurologic injuries. We conducted experiments to determine if expression of an excitatory DREADD

(designer receptors exclusively activation by designer drugs) in the mid-cervical spinal cord would enable

respiratory-related activation of phrenic motoneurons to increase diaphragm activation. Wild type

(C57/bl6) and ChAT-Cre mice received bilateral intraspinal (C4) injections of an adeno-associated virus

(AAV) encoding the hM3D(Gq) excitatory DREADD. In wild type mice, this produced non-specific

DREADD expression throughout the mid-cervical ventral horn. In ChAT-Cre mice, a Cre-dependent viral

construct was used to drive DREADD expression in C4 ventral horn motoneurons, targeting the phrenic

motoneuron pool. Diaphragm EMG was recorded during spontaneous breathing at 6-8 weeks post-AAV

delivery. The selective DREADD ligand JHU37160 (J60) caused a bilateral, sustained (>1 hr) increase in

inspiratory EMG bursting in both groups; the relative increase was greater in ChAT-Cre mice. Additional

experiments in a ChAT-Cre rat model were conducted to determine if spinal DREADD activation could

increase inspiratory tidal volume (VT) during spontaneous breathing without anesthesia. Three to four

months after intraspinal (C4) injection of AAV driving Cre-dependent hM3D(Gq) expression,

intravenous J60 resulted in a sustained (>30 min) increase in VT assessed using whole-body

plethysmography. Subsequently, direct nerve recordings confirmed that J60 evoked a >50% increase in

inspiratory phrenic output. The data show that mid-cervical spinal DREADD expression targeting the

phrenic motoneuron pool enables ligand-induced, sustained increases in the neural drive to the

diaphragm. Further development of this technology may enable application to clinical conditions

associated with impaired diaphragm activation and hypoventilation.

INTRODUCTION

Many respiratory disorders are associated with reduced or impaired activation of respiratory motoneurons. Neurologic injuries (e.g., traumatic spinal cord injury, stroke) and neurodegenerative conditions (e.g., ALS, Pompe disease) will often result in decreased respiratory motor output, including impaired activation of the phrenic motoneurons which innervate the diaphragm[1-6]. Another prominent example is obstructive sleep apnea, in which pharyngeal motoneurons have reduced output during sleep[7]. Treatment options that target increased respiratory motoneuron activation to improve breathing ability are limited. However, designer receptors exclusively activated by designer drugs (DREADDs) may have use in this regard[8]. Structurally derived from naturally occurring G-protein coupled receptors, DREADDs have been engineered to respond exclusively to exogenous ligands that are otherwise biologically inert[9-11]. This provides a means to selectively stimulate cells expressing the DREADD. Prior studies have used DREADDs to stimulate upper airway muscle activation during breathing[8]. For example, following expression of DREADDs in murine hypoglossal motoneurons, tongue electromyogram (EMG) activity can be increased using DREADD ligands[12-16]. This response is functionally beneficial as shown by increased patency of the upper airway[12].

The present study focused on chemogenetic activation of the phrenic neuromuscular system. Phrenic motoneurons provide motor innervation of the diaphragm muscle and are located in the mid-cervical (C3-5) spinal cord[17]. We tested the hypothesis that expressing DREADDs in the mid-cervical spinal cord would enable systemic (intravenous) delivery of a selective DREADD ligand to produce sustained increases in the respiratory-related activation of the diaphragm. In doing so, we addressed two important questions. First, we determined if effective diaphragm activation requires focal DREADD expression targeting phrenic motor neurons, or if non-specific expression in the immediate vicinity of the phrenic motor nucleus would be sufficient. This question derives from prior studies of cervical spinal cord stimulation. A compelling body of work, with studies in multiple species, demonstrates that non-specific activation of cervical spinal networks can be highly effective at increasing diaphragm activation[18-20]. One

theory to explain this result is that a general increase in excitability of cervical propriospinal networks

leads to increased phrenic motoneuron activation[21-23]. There is also evidence that phrenic motoneurons

can integrate multiple synaptic inputs in a manner that produces orderly recruitment[18]. Accordingly,

DREADD-induced activation of mid-cervical neurons or networks may be sufficient for ligand-induced

diaphragm activation. On the other hand, DREADD expression may need to be restricted to phrenic

motoneurons if the goal is to produce inspiratory-related diaphragm activation. To address this question,

we studied diaphragm responses in two mouse models: 1) a wild type model in which DREADDs were

non-specifically expressed in the C3-5 spinal cord, encompassing mid-cervical interneurons and

motoneurons, and 2) a choline acetyltransferase (ChAT)-Cre transgenic model in which DREADD

expression was restricted to ChAT-positive neurons in the anterior C3-5 spinal cord, targeting the phrenic

motoneuron pool.

The second question we addressed was if phrenic motoneuron activation via cervical spinal cord

DREADDs could produce a sustained increase in inspiratory tidal volume in unanesthetized,

spontaneously breathing animals. While the previous results from the hypoglossal motor system[12,14-16]

provide a proof-of-concept that DREADDs can stimulate respiratory motoneuron activity, whether a

sustained increases in tidal volume could be evoked by expressing DREADDs in phrenic motoneuron

was uncertain. For example, during spontaneous breathing, a DREADD-induced increase in phrenic

motoneuron excitability, and thus diaphragm activation, could be rapidly offset by decreases in

bulbospinal neural drive to the phrenic motor pool, secondary to reduced arterial CO_2 or increased vagal-

mediated inhibition. An increase in diaphragm activation could also trigger a decrease in accessory

respiratory muscle activation, thereby attenuating or preventing increases in tidal volume. Lastly, data

from the hypoglossal motor system[12,14-16], as well as our initial results in the anesthetized mouse indicated

that both phasic (i.e., during the inspiratory period) and tonic (i.e., occurring across the respiratory cycle)

activation of the diaphragm would increase after DREADD activation, and how this would impact tidal

volume was not clear. Accordingly, we studied ChAT-Cre rats using whole body plethysmography and

direct measure of phrenic motor output via nerve recordings. The plethysmography studies allowed us to

determine if DREADD activation of phrenic motoneurons causes a sustained increased in tidal volume

and ventilation during spontaneous breathing in the unanesthetized rat. The nerve recordings were done

under anesthesia and enabled direct quantification of the impact of DREADD activation on neural drive

of the diaphragm while controlling variables including arterial CO_2 and lung volume.

Collectively the results of this work demonstrate that mid-cervical spinal DREADD expression

enables the J60 ligand to produce a sustained increase in the neural drive to the diaphragm, producing an

increase in tidal volume during spontaneous breathing. Further development of this technology may

enable application to clinical conditions associated with impaired diaphragm activation and

hypoventilation.

METHODS

Animals. Experiments were carried out using C5/bl6, wild type mice (Taconic), ChAT-Cre transgenic mice (B6.129S6-Chattm2(cre)Lowl/J; Jackson Laboratories), and ChAT-Cre transgenic rats (LE-Tg(Chat-Cre)5.1Deis; Rat Resource & Research Center). Animals were singly housed in a controlled environment (12 h light–dark cycle) with food and water *ad libitum*. All experiments were conducted in accordance with the NIH Guidelines Concerning the Care and Use of Laboratory Animals and were approved by the University of Florida Institutional Animal Care and Usage Committee (protocol #202107438).

Intraspinal injections. An adeno-associated viral vector (AAV) encoding the gene for the excitatory DREADD, hM3D(Gq) was delivered to the mid-cervical spinal cord. Surgery was performed under aseptic conditions. Mice were anesthetized with isoflurane (induction: 3-4% isoflurane; maintenance: 2-3% isoflurane in 100% O_2) while rats were anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) delivered intraperitoneally. Animals were placed prone on a circulating water heating pad to maintain body temperature. A longitudinal incision was made starting at the base of the skull and extending caudally. The underlying back musculature was opened from the base of the skull to spinal segment C6. Using a micro-curette, the muscle and connective tissue overlying laminae C3 to C5 were removed. A laminectomy of the C4 dorsal lamina exposed the dura mater below. A bilateral durotomy was then performed exposing the spinal cord. A Hamilton syringe (34-gauge needle) held in a Kopf stereotaxic frame was as used to inject 1 μl of AAV9-hSyn-DIO-hM3D(Gq)-mCherry (ChAT-Cre mice and rats; titer: 2.07×10^{12} vg/mL) or AAV9-hSyn-HA-hM3D(Gq)-mCherry (C57/bl6 mice; titer: 2.44×10^{13} vg/mL), bilaterally into the ventral horns at C4. Injections were made 0.5 mm lateral to the spinal midline at a depth of 0.9 mm for mice[24] and 1 mm lateral to midline at a depth of 1.5 mm for rats[25].The needle was left to dwell for 5 minutes. Following injections, the overlying muscle and fascia were sutured with absorbable suture, the skin closed, and the animal returned to its home cage. Animals received a post-operative analgesia regiment of subcutaneous buprenorphine (1 mg/kg; slow-release formulation) and carprofen (5 mg/kg) for the first three days after surgery.

Diaphragm EMG recordings. Recordings were conducted using wild type (n= 11; n= 7 females) and

ChAT-Cre mice (n= 9; n= 6 females; n= 1 excluded from analysis), 6-8 weeks following intraspinal

injections of AAV-DREADD. Animals were anesthetized with 2-3% isoflurane in a closed chamber and

then placed supine on a closed loop heating pad to maintain rectal temperature at 37 ± 0.5 °C (model 700

TC-1000, CWE Inc.). Mice spontaneously inhaled 2% isoflurane in 100% O_2 for the duration of the

experiment.

A laparotomy was performed and two sets of 50 μm tungsten wires were placed in the mid-costal

region of the left and right hemi-diaphragm. The tips of each wire were de-insulated, bent into small

hooks, and inserted through the diaphragm approximately 3 mm apart. The recorded EMG signals were

amplified (1000x) and filtered (100–1000 Hz) using a differential amplifier (A–M systems model 1700).

Signals were digitized at 10 kS/s using a Power 1401 (CED, Cambridge, UK).

Once a stable plane of anesthesia was reached, animals underwent a 10-minute recording to

establish baseline diaphragm EMG parameters. Subsequently, animals received injections of vehicle (100

μl of saline delivered intraperitoneally (IP) followed by a 20-minute recording. Animals then received an

intraperitoneal injection of the selective DREADD agonist, JHU37160 (J60; 0.1 mg/kg, HB6261,

HelloBio), and recordings continued for 90 minutes. At the conclusion of each experiment, animals

underwent transcardial perfusion with saline followed by 4% paraformaldehyde. Following perfusion,

spinal cords were harvested for histological analysis.

Whole body plethysmography. ChAT-Cre rats (n= 9; n= 3 females) were studied using flow-through

whole body plethysmography 14-16 weeks after intraspinal delivery of AAV9-hSyn-DIO-hM3D(Gq)-

mCherry, as described above. Unanesthetized rats were sealed into the Plexiglas plethysmograph with

airflow maintained at 6 L/min for the duration of the recording. The recording protocol consisted of a 40-

minute acclimation period (inspired air: 21% O_2, 79% N_2), followed by a 7-minute ventilatory challenge

(10% O_2, 7% CO_2, 83% N_2) and a 10-minute normoxic recovery period (21% O_2, 79% N_2). Subsequently,

rats underwent a 20-minute long, pre-vehicle, baseline under normoxic conditions (21% O_2, 79% N_2)

followed by intravenous infusion of the J60 vehicle (saline). Following vehicle infusion recording

continued for 30-minutes followed by a 7-minute ventilatory challenge (10% O_2, 7% CO_2, 83% N_2) and

10-minutes of normoxic breathing (21% O_2, 79% N_2). After an additional 20-minute pre-J60 baseline

(21% O_2, 79% N_2), an intravenous infusion of the J60 ligand was given (0.1 mg/ml) and recordings

continued for 30-minute followed by a final ventilatory challenge (10% O_2, 7% CO_2, 83% N_2). The

ventilatory challenges were performed to assess the ability to increase breathing.

Phrenic nerve recordings. Two-to-eight-weeks following plethysmography recordings, bilateral phrenic

nerve recordings were performed. This procedure was done to directly assess the effect of DREADD

activation on phrenic motor output under rigorously controlled experimental conditions. Anesthesia was

induced by placing the rat in a closed chamber to inhale 3% isoflurane in 100% O_2. Rats were then moved

onto a closed-loop heating pad set to maintain rectal temperature at $37 \pm 1°C$ (model 700 TC-1000, CWE

Inc.). Isoflurane anesthesia was maintained using a nose cone. Once a surgical plane of anesthesia was

reached as evidenced by loss of corneal reflexes and hindlimb withdrawal, rats were tracheotomized and

ventilated (VentElite, model 55-7040; Harvard Apparatus Inc.) with a gas mixture of 50% O_2, 1% CO_2,

balanced with N_2. End-tidal CO_2 was maintained at 45-47 mmHg throughout the surgery and

experimental protocol (Capnogard; Novametrix). Ventilator frequency was maintained between 65 and 75

breaths/min, and tidal volume was set at 7 mL/kg[26]. The vagus nerves were transected bilaterally to

prevent entrainment of phrenic efferent output with the ventilator.

A tail vein catheter was placed to allow for intravenous infusion of urethane anesthesia,

supplementary fluids, and the J60 ligand. Animals were slowly converted from inhaled isoflurane to

urethane anesthesia (2.1 g/kg at 6 mL/hr; IV). During this conversion, the depth of anesthesia was

consistently monitored by evaluating the pedal withdrawal reflex. Following administration of the full

urethane dose, a mixture of 8.4% sodium bicarbonate and lactated Ringer's was administered (2 mL/h;

IV) to maintain acid-base balance. To prevent movements and EMG contamination of the phrenic

neurogram pancuronium bromide was administered (3 mg/kg IV, Sigma-Aldrich, St Louis) to achieve

neuromuscular blockade. A catheter (polyethylene tubing; PE 50; Intramedic) was placed in the femoral

artery to enable monitoring of arterial blood pressure via a transducer amplifier (TA-100, CWE) and

allow withdrawal of arterial blood samples (65 µL) for measurement of partial pressure of CO_2 ($PaCO_2$)

and O_2 (PaO_2), pH, and base excess (ABL 90 Flex, Radiometer; Copenhagen, Denmark).

The phrenic nerves were exposed bilaterally using a dorsal approach as described previously[27,28].

Briefly, a midline incision was made at the base of the skull extending to spinal level T2. The muscles

connecting the shoulder blades to the spinal column were separated to expose the phrenic nerves. The

phrenic nerves were isolated, cut distal to the spinal cord and suctioned into a custom-made glass

electrodes filled with 0.9% saline solution. Phrenic nerve activity was amplified (10 kHz) using a

differential AC amplifier (Model 1700, A-M systems, Everett, WA), band-pass filtered (100Hz-3 kHz),

and digitized at 25ks/second (Power 1401, CED).

At the beginning of the experiment, the apneic threshold was determined by slowly reducing the

inspired CO_2 until phrenic nerve inspiratory activity ceased for 60 seconds. The recruitment threshold was

established by slowly increasing the inspired CO_2 until phrenic bursting returned. The end tidal CO_2

($ETCO_2$) was then maintained 2-3 mmHg above the recruitment threshold for the duration of the

experiment. After achieving a stable phrenic nerve recording and blood gases a 15-minute-long baseline

recording was collected (50% O_2, 3% CO_2) followed by a brief, 5-minute exposure to hypoxia (11.5% O_2,

3% CO_2) and 10–15-minute recovery period (50% O_2, 3% CO_2). Subsequently, intravenous infusion of

vehicle (saline) was given followed by a 15-minute recording period. The J60 ligand (0.1 mg/kg) was

then administered intravenously over a 2 minute infusion period followed by a 100-minute recording

period.

Arterial blood samples were collected at specific intervals: initially at baseline, during the last

minute of each hypoxia episode, 15 minutes post vehicle administration, and subsequently at 20, 40, 60,

80, and 100 minutes post J60 administration. Baseline blood gas values served as references to assess if

further arterial samples were isocapnic. To keep end-tidal CO_2 and $PaCO_2$ near baseline (within $\pm$ 2.0

mmHg), minor adjustments to inspired CO_2 and ventilation rate were made as needed. PaO_2 was kept

above 150 mmHg, except during hypoxia; if it dropped below, O_2 intake was increased by 5%, and a new

blood sample was analyzed within 5 minutes.

At the end of the experiment animals were exposed to a second 5-min episode of hypoxia (11.5%

O_2) followed by a brief "maximal" chemoreceptor challenge induced by switching off the mechanical

ventilator until the animal exhibited a "gasping-like" phrenic discharge pattern (approximately 20-30

seconds). If the increase in phrenic nerve amplitude in response to the "maximal" challenge was lower

than the response observed during either hypoxic episodes, it was considered a sign of deteriorating

nerve-electrode contact and the preparation was excluded from all formal analysis. Rats were then

perfused transcardially with heparinized saline followed by 4% paraformaldehyde and spinal cords were

removed harvested for histological analysis.

Histology. Spinal cords were harvested and placed in 4% paraformaldehyde for 24 hours. The cords were

subsequently moved to a cryo-protecting solution (30% sucrose in 1x PBS) for a minimum of three days.

Cervical and thoracic spinal cords were blocked in optimal cutting temperature media and cryosectioned

at 20 μm. The viral constructs included a red fluorescent protein (mCherry) fused to the hM3D(Gq)

DREADD which allowed evaluation of DREADD expression by assessing mCherry expression via

fluorescence microscopy.

We performed a qualitative assessment of mCherry expression in the mid-cervical spinal cord.

One intact section from the middle of each spinal segment (C3-C6) was chosen as a representative section

and underwent assessment. Sections were segmented into the following quadrants: left dorsal, right

dorsal, left ventral, and right ventral. The quadrant was scored as "positive" if mCherry positive neurons

or fibers were observed; otherwise, the sub-segment was marked "negative". Results were compiled into a

summary table showing the total positive counts by animal cohort, spinal segment, and quadrant (see

Results section; **Table 1**). Animals that showed no positive mCherry labeling in the C3-C6 cord and no

EMG response to J60 were excluded from analysis.

Data analysis. Custom MATLAB (MathWorks; Natick, MA) scripts were created, and are available upon

request. These scripts were used to analyze diaphragm EMG, phrenic nerve, and plethysmography

waveforms. EMG signals were digitally filtered using a second-order, bandpass Butterworth filter (100–

1000 Hz) and then rectified and integrated by taking the absolute value of the signal followed by applying

a moving median filter (50 ms time constant for mice; 75 ms time constant for rats) and moving average

filter (50 ms time constant for mice; 175 ms time constant for rats). The script identified each EMG burst

and calculated peak amplitude, minimum amplitude (tonic activity), and area under the curve (AUC) for

each burst which was then averaged across animals and compared across experimental conditions.

Differences in mortality between wild type and ChAT-Cre mice post-J60 application was assessed using

Pearson's Chi-squared test with Yates' continuity correction using the chisq.test function in R. In instances

where animals did not survive the entire duration of the anesthetized recording, data up until the time

point preceding their death was included.

Phrenic nerve signals were digitally filtered using a second-order, bandpass Butterworth filter

(100–3 kHz) and then rectified and integrated by taking the absolute value of the signal followed by

applying a moving median filter (50 ms time constant) and moving average filter (50 ms time constant).

The script calculated the peak phrenic burst amplitude and minimum amplitude for each burst which was

then averaged across animals and compared across experimental conditions. Systolic (SP), diastolic (DP),

and mean arterial blood pressure (MAP; formula: MAP = DP + 1/3 (SP - DP)) along with instantaneous

heart rate were calculated from the arterial pressure trace.

In plethysmography experiments, airflow pressure, chamber temperature, chamber humidity,

barometric pressure, and animal's body temperature were used to calculate respiratory frequency, tidal

volume, and ventilation via a custom MATLAB script. Tidal volume was calculated using the Drorbaugh

and Fenn equation[29].

Statistical analyses were performed using SigmaPlot 14 (Systat Software) and R (The R

Foundation for Statistical Computing; version 4.3.1). In mouse diaphragm EMG studies, one-way

repeated measure analysis of variance (ANOVA) was used to statistically compare diaphragm EMG peak

amplitude, area under the curve, tonic activity, and heart rate across time before and after J60 application.

Two-sample t-tests were used to compare left and right hemi-diaphragm EMG peak amplitude, area under

the curve, tonic activity, and heart rate between ChAT-Cre and wild type mice at the 30-minute post-J60

administration time point. In plethysmography experiments, two-way repeated measures ANOVA was

used to compare raw and normalized tidal volume, respiratory frequency, and minute ventilation across

time and treatment (saline vs J60). One-way RM ANOVA was used to compare phrenic peak amplitude,

systolic and diastolic blood pressures, mean arterial blood pressure, and respiratory rate across time for

phrenic nerve recordings. The relationship between time post-AAV injection and average phrenic

response to J60 was assessed for ChAT-Cre rats using the cor.test function in R to run a Pearson's

product moment correlation.

In cases of significant main effects, Tukey post-hoc test was used to assess differences between

individual time points. For instances where data failed to meet general linear model assumptions (i.e.,

normality, homogeneity of variances), nonparametric equivalents of the previously mentioned statistical

tests were used. Data were considered statistically significant when $p \leq 0.05$. The mean data are presented

along with standard error of the mean.

RESULTS

Diaphragm EMG responses in wild type mice. Wild type mice underwent bilateral injections of AAV9-
hSyn-HA-hM3D(Gq)-mCherry into the ventral horns at spinal segment C4. Following a six-to-eight-
weeks incubation period, mice underwent terminal diaphragm EMG recordings before and after
application of the selective DREADD ligand, J60. On average, wild type mice showed increases in
diaphragm EMG output in at least one hemi-diaphragm after J60 administration (**Figure 1**). Diaphragm
EMG AUC was significantly increased after DREADD activation (**Figure 1d**) in both the left (**p = 0.002**;
Table S1) and right (**p = 0.002**; **Table S1**) hemi-diaphragm. Additionally, peak-to-peak amplitude of the
rectified and integrated diaphragm EMG burst activity increased bilaterally following J60 administration
(Left hemi-diaphragm: p = 0.056; Right hemi-diaphragm: **p = 0.01**; **Figure 1e**; **Table S1**). Lastly, the
tonic activity of the diaphragm was assessed (**Figure 1f**). Similar to the previous measure of EMG output
both the left (**p = 0.052**; **Table S1**) and right (**p < 0.001**; **Table S1**) hemi-diaphragm exhibited a
significant increase in tonic activity following J60 administration. Respiratory rate was consistent for the
duration of the experiment. Notably, there was no substantial change in respiratory rate of these
spontaneous breathing mice after J60 administration (p = 0.863; **Figure 1g**; **Table S1**).

   In all experiments, the selective DREADD ligand, J60, produced an increase in diaphragm EMG
burst amplitude during inspiration. However, this increase was not always detected in both the left and
right hemi-diaphragm EMG recording. Five mice showed a bilateral increase in diaphragm output after
J60 administration (**Figure 1**). Four mice showed a response that was limited to the right hemi-diaphragm
while two showed a response that was limited to the left hemi-diaphragm (**Figure 1**).

Diaphragm EMG responses in ChAT-Cre mice. ChAT-Cre mice received bilateral intraspinal injections
of AAV9-hSyn-DIO-hM3D(Gq)-mCherry into the ventral horns at C4. ChAT-Cre mice underwent
terminal diaphragm EMG recording at the same time point and with the same protocol as wild type mice.
All animals (n= 9/9) showed an increase in diaphragm EMG output in response to the J60 DREADD
ligand in at least one hemi-diaphragm (**Figure 2**). On average, both left (**p = 0.011**; **Table S2**) and right

(**p < 0.001**; **Table S2**) diaphragm EMG AUC increased over time after J60 administration (**Figure 2d**).

Diaphragm EMG peak-to-peak amplitude had a similar, bilateral increase following J60 delivery (Left

hemi-diaphragm: **p = 0.013**; Right hemi-diaphragm: **p < 0.001**; **Table S2**; **Figure 2e**). Lastly, tonic

activity also showed an increase over time after J60 administration (**Figure 2f**) for both the left (**p= 0.002**;

Table S2) and right (**p < 0.001**; **Table S2**) hemi-diaphragm. Respiratory rate decreased significantly over

time after J60 administration (**p < 0.001**; **Figure 2g**; **Table S2**). Apart from two animals that showed a

unilateral EMG response that was limited to the right hemi-diaphragm the remaining ChAT-Cre mice (n=

7/9) had bilateral increases in diaphragm EMG output following J60 administration (**Figure 2**).

Wild type vs. ChAT-Cre comparison. Diaphragm EMG responses of wild type and ChAT-Cre mice were

compared at the 30-minute post-J60 time point (**Figure 3**). Left hemi-diaphragm responses to J60 were

similar between the two groups across all outcome measures (AUC: p = 0.998; Peak-to-peak amplitude: p

= 0.771; Tonic activity: p = 0.160; **Table S3**; **Figure 3a-c**). However, right hemi-diaphragm responses to

J60 differed across AUC (**Figure 3d**), peak-to-peak amplitude (**Figure 3e**), and tonic activity (**Figure 3d**)

with ChAT-Cre animals on average having larger magnitude responses compared to wild type animals

(AUC: **p = 0.0417**; Peak-to-peak amplitude: **p = 0.00403**; Tonic activity: **p = 0.00207**; **Table S3**).

Respiratory rate was not different between the two groups (p = 0.382; **Table S3**; **Figure 3g**).

Five of eleven wild type mice died before the end of the experimental recording period. It is

unclear if this was due to the nature of the preparation or due to an unforeseen consequence of DREADD

activation. Three animals died between the 30-minute and 60-minute time points, and two animals died

before the final 90-minute time point. In contrast, all mice in the ChAT-Cre cohort survived the total

duration of the experimental protocol. A chi-squared assessment was run to assess the difference in

survival between the two cohorts. The test did not return a p-value that met the threshold for statistical

significance however, considering the small sample size the resultant p-value does indicate that there was

likely some association between mouse strain (i.e., wild type, ChAT-Cre) and death after DREADD

activation (Chi-squared = 3.2997, df = 1, p = 0.06929).

ChAT-Cre rats – Plethysmography and Phrenic Nerve Recordings. A small cohort of ChAT-Cre rats

underwent diaphragm EMG recordings to ensure DREADD responses similar to mouse cohort could be

obtained in rats. ChAT-Cre rats (n= 4) underwent bilateral, intraspinal injections of AAV9-hSyn-DIO-

hM3D(Gq)-mCherry into the ventral horns at C4 to introduce the hM3D(Gq) DREADD transgene into

phrenic motoneurons. Four of four rats showed increased diaphragm EMG output after DREADD

activation (**Figure S1**). With that knowledge, we used a separate group of ChAT-Cre rats (n= 9; n= 3

females) to assess the effects of DREADD activation on ventilation. Whole-body plethysmography was

used to measure breathing frequency, tidal volume, and minute ventilation before and after intravenous

delivery of saline (sham) and J60 (**Figure 4**). Delivery of the J60 ligand resulted in an increase in

inspiratory tidal volume compared to sham infusion (<u>Normalized to Weight (ml/kg):</u> Main effect of

Treatment: **p= 0.037**; **Figure 4a**; <u>Normalized to Baseline:</u> Main effect of Treatment: p= 0.091; **Figure 4d**;

Table S4). Respiratory rate appeared to be unaffected by DREADD activation and was similar between

sham and J60 conditions (<u>Respiratory Rate:</u> Main effect of Treatment: p= 0.582; **Figure 4b**; <u>Respiratory</u>

<u>Rate Normalized to Baseline:</u> Main effect of Treatment: p= 0.774; **Figure 4e**; **Table S4**). Minute

ventilation was slightly elevated in the J60 condition vs sham; however, this increase did not reach the

threshold for statistical significance (<u>Normalized to Body Weight</u>: Main effect of Treatment: p= 0.194;

Figure 4c; <u>Normalized to Baseline:</u> Main effect of Treatment: p= 0.337; **Figure 4f; Table S4**).

Phrenic nerve recordings were made to directly assess the effects of DREADD activation

on phrenic output. There was no detectable relationship between time post-AAV injection and phrenic

response to DREADD activation (Pearson correlation; <u>Left peak-to-peak response</u>: p= 0.215; <u>Right peak-</u>

<u>to-peak response</u>: p= 0.318).

Application of the J60 ligand caused a rapid, sustained, and bilateral increase in phrenic nerve

efferent burst amplitude (<u>Left phrenic peak-to-peak amplitude (normalized to baseline)</u>: **p< 0.001**; <u>Right</u>

<u>phrenic peak-to-peak amplitude (normalized to baseline)</u>: **p< 0.001**; **Table S5**; **Figure 5b-c**), whereas

saline injection had no impact. The increase in phrenic burst amplitude lasted up to 100 minutes post J60

324   administration, at which point the experiment was terminated. Application of the J60 ligand also resulted

325   in an increase in phrenic tonic activity (<u>Left phrenic tonic activity (normalized to baseline)</u>: **p< 0.001**;

326   <u>Right phrenic tonic activity (normalized to baseline)</u>: **p< 0.001**; **Table S5**; **Figure 5d-e**).

327   Heart rate, systolic and diastolic blood pressure, mean arterial blood pressure (MAP), as

well as respiratory rate were also assessed (**Figure 5f-j**). Application of J60 had no effect on heart rate

(p= 0.587; **Table S5**; **Figure 5h**) or respiratory rate (p= 0.282; **Table S5; Figure 5j**) but did result in an

decrease in both systolic (**p< 0.001**; **Table S5**; **Figure 5f**) and diastolic blood pressure (**p< 0.001**; **Table**

S5; **Figure 5g**) as well as MAP (**p< 0.001**; **Table S5**; **Figure 5i**).

Histological analysis. We performed qualitative analysis of the mid-cervical spinal cord from each animal

to assess the extent of mCherry fluorophore expression (**Figure S2**). All mice from both cohorts showed

evidence of mCherry expression in at least one segment of the mid-cervical spinal cord (**Figure 6**) except

for n= 1 ChAT-Cre mouse. This mouse was excluded from analysis based off *a priori* exclusion criteria

which stipulated that to be included in the final analysis an animal must show evidence of mCherry

expression in the grey matter of at least one spinal segment from C3-C6. A summary of the results is

given in **Table 1**.

Patterns of expression were relatively homogenous in wild type animals. In this cohort, the number of

animals with positive mCherry expression in the grey matter increased on the rostral caudal axis. Positive

mCherry counts were comparable on both the dorsal-ventral and left-right axes, with a majority of

animals expression mCherry in all four quadrants. The diaphragm EMG responses to J60, on average,

exhibited similarity between the left and right hemi-diaphragm in these animals, aligning with the

observed pattern of mCherry expression. (**Figure 1d-f**).

In contrast, mCherry expression in the ChAT-Cre mice cohort was more prevalent in the ventral

horns and the right side of the cord. Like the wild type animals there was a slight trend for increased

mCherry expression moving rostral to caudal. These animals demonstrated a larger average DREADD

response in the right hemi-diaphragm than the left (**Figure 2d-f**), possibly stemming from the fact that a

greater number of animals exhibited mCherry expression on the right side compared to the left (**Figure**

S3).

ChAT-Cre rats showed expression predominately in the ventral horns throughout the mid-cervical

spinal cords with the highest levels of expression in spinal segments C4 and C5. One animal from the

diaphragm EMG cohort showed no mCherry expression in the mid-cervical spinal cord but still had a

robust increase in diaphragm output after J60 administration. All other ChAT-Cre rats showed robust

mCherry expression in the ventral horns of at least one spinal segment from C3-C6. Expression in this

cohort was slightly more prominent on the right side of the cord, and in the ventral horn. These

histological findings were consistent with the physiology results. Although the magnitude of DREADD

response between the left and right phrenic nerves for this cohort was not statistically different, there was

a trend of slightly higher right phrenic tonic response compared to the left (**Figure 5b-e**). This trend is

mirrored in the pattern of mCherry expression, where expression levels were approximately equal

between spinal segments but tended to be slightly higher in the right ventral horns compared to the left.

DISCUSSION

We describe a novel method to increase diaphragm EMG output by expressing the excitatory DREADD, hM3D(Gq), in the mid-cervical spinal cord, targeting phrenic motoneurons. Following AAV-driven expression of the DREADD in the spinal cord, application of the J60 ligand caused sustained increases in diaphragm output as measured through EMG in spontaneously breathing animals. This response was also verified using direct recordings of phrenic nerve discharge. Additionally, the DREADD ligand was able to produce an increase in inspiratory tidal volume in awake, freely behaving animals. These proof-of-concept studies provide a foundation for further development of this technology towards clinical application for restoring diaphragm activation in conditions such as cervical spinal cord injury.

Targeted gene delivery to the phrenic motor pool. The intraspinal AAV delivery used here was based on previous studies demonstrating successful gene delivery to phrenic motoneurons[20][24,30,31]. For example, mid-cervical spinal injections of an AAV5 vector encoding the lysosomal enzyme acid alpha-glucosidase (GAA) in animals with Pompe disease (*Gaa* null) effectively restores spinal GAA enzyme activity [24]. Spinal-delivered viral vectors have also been used to successfully drive local expression of channelrhodopsin-2 to enable light activation of diaphragm output[20], and to drive expression of the astrocyte glutamate transporter GLT1 in the area of the phrenic motor nuclei[30]. Other methods that have been employed to drive gene expression in phrenic motoneurons include intrapleural- and intramuscular diaphragm injection of viral vectors[32]. Intrapleural delivery requires microinjection to the "pleural space" between the visceral pleura that lines the lungs and the parietal pleura which covers the thoracic cavity. This technique[33] effectively targets phrenic motoneurons in rodent models of cervical spinal cord injury[34-36] and Pompe disease[37]. Intramuscular diaphragm injection allows the vector to enter phrenic nerve terminals and reach phrenic motoneuron soma via retrograde movement[32]. Direct diaphragm injection allows for a relatively high target specificity, with the gene of interest expressed almost exclusively in phrenic motoneurons (although expression can also occur in diaphragm myofibers, depending on the promoter sequence used). In pilot experiments, we tested intrapleural and also intramuscular diaphragm

injections using AAV9 vectors encoding GFP (AAV9-CAG-GFP) or DREADD (AAV9-hSyn-HA-

hM3D(Gq)-mCherry & AAV9-hSyn-DIO-hM3D(Gq)-mCherry). We did not, however, observe

histological or physiological evidence of phrenic motoneuron transduction with these AAV9 vectors. We

therefore utilized direct intraspinal injection[24,30] to introduce the hM3D(Gq) into the phrenic motor

nucleus. While this enabled proof-of-concept for targeting DREADDs to the cervical spinal cord and

phrenic motoneurons, the aforementioned methods might ultimately prove better for selective phrenic

motoneuron targeting. We predict that using different AAV serotypes or viruses with better retrograde

movement (e.g., "AAV retro") could optimize targeting of phrenic motoneurons[38].

DREADD-mediated motoneuron activation. DREADD technology is widely used for studying brain and

spinal cord neurons and networks[39,40]. Relatively few studies, however, have studied if and how

DREADDs can be used to activate (or inhibit) lower motoneurons. In regard to the spinal cord, we are

aware of a few prior publications[41-44]. Two of these studies used pharmacologically selective actuator

module (PSAM), a type of ionotropic chemogenetic receptor, to activate thoracic[41] and lumbar[42]

motoneurons, respectively, in mouse models of amyotrophic lateral sclerosis (ALS). In the remaining

studies, excitatory DREADDs were applied to spinal motoneurons as way to improve axon regeneration

follow peripheral nerve injury[43,44]. A small but growing body of work has employed DREADDs to

activate hypoglossal (XII) motoneurons in the brainstem[8]. Collectively, these studies show that once

hM3D(Gq) is expressed in XII motoneurons, DREADD ligands will rapidly produce an increase in the

EMG activation of tongue muscles[14,15]. This increase in tongue muscle output tends to manifest as an

increase in the inspiratory related activation, and tonic discharge across the respiratory cycle. Since

increased tongue muscle activation can promote patency of the upper airway, XII motoneuron DREADD

expression has been suggested as a possible treatment for obstructive sleep apnea[12,14,16]. For the present

study, the primary innovation is the first application of DREADD technology to phrenic motoneurons.

This approach was highly effective at driving sustained activation of the diaphragm muscle, and the

underlying mechanisms are discussed next.

Chemogenetic stimulation of breathing. An important consideration is how DREADD-induced increases

in the excitability of spinal neurons, including phrenic motoneurons, interacts with the endogenous neural

control of breathing. Phrenic motoneurons receive a rhythmic, monosynaptic, glutamatergic synaptic

input from medullary neurons. Acting via NMDA and AMPA receptors, this produces phrenic

motoneuron depolarization and subsequent diaphragm muscle contraction[17]. Activating DREADD

receptors on phrenic motoneurons should lower the threshold for activation via excitatory glutamatergic

synaptic inputs, which would produce a greater output during the inspiratory phase. Alternatively,

DREADD activation could directly lead to phrenic motoneuron action potentials even in the absence of

synaptic input from the brainstem. This latter possibility could explain the tonic discharge (i.e., EMG

output across the entire respiratory cycle) that was noted to occur after delivery of the DREAD ligand.

Non-specific spinal cord DREADD expression, as occurred in the wild type mice (e.g., **Figure 6**), would

likely produce an increase in the excitability and/or activation of phrenic motoneurons as well as

propriospinal neurons in the immediate vicinity. Neurophysiological[45] as well as anatomical data[23,46]

confirm synaptic connections between mid-cervical interneurons and phrenic motoneurons, making it

possible that DREADD activation of these interneurons impacted the diaphragm motor response in the

wild type mice.

The control of breathing is also impacted by well established "closed loop" physiologic feedback

mechanisms regulating lung volume and arterial blood gases[47,48]. For example, if DREADD-induced

activation of the diaphragm leads to increased alveolar ventilation, and metabolic rate is not impacted,

then arterial CO_2 values will decrease and the overall neural drive to breathe will also decrease. Vagal

afferent feedback corresponding to increased lung volume also has a powerful inhibitory impact on

inspiration and therefore diaphragm activation. However, the sustained increase in diaphragm EMG and

tidal volume that we observed following application of the DREADD ligand indicates that these

mechanisms, if activated, were not sufficient to fully inhibit the increased phrenic motoneuron output. In

this regard, our additional experiments in which direct recordings of bilateral phrenic nerve discharge are

informative. These nerve recording experiments were done to enable direct evaluation of the impact of

spinal DREADD activation on phrenic motor output, but while keeping arterial blood gases and lung

volume constant. Under these more rigorously controlled conditions, intravenous delivery of the

DREADD ligand produced a rapid and sustained increase in inspiratory burst amplitude in the phrenic

nerve, and with no impact on the rate of the inspiratory bursts. The relative increase in inspiratory motor

output was considerably greater in the phrenic nerve recording experiments (~250% of baseline) as

compared to the diaphragm EMG response in spontaneously breathing animals (~100% of baseline). This

may indicate that vagal and/or blood gas related inhibitory mechanisms, as mentioned above, somewhat

constrained the response to the DREADD ligand in the spontaneously breathing animal.

Critique of methods. There are a few caveats that should be discussed. First, the precision of the AAV

delivery could be improved by further refining spinal injection surgical techniques. In the current study

we used a stereotaxic frame and previously validated coordinates[24,25] to guide the intraspinal AAV

injections. However, we observed variability in the (left vs. right) laterality of mid-cervical mCherry

expression as well as the physiological response to the DREADD ligand, particularly in the ChAT-Cre

mice (e.g., **Figure 2**; **Table 1**). This could have occurred due to subtle variations of the positioning of the

animal within the stereotaxic frame, and/or placement of the needle tip, leading to slight deviations for the

desired coordinates between the left and right phrenic nuclei. Second, we did not unequivocally verify

that the DREADD was expressed in phrenic motoneurons using retrograde labeling methods[33,49].

However, the phrenic motor nucleus has been well described in the mouse[50] and the rat[51,52], and the

fluorophore (mCherry) expression observed in our experiments is very clearly in the expected location of

phrenic motoneurons (**Figure 6b-c**; **Supplemental Figure 2**). Further, the robust increase in phrenic

motor output after the DREADD ligand, particularly in the ChAT-Cre rat experiment (**Figure 5**) is further

evidence of effective phrenic motoneuron targeting.

Conclusion. Collectively, the data support the conclusion that cervical spinal cord directed chemogenetic

methods can be used to produce a sustained increase in phrenic motor output, diaphragm activation, and

inspiratory tidal volume. However, more direct targeting of DREADD expression to phrenic motoneurons

may likely to be needed for optimal diaphragm activation using this method. In this regard, improvement

of the AAV delivery methods will increase the selectively of the approach for more precise targeting of

phrenic motoneurons. In regard to the "translational value" of this work, spinal cord chemogenetics may

have application to clinical conditions associated with an inability to activate the diaphragm. For example,

incomplete cervical spinal cord injury is a condition in which the bulbospinal synaptic inputs to phrenic

motoneurons are interrupted. After cervical spinal cord injury, expressing an excitatory DREADD in

phrenic motoneuron could be used to increase the excitability of these cells, and thereby improve the

efficacy of the limited bulbospinal synaptic inputs.

Figure 1. *DREADD activation increases diaphragm EMG output in wild type mice.* A representative example of diaphragm EMG activity before and after application of the J60 DREADD ligand is shown in the top panel. Examples of the individual inspiratory EMG bursts at baseline (a), after vehicle (b), and after J60 (c) are shown. The J60 ligand increased diaphragm output but did not impact respiratory rate. The mean responses (n= 11; n= 7 females) for EMG AUC, peak-to-peak amplitude, tonic activity, and respiratory rate are shown in panels d-g. For diaphragm EMG data (panels d-f) left hemi-diaphragm EMG is represented in orange, while right hemi-diaphragm EMG is blue. Error bars depict ± 1 SEM. Statistical reports for all panels are provided in Supplemental Table 1. Dia = diaphragm, AUC = area under the curve, amp = peak amplitude, BL = baseline, SL = saline (sham injection).

Figure 2. *DREADD activation increases diaphragm EMG output in ChAT-Cre mice.* A representative example of diaphragm EMG activity before and after application of the J60 DREADD ligand is shown in the top panel. Examples of the individual inspiratory EMG bursts at baseline (a), after vehicle (b), and after J60 (c) are shown. Mean responses (n= 9; n= 6 females) for EMG AUC, peak-to-peak amplitude, tonic activity, and respiratory rate are shown in panels d-g. The DREADD ligand caused a bilateral increase in diaphragm EMG AUC, peak-to-peak amplitude, and tonic activity. For all EMG parameters, the responses were greater on the right vs. left hemi-diaphragm. Respiratory rate decreased over time. For panels d-f, the left hemi-diaphragm EMG is represented in orange, while right hemi-diaphragm EMG is blue. Error bars depict ± 1 SEM. Statistical reports for all panels are provided in Supplemental Table 2. Dia = diaphragm, AUC = area under the curve, amp = peak amplitude, BL = baseline, SL = saline (sham injection).

Figure 3. *Wild type vs. ChAT-Cre mouse responses to DREADD activation.* Direct comparisons of diaphragm EMG response parameters (a-f) and respiratory rate (g) at 30-minute post-J60 application (Wild type, n= 11; n= 7 females; ChAT-Cre, n= 9; n= 6 females). Left hemi-diaphragm EMG AUC (a), peak-to-peak amplitude (b) and tonic activity (c) were similar between groups. However, the same parameters on the right hemi-diaphragm (d-f) were greater in ChAT-Cre mice. Respiratory rate was similar between groups. Error bars depict ± 1 SEM. Statistical reports for all panels are provided in Supplemental Table 3. * p < 0.05. AUC = area under the curve, amp = peak EMG amplitude, Dia = diaphragm, BL = baseline, resp rate = respiratory rate.

Figure 4. *DREADD activation increases ventilation in unanesthetized ChAT-Cre rats.* Summary plots (n= 9; n= 3 females) showing the impact of the J60 DREADD ligand on tidal volume, respiratory rate, and minute ventilation are shown in panels a-c. The normalized values (% of baseline) are shown in panels d-f. The DREADD ligand increased tidal volume compared to sham infusion (saline). Error bars depict ± 1 SEM. Statistical reports for all panels are provided in Supplemental Table 4. BL = baseline, IV = intravenous infusion period.

Figure 5. *DREADD activation increases phrenic nerve output in ChAT-Cre rats.* Representative data showing that the J60 DREADD ligand causes a rapid increase in phrenic nerve output (a). Mean data (n= 9; n= 3 females) showing the impact of J60 application on phrenic nerve raw (b) and normalized (c) peak-to-peak amplitude, raw (d) and normalized (e) tonic activity, systolic blood pressure (f), diastolic blood pressure (g), heart rate (h), mean arterial blood pressure (i), and respiratory rate (j). The J60 ligand caused an increase in phrenic peak-to-peak amplitude, and tonic activity. Systolic, diastolic, and mean arterial blood pressure all decreased after J60 application. Heart rate and respiratory rate were not statistically different after J60. In panels b-e, the left phrenic is represented in orange, while right phrenic is blue. Error bars depict ± 1 SEM. Statistical reports for all panels are provided in Supplemental Table 6. Phr = phrenic, amp = amplitude, BL = baseline, SP = systolic pressure, DP = diastolic pressure, HR = heart rate, MAP = mean arterial pressure.

Figure 6. *Histological assessment of mCherry expression in the C4/C5 spinal segments.* Representative photomicrographs of mid-cervical spinal sections from a wild type mouse (a-a$_i$), a ChAT-Cre mouse (b-b$_i$) and a ChAT-Cre rat (c-c$_i$). Wild type mice (a-a$_i$) showed nonspecific pattern of expression throughout the mid-cervical grey matter. ChAT-Cre mice and rats (b-c$_i$) showed expression limited to neurons in the ventral horns. Red color indicates positive and mCherry fluorescence. Dashed white line indicates approximate white-gray matter demarcation.

	Wildtype mice (n= 11)				ChAT-Cre mice (n= 9)				ChAT-Cre rats (n= 9)			
	Dorsal		Ventral		Dorsal		Ventral		Dorsal		Ventral	
	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right
C3	4	5	6	6	0	1	3	4	0	1	3	4
C4	7	9	10	9	1	2	5	9	0	1	6	8
C5	11	11	11	11	0	1	6	9	1	1	5	9
C6	10	10	9	9	2	2	5	8	0	0	1	2

Table 1. *Qualitative assessment of mCherry expression in the mid-cervical spinal cord.* Spinal segments C3-C6 were assessed in quadrants broken into dorsal, ventral, left, and right. Spinal segments were counted as "positive" if they showed any evidence of mCherry expression in neuronal soma or fibers. The counts therefore indicate the number of animals of a given cohort that were mCherry positive for a given spinal segment quadrant. All animals showed slight trend for more mCherry expression moving rostral to caudal and for more expression in the ventral vs the dorsal lamina. This trend was more prominent in the ChAT-Cre animals. At the bottom of the table, a heatmap is provided for easier assessment of the distribution of positive mCherry counts across quadrants and spinal segments.

Supplemental Figure 1. *Impact of DREADD activation on diaphragm EMG in ChAT-Cre rats.*
Example trace of rectified and integrated diaphragm EMG activity from a ChAT-Cre rat that had previously undergone injections of AAV9-hSyn-DIO-hM3D(Gq)-mCherry into the C4 ventral horns before and after application of the DREADD ligand, J60. Callout panels below show example activity at baseline (a), after injection of vehicle (b), and after application of the DREADD ligand, J60 (c).

Supplemental Figure 2. *Example quantification of mCherry expression in the C3-C6 spinal cord.* Mid-cervical spinal cord images from ChAT-Cre animal (left) with positive mCherry expression. A grid has been placed on each image to highlight the quadrant system (i.e., left dorsal, right dorsal, left ventral, right ventral) used to score the tissue. The right column shows an outline of each section on the left with a score for each quadrant based on mCherry expression. A given quadrant was scored with a "+" if it contained any mCherry positive neurons or fibers else it was given a "-" signifying no mCherry expression.

a - Unilateral (right) mCherry expression in the C4 spinal cord corresponds to unilateral (right) diaphragm response to the J60 ligand

b - Bilateral mCherry expression in the C4 spinal cord corresponds to a bilateral diaphragm response to the J60 ligand

Supplemental Figure 3. *mCherry expression in the mid-cervical ventral horn corresponds with the laterality of the diaphragm EMG DREADD response.* Example spinal histology (left) from two ChAT-Cre mice showing unilateral (a) and bilateral (b) expression of mCherry (red) in the ventral horn(s) and corresponding diaphragm EMG traces. Unilateral ventral horn mCherry expression (a) results in DREADD responses that were limited to the ipsilateral hemi-diaphragm. Animals with bilateral ventral horn DREADD expression responded to DREADD activation with a bilateral increase in diaphragm output.

Main Effects

Outcome	Hemi-diaphragm	Test	df	Test statistic	p value
Diaphragm EMG AUC	left	One-way RM ANOVA	6,68	F = 4.008	**0.002**
	right	One-way RM ANOVA	6,68	F = 4.054	**0.002**
Diaphragm EMG peak-to-peak amplitude	left	One-way RM ANOVA	6,68	F = 2.213	0.056
	right	One-way RM ANOVA	6,68	F = 3.191	**0.01**
Diaphragm EMG tonic activity	left	One-way RM ANOVA	6,68	F = 2.256	0.052
	right	One-way RM ANOVA	6,68	F = 5.225	**< 0.001**
Respiratory Rate	NA	One-way RM ANOVA	6,68	F = 0.419	0.863

Post-Hoc Tests (Tukey Test)

Outcome	Hemi-diaphragm	Comparison	Diff of Ranks	q	P
Diaphragm EMG AUC	left	15 min vs. Saline	60.488	4.972	**0.015**
		60 min vs. Saline	68.183	5.064	**0.013**
		90 min vs. Saline	66.079	4.446	**0.041**
Diaphragm EMG AUC	right	15 min vs. Saline	41.481	4.41	**0.044**
		60 min vs. Saline	54.104	5.197	**0.01**
		90 min vs. Saline	51.214	4.457	**0.04**
Diaphragm EMG peak-to-peak amplitude	right	90 min vs. Saline	40.925	4.947	**0.016**
Diaphragm EMG tonic activity	right	15 min vs. Baseline	124.737	5.044	**0.013**
		15 min vs. Saline	135.783	5.49	**0.005**
		30 min vs. Baseline	110.583	4.471	**0.039**
		30 min vs. Saline	121.628	4.918	**0.017**
		90 min vs. Baseline	140.208	4.641	**0.029**
		90 min vs. Saline	151.254	5.006	**0.014**

Supplemental Table 1. *Statistical summary for impact of DREADD activation on diaphragm EMG in wild type mice.* Time points are in reference to minutes passed since J60 infusion. Summary data presented in Figure 1. EMG = electromyography, AUC = area under the curve, RM = repeated measures, df = degrees of freedom. Bolded p-values indicates p<0.05.

Main Effects

Outcome	Hemi-diaphragm	Test	df	Test statistic	p value
Diaphragm EMG AUC	left	Friedman RM analysis on ranks	6	Chi-square = 16.619	**0.011**
	right	Friedman RM analysis on ranks	6	Chi-square = 40.143	**< 0.001**
Diaphragm EMG peak-to-peak amplitude	left	One-way RM ANOVA	6,62	F = 3.05	**0.013**
	right	Friedman RM analysis on ranks	6	Chi-square = 36.667	**< 0.001**
Diaphragm EMG tonic activity	left	Friedman RM analysis on ranks	6	Chi-square = 20.952	**0.002**
	right	One-way RM ANOVA	6,62	F = 10.076	**< 0.001**
Respiratory Rate	NA	One-way RM ANOVA	6,62	F = 25.13	**< 0.001**

Post-Hoc Tests (Tukey Test)

Outcome	Hemi-diaphragm	Comparison	Diff of Ranks	q	P
Diaphragm EMG AUC	left	30 min vs. Saline	32	4.938	**0.009**
		15 min vs. Baseline	31	4.783	**0.013**
		15 min vs. Saline	38	5.864	**<0.001**
		30 min vs. Baseline	35	5.401	**0.003**
Diaphragm EMG AUC	right	30 min vs. Saline	42	6.481	**<0.001**
		60 min vs. Baseline	32	4.938	**0.009**
		60 min vs. Saline	39	6.018	**<0.001**
		90 min vs. Baseline	27	4.166	**0.05**
		90 min vs. Saline	34	5.246	**0.004**
Diaphragm EMG peak-to-peak amplitude	left	30 min vs. Saline	32.824	4.447	**0.042**
		15 min vs. Baseline	31	4.783	**0.013**
		15 min vs. Saline	36	5.555	**0.002**
		30 min vs. Baseline	34	5.246	**0.004**
Diaphragm EMG peak-to-peak amplitude	right	30 min vs. Saline	39	6.018	**<0.001**
		60 min vs. Baseline	32	4.938	**0.009**
		60 min vs. Saline	37	5.709	**0.001**
		90 min vs. Saline	31	4.783	**0.013**
Diaphragm EMG tonic activity	left	5 min vs. Baseline	28	4.32	**0.037**
		15 min vs. Baseline	29	4.475	**0.026**
		90 min vs. Baseline	35	5.401	**0.003**
		5 min vs. Baseline	295.193	5.582	**0.005**
		5 min vs. Saline	297.174	5.619	**0.004**
		15 min vs. Baseline	375.713	7.104	**<0.001**
		15 min vs. Saline	377.694	7.142	**<0.001**
Diaphragm EMG tonic activity	right	30 min vs. Baseline	347.212	6.565	**<0.001**
		30 min vs. Saline	349.193	6.603	**<0.001**
		60 min vs. Baseline	317.232	5.998	**0.002**
		60 min vs. Saline	319.213	6.036	**0.002**
		90 min vs. Baseline	358.443	6.778	**<0.001**
		90 min vs. Saline	360.424	6.815	**<0.001**
		5 min vs. 30 min	5.296	4.733	**0.025**
		5 min vs. 60 min	7.296	6.52	**<0.001**
		5 min vs. 90 min	11.889	10.624	**<0.001**
		15 min vs. Baseline	5.778	5.163	**0.011**
		15 min vs. Saline	5.963	5.328	**0.008**
		15 min vs. 90 min	9.222	8.241	**<0.001**
Respiratory Rate	NA	30 min vs. Baseline	8.407	7.513	**<0.001**
		30 min vs. Saline	8.593	7.678	**<0.001**
		30 min vs. 90 min	6.593	5.891	**0.002**
		60 min vs. Baseline	10.407	9.3	**<0.001**
		60 min vs. Saline	10.593	9.465	**<0.001**
		90 min vs. Baseline	15	13.404	**<0.001**
		90 min vs. Saline	15.185	13.569	**<0.001**

Supplemental Table 2. *Statistical summary for impact of DREADD activation on diaphragm EMG in ChAT-Cre mice.* Time points are in reference to minutes passed since J60 infusion. Summary data presented in Figure 2. EMG = electromyography, AUC = area under the curve, RM = repeated measures, df = degrees of freedom. Bolded p-values indicates p<0.05.

Outcome	Hemi-diaphragm	Test	df	Test statistic	p value
Diaphragm EMG AUC	left	Student's t-test	18	t = 0.00310	p = 0.998
	right	Student's t-test	18	t = -2.913	**p = 0.0417**
Diaphragm EMG peak-to-peak amplitude	left	Student's t-test	18	t = -0.295	p = 0.771
	right	Student's t-test	18	t = -3.294	**p = 0.00403**
Diaphragm EMG tonic activity	left	Mann-Whitney Rank Sum test	18	Mann-Whitney U Statistic= 31.000	p = 0.160
	right	Student's t-test	18	t = -3.596	**p = 0.00207**
Respiratory Rate	NA	Mann-Whitney Rank Sum test	18	Mann-Whitney U Statistic= 38.000	p = 0.382

Supplemental Table 3. *Statistical summary for impact of DREADD activation on diaphragm EMG in in wild type mice vs. ChAT-Cre mice at the 30-min post J60 infusion time point.* Summary data presented in Figure 3. EMG = electromyography, AUC = area under the curve, df = degrees of freedom. Bolded p-values indicates p<0.05.

Outcome	Test	Normalization	Main effects	df	Test statistic	p value
Tidal Volume	Two-way RM ANOVA	Normalized to body weight (ml/kg)	Treatment	1,89	F = 6.286	**p= 0.037**
			Time	4,89	F = 9.817	**p < 0.001**
			Interaction	4,89	F = 0.969	p= 0.438
		Normalized to baseline	Treatment	1,89	F = 3.7	p= 0.091
			Time	4,89	F = 9.448	**p< 0.001**
			Interaction	4,89	F = 1.049	p= 0.398
Respiratory Rate	Two-way RM ANOVA	Raw values	Treatment	1,89	F = 0.329	p= 0.582
			Time	4,89	F = 14.326	**p < 0.001**
			Interaction	4,89	F = 1.640	p= 0.188
		Normalized to baseline	Treatment	1,89	F = 0.0885	p= 0.774
			Time	4,89	F = 11.862	**p< 0.001**
			Interaction	4,89	F = 1.310	p= 0.287
Minute Ventilation	Two-way RM ANOVA	Normalized to body weight (ml/kg)	Treatment	1,89	F = 2.011	p= 0.194
			Time	4,89	F = 12.875	**p < 0.001**
			Interaction	4,89	F = 1.215	p= 0.324
		Normalized to baseline	Treatment	1,89	F = 1.043	p= 0.337
			Time	4,89	F = 10.776	**p< 0.001**
			Interaction	4,89	F = 0.490	p= 0.490

Supplemental Table 4. *Statistical summary for impact of DREADD activation on plethysmography outcomes in unanesthetized ChAT-Cre rats using two-way repeated measures ANOVAs.* Each outcome measure is presented normalized to body weight (with the exception of respiratory rate) and normalized to values at baseline. Summary data presented in Figure 4. RM = repeated measures, ml/kg = milliliters of air per kilogram of animal's body weight, df = degrees of freedom. Bolded p-values indicates p<0.05.

Main Effects

Outcome	Nerve	Test	df	Test statistic	p value
Phrenic peak-to-peak amplitude (normalized to baseline)	left	One-way RM ANOVA	6,62	F = 13.355	**p< 0.001**
	right	Friedman Repeated Measures Analysis of Variance on Ranks	6	Chi-square = 35.619	**p< 0.001**
Phrenic tonic activity (normalized to baseline)	left	Friedman Repeated Measures Analysis of Variance on Ranks	6	Chi-square = 36.714	**p< 0.001**
	right	One-way RM ANOVA	6,62	F = 14.544	**p< 0.001**
Heart rate	NA	Friedman Repeated Measures Analysis of Variance on Ranks	6	Chi-square = 4.667	p= 0.587
Systolic blood pressure	NA	One-way RM ANOVA	6,62	F = 5.403	**p< 0.001**
Diastolic blood pressure	NA	One-way RM ANOVA	6,62	F = 12.460	**p< 0.001**
Mean arterial blood pressure	NA	One-way RM ANOVA	6,62	F = 9.422	**p< 0.001**
Respiratory Rate	NA	One-way RM ANOVA	6,62	F = 1.285	p= 0.282

Post-Hoc Tests (Tukey Test)

Outcome	Nerve	Comparison	Diff of Ranks	q	P
Phrenic peak-to-peak amplitude (normalized to baseline)	left	5 min vs. Baseline	111.026	6.52	**<0.001**
		5 min vs. Saline	86.692	5.091	**0.013**
		15 min vs. Baseline	133.337	7.83	**<0.001**
		15 min vs. Saline	109.003	6.401	**<0.001**
		30 min vs. Baseline	152.488	8.955	**<0.001**
		30 min vs. Saline	128.153	7.526	**<0.001**
		60 min vs. Baseline	140.658	8.26	**<0.001**
		60 min vs. Saline	116.324	6.831	**<0.001**
		90 min vs. Baseline	143.291	8.415	**<0.001**
		90 min vs. Saline	118.956	6.986	**<0.001**
	right	15 min vs. Baseline	32	4.938	**0.009**
		15 min vs. Saline	29	4.475	**0.026**
		30 min vs. Baseline	37	5.709	**0.001**
		30 min vs. Saline	34	5.246	**0.004**
		60 min vs. Baseline	35	5.401	**0.003**
		60 min vs. Saline	32	4.938	**0.009**
		90 min vs. Baseline	35	5.401	**0.003**
		90 min vs. Saline	32	4.938	**0.009**
Phrenic tonic activity (normalized to baseline)	left	30 min vs. Baseline	32	4.938	**0.009**
		30 min vs. Saline	42	6.481	**<0.001**
		60 min vs. Baseline	33	5.092	**0.006**
		60 min vs. Saline	43	6.635	**<0.001**
		90 min vs. Saline	31	4.783	**0.013**
	right	15 min vs. Baseline	161.81	6.077	**0.002**
		15 min vs. Saline	167.132	6.277	**0.001**
		30 min vs. Baseline	227.376	8.54	**<0.001**
		30 min vs. Saline	232.698	8.74	**<0.001**
		30 min vs. 5 min	151.316	5.683	**0.004**
		60 min vs. Baseline	222.864	8.37	**<0.001**
		60 min vs. Saline	228.186	8.57	**<0.001**
		60 min vs. 5 min	146.805	5.514	**0.005**
		90 min vs. Baseline	198.832	7.468	**<0.001**
		90 min vs. Saline	204.154	7.668	**<0.001**
		90 min vs. 5 min	122.773	4.611	**0.031**
Systolic blood pressure	NA	5 min vs. Baseline	33.737	6.017	**0.002**
		5 min vs. Saline	33.977	6.06	**0.002**
		15 min vs. Baseline	26.184	4.67	**0.028**
		15 min vs. Saline	26.423	4.713	**0.026**
		90 min vs. Baseline	24.843	4.431	**0.043**
		90 min vs. Saline	25.083	4.474	**0.04**
Diastolic blood pressure	NA	5 min vs. Baseline	27.201	9.186	**<0.001**
		5 min vs. Saline	20.438	6.902	**<0.001**
		15 min vs. Baseline	23.277	7.861	**<0.001**
		15 min vs. Saline	16.515	5.577	**0.005**
		30 min vs. Baseline	17.364	5.864	**0.003**
		60 min vs. Baseline	21.693	7.326	**<0.001**
		60 min vs. Saline	14.931	5.042	**0.014**
		90 min vs. Baseline	26.929	9.094	**<0.001**
		90 min vs. Saline	20.167	6.811	**<0.001**
Mean arterial blood pressure	NA	5 min vs. Baseline	29.379	8.053	**<0.001**
		5 min vs. Saline	24.951	6.839	**<0.001**
		15 min vs. Baseline	24.246	6.646	**<0.001**
		15 min vs. Saline	19.818	5.432	**0.006**
		30 min vs. Baseline	17.025	4.667	**0.028**
		60 min vs. Baseline	20.993	5.754	**0.003**
		60 min vs. Saline	16.565	4.54	**0.036**
		90 min vs. Baseline	26.234	7.191	**<0.001**
		90 min vs. Saline	21.806	5.977	**0.002**

Supplemental Table 5. *Statistical summary for impact of DREADD activation on phrenic nerve activity in ChAT-Cre mice.* Time points are in reference to minutes passed since J60 infusion. Summary data presented in Figure 5. RM = repeated measures, df = degrees of freedom. Bolded p-values indicates p<0.05.

REFERENCES

Berlowitz, D. J., Wadsworth, B. & Ross, J. Respiratory problems and management in people with spinal cord injury. *Breathe (Sheff)* **12**, 328-340 (2016). https://doi.org/10.1183/20734735.012616

Brown, R., DiMarco, A. F., Hoit, J. D. & Garshick, E. Respiratory dysfunction and management in spinal cord injury. *Respir Care* **51**, 853-868;discussion 869-870 (2006).

Perrin, C., Unterborn, J. N., Ambrosio, C. D. & Hill, N. S. Pulmonary complications of chronic neuromuscular diseases and their management. *Muscle Nerve* **29**, 5-27 (2004). https://doi.org/10.1002/mus.10487

Mehta, S. Neuromuscular disease causing acute respiratory failure. *Respir Care* **51**, 1016-1021; discussion 1021-1013 (2006).

Burakgazi, A. Z. & Höke, A. Respiratory muscle weakness in peripheral neuropathies. *J Peripher Nerv Syst* **15**, 307-313 (2010). https://doi.org/10.1111/j.1529-8027.2010.00293.x

Fuller, D. D. *et al.* The respiratory neuromuscular system in Pompe disease. *Respir Physiol Neurobiol* **189**, 241-249 (2013). https://doi.org/10.1016/j.resp.2013.06.007

Jordan, A. S. & White, D. P. Pharyngeal motor control and the pathogenesis of obstructive sleep apnea. *Respir Physiol Neurobiol* **160**, 1-7 (2008). https://doi.org/10.1016/j.resp.2007.07.009

Doyle, B. M. *et al.* Gene delivery to the hypoglossal motor system: preclinical studies and translational potential. *Gene Ther* (2021). https://doi.org/10.1038/s41434-021-00225-1

Zhu, H. & Roth, B. L. DREADD: a chemogenetic GPCR signaling platform. *Int J Neuropsychopharmacol* **18** (2014). https://doi.org/10.1093/ijnp/pyu007

Alexander, G. M. *et al.* Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. *Neuron* **63**, 27-39 (2009). https://doi.org/10.1016/j.neuron.2009.06.014

Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc Natl Acad Sci U S A* **104**, 5163-5168 (2007). https://doi.org/10.1073/pnas.0700293104

Fleury Curado, T. *et al.* Chemogenetic stimulation of the hypoglossal neurons improves upper airway patency. *Scientific reports* **7**, 44392 (2017). https://doi.org/10.1038/srep44392

Fleury Curado, T. A. *et al.* Silencing of Hypoglossal Motoneurons Leads to Sleep Disordered Breathing in Lean Mice. *Front Neurol* **9**, 962 (2018). https://doi.org/10.3389/fneur.2018.00962

Fleury Curado, T. *et al.* Designer Receptors Exclusively Activated by Designer Drugs Approach to Treatment of Sleep-disordered Breathing. *American journal of respiratory and critical care medicine* **203**, 102-110 (2021). https://doi.org/10.1164/rccm.202002-0321OC

Singer, M. L. *et al.* Chemogenetic activation of hypoglossal motoneurons in a mouse model of Pompe disease. *Journal of neurophysiology* **128**, 1133-1142 (2022). https://doi.org/10.1152/jn.00026.2022

Horton, G. A. *et al.* Activation of the Hypoglossal to Tongue Musculature Motor Pathway by Remote Control. *Sci Rep* **7**, 45860 (2017). https://doi.org/10.1038/srep45860

Fuller, D. D., Rana, S., Smuder, A. J. & Dale, E. A. The phrenic neuromuscular system. *Handbook of clinical neurology* **188**, 393-408 (2022). https://doi.org/10.1016/B978-0-323-91534-2.00012-6

DiMarco, A. F. & Kowalski, K. E. Activation of inspiratory muscles via spinal cord stimulation. *Respiratory physiology & neurobiology* **189**, 438-449 (2013). https://doi.org/10.1016/j.resp.2013.06.001

Jensen, V. N., Seedle, K., Turner, S. M., Lorenz, J. N. & Crone, S. A. V2a Neurons Constrain Extradiaphragmatic Respiratory Muscle Activity at Rest. *eNeuro* **6** (2019). https://doi.org/10.1523/ENEURO.0492-18.2019

Alilain, W. J. *et al.* Light-induced rescue of breathing after spinal cord injury. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**, 11862-11870 (2008). https://doi.org/10.1523/JNEUROSCI.3378-08.2008

Satkunendrarajah, K., Karadimas, S. K., Laliberte, A. M., Montandon, G. & Fehlings, M. G. Cervical excitatory neurons sustain breathing after spinal cord injury. *Nature* **562**, 419-422 (2018). https://doi.org/10.1038/s41586-018-0595-z

Jensen, V. N., Alilain, W. J. & Crone, S. A. Role of Propriospinal Neurons in Control of Respiratory Muscles and Recovery of Breathing Following Injury. *Front Syst Neurosci* **13**, 84 (2019). https://doi.org/10.3389/fnsys.2019.00084

Lane, M. A. Spinal respiratory motoneurons and interneurons. *Respiratory physiology & neurobiology* **179**, 3-13 (2011). https://doi.org/10.1016/j.resp.2011.07.004

Qiu, K., Falk, D. J., Reier, P. J., Byrne, B. J. & Fuller, D. D. Spinal delivery of AAV vector restores enzyme activity and increases ventilation in Pompe mice. *Molecular therapy : the journal of the American Society of Gene Therapy* **20**, 21-27 (2012). https://doi.org/10.1038/mt.2011.214

McGuire, M., Zhang, Y., White, D. P. & Ling, L. Phrenic long-term facilitation requires NMDA receptors in the phrenic motonucleus in rats. *J Physiol* **567**, 599-611 (2005). https://doi.org/10.1113/jphysiol.2005.087650

Lee, K. Z., Sandhu, M. S., Dougherty, B. J., Reier, P. J. & Fuller, D. D. Hypoxia triggers short term potentiation of phrenic motoneuron discharge after chronic cervical spinal cord injury. *Exp Neurol* **263**, 314-324 (2015). https://doi.org/10.1016/j.expneurol.2014.10.002

Thakre, P. P. & Fuller, D. D. Pattern sensitivity of ampakine-hypoxia interactions for evoking phrenic motor facilitation in anesthetized rat. *J Neurophysiol* **131**, 216-224 (2024). https://doi.org/10.1152/jn.00315.2023

Thakre, P. P., Sunshine, M. D. & Fuller, D. D. Ampakine pretreatment enables a single hypoxic episode to produce phrenic motor facilitation with no added benefit of additional episodes. *J Neurophysiol* **126**, 1420-1429 (2021). https://doi.org/10.1152/jn.00307.2021

Drorbaug, J. E. & Fenn, W. O. A BAROMETRIC METHOD FOR MEASURING VENTILATION IN NEWBORN INFANTS. *Pediatrics* **16**, 81-87 (1955). https://doi.org/10.1542/peds.16.1.81

Li, K. *et al.* Overexpression of the astrocyte glutamate transporter GLT1 exacerbates phrenic motor neuron degeneration, diaphragm compromise, and forelimb motor dysfunction following cervical contusion spinal cord injury. *J Neurosci* **34**, 7622-7638 (2014). https://doi.org/10.1523/jneurosci.4690-13.2014

Li, K. *et al.* GLT1 overexpression in SOD1(G93A) mouse cervical spinal cord does not preserve diaphragm function or extend disease. *Neurobiology of disease* **78**, 12-23 (2015). https://doi.org/10.1016/j.nbd.2015.03.010

Thakre, P. P., Rana, S., Benevides, E. S. & Fuller, D. D. Targeting drug or gene delivery to the phrenic motoneuron pool. *Journal of neurophysiology* **129**, 144-158 (2023). https://doi.org/10.1152/jn.00432.2022

Mantilla, C. B., Zhan, W. Z. & Sieck, G. C. Retrograde labeling of phrenic motoneurons by intrapleural injection. *J Neurosci Methods* **182**, 244-249 (2009). https://doi.org/10.1016/j.jneumeth.2009.06.016

Gransee, H. M., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Targeted delivery of TrkB receptor to phrenic motoneurons enhances functional recovery of rhythmic phrenic activity after cervical spinal hemisection. *PLoS One* **8**, e64755 (2013). https://doi.org/10.1371/journal.pone.0064755

Martínez-Gálvez, G. *et al.* TrkB gene therapy by adeno-associated virus enhances recovery after cervical spinal cord injury. *Exp Neurol* **276**, 31-40 (2016). https://doi.org/10.1016/j.expneurol.2015.11.007

Gransee, H. M., Gonzalez Porras, M. A., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Motoneuron glutamatergic receptor expression following recovery from cervical spinal hemisection. *J Comp Neurol* **525**, 1192-1205 (2017). https://doi.org/10.1002/cne.24125

Keeler, A. M. *et al.* Intralingual and Intrapleural AAV Gene Therapy Prolongs Survival in a SOD1 ALS Mouse Model. *Mol Ther Methods Clin Dev* **17**, 246-257 (2020). https://doi.org/10.1016/j.omtm.2019.12.007

Tervo, D. G. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. *Neuron* **92**, 372-382 (2016). https://doi.org/10.1016/j.neuron.2016.09.021

Smith, K. S., Bucci, D. J., Luikart, B. W. & Mahler, S. V. Dreadds: Use and application in behavioral neuroscience. *Behav Neurosci* **135**, 89-107 (2021). https://doi.org/10.1037/bne0000433

Roth, B. L. DREADDs for Neuroscientists. *Neuron* **89**, 683-694 (2016). https://doi.org/10.1016/j.neuron.2016.01.040

Ouali Alami, N. *et al.* Multiplexed chemogenetics in astrocytes and motoneurons restore blood-spinal cord barrier in ALS. *Life Sci Alliance* **3** (2020). https://doi.org/10.26508/lsa.201900571

Saxena, S. *et al.* Neuroprotection through excitability and mTOR required in ALS motoneurons to delay disease and extend survival. *Neuron* **80**, 80-96 (2013). https://doi.org/10.1016/j.neuron.2013.07.027

Jaiswal, P. B., Mistretta, O. C., Ward, P. J. & English, A. W. Chemogenetic Enhancement of Axon Regeneration Following Peripheral Nerve Injury in the SLICK-A Mouse. *Brain Sci* **8** (2018). https://doi.org/10.3390/brainsci8050093

Jaiswal, P. B. & English, A. W. Chemogenetic enhancement of functional recovery after a sciatic nerve injury. *Eur J Neurosci* **45**, 1252-1257 (2017). https://doi.org/10.1111/ejn.13550

Streeter, K. A. *et al.* Mid-cervical interneuron networks following high cervical spinal cord injury. *Respir Physiol Neurobiol* **271**, 103305 (2020). https://doi.org/10.1016/j.resp.2019.103305

Lane, M. A. *et al.* Cervical prephrenic interneurons in the normal and lesioned spinal cord of the adult rat. *The Journal of comparative neurology* **511**, 692-709 (2008). https://doi.org/10.1002/cne.21864

Molkov, Y. I., Rubin, J. E., Rybak, I. A. & Smith, J. C. Computational models of the neural control of breathing. *Wiley Interdiscip Rev Syst Biol Med* **9** (2017). https://doi.org/10.1002/wsbm.1371

Dempsey, J. A. & Welch, J. F. Control of Breathing. *Semin Respir Crit Care Med* **44**, 627-649 (2023). https://doi.org/10.1055/s-0043-1770342

Rana, S., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Cervical spinal hemisection alters phrenic motor neuron glutamatergic mRNA receptor expression. *Experimental neurology* **353**, 114030 (2022). https://doi.org/10.1016/j.expneurol.2022.114030

Qiu, K., Lane, M. A., Lee, K. Z., Reier, P. J. & Fuller, D. D. The phrenic motor nucleus in the adult mouse. *Experimental neurology* **226**, 254-258 (2010). https://doi.org/10.1016/j.expneurol.2010.08.026

Rana, S., Sieck, G. C. & Mantilla, C. B. Heterogeneous glutamatergic receptor mRNA expression across phrenic motor neurons in rats. *Journal of neurochemistry* **153**, 586-598 (2020). https://doi.org/10.1111/jnc.14881

Rana, S., Mantilla, C. B. & Sieck, G. C. Glutamatergic input varies with phrenic motor neuron size. *J Neurophysiol* **122**, 1518-1529 (2019). https://doi.org/10.1152/jn.00430.2019