Chemogenetic stimulation of phrenic motor output and diaphragm activity

Ethan S. Benevides[1,2], Prajwal P. Thakre[1,2,3], Sabhya Rana[1,2,3], Michael D. Sunshine[1,2,3], Victoria N. Jensen[1,2,3], Karim Oweiss[3,4] and David D. Fuller[1,2,3]

[1] Department of Physical Therapy, University of Florida, Gainesville, FL, 32601
[2] Breathing Research and Therapeutics Center, University of Florida, Gainesville, FL, 32601
[3] McKnight Brain Institute, University of Florida, Gainesville, FL, 32601
[4] Department of Electrical and Computer Engineering, University of Florida, Gainesville, FL, 32601

Running Title:	Chemogenetic stimulation of phrenic motor output and diaphragm activity
Article Type:	Original Research
Number of Figures:	9
Number of Tables:	6

Corresponding Author: David Fuller, Ph.D
Professor
Department of Physical Therapy
University of Florida
PO Box 100514, UFHSC
Gainesville, FL, 32610-0154
Phone (352) 273-6634
dfuller@phhp.ufl.edu

ORCIDs: Ethan S. Benevides: 0000-0002-2472-0606
Thakre P. Prajwal: 0000-0001-5440-7023
Sabhya Rana: 0000-0002-1303-6614
Michael D. Sunshine: 0000-0002-7133-4185
Victoria N. Jensen: 0000-0002-6374-6323
Karim Oweiss: 0000-0002-1175-7258
David D. Fuller: 0000-0001-5013-5972

Author Contributions: E.S.B.: design of the experiment, data acquisition, analysis, and interpretation; initial draft of the manuscript. T.P.P: design of the study, data acquisition, analysis, and interpretation; manuscript editing. S.R.: design of the study, acquisition of the data, analysis, and interpretation; review of manuscript draft. M.D.S: data analysis and interpretation; review of manuscript draft. V.N.J.: acquisition; review of manuscript draft. K.O.: provided ChAT-Cre rats; review of manuscript draft. D.D.F.: conception and design of the study, interpretation of the data, revisions, and final approval of the manuscript. All authors read and approved the submitted version.

Author's Disclosure Statement: Authors declare no competing financial interests.

Funding Information: Support for this work was provided by the National Institutes of Health: 5R01HD052682-14 (DDF), and a Graduate School Funding Award from the University of Florida (ESB).

Data and Code Availability: The datasets along with MATLAB and R code generated during the current study are available from the corresponding author upon reasonable request.

Animal Studies: All procedures described in this manuscript involving rats, mice, and tissue were approved by the University of Florida Institutional Animal Care and Use Committee (protocol #202107438) and in strict accordance with the US National Institute of Health (NIH) Guide for the Care and Use of Laboratory Animals.

Abbreviations: DREADD (designer receptors exclusively activated by designer drugs); EMG (electromyography), AUC (area under the curve); J60 (JHU37160); AAV (adeno-associated virus); VT (tidal volume); ChAT (choline acetyltransferase); Cre (Cre recombinase)
Keywords: DREADD, Chemogenetics, Phrenic, Diaphragm EMG, Plethysmography, Adeno-associated virus, Respiratory neural control.

ABSTRACT

Impaired respiratory motor output contributes to morbidity and mortality in many neurodegenerative diseases and neurologic injuries. We investigated if expressing designer receptors exclusively activated by designer drugs (DREADDs) in the mid-cervical spinal cord could effectively stimulate phrenic motor output to increase diaphragm activation. Two primary questions were addressed: 1) does effective DREADD-mediated diaphragm activation require focal expression in phrenic motoneurons (vs. non-specific mid-cervical expression), and 2) can this method produce a sustained increase in inspiratory tidal volume? Wild type (C57/bl6) and ChAT-Cre mice received bilateral intraspinal (C4) injections of an adeno-associated virus (AAV) encoding the hM3D(Gq) excitatory DREADD. In wild-type mice, this produced non-specific DREADD expression throughout the mid-cervical ventral horn. In ChAT-Cre mice, a Cre-dependent viral construct was used to drive neuronal DREADD expression in the C4 ventral horn, targeting phrenic motoneurons. Diaphragm EMG was recorded in isoflurane-anesthetized spontaneously breathing mice at 4-9 weeks post-AAV delivery. The DREADD ligand JHU37160 (J60) caused a bilateral, sustained (>1 hour) increase in inspiratory EMG bursting in both groups; the relative increase was greater in ChAT-Cre mice. Additional experiments in ChAT-Cre rats were conducted to determine if spinal DREADD activation could increase inspiratory tidal volume (VT) during spontaneous breathing, assessed using whole-body plethysmography without anesthesia. Three-to-four months after intraspinal (C4) injection of AAV driving Cre-dependent hM3D(Gq) expression, intravenous J60 resulted in a sustained (>30 min) increase in VT. Subsequently, phrenic nerve recordings performed under urethane anesthesia confirmed that J60 evoked a > 200% increase in inspiratory output. We conclude that targeting mid-cervical spinal DREADD expression to the phrenic motoneuron pool enables ligand-induced, sustained increases in phrenic motor output and VT. Further development of this technology may enable application to clinical conditions associated with impaired diaphragm activation and hypoventilation.

INTRODUCTION

Many respiratory disorders are associated with reduced or impaired activation of respiratory motoneurons. Neurologic injuries (e.g., traumatic spinal cord injury, stroke) and neurodegenerative conditions (e.g., ALS, Pompe disease) will often result in decreased respiratory motor output, including impaired activation of the phrenic motoneurons which innervate the diaphragm[1-6]. Another prominent example is obstructive sleep apnea, in which pharyngeal motoneurons have reduced output during sleep[7]. Treatment options that increase respiratory motoneuron activation to improve breathing are limited. However, designer receptors exclusively activated by designer drugs (DREADDs) may have use in this regard[8]. Structurally derived from naturally occurring G-protein coupled receptors, DREADDs have been engineered to respond exclusively to exogenous ligands that are otherwise biologically inert[9-11]. This provides a means to selectively stimulate cells expressing the DREADD. Prior studies have used DREADDs to stimulate upper airway muscle activation during breathing[8]. For example, following expression of DREADDs in murine hypoglossal motoneurons, tongue electromyogram (EMG) activity can be increased using DREADD ligands[12-16]. This response is functionally beneficial as shown by increased patency of the upper airway[12].

The present study focused on chemogenetic activation of the phrenic neuromuscular system. Phrenic motoneurons provide motor innervation of the diaphragm muscle and are located in the mid-cervical (C3-5) spinal cord[17]. We tested the hypothesis that expressing DREADDs in the mid-cervical spinal cord would enable systemic (intravenous or intraperitoneal) delivery of a selective DREADD ligand to produce sustained increases in the respiratory-related activation of the diaphragm. In doing so, we addressed two important questions. First, we determined if effective diaphragm activation requires focal DREADD expression targeting phrenic motoneurons, or if non-specific expression in mid-cervical interneurons and phrenic motoneurons would be sufficient. This question derives from prior studies of cervical spinal cord stimulation. A compelling body of work, with studies in multiple species, demonstrates that non-specific activation of cervical spinal networks can be highly effective at increasing

diaphragm activation[18-20]. One theory to explain this result is that a general increase in the excitability of

cervical propriospinal networks leads to increased phrenic motoneuron activation[21-23]. There is also

evidence that phrenic motoneurons can integrate multiple synaptic inputs in a manner that produces

orderly recruitment[18]. Accordingly, DREADD-induced activation of mid-cervical neurons or networks

may be sufficient for ligand-induced diaphragm activation. On the other hand, DREADD expression may

need to be restricted to phrenic motoneurons if the goal is to produce inspiratory-related diaphragm

activation. To address this question, we studied diaphragm responses in two mouse models: 1) a wild-type

model in which DREADDs were non-specifically expressed in the C3-5 spinal cord, encompassing

interneurons populations and motoneurons, and 2) a choline acetyltransferase (ChAT)-Cre transgenic

model in which DREADD expression was restricted to ChAT-positive neurons in the ventral C3-5 spinal

cord, targeting the phrenic motoneuron pool.

The second question we addressed was if phrenic motoneuron activation via cervical spinal cord

DREADDs could produce a sustained increase in inspiratory tidal volume in unanesthetized,

spontaneously breathing animals. While the previous results from the hypoglossal motor system[12,14-16]

provide a proof-of-concept that DREADDs can stimulate respiratory motoneuron activity, whether a

sustained increase in tidal volume could be evoked by expressing DREADDs in phrenic motoneuron was

uncertain. For example, during spontaneous breathing, a DREADD-induced increase in phrenic

motoneuron excitability, and thus diaphragm activation, could be rapidly offset by decreases in

bulbospinal neural drive to the phrenic motor pool, secondary to reduced arterial CO_2 or increased vagal-

mediated inhibition. An increase in diaphragm activation could also trigger a decrease in accessory

respiratory muscle activation, thereby attenuating or preventing increases in tidal volume. Lastly, data

from the hypoglossal motor system[12,14-16], as well as our initial results in the anesthetized mouse indicated

that both phasic (i.e., during the inspiratory period) and tonic (i.e., occurring across the respiratory cycle)

activation of the diaphragm would increase after DREADD activation, and how this would impact tidal

volume was not clear. Accordingly, we studied ChAT-Cre rats using whole-body plethysmography and a

direct measure of phrenic motor output via nerve recordings. The plethysmography studies allowed us to

determine if DREADD activation of phrenic motoneurons causes a sustained increase in tidal volume and

ventilation during spontaneous breathing in the unanesthetized rat. The nerve recordings were done under

urethane anesthesia and enabled direct quantification of DREADD activation on the neural drive of the

diaphragm while controlling variables including arterial CO_2 and lung volume.

Collectively the results of this work demonstrate that mid-cervical spinal DREADD expression

enables the J60 ligand to produce a sustained increase in the neural drive to the diaphragm, producing an

increase in tidal volume during spontaneous breathing. Further development of this technology may

enable application to clinical conditions associated with impaired diaphragm activation and

hypoventilation.

RESULTS

Diaphragm EMG responses in wild-type mice. Wild-type mice underwent bilateral injections of AAV9-

hSyn-HA-hM3D(Gq)-mCherry into the ventral horns at spinal segment C4. Following a four-to-five-

weeks incubation period, mice underwent terminal diaphragm EMG recordings before and after

application of the selective DREADD ligand, J60. On average, wild-type mice showed increases in

diaphragm EMG output in at least one hemidiaphragm after J60 administration (**Figure 1**). The area

under the curve (AUC) of the rectified and integrated diaphragm EMG was significantly increased after

DREADD activation (**Figure 1d**) in both the left (**p = 0.002**; **Table S1**) and right (**p = 0.002**; **Table S1**)

hemidiaphragm. Additionally, the peak-to-peak amplitude of the rectified and integrated diaphragm EMG

burst activity increased bilaterally following J60 administration (<u>Left hemidiaphragm</u>: p = 0.056; <u>Right</u>

<u>hemidiaphragm</u>: **p = 0.01**; **Figure 1e**; **Table S1**). Lastly, the tonic activity of the diaphragm was assessed

(**Figure 1f**). Similar to the previous measures of EMG output both the left (**p = 0.052**; **Table S1**) and right

(**p < 0.001**; **Table S1**) hemidiaphragm exhibited a significant increase in tonic activity following J60

administration. Respiratory rate was consistent for the duration of the experiment. Notably, there was no

substantial change in the respiratory rate of these spontaneous breathing mice after J60 administration (p

= 0.863; **Figure 1g**; **Table S1**).

In all experiments, the selective DREADD ligand, J60, produced an increase in diaphragm EMG

burst amplitude during inspiration. However, this increase was not always detected in both the left and

right hemidiaphragm EMG recordings. Five mice showed a bilateral increase in diaphragm output after

J60 administration (**Figure 1**), four mice showed a response that was limited to the right hemidiaphragm,

and two showed a response that was limited to the left hemidiaphragm (**Figure 1**).

Diaphragm EMG responses in ChAT-Cre mice. ChAT-Cre mice received bilateral intraspinal injections

of AAV9-hSyn-DIO-hM3D(Gq)-mCherry into the ventral horns at C4. ChAT-Cre mice underwent

terminal diaphragm EMG recording using the same protocol as wild-type mice, following a four-to-nine-

week incubation. All mice (n = 9/9) showed an increase in diaphragm EMG output in response to the J60

DREADD ligand in at least one hemidiaphragm (**Figure 2**). On average, both left (**p = 0.011**; **Table S2**)

and right (**p < 0.001**; **Table S2**) diaphragm EMG AUC increased over time after J60 administration

(**Figure 2d**). Diaphragm EMG peak-to-peak amplitude had a similar, bilateral increase following J60

delivery (Left hemidiaphragm: **p = 0.013**; Right hemidiaphragm: **p < 0.001**; **Table S2**; **Figure 2e**).

Lastly, tonic activity also showed an increase over time after J60 administration (**Figure 2f**) for both the

left (**p = 0.002**; **Table S2**) and right (**p < 0.001**; **Table S2**) hemidiaphragm. Respiratory rate decreased

significantly over time after J60 administration (**p < 0.001**; **Figure 2g**; **Table S2**). Apart from two mice

that showed a unilateral EMG response that was limited to the right hemidiaphragm the remaining ChAT-

Cre mice (n = 7/9) had bilateral increases in diaphragm EMG output following J60 administration

(**Figure 2**).

Wild type vs. ChAT-Cre comparison. Diaphragm EMG responses of wild-type and ChAT-Cre mice were

compared at the 30-minute post-J60 time point (**Figure 3**). Left hemidiaphragm responses to J60 were

similar between the two groups across all outcome measures (AUC: p = 0.998; Peak-to-peak amplitude: p

= 0.771; Tonic activity: p = 0.160; **Table S3**; **Figure 3a-c**). However, right hemidiaphragm responses to

J60 differed across AUC (**Figure 3d**), peak-to-peak amplitude (**Figure 3e**), and tonic activity (**Figure 3f**)

with ChAT-Cre mice on average having larger magnitude responses compared to wild-type mice (AUC: **p**

= 0.0417; Peak-to-peak amplitude: **p = 0.00403**; Tonic activity: **p = 0.00207**; **Table S3**). Respiratory rate

was not different between the two groups (p = 0.382; **Table S3**; **Figure 3g**).

The *a priori* expected recording duration for these experiments in anesthetized and spontaneously

breathing mice was 90-minute. However, five of the eleven total wild-type mice in this experiment did

not survive for this duration. It is unclear if this was a non-specific result associated with prolonged

anesthesia, or if this was physiologically related to DREADD activation. No mice had evidence of

adverse reaction in the initial 30-minutes following delivery of J60. Of the five mice which did not

survive, three mice died between the 30- and 60-minute time points after J60, and two mice died just prior

to the 90-minute time point. In contrast, all mice in the ChAT-Cre cohort (9 of 9) survived the total

duration of the experimental protocol. A chi-square evaluation of the survival proportions was not

statistically significant (Chi-squared = 3.2997, df = 1, p = 0.06929). However, considering the sample

size, the results suggest some association between mouse strain (i.e., wild type, ChAT-Cre) and death,

suggesting that non-specific DREADD activation may be contraindicated.

J60 control experiments. The DREADD ligand was administered to wild-type animals with no

hM3D(Gq) expression in the mid-cervical spinal cord (**Figure S1**). This was done to determine the impact

of J60 administration on diaphragm EMG in the absence of DREADD expression (n= 2 C57/bl mice; n=

3 Sprague Dawley rats). There was no discernable impact of J60 on the diaphragm EMG burst amplitude

(mV) (**Figure S1a-b**). Responses were also not different between sham (saline) and J60 when normalized

to baseline activity (**Figure S1c-d**).

ChAT-Cre rats – Plethysmography and Phrenic Nerve Recordings. A small cohort of ChAT-Cre rats

underwent anesthetized diaphragm EMG recordings to ensure DREADD responses similar to the mouse

cohorts could be obtained in rats. ChAT-Cre rats (n = 4) underwent bilateral, intraspinal injections of

AAV9-hSyn-DIO-hM3D(Gq)-mCherry into the ventral horns at C4 to introduce the hM3D(Gq)

DREADD transgene into phrenic motoneurons. Four of four rats showed increased diaphragm EMG

output after DREADD activation (**Figure S2**). With that knowledge, we used a separate group of ChAT-

Cre rats (n = 9; n = 3 females) to assess the effects of DREADD activation on ventilation. Whole-body

plethysmography was used to measure breathing frequency, tidal volume, and minute ventilation before

and after intravenous delivery of saline (sham) and J60 (**Figure 4**). Delivery of the J60 ligand resulted in

an increase in inspiratory tidal volume compared to sham infusion (<u>Normalized to Weight (ml/kg)</u>: Main

effect of Treatment: **p = 0.037**; **Figure 4a**; <u>Normalized to Baseline</u>: Main effect of Treatment: p = 0.091;

Figure 4d; **Table S4**). Respiratory rate appeared to be unaffected by DREADD activation and was

similar between sham and J60 conditions (<u>Respiratory Rate</u>: Main effect of Treatment: p = 0.582; **Figure**

4b; <u>Respiratory Rate Normalized to Baseline</u>: Main effect of Treatment: p = 0.774; **Figure 4e**; **Table S4**).

Minute ventilation was slightly elevated in the J60 condition vs sham; however, this increase did not

160 reach the threshold for statistical significance (<u>Normalized to Body Weight</u>: Main effect of Treatment: $p =$

161 0.194; **Figure 4c**; <u>Normalized to Baseline</u>: Main effect of Treatment: $p = 0.337$; **Figure 4f; Table S4**).

162 Responses to a hypercapnic-hypoxia ventilatory challenge were also assessed (**Figure S3**). Tidal volume

163 (**Figure S3a**; <u>Normalized to Weight (ml/kg)</u>: $p = 0.845$; **Figure S3d**; <u>Normalized to baseline</u>: $p = 0.643$),

164 respiratory rate (**Figure S3b**; <u>Respiratory Rate</u>: $p = 0.262$; **Figure S3e**; <u>Rate Normalized to Baseline</u>: $p =$

165 0.734), and minute ventilation (**Figure S3c**; <u>Normalized to Body Weight</u>: $p = 0.697$; **Figure S3f**; <u>Rate</u>

166 <u>Normalized to Baseline</u>: $p = 0.912$) did not differ between J60 vs sham condition during hypercapnic-

167 hypoxic ventilatory challenges.

168   Phrenic nerve recordings were made to directly assess the effects of DREADD activation

169 on phrenic output. There was no detectable relationship between time post-AAV injection and phrenic

170 response to DREADD activation (Pearson correlation; <u>Left peak-to-peak response</u>: $p = 0.215$; <u>Right peak-</u>

171 <u>to-peak response</u>: $p = 0.318$).

172   Application of the J60 ligand caused a rapid, sustained, and bilateral increase in phrenic nerve

173 efferent burst amplitude (<u>Left phrenic peak-to-peak amplitude (normalized to baseline)</u>: **$p < 0.001$**; <u>Right</u>

174 <u>phrenic peak-to-peak amplitude (normalized to baseline)</u>: **$p < 0.001$**; **Table S5**; **Figure 5b-c**), whereas

175 saline injection had no impact. The increase in phrenic burst amplitude lasted up to 100 minutes post-J60

176 administration, at which point the experiment was terminated. Application of the J60 ligand also resulted

177 in an increase in phrenic tonic activity (<u>Left phrenic tonic activity (normalized to baseline)</u>: **$p < 0.001$**;

178 <u>Right phrenic tonic activity (normalized to baseline)</u>: **$p < 0.001$**; **Table S5**; **Figure 5d-e**).

179   Heart rate, systolic and diastolic blood pressure, mean arterial blood pressure (MAP), as

180 well as respiratory rate, were also assessed (**Figure 5f-j**). Application of J60 did not affect heart rate ($p =$

181 0.587; **Table S5**; **Figure 5h**) or respiratory rate ($p = 0.282$; **Table S5; Figure 5j**) but did result in a

182 decrease in both systolic (**$p < 0.001$**; **Table S5**; **Figure 5f**) and diastolic blood pressure (**$p < 0.001$**; **Table**

183 **S5**; **Figure 5g**) as well as MAP (**$p < 0.001$**; **Table S5**; **Figure 5i**).

Histological analysis. We performed a qualitative analysis of the mid-cervical spinal cord from each

animal to assess the extent of mCherry fluorophore expression (**Figure S4**). All mice from both cohorts

showed evidence of mCherry expression in at least one segment of the mid-cervical spinal cord (**Figure

6**) with the exception n = 1 ChAT-Cre mouse. This mouse was excluded from all analyses based on *a

priori* exclusion criteria, which stipulated animals must show evidence of mCherry expression in the grey

matter of at least one spinal segment from C3-C6 to be included in the final analysis. A summary of the

results is given in **Table 1**.

Patterns of expression were relatively homogenous in wild-type animals. In this cohort, the number of

mice with positive mCherry expression in the grey matter increased on the rostral-caudal axis. Positive

mCherry counts were comparable on both the dorsal-ventral and left-right axes, with a majority of mice

expressing mCherry in all four quadrants. The diaphragm EMG responses to J60, on average, exhibited

similarity between the left and right hemidiaphragm in these mice, aligning with the observed pattern of

mCherry expression. (**Figure 1d-f**).

In contrast, mCherry expression in the ChAT-Cre mice cohort was more prevalent in the ventral

horns and the right side of the cord. Like the wild-type mice, there was a slight trend for increased

mCherry expression moving rostral to caudal. Clear mCherry expression was detectable in the spinal cord

in of all nine ChAT-Cre mice included in the final data set. One additional ChAT-Cre mouse was

excluded from analysis as it showed no evidence of mCherry in the mid-cervical spine. Interestingly, this

particular mouse appeared to show a modest increase in diaphragm output in response to the J60 ligand in

the left-hemidiaphragm only (~45% increase compared to baseline activity). While this animal was

ultimately excluded from our analysis, it is possible that this mouse did express hM3D(Gq) in the mid-

cervical spinal cord but an issue in tissue processing resulted in an inability to visualize the mCherry

fluorophore in the spinal tissue. All other ChAT-Cre mice showed robust mCherry expression in the

ventral horns of at least one spinal segment from C3-C6. These mice demonstrated a larger average

DREADD response in the right hemidiaphragm than the left (**Figure 2d-f**), possibly stemming from the

fact that a greater number of mice exhibited mCherry expression on the right side compared to the left

(**Figure S5**).

ChAT-Cre rats showed expression predominately in the ventral horns throughout the mid-cervical

spinal cords with the highest levels of expression in spinal segments C4 and C5. Expression in this cohort

was slightly more prominent on the right side of the cord and in the ventral horn. These histological

findings were consistent with the physiological results. Although the magnitude of DREADD response

between the left and right phrenic nerves for this cohort was not statistically different, there was a trend of

slightly higher right phrenic tonic response compared to the left (**Figure 5b-e**). This trend is mirrored in

the pattern of mCherry expression, where expression levels were approximately equal between spinal

segments but tended to be slightly higher in the right ventral horns compared to the left.

DISCUSSION

We describe a novel method to increase diaphragm EMG output by expressing the excitatory

DREADD, hM3D(Gq), in the mid-cervical spinal cord, targeting phrenic motoneurons. Following AAV-

driven expression of the DREADD in the spinal cord, application of the J60 ligand caused sustained

increases in diaphragm output as measured through EMG in spontaneously breathing animals. This

response was also verified using direct recordings of phrenic nerve discharge. Additionally, the DREADD

ligand was able to produce an increase in inspiratory tidal volume in awake, freely behaving animals.

These proof-of-concept studies provide a foundation for further development of this technology towards

clinical application for restoring diaphragm activation in conditions such as cervical spinal cord injury.

Targeted gene delivery to the phrenic motor pool. The intraspinal AAV delivery used here was based on

previous studies demonstrating successful gene delivery to phrenic motoneurons[20 24-26]. For example, mid-

cervical spinal injections of an AAV5 vector encoding the lysosomal enzyme acid alpha-glucosidase

(GAA) in animals with Pompe disease (*Gaa* null) effectively restores spinal GAA enzyme activity [26].

Spinal-delivered viral vectors have also been used to successfully drive local expression of

channelrhodopsin-2 to enable light activation of diaphragm output[20], and to drive expression of the

astrocyte glutamate transporter GLT1 in the area of the phrenic motor nuclei[24]. Other methods that have

been employed to drive gene expression in phrenic motoneurons include intrapleural- and intramuscular

diaphragm injection of viral vectors[27]. Intrapleural delivery requires microinjection to the "pleural space"

between the visceral pleura that lines the lungs and the parietal pleura that covers the thoracic cavity. This

technique[28] effectively targets phrenic motoneurons in rodent models of cervical spinal cord injury[29-31]

and Pompe disease[32]. Intramuscular diaphragm injection allows the vector to enter phrenic nerve

terminals and reach phrenic motoneuron soma via retrograde movement[27]. Direct diaphragm injection

allows for a relatively high target specificity, with the gene of interest expressed almost exclusively in

phrenic motoneurons (although expression can also occur in diaphragm myofibers, depending on the

promoter sequence used). In pilot experiments, we tested intrapleural and intramuscular diaphragm

injections using AAV9 vectors encoding GFP (AAV9-CAG-GFP) or DREADD (AAV9-hSyn-HA-

hM3D(Gq)-mCherry & AAV9-hSyn-DIO-hM3D(Gq)-mCherry). We did not, however, observe

histological or physiological evidence of phrenic motoneuron transduction with these AAV9 vectors.

Direct intraspinal injection[24,26] was therefore used to introduce the hM3D(Gq) into the phrenic motor

nucleus. While this enabled proof-of-concept for targeting DREADDs to the cervical spinal cord and

phrenic motoneurons, the intrapleural or diaphragmatic injection delivery routes might ultimately prove

better for selective phrenic motoneuron targeting. We predict that using different AAV serotypes or

viruses with better retrograde movement (e.g., "AAV retro") could optimize the targeting of phrenic

motoneurons[33].

DREADD-mediated motoneuron activation. DREADD technology is widely used for studying brain and

spinal cord neurons and networks[34,35]. Relatively few studies, however, have examined if and how

DREADDs can be used to activate (or inhibit) lower motoneurons. Regarding the spinal cord, we are

aware of only a few prior publications[36-39]. Two of these studies used pharmacologically selective actuator

module (PSAM), a type of ionotropic chemogenetic receptor, to activate lumbar[36,37] motoneurons, in

mouse models of amyotrophic lateral sclerosis (ALS). In the remaining studies, excitatory DREADDs

were applied to spinal motoneurons to improve axon regeneration following peripheral nerve injury[38,39]. A

small but growing body of work has employed DREADDs to activate hypoglossal (XII) motoneurons in

the brainstem[8]. Collectively, these studies show that once hM3D(Gq) is expressed in XII motoneurons,

DREADD ligands will rapidly produce an increase in the EMG activation of tongue muscles[14,15]. This

increase in tongue muscle output tends to manifest as an increase in the inspiratory-related activation and

tonic discharge across the respiratory cycle. Since increased tongue muscle activation can promote

patency of the upper airway, XII motoneuron DREADD expression has been suggested as a possible

treatment for obstructive sleep apnea[12,14,16]. For the present study, the primary innovation is the first

application of DREADD technology to phrenic motoneurons. This approach was highly effective at

driving sustained activation of the diaphragm muscle, and the underlying mechanisms are discussed next.

Chemogenetic stimulation of breathing. An important consideration is how DREADD-induced increases

in the excitability of spinal neurons, including phrenic motoneurons, interacts with the endogenous neural

control of breathing. Phrenic motoneurons receive a rhythmic, monosynaptic, glutamatergic synaptic

input from medullary neurons. Acting via NMDA and AMPA receptors, this produces phrenic

motoneuron depolarization and subsequent diaphragm muscle contraction[17]. Activating DREADDs on

phrenic motoneurons should lower the threshold for activation via excitatory glutamatergic synaptic

inputs, which would produce a greater output during the inspiratory phase. Alternatively, DREADD

activation could directly lead to phrenic motoneuron action potentials even in the absence of synaptic

input from the brainstem. This latter possibility could explain the tonic discharge (i.e., EMG output across

the entire respiratory cycle) that was noted to occur after delivery of the DREADD ligand. Non-specific

spinal cord DREADD expression, as occurred in the wild-type mice (e.g., **Figure 6**), would likely

produce an increase in the excitability and/or activation of phrenic motoneurons as well as propriospinal

neurons in the immediate vicinity. Neurophysiological[40], as well as anatomical data[23,41], confirm synaptic

connections between mid-cervical interneurons and phrenic motoneurons, making it possible that

DREADD activation of these interneurons impacted the diaphragm motor response in the wild-type mice.

The control of breathing is also impacted by well-established "closed loop" physiologic feedback

mechanisms regulating lung volume and arterial blood gases[42,43]. For example, if DREADD-induced

activation of the diaphragm leads to increased alveolar ventilation, and metabolic rate is not impacted,

then arterial CO_2 values will decrease and the overall neural drive to breathe will also decrease. Vagal

afferent feedback corresponding to increased lung volume also has a powerful inhibitory impact on

inspiration and therefore diaphragm activation. However, the sustained increase in diaphragm EMG and

tidal volume that we observed following application of the DREADD ligand indicates that these

mechanisms, if activated, were not sufficient to fully inhibit the increased phrenic motoneuron output. In

this regard, our additional experiments in which direct recordings of bilateral phrenic nerve discharge are

informative. These nerve recording experiments were done to enable direct evaluation of the impact of

spinal DREADD activation on phrenic motor output while keeping arterial blood gases and lung volume

constant. Under these more rigorously controlled conditions, intravenous delivery of the DREADD ligand

produced a rapid and sustained increase in inspiratory burst amplitude in the phrenic nerve, and with no

impact on the rate of the inspiratory bursts. The relative increase in inspiratory motor output was

considerably greater in the phrenic nerve recording experiments (~250% of baseline) as compared to the

diaphragm EMG response in spontaneously breathing animals (~100% of baseline). This may indicate

that vagal and/or blood gas-related inhibitory mechanisms, as mentioned above, somewhat constrained

the response to the DREADD ligand in the spontaneously breathing animal.

Critique of methods. There are a few caveats that should be discussed. First, the precision of the AAV

delivery could be improved by further refining spinal injection surgical techniques. In the current study,

we used a stereotaxic frame and previously validated coordinates[26,44] to guide the intraspinal AAV

injections. However, we observed variability in the laterality (i.e., left vs. right side of the spinal cord) of

mid-cervical mCherry expression as well as the physiological response to the DREADD ligand,

particularly in the ChAT-Cre mice (e.g., **Figure 2**; **Table 1**). This could have occurred due to subtle

variations of the positioning of the animal within the stereotaxic frame, and/or placement of the needle

tip, leading to slight deviations for the desired coordinates between the left and right phrenic nuclei.

Second, we did not unequivocally verify that the DREADD was expressed in phrenic motoneurons using

retrograde labeling methods[28,45]. However, the phrenic motor nucleus has been well described in the

mouse[46] and the rat[47,48], and the fluorophore (mCherry) expression observed in our experiments is very

clearly in the expected location of phrenic motoneurons (**Figure 6b-c$_i$; Figure S4**). Further, the robust

increase in phrenic motor output after the DREADD ligand, particularly in the ChAT-Cre rat experiment

(**Figure 5**) is further evidence of effective phrenic motoneuron targeting.

Conclusion. Our data support the conclusion that cervical spinal cord directed chemogenetic methods can

be used to produce sustained increases in phrenic motor output, diaphragm activation, and inspiratory

tidal volume. Collectively, the data indicate that DREADDs should be directed exclusively to phrenic

motoneurons vs. non-specific expression in the immediate vicinity. In this regard, improvement of the

AAV delivery methods will increase the selectivity of the approach for more precise targeting of phrenic

motoneurons. Concerning the "translational value" of this work, spinal cord chemogenetics may have

application to clinical conditions associated with an inability to activate the diaphragm. For example,

incomplete cervical spinal cord injury is a condition in which the bulbospinal synaptic inputs to phrenic

motoneurons are interrupted. After incomplete cervical spinal cord injury, focal expression of an

excitatory DREADD in phrenic motoneurons could be used to increase the excitability of these cells,

thereby improving the efficacy of spared bulbospinal synaptic inputs which convey "inspiratory drive".

METHODS

Animals. Experiments were carried out using C5/bl6, wild-type mice (Taconic), ChAT-Cre transgenic mice (B6.129S6-Chattm2(cre)Lowl/J; Jackson Laboratories), and ChAT-Cre transgenic rats (LE-Tg(Chat-Cre)5.1Deis; Rat Resource & Research Center). Animals were singly housed in a controlled environment (12 h light-dark cycle) with food and water *ad libitum*. All experiments were conducted in accordance with the NIH Guidelines Concerning the Care and Use of Laboratory Animals and were approved by the University of Florida Institutional Animal Care and Usage Committee (protocol #202107438). A full experimental timeline for mouse and rat experiments is shown in **Figure S6**, panels a and b, respectively.

Adeno-associated viral vectors. All animals underwent intraspinal injections (see section below) of an AAV vector encoding the excitatory DREADD (hM3D(Gq)) under a human synapsin promoter. Wildtype mice received injections of AAV9-hSyn-HA-hM3D(Gq)-mCherry (titer: 2.44×10^{13} vg/mL) while ChAT-Cre mice and rats received injections of a similar construct with a double-floxed inverted open-reading frame (DIO) allowing for Cre-dependent transgene expression (AAV9-hSyn-DIO-hM3D(Gq)-mCherry; titer: 2.07×10^{12} vg/mL). The pAAV-hSyn-hM3D(Gq)-mCherry (Addgene plasmid # 50474; http://n2t.net/addgene:50474; RRID: Addgene_50474) and pAAV-hSyn-DIO-hM3D(Gq)-mCherry (Addgene plasmid # 44361; http://n2t.net/addgene:44361 ; RRID:Addgene_44361) transgene plasmids were gifts from the laboratory of Dr. Brian Roth at the University of North Carolina. Viral preparations were generated and titered by the University of Florida Powell Gene Therapy Center Vector Core Lab. Vectors were purified by iodixanol gradient centrifugation and anion-exchange chromatography as previously described[49].

Intraspinal injections. An adeno-associated viral vector (AAV) encoding the gene for the excitatory DREADD, hM3D(Gq) was delivered to the mid-cervical spinal cord. Mice were 6-10 weeks old (WT cohort: 7-9 weeks; ChAT-Cre cohort: 6-10 weeks) at the time of injection while ChAT-Cre rats were 2-5 months old. Surgery was performed under aseptic conditions. Mice were anesthetized with isoflurane (induction: 3-4% isoflurane; maintenance: 2-3% isoflurane in 100% O_2) while rats were anesthetized with

a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) delivered intraperitoneally. Animals were

placed prone on a circulating water heating pad to maintain body temperature. A longitudinal incision was

made starting at the base of the skull and extending caudally. The underlying back musculature was

opened from the base of the skull to spinal segment C6. Using a micro-curette, the muscle and connective

tissue overlying laminae C3 to C5 were removed. A laminectomy of the C4 dorsal lamina exposed the

dura mater below. A bilateral durotomy was then performed exposing the spinal cord. A Hamilton syringe

(34-gauge needle) held in a Kopf stereotaxic frame was used to inject 1 µl of AAV9-hSyn-DIO-

hM3D(Gq)-mCherry (ChAT-Cre mice and rats) or AAV9-hSyn-HA-hM3D(Gq)-mCherry (C57/bl6

mice), bilaterally into the ventral horns at C4. Injections were made 0.5 mm lateral to the spinal midline at

a depth of 0.9 mm for mice[26] and 1 mm lateral to midline at a depth of 1.5 mm for rats[44]. The needle was

left to dwell for 5 minutes. Following injections, the overlying muscle and fascia were sutured with

absorbable suture, the skin closed, and the animal returned to its home cage. Animals received a post-

operative analgesia regiment of subcutaneous buprenorphine (1 mg/kg; slow-release formulation) and

carprofen (5 mg/kg) for the first three days after surgery.

Diaphragm EMG recordings. Recordings were conducted using wild-type (n = 11; n = 7 females) and

ChAT-Cre mice (n = 9; n = 6 females; n = 1 excluded from analysis), 4-9 weeks following intraspinal

injections of AAV-DREADD. Mice were anesthetized with 2-3% isoflurane in a closed chamber and then

placed supine on a closed loop heating pad to maintain rectal temperature at 37 ± 0.5 °C (model 700 TC-

1000, CWE Inc.). Mice spontaneously inhaled 2% isoflurane in 100% O_2 for the duration of the

experiment.

A laparotomy was performed and two sets of 50 µm tungsten wires were placed in the mid-costal

region of the left and right hemidiaphragm. The tips of each wire were de-insulated, bent into small

hooks, and inserted through the diaphragm approximately 3 mm apart. The recorded EMG signals were

amplified (1000x) and filtered (100–1000 Hz) using a differential amplifier (A–M systems model 1700).

Signals were digitized at 10 kS/s using a Power 1401 (CED, Cambridge, UK).

Once a stable plane of anesthesia was reached, mice underwent a 10-minute recording to establish

baseline diaphragm EMG parameters. Subsequently, mice received injections of vehicle (100 µl of saline

delivered intraperitoneally (IP)) followed by a 20-minute recording. Mice then received an intraperitoneal

injection of the selective DREADD agonist, JHU37160 (J60; 0.1 mg/kg, HB6261, HelloBio), and

recordings continued for 90 minutes. At the conclusion of each experiment, mice underwent transcardial

perfusion with saline followed by 4% paraformaldehyde. Following perfusion, spinal cords were

harvested for histological analysis.

J60 control experiments. A small cohort of animals (n = 2 C57/bl mice; n = 3 Sprague Dawley rats) was

used to assess the impact of J60 (0.1 mg/kg) on diaphragm EMG activity in the absence of hM3D(Gq)

expression. The animals used in this study include n = 2 C57/bl mice that had undergone intrapleural

injection (i.e., injection to the thoracic cavity) of an AAV9 construct encoding the red fluorescent protein,

mCherry and n = 3 vector naïve Sprague Dawley rats.

Recordings in mice proceeded as described above (see *Diaphragm EMG recordings*). In rat

recordings, rats were induced with 3% isoflurane in 100% O_2 and moved onto a closed-loop heating pad

set to maintain rectal temperature at $37 \pm 1°C$ (model 700 TC-1000, CWE Inc.). Rats were

tracheotomized and ventilated (Model 683; Harvard Apparatus Inc.) with a gas mixture of 50% O_2, and

1% CO_2, balanced with N_2. End-tidal CO_2 was maintained at 45-47 mmHg throughout the experimental

protocol (Capnogard; Novametrix). Rats were converted from isoflurane to urethane anesthesia (2.1 g/kg

at 6 mL/hr; IV). At the competition of urethane dosing lactated Ringer's was administered (2 mL/h; IV) to

keep the animal hydrated and ensure the catheter remained viable for J60 administration. A femoral artery

catheter (polyethylene tubing; PE 50; Intramedic) was placed to enable monitoring of arterial blood

pressure via a transducer amplifier (TA-100, CWE).

At the beginning of the experimental period, rats underwent a 10-minute recording to establish

baseline diaphragm EMG parameters. This was followed by an IV injection of vehicle (0.6 mL of saline)

and a subsequent 20-minute recording. Next, rats received an IV infusion of the J60 agonist (0.1 mg/kg),

and the recording continued for 90 minutes. At the end of the experiment, rats were euthanized via an

overdose of pentobarbital sodium and phenytoin sodium (150 mg/kg) given intravenously. Death was

confirmed by thoracotomy once breathing had ceased, and a heartbeat was no longer detectable.

Whole body plethysmography. ChAT-Cre rats (n = 9; n = 3 females) were studied using flow-through

whole-body plethysmography 14-16 weeks after intraspinal delivery of AAV9-hSyn-DIO-hM3D(Gq)-

mCherry, as described above. A tail vein catheter was placed to allow for intravenous infusion (IV) of the

J60 ligand and vehicle. An IV catheter was externalized via a port in the plethysmograph allowing for IV

infusion during recording without handling the animal or opening the plethysmograph. Unanesthetized

rats were sealed into the Plexiglas plethysmograph with airflow maintained at 6 L/min for the duration of

the recording. The recording protocol consisted of a 40-minute acclimation period (inspired air: 21% O_2,

79% N_2), followed by a 7-minute ventilatory challenge (10% O_2, 7% CO_2, 83% N_2) and a 10-minute

normoxic recovery period (21% O_2, 79% N_2). Subsequently, rats underwent a 20-minute long, pre-

vehicle, baseline under normoxic conditions (21% O_2, 79% N_2) followed by a 2-minute-long intravenous

infusion of the J60 vehicle (saline; 0.6 mL). Following vehicle infusion recording continued for 30

minutes followed by a 7-minute ventilatory challenge (10% O_2, 7% CO_2, 83% N_2) and 10 minutes of

normoxic breathing (21% O_2, 79% N_2). After an additional 20-minute pre-J60 baseline (21% O_2, 79%

N_2), an intravenous infusion of the J60 ligand was given (0.1 mg/ml dose; 2 minutes long; final volume

standardized to 0.6 mL) and recordings continued for 30-minute followed by a final ventilatory challenge

(10% O_2, 7% CO_2, 83% N_2). The ventilatory challenges were performed to assess the ability to increase

breathing.

Phrenic nerve recordings. Two-to-eight-weeks following plethysmography recordings, bilateral phrenic

nerve recordings were performed. This procedure was done to directly assess the effect of DREADD

activation on phrenic motor output under rigorously controlled experimental conditions. Anesthesia was

induced by placing the rat in a closed chamber to inhale 3% isoflurane in 100% O_2. Rats were then moved

onto a closed-loop heating pad set to maintain rectal temperature at 37 ± 1°C (model 700 TC-1000, CWE

Inc.). Isoflurane anesthesia was maintained using a nose cone. Once a surgical plane of anesthesia was

reached as evidenced by loss of corneal reflexes and hindlimb withdrawal, rats were tracheotomized and

ventilated (VentElite, model 55-7040; Harvard Apparatus Inc.) with a gas mixture of 50% O_2, 1% CO_2,

balanced with N_2. End-tidal CO_2 was maintained at 45-47 mmHg throughout the surgery and

experimental protocol (Capnogard; Novametrix). Ventilator frequency was maintained between 65 and 75

breaths/min, and tidal volume was set at 7 mL/kg[50]. The vagus nerves were transected bilaterally to

prevent entrainment of phrenic efferent output with the ventilator.

A tail vein catheter was placed to allow for intravenous infusion of urethane anesthesia,

supplementary fluids, and the J60 ligand. Rats were slowly converted from inhaled isoflurane to urethane

anesthesia (2.1 g/kg at 6 mL/hr; IV). During this conversion, the depth of anesthesia was consistently

monitored by evaluating the pedal withdrawal reflex. Following administration of the full urethane dose, a

mixture of 8.4% sodium bicarbonate and lactated Ringer's was administered (2 mL/h; IV) to maintain

acid-base balance. To prevent movements and EMG contamination of the phrenic neurogram

pancuronium bromide was administered (3 mg/kg IV, Sigma-Aldrich, St Louis) to achieve neuromuscular

blockade. A catheter (polyethylene tubing; PE 50; Intramedic) was placed in the femoral artery to enable

monitoring of arterial blood pressure via a transducer amplifier (TA-100, CWE) and allow withdrawal of

arterial blood samples (65 μL) for measurement of partial pressure of CO_2 ($PaCO_2$) and O_2 (PaO_2), pH,

and base excess (ABL 90 Flex, Radiometer; Copenhagen, Denmark).

The phrenic nerves were exposed bilaterally using a dorsal approach as described previously[51,52].

Briefly, a midline incision was made at the base of the skull extending to spinal level T2. The muscles

connecting the shoulder blades to the spinal column were separated to expose the phrenic nerves. The

phrenic nerves were isolated, cut distal to the spinal cord, and suctioned into custom-made glass

electrodes filled with 0.9% saline solution. Phrenic nerve activity was amplified (10 kHz) using a

differential AC amplifier (Model 1700, A-M systems, Everett, WA), band-pass filtered (100Hz-3□kHz),

and digitized at 25ks/second (Power 1401, CED).

At the beginning of the experiment, the apneic threshold was determined by slowly reducing the

inspired CO_2 until phrenic nerve inspiratory activity ceased for 60 seconds. The recruitment threshold was

established by slowly increasing the inspired CO_2 until phrenic bursting returned. The end-tidal CO_2

($ETCO_2$) was then maintained 2-3 mmHg above the recruitment threshold for the duration of the

experiment. After achieving a stable phrenic nerve recording and blood gases a 15-minute-long baseline

recording was collected (50% O_2, 3% CO_2) followed by a brief, 5-minute exposure to hypoxia (11.5% O_2,

3% CO_2) and 10–15-minute recovery period (50% O_2, 3% CO_2). Subsequently, intravenous infusion of

vehicle (saline) was given followed by a 15-minute recording period. The J60 ligand (0.1 mg/kg) was

then administered intravenously over a 2-minute infusion period followed by a 100-minute recording

period.

Arterial blood samples were collected at specific intervals: initially at baseline, during the last

minute of each hypoxia episode, 15 minutes post vehicle administration, and subsequently at 20-, 40,- 60-

, 80-, and 100-minutes post J60 administration. Baseline blood gas values served as references to assess if

further arterial samples were isocapnic. To keep end-tidal CO_2 and $PaCO_2$ near baseline (within $\pm$ 2.0

mmHg), minor adjustments to inspired CO_2 and ventilation rate were made as needed. PaO_2 was kept

above 150 mmHg, except during hypoxia; if it dropped below, O_2 intake was increased by 5%, and a new

blood sample was analyzed within 5 minutes.

At the end of the experiment, rats were exposed to a second 5-min episode of hypoxia (11.5% O_2)

followed by a brief "maximal" chemoreceptor challenge induced by switching off the mechanical

ventilator until the animal exhibited a "gasping-like" phrenic discharge pattern (approximately 20-30

seconds). If the increase in phrenic nerve amplitude in response to the "maximal" challenge was lower

than the response observed during either hypoxic episode, it was considered a sign of deteriorating nerve-

electrode contact, and the preparation was excluded from all formal analyses. Rats were then perfused

transcardially with heparinized saline followed by 4% paraformaldehyde and spinal cords were harvested

for histological analysis.

Histology. Spinal cords were harvested and placed in 4% paraformaldehyde for 24 hours. The cords were

subsequently moved to a cryo-protecting solution (30% sucrose in 1x PBS) for a minimum of three days.

Cervical and thoracic spinal cords were blocked in optimal cutting temperature media and cryosectioned

at 20 µm. The viral constructs included a red fluorescent protein (mCherry) fused to the hM3D(Gq)

DREADD which allowed evaluation of DREADD expression by assessing mCherry expression via

fluorescence microscopy.

We performed a qualitative assessment of mCherry expression in the mid-cervical spinal cord.

One intact section from the middle of each spinal segment (C3-C6) was chosen as a representative section

and underwent assessment. Sections were segmented into the following quadrants: left dorsal, right

dorsal, left ventral, and right ventral. The quadrant was scored as "positive" if mCherry positive neurons

or fibers were observed; otherwise, the sub-segment was marked "negative" (see **Figure S4** for example).

The entirety of the grey matter from each section was analyzed for all animals, whether wild-type or

ChAT-Cre. Although ChAT-Cre expression was expected to be limited primarily to motoneurons, which

are the predominant ChAT-positive neuronal subtype in the spinal cord, there is also evidence of ChAT-

positive interneuron populations[53-55] which we also wished to capture in our analysis. Results were

compiled into a summary table showing the total positive counts by animal cohort, spinal segment, and

quadrant (see Results section; **Table 1**). Animals that showed no positive mCherry labeling in the C3-C6

cord were excluded from analysis.

Data analysis. Custom MATLAB (MathWorks; Natick, MA) scripts were created, and are available upon

request. These scripts were used to analyze diaphragm EMG, phrenic nerve, and plethysmography

waveforms. EMG signals were digitally filtered using a second-order, bandpass Butterworth filter (100–

1000 Hz) and then rectified and integrated by taking the absolute value of the signal followed by applying

a moving median filter (50 ms time constant for mice; 75 ms time constant for rats) and moving average

filter (50 ms time constant for mice; 175 ms time constant for rats). The script identified each EMG burst

and calculated peak amplitude, minimum amplitude (tonic activity), and AUC for each burst which was

then averaged across animals and compared across experimental conditions.

Phrenic nerve signals were digitally filtered using a second-order, bandpass Butterworth filter

(100–3 kHz) and then rectified and integrated by taking the absolute value of the signal followed by

applying a moving median filter (50 ms time constant) and moving average filter (50 ms time constant).

The analysis script calculated the peak phrenic burst amplitude and minimum amplitude for each burst

which was then averaged across animals and compared across experimental conditions. Systolic (SP),

diastolic (DP), and mean arterial blood pressure (MAP; formula: MAP = DP + 1/3 (SP - DP)) along with

instantaneous heart rate were calculated from the arterial pressure trace.

In plethysmography experiments, airflow pressure, chamber temperature, chamber humidity,

barometric pressure, and animal body temperature were used to calculate respiratory frequency, tidal

volume, and ventilation via a custom MATLAB script. Tidal volume was calculated using the Drorbaugh

and Fenn equation[56].

Statistical analyses were performed using SigmaPlot 14 (Systat Software) and R (The R Foundation for

Statistical Computing; version 4.3.1). In mouse diaphragm EMG studies, one-way repeated measure

analysis of variance (ANOVA) was used to statistically compare diaphragm EMG peak amplitude, area

under the curve, tonic activity, and heart rate across time before and after J60 application. Paired t-tests

were used to compare left and right hemidiaphragm EMG peak amplitude, area under the curve, tonic

activity, and heart rate between ChAT-Cre and wild-type mice at the 30-minute post-J60 administration

time point. Differences in mortality between wild-type and ChAT-Cre mice post-J60 application were

assessed using Pearson's Chi-squared test with Yates' continuity correction using the chisq.test function in

R. In instances where animals did not survive the entire duration of the anesthetized recording, data up

until the time point preceding their death was included. In control EMG experiments, one-way RM

ANOVA was used to compare EMG peak responses across baseline, sham injection, and J60

administration. These data were also assessed normalized to baseline, in which case EMG peak responses

after sham injection and J60 application were compared using paired t-tests. In plethysmography

experiments, two-way repeated measures ANOVA was used to compare raw and normalized tidal

volume, respiratory frequency, and minute ventilation across time and treatment (saline vs J60). Paired t-

tests were used to compare responses to hypercapnic-hypoxic ventilatory challenges across treatments.

One-way RM ANOVA was used to compare phrenic peak amplitude, systolic and diastolic blood

pressures, mean arterial blood pressure, and respiratory rate across time for phrenic nerve recordings. The

relationship between time post-AAV injection and average phrenic response to J60 was assessed for

ChAT-Cre rats using the cor.test function in R to run a Pearson's product moment correlation. Both male

and female animals were included in this study to improve the generalizability of the results. However,

we were not adequately powered for sex comparisons and therefore did not perform any statistical

analysis to assess sex differences.

In cases of significant main effects, the Tukey post-hoc test was used to assess differences

between individual time points. For instances where data failed to meet general linear model assumptions

(i.e., normality, homogeneity of variances), nonparametric equivalents of the previously mentioned

statistical tests were used. Data were considered statistically significant when $p \leq 0.05$. The mean data are

presented along with the standard error of the mean.

Figure 1. *DREADD activation increases diaphragm EMG output in wild-type mice.* A representative example of diaphragm EMG activity before and after application of the J60 DREADD ligand is shown in the top panel. Examples of the individual inspiratory EMG bursts at baseline (a), after vehicle (b), and after J60 (c) are shown. The J60 ligand increased diaphragm output but did not impact respiratory rate. The mean responses (n = 11; n = 7 females) for EMG AUC, peak-to-peak amplitude, tonic activity, and respiratory rate are shown in panels d-g. For diaphragm EMG data (panels d-f) left hemidiaphragm EMG is represented in orange, while right hemidiaphragm EMG is blue. Error bars depict ±□1 SEM. Statistical reports for all panels are provided in Supplemental Table 1. * and ‡ symbols indicate significant main effects (p < 0.05) on One-Way RM ANOVA for the left and right hemidiaphragm, respectively. Dia = diaphragm, AUC = area under the curve, amp = peak amplitude, BL = baseline, SL = saline (sham injection).

Figure 2. *DREADD activation increases diaphragm EMG output in ChAT-Cre mice.* A representative example of diaphragm EMG activity before and after application of the J60 DREADD ligand is shown in the top panel. Examples of the individual inspiratory EMG bursts at baseline (a), after vehicle (b), and after J60 (c) are shown. Mean responses (n = 9; n = 6 females) for EMG AUC, peak-to-peak amplitude, tonic activity, and respiratory rate are shown in panels d-g. The DREADD ligand caused a bilateral increase in diaphragm EMG AUC, peak-to-peak amplitude, and tonic activity. For all EMG parameters, the responses were greater on the right vs. left hemidiaphragm. Respiratory rate decreased over time. For panels d-f, the left hemidiaphragm EMG is represented in orange, while right hemidiaphragm EMG is blue. Error bars depict ±□1 SEM. Statistical reports for all panels are provided in Supplemental Table 2.

* and ‡ symbols indicate significant main effects (p < 0.05) on One-Way RM ANOVA for the left and right hemidiaphragm, respectively. # indicates a significant main effect (p < 0.05) on One-Way RM ANOVA for respiratory rate data. Dia = diaphragm, AUC = area under the curve, amp = peak amplitude, BL = baseline, SL = saline (sham injection).

Figure 3. *Wild type vs. ChAT-Cre mouse responses to DREADD activation.* Direct comparisons of diaphragm EMG response parameters (a-f) and respiratory rate (g) at 30-minute post-J60 application (Wild type, n = 11; n = 7 females; ChAT-Cre, n = 9; n = 6 females). Left hemidiaphragm EMG AUC (a), peak-to-peak amplitude (b), and tonic activity (c) were similar between groups. However, the same parameters on the right hemidiaphragm (d-f) were greater in ChAT-Cre mice. Respiratory rate was similar between groups. Error bars depict ±□1 SEM. Statistical reports for all panels are provided in Supplemental Table 3. * p < 0.05. AUC = area under the curve, amp = peak EMG amplitude, Dia = diaphragm, BL = baseline, resp rate = respiratory rate.

Figure 4. *DREADD activation increases ventilation in unanesthetized ChAT-Cre rats.* Summary plots (n = 9; n = 3 females) showing the impact of the J60 DREADD ligand on tidal volume, respiratory rate, and minute ventilation are shown in panels a-c. The normalized values (% of baseline) are shown in panels d-f. The DREADD ligand increased tidal volume compared to sham infusion (saline). Error bars depict ±☐1 SEM. Statistical reports for all panels are provided in Supplemental Table 4. BL = baseline, IV = intravenous infusion period.

Figure 5. *DREADD activation increases phrenic nerve output in ChAT-Cre rats.* Representative data showing that the J60 DREADD ligand causes a rapid increase in phrenic nerve output (a). Mean data (n = 9; n = 3 females) showing the impact of J60 application on phrenic nerve raw (b) and normalized (c) peak-to-peak amplitude, raw (d) and normalized (e) tonic activity, systolic blood pressure (f), diastolic blood pressure (g), heart rate (h), mean arterial blood pressure (i), and respiratory rate (j). The J60 ligand caused an increase in phrenic peak-to-peak amplitude and tonic activity. Systolic, diastolic, and mean arterial blood pressure all decreased after J60 application. Heart rate and respiratory rate were not statistically different after J60. In panels b-e, the left phrenic is represented in orange, while right phrenic is blue. Error bars depict ±□1 SEM. Statistical reports for all panels are provided in Supplemental Table 6. * and ‡ symbols indicate significant main effects (p < 0.05) on One-Way RM ANOVA for the left and right hemidiaphragm, respectively. # indicates a significant (p < 0.05) effect on One-Way RM ANOVA for respiratory rate data. Phr = phrenic, amp = amplitude, BL = baseline, SP = systolic pressure, DP = diastolic pressure, HR = heart rate, MAP = mean arterial pressure.

Figure 6. *Histological assessment of mCherry expression in the C4/C5 spinal segments.* Representative photomicrographs of mid-cervical spinal sections from a wild-type mouse (a-a$_i$), a ChAT-Cre mouse (b-b$_i$), and a ChAT-Cre rat (c-c$_i$). Wild-type mice (a-a$_i$) showed a nonspecific pattern of expression throughout the mid-cervical grey matter. ChAT-Cre mice and rats (b-c$_i$) showed expression limited to neurons in the ventral horns. Red color indicates positive and mCherry fluorescence. Dashed white line indicates the approximate white-gray matter demarcation.

	Wildtype mice (n= 11)				ChAT-Cre mice (n= 9)				ChAT-Cre rats (n= 9)			
	Dorsal		Ventral		Dorsal		Ventral		Dorsal		Ventral	
	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right
C3	4	5	6	6	0	1	3	4	0	1	3	4
C4	7	9	10	9	1	2	5	9	0	1	6	8
C5	11	11	11	11	0	1	6	9	1	1	5	9
C6	10	10	9	9	2	2	5	8	0	0	1	2

Table 1. *Qualitative assessment of mCherry expression in the mid-cervical spinal cord.* Spinal segments C3-C6 were assessed in quadrants broken into dorsal, ventral, left, and right. Spinal segments were counted as "positive" if they showed any evidence of mCherry expression in neuronal soma or fibers. The counts therefore indicate the number of animals of a given cohort that were mCherry positive for a given spinal segment quadrant. All animals showed a slight trend for more mCherry expression moving rostral to caudal and for more expression in the ventral vs the dorsal lamina. This trend was more prominent in the ChAT-Cre animals. At the bottom of the table, a heatmap is provided for easier assessment of the distribution of positive mCherry counts across quadrants and spinal segments.

REFERENCES

1   Berlowitz, D. J., Wadsworth, B. & Ross, J. Respiratory problems and management in people with
spinal cord injury. *Breathe (Sheff)* **12**, 328-340 (2016). https://doi.org/10.1183/20734735.012616
2   Brown, R., DiMarco, A. F., Hoit, J. D. & Garshick, E. Respiratory dysfunction and management
in spinal cord injury. *Respir Care* **51**, 853-868;discussion 869-870 (2006).
3   Perrin, C., Unterborn, J. N., Ambrosio, C. D. & Hill, N. S. Pulmonary complications of chronic
neuromuscular diseases and their management. *Muscle Nerve* **29**, 5-27 (2004).
https://doi.org/10.1002/mus.10487
4   Mehta, S. Neuromuscular disease causing acute respiratory failure. *Respir Care* **51**, 1016-1021;
discussion 1021-1013 (2006).
5   Burakgazi, A. Z. & Höke, A. Respiratory muscle weakness in peripheral neuropathies. *J Peripher*
Nerv Syst **15**, 307-313 (2010). https://doi.org/10.1111/j.1529-8027.2010.00293.x
6   Fuller, D. D. *et al.* The respiratory neuromuscular system in Pompe disease. *Respir Physiol*
Neurobiol **189**, 241-249 (2013). https://doi.org/10.1016/j.resp.2013.06.007
7   Jordan, A. S. & White, D. P. Pharyngeal motor control and the pathogenesis of obstructive sleep
apnea. *Respir Physiol Neurobiol* **160**, 1-7 (2008). https://doi.org/10.1016/j.resp.2007.07.009
8   Doyle, B. M. *et al.* Gene delivery to the hypoglossal motor system: preclinical studies and
translational potential. *Gene Ther* (2021). https://doi.org/10.1038/s41434-021-00225-1
9   Zhu, H. & Roth, B. L. DREADD: a chemogenetic GPCR signaling platform. *Int J*
Neuropsychopharmacol **18** (2014). https://doi.org/10.1093/ijnp/pyu007
10   Alexander, G. M. *et al.* Remote control of neuronal activity in transgenic mice expressing
evolved G protein-coupled receptors. *Neuron* **63**, 27-39 (2009).
https://doi.org/10.1016/j.neuron.2009.06.014
11   Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the lock to fit the
key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc*
Natl Acad Sci U S A **104**, 5163-5168 (2007). https://doi.org/10.1073/pnas.0700293104
12   Fleury Curado, T. *et al.* Chemogenetic stimulation of the hypoglossal neurons improves upper
airway patency. *Scientific reports* **7**, 44392 (2017). https://doi.org/10.1038/srep44392
13   Fleury Curado, T. A. *et al.* Silencing of Hypoglossal Motoneurons Leads to Sleep Disordered
Breathing in Lean Mice. *Front Neurol* **9**, 962 (2018). https://doi.org/10.3389/fneur.2018.00962
14   Fleury Curado, T. *et al.* Designer Receptors Exclusively Activated by Designer Drugs Approach
to Treatment of Sleep-disordered Breathing. *American journal of respiratory and critical care*
medicine **203**, 102-110 (2021). https://doi.org/10.1164/rccm.202002-0321OC
15   Singer, M. L. *et al.* Chemogenetic activation of hypoglossal motoneurons in a mouse model of
Pompe disease. *Journal of neurophysiology* **128**, 1133-1142 (2022).
https://doi.org/10.1152/jn.00026.2022
16   Horton, G. A. *et al.* Activation of the Hypoglossal to Tongue Musculature Motor Pathway by
Remote Control. *Sci Rep* **7**, 45860 (2017). https://doi.org/10.1038/srep45860
17   Fuller, D. D., Rana, S., Smuder, A. J. & Dale, E. A. The phrenic neuromuscular system.
Handbook of clinical neurology **188**, 393-408 (2022). https://doi.org/10.1016/B978-0-323-91534-
2.00012-6
18   DiMarco, A. F. & Kowalski, K. E. Activation of inspiratory muscles via spinal cord stimulation.
Respiratory physiology & neurobiology **189**, 438-449 (2013).
https://doi.org/10.1016/j.resp.2013.06.001
19   Jensen, V. N., Seedle, K., Turner, S. M., Lorenz, J. N. & Crone, S. A. V2a Neurons Constrain
Extradiaphragmatic Respiratory Muscle Activity at Rest. *eNeuro* **6** (2019).
https://doi.org/10.1523/ENEURO.0492-18.2019
20   Alilain, W. J. *et al.* Light-induced rescue of breathing after spinal cord injury. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **28**, 11862-11870 (2008).
https://doi.org/10.1523/JNEUROSCI.3378-08.2008

Satkunendrarajah, K., Karadimas, S. K., Laliberte, A. M., Montandon, G. & Fehlings, M. G. Cervical excitatory neurons sustain breathing after spinal cord injury. *Nature* **562**, 419-422 (2018). https://doi.org/10.1038/s41586-018-0595-z

Jensen, V. N., Alilain, W. J. & Crone, S. A. Role of Propriospinal Neurons in Control of Respiratory Muscles and Recovery of Breathing Following Injury. *Front Syst Neurosci* **13**, 84 (2019). https://doi.org/10.3389/fnsys.2019.00084

Lane, M. A. Spinal respiratory motoneurons and interneurons. *Respiratory physiology & neurobiology* **179**, 3-13 (2011). https://doi.org/10.1016/j.resp.2011.07.004

Li, K. *et al.* Overexpression of the astrocyte glutamate transporter GLT1 exacerbates phrenic motor neuron degeneration, diaphragm compromise, and forelimb motor dysfunction following cervical contusion spinal cord injury. *J Neurosci* **34**, 7622-7638 (2014). https://doi.org/10.1523/jneurosci.4690-13.2014

Li, K. *et al.* GLT1 overexpression in SOD1(G93A) mouse cervical spinal cord does not preserve diaphragm function or extend disease. *Neurobiology of disease* **78**, 12-23 (2015). https://doi.org/10.1016/j.nbd.2015.03.010

Qiu, K., Falk, D. J., Reier, P. J., Byrne, B. J. & Fuller, D. D. Spinal delivery of AAV vector restores enzyme activity and increases ventilation in Pompe mice. *Molecular therapy : the journal of the American Society of Gene Therapy* **20**, 21-27 (2012). https://doi.org/10.1038/mt.2011.214

Thakre, P. P., Rana, S., Benevides, E. S. & Fuller, D. D. Targeting drug or gene delivery to the phrenic motoneuron pool. *Journal of neurophysiology* **129**, 144-158 (2023). https://doi.org/10.1152/jn.00432.2022

Mantilla, C. B., Zhan, W. Z. & Sieck, G. C. Retrograde labeling of phrenic motoneurons by intrapleural injection. *J Neurosci Methods* **182**, 244-249 (2009). https://doi.org/10.1016/j.jneumeth.2009.06.016

Gransee, H. M., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Targeted delivery of TrkB receptor to phrenic motoneurons enhances functional recovery of rhythmic phrenic activity after cervical spinal hemisection. *PLoS One* **8**, e64755 (2013). https://doi.org/10.1371/journal.pone.0064755

Martínez-Gálvez, G. *et al.* TrkB gene therapy by adeno-associated virus enhances recovery after cervical spinal cord injury. *Exp Neurol* **276**, 31-40 (2016). https://doi.org/10.1016/j.expneurol.2015.11.007

Gransee, H. M., Gonzalez Porras, M. A., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Motoneuron glutamatergic receptor expression following recovery from cervical spinal hemisection. *J Comp Neurol* **525**, 1192-1205 (2017). https://doi.org/10.1002/cne.24125

Keeler, A. M. *et al.* Intralingual and Intrapleural AAV Gene Therapy Prolongs Survival in a SOD1 ALS Mouse Model. *Mol Ther Methods Clin Dev* **17**, 246-257 (2020). https://doi.org/10.1016/j.omtm.2019.12.007

Tervo, D. G. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. *Neuron* **92**, 372-382 (2016). https://doi.org/10.1016/j.neuron.2016.09.021

Smith, K. S., Bucci, D. J., Luikart, B. W. & Mahler, S. V. Dreadds: Use and application in behavioral neuroscience. *Behav Neurosci* **135**, 89-107 (2021). https://doi.org/10.1037/bne0000433

Roth, B. L. DREADDs for Neuroscientists. *Neuron* **89**, 683-694 (2016). https://doi.org/10.1016/j.neuron.2016.01.040

Ouali Alami, N. *et al.* Multiplexed chemogenetics in astrocytes and motoneurons restore blood-spinal cord barrier in ALS. *Life Sci Alliance* **3** (2020). https://doi.org/10.26508/lsa.201900571

Saxena, S. *et al.* Neuroprotection through excitability and mTOR required in ALS motoneurons to delay disease and extend survival. *Neuron* **80**, 80-96 (2013). https://doi.org/10.1016/j.neuron.2013.07.027

Jaiswal, P. B., Mistretta, O. C., Ward, P. J. & English, A. W. Chemogenetic Enhancement of Axon Regeneration Following Peripheral Nerve Injury in the SLICK-A Mouse. *Brain Sci* **8** (2018). https://doi.org/10.3390/brainsci8050093

Jaiswal, P. B. & English, A. W. Chemogenetic enhancement of functional recovery after a sciatic nerve injury. *Eur J Neurosci* **45**, 1252-1257 (2017). https://doi.org/10.1111/ejn.13550

Streeter, K. A. *et al.* Mid-cervical interneuron networks following high cervical spinal cord injury. *Respir Physiol Neurobiol* **271**, 103305 (2020). https://doi.org/10.1016/j.resp.2019.103305

Lane, M. A. *et al.* Cervical prephrenic interneurons in the normal and lesioned spinal cord of the adult rat. *The Journal of comparative neurology* **511**, 692-709 (2008). https://doi.org/10.1002/cne.21864

Molkov, Y. I., Rubin, J. E., Rybak, I. A. & Smith, J. C. Computational models of the neural control of breathing. *Wiley Interdiscip Rev Syst Biol Med* **9** (2017). https://doi.org/10.1002/wsbm.1371

Dempsey, J. A. & Welch, J. F. Control of Breathing. *Semin Respir Crit Care Med* **44**, 627-649 (2023). https://doi.org/10.1055/s-0043-1770342

McGuire, M., Zhang, Y., White, D. P. & Ling, L. Phrenic long-term facilitation requires NMDA receptors in the phrenic motonucleus in rats. *J Physiol* **567**, 599-611 (2005). https://doi.org/10.1113/jphysiol.2005.087650

Rana, S., Zhan, W. Z., Sieck, G. C. & Mantilla, C. B. Cervical spinal hemisection alters phrenic motor neuron glutamatergic mRNA receptor expression. *Experimental neurology* **353**, 114030 (2022). https://doi.org/10.1016/j.expneurol.2022.114030

Qiu, K., Lane, M. A., Lee, K. Z., Reier, P. J. & Fuller, D. D. The phrenic motor nucleus in the adult mouse. *Experimental neurology* **226**, 254-258 (2010). https://doi.org/10.1016/j.expneurol.2010.08.026

Rana, S., Sieck, G. C. & Mantilla, C. B. Heterogeneous glutamatergic receptor mRNA expression across phrenic motor neurons in rats. *Journal of neurochemistry* **153**, 586-598 (2020). https://doi.org/10.1111/jnc.14881

Rana, S., Mantilla, C. B. & Sieck, G. C. Glutamatergic input varies with phrenic motor neuron size. *J Neurophysiol* **122**, 1518-1529 (2019). https://doi.org/10.1152/jn.00430.2019

Zolotukhin, S. *et al.* Production and purification of serotype 1, 2, and 5 recombinant adeno-associated viral vectors. *Methods* **28**, 158-167 (2002). https://doi.org/10.1016/s1046-2023(02)00220-7

Lee, K. Z., Sandhu, M. S., Dougherty, B. J., Reier, P. J. & Fuller, D. D. Hypoxia triggers short term potentiation of phrenic motoneuron discharge after chronic cervical spinal cord injury. *Exp Neurol* **263**, 314-324 (2015). https://doi.org/10.1016/j.expneurol.2014.10.002

Thakre, P. P. & Fuller, D. D. Pattern sensitivity of ampakine-hypoxia interactions for evoking phrenic motor facilitation in anesthetized rat. *J Neurophysiol* **131**, 216-224 (2024). https://doi.org/10.1152/jn.00315.2023

Thakre, P. P., Sunshine, M. D. & Fuller, D. D. Ampakine pretreatment enables a single hypoxic episode to produce phrenic motor facilitation with no added benefit of additional episodes. *J Neurophysiol* **126**, 1420-1429 (2021). https://doi.org/10.1152/jn.00307.2021

Gotts, J., Atkinson, L., Yanagawa, Y., Deuchars, J. & Deuchars, S. A. Co-expression of GAD67 and choline acetyltransferase in neurons in the mouse spinal cord: A focus on lamina X. *Brain Res* **1646**, 570-579 (2016). https://doi.org/10.1016/j.brainres.2016.07.001

Alkaslasi, M. R. *et al.* Single nucleus RNA-sequencing defines unexpected diversity of cholinergic neuron types in the adult mouse spinal cord. *Nat Commun* **12**, 2471 (2021). https://doi.org/10.1038/s41467-021-22691-2

Mesnage, B. *et al.* Morphological and functional characterization of cholinergic interneurons in the dorsal horn of the mouse spinal cord. *J Comp Neurol* **519**, 3139-3158 (2011). https://doi.org/10.1002/cne.22668

Drorbaug, J. E. & Fenn, W. O. A BAROMETRIC METHOD FOR MEASURING
      VENTILATION IN NEWBORN INFANTS. *Pediatrics* **16**, 81-87 (1955).
      https://doi.org/10.1542/peds.16.1.81