1 **A stress-dependent TDP-43 SUMOylation program preserves neuronal function**

3 **Authors:** Terry R. Suk[1-4], Caroline E. Part[1-4], Trina T. Nguyen[1-4], Jenny L. Zhang[1-4], Meghan M.
4 Heer[1-4], Alejandro Caballero-Gómez[1-4], Veronica S. Grybas[1-4], Paul M. McKeever[5], Benjamin
5 Nguyen[1-4], Steve M. Callaghan[1-4], John M. Woulfe[1,2,6-8], Janice Robertson[5],
6   & Maxime W.C. Rousseaux[1-4]*.

[1]University of Ottawa Brain and Mind Research Institute, ON, Canada

[2]Department of Cellular and Molecular Medicine, University of Ottawa, ON, Canada

[3]Eric Poulin Center for Neuromuscular Diseases, ON, Canada

[4]Ottawa Institute of Systems Biology, ON, Canada

[5]Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, ON, Canada

[6]The Ottawa Hospital Research Institute, The Ottawa Hospital, ON, Canada

[7]Department of Pathology and Laboratory Medicine, University of Ottawa, ON, Canada

[8]Department of Biochemistry, Microbiology, and Immunology, University of Ottawa, ON,
Canada

*Corresponding Author. Email: max.rousseaux@uottawa.ca

Abstract: Amyotrophic Lateral Sclerosis (ALS) and Frontotemporal Dementia (FTD) are
overwhelmingly linked to TDP-43 dysfunction. Mutations in TDP-43 are rare, indicating that
exogenous factors – such as cellular stressors – converge on TDP-43 to play a key role in disease
pathogenesis. Post translational modifications such as SUMOylation play essential roles in
response to such exogenous stressors. We therefore set out to understand how SUMOylation may
regulate TDP-43 in health and disease. We find that TDP-43 is regulated dynamically via
SUMOylation in response to cellular stressors. When this process is blocked *in vivo*, we note
age-dependent TDP-43 pathology and sex-specific behavioral deficits linking TDP-43
SUMOylation with aging and disease. Collectively, this work presents TDP-43 SUMOylation as
an early physiological response to cellular stress, disruption of which may confer a risk for TDP-
43 proteinopathy.

Introduction

TDP-43 is an essential RNA binding protein notorious for its involvement in neurodegenerative diseases. Most notably in Amyotrophic Lateral Sclerosis (ALS) and Frontotemporal Dementia (FTD), TDP-43 is pathologically found mislocalized from the nucleus and aggregated in the cytoplasm of neurons within the central nervous system (CNS) in ~97% and ~45% of disease cases, respectively(*1*, *2*). Furthermore, TDP-43 pathology is increasingly linked to other neurological disorders including Limbic-predominant Age-related TDP-43 Encephalopathy (LATE), Alzheimer's Disease (AD), Chronic Traumatic Encephalopathy (CTE), and Stroke(*3–6*). It is becoming evident that not one, but many mechanisms, including ALS/FTD linked mutations in the gene encoding TDP-43 (*TARDBP*), genetic variants affecting related pathways, and/or environmental factors converge on TDP-43 to cause pathology(*7–12*). Therefore, it is critical to identify molecular pathways that regulate TDP-43. This will yield a better understanding of how their dysfunction contributes to early disease pathogenesis and help uncover novel biomarkers and therapeutic targets directed towards TDP-43.

The formation of cytoplasmic TDP-43 aggregates is increasingly considered a late-stage event in ALS, FTD, and related proteinopathies (herein referred to as ALS/FTD)(*7*). Increasing evidence suggests that the partial mislocalization or complete depletion of TDP-43 from the nucleus to the cytoplasm is an early event in ALS/FTD pathogenesis(*7, 13–18*). As a result, more efforts have focused on the outcomes of TDP-43 mislocalization as early markers of dysfunction and/or mechanisms driving disease. Indeed, loss of function due to mislocalization of TDP-43 causes aberrant cryptic splicing in genes including *STMN2* and *UNC13A* that can drive the progression of ALS(*14–17, 19*). Cryptic mis-splicing of *STMN2* was recently found to show better correlation with disease progression compared to classical TDP-43 cytoplasmic aggregates further supporting the need to understand early mechanisms of TDP-43 dysfunction(*20, 21*).

The earliest events in TDP-43 pathogenesis, upstream of mislocalization, remain convoluted likely due to the coincidence of pathological TDP-43 mislocalization with its normal physiological roles in response to stress(*7*). Thus, mechanisms that may tip this physiological balance towards a pathogenic state likely underlie susceptibility to neurodegeneration. The cellular stress response is a critical pathway tightly linked to ALS/FTD and TDP-43 supported via genetic, experimental, and epidemiological evidence(*22–35*). On one hand, ALS/FTD-linked mutations disrupting various steps of the stress response pathways converge on the dysregulation of TDP-43 resulting in pathology(*22–24, 29–32, 35*). On the other hand, exogenous insults such as head injuries, viral infections, and other exposures may confer susceptibility or precipitate ALS/FTD and can serve as pre-clinical models of TDP-43 proteinopathy(*5, 10, 11, 36*). Indeed, prolonged cellular stress due to chronic stress exposure or failures in stress recovery can result in TDP-43 pathology(*27, 37*). However, much less is known about the roles and direct mechanisms regulating TDP-43 in the cell stress response and how they might be linked to neurodegeneration.

Post translational modifications (PTMs) play key roles in regulating protein function and have been tightly linked to TDP-43 (dys)function and disease. TDP-43 is modified by an array of PTMs including phosphorylation, ubiquitination, acetylation, and polyADP-ribosylation(*38–42*). Abnormal TDP-43 phosphorylation and ubiquitination are pathognomonic of ALS and TDP-43 related FTD(*38*). Recent studies have begun to suggest a role for SUMOylation – the covalent addition of small ubiquitin like modifiers (SUMOs) to lysine residues – in ALS and TDP-43 function, highlighting roles in regulating stress granule disassembly, and TDP-43 splicing and

aggregation(*43–47*). SUMOylation is a highly conserved and essential mechanism whereby
SUMO proteins are covalently bound to target substrate through a cascade mirroring
ubiquitination(*48, 49*). SUMOylation plays a wide variety of roles in the cell including
regulation of cellular stress responses, nucleocytoplasmic transport, and protein solubility(*50*).
As these functions strongly overlap with TDP-43 pathobiology and stress response, we posited
that SUMOylation has a direct role in regulating TDP-43-mediated stress responses.

TDP-43 is SUMOylated in the nucleus in a context specific manner

Various acute stressors including oxidative (sodium arsenite, $NaAsO_2$), hyperosmotic
(D-sorbitol and sodium chloride), and heat shock have been demonstrated to induce TDP-43
mislocalization and aggregation *in vitro*, helping to uncover key mechanisms affected in
disease(*23, 51, 52*). As sodium arsenite is a commonly used stressor to interrogate TDP-43
alterations, we examined whether TDP-43 became SUMOylated in response to sodium arsenite
treatment. Using an immunoprecipitation assay ("SUMOylation Assay", Fig. 1A) to
immunoprecipitate HA-SUMO bound proteins under denaturing conditions to disrupt non-
covalent protein interactions, we found that TDP-43 becomes SUMOylated by SUMO2 and can
form polySUMO chains specifically in response to sodium arsenite (Fig. 1B). Interestingly we
found that sodium arsenite and heat shock stressors, but not hyperosmotic stress, induced
TDP-43 SUMOylation suggesting that SUMOylation is a context specific modifier of TDP-43
(fig. S1A). We further validated that TDP-43 is SUMOylated by SUMO2 and SUMO3, but not
SUMO1, consistent with the roles of SUMO2/3 in response to stress (fig. S1B). Mature
(covalently bound) SUMO2 and SUMO3 share nearly 100% amino acid homology making them
difficult to differentiate and thus referred together as SUMO2/3. Since SUMO2 is the only
essential and most abundantly expressed SUMO paralog in the CNS and sodium arsenite is
commonly used to study TDP-43 pathobiology, we focused on TDP-43 SUMOylation by
SUMO2 in response to sodium arsenite stress(*53, 54*).

TDP-43 and SUMO2 are widely expressed throughout the nucleus however interactions
only occur in response to stress, making co-localization analysis challenging to infer
SUMOylation events (fig. S1C). To visualize the interaction between TDP-43 and SUMO2/3 we
performed a proximity ligation assay (PLA) in murine primary cortical neurons cultures
(Fig. 1C, S1D). We observed that in response to sodium arsenite there is a significant increase in
PLA signal between endogenous TDP-43 and SUMO2/3, consistent with an increased TDP-43
SUMOylation in response to stress. To connect TDP-43 SUMOylation to disease-like states, we
expressed ALS-causing TDP-43^{Q331K} and disrupted the nuclear localization sequence in TDP-43
(TDP-43^{NLSmut}) to mislocalize TDP-43 to the cytoplasm and found that in response to sodium
arsenite TDP-43^{Q331K} leads to increased TDP-43 SUMOylation, while TDP-43^{NLSmut} blocks
SUMOylation indicating that nuclear localization is essential for TDP-43 SUMOylation
(Fig. 1D). The latter finding is supported by the nuclear PLA signal indicating TDP-43 and
SUMO2/3 interact in the nucleus (Fig. 1C). Together, this demonstrates that TDP-43
SUMOylation occurs in the nucleus in a stress-specific manner, suggesting that this modification
may be an important early mechanism upstream of proteinopathy.

SUMOylation is an early event in response to stress and helps clear TDP-43 through the
ubiquitin proteasome system during recovery

Prolonged cellular stress *in vitro* leads to mislocalization of TDP-43 and progressive
transition into insoluble aggregates resembling aspects of pathology observed in disease(*18, 27,*

30, *32*, *55*). As TDP-43 SUMOylation occurs within the nucleus upstream of nuclear egress, we
postulated that SUMOylation would occur upstream of aggregation during prolonged sodium
arsenite stress. Indeed, we observed that TDP-43 becomes readily SUMOylated in the first 15-30
minutes of the acute stress response upstream of the accumulation of RIPA-insoluble,
phosphorylated TDP-43 (fig. S1E). Furthermore, we observed an increase in TDP-43
SUMOylation that correlated with the duration of stress. To determine whether SUMOylation
occurs in response to relative stress intensity, we performed a dose response assay and found that
TDP-43 SUMOylation is proportional to relative stress intensity (fig. S1F). Therefore, the
increase in TDP-43 SUMOylation during prolonged stress is likely due to an increased stress
burden over time.

To determine the fate of SUMOylated TDP-43, we performed a stress-recovery time-
course assay where cells were stressed with sodium arsenite for up to 1 hour, then allowed cells
to recover. For the first hour during stress recovery, we observed a significant peak in TDP-43
SUMOylation indicating that TDP-43 is increasingly SUMOylated and polySUMOylated during
stress recovery (Fig. 1E). SUMOylated TDP-43 was not detectable after 3 hours of recovery,
suggesting that most of the clearance occurs between hours 2-3 post stress. TDP-43 is known to
interact with the SUMO-Targeted Ubiquitin Ligase RNF4 which can polyUbiquitinate
SUMOylated proteins to be degraded through the UPS(*56*). To determine whether the UPS
functions to clear SUMOylated TDP-43 during stress recovery, we treated cells with the
proteasome inhibitor MG132 and monitored TDP-43 SUMOylation during stress and recovery.
We observed that SUMOylated TDP-43 was not cleared during post-stress recovery with the
addition of MG132 indicating that SUMOylated TDP-43 is marked for degradation by the UPS
pathway. Importantly, treatment with MG132 for 1 or 4 hours was not sufficient to induce
TDP-43 SUMOylation by itself, consistent with the stress-selective nature of TDP-43
SUMOylation. Taken together, TDP-43 SUMOylation occurs early in the stress response which
then leads to ubiquitin-targeted clearance during stress recovery demonstrating SUMOylation
may be an important mechanism regulating nuclear TDP-43 proteostasis.

Stress-dependent TDP-43 SUMOylation is mediated by select E3 SUMO ligases
E3 SUMO ligases can be essential for SUMOylation of select substrates and enable
spatiotemporal regulation. Thus, they may represent critical modulators of TDP-43
SUMOylation. We used a dual sgRNA/Cas9 approach to knockout 15 candidate E3 SUMO
ligases to screen for loss of TDP-43 SUMOylation (Fig. 1F). As a positive control, knockout of
the sole E2 SUMO ligase, UBC9 (encoded by *UBE2I*), led to a near complete loss of TDP-43
SUMOylation demonstrating the functionality of the approach. We identified four SUMO E3
ligases whose knockout consistently reduced the levels of stress-induced TDP-43 SUMOylation:
EGR2 (KROX20), PIAS1, TRIM28, and *ZNF451*. EGR2 is linked to the neuromuscular disorder
Charcot Marie Tooth Disease and is an immediate early gene rapidly reacting to external cellular
stimuli aligning with the early response of TDP-43 SUMOylation upon stress (*57*, *58*). PIAS1 is
a canonical, highly conserved E3 SUMO ligase known to play roles in cellular stress
responses(*59*). TRIM28 is also a well characterized E3 SUMO ligase which is a predicted
interactor of SUMOylated TDP-43 based on the GPS-SUMO2.0 algorithm(*60–62*). ZNF451 is
characterized as a stress dependent E3 SUMO ligase specifically promoting polySUMOylation
with SUMO2/3. It is sometimes referred to as an "E4 SUMO elongase" which complements our
data that TDP-43 is polySUMOylated in response to stress(*63*, *64*). Identification of these
putative TDP-43 SUMO ligases supports the robustness of stress-induced TDP-43 SUMOylation

and adds layers of nuance into the regulation of TDP-43 with variable routes of modulating
TDP-43 functions.

TDP-43 SUMOylation occurs in a conserved region of the C-terminal domain

SUMOylation occurs at lysine (K) residues often residing within a consensus
SUMOylation motif, Ψ-K-x-D/E (Ψ=large hydrophobic residue, x = any amino acid), however it
is increasingly recognized that SUMOylation at lysine residues in non-consensus motifs is not
uncommon and plays significant roles in regulating protein function[61]. Using GPS-SUMO
(https://sumo.biocuckoo.cn/) to predict likely SUMOylation sites on TDP-43, we identified a
consensus SUMOylation motif within the first RNA recognition motif at K136, and a
non-consensus SUMOylation motif within the C-terminal domain at K408[61, 62]. We
expressed mutant forms of TDP-43-GFP blocking the predicted sites by mutating the lysine
residue to similarly charged arginine residues: K136R, K408R, and double mutant 2KR. We
observed that TDP-43[K136R] did not block sodium arsenite-induced TDP-43 SUMOylation,
however the TDP-43[K408R] mutant significantly reduced TDP-43 SUMOylation, suggesting K408
is the primary site of SUMOylation in response to sodium arsenite stress (Fig. 2A). Importantly,
mining two independent unbiased mass spectrometry studies identifying sites of SUMOylation
across the proteome also revealed TDP-43 SUMOylation at K408, supporting our findings (65,
66).

To address the evolutionary conservation of TDP-43 SUMOylation, we aligned the
amino acid sequences of TDP-43 with 300 orthologs from OrthoDB and found that human K408
is nearly completely conserved throughout jawed-vertebrates concurrent with the evolution of
the C-terminal domain (Fig. 2B, table S1). While this domain harbors the majority of ALS/FTD
causing mutations, the residues surrounding K408 display considerably high levels of
conservation (~86% pairwise identity) indicating that this motif may play important roles in
regulating TDP-43 in vertebrates that may be altered by disease causing mutations (Fig. 2C, 1D,
table S1). Consistent with these findings, no missense variants have been identified that disrupt
the SUMOylation motif (based on gnomAD v4.0, [67]) supporting its robust conservation in the
healthy human population. Furthermore, the residues surrounding TDP-43 K408 are enriched for
PTM sites, specifically phosphorylation at S403, S404, S409, and S410 which are often
phosphorylated in patients with TDP-43 proteinopathy, potentially linking TDP-43
SUMOylation with phosphorylation and disease pathogenesis[38]. To determine whether
phosphorylation of TDP-43 at these serine residues interact with SUMOylation at K408, we
expressed phospho-dead "4SA" TDP-43 (S403/404/409/410A) and observed that TDP-43[4SA]
exhibits increased SUMOylation following stress (Fig. 2D). Whether phosphorylation functions
allosterically to remove SUMO2/3 at K408 or antagonistically to block the SUMOylation
recognition site remains unclear. The independent modification by either phosphorylation or
SUMOylation may serve as a mechanism to differentially regulate TDP-43 function
independently or sequentially in response to stimuli. Thus, while TDP-43 phosphorylation has
gained significant attention since its initial discovery[38] – in part due to the availability of
phospho-specific antibodies – we now show that the C-terminal domain contains a conserved
PTM-enriched region where SUMOylation at K408 precedes phosphorylation, revealing
additional complexity and nuance to the stimulus-induced regulation of TDP-43.

TDP-43 SUMOylation at K408 supports cellular stress response and recovery

To understand the cellular and physiological roles of TDP-43 SUMOylation, we
generated a knock in mouse model harboring a missense (c.1223 A>G) point mutation in the
endogenous mouse *Tardbp* locus resulting in the expression of TDP-43 bearing a p.K408R
missense mutation blocking endogenous TDP-43 SUMOylation (Fig. 2E, S2A). For simplicity
herein, all genes/proteins will be referred to following their human nomenclature. Although
TDP-43 and SUMO2 are independently essential for embryonic development, the "TDP-43^{K408R}"
mice are born at normal mendelian ratios indicating that TDP-43 SUMOylation at K408 is not
essential for murine embryonic development (fig. S2B)(*53*, *68*).

As TDP-43 SUMOylation responds to, and is cleared during cellular stress and recovery
respectively, we posited that SUMOylated TDP-43 plays key roles in these processes. We
cultured TDP-43$^{+/+}$ and TDP-43$^{K408R/K408R}$ primary cortical neurons and performed stress and
recovery assays to determine how the cell stress response may be affected. Since TDP-43 is
critical for the formation of G3BP1+ stress granules, we examined their formation and resolution
following stress and recovery(*69*, *70*). We found that stress granules could form in
TDP-43$^{K408R/K408R}$ neurons to the same degree as TDP-43$^{+/+}$ neurons, but there was a significant
delay in stress granule clearance during recovery (Fig. 2F, S3A, S3B). Despite this deficiency,
there was no significant change in neuron viability during recovery (fig. S3E). To our surprise,
we also observed a significant increase in RIPA insoluble phosphorylated TDP-43 at 1 hour
recovery that was cleared by 3 hours post-stress recovery (Fig. 2G). Based on our previous
results that SUMOylation is cleared by the UPS pathway and phosphorylation may function
antagonistically with SUMOylation, the increase in phosphorylation during recovery may serve
as an alternative method of proteostasis to compensate for loss of SUMOylation. We further
observed a significant increase in nuclear TDP-43 foci that were specific to the recovery phase
and were exacerbated after 3 hours recovery in the TDP-43$^{K408R/K408R}$ neurons (Fig. 2F, S2C,
S2D). Various forms of nuclear TDP-43 bodies are increasingly recognized to be aberrantly
regulated in disease and stress states (*21*, *71–73*). Together, this may indicate that
TDP-43$^{K408R/K408R}$ neurons appropriately respond to, but not efficiently resolve cellular stress
which can render cells vulnerable to further insult.

237 Blocking TDP-43 SUMOylation *in vivo* leads to sex-specific social and cognitive deficits and
238 TDP-43 proteinopathy reminiscent of FTD

Aging is one of the strongest risk factors for many neurodegenerative diseases including
ALS/FTD. The process of aging is known to progressively induce stress on cells, particularly in
the nervous system, as cellular functions become impaired or as environmental triggers
accumulate over time(*74*). As TDP-43 SUMOylation is involved in supporting the normal
cellular stress response and maintaining TDP-43 proteostasis, we hypothesized that disrupting
this process leads to age-dependent physiological changes accelerated by impaired stress
responses and TDP-43 accumulation.

In young pups, we did not observe any significant differences between TDP-43$^{+/+}$,
TDP-43$^{K408R/+}$, or TDP-43$^{K408R/K408R}$ with respect to body weight or motor abilities indicating
that blocking TDP-43 SUMOylation does not lead to gross developmental abnormalities (fig.
S4). We performed a battery of behavioral tests focusing on motor, social, and cognitive function
related to ALS and FTD at three time points (2, 9 and 16 months, fig. S5-S8, table S2). We
observed sex-specific cognitive and social effects biased to female animals that generally
worsened with aging. Starting around 2 months of age, we observed significant barbering in
female mice leading to alopecia in the K408R mutants (Fig. 3A). Consistent with this finding we

found that aged female TDP-43$^{K408R/K408R}$ mice were significantly more submissive to their
TDP-43$^{+/+}$ counterparts in the tube test (Fig. 3B). Additionally, we observed age dependent
changes in behavior, such as significant risk taking and hyperactivity in the open field test at 2
months of age (Fig. 3C). This hyperactivity was age-dependent and selectively affected young
female mice as they were significantly less active than their TDP-43$^{+/+}$ counterparts at 9 months
of age in the open field tests (Fig. 3C). Additionally, these female TDP-43$^{K408R/K408R}$ mice
presented with significant impairment in habituation at 9 months in the beam break test
(Fig. 3D). By 16 months of age, female TDP-43$^{K408R/K408R}$ mice performed significantly worse
than TDP-43$^{+/+}$ mice in the spontaneous Y-maze, inferring age-dependent cognitive decline in
the domain of working memory (Fig. 3E). There were no significant cognitive effects in male
TDP-43$^{K408R/K408R}$ mice, however impairments were observed on the rotarod test at 9 months of
age (fig. S6). This deficiency was not observed at 16 months of age, largely due to all mice
performing significantly worse at the rotarod task, limiting interpretation. Taken together, the
age-dependent cognitive and social impairment observed in female TDP-43 SUMO-deficient
mutants highlight interesting and unexplored aspects of TDP-43 with important implications for
understanding sex-specific roles of TDP-43 SUMOylation.

Cellular and molecular changes are generally thought to precede behavioral changes in
mouse models and human disease. Consistent with the behavioral findings described above, we
find that insoluble phosphorylated TDP-43 accumulated in the cortex of female, but not male
mice, and mislocalized and phosphorylated TDP-43 could be observed in the lumbar spinal cord
of both male and female mice at 9 months of age (Fig. 3F, S9, S10A). At 16 months of age, male
TDP-43$^{K408R/K408R}$ mice displayed a significant denervation of neuromuscular junctions in the
tibialis anterior and a reduction in ChAT positive neurons in the spinal cord compared to
littermate controls (Fig. 3G, fig S10B, S10C). These data support that TDP-43 SUMOylation
plays a protective role in mediating TDP-43 proteostasis in the CNS and that its blockade may
confer a risk for ALS/FTD-like pathogenesis in an age-dependent, and sex-specific manner
providing insights into molecular substrates underlying sexually dimorphic clinical features of
ALS and FTD.

Aberrant SUMOylation is a Feature of Human Aging and Disease

Although SUMOylation is well understood to play an essential role in the CNS, its
involvement in human aging and neurodegenerative diseases remains largely unknown. To
assess how SUMOylation is normally affected in human aging, we processed samples from
human temporal lobe tissue ranging between 0.5 and 88 years of age from individuals unaffected
by neurological diseases (Fig. 4A, S11A, table S3, n=55). Complementing our findings that
SUMOylation serves as a protective feature of aging in our TDP-43^{K408R} model, we observed
that global SUMOylation was also positively correlated with age. This supports a link between
increased SUMOylation and normal aging in the human brain, possibly as a protective
mechanism due to increased proteostasis demand (Fig. 4B).

To explore the extent of TDP-43 SUMOylation in human ALS/FTD cases, we performed
proximity ligation assays against TDP-43 and SUMO2/3 in prefrontal cortex samples from
ALS/FTD cases positive for TDP-43 pathology and age/sex matched unaffected controls
(fig. S11B, table S3). Surprisingly, we observed a significant increase in TDP-43 – SUMO2/3
interactions in the prefrontal cortex of ALS/FTD patients with TDP-43 pathology (Fig. 4C). This
may be a result of increased stress burden throughout the prefrontal cortex in those affected by

ALS/FTD with TDP-43 pathology compared to age and sex matched counterparts. Taken
together, SUMOylation as a stress responsive modification is positively correlated with aging in
the human brain, potentially linked to the increase demand imposed by age-associated stress.
Additionally, the SUMOylation response is enhanced in the frontal cortex of those affected by
ALS/FTD providing further *in vivo* support for its role in disease.

Discussion

Although ALS and FTD are highly heterogenous diseases, the nearly universal
convergence on TDP-43 emphasizes a need to understand early drivers of TDP-43 pathogenesis.
Despite this, there remains a gap in knowledge about the mechanisms maintaining proteostatic
balance and how their disruption ultimately leads to dysfunction of TDP-43. We found that
SUMOylation is a conserved mechanism regulating nuclear TDP-43 function selectively during
cellular stress that subsequently promotes UPS-mediated clearance during recovery. Indeed,
recent reports have found that other stress related proteins such as G3BP1 are differentially
modified by PTMs according to the type of stress that cells are responding to, highlighting
unique ways stress-related proteins can execute different functions in a context-specific
manner(*75, 76*). This context-specific regulation provides unique opportunities to understand
specific pathways activated in disease that may be therapeutically targeted. Future studies
modulating TDP-43 E3 SUMO ligases may help alter the cell stress response, potentially by
decreasing aberrant TDP-43 activity or by promoting stress-specific responses to ease the burden
in vulnerable cells.

SUMOylation of TDP-43 occurs at K408 in a highly conserved region of the C-terminal
domain directly adjacent to phosphorylation sites that are characteristically phosphorylated in
aggregates in TDP-43 proteinopathies(*38*). We found that phosphorylation and SUMOylation act
antagonistically on TDP-43, indicating potential distinct mechanisms of regulation. As
SUMOylated TDP-43 is cleared by the UPS-pathway during stress recovery we hypothesized
that blocking TDP-43 SUMOylation with K408R mutation would result in its accumulation
during stress recovery. Interestingly, we observed a significant increase in insoluble
phosphorylated TDP-43 in TDP-43$^{K408R/K408R}$ cortical neurons and a delay in stress granule
disassembly during stress recovery indicating TDP-43 accumulation. However, this was resolved
3 hours post-stress indicating that the neurons could compensate to clear the aberrantly
accumulated TDP-43 within this acute stress paradigm. Future studies exploring repeated and/or
divergent stressors will be crucial in determining the degree to which TDP-43 SUMOylation
safeguards from cellular demise.

Previous reports have emphasized the status of altered proteostasis as a hallmark of aging
in the CNS, culminating in impaired removal of damaged proteins with age(*74*). In
TDP-43$^{K408R/K408R}$ mice, TDP-43 was found to accumulate in the CNS in an age-dependent
manner. TDP-43 accumulation within the CNS progresses with age, inferring an age-dependent
requirement for SUMOylation to maintain proteostasis. Indeed, in human temporal lobe samples,
we similarly observed a positive correlation between global SUMOylation and age supporting
SUMOylation as a potential protective mechanism during aging in the CNS. However, directly
monitoring SUMOylation events remains challenging due to the dynamic nature of the
modification and lack of specific tools to study its interaction. We employed a PLA approach to
assess the interaction between TDP-43 and SUMO in ALS/FTD patients and found significantly
increased interactions in the prefrontal cortex relative to age/sex matched unaffected controls.
Recent studies have used RNA aptamers to reveal nuclear changes in TDP-43 in spinal and

cortical neurons of ALS patients prior to traditional cytoplasmic pathology in vulnerable cells(*21*). Further development of these technologies with respect to TDP-43 SUMOylation may help advance biomarker development that can provide information about TDP-43 stress-response and pathogenesis in ALS/FTD patients.

The emergence of predominantly cognitive FTD-like phenotypes in the TDP-43[K408R] mice is not unusual compared to other TDP-43 knock in models such as TDP-43[Q331K] and TDP-43[K145R] mice which present with features of cognitive dysfunction but lacking the entire constellation of ALS/FTD-like phenotypes(*41, 77*). Though we initially hypothesized that aging may be a sufficient stressor to interrogate TDP-43 SUMOylation in mice, the various stressors that individuals with ALS/FTD were previously exposed to during their lifetime (i.e. the "exposome") are not experienced by mice in vivaria on a short timescale(*69*). The TDP-43[K408R] model will therefore facilitate the exploration of how the exposome synergizes with TDP-43 to better understand how ALS/FTD relevant stressors (e.g. *C9ORF72* expansions, viral infections, and traumatic brain injury) converge on TDP-43 and uniquely drive neurodegeneration.

References

1. O. Hardiman, A. Al-Chalabi, A. Chio, E. M. Corr, G. Logroscino, W. Robberecht, P. J. Shaw, Z. Simmons, L. H. Van Den Berg, Amyotrophic lateral sclerosis. *Nature Reviews Disease Primers 2017 3:1* **3**, 1–19 (2017).

2. M. Grossman, W. W. Seeley, A. L. Boxer, A. E. Hillis, D. S. Knopman, P. A. Ljubenov, B. Miller, O. Piguet, R. Rademakers, J. L. Whitwell, H. Zetterberg, J. C. van Swieten, Frontotemporal lobar degeneration. *Nature Reviews Disease Primers 2023 9:1* **9**, 1–19 (2023).

3. P. T. Nelson, D. W. Dickson, J. Q. Trojanowski, C. R. Jack, P. A. Boyle, K. Arfanakis, R. Rademakers, I. Alafuzoff, J. Attems, C. Brayne, I. T. S. Coyle-Gilchrist, H. C. Chui, D. W. Fardo, M. E. Flanagan, G. Halliday, S. R. K. Hokkanen, S. Hunter, G. A. Jicha, Y. Katsumata, C. H. Kawas, C. D. Keene, G. G. Kovacs, W. A. Kukull, A. I. Levey, N. Makkinejad, T. J. Montine, S. Murayama, M. E. Murray, S. Nag, R. A. Rissman, W. W. Seeley, R. A. Sperling, C. L. White Iii, L. Yu, J. A. Schneider, Limbic-predominant age-related TDP-43 encephalopathy (LATE): consensus working group report. doi: 10.1093/brain/awz099.

4. A. Meneses, S. Koga, J. O'Leary, D. W. Dickson, G. Bu, N. Zhao, TDP-43 Pathology in Alzheimer's Disease. *Molecular Neurodegeneration 2021 16:1* **16**, 1–15 (2021).

5. R. Nicks, N. F. Clement, V. E. Alvarez, Y. Tripodis, Z. H. Baucom, B. R. Huber, J. Mez, M. L. Alosco, N. Aytan, J. D. Cherry, K. A. Cormier, C. Kubilius, R. Mathias, S. E. Svirsky, M. J. Pothast, A. M. Hildebrandt, J. Chung, X. Han, J. F. Crary, A. C. McKee, M. P. Frosch, T. D. Stein, Repetitive head impacts and chronic traumatic encephalopathy are associated with TDP-43 inclusions and hippocampal sclerosis. *Acta Neuropathol* **145**, 395–408 (2023).

6. S. S. Thammisetty, J. Pedragosa, Y. C. Weng, F. Calon, A. Planas, J. Kriz, Age-related deregulation of TDP-43 after stroke enhances NF-κB-mediated inflammation and neuronal damage. *J Neuroinflammation* **15**, 1–15 (2018).

7. T. R. Suk, M. W. C. Rousseaux, The role of TDP-43 mislocalization in amyotrophic lateral sclerosis. *Mol Neurodegener* **15**, 45 (2020).

8.   J. Chew, T. F. Gendron, M. Prudencio, H. Sasaguri, Y. J. Zhang, M. Castanedes-
Casey, C. W. Lee, K. Jansen-West, A. Kurti, M. E. Murray, K. F. Bieniek, P. O.
Bauer, E. C. Whitelaw, L. Rousseau, J. N. Stankowski, C. Stetler, L. M. Daughrity,
E. A. Perkerson, P. Desaro, A. Johnston, K. Overstreet, D. Edbauer, R.
Rademakers, K. B. Boylan, D. W. Dickson, J. D. Fryer, L. Petrucelli, C9ORF72
repeat expansions in mice cause TDP-43 pathology, neuronal loss, and behavioral
deficits. *Science (1979)* **348**, 1151–1154 (2015).
9.   W. Shao, T. W. Todd, Y. Wu, C. Y. Jones, J. Tong, K. Jansen-West, L. M.
Daughrity, J. Park, Y. Koike, A. Kurti, M. Yue, M. Castanedes-Casey, G. del
Rosso, J. A. Dunmore, D. Z. Alepuz, B. Oskarsson, D. W. Dickson, C. N. Cook, M.
Prudencio, T. F. Gendron, J. D. Fryer, Y. J. Zhang, L. Petrucelli, Two FTD-ALS
genes converge on the endosomal pathway to induce TDP-43 pathology and
degeneration. *Science (1979)* **378**, 94–99 (2022).
10. G. Fung, J. Shi, H. Deng, J. Hou, C. Wang, A. Hong, J. Zhang, W. Jia, H. Luo,
Cytoplasmic translocation, aggregation, and cleavage of TDP-43 by enteroviral
proteases modulate viral pathogenesis. *Cell Death & Differentiation 2015 22:12* **22**,
2087–2097 (2015).
11. P. E. A. Ash, E. A. Stanford, A. Al Abdulatif, A. Ramirez-Cardenas, H. I. Ballance,
S. Boudeau, A. Jeh, J. M. Murithi, Y. Tripodis, G. J. Murphy, D. H. Sherr, B.
Wolozin, Dioxins and related environmental contaminants increase TDP-43 levels.
Mol Neurodegener **12**, 1–14 (2017).
12. J. Sreedharan, I. P. Blair, V. B. Tripathi, X. Hu, C. Vance, B. Rogelj, S. Ackerley,
J. C. Durnall, K. L. Williams, E. Buratti, F. Baralle, J. De Belleroche, J. D.
Mitchell, P. N. Leigh, A. Al-Chalabi, C. C. Miller, G. Nicholson, C. E. Shaw, TDP-
43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science (1979)*
319, 1668–1672 (2008).
13. A. N. Coyne, V. Baskerville, B. L. Zaepfel, D. W. Dickson, F. Rigo, F. Bennett, C.
Patrick Lusk, J. D. Rothstein, Nuclear accumulation of CHMP7 initiates nuclear
pore complex injury and subsequent TDP-43 dysfunction in sporadic and familial
ALS. *Sci Transl Med* **13**, 1923 (2021).
14. X. R. Ma, M. Prudencio, Y. Koike, S. C. Vatsavayai, G. Kim, F. Harbinski, A.
Briner, C. M. Rodriguez, C. Guo, T. Akiyama, H. B. Schmidt, B. B. Cummings, D.
W. Wyatt, K. Kurylo, G. Miller, S. Mekhoubad, N. Sallee, G. Mekonnen, L.
Ganser, J. D. Rubien, K. Jansen-West, C. N. Cook, S. Pickles, B. Oskarsson, N. R.
Graff-Radford, B. F. Boeve, D. S. Knopman, R. C. Petersen, D. W. Dickson, J.
Shorter, S. Myong, E. M. Green, W. W. Seeley, L. Petrucelli, A. D. Gitler, TDP-43
represses cryptic exon inclusion in the FTD–ALS gene UNC13A. *Nature 2022*
603:7899 **603**, 124–130 (2022).
15. A. L. Brown, O. G. Wilkins, M. J. Keuss, S. E. Hill, M. Zanovello, W. C. Lee, A.
Bampton, F. C. Y. Lee, L. Masino, Y. A. Qi, S. Bryce-Smith, A. Gatt, M.
Hallegger, D. Fagegaltier, H. Phatnani, H. Phatnani, J. Kwan, D. Sareen, J. R.
Broach, Z. Simmons, X. Arcila-Londono, E. B. Lee, V. M. Van Deerlin, N. A.
Shneider, E. Fraenkel, L. W. Ostrow, F. Baas, N. Zaitlen, J. D. Berry, A.
Malaspina, P. Fratta, G. A. Cox, L. M. Thompson, S. Finkbeiner, E. Dardiotis, T.
M. Miller, S. Chandran, S. Pal, E. Hornstein, D. J. MacGowan, T. Heiman-
Patterson, M. G. Hammell, N. A. Patsopoulos, O. Butovsky, J. Dubnau, A. Nath, R.

Bowser, M. Harms, E. Aronica, M. Poss, J. Phillips-Cremins, J. Crary, N. Atassi, D. J. Lange, D. J. Adams, L. Stefanis, M. Gotkine, R. H. Baloh, S. Babu, T. Raj, S. Paganoni, O. Shalem, C. Smith, B. Zhang, B. Harris, I. Broce, V. Drory, J. Ravits, C. McMillan, V. Menon, L. Wu, S. Altschuler, Y. Lerner, R. Sattler, K. Van Keuren-Jensen, O. Rozenblatt-Rosen, K. Lindblad-Toh, K. Nicholson, P. Gregersen, J. H. Lee, S. Kokos, S. Muljo, J. Newcombe, E. K. Gustavsson, S. Seddighi, J. F. Reyes, S. L. Coon, D. Ramos, G. Schiavo, E. M. C. Fisher, T. Raj, M. Secrier, T. Lashley, J. Ule, E. Buratti, J. Humphrey, M. E. Ward, P. Fratta, TDP-43 loss and ALS-risk SNPs drive mis-splicing and depletion of UNC13A. *Nature 2022 603:7899* **603**, 131–137 (2022).

16. Z. Melamed, J. López-Erauskin, M. W. Baughn, O. Zhang, K. Drenner, Y. Sun, F. Freyermuth, M. A. McMahon, M. S. Beccari, J. W. Artates, T. Ohkubo, M. Rodriguez, N. Lin, D. Wu, C. F. Bennett, F. Rigo, S. Da Cruz, J. Ravits, C. Lagier-Tourenne, D. W. Cleveland, Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of TDP-43-dependent neurodegeneration. *Nat Neurosci*, 1 (2019).

17. J. R. Klim, L. A. Williams, F. Limone, I. Guerra San Juan, B. N. Davis-Dusenbery, D. A. Mordes, A. Burberry, M. J. Steinbaugh, K. K. Gamage, R. Kirchner, R. Moccia, S. H. Cassel, K. Chen, B. J. Wainger, C. J. Woolf, K. Eggan, ALS-implicated protein TDP-43 sustains levels of STMN2, a mediator of motor neuron growth and repair. *Nat Neurosci* **22**, 167–179 (2019).

18. F. Gasset-Rosa, S. Lu, H. Yu, C. Chen, Z. Melamed, L. Guo, J. Shorter, S. Da Cruz, D. W. Cleveland, Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron* **102**, 339-357.e7 (2019).

19. M. W. Baughn, Z. Melamed, J. López-Erauskin, M. S. Beccari, K. Ling, A. Zuberi, M. Presa, E. Gonzalo-Gil, R. Maimon, S. Vazquez-Sanchez, S. Chaturvedi, M. Bravo-Hernández, V. Taupin, S. Moore, J. W. Artates, E. Acks, I. Sandra Ndayambaje, A. R. Agra de Almeida Quadros, P. Jafar-Nejad, F. Rigo, C. Frank Bennett, C. Lutz, C. Lagier-Tourenne, D. W. Cleveland, Mechanism of STMN2 cryptic splice-polyadenylation and its correction for TDP-43 proteinopathies. *Science (1979)* **379**, 1140–1149 (2023).

20. A. R. Agra Almeida Quadros, Z. Li, X. Wang, I. S. Ndayambaje, S. Aryal, N. Ramesh, M. Nolan, R. Jayakumar, Y. Han, H. Stillman, C. Aguilar, H. J. Wheeler, T. Connors, J. Lopez-Erauskin, M. W. Baughn, Z. Melamed, M. S. Beccari, L. Olmedo Martínez, M. Canori, C. Z. Lee, L. Moran, I. Draper, A. S. Kopin, D. H. Oakley, D. W. Dickson, D. W. Cleveland, B. T. Hyman, S. Das, N. Ertekin-Taner, C. Lagier-Tourenne, Cryptic splicing of stathmin-2 and UNC13A mRNAs is a pathological hallmark of TDP-43-associated Alzheimer's disease. *Acta Neuropathologica 2024 147:1* **147**, 1–18 (2024).

21. H. Spence, F. M. Waldron, R. S. Saleeb, A. L. Brown, O. M. Rifai, M. Gilodi, F. Read, K. Roberts, G. Milne, D. Wilkinson, J. O'Shaughnessy, A. Pastore, P. Fratta, N. Shneider, G. G. Tartaglia, E. Zacco, M. H. Horrocks, J. M. Gregory, RNA aptamer reveals nuclear TDP-43 pathology is an early aggregation event that coincides with STMN-2 cryptic splicing and precedes clinical manifestation in ALS. *Acta Neuropathol* **147**, 1–15 (2024).

22. I. R. Mackenzie, A. M. Nicholson, M. Sarkar, J. Messing, M. D. Purice, C. Pottier,
K. Annu, M. Baker, R. B. Perkerson, A. Kurti, B. J. Matchett, T. Mittag, J.
Temirov, G. Y. R. Hsiung, C. Krieger, M. E. Murray, M. Kato, J. D. Fryer, L.
Petrucelli, L. Zinman, S. Weintraub, M. Mesulam, J. Keith, S. A. Zivkovic, V.
Hirsch-Reinshagen, R. P. Roos, S. Züchner, N. R. Graff-Radford, R. C. Petersen, R.
J. Caselli, Z. K. Wszolek, E. Finger, C. Lippa, D. Lacomis, H. Stewart, D. W.
Dickson, H. J. Kim, E. Rogaeva, E. Bigio, K. B. Boylan, J. P. Taylor, R.
Rademakers, TIA1 Mutations in Amyotrophic Lateral Sclerosis and Frontotemporal
Dementia Promote Phase Separation and Alter Stress Granule Dynamics. *Neuron*
95, 808-816.e9 (2017).
23. A. K. Walker, K. Y. Soo, V. Sundaramoorthy, S. Parakh, Y. Ma, M. A. Farg, R. H.
Wallace, P. J. Crouch, B. J. Turner, M. K. Horne, J. D. Atkin, ALS-associated
TDP-43 induces endoplasmic reticulum stress, which drives cytoplasmic TDP-43
accumulation and stress granule formation. *PLoS One* **8**, e81170 (2013).
24. O. H. Tam, N. V. Rozhkov, R. Shaw, D. Kim, I. Hubbard, S. Fennessey, N. Propp,
H. Phatnani, J. Kwan, D. Sareen, J. R. Broach, Z. Simmons, X. Arcila-Londono, E.
B. Lee, V. M. Van Deerlin, N. A. Shneider, E. Fraenkel, L. W. Ostrow, F. Baas, N.
Zaitlen, J. D. Berry, A. Malaspina, P. Fratta, G. A. Cox, L. M. Thompson, S.
Finkbeiner, E. Dardiotis, T. M. Miller, S. Chandran, S. Pal, E. Hornstein, D. J.
MacGowan, T. Heiman-Patterson, M. G. Hammell, N. A. Patsopoulos, O.
Butovsky, J. Dubnau, A. Nath, R. Bowser, M. Harms, E. Aronica, M. Poss, J.
Phillips-Cremins, J. Crary, N. Atassi, D. J. Lange, D. J. Adams, L. Stefanis, M.
Gotkine, R. Baloh, S. Babu, T. Raj, S. Paganoni, O. Shalem, C. Smith, B. Zhang, B.
T. Harris, D. Fagegaltier, J. Ravits, J. Dubnau, M. Gale Hammell, Postmortem
Cortex Samples Identify Distinct Molecular Subtypes of ALS: Retrotransposon
Activation, Oxidative Stress, and Activated Glia. *Cell Rep* **29**, 1164-1177.e5
(2019).
25. B. Wolozin, P. Ivanov, Stress granules and neurodegeneration. Nature Publishing
Group [Preprint] (2019). https://doi.org/10.1038/s41583-019-0222-5.
26. M. T. Lin, M. F. Beal, Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature Publishing Group [Preprint] (2006).
https://doi.org/10.1038/nature05292.
27. A. Ratti, V. Gumina, P. Lenzi, P. Bossolasco, F. Fulceri, C. Volpe, D. Bardelli, F.
Pregnolato, A. M. Maraschi, F. Fornai, V. Silani, C. Colombrita, Chronic stress
induces formation of stress granules and pathological TDP-43 aggregates in human
ALS fibroblasts and iPSC-motoneurons. *Neurobiol Dis* **145**, 105051 (2020).
28. E. D'Amico, P. Factor-Litvak, R. M. Santella, H. Mitsumoto, Clinical perspective
on oxidative stress in sporadic amyotrophic lateral sclerosis. NIH Public Access
[Preprint] (2013). https://doi.org/10.1016/j.freeradbiomed.2013.06.029.
29. C. Colombrita, E. Zennaro, C. Fallini, M. Weber, A. Sommacal, E. Buratti, V.
Silani, A. Ratti, TDP-43 is recruited to stress granules in conditions of oxidative
insult. *J Neurochem* **111**, 1051–1061 (2009).
30. X. Zuo, J. Zhou, Y. Li, K. Wu, Z. Chen, Z. Luo, X. Zhang, Y. Liang, M. A.
Esteban, Y. Zhou, X. D. Fu, TDP-43 aggregation induced by oxidative stress causes
global mitochondrial imbalance in ALS. *Nature Structural & Molecular Biology*
2021 28:2 **28**, 132–142 (2021).

31. T. Ueda, T. Takeuchi, N. Fujikake, M. Suzuki, E. N. Minakawa, M. Ueyama, Y. Fujino, N. Kimura, S. Nagano, A. Yokoseki, O. Onodera, H. Mochizuki, T. Mizuno, K. Wada, Y. Nagai, Dysregulation of stress granule dynamics by DCTN1 deficiency exacerbates TDP-43 pathology in Drosophila models of ALS/FTD. *Acta Neuropathol Commun* **12**, 1–15 (2024).

32. Y. Iguchi, M. Katsuno, S. Takagi, S. Ishigaki, J. ichi Niwa, M. Hasegawa, F. Tanaka, G. Sobue, Oxidative stress induced by glutathione depletion reproduces pathological modifications of TDP-43 linked to TDP-43 proteinopathies. *Neurobiol Dis* **45**, 862–870 (2012).

33. L. Liu-Yesucevitz, A. Bilgutay, Y. J. Zhang, T. Vanderwyde, A. Citro, T. Mehta, N. Zaarur, A. McKee, R. Bowser, M. Sherman, L. Petrucelli, B. Wolozin, Tar DNA binding protein-43 (TDP-43) associates with stress granules: Analysis of cultured cells and pathological brain tissue. *PLoS One* **5**, e13250 (2010).

34. K. K. McDonald, A. Aulas, L. Destroismaisons, S. Pickles, E. Beleac, W. Camu, G. A. Rouleau, C. Vande Velde, TAR DNA-binding protein 43 (TDP-43) regulates stress granule dynamics via differential regulation of G3BP and TIA-1. *Hum Mol Genet* **20**, 1400–1410 (2011).

35. W. Luan, A. L. Wright, H. Brown-Wright, S. Le, R. San Gil, L. Madrid San Martin, K. Ling, P. Jafar-Nejad, F. Rigo, A. K. Walker, Early activation of cellular stress and death pathways caused by cytoplasmic TDP-43 in the rNLS8 mouse model of ALS and FTD. *Molecular Psychiatry 2023 28:6* **28**, 2445–2461 (2023).

36. A. Dubinski, M. Gagne, S. Peyrard, D. Gordon, K. Talbot, C. Vande Velde, Stress granule assembly in vivo is deficient in the CNS of mutant TDP-43 ALS mice. *Hum Mol Genet* **32**, 319–332 (2023).

37. P. Zhang, B. Fan, P. Yang, J. Temirov, J. Messing, H. J. Kim, J. P. Taylor, Chronic optogenetic induction of stress granules is cytotoxic and reveals the evolution of ALS-FTD pathology. *Elife* **8** (2019).

38. M. Neumann, L. K. Kwong, E. B. Lee, E. Kremmer, A. Flatley, Y. Xu, M. S. Forman, D. Troost, H. A. Kretzschmar, J. Q. Trojanowski, V. M. Y. Lee, Phosphorylation of S409/410 of TDP-43 is a consistent feature in all sporadic and familial forms of TDP-43 proteinopathies. *Acta Neuropathol* **117**, 137–149 (2009).

39. M. Neumann, D. M. Sampathu, L. K. Kwong, A. C. Truax, M. C. Micsenyi, T. T. Chou, J. Bruce, T. Schuck, M. Grossman, C. M. Clark, L. F. McCluskey, B. L. Miller, E. Masliah, I. R. Mackenzie, H. Feldman, W. Feiden, H. A. Kretzschmar, J. Q. Trojanowski, V. M.-Y. Lee, Ubiquitinated TDP-43 in Frontotemporal Lobar Degeneration and Amyotrophic Lateral Sclerosis. *Science (1979)* **314**, 130–133 (2006).

40. T. Arai, M. Hasegawa, H. Akiyama, K. Ikeda, T. Nonaka, H. Mori, D. Mann, K. Tsuchiya, M. Yoshida, Y. Hashizume, T. Oda, TDP-43 is a component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Biochem Biophys Res Commun* **351**, 602–611 (2006).

41. J. Necarsulmer, J. Simon, B. Evangelista, Y. Chen, X. Tian, S. Nafees, A. M. Gonzalez, H. Jiang, P. Wang, D. Ajit, V. Nikolova, K. Harper, J. Ezzell, F.-C. Lin, A. Beltran, S. Moy, T. Cohen, RNA-binding deficient TDP-43 drives cognitive decline in a mouse model of TDP-43 proteinopathy. *Elife* **12** (2023).

42. L. McGurk, E. Gomes, L. Guo, J. Mojsilovic-Petrovic, V. Tran, R. G. Kalb, J. Shorter, N. M. Bonini, Poly(ADP-Ribose) Prevents Pathological Phase Separation of TDP-43 by Promoting Liquid Demixing and Stress Granule Localization. *Mol Cell* **71**, 703-717.e9 (2018).

43. H. Marmor-Kollet, A. Siany, N. Kedersha, N. Knafo, N. Rivkin, Y. M. Danino, T. G. Moens, T. Olender, D. Sheban, N. Cohen, T. Dadosh, Y. Addadi, R. Ravid, C. Eitan, B. Toth Cohen, S. Hofmann, C. L. Riggs, V. M. Advani, A. Higginbottom, J. Cooper-Knock, J. H. Hanna, Y. Merbl, L. Van Den Bosch, P. Anderson, P. Ivanov, T. Geiger, E. Hornstein, Spatiotemporal Proteomic Analysis of Stress Granule Disassembly Using APEX Reveals Regulation by SUMOylation and Links to ALS Pathogenesis. *Mol Cell* **80**, 876-891.e6 (2020).

44. A. Osmanovic, A. Förster, M. Widjaja, B. Auber, · Anibh, M. Das, A. Christians, · Frank Brand, S. Petri, R. G. Weber, A SUMO4 initiator codon variant in amyotrophic lateral sclerosis reduces SUMO4 expression and alters stress granule dynamics. *Journal of Neurology 2022* **1**, 1–9 (2022).

45. A. M. Maraschi, V. Gumina, J. Dragotto, C. Colombrita, M. Mompeán, E. Buratti, V. Silani, M. Feligioni, A. Ratti, SUMOylation Regulates TDP-43 Splicing Activity and Nucleocytoplasmic Distribution. *Mol Neurobiol* **58**, 5682–5702 (2021).

46. C. Maurel, A. A. Chami, R. A. Thépault, S. Marouillat, H. Blasco, P. Corcia, C. R. Andres, P. Vourc'h, A role for SUMOylation in the Formation and Cellular Localization of TDP-43 Aggregates in Amyotrophic Lateral Sclerosis. *Mol Neurobiol* **57**, 1361–1373 (2020).

47. N. T. Seyfried, Y. M. Gozal, E. B. Dammer, Q. Xia, D. M. Duong, D. Cheng, J. J. Lah, A. I. Levey, J. Peng, Multiplex SILAC analysis of a cellular TDP-43 proteinopathy model reveals protein inclusions associated with SUMOylation and diverse polyubiquitin chains. *Molecular and Cellular Proteomics* **9**, 705–718 (2010).

48. J. M. Henley, T. J. Craig, K. A. Wilkinson, Neuronal SUMOylation: mechanisms, physiology, and roles in neuronal dysfunction. American Physiological Society Bethesda, MD [Preprint] (2014). https://doi.org/10.1152/physrev.00008.2014.

49. K. A. Wilkinson, J. M. Henley, Mechanisms, regulation and consequences of protein SUMOylation. *Biochemical Journal* **428**, 133–145 (2010).

50. J. M. Enserink, Sumo and the cellular stress response. BioMed Central Ltd. [Preprint] (2015). https://doi.org/10.1186/s13008-015-0010-1.

51. C. M. Dewey, B. Cenik, C. F. Sephton, D. R. Dries, P. Mayer, S. K. Good, B. A. Johnson, J. Herz, G. Yu, TDP-43 Is Directed to Stress Granules by Sorbitol, a Novel Physiological Osmotic and Oxidative Stressor. *Mol Cell Biol* **31**, 1098–1108 (2011).

52. H. Y. Chang, S. C. Hou, T. Der Way, C. H. Wong, I. F. Wang, Heat-shock protein dysregulation is associated with functional and pathological TDP-43 aggregation. *Nat Commun* **4**, 1–11 (2013).

53. L. Wang, C. Wansleeben, S. Zhao, P. Miao, W. Paschen, W. Yang, SUMO 2 is essential while SUMO 3 is dispensable for mouse embryonic development . *EMBO Rep* **15**, 878–885 (2014).

54. T. R. Suk, T. T. Nguyen, Z. A. Fisk, M. Mitkovski, H. M. Geertsma, J. L. A. Parmasad, M. M. Heer, S. M. Callaghan, F. Benseler, N. Brose, M. Tirard, M. W.

C. Rousseaux, Characterizing the differential distribution and targets of Sumo1 and Sumo2 in the mouse brain. *iScience* **26**, 106350 (2023).

55. J. R. Mann, A. M. Gleixner, J. C. Mauna, E. Gomes, M. R. DeChellis-Marks, P. G. Needham, K. E. Copley, B. Hurtle, B. Portz, N. J. Pyles, L. Guo, C. B. Calder, Z. P. Wills, U. B. Pandey, J. K. Kofler, J. L. Brodsky, A. Thathiah, J. Shorter, C. J. Donnelly, RNA Binding Antagonizes Neurotoxic Phase Transitions of TDP-43. *Neuron* **102**, 321-338.e8 (2019).

56. J. Keiten-Schmitz, K. Wagner, T. Piller, M. Kaulich, S. Alberti, S. Müller, The Nuclear SUMO-Targeted Ubiquitin Quality Control Network Regulates the Dynamics of Cytoplasmic Stress Granules. *Mol Cell* **79**, 54-67.e7 (2020).

57. L. E. Warner, P. Mancias, I. J. Butler, C. M. McDonald, L. Keppen, K. G. Koob, J. R. Lupski, Mutations in the early growth response 2 (EGR2) gene are associated with hereditary myelinopathies. *Nature Genetics 1998 18:4* **18**, 382–384 (1998).

58. P. Garcáa-Gutiérrez, F. Juárez-Vicente, F. Gallardo-Chamizo, P. Charnay, M. García-Domínguez, The transcription factor Krox20 is an E3 ligase that sumoylates its Nab coregulators. *EMBO Rep* **12**, 1018–1023 (2011).

59. T. Nishida, H. Yasuda, PIAS1 and PIASxα Function as SUMO-E3 Ligases toward Androgen Receptor and Repress Androgen Receptor-dependent Transcription. *Journal of Biological Chemistry* **277**, 41311–41317 (2002).

60. Q. Liang, H. Deng, X. Li, X. Wu, Q. Tang, T.-H. Chang, H. Peng, F. J. Rauscher, K. Ozato, F. Zhu, Tripartite Motif-Containing Protein 28 Is a Small Ubiquitin-Related Modifier E3 Ligase and Negative Regulator of IFN Regulatory Factor 7. *The Journal of Immunology* **187**, 4754–4763 (2011).

61. Q. Zhao, Y. Xie, Y. Zheng, S. Jiang, W. Liu, W. Mu, Z. Liu, Y. Zhao, Y. Xue, J. Ren, GPS-SUMO: A tool for the prediction of sumoylation sites and SUMO-interaction motifs. *Nucleic Acids Res* **42**, W325–W330 (2014).

62. J. Ren, X. Gao, C. Jin, M. Zhu, X. Wang, A. Shaw, L. Wen, X. Yao, Y. Xue, Systematic study of protein sumoylation: Development of a site-specific predictor of SUMOsp 2.0. *Proteomics* **9**, 3409–3412 (2009).

63. U. Karvonen, T. Jääskeläinen, M. Rytinki, S. Kaikkonen, J. J. Palvimo, ZNF451 Is a Novel PML Body- and SUMO-Associated Transcriptional Coregulator. *J Mol Biol* **382**, 585–600 (2008).

64. N. Eisenhardt, V. K. Chaugule, S. Koidl, M. Droescher, E. Dogan, J. Rettich, P. Sutinen, S. Y. Imanishi, K. Hofmann, J. J. Palvimo, A. Pichler, A new vertebrate SUMO enzyme family reveals insights into SUMO-chain assembly. *Nature Structural & Molecular Biology 2015 22:12* **22**, 959–967 (2015).

65. R. J. Lumpkin, H. Gu, Y. Zhu, M. Leonard, A. S. Ahmad, K. R. Clauser, J. G. Meyer, E. J. Bennett, E. A. Komives, Site-specific identification and quantitation of endogenous SUMO modifications under native conditions. *Nature Communications 2017 8:1* **8**, 1–11 (2017).

66. I. A. Hendriks, D. Lyon, D. Su, N. H. Skotte, J. A. Daniel, L. J. Jensen, M. L. Nielsen, Site-specific characterization of endogenous SUMOylation across species and organs. *Nature Communications 2018 9:1* **9**, 1–17 (2018).

67. S. Chen, L. C. Francioli, J. K. Goodrich, R. L. Collins, M. Kanai, Q. Wang, J. Alföldi, N. A. Watts, C. Vittal, L. D. Gauthier, T. Poterba, M. W. Wilson, Y. Tarasova, W. Phu, R. Grant, M. T. Yohannes, Z. Koenig, Y. Farjoun, E. Banks, S.

Donnelly, S. Gabriel, N. Gupta, S. Ferriera, C. Tolonen, S. Novod, L. Bergelson, D. Roazen, V. Ruano-Rubio, M. Covarrubias, C. Llanwarne, N. Petrillo, G. Wade, T. Jeandet, R. Munshi, K. Tibbetts, M. Abreu, C. A. Aguilar Salinas, T. Ahmad, C. M. Albert, D. Ardissino, I. M. Armean, E. G. Atkinson, G. Atzmon, J. Barnard, S. M. Baxter, L. Beaugerie, E. J. Benjamin, D. Benjamin, M. Boehnke, L. L. Bonnycastle, E. P. Bottinger, D. W. Bowden, M. J. Bown, H. Brand, S. Brant, T. Brookings, S. Bryant, S. E. Calvo, H. Campos, J. C. Chambers, J. C. Chan, K. R. Chao, S. Chapman, D. I. Chasman, R. Chisholm, J. Cho, R. Chowdhury, M. K. Chung, W. K. Chung, K. Cibulskis, B. Cohen, K. M. Connolly, A. Correa, B. B. Cummings, D. Dabelea, J. Danesh, D. Darbar, P. Darnowsky, J. Denny, R. Duggirala, J. Dupuis, P. T. Ellinor, R. Elosua, J. Emery, E. England, J. Erdmann, T. Esko, E. Evangelista, D. Fatkin, J. Florez, A. Franke, J. Fu, M. Färkkilä, K. Garimella, J. Gentry, G. Getz, D. C. Glahn, B. Glaser, S. J. Glatt, D. Goldstein, C. Gonzalez, L. Groop, S. Gudmundsson, A. Haessly, C. Haiman, I. Hall, C. L. Hanis, M. Harms, M. Hiltunen, M. M. Holi, C. M. Hultman, C. Jalas, M. Kallela, D. Kaplan, J. Kaprio, S. Kathiresan, E. E. Kenny, B. J. Kim, Y. J. Kim, D. King, G. Kirov, J. Kooner, S. Koskinen, H. M. Krumholz, S. Kugathasan, S. H. Kwak, M. Laakso, N. Lake, T. Langsford, K. M. Laricchia, T. Lehtimäki, M. Lek, E. Lipscomb, R. J. F. Loos, W. Lu, S. A. Lubitz, T. T. Luna, R. C. W. Ma, G. M. Marcus, J. Marrugat, K. M. Mattila, S. McCarroll, M. I. McCarthy, J. L. McCauley, D. McGovern, R. McPherson, J. B. Meigs, O. Melander, A. Metspalu, D. Meyers, E. V. Minikel, B. D. Mitchell, V. K. Mootha, A. Naheed, S. Nazarian, P. M. Nilsson, M. C. O'Donovan, Y. Okada, D. Ongur, L. Orozco, M. J. Owen, C. Palmer, N. D. Palmer, A. Palotie, K. S. Park, C. Pato, A. E. Pulver, D. Rader, N. Rahman, A. Reiner, A. M. Remes, D. Rhodes, S. Rich, J. D. Rioux, S. Ripatti, D. M. Roden, J. I. Rotter, N. Sahakian, D. Saleheen, V. Salomaa, A. Saltzman, N. J. Samani, K. E. Samocha, A. Sanchis-Juan, J. Scharf, M. Schleicher, H. Schunkert, S. Schönherr, E. G. Seaby, S. H. Shah, M. Shand, T. Sharpe, M. B. Shoemaker, T. Shyong, E. K. Silverman, M. Singer-Berk, P. Sklar, J. T. Smith, J. G. Smith, H. Soininen, H. Sokol, R. G. Son, J. Soto, T. Spector, C. Stevens, N. O. Stitziel, P. F. Sullivan, J. Suvisaari, E. S. Tai, K. D. Taylor, Y. Y. Teo, M. Tsuang, T. Tuomi, D. Turner, T. Tusie-Luna, E. Vartiainen, M. Vawter, L. Wang, A. Wang, J. S. Ware, H. Watkins, R. K. Weersma, B. Weisburd, M. Wessman, N. Whiffin, J. G. Wilson, R. J. Xavier, A. O'Donnell-Luria, M. Solomonson, C. Seed, A. R. Martin, M. E. Talkowski, H. L. Rehm, M. J. Daly, G. Tiao, B. M. Neale, D. G. MacArthur, K. J. Karczewski, A genomic mutational constraint map using variation in 76,156 human genomes. *Nature 2023 625:7993* **625**, 92–100 (2023).

68. B. C. Kraemer, T. Schuck, J. M. Wheeler, L. C. Robinson, J. Q. Trojanowski, V. M. Y. Lee, G. D. Schellenberg, Loss of Murine TDP-43 disrupts motor function and plays an essential role in embryogenesis. *Acta Neuropathol* **119**, 409–419 (2010).

69. Y. Khalfallah, R. Kuta, C. Grasmuck, A. Prat, H. D. Durham, C. Vande Velde, TDP-43 regulation of stress granule dynamics in neurodegenerative disease-relevant cell types. *Sci Rep* **8**, 7551 (2018).

70. H. Sidibé, Y. Khalfallah, S. Xiao, N. B. Gómez, H. Fakim, E. M. H. Tank, G. Di Tomasso, E. Bareke, A. Aulas, P. M. McKeever, Z. Melamed, L. Destroimaisons,

J.-E. Deshaies, L. Zinman, J. A. Parker, P. Legault, M. Tétreault, S. J. Barmada, J. Robertson, C. Vande Velde, TDP-43 stabilizes G3BP1 mRNA: relevance to amyotrophic lateral sclerosis/frontotemporal dementia . *Brain*, doi: 10.1093/brain/awab217 (2021).

71. T. A. Shelkovnikova, M. S. Kukharsky, H. An, P. Dimasi, S. Alexeeva, O. Shabir, P. R. Heath, V. L. Buchman, Protective paraspeckle hyper-assembly downstream of TDP-43 loss of function in amyotrophic lateral sclerosis. *Mol Neurodegener* **13**, 1–17 (2018).

72. C. Wang, Y. Duan, G. Duan, Q. Wang, K. Zhang, X. Deng, B. Qian, J. Gu, Z. Ma, S. Zhang, L. Guo, C. Liu, Y. Fang, Stress Induces Dynamic, Cytotoxicity-Antagonizing TDP-43 Nuclear Bodies via Paraspeckle LncRNA NEAT1-Mediated Liquid-Liquid Phase Separation. *Mol Cell* **79**, 443-458.e7 (2020).

73. H. Yu, S. Lu, K. Gasior, D. Singh, S. Vazquez-Sanchez, O. Tapia, D. Toprani, M. S. Beccari, J. R. Yates, S. Da Cruz, J. M. Newby, M. Lafarga, A. S. Gladfelter, E. Villa, D. W. Cleveland, HSP70 chaperones RNA-free TDP-43 into anisotropic intranuclear liquid spherical shells. *Science (1979)* **371** (2021).

74. M. S. Hipp, P. Kasturi, F. U. Hartl, The proteostasis network and its decline in ageing. *Nature Reviews Molecular Cell Biology 2019 20:7* **20**, 421–435 (2019).

75. Y. Gwon, B. A. Maxwell, R.-M. Kolaitis, P. Zhang, H. J. Kim, J. P. Taylor, Ubiquitination of G3BP1 mediates stress granule disassembly in a context-specific manner. *Science (1979)* **372**, eabf6548 (2021).

76. B. A. Maxwell, Y. Gwon, A. Mishra, J. Peng, H. Nakamura, K. Zhang, H. J. Kim, J. P. Taylor, Ubiquitination is essential for recovery of cellular activities after heat shock. *Science (1979)* **372** (2021).

77. M. A. White, E. Kim, A. Duffy, R. Adalbert, B. U. Phillips, O. M. Peters, J. Stephenson, S. Yang, F. Massenzio, Z. Lin, S. Andrews, A. Segonds-Pichon, J. Metterville, L. M. Saksida, R. Mead, R. R. Ribchester, Y. Barhomi, T. Serre, M. P. Coleman, J. R. Fallon, T. J. Bussey, R. H. Brown, J. Sreedharan, TDP-43 gains function due to perturbed autoregulation in a Tardbp knock-in mouse model of ALS-FTD. *Nat Neurosci* **21**, 1138 (2018).

78. A. Dereeper, V. Guignon, G. Blanc, S. Audic, S. Buffet, F. Chevenet, J. F. Dufayard, S. Guindon, V. Lefort, M. Lescot, J. M. Claverie, O. Gascuel, Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* **36**, W465–W469 (2008).

79. M. Gertsenstein, L. M. J. Nutter, Engineering Point Mutant and Epitope-Tagged Alleles in Mice Using Cas9 RNA-Guided Nuclease. *Curr Protoc Mouse Biol* **8**, 28–53 (2018).

80. C. J. Miedel, J. M. Patton, A. N. Miedel, E. S. Miedel, J. M. Levenson, Assessment of Spontaneous Alternation, Novel Object Recognition and Limb Clasping in Transgenic Mouse Models of Amyloid-β and Tau Neuropathology. *JoVE (Journal of Visualized Experiments)* **2017**, e55523 (2017).

81. S. J. Guyenet, S. A. Furrer, V. M. Damian, T. D. Baughan, A. R. la Spada, G. A. Garden, A Simple Composite Phenotype Scoring System for Evaluating Mouse Models of Cerebellar Ataxia. *JoVE (Journal of Visualized Experiments)*, e1787 (2010).

753 82. S. Lê, J. Josse, F. Husson, FactoMineR: An R Package for Multivariate Analysis. *J*
754   *Stat Softw* **25**, 1–18 (2008).
755 83. K. Alboukadel, M. Fabian, Extract and Visualize the Results of Multivariate Data
756   Analyses [R package factoextra version 1.0.7] (2020). https://CRAN.R-
757   project.org/package=factoextra.
758

Acknowledgements:

Human tissue was received from the NIH NeuroBioBank at the University of Maryland,
University of Pittsburgh, and Sepulveda sites. The authors also thank all members of the
Rousseaux lab for important discussions and critical feedback on the manuscript. The authors
also thank the following Core facilities from the University of Ottawa and the Ottawa Hospital
Research Institute (OHRI) for use of their facility, equipment, and expertise: the Cell Biology
and Imaging Acquisition Core (RRID:SCR_021845), STEMCore Laboratories
(RRID:SCR_012601), Genome Engineering and Molecular Biology (GEM) Core
(RRID:SCR_022954), Animal Behaviour and Physiology Core (RRID:SCR_022882), Louise
Pelletier Histology Core (RRID: SCR_021737), Figs. 1, 2, 4, and S2 were generated in part with
Biorender.com.

Funding:

This project was supported in part by an NSERC Discovery Grant and Discovery Launch
Supplement to M.W.C.R. (RGPIN-2019-04133 and DGECR-2019-00369); the Canada Research
Chairs program to M.W.C.R; the ALS Society of Canada in partnership with the Brain Canada
Foundation through the Brain Canada Research Fund, with the financial support of Health
Canada, for financial support through the ALS Trainee Award Program 2019 (T.R.S.) and the
Discovery Grants Program 2021 (M.W.C.R.); the J.P. Bickell Medical Research Fund
(M.W.C.R.); The Eric Poulin Center for Neuromuscular Disease through the Student
Translational Research Awards (T.R.S., V.S.G.); The Canadian Institute of Health Research
through the CGS-M award (C.E.P., V.S.G.); The Ontario Graduate Scholarship Program Award
(C.E.P.); NSERC Undergraduate Summer Research Award (T.T.N.); Medical Student Summer
Research Program Award (T.T.N); Christopher Chiu senior postdoctoral fellowship (P.M.M.);
James Hunter and Family ALS Initiative (J.R.).

Author Contributions

Conceptualization: TRS, CEP, MWCR

Methodology: TRS, CEP, TTN, JLZ, VSG, PMM, BN, PMM, SMC, JLR

Investigation: TRS, CEP, TTN, JLZ, MMH, ACG, VSG, PMM, BN, JLR

Visualization: TRS, CEP, MWCR

Funding acquisition: TRS, CEP, MWCR

Project administration: JMW, JR, MWCR

Supervision: JR, MWCR

Writing – original draft: TRS, MWCR

Writing – review & editing: TRS, MWCR

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials.

Supplementary Materials

Materials and Methods
Figs. S1-S11
Tables S1-S5

Figure Captions

Fig. 1: SUMOylation dynamically regulates nuclear TDP-43 in a stress-responsive manner.
(A) Schematic of immunoprecipitation assays to detect TDP-43 SUMOylation. **(B)**
Representative SUMOylation Assay western blot detecting TDP-43 SUMOylation specifically in
response to 1 hour sodium arsenite (250 µM) stress. * = SUMOylated TDP-43, ** =
PolySUMOylated TDP-43. **(C)** Representative image and quantification of relative proximity
ligation signal between TDP-43 and SUMO2/3 in murine primary cortical neuron cultures in
response to 1 hour sodium arsenite (250 µM) treatment. Scalebar = 20 um. (N = 4 per condition)
Unpaired T-test, *** p<0.0005 **(D)** Representative GFP-Trap SUMOylation assay depicting
increased SUMOylation with ALS-causing Q331K mutation and blocked SUMOylation by
disruption of the Nuclear Localization Sequence (NLS^{mut}) in response to 250 µM sodium
arsenite stress. **(E)** Representative SUMOylation Assay of TDP-43 SUMOylation dynamics
during stress (250 µM sodium arsenite) and recovery demonstrating a dependency on the
ubiquitin proteosome for clearance of polySUMOylated TDP-43 by treatment with proteasome
inhibitor MG132 (2uM). **(F)** Representative GFP-Trap SUMOylation assay screening for
potential E3 ligases regulating TDP-43 SUMOylation in response to 1 hour sodium arsenite
stress (250 µM). (N = 3) RM One-Way ANOVA with Fisher's LSD test, * p<0.05, ** p<0.005,
*** p<0.0005, **** p<0.0001.

Fig. 2: Blocking TDP-43 SUMOylation at conserved K408 impairs the cellular stress
response. (A) Representative GFP-Trap SUMOylation assay to map the site of TDP-43
SUMOylation in response to 1 hour sodium arsenite (250 µM) stress. **(B)** Phylogenetic
representation from alignment of representative TDP-43 paralogs highlighting the emergence
and conservation of [human] K408 (Blue). **(C)** Pairwise identity of the [human] TDP-43 C-
terminal domain from MUSCLE alignment of 300 amino acid sequences for TDP-43 paralogs
from Humans to Actinopterygii from OrthoDB. Red represents the TDP-43 C-terminal domain
(Intrinsically disordered domain). Cyan represents PTM- enriched region at the extreme C-
terminus. **(D)** Representative GFP-Trap SUMOylation assay with phospho-dead TDP-43 "4SA"
(S403/404/409/410A) highlighting antagonism between SUMOylation and phosphorylation. **(E)**
Schematic and sanger sequencing of the TDP-43^{K408R} knock in mouse line. **(F)** Representative
images and quantification of G3BP1 stress granule dynamics and TDP-43 nuclear foci formation
in mouse primary cortical neurons (7 DIV) during stress (1 hour 250 µM sodium arsenite) and

recovery. Cyan arrowheads denote cells with nuclear TDP-43 foci, magenta arrow in insets

denote TDP-43 foci. G3BP1 contrast was set for optimal visualization of stress granules. Scale

Bar = 25 μm. N=5, 2-Way ANOVA with Fisher's LSD test, ** $p<0.005$. **(G)** Representative

western blot and quantification of RIPA and UREA fractions from mouse primary cortical

neurons (7 DIV) during stress (1 hour 250 μM sodium arsenite) and recovery. N=3-5, 2-Way

ANOVA with Fisher's LSD test, * $p<0.05$, ** $p<0.005$.

Fig. 3: Blocking TDP-43 SUMOylation *in vivo* leads to sex-specific and age-dependent

impairment and pathology. (A) Representative image of a barbered female mouse and

quantification of cumulative barbering probability. 2-Way ANOVA with Tukey's Multiple

Comparisons, *** $p<0.0005$. **(B)** Tube test of social dominance in female mice 16 months of age

(16MO) displaying the total head-to-head battles and win percentage by genotype comparing

Wild Type against K408R (K408R/+ and K408R/K408R animals). Binomial Test of Observed

vs. Expected with Expected due to random chance set to 50%, * $p<0.05$. **(C)** Quantification of

total distance traveled (cm) and total time in center (s) of female mice in the open field test.

Mixed Effects Analysis with Tukey's multiple comparison, *** $p<0.0005$, * $p<0.05$. **(D)**

Quantification of percent alternations in the spontaneous Y-maze test in female mice at 2, 9, and

16 months of age (2MO, 9MO, 16MO, respectively). Mixed-Effects Analysis with Tukey's

multiple comparison, * $p<0.05$. (E) Quantification of relative ambulatory activity normalized to

Wild Type at 1 hour in female mice at 2, 9, and 16 months of age (2MO, 9MO, and 16MO,

respectively). Grey shading between hours 2 and 14 indicate "lights off" with respect to

day/night light cycle. 2-Way ANOVA with Tukey's multiple comparison, * $p<0.05$. **(F)**

Representative images and quantification of TDP-43 mislocalization in the lumbar spinal cord of

9-month-old TDP-43^{K408R} mice. Each datapoint represents the average of 4 serial sections 40 μm

apart per individual mouse. Scalebar = 100 μm. (N = 8 per genotype, 4 per sex/genotype)

Unpaired t-test, *** $p<0.0005$. See fig. S10A for sex comparisons. **(G)** Representative images

and quantification of neuromuscular junction (NMJ) innervation in the tibialis anterior of 16-

month-old male TDP-43^{K408R} mice. Scale bar = 30 μm. (N = 4 per genotype) 2-Way ANOVA

with Tukey's multiple comparison analysis, * $p<0.05$, **** $p<0.0001$.

Fig. 4: SUMOylation correlates with aging and is enriched in the prefrontal cortex of

ALS/FTD patients. (A) Demographic composition of temporal lobe samples from aging cohort.

(B) Representative western blot and box plot of SUMO2/3 across binned age groups. Average

Age: "Young" = 6.32 yrs, N = 14; "Adult" = 31.11 yrs, N = 20; "Aged" = 64.00 yrs, N = 19.

Error bars indicate 10-90 percentile. Kruskall-Wallis with Uncorrected Dunn's multiple

comparisons test. * $p<0.05$, ** $p<0.005$. (C) Representative images and quantification from

proximity ligation assay (PLA) between SUMO2/3 and TDP-43 in the prefrontal cortex.

Unaffected patient ID: NP23-02, Female, 74 yrs. ALS/FTD patient ID: A14-101, Female, 67 yrs.

Scale bar = 25 μm. (N = 7-8) Unpaired T-test, * $p<0.05$.

Supplemental Figure Captions

fig. S1: Characterizing stress responsive TDP-43 SUMOylation. (A) Representative western

blot from SUMOylation assays testing whether various stressors that cause TDP-43 aggregation

induce TDP-43 SUMOylation: NaAsO2 (250 μM), D-Sorbitol (400 mM), NaCl (300 mM), Heat
Shock (42 °C). **(B)** Representative western blot from SUMOylation assays testing the selectivity
of SUMO paralogs to SUMOylate TDP-43 in response to 1 hour sodium arsenite (250 μM)
stress. **(C)** Representative images of SUMO2/3 and TDP-43 in the nucleus of mouse cortical
motor neurons (7 DIV) in unstressed and 1 hour sodium arsenite (250μM) stressed conditions.
(D) Representative control imaged from Proximity Ligation Assay (PLA) in mouse primary
cortical neurons. Scalebar = 20 μm. (N = 4). **(E)** Representative western blot and quantification
from GFP-Trap SUMOylation assays testing the kinetics of TDP-43 SUMOylation in response to
sodium arsenite stress (250 μM) and its relationship to insoluble phosphorylated TDP-43. (N =
3) **(F)** Representative dose-response assay to detect TDP-43 SUMOylation after 1 hour of
sodium arsenite stress (1 = 250 μM, 1/2 = 125 μM, 1/4 = 62.5 μM, etc.)

fig. S2: Generation of the TDP-43^{K408R} mouse line. (A) Schematic of the knock in approach to
edit the endogenous *Tardbp* to express TDP-43^{K408R}. **(B)** Schematic of LNA probe based
genotyping approach and allelic discrimination assay to genotype TDP-43^{K408R} mouse line. **(C)**
Sex and mendelian ratios of F3 generation TDP-43^{K408R} mice from K408R/+ and K408R/+
breeding pairs used to generate behavior and histology cohorts. Observed vs Expected statistical
analysis assuming an expected 1:1 ratio sex ratio and 1:2:1 mendelian ratio.

**fig. S3: Characterizing the stress response in TDP-43^{K408R} primary cortical neuron
cultures. (A)** Classification and quantification of G3BP1 stress granule (SG) intensity in
response to 1 hour sodium arsenite stress (250 μM) and recovery. (N = 5 per genotype) 2-Way
ANOVA with Tukey's multiple comparisons analysis. *p<0.05, ***p<0.0005. **(B)**
Quantification of the localization of G3BP1 stress granules (SGs) in response to 1 hour sodium
arsenite stress (250 μM) and recovery. (N = 5 per genotype) 2-Way ANOVA with Tukey's
multiple comparisons analysis. ***p<0.0005. **(C)** Quantification of the percentage of neurons
with positive TDP-43 nuclear foci in G3BP1 stress granule positive and G3BP1 stress granule
negative neurons in response to 1 hour sodium arsenite stress (250 μM) and recovery. (N = 5 per
genotype) 2-Way ANOVA with Tukey's multiple comparisons analysis. ***p<0.0005. **(D)**
Quantification of the average number of TDP-43 nuclear foci in neurons positive for TDP-43
nuclear foci in response to 1 hour sodium arsenite stress (250 μM) and recovery. (N = 5 per
genotype) 2-Way ANOVA with Tukey's multiple comparisons analysis. *p<0.05.

fig. S4: Characterization of TDP-43^{K408R} during development. (A) Genotype and sex
composition of mice subject to behavior testing. **(B)** Weight of mice at P10-P11. Unpaired T-test
(Males); Mann-Whitney test (Females and All) **(C)** Quantification of the duration hanging on a
wire at P10-P11. Mann-Whitney test (Males, Females, and All). **(D)** Survival curve for male and
female TDP-43^{K408R} mice. Curve comparisons analyzed using Log-Rank test and Gehan-
Breslow-Wilcoxon test. **(E)** Weights of male and female TDP-43^{K408R} mice from 3 weeks to 64
weeks of age. Mixed-effects analysis. **(F)** Hindlimb clasping scores of male and female TDP-
43^{K408R} mice from 3 weeks to 64 weeks of age. Mixed-effects analysis. See table S2 for raw data
details on statistical tests.

fig. S5: Characterizing the cognitive and social behavior of TDP-43^{K408R} mice.
Quantification of cognitive and social behaviors in the **(A)** Light/Dark Box, **(B)** Y-maze, **(C)** 3
Chamber Social Interaction, **(D)** Marble Burying, and **(E)** Tail Suspension behavior tasks at 2, 9,
and 16 months of age (2MO, 9MO, and 16MO, respectively) in male and female TDP-43^{K408R}
mice. All behavior tests were analyzed with a Mixed-Effects Model with Tukey's multiple
comparison analysis, see table S2.

fig. S6: Characterizing the motor behavior of TDP-43^{K408R} mice. Quantification of motor abilities in the **(A)** Hanging Wire, **(B)** Grip Strength, and **(C)** Rotarod behavior tasks at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively) in male and female TDP-43^{K408R} mice. Behavior tests were analyzed with 2-Way ANOVA or Mixed-Effects Models, see table S2.

fig. S7: Digigait analysis of TDP-43^{K408R} mice. Principle component analyses of all outputs from the Digigait behavior test for male and female TDP-43^{K408R} mice at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively).

fig. S8: Characterizing the activity of TDP-43^{K408R} mice. Quantification of **(A)** total distance traveled (cm) and time in center (s) during the Open Field behavior test and **(B)** the relative ambulatory activity (males) of TDP-43^{K408R} mice at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively). Behavior tests were analyzed with 2-Way ANOVA or Mixed-Effects models, see table S2.

fig. S9: Molecular characterization of TDP-43^{K408R} mice. (A) Representative western blot and quantification of TDP-43 and phosphorylated TDP-43 in the RIPA and UREA fractions of the cortex from male mice at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively. N = 4 per genotype) 2-Way ANOVA with Fisher's LSD test. **(B)** Representative western blot and quantification of TDP-43 and phosphorylated TDP-43 in the RIPA and UREA fractions of the cortex from male mice at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively). (N = 4 per genotype) 2-Way ANOVA with Fisher's LSD test, *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$. **(C)** Representative western blot and quantification of TDP-43 and phosphorylated TDP-43 in the RIPA and UREA fractions of the lumbar spinal cord at 2, 9, and 16 months of age (2MO, 9MO, and 16MO, respectively. N = 8 per genotype; 4 per sex/genotype) 2-Way ANOVA with Fisher's LSD test, *$p < 0.05$, ***$p < 0.0005$, ****$p < 0.0001$.

fig. S10: Histological characterization of TDP-43^{K408R} lumbar spinal cord. (A) Quantification of TDP-43 mislocalization in the lumbar spinal cord of 9-month-old from Fig. 3F split by sex. Each datapoint represents the average of 4 serial sections 40 μm apart from an individual mouse. Unpaired t-test. * = $p < 0.05$, ** = $p < 0.005$. **(B)** Representative images and quantification of ChAT+ motor neurons in the ventral horn of the lumbar spinal cord of male and female mice. Each datapoint is the average of 4 serial sections spaced 40 μm apart through the lumbar enlargement of the lumbar spinal cord. Unpaired t-test, ** $p < 0.005$. **(C)** Example images of neuromuscular junctions (NMJs) for scoring criteria to assess fully innervated, partially innervated, and fully denervated NMJs.

fig. S11: Optimization of proximity ligation assay in human tissue. Representative images of proximity ligation assay (PLA) between TDP-43 and SUMO2/3 in human PPFE hippocampus tissue sections (5um). Yellow arrows indicate TDP-43:SUMO2/3 interactions where the signal specific to the PLA channel and not identified in the autofluorescence channel. Scalebar = 18 μm.

Materials and Methods

Cloning of pEGFP-TDP-43 plasmid and variants

For expression of C-terminally tagged TDP-43-EGFP, human TDP-43 was subcloned out of a wtTDP-43tdTOMATOHA (gift from Zoushang Xu, Addgene # 28205) via XhoI and Kpn1 double digest and inserted into a pEGFP-N3 (Clontech) backbone and validated by sanger

sequencing. Site directed mutagenesis was performed using QuikChange II XL (Agilent 200521)
following manufacturer instructions. Primers for site directed mutagenesis can be found in table
S4. Several TDP-43 variants were generated through the uOttawa Genome Engineering and
Molecular Biology facility or designed and ordered through twist bioscience and were subcloned
into the pEGFP-TDP-43 backbone for consistency using XhoI and HindIII double digest, see
table S4 for plasmids used/generated in this study.

Generation of pL302-HA-SUMO2 stable HEK293T cell line

To generate a pL302-HA-SUMO2 lentivirus-compatible vector for stable HA-SUMO2
expression, HA-SUMO2 was subcloned from pcDNA-HA-SUMO2 (gift from Guy Salvesen,
Addgene #48967) and inserted into the pL302 lentiviral backbone (gift from Jacqueline Burré
and Thomas Südhof at Stanford). Briefly, extension PCR was performed to add an Xba1
restriction site to the 5' and 3' ends of the HA-SUMO2 insert using PCR amplification using
forward primer sequence: 5'- ggctgcaggtcgactctagaaagcttatggatgcctaccc-3', and reverse primer
sequence: 5'- tggctgcaggtcgactctagatgcatgctcgagtcaacct-3'. The PCR amplicon and pL302
backbone were digested with XbaI and the amplicon was ligated into the pL302 backbone.
Colonies were screened by sanger sequencing to ensure proper orientation and copy number of
the insert. Next the pL302-HA-SUMO2 plasmid was packaged into lentivirus by co-transfection
of psPAX2 (Gift from Didier Trono, Addgene #12259), and pMD2-G (Gift from Didier Trono,
Addgene #12260) into HEK293T cells and subsequent media collections. Viral media was
filtered through a 0.45 μm filter then concentrated by centrifugation at 100,000G for 2 hours at 4
°C and the pellet was resuspended in 1X PBS. Fresh HEK293T cells were transduced with
pL302-HA-SUMO2 virus to generate a stable polyclonal cell line.

GFP-trap SUMOylation assay

~250,000 HEK293T and/or pL302-HA-SUMO2 HEK293T cells were seeded (experiment
dependent) in a 6-well plate. The following day, plasmids were transfected to express TDP-43-
GFP (or mutant variants) and/or HA-SUMO variants and allowed to incubate for 48 hours for
optimal expression. Cells were treated with 250 μM NaAsO$_2$ for one hour (or variations of time,
concentration, or chemical depending on assay), after which cells were immediately collected by
scraping the bottom of the well and pelleting the samples by centrifugation at 2500G for 5 minutes
at room temperature. The supernatant was then aspirated, and the cells were flash frozen or
immediately lysed in 450 uL of ice cold 1X Denaturing Lysis buffer (1X RIPA, supplemented
with 1X protease inhibitor, 1X phosphatase inhibitor, 50mM N-Ethylmalamide, and 5% 2-
mercaptoethanol) and boiled at 95 °C for 5 minutes. After boiling, 450 uL of ice cold 1X Lysis
buffer (1X RIPA, supplemented with 1X protease inhibitor, 1X phosphatase inhibitor, 50mM N-
Ethylmalamide) was then added to the sample, and cells were left on ice for 20 minutes with
vortexing every 5 minutes for 10 second. Next, the cell lysates are centrifuged at ~21000G for 20
minutes at 4 °C and during this time, 30 uL of GFP-Trap beads (Bulldog Bio) per sample were
washed 3 times in cold 1X RIPA buffer then aliquoted equally into microcentrifuge tubes for each
sample. Following the centrifugation, 1-3% of the supernatant was transferred into a
microcentrifuge tube to be used as the *Input* sample. The remainder of the supernatant was
transferred into a microcentrifuge tube containing the GFP-Trap beads as the immunoprecipitation
sample and was then placed on a rotator for 45 minutes at 4 °C. After the incubation, the solution
was aspirated, and the beads were washed using cold 1X RIPA buffer 5 times. Samples were eluted
in 2X Laemmli buffer containing 10% Beta-Mercaptoethanol and boiled at 85 °C for 10 minutes
at 1250 rpm on a thermomixer. The input samples were prepared with 4X Laemmli buffer

containing 10% Beta-Mercaptoethanol and then boiled at 85 °C for 10 minutes. Samples were then
analyzed by Western Blot.

Endogenous SUMOylation assay

~500,000 HEK293T and/or pL302-HA-SUMO2 HEK293T cells were seeded (experiment
dependent) in a 6-well plate. The following day, cells were treated respective of their experiment
after which cells were immediately collected by scraping the bottom of the well and pelleting the
samples by centrifugation at 2500G for 5 minutes at room temperature. The supernatant was then
aspirated, and the cells were flash frozen or immediately lysed in 100 uL of 1% SDS buffer (1%
SDS in IP Lysis Buffer supplemented with 1X protease inhibitor, 1X phosphatase inhibitor,
100mM N-Ethylmalamide), vortexed for 20 seconds then boiled at 95C for 10 minutes. After
boiling, 900 µL of ice-cold IP Lysis buffer was then added to the sample, and cells were left on
ice for 20 minutes with vortexing every 5 minutes for 10 second. Next, the cell lysates are
centrifuged at ~21000G for 20 minutes at 4 C. Following the centrifugation, 1-3% of the
supernatant was transferred into a microcentrifuge tube to be used as the *Input* sample. The
remainder of the supernatant was transferred into a microcentrifuge tube containing 1 to 10 ug of
antibody per sample (See table S5 for concentrations) and incubated overnight rotating at 4 °C.
The next day, the solution was added to a magnetic rack kept at 4 °C and the supernatant was
collected for *flowthrough* to test immunodepletion or aspirated. The beads were washed 5X in ice
cold IP Lysis buffer. Samples were eluted using 5 ug of synthetic HA peptide (Sino Biological)
per 1ug of antibody and incubating at 37 °C for 15 minutes at 900rpm on a thermomixer (for anti-
HA immunoprecipitations) or by boiling in 2X Laemmli buffer containing 10% Beta-
Mercaptoethanol at 85C for 10 minutes at 1250rpm on a thermomixer (for other endogenous
immunoprecipitations). The input/flowthrough and HA immunoprecipitation samples were
prepared with 4X Laemmli buffer containing 10% Beta-Mercaptoethanol and then boiled at 85 °C
for 10 minutes. Samples were then analyzed by Western Blot using light chain specific secondary
antibodies.

Serial protein extraction from cell culture

Soluble protein was extracted using RIPA buffer (supplemented with protease inhibitor,
phosphatase inhibitor, and 50mM N-Ethylmalamide) during immunoprecipitation assays (taken
as *Input* prior to immunoprecipitation) or from lysate in cortical neuron experiments. The
remaining pellet after the centrifugation step post-lysis was washed in 1mL RIPA buffer with
vortexing for 10 seconds followed by centrifugation at ~21000G for 20 minutes at 4 °C. The
supernatant was carefully removed, and the pellet was resuspended in 300 µL of 2% SDS buffer
in PBS for immunoprecipitation assays, or in 300 µL of 8M UREA in PBS with 10mM Tris-HCl
pH 7.4. Samples were left to solubilize overnight at room temperature before 2X Laemmli was
added prior to Western Blot analysis.

Western blot analysis

Protein sample prepared in Laemmli loading buffer was loaded onto 8% polyacrylamide gel,
TGx Mini-PROTEAN 4-15% precast gel (BioRad), or Bolt Bis-Tris Plus mini-Protein Gel 4-
12% (Invitrogen) and run at 100-140 constant voltage in Tris-Glycine or MES buffer for their
respective gels. Proteins were transferred onto a 0.45 µm nitrocellulose membrane at a constant
340 mA for 2 hours at 4 °C. The membranes were blocked in 10% milk diluted in TBS-T,
washed 5 x 5 minutes in TBS-T then incubated in primary antibody overnight (table S5). The
following day the membranes were washed for 5 x 5 minutes in TBS-T, then incubated in

secondary antibody for 1-2 hours at room temperature. Finally, the secondary antibody is washed
5 x 5 minutes in TBS-T before being imaged using chemiluminescence Clarity Western ECL or
Clarity Max Western ECL on an LAS4000 (GE). Densitometry was performed using the
volumes function of the ImageLab (BioRad) software.

Conservation analysis of TDP-43 K408

1059
Representative FASTA amino acid sequences of TDP-43 paralogs were aligned for phylogenetic
analysis using MUSCLE multiple alignment and PhyML maximum likeliness for phylogenetic
tree generation(*78*). Pairwise alignment of the C-terminal domain was performed using Geneious
Prime 2022.1.1 (https://geneious.com) performing a MUSCLE multiple alignment of 300 FASTA
amino acid sequences of TDP-43 paralogs between Humans and Actinopterygii representing the
emergence of [human] K408 curated from the OrthoDB database (table S1).

Mouse lines

1067 *C57BL6/N, TDP-43^{K408R} (C57BL6/N Background)*

Sex and age of experimental mice

Experiments were performed using both male and female mice sufficiently powered to detect sex
differences. In some instances where no sex differences were observed, male and female mice
were grouped together to assess genotypic differences independent of sex. Experiments
involving primary cortical neurons cultures dissected from individual embryos were used
involving a mixture of male and female embryos. Experimental animals underwent behavioral
testing as early as P10 and as late as 16 months of age. See table S2 for raw data and statistical
analyses.

Mouse husbandry

All mouse procedures were carried out in accordance with the Canadian Council on Animal Care
and approved by the University of Ottawa Animal Care Committee. All mice were group housed
(3-5 per mice cage) with access to food and water *ad libitum* on a standard 12-hour light dark
cycle. Exceptions included adult male mice who were separated and single housed if persistent
fighting and fight wounds were observed in the cage. All experimental mice were given crinkle
paper in addition to the standard nestlet and hut enrichment material. Husbandry was completed
by University of Ottawa Animal Care and Veterinary Services except for cohorts actively
undergoing behavior testing for which husbandry was completed by the experimenter.

TDP-43^{K408R} mouse line generation

At The Centre of Phenogenomics (SickKids, Toronto, ON, Canada), the *TDP-43^{K408R}* mice were
generated on a C57BL6/N background based on previous methods (*79*) using CRISPR/Cas9-
mediated gene editing of the endogenous *Tardbp* locus. In brief, spCas9 loaded with an sgRNA
to target exon 6 of *Tardbp (MGI:2387629; 5'-TGGGGGCTTTGGCTCGAGCA-3')* was
electroporated into embryos alongside a single stranded oligonucleotide (ssODN) repair template
(5'-
C*TAAATCTACCTAACCTAATAACCAACCTACTAACCACCCCCCACCACCTACATTCCCCAG*
CCAGAAGACcTAGAATCCATGgaCGAGCCAAAGCCCCCATTAAAACCACTGCCCGATCCTG
CATTTGATGCTGACCCCCAACCAAGGGGGGC-3'). Embryos were screened via allelic
discrimination (see "TDP-43^{K408R} genotyping" below) to identify founders. Four founders were
crossed to C57BL6/N mice to ensure germline transmission of knock-in allele.

TDP-43^(K408R) genotyping

Tail samples were collected prior to weaning and again postmortem for genomic DNA (gDNA) isolation and genotyping. Tails were solubilized in 300 μL solubilization buffer (10X SET, 100 mM NaCl, 100 μg/mL Proteinase K (Bio Basic PB0451-250)) at 55 °C overnight. Cell debris was precipitated by adding 150 μL of "Tail Salts Buffer" (4.31M NaCl, 0.63M HCl, 10 mM Tris-HCl, pH7.4) then samples were centrifuged at ~21,000*g* for 15 minutes at 4 °C. Supernatant containing gDNA was transferred into 600 μL of chilled 100% ethanol to precipitate nucleic acids which were then pelleted by centrifugation for 10 minutes at ~21,000*g* at 4 °C. The supernatant was carefully removed, and the pellet was washed in 600 μL of chilled 70% ethanol then centrifuged at ~21,000*g* at 4 °C for 5 minutes. The supernatant was carefully removed, and residual supernatant was left to evaporate for 5 minutes. The gDNA pellet was resuspended in double distilled H2O at 55 °C for 10 minutes. Genotyping reaction was prepared in a 10 μL reaction containing ~5ng of gDNA, 250nM *K408R Genotyping Forward* primer (5'-*CCACCATTCTAAATCTACCTAACCTAATA-3'*), 250nM *K408R Genotyping Reverse* primer (5'-*GGATCGGGCAGTGGTTTTA-3'*), 125nM Wild Type Locked Nucleic Acid (LNA) probe (*HEX-TCT+A+A+GT+CT+TCT+GGC-IowaBlack FQ),* 125nM K408R LNA probe (*FAM-TCT+A+G+GT+CT+T+CT-IowaBlack FQ),* and 2X PerfeCTa qPCR Tough Mix (QuantaBio). Reactions were run on a BioRad CFX96 qPCR thermocycler with the following cycling parameters: Initial annealing at 95 °C for 2 minutes, followed by 40 cycles of 95 °C for 15 seconds then 60 °C for 60 seconds. Results were analyzed via allelic discrimination in the BioRad Maestro software. See fig. S2B.

Mouse wellness monitoring

Mice were monitored weekly starting at age P21 and included assessment of hindlimb clasping, kyphosis, weight, and general health. The hindlimb function was assessed following previous methods (*80*). Briefly, the mouse was suspended in the air for approximately 5-10 seconds by the base of the tail. Features of the limbs were assessed to give a score from 0-4. Kyphosis was assessed by allowing the mice to briefly walk on the flat table top in the housing room and visually observing the straightness of the spine. A score of 0-3 was given based on previously described methods(*81*). Weight was measured every other week by placing individual mice on a digital weigh scale. Notably, mice exhibiting barbered patches were monitored and recorded.

Primary cortical neuron cultures

Pregnant mice were euthanized between gestation E13.5–15.5 with 120 mg/kg Pentobarbital Sodium (Bimeda-MTC, 8015E) delivered via intraperitoneal injection. The embryos were removed and placed into ice cold 1X PBS. For each embryo, the cortices were carefully isolated and meninges removed and then placed in ice cold HBSS (Sigma Aldrich). Cortices were dissociated for 20 minutes with trypsin (Thermo Scientific) at room temperature on a rotator before trypsin inhibitor with DNase solution was added to quench the reaction. Cells were pelleted at 2,500 x *g* for 5 minutes at 4°C. The supernatant was carefully removed and the pellet was washed with trypsin inhibitor plus DNase solution. Cortical neurons were pelleted at 2,500 x *g* for 5 minutes at 4°C. The supernatant was carefully removed and the neurons were resuspended in 1 mL of ice cold outgrowth media (Neurobasal media (Thermo Scientific), supplemented with 1X B-27 (Thermo Scientific), 1X N-2 (Thermo Scientific), 500 μM L-Glutamine (Wisent Bioproducts), and 0.5% penicillin/streptomycin (GE Healthcare Life Sciences)). Culture dishes and coverslips were prepared in advanced coated with 50 μg/mL Poly-

D-Lysine overnight at 37 °C specific to each experiment detailed below. Cultures were
maintained for 7-9 days *in vitro* prior (DIV) to experimentation.

Immunofluorescence in primary cortical neurons

Micro Coverglass #1.5 coverslips (Electron Microscopy Sciences) coverslips were washed in 2M
HCl overnight at 55 °C, washed 5 times in sterile H_2O, then pre-coated with poly-D-lysine
(50 μg/mL) overnight at 37 °C, then washed with distilled water three times and air-dried at
room temperature for at least 2 hours. Primary mouse cortical neurons were seeded at 75,000-
100,000 cells per coverslip and cultured as described above. On day 7, neurons were fixed for 10
minutes using 10% phosphate buffered formalin for (Fisher Chemical, SF100-4) followed by 3 x
5-minutes washes in 1 mL of 1X PBS. Neurons were blocked in 500 μL of blocking buffer (1%
Triton X-100, 10% cosmic calf serum in 1X PBS) for 1 hour, then incubated in 300 μL of
primary antibody diluted in blocking buffer overnight at 4 °C. The following day, the neurons
were washed for 5 x 5-minutes in 1 mL of 1X PBS then incubated in 300 μL of secondary
antibody diluted in blocking buffer for 2 hours at room temperature. Neurons were then washed
in 1 mL 1X PBS, then stained with DAPI for 10 minutes at room temperature followed by 4 x 5-
minutes washes in 1X PBS before being mounted using antifade fluorescence mounting media
(Dako, S3023).

Proximity ligation assay in primary cortical neurons

Primary mouse cortical neurons were cultured and fixed as described in *primary cortical neuron*
cultures and *immunofluorescence in primary cortical neurons*. Fixed coverslips were transferred
into 12-well plates and outlined with a hydrophobic pen and blocked using 40 μL of Duolink
blocking buffer (Sigma Aldrich, DUO82007) at 37 °C for 1 hour. Coverslips were then washed
for 3 × 5 minutes in 1 mL 1X PBS. Next, coverslips were incubated in 300 μL of primary
antibody (anti-TDP-43 1:750, PTGLabs; anti-SUMO2/3 [8A2] 1:500 Sigma) diluted in blocking
buffer (1.5% Triton X-100, 10% cosmic calf serum in 1X PBS) overnight at 4 °C. The next day,
the coverslips were washed for 3 × 5 minutes in 1 mL of Duolink Wash Buffer A (0.01 M Tris-
Base, 0.15 M NaCl, 0.05% Tween-20, pH 7.4) followed by incubation at 37°C for 1 hour in
40 μL of dilute probe mixture containing a 1:5 dilution of Duolink PLA MINUS (Sigma Aldrich,
DUO82004) and PLUS probes (Sigma Aldrich, DUO82002) in antibody diluent (Sigma Aldrich,
DUO82008). Duolink PLA probes were diluted in antibody diluent (Sigma Aldrich, DUO82008)
at a 1:5 dilution. Coverslips were washed for 3 × 5 minutes in Duolink Wash Buffer A and then
incubated in 40 μL of ligase (Sigma Aldrich, DUO82027) diluted in 1X ligation buffer (Sigma
Aldrich, DUO82009) at a 1:40 dilution at 37 °C for 30 minutes. The coverslips were then
washed for 3 x 5-minutes washes in Duolink Wash Buffer A and then incubated in 40 μL of
polymerase (Sigma Aldrich, DUO82028) diluted 1:80 in 1X amplification buffer (Sigma
Aldrich, DUO82011) at 37°C for 90 minutes. Next, the coverslips were washed 2 × 10 minutes
in Duolink Wash Buffer B (0.2 M Tris-Base, 0.1 M NaCl, pH 7.5) and then again in 1 mL of
Duolink Wash Buffer B diluted at 1:100 in 1X PBS for 1 minute. Coverslips were briefly air
dried and then mounted on slides using Vectashield Antifade Mounting Medium with DAPI. Z-
stack images were obtained on a Zeiss AxioObserverZ1 LSM800 Confocal Microscope at 63×
magnification with a 5× digital zoom through a Z distance of 10-12 μm per image using optimal
0.27 μm spacing per slice with dimensions set to 512 × 512 pixels with 2X averaging per frame.
~5 images were randomly obtained around the coverslip sampling >15 cells per condition per
replicate. Images were analyzed and quantified using the Spots function on the Imaris (ver. 9.9.1
Bitplane, Switzerland) software.

Cell viability in primary cortical neurons

Mouse primary cortical neurons were collected and cultured as described in "Primary cortical neuron cultures". 20,000 cells were seeded in white opaque 96 well plates in technical quadruplicates per embryo per condition. After 7 days *in vitro* neurons were stressed with 250 μM sodium arsenite for 1 hour and recovered respectively. Cell Titer Glo 2.0 Cell Viability Assay (Promega) was performed on primary cortical neurons following manufacturer instructions including a standard curve from 0.1 nM to 100 μM ATP in each plate for standardization. Luminescence was quantified using GLOMAX plate reader and values were standardized to the luminescence values of the standard curve. The values were then normalized to the unstressed condition to quantify relative viability during stress and recovery.

Behavior testing general information

All adult behavior testing was performed in the University of Ottawa Behaviour and Physiology Core. During behavior testing periods the mice were minimally disturbed: food and water were only added/changed as needed and cages were changed by the experimenter with a transfer of old bedding material. Apart from nesting and Beam Break testing, all behavior tests were performed in the light phase of the cycle. The experimenter was blinded to the genotype of the mice during behavior testing and until the mice were end pointed. Mice were brought to the testing room to habituate in dim white light at least 30 minutes prior to commencing testing with the exceptions of nest building and fear conditioning for which there was no habituation period. The behavior cohort consisted of males and females with 12-15 animals per sex per genotype. These mice had been backcrossed three times to C57BL/6N background. All mice used for histology, biochemistry, and RNA analyses performed the fear conditioning test 1 week prior to dissections to control for potential molecular changes induced by this behavior test.

Developmental testing

A small battery of behavior tests was completed at P8 and P21. At P8, these tested included (I) weight, (II) righting reflex, (III) hindlimb suspension, and (IV) forelimb suspension. (I) Weight was measured every other week by placing individual mice on a digital weigh scale. (II) Mice were placed on their back on a tabletop and timed for how long it took them to right themselves onto their paws. (III) Mice were placed facing downward in a 50mL Falcon tube (VWR 21008-940) and timed for their latency to fall. (IV) Mice were placed to grasp a wire suspended by a pencil holder and timed for latency to fall. At P21, hanging wire was performed in which the mouse was allowed to grasp a wire 30 cm off the ground and time for its latency to fall. Each mouse completed three trials with a maximum time of 60 seconds.

Beam break

The Beam Break test was used to assess general locomotor activity and habituation to a novel cage environment. The apparatuses used to record and analyze the mice activity included the Micromax analyzer/Fusion Software (Omnitech Electronics; Columbus, OH, USA) at the 2-month time point and the Photobeam Activity System (San Diego Instruments; San Diego, CA, USA) at the 9 and 16-month timepoints. Clean cages with only a thin layer of corncob bedding, food, and water were loaded into the recording frame. Mice were singly housed in one of these cages 2 hours prior to the start of their dark cycle. For a 24-hour period at the standard 12-hour dark/light cycle, the activity of the mice was monitored by infrared beam emitters and receptors. When the test was completed, the sum of the infrared beam breaks in 5-minute, 1-hour, and 24-

1229 hour sampling bins was analyzed.

Digigait

The Digigait treadmill Imager and Analysis Software (Mouse Specifics Inc.) were used to record
and analyze parameters of gait, respectively. Mice were placed on the unmoving treadmill
surface and once recording was started the speed was increased to 18 cm/s with 0 degree incline.
A 3 second video of each mouse walking continuously, with no stopping or starting, was
captured. A mouse was excluded from the test if it was unable to walk for 3 seconds at the 18
cm/s speed. Digiait data was analyzed using FactoMinR (*82*) and visualized using FactoExtra
(*83*) packages in R (version 4.2.3).

Fear conditioning

Contextual and cued fear conditioning were tested using a 3-day protocol to assess associative
fear learning and memory. Naïve age and sex matched mice were initially used to determine the
optimal shock value. On day 1 (training), the mice were placed into a Phenotyper box (Noldus
Information Technology) with a grid shock floor (Med Associates). The testing room was set to
60 lux light level and 70 dB of white noise (context A). The mice were left in the apparatus for 6
minutes during which they receive 3 tone-shock pairings (30 seconds tone co-terminated with a 2
second foot shock). On day 2 (Context), the mice were placed in the same apparatus (Context A)
with no tone or foot shock delivered for the 6-minute trial. On day 3 (cue), the animals were
placed in the test apparatus with an altered context (context B) including red light, no white
noise, vanilla scent, textured mat covering the shock floor and plastic inserts in the apparatus.
The mice are allowed to explore this context with no tone for 3 minutes, and then are presented
with the same tone from Day 1 for the last 3 minutes. The time freezing was analyzed with
EthoVision software for all 3 testing days.

Grip strength

A grip strength meter (Chatillon DFE II, Columbus Instruments) was used to assess the maximal
forelimb grip strength of the mice. The grip strength meter was rotated vertically and temporarily
mounted on a flat surface prior to testing. The mouse was brought near the triangular attachment
and allowed to grasp the lower bar with its forepaws. The mouse was pulled directly downward,
by the tail, in one smooth motion. Each mouse was tested, and the grip strength recorded for 5
consecutive trials with a 1-minute intertrial interval. If a trial was deemed unsuccessful by the
blinded experimenter, the trial was redone. Causes of an unsuccessful trial included the mouse
prematurely losing its grip on the bar or grasping the bar with its hindfoot. The average grip
strength for the 5 successful trials was analyzed.

Hanging wire

The hanging wire test was used to assess muscle strength and coordination. A metal wire was
secured to the top of a tall plastic box with padding on the bottom. The mouse was brought near
and allowed to grasp the wire with its forepaws. A timer was started once the experimenter
released the mouse to allow it to hang freely. When the mouse fell from the wire the time was
recorded. Mice had three trials and were allowed to hang for a maximum of 600 seconds with a
60 second intertrial interval in between. If a mouse hung for 600 seconds, it did not complete any
additional trials. The maximum hanging time from the trials was used for analysis.

Light/dark box

The light dark paradigm was used to assess anxiety-like or disinhibited, exploratory behavior.
The testing apparatus (Med Associates) has two equal sized rectangular compartments the mouse
can move freely between. The one is fully illuminated whereas the other is covered by a black
plastic insert. The position of the mouse in the apparatus testing field is recorded using infrared
beams. To start the test, the mice are placed into the lit side and allowed to explore for 10
minutes. The time spent in each compartment and number of entries into each compartment was
recorded and analyzed by Activity Monitor software (Med Associates).

Marble burying

The marble burying assay can assess the motor function required to bury objects as well as
cognitive changes related to apathy or perseverance. Cages were filled with 10cm of fresh
corncob bedding and one mouse was placed to habituate for 5 minutes. After habituation, 20
glass marbles were laid out evenly in a 4 by 5 pattern. The mouse was returned to the cage and
left alone for 30 minutes in 60 lux light. With the completion of the trial, the number of marbles
buried by at least two thirds was scored by a blinded experimenter.

Nest building

Nest building is an innate behavior of mice in their daily lives. This complex behavior requires
both executive planning and sensorimotor coordination. Directly following Beam Break testing,
a single square nestlet (5 cm^2 cotton pad) was placed in each Beam Break cage for 16 – 18 hours,
of which 12 of these hours was during the dark cycle. At the end of the test, images were taken
of the nests and scored blindly for quality as previously described on a scale from 1-5.

Open field

The open field test was used to assess anxiety and locomotor activity in a novel environment.
The apparatus consists of white plastic square arenas measuring 45cm on each side. The mice
were placed in a corner of the arena and allowed to freely explore for 10 minutes with light
levels at 300 lux. The distance travelled, time spent in the corners and center of the field was
recorded and analyzed by EthoVision software (Noldus Information Technology).

Rotarod

The rotarod (IITC Life Science, Ugo Basile) was used to test the motor performance of mice
including coordination and resistance to fatigue. Mice were placed in the stationary rotarod bar
for 10 seconds before the rotarod program was initiated. The bar accelerated from 4 rpm to 40
rpm for 5 minutes and the latency to fall for each mouse was recorded. The time was stopped
when the mouse fell from the bar or rotated passively. Mice were completed four trials per day,
with a 10-minute intertrial interval in their home cage, for 3 consecutive days.

Spontaneous Y-maze

The spontaneous Y-Maze test was used to assess spatial working memory. The Y-shaped maze
has three identical arms at 120°C around a center point triangle. The mice were placed in the
center point and allowed to freely explore the arms for 8 minutes. The movement of the mouse
was tracked including the sequence of arm entries by EthoVision (Noldus Information
Technology). An alternation is defined as the mouse making consecutive, sequential entries into
each of the three arms without revisiting an arm. The alternation index was calculated as
(number of alternations/(total number of arm entries minus two)) and reported as a percent.

Tail suspension

The automated Tail Suspension apparatus (Med Associates) was used to assess apathetic-like behaviors of the mice. Mice were taped and suspended by the tail to a vertical steel bar which measures strain gauge as the mouse moves during a 6-minute trial. The cumulative time spent immobile, hanging passively (below lower threshold) was measured by the Tail Suspension software.

Three chamber sociability

The Three Chamber test was conducted to measure the sociability of mice when allowed to interact with a novel mouse or a similarly sized inanimate object. The testing apparatus is a 19 x 45 cm plastic box divided into three equal chambers with clear plastic wall dividers. The two external chambers each have a single weighted metallic mesh pencil holder and the central chamber is empty. The mice are habituated to the apparatus for 5 minutes by being placed in the central chamber and allowed to freely explore and enter all chambers. The mouse has a 5-minute intertrial interval in its home cage. In the test trial, a sex and age matched wild-type mouse (social target) is placed beneath one mesh pencil holder, and an inanimate plastic toy (non-social target) is placed beneath the other. To commence the test trial, the mouse is paced in the central chamber. The time spent in each chamber and interacting with the social or non-social target is recorded and analyzed by EthoVision software (Noldus Information Technology).

Tube test of social dominance

Tube test was used to assess social dominance and within cage social hierarchy. Cage mates were paired against each other for testing in a round-robin design. The tube test was conducted on a flat tabletop, which the mice were allowed to run around on for 5 minutes prior to commencing the testing. One foot of vinyl tubing was used. Mice were habituated to the tube prior to testing on the same day by encouraging them to run through the empty tube from either side 5 times. A blinded experimenter placed a mouse on either end of the tube, and released their tales when they completely entered the tube. The first mouse to step its hind paws out of the tube lost the battle. The battle was redone if after 2 minutes no mouse had won.

Collection and preparation of mouse tissue

Mice were harvested for both histology and biochemistry/molecular analyses to reduce total animal numbers. Mice were first acclimated for 2 hours prior to dissection to help control for activity-induced changes and all dissections were performed at zeitgeber hours ZT6 to ZT11 to control for potential circadian affects. Mice were randomly dissected by cage to limit batch effects and additional stress. Mice were anesthetized using isoflurane inhalation, blood was collected, and then mice were sacrificed via decapitation and the brains were quickly isolated, weighted, then cut into 2 hemispheres where the left hemisphere was drop fixed in 10% buffered formalin and the right hemisphere was dissected by region and flash frozen on dry ice. Spinal cords were removed using hydraulic extrusion and the lumbar spinal cord was isolated. The lumbar enlargement was isolated for histology whereas the remaining lumbar spinal cord was used for molecular analysis. Tibialis anterior and soleus muscles were isolated and drop fixed in 10% buffered formalin for imaging studies and gastrocnemius muscle was flash frozen.

Immunofluorescence analysis in mouse tissue

Tissue was collected as described in *Collection and preparation of mouse tissue*. Following fixation for 48 hours, spinal cord samples were transferred to a 70% ethanol solution and sent to the Louise Pelletier Histology Core facility at the University of Ottawa for paraffin embedding

and microtome sectioning. The samples were sectioned at a thickness of 5 µm. Serial sectioning
was performed at four levels within the tissue each separated by 40 µm. For staining, slide-
mounted sections were deparaffinized in 100% xylenes (Fisher Scientific X3P-1GAL) for 10
minutes, then transferred into a second container of fresh 100% xylenes for an additional 10
minutes. The slides were then rehydrated in descending ethanol solutions: two 5 minutes
treatments in 100% ethanol, 5 minutes in 70% ethanol, and 5 minutes in 50% ethanol. The slides
were then immersed in 1X PBS for at least 5 minutes to rinse residual ethanol. Antigen was
performed by placing slides in 1X sodium citrate buffer (2.94g sodium citrate 0.5mL Tween-20
in 1L 1X PBS, pH6) at 95°C for 30 minutes then rinsed in 1X PBS. Tissue sections were then
blocked in blocking buffer containing 1% Triton-X and serum (5% horse serum, 5% cosmic calf
serum, or 1% bovine serum albumin) in 1X PBS for 2 hours at room temperature. Next the tissue
sections were incubated with primary antibody (table S5) overnight at 4°C. The next day, the
sections were washed twice in PBS 1X + 0.1% Triton-X for 5 minutes each and then 3X in 1X
PBS for 5 minutes each. The slides were then incubated in secondary antibody at room
temperature for 2 hours. The slides were then washed in 1X PBS + 0.1% Triton-X for 5 x 5-
minutes, before mountin with #1.5 coverslips (Thermo Fisher Scientific 12-544E) and
fluorescent mounting media (Dako S3023). The coverslips were sealed with clear nail polish and
allowed to dry before imaging. Images of each ventral horn were taken on a Zeiss Axio Imager 2
at 20X magnification. Cells were counted manually per image using Fiji (ImageJ).

Neuromuscular junction analysis

Tissue was collected as described in *Collection and preparation of mouse tissue*. After 24 hours
of fixation, tibialis anterior muscles were stored in a solution of 1X PBS. Muscle bundles were
carefully teased apart from the tissue into 4-5 bundles with forceps. The muscle bundles were
transferred to 6-well plate cell culture plate containing 1% Triton-X in 1X PBS overnight at 4°C
under gentile agitation for permeabilization. The next day, the muscle bundles were washed three
times in 1X PBS for 5 minutes at room temperature. They were placed in a blocking buffer
solution of 4% bovine serum albumin and 1% Triton-X in 1X PBS at 4°C overnight. The
following day, the muscles were incubated in primary antibody (BTX and NfL/SV2 cocktail, see
table S5) diluted in blocking buffer solution overnight at 4°C. The next day, the muscle bundles
were washed with 1X PBS three times for 5 minutes and incubated with secondary antibody
diluted in blocking buffer solution at room temperature for 2 hours. The muscle bundles were
then washed 3 times in 1X PBS for 5 minutes. The bundles were then placed on slides, mounted
with VECTASHIELD® HardSet™ Antifade Mounting Medium (#H-1400), and #1.5 coverslips
placed above the muscle bundles. The slides were left at room temperature overnight in the dark
to allow the mounting media to cure. Images were taken on a Zeiss LSM800 Confocal
Microscope at 20X through a Z-stack with optimal spacing (0.61 µm). Images were taken
throughout the entirety of the muscle tissue such that at least 80 neuromuscular junctions per
animal were captures. Images were scored manually per image using Fiji (ImageJ). See fig.
S10C for scoring examples.

SUMOylation analysis in human brain tissue

20 mg of frozen human brain tissue was homogenized in 300 µL of RIPA buffer supplemented
with protease inhibitor, phosphatase inhibitor, and 50mM N-Ethylmalamide and vortexed every
5 minutes for 20 minutes while incubating on ice. The samples were centrifuged at 21000G for
20 minutes at 4°C and the supernatant (RIPA soluble fraction) was carefully removed. 4X
laemmli buffer was added and samples were boiled for 5 minutes at 95 °C then stored overnight

at 4 °C. The remaining pellet after the centrifugation step post-lysis was washed in 1mL RIPA
buffer with vortexing for 10 seconds followed by centrifugation at ~21000G for 20 minutes at 4
°C. The supernatant was carefully removed, and the pellet was resuspended in 300 μL of of 8M
UREA in PBS with 10mM Tris-HCl pH 7.4. Samples were vortexed and left to solubilize
overnight at room temperature before 4X Laemmli was added prior to Western Blot analysis.
RIPA and UREA samples were randomly loaded onto gels (separated by fraction) with 4 control
samples consistently loaded (aged 20-30) on each gel for normalization. Western blot was
performed as described above. Samples were standardized to ponceau signal then normalized to
control samples.

Collection and preparation of FFPE human brain tissue

Frontal cortex tissues were collected antemortem from three sporadic ALS/FTLD patients (a 74-
year-old male, 67-year-old female and 59-year-old female) obtained through the ALS Clinic at
Sunnybrook Health Sciences Centre, Toronto. ALS was diagnosed using the revised El Escorial
Criteria (Brooks et al., 2000) and informed consent was obtained with approval from the local
ethical review board. The presence of TDP-43 proteinopathy within the frontal cortex was
verified through immunohistochemical labeling using rat anti-phosphorylated TDP-43
(p409/410) antibody on formalin fixed paraffin embedded sections. Genetic analyses confirmed
absence of mutations in key ALS/FTD-associated genes: *C9orf72*, *SOD1*, *FUS* and *TARDBP*.

Proximity ligation assay in human FFPE tissue

Formalin fixed paraffin-embedded human brain sections (5 μm sections) were deparaffinized
using two 10-minute washes with xylenes, two 5-minute washes with 100% ethanol, one 5-
minute wash with 70% ethanol, and one 5-minute was with 50% ethanol. Deparaffinized sections
then underwent sodium citrate antigen retrieval (10mM sodium citrate, 0.05% Tween-20, pH 6)
for 2 hours at 80°C. Sections were then blocked with Duolink blocking buffer (Sigma Aldrich,
DUO82007) for 1 hour at room temperature. Next, sections were incubated with primary
antibodies (anti-TDP-43 1:750, PTGLabs; anti-SUMO2/3 [8A2] 1:500 Sigma) diluted in
blocking buffer (1.5% Triton X-100, 10% cosmic calf serum in 1X PBS) overnight at 4°C. The
following day, the sections were washed in Duolink Wash Buffer A (0.01M Tris-Base, 0.15M
NaCl, 0.05% Tween-20, pH 7.4), followed by incubation in Duolink PLA MINUS (Sigma
Aldrich, DUO82004) and PLUS probes (Sigma Aldrich, DUO82002) at 37°C for 1 hour.
Duolink PLA probes were diluted in antibody diluent (Sigma Aldrich, DUO82008) at a 1:5
dilution. Sections were washed in Duolink Wash Buffer A and then incubated in ligase (Sigma
Aldrich, DUO82027) at 37°C for 30 minutes. Ligase was diluted in 1X ligation buffer (Sigma
Aldrich, DUO82009) at a 1:40 dilution. Then, the sections were washed in Duolink Wash Buffer
A and then incubated in polymerase (Sigma Aldrich, DUO82028) at 37°C for 90 minutes.
Polymerase was diluted in 1X amplification buffer (Sigma Aldrich, DUO82011) at a 1:80
solution. The sections were then washed in Duolink Wash Buffer B (0.2M Tris-Base, 0.1M
NaCl, pH 7.5) and then again in Duolink Wash Buffer B diluted at 1:1000. Finally, sections were
incubated with DAPI at room temperature for 15 minutes. DAPI was diluted at 1:1000 in PBS.
Fluorescent mounting media (Dako, S3023) and #1.5 coverslips were then placed over the
sections and sealed using clear nail polish. Images were taken on a Zeiss LSM800 Confocal
Microscope. 5 images per tissue were randomly collected from grey matter at 63X magnification
and 2X digital zoom through a Z stack of 10-12 μm with optimal spacing (0.27 μm) at a
resolution of 512 x 512 pixels. PLA foci were manually counted per image using Fiji (ImageJ).

Fig. 1: SUMOylation dynamically regulates nuclear TDP-43 in a stress-responsive manner.

Fig. 2: Blocking TDP-43 SUMOylation at conserved K408 impairs the cellular stress response.

Fig. 3: Blocking TDP-43 SUMOylation *in vivo* leads to sex-specific and age-dependent impairment and pathology.

Fig. 4: SUMOylation correlates with aging and is enriched in the prefrontal cortex of ALS/FTD patients.