A single-nucleus RNA sequencing atlas of the postnatal retina of the shark *Scyliorhinus canicula*

Authors

Nicolás Vidal-Vázquez[1,2,*], Ismael Hernández-Núñez[1,*], Pablo Carballo-Pacoret[1,3], Sarah Salisbury[3], Paula R. Villamayor[3,4], Francisca Hervas-Sotomayor[5,6], Xuefei Yuan[5], Francesco Lamanna[5], Céline Schneider[5], Julia Schmidt[5], Sylvie Mazan[7], Henrik Kaessmann[5], Fátima Adrio[1], Diego Robledo[3,8], Antón Barreiro-Iglesias[1,2], Eva Candal[1,2]

Affiliations
1. Departamento de Bioloxía Funcional, Facultade de Bioloxía, Universidade de Santiago de Compostela, 15782, Santiago de Compostela, Spain
2. Aquatic One Health Research Center (ARCUS), Universidade de Santiago de Compostela, 15782, Santiago de Compostela, Spain
3. The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh EH25 9RG, UK
4. Departamento de Zooloxía, Xenética e Antropoloxía Física, Facultade de Veterinaria, Universidade de Santiago de Compostela, 27002, Lugo, Spain
5. Center for Molecular Biology (ZMBH), DKFZ-ZMBH Alliance, Heidelberg University, Heidelberg, Germany
6. Current affiliation: INRAE, LPGP, Rennes, France
7. CNRS, Sorbonne Université, Biologie Intégrative des Organismes Marins, UMR7232-BIOM, Banyuls-sur-Mer, France
8. Departamento de Zooloxía, Xenética e Antropoloxía Física, Facultade de Bioloxía, Universidade de Santiago de Compostela, 15782, Santiago de Compostela, Spain
* These authors contributed equally

corresponding author: Eva Candal (eva.candal@usc.es)

Abstract

The retina, whose basic cellular structure is highly conserved across vertebrates, constitutes an accessible system for studying the central nervous system. In recent years, single-cell RNA-sequencing studies have uncovered cellular diversity in the retina of a variety of species, providing new insights on retinal evolution and development. However, similar data in cartilaginous fishes, the sister group to all other extant jawed vertebrates, are still lacking. Here, we present a single-nucleus RNA-sequencing atlas of the postnatal retina of the catshark *Scyliorhinus canicula*, consisting of the expression profiles for 17,438 individual cells from three female, juvenile catshark specimens. Unsupervised clustering revealed 22 distinct cell types comprising all major retinal cell classes, as well as retinal progenitor cells (whose presence reflects the persistence of proliferative activity in postnatal stages in sharks) and oligodendrocytes. Thus, our dataset serves as a foundation for further studies on the development and function of the catshark retina. Moreover, integration of our atlas with data from other species will allow for a better understanding of vertebrate retinal evolution.

Background & Summary

The neural retina shows a remarkable degree of conservation in its cellular structure in all extant vertebrates[1–3]. This basic plan consists of five neuronal cell classes and one glial cell class, whose somata are arrayed in three nuclear layers, interspersed with two plexiform layers, where synapses occur (**Fig. 1a**). The outer nuclear layer (ONL) contains photoreceptors (PRs), the light-sensitive cells of the retina, which can usually be classified in two morphologically and functionally distinct major types, cones and rods, responsible for photopic and scotopic vision, respectively. In turn, the inner nuclear layer (INL) hosts three types of interneurons, namely horizontal cells (HCs), bipolar cells (BCs) and amacrine cells (ACs), which receive, integrate, modulate and transmit the signals coming from photoreceptors to retinal ganglion cells (RGCs). RGCs are located in the ganglion cell layer (GCL) and project their axons through the optic nerve to the visual processing centres of the brain. Besides, the retina contains a major glial cell class, Müller glia (MG), a type of radial glial cells whose nuclei are located within the INL. Other glial cell types, such as oligodendrocytes, microglia or astrocytes, may be present in the innermost layers of the retina, but, unlike MG, these are not derived from the optic cup; instead, they originate in other parts of the brain and migrate into the eye through the optic nerve[4].

Most of these cell classes can be subdivided into a number of cell types with diverse morphological, functional and molecular properties[5–7]. In recent years, high-throughput single-cell RNA sequencing (scRNA-seq) studies have provided unprecedented insights into molecular cell type diversity in the mature and developing vertebrate retina[6,8–10]. Although most studies have focused on the retina of mammals[11–32] (particularly that of rodents[12–18] and primates[19–32]), cellular atlases of the chick[13,33], brown anole lizard[11], zebrafish[9,13,34–40] and sea lamprey[11,41] retinas have also been generated to date, providing new resources to study the evolution and development of vertebrate retinal cell types. However, scRNA-seq studies on the retina of cartilaginous fishes are still lacking.

Given their phylogenetic position as the sister group to all other extant gnathostomes (**Fig. 1b**), chondrichthyans (sharks, rays and chimaeras) constitute a particularly interesting group to study retinal evolution[42–44]. Among cartilaginous fishes, the shark *Scyliorhinus canicula* (Linnaeus, 1758), also known as the small-spotted catshark or the lesser-spotted dogfish, stands as a suitable model for experimental studies, owing to its abundance, relatively small size and accessible oviparous development[43]. Furthermore, the catshark retina shows persistent cell proliferation in postnatal stages[45,46] (similar to other fishes[47]), making *S. canicula* an interesting species for the study of postnatal retinal neurogenesis.

In this study, we generated a single-nucleus RNA-sequencing (snRNA-seq) atlas of the postnatal retina of *S. canicula* (**Fig. 1c**). Early juvenile specimens were chosen to allow profiling of both mature and progenitor cell types, since previous work has shown that cell proliferation decreases as the animal grows, with mitotic activity being virtually absent in sexually mature adults[45,48]. Thus, we used three retinas from three female juvenile catsharks to generate a dataset consisting of 17,438 high quality nuclei. Unsupervised clustering revealed 22 cell types representing all major retinal cell classes, as well as retinal progenitor cells and oligodendrocytes. This constitutes, to the best of our knowledge, the first single-cell transcriptomic study of the retina of a chondrichthyan, providing a groundwork for comparative studies on the evolution of both retinal cell type diversity and retinal neurogenesis.

Methods

Animals
Three female, juvenile specimens of *S. canicula*, with a total length of 10.5 to 11.1 cm (**Table 1**) were kindly provided by the Aquarium Finisterrae in A Coruña (Spain) and kept in artificial

99   seawater tanks under standard conditions of temperature (15-16 °C), pH (7.5-8.5) and salinity
100   (35 g/L). All procedures were performed in accordance with the guidelines for animal
101   experimentation established by the European Union and the Spanish government and were
102   approved by the Bioethics Committee of the University of Santiago de Compostela (license
103   number 15004/2022/001).
104
Retina sampling
Animals were deeply anaesthetised with 0.5% tricaine methanesulfonate (MS-222; Thermo
Fisher Scientific, 118000500) in seawater. The animals were removed from water, the eyes
enucleated, and the retinas (one from each specimen, n = 3) were dissected out under a
stereomicroscope. Retinas were immediately put in Eppendorf tubes and frozen in liquid
nitrogen. Retinal samples were kept at -80 °C until they were further processed for nuclei
isolation and RNA sequencing.
112
Nuclei isolation and snRNA-seq data generation
Retina nuclei were extracted following a published protocol[49] with small modifications. The
frozen retinas were homogenised using a micropestle in 400 µL ice-cold homogenisation
buffer (250 mM sucrose, 25 mM KCl, 5 mM $MgCl_2$, 10 mM Tris-HCl [pH 8], 0.1% IGEPAL, 1 µM
dithiothreitol [DTT], 0.4 U/µL Murine RNase Inhibitor [New England BioLabs, M0314L], and 0.2
U/µL SUPERase-In [Ambion, AM2694]). The homogenates were triturated gently using a p1000
tip for 10 times, incubated on ice for 5 min and then centrifuged at 100 g for 1 min at 4 °C to
pellet any unlysed tissue chunks. The supernatant was transferred into another 1.5 mL
Eppendorf tube and centrifuged at 400 g for 4 min at 4 °C to collect nuclei. The nuclei were
washed twice in 400 µL homogenisation buffer and strained using a 40 µm Flowmi strainer
(Sigma, BAH136800040) during the second wash step to remove nuclei aggregates. The final
nuclei pellet was resuspended in 30-50 µL Nuclei Buffer (10x Genomics, PN-2000207). To
estimate the nuclei concentration, nuclei aliquots were diluted in phosphate-buffered-saline
(PBS) with Hoechst and propidium iodide (PI) DNA dyes and counted on Countless II FL
Automated Cell Counter (Thermo Fisher Scientific, RRID: SCR_020236). Around 15,000 nuclei
were used as input for the snRNA-seq experiment. The Chromium Next GEM Single Cell 3'
Reagent Kits v3.1 (PN-1000121, PN-1000120, and PN-1000213) were used to make snRNA-seq
libraries. Libraries were quantified on a Qubit Fluorometer (Thermo Fisher Scientific; RRID:
SCR_018095) and quality checked on a Fragment Analyzer (Agilent; RRID: SCR_019417).
Libraries were sequenced on NextSeq550 (Illumina; RRID: SCR_016381; 28 cycles for Read 1,
56 cycles for Read 2, 8 cycles for i7 index).
134
Genome indexing and read alignment
Genome indexing and library mapping was performed with STAR (v2.7.10a)[50]. The *S. canicula*
genome   assembly   sScyCan 1.1[51]   (GCF_902713615.1;
GCF_902713615.1_sScyCan1.1_genomic.fna) and its associated annotation in GFF format
(*GCF_902713615.1_sScyCan1.1_genomic.gff*) were downloaded from the National Center for
Biotechnology Information (NCBI). The mitochondria (NC_001950.1) annotations within the
GFF file were manually edited to convert "CDS" annotations to "exon" annotations and to
convert all annotations of "tRNA" and "rRNA" to "gene" annotation. This was done to ensure
mitochondrial gene annotations were consistent with those of the nuclear genes as STAR
assigns transcripts to "exon" annotations in the .gff file. The GFF annotation file was converted
to GTF format using gffread (v0.10.1)[52]. The genome and its annotation (GTF) were indexed
using STAR (--runMode genomeGenerate). Each library was then mapped against the genome
with the 10x V3 cell barcode whitelist (*3M-february-2018.txt*) and using standard parameters
for single cell libraries (--soloMultiMappers Unique --soloBarcodeReadLength 28 --soloType
CB_UMI_Simple --soloUMIlen 12 -- soloCBwhitelist 3M-february-2018.txt --soloFeatures
GeneFull --clipAdapterType CellRanger4 --outFilterScoreMin 20 --soloCBmatchWLtype

1MM_multi_Nbase_pseudocounts   --soloUMIfiltering   MultiGeneUMI_CR   --soloUMIdedup
1MM_CR --readFilesCommand zcat --outSAMtype BAM Unsorted). The raw (unfiltered) files
(*genes.tsv*, *barcodes.tsv*, and *matrix.mtx*) generated for each sample were then used for
downstream analysis. On average, there were 274 million reads per sample with 94% of reads
with valid barcodes, and a 45% saturation. A summary of the STAR output statistics for each
sample can be found in **Table 2**.
157
Bioinformatic quality control
Samples were then analysed in an R (v4) environment using Seurat (v4.3.0)[53]. We created
Seurat objects for each library after removing nuclei with less than 200 features and features
occurring in fewer than three nuclei. Nuclei where mitochondrial (mtDNA) features accounted
for 10% or more of their total unique molecular identifiers (UMIs) were removed before
removing all mtDNA features. After sub-setting the Seurat object into individual samples,
upper and lower thresholds for UMI and feature counts per nuclei were then applied
individually to each sample based on knee plot visualisation. In sample 1, nuclei with more
than 1,000 but less than 20,000 UMIs and more than 500 but less than 5,500 features were
retained. In sample 2, nuclei with more than 500 but less than 24,000 UMIs and more than
500 but less than 6,500 features were retained. Finally, in sample 3, nuclei with more than 750
but less than 20,000 UMIs and more than 300 but less than 7,000 features were retained.
Samples were then merged into a single Seurat object before splitting samples again into
individual sample datasets. This was done to ensure that the same features were considered
across samples. Counts were then normalised for each sample using the "NormalizeData"
function prior to calculating cell cycle scores using the "CellCycleScoring" function (see
Supplementary Table 1 for list of genes used). The "v2" SCTransform version with the
glmGamPoi method (v1.9.0)[54] was used to normalise RNA counts for each sample, regressing
out scores for the S and G2M cell cycle stages. Linear dimension reduction was conducted for
each sample using the "RunPCA" function with 50 PCs. After consulting Elbowplots for each
sample, a Uniform Manifold Approximation and Projection (UMAP) using 20 principal
components (PCs) was run for each sample and the "FindNeighbours" function was applied
using 20 PCs, before using the "FindClusters" function with a resolution of 0.5. DoubletFinder
(v2.0.3)[55] was then applied independently to each sample selecting pK values with the highest
associated mean-variance normalised bimodality coefficient (BCmvn) value. We assumed a 4%
doublet formation rate (based on the Chromium instrument specifications) and adjusted for
homotypic doublets.
185
Clustering and differential gene expression analyses
Samples were integrated using 2000 features and anchors that were identified with the "rpca"
reduction method and the "FindIntegrationAnchors" function. A principal component analysis
(PCA) was rerun on the integrated dataset using 50 PCs, and 30 PCs were used for subsequent
UMAP generation and clustering with a resolution of 0.5. Markers for each cluster were
assessed using the logistic regression method and the FindAllMarkers function on the "SCT"
assay and "data" slot, using sample ID as a latent variable to help reduce batch effects among
samples. We used a pseudocount of 0.001, set a p-value threshold of 0.01, and only considered
genes that were upregulated, expressed in at least 25% of all nuclei (in either of the compared
groups), and demonstrated a threshold of 0.25 X difference (log-scale) between the two
compared groups. Three clusters that were mostly composed of cells from a single sample and
did not show differential expression of typical retinal cell marker genes were removed (see
below; **Supplementary Table 2** shows differentially expressed genes in the removed clusters),
and all remaining cells were re-clustered using the same parameters. The same differential
expression analysis was applied to various groups of clusters (**Table 3**) to identify marker genes
for cell classes composed of several clusters.
202

Cluster annotation

To assign a cell class identity to each cluster, we elaborated a list of marker genes reported in previous studies on other vertebrate species[11–15,18–21,23,25–31,33,35,41]. Putative orthologues in the *S. canicula* genome were identified by comparing the sequence of the mouse gene (genome assembly GRCm39, GCF_000001635.27)[56] with the *S. canicula* genome assembly sScyCan1.1 (GCF_902713615.1)[51] using Blastn[57]. For marker genes absent from the mouse genome, either the human (genome assembly GRCh38.p14, GCF_000001405.40)[58] or zebrafish (genome assembly GRCz11, GCF_000002035.6)[59] genes were used. All marker gene sequences were obtained from GenBank[60]. Marker genes present in the *S. canicula* genome where then visualised in our dataset using the "FeaturePlot" and "DotPlot" functions of Seurat. This, coupled with the examination of the top differentially expressed genes in each cluster (see above), allowed us to assign all clusters in the final object to known retinal cell classes (see Technical Validation).

Gene nomenclature

For the sake of readability, we decided to change the symbols of genes annotated with a LOC number in the main text and figures of the present article. We replaced LOC numbers for symbols based on the identification of their protein products, adding an "-*l*" at the end for those annotated as "-like", except for uncharacterised genes, which were left with their LOC number. **Supplementary Table 3** shows the correspondence between the symbols used in the text and the sScyCan1.1 annotation, which is the one used in the dataset.

It should also be noted that many *S. canicula* genes follow the nomenclature used in zebrafish, including a letter or number at the end of their symbol. Since teleosts experienced an additional whole genome duplication after diverging from cartilaginous fishes[61], many gene paralogues present in zebrafish are absent from the catshark genome. Therefore, current gene symbols do not accurately reflect the existence of paralogues in *S. canicula*, nor their correspondence to specific zebrafish paralogues.

Data Records

Raw sequencing data for each sample (*.fastq* files) have been uploaded to the NCBI Sequence Read Archive (SRA) under BioProject accession number PRJNA1056918[62]. For each sample, STAR raw output files, i.e. expression matrices for each gene in each cell (*matrix.mtx*), barcodes (*barcodes.tsv*) and genes (*features.tsv*), have been uploaded to Figshare (doi: XXX). The final annotated Seurat object (*Scyliorhinus_retina.rds*) is also available on Figshare.

Two *.xlsx* files containing the list of differentially expressed genes in each cluster and each cell class (see **Table 3**) have been uploaded to Figshare. For each gene, the p-value ("p_val"), adjusted p-value ("p_val_adj"), average $\log_2$ fold change ("avg_log2FC"), the percentage of cells expressing that gene in the present cluster ("pct.1") and the percentage of cells expressing said gene in the rest of the dataset ("pct.2") are provided.

Technical Validation

Quality control

Retinas from three similarly sized, female, juvenile *Scyliorhinus canicula* specimens (**Table 1**) were used to generate the snRNA-seq dataset. To ensure that all the obtained barcodes correspond to viable nuclei, we established selection criteria based on feature counts, UMI counts and expression of mitochondrial genes (**Fig. 2a, b**). Prior to filtering, the three samples showed a similar number of nuclei (**Table 4**), features and UMIs, with the proportion of mitochondrial transcripts being higher in sample 1 and lower in sample 3 (**Fig. 2a**). As expected, there was a positive correlation between the number of detected genes and the number of UMIs, whereas a negative correlation was observed between the expression levels

of mitochondrial genes and both gene counts and UMI counts, with no major differences
between samples (**Fig. 2c**). Quality control filtering was performed by removing nuclei with
mitochondrial transcripts representing more than 10% of the total counts, and upper and
lower thresholds for UMI and number of unique genes were defined for each sample (see
Methods). After filtering, initial clustering analyses revealed three clusters that were mostly
composed of cells from a single sample that did not show expression of typical retinal cell
marker genes (**Supplementary Table 2**). Since these nuclei most likely correspond to either
low quality cells or non-retinal cells, they were removed from the dataset. The final object was
composed of 17,438 cells containing 23,489 features (unique genes). There were no major
differences in the number of unique genes, UMI count (number of transcripts) or
mitochondrial feature levels between samples (**Fig. 2b, d**). However, sample 3 yielded a higher
number of cells (**Table 4**), which can be attributed to minor technical differences in retinal
dissection and/or library preparation.
268
Cluster annotation
Unsupervised clustering of filtered cells revealed 22 clusters (**Fig. 3a**), all of which were present
in generally similar proportions in the three samples (**Fig. 3b**), with cells from the three retinas
showing similar distributions across the UMAP (**Fig. 3c**), thus confirming the similarity of the
samples. To annotate the clusters, we analysed the expression levels of known marker genes
from other species (see Methods) for the major retinal cell classes or subclasses (**Fig. 4a, b**).
This allowed us to divide cells into the six major retinal cell classes (PRs, HCs, BCs, ACs, RGCs
and MG), as well as to identify two clusters of retinal progenitor cells (RPCs) and a single cluster
of oligodendrocytes (OLs) (**Fig. 3a**).
Three clusters could be identified as rod PRs by the expression of genes such as *rho*, *pdc-l* or
pde6a[63]. Expression of established cone PR marker genes (not shown) was either present in
rod clusters or virtually non-existing. This is in accordance with previous observations that *S.*
canicula has either scarce cones[64] or a pure-rod retina[46,65] (reviewed in Ferreiro-Galve *et al.*[46]),
similar to some other benthic sharks[65,66] and the skate[67]. A fourth cluster that did not show
high levels of most rod markers but showed significant expression of *nr2e3* and *neurod1*,
known markers of rod precursors/progenitors[68], was thus annotated as PR precursor cells.
Since rod PRs usually comprise a single cell type[33,38,69] (with some exceptions, such as those of
the amphibian retina[70]), the four PR clusters identified here could correspond to different
states of PR differentiation, but further research is needed to confirm this.
Regarding retinal interneurons, a single cluster of HCs (expressing the canonical marker
onecut3a[33,71]) and 3 clusters of BCs (expressing *vsx1* and *vsx2*[11,14,72], as well as high levels of
isl1 in one of the BC clusters, confirming previous immunohistochemical results[73]) could be
identified in our dataset. We also found eight clusters of ACs, most of which showed high levels
of *pax6*, confirming previous results[74]. The AC clusters could be further divided into three
subclasses according to the expression of small molecule neurotransmitter-associated genes.
Thus, we found four clusters of GABAergic ACs expressing *gad1* and/or *gad2*[18,75]; a single
cluster of cholinergic (starburst) ACs, which show high levels of *chata* (similar to those
described in *Squalus*[76]); and three clusters of glycinergic ACs, which show high levels of glycine
transporters *slc6a5* and *slc6a9*[18]. Furthermore, two clusters of RGCs (expressing canonical
markers, such as *nefl-l*[15]) could be recognised.
MG were represented by a single cluster with high levels of *rlbp1b*[11,77] and *glul* (the gene
encoding glutamine synthetase, whose expression in *S. canicula* MG has previously been
described by means of immunohistochemistry[73,78]). Two clusters of RPCs could also be
identified by the expression of cell proliferation markers (*mki67-l*[19,79], *pcna*[45,46,78]) and typical
markers of retinal progenitors (*btg2-l*, *rx3*[12]). Finally, a small cluster of OLs was identified by
high levels of *olig2-l* and *mbp* expression[13,33], probably corresponding to cells from the optic
nerve head or the optic nerve fibres layer, where myelinated fibres are present in other
elasmobranchs[42]. A remarkable absence from our dataset are microglial cells, which are known

to be present in the innermost retinal layers of the postnatal retina of *S. canicula*[80]. This is
most likely due to microglia being lost in the nuclei dissociation or quality control filtering
processes.
To further validate cluster annotation and find novel markers specific to the *S. canicula* retinal
cell classes, we identified marker genes (differential gene expression analyses vs the remaining
cells in the dataset), both for individual clusters and for clusters grouped into
classes/subclasses (**Table 3**). **Fig. 4c** shows the top marker genes for each class-level group,
including both established and novel marker genes. For a complete list, see Data Records.
Overall, the expression of marker genes confirms that our atlas comprises all major cell classes
expected in the juvenile catshark retina, although some low abundant cell types (e.g.
microglia) could be absent from the dataset. Thus, this dataset will provide a groundwork for
studies on cell type diversity in the shark retina, allowing a better understanding of vertebrate
retinal evolution and development.
320
Usage Notes
321
322
All the code used to analyse the dataset is available in GitHub (see Code Availability). Raw
.fastq files are standard Illumina sequencing files for 10x Genomics single-cell RNA sequencing
libraries, and as such they can be processed using any typical single-cell analysis software
(STAR[50], as in our work, or Cell Ranger[81], Kallisto[82], Alevin[83], etc.). STAR output files for each
sample can be loaded into Seurat using the "ReadSTARsolo" function, and then merged into a
single Seurat object using the "merge" function. Finally, the *Scyliorhinus_retina.rds* file is a
Seurat object with the processed dataset that can be loaded into R with "ReadRDS".
330
Code Availability
331
332
333　The code used to process the raw sequencing files and generate all the results presented in
334　this study can be found in https://github.com/Roslin-Aquaculture/SHARK_retina.
335

Acknowledgements
336
337
This work was supported by grant ED431C 2021/18 funded by Xunta de Galicia to E.C., and
grant PID2020-115121GB-I00 funded by MICIU/AEI/10.13039/501100011033 to A.B.-I. N.V.-V.
was supported by grant FPU21/03076 funded by Ministerio de Ciencia, Innovación y
Universidades. D.R. was supported by the Axencia Galega the Innovación (GAIN, Xunta de
Galicia) as part of the Oportunius programme, and by BBSRC Institute Strategic Grants to the
Roslin Institute (BBS/E/20002172, BBS/E/D/30002275, BBS/E/D/10002070 and
BBS/E/RL/230002A). S.S. gratefully acknowledges an NSERC PDF award. This project has
received funding from the European Research Council (ERC) under the European Union's
Horizon 2020 research and innovation programme (VerteBrain to H.K., grant agreement no.
101019268).

348
Author contributions
349
350
Conceptualisation: N.V.-V., I.H.-N., A.B.-I., E.C. Data acquisition: I.H.-N., P.C.-P., S.S., P.R.V.,
F.H.-S., X.Y., C.S., J.S., S.M. (provided early access to the catshark genome), H.K. (supervision),
D.R. Data analysis: N.V.-V., P.C.-P., S.S., P.R.V., F.L., D.R., E.C. Data interpretation: N.V.-V., P.C.-
P., A.B.I., E.C. Writing (original draft): N.V.-V., P.C.-P., S.S., D.R., A.B.-I., E.C. Writing (review and
editing): N.V.-V., I.H.-N., P.C.-P., S.S., P.R.V., S.M., H.K., F.A., D.R., A.B.-I., E.C. Funding
acquisition: H.K., D.R., A.B.-I., E.C. Project supervision: E.C.
357

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. Background and experimental workflow. (**a**) Structure of the juvenile catshark retina, based on microscopy images kindly provided by Dr J. Francisco-Morcillo; cell drawings based on Neumayer[64]. (**b**) Cladogram showing relationships among major extant vertebrate groups, highlighting the position of chondrichthyans as the sister group to all other gnathostomes. Silhouettes from PhyloPic[84]. (**c**) Experimental workflow. Eye diagram after Collin[42]. GCL, ganglion cell layer; INL, inner nuclear layer; IPL, inner plexiform layer; ONL, outer nuclear layer; OPL, outer plexiform layer; OS, photoreceptor outer segments.

Figure 2. Quality control for snRNA-seq data. (**a, b**) Violin plots showing: left, the number of detected unique genes ("nFeature_RNA"); middle, the number of unique molecular identifiers (UMIs, "nCount_RNA"); and right, percentage of mitochondrial genes ("percent.mt") in each cell from each sample before (**a**) and after (**b**) quality control. Note that all mitochondrial features were removed from the analyses after quality control (see Methods). (**c**) Scatter plots showing the relationship between gene counts and UMI counts (left), UMI counts and percentage of mitochondrial genes (middle), together with gene counts and percentage of mitochondrial genes (right) in the three samples before quality control. (**d**) Gene counts (left), UMI counts (middle) and percentage of mitochondrial genes (right) mapped onto the UMAP.

Figure 3. Clustering of snRNA-seq data reveals cell heterogeneity in the juvenile catshark retina. (**a**) UMAP of retinal cells showing their unbiased assignment to the 22 clusters identified in this study. For abbreviations, see **Table 3**. (**b**) Barplots showing the proportions of each cluster for each sample. (**c**) Separate UMAPs of cells from each sample, coloured according to their assigned identities.

Figure 4. Expression of marker genes used for cluster annotation. (**a**) Dot plot showing expression levels of marker genes selected for identification of retinal cell classes (cluster annotation). (**b**) UMAPs showing expression of selected canonical marker genes. (**c**) Dot plot showing expression of the top three differentially expressed genes from class-level groupings (see text and **Table 3** for details). See Methods and **Supplementary Table 3** for information on gene nomenclature.

Tables

Sample	Total length (cm)	Sex
Sample 1	11.0	Female
Sample 2	10.5	Female
Sample 3	11.1	Female

Table 1. Sample information.

Sample	Sample 1	Sample 2	Sample 3
Number of reads	229,861,905	249,344,842	343,563,083
Reads with valid barcodes	0.960049	0.932741	0.917902
Sequencing saturation	0.502799	0.521001	0.338812
Q30 bases in CB + UMI	0.957978	0.95938	0.961753
Q30 bases in RNA read	0.89873	0.90096	0.888598
Reads mapped to genome: unique + multiple	0.88851	0.846251	0.647372
Reads mapped to genome: unique	0.781012	0.711432	0.518868
Reads mapped to GeneFull: unique + multiple GeneFull	0.693977	0.471452	0.354986
Reads mapped to GeneFull: unique GeneFull	0.642748	0.424422	0.314171
Estimated number of cells	3,028	1,074	2,521
Unique reads in cells mapped to GeneFull	62,372,962	34,713,678	39,214,416
Fraction of unique reads in cells	0.422171	0.328021	0.363307
Mean reads per cell	20,598	32,321	15,555
Median reads per cell	15,254	21,413	11,331
UMIs in cells	30,760,018	16,433,368	25,334,077
Mean UMIs per cell	10,158	15,301	10,049
Median UMIs per cell	7,593	10,144	7,340
Mean GeneFull per cell	3,508	4,872	3,988
Median GeneFull per cell	3,170	4,285	3,540
Total GeneFull Detected	23,271	23,076	23,449

Table 2. Summary statistics for STAR outputs for each sample. Note that raw files were used for downstream analysis (which includes all barcodes, not just those estimated to be viable cells by STAR). Therefore, the statistics relating to STAR-assigned cells are not necessarily relevant but are provided for the reader's interest.

Group	Abbreviation	Clusters
Rod photoreceptors	PR_Rod	PR_Rod1, PR_Rod2, PR_Rod3
Photoroceptor precursors	PR_Precursor	PR_Precursor
Horizontal cells	HC	HC
Bipolar cells	BC	BC1, BC2, BC3
GABAergic amacrine cells	AC_GABA	AC_GABA1, AC_GABA2, AC_GABA3, AC_GABA4
Cholinergic amacrine cells	AC_ACh	AC_ACh
Glycinergic amacrine cells	AC_Gly	AC_Gly1, AC_Gly2, AC_Gly3
Retinal ganglion cells	RGC	RGC1, RGC2
Müller glia	MG	MG
Retinal progenitor cells	RPC	RPC1, RPC2
Oligodendrocytes	OL	OL

404 **Table 3.** Class/subclass-level cluster groups for differential expression analysis.
405

Sample	Sample 1	Sample 2	Sample 3	Total
Unfiltered nuclei	69,723	52,932	64,138	186,793
Filtered nuclei	4,869	4,130	8,439	17,438
Filtered features	22,472	22,458	23,014	23,489
Mean UMIs per cell	4,527	3,716	3,486	3,831
Median UMIs per cell	3,333	2,234	2,359	2,572
Mean features per cell	2,153	1,885	1,852	1,944
Median features per cell	1,892	1,434	1,502	1604
Mean percent of mitochondrial features	3.55	2.66	2.13	2.65
Median percent of mitochondrial features	3.06	2.05	1.64	2.05

406 **Table 4.** Final cell quantification and sequencing statistics. "Unfiltered nuclei" refers to the
407 number of barcodes after importing the STAR files to Seurat, excluding cells with less than 200
408 UMIs. All other parameters refer to the final object.
409
410 **Supplementary Table 1.** Genes used in cell cycle scoring.
411 **Supplementary Table 2.** Differentially expressed genes from removed clusters.
412 **Supplementary Table 3.** Gene nomenclature.
413

References

1. Baden, T. Vertebrate vision: lessons from non-model species. *Semin. Cell Dev. Biol.* **106**, 1–4 (2020).

2. Kolb, H., Nelson, R., Ahnelt, P. & Cuenca, N. Cellular organization of the vertebrate retina. *Prog. Brain Res.* **131**, 3–26 (2001).

3. Lamb, T. D. Evolution of phototransduction, vertebrate photoreceptors and retina. *Prog. Retin. Eye Res.* **36**, 52–119 (2013).

4. Vecino, E., Rodriguez, F. D., Ruzafa, N., Pereiro, X. & Sharma, S. C. Glia–neuron interactions in the mammalian retina. *Prog. Retin. Eye Res.* **51**, 1–40 (2016).

5. Goetz, J. *et al.* Unified classification of mouse retinal ganglion cells using function, morphology, and gene expression. *Cell Rep.* **40**, 111040 (2022).

6. Shekhar, K. & Sanes, J. R. Generating and using transcriptomically based retinal cell atlases. *Annu. Rev. Vis. Sci.* **7**, 43–72 (2021).

7. Zeng, H. & Sanes, J. R. Neuronal cell-type classification: challenges, opportunities and the path forward. *Nat. Rev. Neurosci.* **18**, 530–546 (2017).

8. Shiau, F., Ruzycki, P. A. & Clark, B. S. A single-cell guide to retinal development: cell fate decisions of multipotent retinal progenitors in scRNA-seq. *Dev. Biol.* **478**, 41–58 (2021).

9. Vöcking, O. & Famulski, J. K. Single cell transcriptome analyses of the developing zebrafish eye— perspectives and applications. *Front. Cell Dev. Biol.* **11**, 1213382 (2023).

10. Zhang, X., Leavey, P., Appel, H., Makrides, N. & Blackshaw, S. Molecular mechanisms controlling vertebrate retinal patterning, neurogenesis, and cell fate specification. *Trends Genet.* **39**, 736–757 (2023).

11. Hahn, J. *et al.* Evolution of neuronal cell classes and types in the vertebrate retina. *Nature* **624**, 415–424 (2023).

12. Clark, B. S. *et al.* Single-cell RNA-seq analysis of retinal development identifies NFI factors as regulating mitotic exit and late-born cell specification. *Neuron* **102**, 1111-1126.e5 (2019).

13. Hoang, T. *et al.* Gene regulatory networks controlling vertebrate retinal regeneration. *Science* **370**, eabb8598 (2020).

14. Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).

15. Rheaume, B. A. *et al.* Single cell transcriptome profiling of retinal ganglion cells identifies cellular subtypes. *Nat. Commun.* **9**, 2759 (2018).

16. Shekhar, K. *et al.* Comprehensive classification of retinal bipolar neurons by single-cell transcriptomics. *Cell* **166**, 1308-1323.e30 (2016).

17. Tran, N. M. *et al.* Single-cell profiles of retinal ganglion cells differing in resilience to injury reveal neuroprotective genes. *Neuron* **104**, 1039-1055.e12 (2019).

18. Yan, W. *et al.* Mouse retinal cell atlas: molecular identification of over sixty amacrine cell types. *J. Neurosci.* **40**, 5177–5195 (2020).

19. Cowan, C. S. *et al.* Cell types of the human retina and its organoids at single-cell resolution. *Cell* **182**, 1623-1640.e34 (2020).

20. Gautam, P. *et al.* Multi-species single-cell transcriptomic analysis of ocular compartment regulons. *Nat. Commun.* **12**, 5675 (2021).

21. Hu, F. *et al.* Single-cell RNA-seq reveals the cellular diversity and developmental characteristics of the retinas of an infant and a young child. *Front. Cell Dev. Biol.* **10**, 803466 (2022).

22. Hu, Y. *et al.* Dissecting the transcriptome landscape of the human fetal neural retina and retinal pigment epithelium by single-cell RNA-seq analysis. *PLoS Biol.* **17**, e3000365 (2019).

23. Liang, Q. *et al.* Single-nuclei RNA-seq on human retinal tissue provides improved transcriptome profiling. *Nat. Commun.* **10**, 5743 (2019).

24. Lu, Y. *et al.* Single-cell analysis of human retina identifies evolutionarily conserved and
species-specific mechanisms controlling development. *Dev. Cell* **53**, 473-491.e9 (2020).
25. Lukowski, S. W. *et al.* A single-cell transcriptome atlas of the adult human retina. *EMBO*
J. **38**, e100811 (2019).
26. Lyu, Y. *et al.* Implication of specific retinal cell-type involvement and gene expression
changes in AMD progression using integrative analysis of single-cell and bulk RNA-seq
profiling. *Sci. Rep.* **11**, 15612 (2021).
27. Menon, M. *et al.* Single-cell transcriptomic atlas of the human retina identifies cell
types associated with age-related macular degeneration. *Nat. Commun.* **10**, 4902
(2019).
28. Orozco, L. D. *et al.* Integration of eQTL and a single-cell atlas in the human eye identifies
causal genes for age-related macular degeneration. *Cell Rep.* **30**, 1246-1259.e6 (2020).
29. Peng, Y.-R. *et al.* Molecular classification and comparative taxonomics of foveal and
peripheral cells in primate retina. *Cell* **176**, 1222-1237.e22 (2019).
30. Voigt, A. P. *et al.* Molecular characterization of foveal versus peripheral human retina
by single-cell RNA sequencing. *Exp. Eye Res.* **184**, 234–242 (2019).
31. Yan, W. *et al.* Cell atlas of the human fovea and peripheral retina. *Sci. Rep.* **10**, 9802
(2020).
32. Zhang, L. *et al.* Evolutionary and developmental specialization of foveal cell types in the
marmoset. Preprint at https://biorxiv.org/content/10.1101/2023.12.10.570996v1
(2023).
33. Yamagata, M., Yan, W. & Sanes, J. R. A cell atlas of the chick retina based on single-cell
transcriptomics. *eLife* **10**, e63907 (2021).
34. Celotto, L. *et al.* Single cell RNA sequencing unravels the transcriptional network
underlying zebrafish retina regeneration. *eLife* **12**, RP86507 (2023).
35. Emmerich, K. *et al.* Molecular regulation of retinal regeneration is context specific.
Preprint at https://biorxiv.org/content/10.1101/2023.11.20.567904v1 (2023).
36. Kölsch, Y. *et al.* Molecular classification of zebrafish retinal ganglion cells links genes to
cell types to behavior. *Neuron* **109**, 645-662.e9 (2021).
37. Lyu, P. *et al.* Common and divergent gene regulatory networks control injury-induced
and developmental neurogenesis in zebrafish retina. *Nat. Commun.* **14**, 8477 (2023).
38. Ogawa, Y. & Corbo, J. C. Partitioning of gene expression among zebrafish photoreceptor
subtypes. *Sci. Rep.* **11**, 17340 (2021).
39. Santhanam, A., Shihabeddin, E., Wei, H., Wu, J. & O'Brien, J. Molecular basis of retinal
remodeling in a zebrafish model of retinitis pigmentosa. *Cell. Mol. Life Sci.* **80**, 362
(2023).
40. Xu, B. *et al.* Unifying developmental programs for embryonic and post-embryonic
neurogenesis in the zebrafish retina. *Development* **147**, dev185660 (2020).
41. Wang, J. *et al.* Molecular characterization of the sea lamprey retina illuminates the
evolutionary origin of retinal cell types. Preprint at
https://biorxiv.org/content/10.1101/2023.12.10.571000v1 (2023).
42. Collin, S. P. Scene through the eyes of an apex predator: a comparative analysis of the
shark visual system. *Clin. Exp. Optom.* **101**, 624–640 (2018).
43. Coolen, M. *et al.* The dogfish *Scyliorhinus canicula*: a reference in jawed vertebrates.
Cold Spring Harb. Protoc. **3**, pdb.emo111 (2008).
44. Hart, N. S. Vision in sharks and rays: opsin diversity and colour vision. *Semin. Cell Dev.*
Biol. **106**, 12–19 (2020).
45. Hernández-Núñez, I. *et al.* Loss of active neurogenesis in the adult shark retina. *Front.*
Cell Dev. Biol. **9**, 628721 (2021).
46. Ferreiro-Galve, S., Rodríguez-Moldes, I., Anadón, R. & Candal, E. Patterns of cell
proliferation and rod photoreceptor differentiation in shark retinas. *J. Chem.*
Neuroanat. **39**, 1–14 (2010).

47.   Stenkamp, D. L. Neurogenesis in the fish retina. *Int. Rev. Cytol.* **259**, 173–224 (2007).

48.   Rodríguez-Arrizabalaga, M., Hernández-Núñez, I., Candal, E. & Barreiro-Iglesias, A. Use
of vivo-morpholinos for gene knockdown in the postnatal shark retina. *Exp. Eye Res.*
226, 109333 (2023).

49.   Krishnaswami, S. R. *et al.* Using single nuclei for RNA-seq to capture the transcriptome
of postmortem neurons. *Nat. Protoc.* **11**, 499–524 (2016).

50.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

51.   Genome assembly sScyCan1.1. *NCBI Genome assemby database*
https://identifiers.org/refseq.gcf:GCF_902713615.1 (2020).

52.   Pertea, G. & Pertea, M. GFF Utilities: GffRead and GffCompare. *F1000Res.* **9**, 304
(2020).

53.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).

54.   Ahlmann-Eltze, C. & Huber, W. glmGamPoi: fitting Gamma-Poisson generalized linear
models on single cell count data. *Bioinformatics* **36**, 5701–5702 (2021).

55.   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: doublet detection in
single-cell RNA sequencing data using artificial nearest neighbors. *Cell Syst.* **8**, 329-
337.e4 (2019).

56.   Genome assembly GRCm39. *NCBI Genome assembly database*
https://identifiers.org/refseq.gcf:GCF_000001635.27 (2020).

57.   Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).

58.   Genome assembly GRCh38.p14. *NCBI Genome assembly database*
https://identifiers.org/refseq.gcf:GCF_000001405.40 (2022).

59.   Genome assembly GRCz11. *NCBI Genome assembly database*
https://identifiers.org/refseq.gcf:GCF_000002035.6 (2017).

60.   Benson, D. A. *et al.* GenBank. *Nucleic Acids Res.* **41**, D36–D42 (2012).

61.   Glasauer, S. M. K. & Neuhauss, S. C. F. Whole-genome duplication in teleost fishes and
its evolutionary consequences. *Mol. Genet. Genomics* **289**, 1045–1060 (2014).

62.   *Scyliorhinus canicula* retina single-nuclei RNA sequencing. *NCBI Sequence Read Archive*
https://identifiers.org/insdc.sra:SRP480047 (2023).

63.   Lamb, T. D. Evolution of the genes mediating phototransduction in rod and cone
photoreceptors. *Prog. Retin. Eye Res.* **76**, 100823 (2020).

64.   Neumayer, L. Der feinere Bau der Selachier-Retina. *Arch. Mikr. Anat.* **48**, 83–111 (1896).

65.   Bozzano, A., Murgia, R., Vallerga, S., Hirano J & Archer, S. The photoreceptor system in
the retinae of two dogfishes, *Scyliorhinus canicula* and *Galeus melastomus*: possible
relationship with depth distribution and predatory lifestyle. *J. Fish Biol.* **59**, 1258–1278
(2001).

66.   Schieber, N. L., Collin, S. P. & Hart, N. S. Comparative retinal anatomy in four species of
elasmobranch. *J. Morphol.* **273**, 423–440 (2012).

67.   Dowling, J. E. & Ripps, H. On the duplex nature of the skate retina. *J. Exp. Zool.* **256**, 55–
65 (1990).

68.   Nelson, S. M., Frey, R. A., Wardwell, S. L. & Stenkamp, D. L. The developmental
sequence of gene expression within the rod photoreceptor lineage in embryonic
zebrafish. *Dev. Dyn.* **237**, 2903–2917 (2008).

69.   Baden, T. & Osorio, D. The retinal basis of vertebrate color vision. *Annu. Rev. Vis. Sci.* **5**,
177–200 (2019).

70.   Yovanovich, C. A. M. *et al.* The dual rod system of amphibians supports colour
discrimination at the absolute visual threshold. *Philos. Trans. R. Soc. B-Biol. Sci.* **372**,
20160066 (2017).

71. Kreplova, M. *et al.* Dose-dependent regulation of horizontal cell fate by Onecut family
of transcription factors. *PLoS One* **15**, e0237403 (2020).
72. Passini, M. A., Levine, E. M., Canger, A. K., Raymond, P. A. & Schechter, N. Vsx-1 and
Vsx-2: differential expression of two Paired-like homeobox genes during zebrafish and
goldfish retinogenesis. *J. Comp. Neurol.* **388**, 495–505 (1997).
73. Bejarano-Escobar, R. *et al.* Retinal histogenesis and cell differentiation in an
elasmobranch species, the small-spotted catshark *Scyliorhinus canicula*. *J. Anat.* **220**,
318–335 (2012).
74. Ferreiro-Galve, S., Rodríguez-Moldes, I. & Candal, E. Pax6 expression during
retinogenesis in sharks: comparison with markers of cell proliferation and neuronal
differentiation. *J. Exp. Zool. B Mol. Dev. Evol.* **318**, 91–108 (2012).
75. Ferreiro-Galve, S., Candal, E., Carrera, I., Anadón, R. & Rodríguez-Moldes, I. Early
development of GABAergic cells of the retina in sharks: an immunohistochemical study
with GABA and GAD antibodies. *J. Chem. Neuroanat.* **36**, 6–16 (2008).
76. Brandon, C. Cholinergic amacrine neurons of the dogfish retina. *Vis. Neurosci.* **6**, 553–
562 (1991).
77. Saari, J. C. & Crabb, J. W. Focus on molecules: cellular retinaldehyde-binding protein
(CRALBP). *Exp. Eye Res.* **81**, 245–246 (2005).
78. Sánchez-Farías, N. & Candal, E. Identification of radial glia progenitors in the developing
and adult retina of sharks. *Front. Neuroanat.* **10**, 65 (2016).
79. Kee, N., Sivalingam, S., Boonstra, R. & Wojtowicz, J. M. The utility of Ki-67 and BrdU as
proliferative markers of adult neurogenesis. *J. Neurosci. Methods* **115**, 97–105 (2002).
80. Bejarano-Escobar, R., Blasco, M., Durán, A. C., Martín-Partido, G. & Francisco-Morcillo,
J. Chronotopographical distribution patterns of cell death and of lectin-positive
macrophages/microglial cells during the visual system ontogeny of the small-spotted
catshark *Scyliorhinus canicula*. *J. Anat.* **223**, 171–184 (2013).
81. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nat. Commun. **8**, 14049 (2017).
82. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).
83. Srivastava, A., Malik, L., Smith, T., Sudbery, I. & Patro, R. Alevin efficiently estimates
accurate gene abundances from dscRNA-seq data. *Genome Biol.* **20**, 65 (2019).
84. Keesey, M. T. *PhyloPic* https://www.phylopic.org/ (2024).

a

OS
ONL
OPL
INL
IPL
GCL

b

Cyclostomi

Chondrichthyes

Actinopterygii

Actinistia

Dipnoi

Lissamphibia

Sauropsida

Mammalia

Amniota

Tetrapoda

Rhipidistia

Sarcopterygii

Osteichthyes

Gnathostomata

Vertebrata

c

Retinal dissection → Nuclei extraction → Nuclei capture and barcoding → cDNA library preparation → Sequencing → Analysis