1   **Chromatin accessibility variation provides insights into**
2   **missing regulation underlying immune-mediated diseases**
3

Raehoon Jeong[1,2] and Martha L. Bulyk[1,2,3†]

[1] Division of Genetics, Department of Medicine, Brigham and Women's Hospital and Harvard
Medical School, Boston, MA 02115, USA.

[2] Bioinformatics and Integrative Genomics Graduate Program, Harvard University, Cambridge,
MA 02138, USA.

[3] Department of Pathology, Brigham and Women's Hospital and Harvard Medical School,
Boston, MA 02115, USA.

† Correspondence: mlbulyk@genetics.med.harvard.edu

Abstract

Most genetic loci associated with complex traits and diseases through genome-wide association
studies (GWAS) are noncoding, suggesting that the causal variants likely have gene regulatory
effects. However, only a small number of loci have been linked to expression quantitative trait loci
(eQTLs) detected currently. To better understand the potential reasons for many trait-associated
loci lacking eQTL colocalization, we investigated whether chromatin accessibility QTLs (caQTLs)
in lymphoblastoid cell lines (LCLs) explain immune-mediated disease associations that eQTLs in
LCLs did not. The power to detect caQTLs was greater than that of eQTLs and was less affected
by the distance from the transcription start site of the associated gene. Meta-analyzing LCL eQTL
data to increase the sample size to over a thousand led to additional loci with eQTL colocalization,
demonstrating that insufficient statistical power is still likely to be a factor. Moreover, further
eQTL colocalization loci were uncovered by surveying eQTLs of other immune cell types.
Altogether, insufficient power and context-specificity of eQTLs both contribute to the 'missing
regulation.'

Introduction

More than a decade of genome-wide association studies (GWAS) has revealed several properties of the genetic architecture of complex traits and diseases[1,2]. Most (~93%) of the genetic associations are detected in the noncoding portion of the genome[3], and disease heritability is concentrated in putative regulatory regions[4]. Expression quantitative trait loci (eQTLs), which are loci associated with gene expression levels, are enriched for trait associations[5,6]. Complex traits are also characterized by their extreme polygenicity, where individual genetic association has only a small effect on the trait[7]. Altogether, these observations have led to a prevalent theory that causal genetic variants affect regulation of key genes across the genome, where each gene explains a modest proportion of trait variation[8]. There are experimental strategies aimed at nominating putative causal genes at noncoding GWAS loci[9–11]. As an alternate approach, an eQTL signal colocalizing with the GWAS signal illustrates the effect of the causal variant on gene expression and suggests that the affected gene contributes to the trait. Detection of disease-associated eQTLs thus can identify putative disease genes, helping to elucidate disease mechanisms and develop therapeutics targeting them[12].

Although it has become expected that eQTLs will be discovered in most noncoding GWAS loci, only a minority of trait-associated loci have been explained by eQTLs[13–16]. The Genotype-Tissue Expression (GTEx) eQTL study across 49 human tissues recognized that, for a typical complex trait, about 20% of GWAS loci contained a colocalized eQTL in the *cis* region (*i.e.*, 1 Mb) around the gene (*i.e.*, *cis*-eQTL)[14]. Even when focusing just on putatively causal genes, the rate of colocalization was very low (8%)[15]. Furthermore, the proportion of trait heritability mediated by *cis*-eQTLs (h^2_{med}/h^2_{SNP}) of assayed gene expression was estimated to be only about 11% on average[16]. We will call the missing link between genetic association to traits and regulatory function of the associated noncoding variants as 'missing regulation', as Connally and colleagues introduced[15]. To be able to detect eQTLs in the unexplained disease-associated loci, a better understanding of the possible reasons for why they have been missing is essential.

There are many possible explanations for why disease-associated loci are missing colocalized eQTLs[15,17–19] and for why h^2_{med}/h^2_{SNP} estimates for eQTLs are relatively low[16]. First, statistical power to detect disease-associated eQTLs may be insufficient[19]. For example, negative selection against gene expression variation may lead to challenges in detecting eQTLs for trait-relevant genes[18,20]. Since disease genes are likely to be dosage sensitive, there would be selection

against their having large eQTL effects[20]. Consequently, only weak eQTL variants, often in

regions distal to the gene's promoter[21], may remain and affect traits. In fact, eQTL-mediated

heritability was enriched in genes showing mutational constraint and those with lower *cis*-

heritability[16]. These weaker eQTL effects would require larger sample sizes to be detected with

statistical significance and to show colocalization[19]. Second, causal eQTL effects may be specific

to cell types that have not been assayed[17]. For example, immune cell eQTLs are highly cell-type

specific, and eQTL effects specific to some immune cell types may mediate immune disease risk[22].

Specificity of eQTL effects can also be limited to specific cell states[23,24]. Detecting cell-type or

cell-state specific eQTL effects requires the necessary gene expression data sets from the relevant

cell types and states[17], which has been a limiting resource for such analyses.

To investigate why disease-associated eQTL signals have been missing, we focused on

immune-mediate diseases (IMDs) as a model set of complex traits. We aimed to collect IMD-

associated loci that are expected to show eQTL signals in some cell type. Since active regulatory

elements coordinate target gene expression[25], we reasoned that variants that affect chromatin

phenotypes at regulatory elements, such as transcription factor (TF) binding[26–28] and chromatin

accessibility[29,30], have the potential to impact gene expression[31]. These chromatin phenotypes may

show detectable genetic effects even when an eQTL effect in the same cell type was not identified

in the locus[32]. For example, only about 20% of lymphoblastoid cell lines' (LCLs') PU.1 binding

QTLs (bQTLs) that colocalized with blood cell traits' association showed an eQTL effect for a

nearby gene in LCLs[33].

Here, we analyzed genetic and functional genomic (*i.e.*, ATAC-seq and RNA-seq) data in

LCLs. LCLs are derived from B lymphocytes, and their *cis*-regulatory elements were enriched for

variants associated with some IMDs[34,35]. We evaluated whether chromatin accessibility QTLs

(caQTLs) in LCLs potentially explain IMD associations using mediated heritability analysis[16] and

colocalization[36,37]. Then, we searched for disease-associated loci that were significant caQTLs, but

not eQTLs.

We examined whether the various potential reasons for missing eQTLs can account for

IMD-associated loci that are explained by caQTLs but not eQTLs. First, we explored the extent to

which eQTLs may have been missed because of limited statistical power. We compared *cis*-

heritability of colocalized caQTLs and eQTLs stratified by distance between the accessible region

and the transcription start site of the associated gene. We also investigated whether meta-analysis

of published LCL eQTL summary statistics, in order to effectively increase the sample size, can

uncover previously missed eQTLs. Second, we surveyed whether cell-type specificity of

regulatory variant effect may account for the missing regulation. We surveyed various immune

cell eQTL data to identify loci with which they colocalize even if LCL eQTLs did not colocalize

with those loci.

Through this study, we present how regulatory QTLs beyond eQTLs, such as caQTLs, can

be effective in detecting the potential molecular consequences of disease-associated variants.

Moreover, results from inspecting disease-associated loci where genetic effects are detected on

chromatin accessibility but not on expression suggest reasons why the effects on gene expression

may have been missed. These results provide insights on which strategies may be effective in

uncovering more genes that underlie diseases.

Results

Accessible chromatin in LCLs explains a significant proportion of immune-mediated

disease heritability

We aimed to evaluate whether variants that alter chromatin accessibility in LCLs may explain

genetic associations to IMD. First, we verified whether accessible regions in LCLs are enriched

for IMD heritability. We reanalyzed 100 LCL ATAC-seq samples[30] to define accessible regions

in this cell type. With stratified LD score regression (S-LDSC)[38], we estimated their heritability

enrichment across 13 IMDs, including 11 autoimmune diseases — autoimmune thyroid disease

(ATD)[39], celiac disease (CEL)[40], Crohn's disease (CD)[41], inflammatory bowel disease (IBD)[41],

juvenile idiopathic arthritis (JIA)[42], multiple sclerosis (MS)[43], primary biliary cholangitis (PBC)[39],

rheumatoid arthritis (RA)[44], systemic lupus erythematosus (SLE)[45], ulcerative colitis (UC)[41], and

vitiligo (VIT)[46] — and 2 allergic diseases — allergy (ALL)[47] and asthma (AST)[47]. We also

analyzed GWAS data for 3 non-immune diseases — type 2 diabetes (T2D)[48], coronary artery

disease (CAD)[49] and schizophrenia (SCZ)[50] — for comparison. Single nucleotide polymorphisms

(SNPs) in accessible regions in LCLs were significantly enriched for IMD heritability ($p <$

0.003125 (Bonferroni-corrected), S-LDSC; **Figure 1A**) and there was no significant enrichment

for non-immune diseases ($p > 0.05$, S-LDSC). These results indicate that accessible regions in

LCLs harbor many variants specifically associated with IMDs, and therefore that LCLs share

IMD-associated accessible regions with those of the causal cell type(s).

Figure 1. Immune disease heritability mediated by chromatin accessibility in LCLs. (A-B) Heritability enrichment in accessible regions in LCLs, based on (**A**) stratified LD score regression (S-LDSC) and (**B**) the proportion of heritability mediated by caQTLs in LCLs based on mediated expression score regression (MESC). For both, error bars represent jackknife standard errors of the mean. The color of the bars indicates disease type. *: $p < 0.003125$ (Bonferroni-corrected). (**C**) Mediated heritability enrichment of accessible regions by peak strength quintile. The strongest peaks are in the 1st quintile, and the weakest peaks are in the 5th quintile. Only enrichment values with FDR < 5% (based on q-value) are shown. (**D**) Mediated heritability enrichment of accessible regions by histone mark annotation. The percentages in parentheses represent the proportion of accessible regions with the indicated histone mark. Only enrichment values with FDR < 5% (based on q-value) are shown. Color and size of the points are on the same scale as in (**C**). PBC, primary biliary cholangitis; MS, multiple sclerosis; CEL, celiac disease; RA, rheumatoid arthritis; JIA, juvenile idiopathic arthritis; SLE, systemic lupus erythematosus; CD, Crohn's disease; UC, ulcerative colitis; IBD, inflammatory bowel disease; ATD, autoimmune thyroid disease; VIT, vitiligo; AST, asthma; ALL, allergies; SCZ, schizophrenia; T2D, type 2 diabetes; CAD, coronary artery disease.

Next, we applied mediated expression score regression (MESC)[16] to investigate the causal

relationship between caQTLs in LCLs and IMD associations. Compared to S-LDSC analysis that

tests for heritability enrichment of SNPs with some functional annotation (*e.g.*, accessible regions),

MESC analysis specifically estimates the heritability that is mediated (*i.e.*, h^2_{med}) by the SNPs' *cis*-

effects on a molecular phenotype (*e.g.*, caQTLs). We estimated that caQTLs in LCLs mediate

16.3–42.7 % of autoimmune disease heritability and 8.5–9.4 % of allergic disease heritability

(**Figure 1B**). For non-immune diseases, the estimates were lower and not significant ($p > 0.003125$

(Bonferroni-corrected), MESC). Interestingly, schizophrenia showed a nominally significant

proportion of caQTL-mediated heritability in LCLs ($p < 0.05$, MESC), consistent with the

hypothesis that B cells may play some role in schizophrenia pathogenesis[50,51]. Our results indicate

that LCLs are a valid cell type in which to search for caQTLs that mediate genetic risk for 7 IMDs

— CD, IBD, MS, PBC, RA, SLE, and UC — but not for allergic diseases. In subsequent analyses,

we focused on 7 IMDs – CD, IBD, MS, PBC, RA, SLE, and UC – that showed significant caQTL-

mediated heritability ($p < 0.003125$ (Bonferroni-corrected), MESC).

**Regions with higher levels of accessibility and active histone marks explain most of caQTL-
mediated heritability**

To understand which features characterize accessible regions that mediate IMD heritability, we

estimated h^2_{med} enrichment (proportion of h^2_{med} / proportion of peaks)[16] in specific sets of

accessible regions. We found that peaks with a larger number of nonredundant sequencing reads

(*i.e.*, "stronger" peaks) in LCLs showed stronger h^2_{med} enrichment (**Figure 1C**) and thus likely

affect IMD-relevant gene expression more than "weaker" peaks do. This observation is consistent

with the "Activity-by-Contact" model[9], in which peaks with greater chromatin accessibility and

H3K27ac ChIP-seq signal are predicted to have proportional effects on target gene expression.

Next, we considered peaks with the active histone marks H3K27ac, H3K4me1, or

H3K4me3[28,52]. Consistent with prior observations that putative cell-type-specific regulatory

elements marked with H3K27ac and H3K4me1 are enriched for relevant disease associations[34,35],

we found that caQTLs with H3K27ac and H3K4me1 marks in LCLs were significantly enriched

for mediated IMD heritability (q-value < 0.05, MESC; **Figure 1D**). Strikingly, both peak sets

explained almost all of caQTL-mediated IMD heritability (**Supplementary Figure 1**). Peaks with

H3K4me3 marks, representative of promoters[53], also showed significant h^2_{med} enrichment for most

IMDs (q-value < 0.05, MESC). Peaks with promoter-like signatures (*i.e.*, H3K27ac and

H3K4me3)[54] and those with enhancer-like signatures (*i.e.*, H3K27ac, but no H3K4me3)[54] were

also enriched for all IMD heritability (q-value < 0.05, MESC). Conversely, peaks without any of

the three active histone marks were completely depleted of caQTL-mediated IMD heritability

(**Supplementary Figure 1**). These 'ATAC only' peaks were shorter, weaker, and further away

from the TSS compared to peaks with active histone marks (**Supplementary Figure 2**). Altogether,

these results indicate that peaks characterized as putatively active regulatory elements explain

nearly all of caQTL-mediated IMD heritability.

caQTLs share IMD heritability with eQTLs and explain more of IMD heritability than do

eQTLs

The model that gene regulatory activity explains a significant fraction of noncoding genetic

associations to IMDs is supported by our findings that caQTLs mediate a significant proportion of

IMD heritability and that those with active histone marks show strong h^2_{med} enrichment. This is in

contrast with relatively low average h^2_{med} / h^2_{SNP} estimates (~11%) previously having been

observed for eQTLs across 48 human tissues in GTEx and various human traits[55]. To directly

compare the proportion of IMD heritability mediated by caQTLs and eQTLs in the same cell type

(*i.e.*, LCLs), we additionally applied MESC to gene expression data from LCLs (*i.e.*, Geuvadis

data)[56].

Figure 2. IMD heritability mediated by caQTLs and eQTLs. (**A**) h^2_{med} / h^2_{SNP} estimates of various IMDs for caQTLs, eQTLs, and their union. AID: autoimmune disease. Disease abbreviations along the *x*-axis are as in **Figure 1**. (**B**) Schema of the potential causal relationships between genetic variants, caQTLs, eQTLs, and IMD risk. The two diagrams depict possible h^2_{med} / h^2_{SNP} trends depending on the causal relationship between caQTLs and eQTLs. (**C**) Number of IMD-associated loci colocalized with caQTLs or eQTLs. The proportion is out of the total number of IMD-associated ($p < 10^{-6}$) loci. (**D**) *ELMO1* locus plot showing association to RA, PBC, MS, chromatin accessibility, and *ELMO1* expression in LCLs. Purple shading in the gene plot at the bottom indicates the caQTL peak, and the purple diamond is the lead variant (rs60600003) that is within that peak. The other variants are colored by the degree of linkage disequilibrium (LD) with the annotated variant. (**E**) Enrichment of the Biological Process Gene Ontology (GO) terms of genes in proximity to IMD-colocalized caQTLs without eQTL colocalization.

Across the 7 autoimmune diseases, the estimated proportion of heritability mediated by

eQTLs ($h^2_{med; eQTL} / h^2_{SNP}$) ranged from 9% to 22% (**Figure 2A**). For all 7 diseases, we estimated

that eQTLs mediated less heritability than did caQTLs, even though the caQTLs' smaller sample

size would potentially bias the estimates toward zero[16]. A possible explanation is that some IMD-
associated regulatory variants may show detectable effects on chromatin accessibility, but not on
gene expression, in LCLs at the current sample size ($n = 373$); such loci may account for the
missing regulation[15].

We anticipated that IMD-associated variants that affect gene expression in *cis* do so by
modulating regulatory element's activity. Therefore, we investigated whether eQTL-mediated
IMD heritability is shared by caQTL-mediated signals (**Figure 2B**). We performed MESC on both
caQTLs and eQTLs together to estimate the amount of IMD heritability mediated by both
collectively (<u>Methods</u>). For the 7 IMDs, the combined $h^2_{med; \text{ caQTL} \cup \text{eQTL}} / h^2_{SNP}$ was only slightly
higher (2.2–9.0 %) than the estimates for just caQTLs ($h^2_{med; \text{ just caQTL}} / h^2_{SNP}$; **Figure 2A**),
suggesting that approximately 56–82% of eQTL-mediated heritability is shared with caQTL-
mediated heritability (*i.e.*, $h^2_{med; \text{ caQTL} \cap \text{eQTL}} / h^2_{med; \text{ eQTL}}$; **Supplementary Figure 3**). These
estimates are consistent with substantial sharing of caQTL- and eQTL-mediated IMD heritability.
Nevertheless, 9–27% of IMD heritability is explained just by caQTLs, while only 2–9% of IMD
heritability is explained just by eQTLs.

Many IMD-associated loci show colocalization with caQTLs but not with eQTLs

We applied colocalization analysis[37] to identify IMD-associated loci that share genetic signals with
caQTLs or eQTLs in LCLs. We selected candidate loci of 200-kb windows for each IMD with the
following conditions: (1) lead IMD association at $p < 10^{-6}$, (2) lead caQTL or eQTL association at
$p < 10^{-4}$, and (3) at least one variant simultaneously showed caQTL or eQTL χ^2 statistics greater
than $0.8 \times$ lead χ^2 statistics for the caQTL or eQTL, respectively, and IMD association χ^2 statistics
greater than $0.8 \times \chi^2$ statistics for the IMD lead variant in the locus. We applied gwas-pw[37] and
considered loci with posterior probability of colocalization (PPA3) > 0.98 to be colocalized[57].
Some loci colocalized with only either a caQTL or an eQTL, while others colocalized with both
(**Figure 2C** and **Supplementary Tables 1-2**).

To confirm that the colocalized genes are relevant to IMD, we tested for their enrichment
of Gene Ontology (GO) annotation terms for specific biological processes[58]. Considering all genes
within 500 kb of the IMD GWAS lead variants at colocalized loci as background, the genes that
showed eQTL colocalization for any IMD were enriched for various immune responses and
signaling processes, such as "positive regulation of immune system process" and "regulation of

lymphocyte activation" (**Supplementary Figure 4**), indicating that the colocalized genes in LCLs
are involved in immune function. For example, *IL6R* and *IL12A* encode direct or indirect targets
of approved drugs – Tocilizumab and Ustekinumab – for autoimmune diseases like RA[59] and CD[60].
These two genes showed colocalization with both caQTLs and eQTLs in CD and PBC GWAS,
respectively (**Supplementary Figure 5A-B**). Increased *IL6R* expression was associated with
higher risk for CD, and increased *IL12A* expression was associated with lower risk for PBC and
SLE. The former observation is in line with Tocilizumab, a monoclonal antibody to IL-6 receptor,
showing efficacy in CD patients[59], although it is not pursued for approval because of potential side
effects[61]. Interestingly, *IL6R* and *IL12A* eQTLs did not colocalize with the association signals of
RA and CD, respectively, which are the diseases for which these drugs are approved
(**Supplementary Figure 5C-D**). Moreover, *ELMO1*, which previously had not been associated
with autoimmune diseases, showed eQTL colocalization with RA, PBC, and MS association
signals (**Figure 2D**). In all three, decreased *ELMO1* expression was associated with increased
disease risk. In mice, *Elmo1* was required for polarization and migration of B and T lymphocytes[62].
Across the IMDs, there were many loci that colocalized with a caQTL but not with an
eQTL (**Figure 2C**). These "caQTL only" loci showed enrichment for immune response genes in
cis compared to all accessible regions in LCLs[63,64], even though the colocalized eQTLs were
enriched for immune response genes as well (**Figure 2E** and **Supplementary Figure 4**), indicating
that IMD-relevant genes without eQTL colocalization in Geuvadis LCL data[56] are likely found in
these loci.

Distance to transcription start sites affect eQTLs but not caQTLs

Why might there be loci with caQTL colocalization only, despite the caQTL data having fewer
samples than the eQTL data (100 vs. 373)? Limited statistical power can prevent some eQTLs
from being detected and showing significant colocalization[19]. As hypothesized by Mostafavi and
colleagues, disease-relevant eQTLs may be weaker and more distal[18]. To understand the extent to
which this effect may result in many loci showing colocalization only with caQTLs, we compared
the *cis*-heritability (h^2_{cis}) of caQTLs and eQTLs depending on the distance from the ATAC peak
to the transcription start site (TSS) of the gene (*i.e.*, peak-to-TSS distance). We considered all
caQTLs and eQTLs regardless of disease association.

We identified pairs of caQTLs and eQTLs that colocalized with each other[37], which implies

that the regulatory variant modulating chromatin accessibility also affects gene expression. Then,

the distance between the ATAC peak and the TSS of the eQTL gene (*i.e.*, eGene) is the distance

between a regulatory element and its target gene's TSS. We stratified the pairs into peak-to-TSS

distance quintiles and compared the eQTL h^2_{cis} distribution of the first quintile (*i.e.*, closest pairs)

with that of the later quintiles. We observed that eQTL h^2_{cis} distribution decreased with increasing

distance of the paired ATAC peaks from the TSS ($p = 1.0\times10^{-4}$, 2.1×10^{-10}, 1.6×10^{-15}, and 1.7×10^{-20},

respectively, one-sided Wilcoxon rank-sum test; **Figure 3A**), consistent with the negative

relationship between promoter-enhancer genomic distance and impact on gene expression[65,66].

This result also explains why discovered eQTLs are concentrated near the promoter, where the

variants are more likely to show stronger effects[67]. In contrast, caQTL h^2_{cis} distribution was similar

across peak-to-TSS distances ($p > 0.05$, one-sided Wilcoxon rank-sum test; **Figure 3A**). For all

distance quintiles, caQTL h^2_{cis} was significantly higher than that of the paired eQTLs ($p = 4.0\times10^{-23}$,

1.3×10^{-24}, 1.5×10^{-19}, 1.1×10^{-28}, and 2.4×10^{-49}, respectively, one-sided paired Wilcoxon rank-

sum test; **Figure 3A**), and the contrast between them was greater at more distant quintiles,

suggesting that the statistical power to detect and colocalize eQTLs is increasingly lower than that

for caQTLs for regulatory effects far from the TSS.

Overall, caQTL h^2_{cis} had a significant positive effect ($p = 6.6\times10^{-11}$, linear regression) and

peak-to-TSS distance had a negative effect ($p = 2.4\times10^{-23}$, linear regression) on eQTL h^2_{cis} (**Figure

3B**). Thus, for regulatory variants that showed both caQTL and eQTL signals, those with larger

effects on chromatin accessibility tended to exhibit larger effects on gene expression, but their

eQTL effects diminished with increasing distance from TSSs.

Next, we investigated how caQTL-mediated IMD heritability is distributed with respect to

TSS. If caQTLs beyond the typical *cis*-eQTL window of 1 megabase (Mb) around the genes' TSS

explain some proportion of IMD heritability, then *cis*-eQTL analyses might require a wider

window to detect disease-associated eQTLs. Across the 7 diseases, caQTLs within 500 kb of the

TSS of expressed genes explained almost all of the caQTL-mediated IMD heritability (92–100%;

Figure 3C), indicating that regulatory variants are most likely within 500 kb of the target gene's

TSS and supporting the use of a 1 Mb window for *cis*-eQTL analyses. Depending on the disease,

41–66% of the caQTL-mediated IMD heritability was detected in distal peaks further than 10 kb

from the TSS of expressed genes, further supporting the analysis of regulatory variants beyond

promoter regions.

261   In sum, the power to detect eQTLs diminishes with increasing distance of the variant from

262 the TSS, but the power to detect caQTLs is largely invariant regardless of peak-to-TSS distance

263 (**Figure 3D**). Since $h^2_{med;\ caQTL}$ are distributed mostly within 500 kb of genes' TSS, the IMD loci

264 colocalizing only with caQTLs could still be weak, undetected eQTLs. Under this model, we

265 predicted that increasing the power to detect eQTLs in LCLs, such as increasing sample size, may

266 lead to further eQTL colocalizations in loci in which we observed only caQTL colocalization.

Figure 3. Effect of peak-to-TSS distance on *cis*-heritability of caQTLs and eQTLs and IMD heritability. (**A**) Distribution of *cis*-heritability (h^2_{cis}) of caQTLs and eQTLs by peak-to-TSS distance quintiles. The ranges of peak-to-TSS distance are shown in parentheses. The comparisons shown on the top (in respective colors) are between the nearest and each subsequent quintile of the respective QTL h^2_{cis} distribution (*i.e.*, one-sided Wilcoxon rank-sum test). The comparisons shown on the bottom (in black) are between caQTL and eQTL h^2_{cis} distribution (one-sided paired Wilcoxon rank-sum test). *: $p < 10^{-4}$, **: $p < 10^{-10}$, ***: $p < 10^{-20}$, and ns: $p > 0.05$. (**B**) Regression estimates and their standard errors of the linear regression model testing the effects of caQTL h^2_{cis} and peak-to-TSS distance on eQTL h^2_{cis}. Peak-to-TSS distance was expressed in units of 100 kb to neatly visualize the effect size estimates. SE: standard error. (**C**) Proportion of caQTL-mediated IMD heritability explained by ATAC peaks within various TSS windows. Percentage for each TSS window denotes the proportion of ATAC peaks in that window. Disease abbreviations along the *x*-axis are as in **Figure 1**. (**D**) A model of the relationship between peak-to-TSS distance and power to detect a corresponding caQTL or eQTL. The thickness of the arrows indicates the variant effect size on chromatin accessibility (yellow) or gene expression (blue). TSS, transcription start site; CRE, *cis*-regulatory element.

Increasing the sample size reveals some eQTL colocalization

For genetic association studies, increasing the sample size is a way to increase statistical power.

Therefore, we meta-analyzed four LCL eQTL summary statistics[55,56,68,69], leading to a total sample

size of 1,128 individuals. We performed colocalization analysis using the meta-analyzed summary

statistics to evaluate whether effectively increasing the sample sizes would uncover more disease-

associated eQTLs in LCLs, especially in loci where a caQTL already showed IMD colocalization.

Up to 6 additional loci showing eQTL colocalization were thus detected for each IMD (**Figure 4A**

and **Supplementary Table 3**). For example, *CIITA* is the class II major histocompatibility

complex transactivator, which causes severe immunodeficiency if dysfunctional[70]. The *CIITA*

locus is associated with IBD right below the genome-wide significance level (rs10445003, $p =$

7.5×10^{-8}), and it colocalized with a caQTL signal, but initially not with any eQTL (**Supplementary**

Figure 6A). However, the meta-analyzed statistics showed a stronger association to *CIITA*

expression ($p = 6.7 \times 10^{-8}$) than without meta-analysis ($p = 5.4 \times 10^{-4}$) and exhibited a significant

colocalization. Interestingly, two of the causal CD genes that previously lacked colocalized

eQTLs[15], *CARD9* and *ATG16L1*, showed significant colocalization in the meta-analyzed LCL

eQTL data (**Supplementary Figure 6B-C**).

Figure 4. Additional colocalization of caQTL-colocalized IMD loci with meta-analyzed LCL eQTL data. (**A**) Number of caQTL-colocalized IMD loci that showed eQTL colocalization in LCLs. Disease abbreviations along the *x*-axis are as in **Figure 1**. (**B**) Distribution of peak-to-TSS distance of all caQTL-eQTL pairs and of those colocalized with IMD association. The number of loci in each category is shown in parentheses. *: $p < 0.01$, ns: $p > 0.05$. (**C**) *POU3F1* eQTL that became significantly colocalized with RA association by meta-analyzing LCL eQTL data. Purple shading in the gene plot at the bottom indicates the caQTL peak, and the purple diamond is the lead variant (rs60600003) that is within that peak. The other variants are colored by the degree of linkage disequilibrium (LD) with the annotated variant.

283   We hypothesized that increased sample size would improve the power to detect weaker
284 and distal eQTL colocalization. Comparison of the accessibility peak's distance to the paired eQTL
285 gene's (eGene) TSS showed that the newly detected eQTLs tended to be more distal ($p = 0.06$,
286 one-sided Wilcoxon rank-sum test; **Figure 4B**). However, compared to the distribution of the
287 peak-to-TSS distance for all caQTL-eQTL pairs showing colocalization, IMD-associated loci that
288 showed caQTL and eQTL colocalization had greater peak-to-TSS distance on average ($p = 0.002$
289 for Geuvadis and $p = 5.9 \times 10^{-6}$ for the meta-analyzed data; **Figure 4B**). For example, a RA-
290 associated locus near *POU3F1*, a neuronal transcription factor that is also induced by interferon[71],
291 colocalized with a distal eQTL located about 126 kb upstream of its promoter, after meta-analysis
292 strengthened the eQTL association ($p < 10^{-10}$; **Figure 4C**). These results suggest that additional
293 IMD-associated loci with distal, weaker eQTLs in LCLs might be found if eQTL data were
294 generated for a larger number of individuals.

IMD loci that colocalized with caQTLs but not eQTLs showed lower levels of active histone marks

Despite uncovering more eQTL colocalizations through meta-analysis, more than 40% of the caQTL-colocalized loci nevertheless showed no colocalization with an eQTL in LCLs (**Figure 4B**). We investigated whether these "caQTL only" peaks might be inactive regulatory elements. We quantified the active histone mark levels for H3K27ac, H3K4me1, and H3K4me3 at colocalized caQTL peaks in LCLs and then compared their levels between the "caQTL and eQTL" and "caQTL only" loci. On average, H3K27ac marks were stronger at "caQTL and eQTL" peaks, supporting that the corresponding regulatory elements might be more active (**Supplementary Figure 7**). H3K4me3 marks were detected more often at "caQTL and eQTL" peaks, leading to stronger average signal. In contrast, H3K4me1 levels were highly similar between the two sets of peaks. Although "caQTL only" peaks generally showed lower levels of active histone marks, several individual "caQTL only" peaks showed comparable levels. These peaks could be inactive *cis*-regulatory elements in LCLs that affect gene expression in a different cellular context. Therefore, we next examined whether those caQTLs appear as eQTLs in other immune cell types.

Various immune cell types exhibit eQTL colocalization where LCLs did not

We downloaded eQTL summary statistics generated from 26 naïve and stimulated immune cell types[22,23,72–74] to search for eQTLs that may correspond to the remaining, IMD-colocalized caQTLs (**Supplementary Table 4**). The profiled cell types range from B cells and monocytes to subtypes of T cells, as well as stimulated T cells and macrophages. We tested for colocalization of these eQTLs to IMD associations.

　　　　25–42% of the caQTL-colocalized IMD loci that were missing eQTL colocalizations in LCLs showed eQTL colocalizations in at least one immune cell type (**Figure 5A** and **Supplementary Table 5**). The overlap of LCL caQTLs with non-LCL immune cell eQTLs was greater than expected by chance for 5 of the 7 IMDs ($p < 0.00714$ (Bonferroni-corrected), Fisher's exact test, for CD, IBD, PBC, RA, and UC; **Supplementary Table 6**), suggesting that the caQTLs found in LCLs may also show regulatory function in those immune cell types. Comparing across the datasets, we found that the number of loci with eQTL colocalization varied depending on the cell type, but that the effect of the sample size was more profound ($r^2 = 0.60$–0.79; **Figure 5B** and **Supplementary Table 7**). We meta-analyzed eQTL data of 3 immune cell types with multiple

sources – naïve CD4$^+$ T cell, monocyte, and memory regulatory T cell (Treg) – and this also

increased the number of loci with significant eQTL colocalization (**Supplementary Table 8**).

Altogether, these results suggest that although generating eQTL data in more cell types and cell

states uncovers context-specific eQTLs, increasing the sample size should also be a priority to

ensure sufficient statistical power.

We investigated the potential reasons why IMD loci that colocalized with caQTLs in LCLs

showed eQTLs not in LCLs but in other immune cells. First, LCL caQTLs may correspond to gene

regulatory elements that exert their effects on gene expression in a different cellular context. For

instance, monocyte H3K27ac levels in the "caQTL only" loci where monocyte eQTLs colocalized

were higher than those with no monocyte eQTLs (**Supplementary Figure 8**). Second, some

examples, such as for *TNFSF15*, were due to cell-type specific gene expression (**Figure 6**): despite

a significant colocalization of IBD with a caQTL in LCLs in the *TNFSF15* locus, disease-

associated eQTL signal was detected only in monocytes (**Figure 6A**). Tumor necrosis factor-like

cytokine 1A (TL1A), the protein encoded by *TNFSF15*, is secreted by monocytes and many other

cells to activate helper T cells, regulatory T cells, and B cells[75]. *TNFSF15* expression was low in

LCLs (mean transcript per million [TPM] = 0.30), but higher in monocytes (mean TPM = 2.23).

Of the genes that showed exclusively monocyte eQTL colocalization, those with low expression

(mean TPM < 1) in LCLs generally showed higher expression in monocytes (**Figure 6B-C**).

Figure 5. Added utility of various immune cell eQTL data. (**A**) Number of loci that additionally colocalized with eQTLs by LCL meta-analysis (orange) and immune cell data (cyan) compared to the original analysis with Geuvadis LCL eQTL data (purple). The height of the bar is the proportion of loci with eQTL colocalization out of the total IMD loci with caQTL colocalization in the earlier analysis. Disease abbreviations along the *x*-axis are as in **Figure 1**. (**B**) Relationship between the number of MS GWAS loci with eQTL colocalization and sample size for each eQTL dataset. Meta-analyzed eQTL data are labeled with their cell types. NK cell, natural killer cell; Treg, regulatory T cell.

Figure 6. IMD loci that colocalized with an eQTL in monocytes, but not in LCLs, even though they colocalized with caQTLs in LCLs. (**A**) *TNFSF15* locus plot showing genetic association to IBD, chromatin accessibility in LCLs, and *TNFSF15* expression in LCLs and monocytes. Purple shading in the gene plot at the bottom indicates the caQTL peak, and the purple diamond shows a strongly associated variant (rs7848647) that is within that peak. The other variants are colored by the degree of LD with the annotated variant. (**B**) Expression levels of genes in LCLs and monocytes colored according to their eQTL colocalization outcome. TPM: transcripts per million. (**C**) Number of genes with eQTL colocalization in monocytes, but not in LCLs, separated by the gene's expression level in LCLs (column) and whether it is lower or higher than that in monocytes (row).

However, low expression in LCLs was likely not the explanation for most cases of "monocyte only" eQTLs (blue points in **Figure 6B**) because most "monocyte only" eQTL genes were expressed at level higher than 1 TPM in LCLs.

Overall, expanding the eQTL search to various immune cell types increased the number of eQTL-colocalized loci among those that previously colocalized only with caQTLs (**Figure 5A**). On average, approximately 75% of the caQTL-colocalized loci ultimately showed eQTL colocalization. These results highlight the utility of eQTL data across a range of immune cell types for discovery of IMD-associated eQTLs.

Discussion

A lack of the link between many noncoding GWAS loci to the associated variants' gene regulatory effects have posed challenges in understanding their genetic mechanism. A better evaluation of

the potential reasons for the missing regulation will guide future data generation projects to

elucidate disease-associated loci. To determine why some loci might lack colocalized eQTLs, we

focused on chromatin accessibility, which is a molecular phenotype affected by regulatory variants

more directly than are eQTLs. We approached this question with mediated heritability analysis[16]

and colocalization analysis[36,37].

We found that caQTLs in LCLs mediate a significant proportion of heritability for many

autoimmune diseases. In contrast, LCLs did not appear to be an effective cell type to model gene

regulatory effects in allergic diseases. The h^2_{med} / h^2_{SNP} estimates for caQTLs were higher than

those of eQTLs in most autoimmune diseases, even though the smaller sample size of caQTL data

(*i.e.*, 100 vs. 373) could bias caQTLs' estimates toward zero[16]. We also showed that disease-

associated chromatin accessibility effects often share the genetic signal with gene expression

effects but that there are also many loci without an eQTL detected in LCLs.

By focusing on disease-associated caQTL that lacked significant eQTLs, we explored how

additional colocalized eQTL effects could be uncovered. First, increasing the sample sizes for the

eQTL statistics via meta-analysis demonstrated that more eQTL colocalizations can be detected

with increased statistical power and robustness[19]. These results are consistent with the hypothesis

that disease-associated eQTLs are typically weaker and distal due to negative selection against

large expression changes for causal genes[18]. Second, many caQTLs in LCLs without eQTLs in

LCLs showed eQTL colocalization in other immune cell types. Context specificity of eQTLs has

been widely considered to be the primary explanation for the difficulty of pinpointing disease-

associated eQTLs[17,22,23,73]. Our observation that many IMD-colocalized caQTLs in LCLs show

eQTLs in other immune cell types suggests that caQTL effects may be shared across cell types

whereas eQTL effects are more context-specific[23]. eQTL data for the specific cellular contexts

need to be generated in future studies to uncover more such examples. Our results also showed

that having a large sample size in eQTL data is likely a key factor for discoveries of more disease-

associated eQTLs.

Limitations of the study

The h^2_{med} / h^2_{SNP} estimates can be biased because of insufficient sample size of the QTL data. Thus,

the proportion of mediated IMD heritability could change based on the specific caQTL and eQTL

data. Colocalization analysis tests whether the QTL and GWAS data likely share genetic signal,

and such shared signal could arise from either causal mediation or pleiotropy. Therefore, further

experiments are needed to establish causality of colocalized eQTL genes. Lastly, we hypothesized

that caQTL effects are often shared across cell types, but we did not have access to relevant data

to test this hypothesis.

Methods

ATAC-seq data processing and peak calling

We downloaded LCL ATAC-seq data of British (GBR) samples (n = 100) from European

Nucleotide Archive (ENA) under accession ERP110508[30]. The available files were cram

alignment files mapped to the b37 reference genome, so we extracted unique read pairs using

SAMtools[76] and bamtofastq command from bedtools[77]. The reads were paired-end and each 75

base pairs (bp) long. The data contained reads with Nextera transposase adapters, so we removed

the adapter sequences and bases of poor quality at the 3' end using cutadapt. Trimmed reads with

both pairs shorter than 20 bp were discarded. The command was "cutadapt -a file:${forward} -A

file:${reverse} -e 0.25 -j 2 -q 15 --pair-filter=both -m 20". (${forward} and ${reverse} files

contain the forward and reverse Nextera transposase adapters). We mapped the reads to the

GRCh38 reference genome using Bowtie 2[78] with the "GRCh38_noalt_decoy_as" index provided

on the tool's website. The command was "bowtie2 --very-sensitive --no-mixed --no-discordant -I

20 -X 2000". We kept only read alignments with mapping quality greater than 1. We also removed

reads aligning to the mitochondrial genome, those overlapping ENCODE exclusion regions (file

ID: ENCFF356LFX)[79] and potential PCR duplicates using scripts from WASP[80].

To represent peaks across the samples, we subsampled 3 million read pairs from each and

pooled them. Then, we used MACS2[81] with the BAMPE option for peak calling. The command

was "macs2 callpeak -f BAMPE -g hs -q 0.05". We further used the 'bdgcmp' and 'bdgpeakcall'

subcommand to find peaks that are at least 100 bp long (-l) and merge those that are less than 100

bp apart (-g). We also merged peaks similarly derived from individual samples using the 'merge'

command in bedtools.

Furthermore, for the sake of comprehensiveness, we repeated these steps with LCL ATAC-

seq data of Yoruban (YRI) samples[82] and merged the peaks with the earlier peak set derived from

GBR samples. In total, there were 443,403 peaks genome-wide.

RNA expression data preparation

We downloaded RNA expression level data of the LCL samples[56] from the Expression Atlas[83].

The source data is available at European Nucleotide Archive under accession ERP001942. We

retained TPM values of protein-coding and long noncoding RNAs in European samples for

downstream analyses, including QTL analysis, mediated heritability analysis, and colocalization.

LCL samples' genotype preparation

To utilize the genotype calls of the highest quality, we downloaded the high-coverage 1000 Genomes (1kG) Project data[84]. There was a total of 493 samples that had either ATAC-seq or RNA-seq data in this study. Of those, 479 samples had genotypes derived from the high-coverage whole-genome sequencing data. 14 samples had genotypes derived from microarrays[85]. The samples that needed imputation are listed in **Supplementary Table 9**. We lifted over the microarray data based on the hg19 reference genome to GRCh38 and filtered for variants present in the high-coverage 1kG data. We first imputed microarray genotype data using the TOPMed imputation server[86] and extracted SNPs with imputation $R^2 \geq 0.5$ and imputed the rest of the variants, including short indels, using high-coverage data with Beagle5.2[87,88]. After keeping only variants with $DR2 \geq 0.7$, we merged the imputed genotypes with the high-coverage 1kG genotypes.

Curation of IMD GWAS data

We downloaded GWAS summary statistics for 13 IMDs, including 11 autoimmune diseases (autoimmune thyroid disease (ATD)[39], celiac disease (CEL)[40], Crohn's disease (CD)[41], inflammatory bowel disease (IBD)[41], juvenile idiopathic arthritis (JIA)[42], multiple sclerosis (MS)[43], primary biliary cholangitis (PBC)[39], rheumatoid arthritis (RA)[44], systemic lupus erythematosus (SLE)[45], ulcerative colitis (UC)[41], and vitiligo (VIT)[46]) and 2 allergic diseases (allergy (ALL)[47] and asthma (AST)[47]). For each disease, we searched for more recent studies with larger sample sizes and prioritized those with genome-wide statistics, rather than those with only Immunochip variants[89]. To compare with non-immune diseases, we also downloaded summary statistics for 3 non-immune diseases (type 2 diabetes (T2D)[48], coronary artery disease (CAD)[49] and schizophrenia (SCZ)[50]). In this study, we analyzed only those GWAS summary statistics derived from cohorts of individuals with European ancestries.

Since the LCL samples' genotypes are based on the GRCh38 reference genome, we lifted over any GWAS data based on b37 reference genome to the GRCh38 genomic coordinates. Briefly, we formatted the summary statistics as bed files and used the liftOver tool[90] to convert them to GRCh38 genomic coordinates. Then, to ensure that the reference alleles match the sequences of the GRCh38 reference genome, we used gwas2vcf tool[91].

Chromatin accessibility QTLs in lymphoblastoid cell lines

First, we quantified the chromatin accessibility levels at ATAC-seq peaks identified earlier. We counted the number of read fragments overlapping each peak using featureCounts[92]. For each sample, the read counts were normalized for library size using trimmed mean of M-values[93] so that the values are comparable across the samples. Then, the phenotype values were further normalized to follow a standard normal distribution across the samples, using quantile normalization. Peaks with counts per million (CPM) < 0.8 or counts < 10 for more than 20% of the samples were discarded.

Next, we performed a principal component analysis (PCA) on the phenotype matrix to derive potential latent covariates. We selected the number of principal components (PC) to incorporate in the regression model based on the Buja and Eyuboglu algorithm[94] that is implemented in PCAForQTL[95]. Ultimately, we accounted for sex, library size, 3 genotype PCs and 13 phenotype PCs in the QTL analysis. We performed genetic association tests on variants within 200 kilobases (kb) of the peak using tensorQTL[96]. We discarded variants with minor allele frequency less than 5%.

IMD heritability enrichment in accessible regions of LCLs

We evaluated the relevance of accessible regions in LCLs to IMD heritability using stratified LD score regression (S-LDSC)[38]. We used the baselineLD v2.2 annotation in hg38 and the European LD reference from the 1000 Genomes Project (downloaded from the S-LDSC website, https://alkesgroup.broadinstitute.org/LDSCORE/GRCh38/). We used the set of filtered ATAC-seq peaks that we tested for QTL associations. We accounted for 16 diseases (13 IMDs and 3 non-IMDs) for the Bonferroni-corrected *p*-value threshold.

Mediated heritability estimation of QTLs

We estimated the heritability mediated by QTLs (h^2_{med}) using mediated expression score regression (MESC)[16]. We denote heritability of tested SNPs as h^2_{SNP}.

caQTL-mediated heritability. We estimated the 'expression scores' for chromatin accessibility in LCLs using individual genotypes and phenotypes. We analyzed the same set of peaks and accounted for the same covariates as we did for the QTL analysis above. For mediated heritability

estimation, we accounted for baselineLD v2.2 annotation in hg38 without the QTL annotations, as

they could be redundant. The estimand of interest is the proportion of heritability mediated by

caQTLs (h^2_{med} / h^2_{SNP}).

To evaluate whether certain peak sets are enriched for mediated heritability, we utilized

the gene set analysis functionality. For peak strength, we considered the 95% percentile CPM value

of each peak and stratified the peaks into quintiles. For histone marks, we first generated histone

ChIP-seq peaks using data from Delaneau et $al.$[52]. Similar to calling ATAC-seq peaks, we sampled

3 million reads per sample and merged them before applying MACS2[81]. We downloaded 3 control

ChIP-seq data from ENCODE to use as input (File IDs: ENCFF066RCS, ENCFF159XTB, and

ENCFF850RIE)[79]. Then, we curated sets of ATAC-seq peaks that overlapped H3K27ac,

H3K4me1, and H3K4me3 ChIP-seq peaks by at least 1 bp. ATAC-seq peaks that overlapped

H3K27ac but not H3K4me3 regions were labeled as 'enhancer-like signature', while those that

overlapped H3K27ac and H3K4me3 regions were labeled as 'promoter-like signature'. The

estimand of interest is the proportion of mediated heritability explained by the peak set (peak set

$h^2_{med} /$ total h^2_{med}).

Comparison with eQTL-mediated heritability. First, we estimated h^2_{med} / h^2_{SNP} of eQTLs (*i.e.*,

$h^2_{med; eQTL} / h^2_{SNP}$) the same way as we did for that of caQTLs. Then, we also estimated h^2_{med} / h^2_{SNP}

of caQTLs and eQTLs together (*i.e.*, $h^2_{med; caQTL \cup eQTL}$) with MESC meta-analysis[16]. caQTLs and

eQTLs were also stratified as separate sets to account for potential differences in the relationship

of QTL *cis*-heritability and h^2_{med}. This meta-analyzed estimate is effectively the amount of

heritability mediated by either caQTLs or eQTLs in LCLs. This estimate reveals the overall

relationship between heritability mediated by caQTLs and eQTLs. For instance, the estimate of

the heritability mediated exclusively by caQTLs would be $h^2_{med; just caQTL} = h^2_{med; caQTL \cup eQTL} - h^2_{med;}$

$_{caQTL}$. The estimate of mediated heritability shared by caQTLs and eQTLs is $h^2_{med; caQTL \cap eQTL} =$

$(h^2_{med; caQTL} + h^2_{med; eQTL}) - h^2_{med; caQTL \cup eQTL}$.

Colocalization analyses

caQTL and eQTL colocalization with IMD GWAS. First, we selected candidate colocalization

loci by filtering for overlapping 'significant' variants. The candidate loci met the following

conditions: (1) lead IMD association at $p < 10^{-6}$, (2) lead caQTL or eQTL association at $p < 10^{-4}$,

and (3) at least one variant simultaneously showed caQTL or eQTL χ^2 statistics greater than 0.8 ×

lead χ^2 statistics for the caQTL or eQTL and IMD association χ^2 statistics greater than $0.8 \times \chi^2$

statistics for the IMD lead variant in the locus. Then, we applied gwas-pw[37] on the variants within

100 kb of the lead variant. We considered loci with posterior probability of colocalization (PP3) >

0.98 to be colocalized[57].

Colocalization of caQTL with eQTL. We performed a colocalization analysis of caQTLs and

eQTLs to curate a set of loci where the same genetic signal likely explains both associations. The

pairs of caQTLs and eQTLs reveal the distance between the regulatory element and the target

gene's TSS. We selected candidate colocalization loci with: (1) IMD association at $p < 10^{-6}$, (2)

QTL association at $p < 10^{-4}$, and (3) the two lead variants showed LD $r^2 > 0.8$. We applied gwas-

pw[37] on the variants within 200 kb of the tested caQTL peak. We considered loci with posterior

probability of colocalization (PP3) > 0.98 to be colocalized[57].

Enrichment of biological processes

To test whether colocalized genes are likely relevant to autoimmune diseases, we surveyed which

Biological Process Gene Ontology (GO) terms were overrepresented compared to all the genes within

500 kb of each IMD association signal. Enrichment of biological processes was evaluated using

Protein Analysis Through Evolutionary Relationships (PANTHER)[97]. The foreground list

comprised all of the genes whose eQTL signal colocalized with one of the 7 autoimmune diseases

(CD, IBD, MS, PBC, RA, SLE, and UC). The background list of genes was all of the genes within

500 kb of each IMD lead variant for which we observed colocalization. Moreover, we tested

whether colocalized caQTLs without eQTLs are closer to genes related to immune processes than

expected by chance. For this, we used Genomic Regions Enrichment of Annotations Tool

(GREAT)[63]. The foreground list comprised ATAC-seq peaks at IMD loci showing colocalization

only with caQTLs and not with eQTLs. The background list is all ATAC-seq peaks identified in

LCLs that we tested for caQTL association.

Relationship between peak-to-TSS distance and *cis*-heritability of caQTL and eQTL

The distance between the colocalized caQTL peak and the eGene's TSS (*i.e.*, peak-to-TSS distance)

was determined to be the shortest distance from one end of the caQTL peak to the TSS. The pairs

of caQTLs and eQTLs were split into quintiles based on their peak-to-TSS distance from closest

to farthest.

MESC analysis uses REML implemented in GCTA[98] to estimate QTL *cis*-heritability. We
referred to its output and compared the *cis*-heritability of caQTLs and eQTLs based on peak-to-
TSS distance. To visualize the distribution of *cis*-heritability estimates, we grouped pairs of
colocalized caQTLs and eQTLs based on peak-to-TSS distance quintiles.

LCL eQTL meta-analysis

We downloaded LCL eQTL data from 3 studies through eQTL Catalogue release 6[99]. The sample
sizes were 190, 147, and 418 for GENCORD[68], GTEx[55], and TwinsUK[69], respectively. We meta-
analyzed the summary statistics using the inverse variance weighted fixed effects model. If a
variant was missing in a subset of the studies, then only the available statistics were meta-analyzed.
We used these meta-analyzed statistics to perform colocalization the same way as earlier
colocalization analyses.

Colocalization analysis with immune cell eQTL data

We downloaded eQTL data for various immune cell types from the eQTL Catalogue release 6[99].
The source studies from the eQTL Catalogue include BLUEPRINT[72], DICE[22], Alasoo *et al.*[23] and
Bossini-Castillo *et al.*[74]. The represented immune cell types include T cell subtypes, B cells,
neutrophils, and monocytes. We also separately downloaded data for CD4+ T cells with and
without stimulation[73]. The selection of candidate loci and colocalization analysis on them followed
the same procedure as that for other QTLs.

We also evaluated whether meta-analysis of eQTL data can increase the number of loci
with eQTL colocalization. We meta-analyzed eQTL data for 3 immune cell types with multiple
sources – naïve CD4+ T cell, monocyte, and memory regulatory T cell (Treg) – using the inverse
variance weighted fixed effects model. Specifically, we meta-analyzed naïve CD4+ T cell eQTL
summary statistics from Soskic *et al.*[73], BLUEPRINT[72], and DICE[22]. We meta-analyzed monocyte
eQTL summary statistics from BLUEPRINT[72] and DICE[22]. We meta-analyzed memory Treg
eQTL summary statistics from Bossini-Castillo *et al.*[74] and DICE[22]. We used these meta-analyzed
statistics to perform colocalization the same way as earlier colocalization analyses.

Data availability

Processed data and code for generating the figures presented in the manuscript are available at https://github.com/BulykLab/IMD-colocalization-manuscript-figures. Genotype data are downloaded from the 1000 Genomes Project (https://www.internationalgenome.org/data-portal/data-collection/30x-grch38). LCL ATAC-seq data are downloaded from ENA (https://www.ebi.ac.uk/ena/browser/view/PRJEB28318). LCL RNA-seq count matrix is downloaded from Expression Atlas[83] (https://www.ebi.ac.uk/gxa/experiments/E-GEUV-1/Results). GWAS data for the studies listed in the Methods section are downloaded from the GWAS catalog[100] (https://www.ebi.ac.uk/gwas/). eQTL summary statistics are downloaded either from the eQTL Catalogue[99] (https://www.ebi.ac.uk/eqtl/) or the source publication.

Acknowledgments

We thank Vijay Sankaran, Shamil Sunyaev, Alexander Gusev, and members of the Raychaudhuri lab for helpful discussion. This work was funded by NIH grant R01 HG010501.

Author contributions

R.J. and M.L.B. conceived and designed the research project. R.J. performed all analyses and prepared the figures. M.L.B. supervised the research. R.J. and M.L.B. wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

References

1. Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *Am. J.*
Hum. Genet. **101**, 5–22 (2017).

2. Claussnitzer, M. *et al.* A brief history of human disease genetics. *Nature* **577**, 179–189 (2020).

3. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in
regulatory DNA. *Science* **337**, 1190–1195 (2012).

4. Gusev, A. *et al.* Partitioning heritability of regulatory and cell-type-specific variants across 11
common diseases. *Am. J. Hum. Genet.* **95**, 535–552 (2014).

5. Nicolae, D. L. *et al.* Trait-associated SNPs are more likely to be eQTLs: Annotation to enhance
discovery from GWAS. *PLoS Genet.* **6**, (2010).

6. Hormozdiari, F. *et al.* Leveraging molecular quantitative trait loci to understand the genetic
architecture of diseases and complex traits. *Nat. Genet.* **50**, 1041–1047 (2018).

7. O'Connor, L. J. *et al.* Extreme Polygenicity of Complex Traits Is Explained by Negative
Selection. *Am. J. Hum. Genet.* **105**, 456–476 (2019).

8. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From Polygenic
to Omnigenic. *Cell* **169**, 1177–1186 (2017).

9. Nasser, J. *et al.* Genome-wide enhancer maps link risk variants to disease genes. *Nature* **593**, 238–
243 (2021).

10. Weeks, E. M. *et al.* Leveraging polygenic enrichments of gene features to predict genes
underlying complex traits and diseases. *Nat. Genet.* **55**, 1267–1276 (2023).

11. Morris, J. A. *et al.* Discovery of target genes and pathways at GWAS loci by pooled single-cell
CRISPR screens. *Science* **380**, eadh7699 (2023).

12. Plenge, R. M., Scolnick, E. M. & Altshuler, D. Validating therapeutic targets through human
genetics. *Nat. Rev. Drug Discov.* **12**, 581–594 (2013).

13. Chun, S. *et al.* Limited statistical evidence for shared genetic effects of eQTLs and autoimmune-
disease-associated loci in three major immune-cell types. *Nat. Genet.* **49**, 600–605 (2017).

14. Barbeira, A. N. *et al.* Exploiting the GTEx resources to decipher the mechanisms at GWAS loci.
Genome Biol. **22**, 49 (2021).

15. Connally, N. J. *et al.* The missing link between genetic association and regulatory function. *Elife*
11, (2022).

16. Yao, D. W., O'Connor, L. J., Price, A. L. & Gusev, A. Quantifying genetic effects on disease
mediated by assayed gene expression levels. *Nat. Genet.* 1–8 (2020) doi:10.1038/s41588-020-
0625-2.

17.   Umans, B. D., Battle, A. & Gilad, Y. Where Are the Disease-Associated eQTLs? *Trends Genet.*
612   **37**, 109–124 (2021).

613   18.   Mostafavi, H., Spence, J. P., Naqvi, S. & Pritchard, J. K. Systematic differences in discovery of
614   genetic effects on gene expression and complex traits. *Nat. Genet.* **55**, 1866–1875 (2023).

615   19.   Hukku, A. *et al.* Probabilistic Colocalization of Genetic Variants from Complex and Molecular
616   Traits: Promise and Limitations. *Am. J. Hum. Genet.* **108**, 25–35 (2021).

617   20.   Glassberg, E. C., Gao, Z., Harpak, A., Lan, X. & Pritchard, J. K. Evidence for weak selective
618   constraint on human gene expression. *Genetics* **211**, 757–772 (2019).

619   21.   Dimas, A. S. *et al.* Common regulatory variation impacts gene expression in a cell type-dependent
620   manner. *Science* **325**, 1246–1250 (2009).

621   22.   Schmiedel, B. J. *et al.* Impact of Genetic Polymorphisms on Human Immune Cell Gene
622   Expression. *Cell* **175**, 1701–1715 (2018).

623   23.   Alasoo, K. *et al.* Shared genetic effects on chromatin and gene expression indicate a role for
624   enhancer priming in immune response. *Nat. Genet.* **50**, 424–431 (2018).

625   24.   Nathan, A. *et al.* Single-cell eQTL models reveal dynamic T cell state dependence of disease loci.
626   *Nature* **606**, 120–128 (2022).

627   25.   Field, A. & Adelman, K. Evaluating Enhancer Function and Transcription. *Annu. Rev. Biochem.*
628   **89**, 213–234 (2020).

629   26.   Kasowski, M. *et al.* Variation in transcription factor binding among humans. *Science* **328**, 232–
630   235 (2010).

631   27.   Kilpinen, H. *et al.* Coordinated effects of sequence variation on DNA binding, chromatin
632   structure, and transcription. *Science* **342**, 744–747 (2013).

633   28.   Waszak, S. M. *et al.* Population Variation and Genetic Control of Modular Chromatin Architecture
634   in Humans. *Cell* **162**, 1039–1050 (2015).

635   29.   Degner, J. F. *et al.* DNase-I sensitivity QTLs are a major determinant of human expression
636   variation. *Nature* **482**, 390–394 (2012).

637   30.   Kumasaka, N., Knights, A. J. & Gaffney, D. J. High-resolution genetic mapping of putative causal
638   interactions between regions of open chromatin. *Nat. Genet.* **51**, 128–137 (2019).

639   31.   Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nat.*
640   *Rev. Genet.* **16**, 197–212 (2015).

641   32.   Wu, Y. *et al.* Joint analysis of GWAS and multi-omics QTL summary statistics reveals a large
642   fraction of GWAS signals shared with molecular phenotypes. *Cell Genomics* 100344 (2023)
643   doi:10.1016/j.xgen.2023.100344.

644   33.   Jeong, R. & Bulyk, M. L. Blood cell traits' GWAS loci colocalization with variation in PU.1

645 genomic occupancy prioritizes causal noncoding regulatory variants. *Cell genomics* **3**, 100327
646 (2023).

34. Roadmap Epigenomics Consortium *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–329 (2015).

35. Farh, K. K. H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* **518**, 337–343 (2015).

36. Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).

37. Pickrell, J. K. *et al.* Detection and interpretation of shared genetic influences on 42 human traits. *Nat. Genet.* **48**, 709–717 (2016).

38. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).

39. Cordell, H. J. *et al.* An international genome-wide meta-analysis of primary biliary cholangitis: Novel risk loci and candidate drugs. *J. Hepatol.* **75**, 572–581 (2021).

40. Dubois, P. C. A. *et al.* Multiple common variants for celiac disease influencing immune gene expression. *Nat. Genet.* **42**, 295–302 (2010).

41. de Lange, K. M. *et al.* Genome-wide association study implicates immune activation of multiple integrin genes in inflammatory bowel disease. *Nat. Genet.* **49**, 256–261 (2017).

42. López-Isac, E. *et al.* Combined genetic analysis of juvenile idiopathic arthritis clinical subtypes identifies novel risk loci, target genes and key regulatory mechanisms. *Ann. Rheum. Dis.* **80**, 321–328 (2021).

43. Patsopoulos, N. A. *et al.* Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science* **365**, (2019).

44. Ishigaki, K. *et al.* Multi-ancestry genome-wide association analyses identify novel genetic mechanisms in rheumatoid arthritis. *Nat. Genet.* **54**, 1640–1651 (2022).

45. Bentham, J. *et al.* Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nat. Genet.* **47**, 1457–1464 (2015).

46. Jin, Y. *et al.* Genome-wide association studies of autoimmune vitiligo identify 23 new risk loci and highlight key pathways and regulatory variants. *Nat. Genet.* **48**, 1418–1424 (2016).

47. Loh, P. R., Kichaev, G., Gazal, S., Schoech, A. P. & Price, A. L. Mixed-model association for biobank-scale datasets. *Nat. Genet.* **50**, 906–908 (2018).

48. Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat. Genet.* **44**, 981–990 (2012).

49.   Schunkert, H. *et al.* Large-scale association analysis identifies 13 new susceptibility loci for

coronary artery disease. *Nat. Genet.* **43**, 333–8 (2011).

50.   Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological insights from

108 schizophrenia-associated genetic loci. *Nature* **511**, 421–7 (2014).

51.   van Mierlo, H. C., Broen, J. C. A., Kahn, R. S. & de Witte, L. D. B-cells and schizophrenia: A

promising link or a finding lost in translation? *Brain. Behav. Immun.* **81**, 52–62 (2019).

52.   Delaneau, O. *et al.* Chromatin three-dimensional interactions mediate genetic effects on gene

expression. *Science* **364**, (2019).

53.   Heintzman, N. D. *et al.* Distinct and predictive chromatin signatures of transcriptional promoters

and enhancers in the human genome. *Nat. Genet.* **39**, 311–318 (2007).

54.   ENCODE Project Consortium *et al.* Expanded encyclopaedias of DNA elements in the human and

mouse genomes. *Nature* **583**, 699–710 (2020).

55.   GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues.

Science **369**, 1318–1330 (2020).

56.   Lappalainen, T. *et al.* Transcriptome and genome sequencing uncovers functional variation in

humans. *Nature* **501**, 506–511 (2013).

57.   Kundu, K. *et al.* Genetic associations at regulatory phenotypes improve fine-mapping of causal

variants for twelve immune-mediated diseases. *Nat. Genet.* **44**, 0–23 (2022).

58.   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all. *Protein*

Sci. **31**, 8–22 (2022).

59.   Sanmartí, R., Ruiz-Esquide, V., Bastida, C. & Soy, D. Tocilizumab in the treatment of adult

rheumatoid arthritis. *Immunotherapy* **10**, 447–464 (2018).

60.   Khanna, R. & Feagan, B. G. Ustekinumab for the treatment of Crohn's disease. *Immunotherapy* **5**,

803–15 (2013).

61.   Monemi, S. *et al.* Incidence of Gastrointestinal Perforations in Patients with Rheumatoid Arthritis

Treated with Tocilizumab from Clinical Trial, Postmarketing, and Real-World Data Sources.

Rheumatol. Ther. **3**, 337–352 (2016).

62.   Stevenson, C. *et al.* Essential role of Elmo1 in Dock2-dependent lymphocyte migration. *J.*

Immunol. **192**, 6062–70 (2014).

63.   McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions. *Nat.*

Biotechnol. **28**, 495–501 (2010).

64.   Tanigawa, Y., Dyer, E. S. & Bejerano, G. WhichTF is functionally important in your open

chromatin data? *PLoS Comput. Biol.* **18**, e1010378 (2022).

65.   Fulco, C. P. *et al.* Activity-by-contact model of enhancer–promoter regulation from thousands of

713   CRISPR perturbations. *Nat. Genet.* **51**, (2019).

714   66.   Zuin, J. *et al.* Nonlinear control of transcription through enhancer-promoter interactions. *Nature*
715   **604**, 571–577 (2022).

716   67.   Võsa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and
717   polygenic scores that regulate blood gene expression. *Nat. Genet.* **53**, (2021).

718   68.   Gutierrez-Arcelus, M. *et al.* Passive and active DNA methylation and the interplay with genetic
719   variation in gene regulation. *Elife* **2013**, 1–18 (2013).

720   69.   Buil, A. *et al.* Gene-gene and gene-environment interactions detected by transcriptome sequence
721   analysis in twins. *Nat. Genet.* **47**, 88–91 (2015).

722   70.   Dziembowska, M. *et al.* Three novel mutations of the CIITA gene in MHC class II-deficient
723   patients with a severe immunodeficiency. *Immunogenetics* **53**, 821–9 (2002).

724   71.   Hofmann, E. *et al.* Octamer-binding factor 6 (Oct-6/Pou3f1) is induced by interferon and
725   contributes to dsRNA-mediated transcriptional responses. *BMC Cell Biol.* **11**, 61 (2010).

726   72.   Chen, L. *et al.* Genetic Drivers of Epigenetic and Transcriptional Variation in Human Immune
727   Cells. *Cell* **167**, 1398-1414.e24 (2016).

728   73.   Soskic, B. *et al.* Immune disease risk variants regulate gene expression dynamics during CD4+ T
729   cell activation. *Nat. Genet.* **54**, 817–826 (2022).

730   74.   Bossini-Castillo, L. *et al.* Immune disease variants modulate gene expression in regulatory CD4+
731   T cells. *Cell Genomics* **2**, 100117 (2022).

732   75.   Xu, W.-D., Li, R. & Huang, A.-F. Role of TL1A in Inflammatory Autoimmune Diseases: A
733   Comprehensive Review. *Front. Immunol.* **13**, 891328 (2022).

734   76.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

735   77.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
736   features. *Bioinformatics* **26**, 841–2 (2010).

737   78.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–
738   359 (2012).

739   79.   Dunham, I. *et al.* An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**,
740   57–74 (2012).

741   80.   Van De Geijn, B., Mcvicker, G., Gilad, Y. & Pritchard, J. K. WASP: Allele-specific software for
742   robust molecular quantitative trait locus discovery. *Nat. Methods* **12**, 1061–1063 (2015).

743   81.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

744   82.   Banovich, N. E. *et al.* Impact of regulatory variation across human iPSCs and differentiated cells.
745   *Genome Res.* **28**, 122–131 (2018).

746   83.   Papatheodorou, I. *et al.* Expression Atlas update: from tissues to single cells. *Nucleic Acids Res.*

747 **48**, D77–D83 (2020).

748 84. Byrska-Bishop, M. *et al.* High-coverage whole-genome sequencing of the expanded 1000
749  Genomes Project cohort including 602 trios. *Cell* **185**, 3426-3440.e19 (2022).

750 85. Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).

751 86. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.* **48**, 1284–
752  1287 (2016).

753 87. Browning, B. L., Zhou, Y. & Browning, S. R. A One-Penny Imputed Genome from Next-
754  Generation Reference Panels. *Am. J. Hum. Genet.* **103**, 338–348 (2018).

755 88. Browning, B. L., Tian, X., Zhou, Y. & Browning, S. R. Fast two-stage phasing of large-scale
756  sequence data. *Am. J. Hum. Genet.* **108**, 1880–1890 (2021).

757 89. Cortes, A. & Brown, M. A. Promise and pitfalls of the Immunochip. *Arthritis Res. Ther.* **13**, 101
758  (2011).

759 90. Kent, W. J. *et al.* The human genome browser at UCSC. *Genome Res.* **12**, 996–1006 (2002).

760 91. Lyon, M. S. *et al.* The variant call format provides efficient and robust storage of GWAS summary
761  statistics. *Genome Biol.* **22**, 32 (2021).

762 92. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
763  assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–30 (2014).

764 93. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression
765  analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

766 94. Buja, A. & Eyuboglu, N. Remarks on Parallel Analysis. *Multivariate Behav. Res.* **27**, 509–40
767  (1992).

768 95. Zhou, H. J., Li, L., Li, Y., Li, W. & Li, J. J. PCA outperforms popular hidden variable inference
769  methods for molecular QTL mapping. *Genome Biol.* **23**, 210 (2022).

770 96. Taylor-Weiner, A. *et al.* Scaling computational genomics to millions of individuals with GPUs.
771  *Genome Biol.* **20**, 228 (2019).

772 97. Mi, H. *et al.* Protocol Update for large-scale genome and gene function analysis with the
773  PANTHER classification system (v.14.0). *Nat. Protoc.* **14**, 703–721 (2019).

774 98. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex
775  trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).

776 99. Kerimov, N. *et al.* A compendium of uniformly processed human gene expression and splicing
777  quantitative trait loci. *Nat. Genet.* **53**, 1290–1299 (2021).

778 100. Sollis, E. *et al.* The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. *Nucleic*
779  *Acids Res.* **51**, D977–D985 (2023).