Complex bacterial diversity of Guaymas Basin hydrothermal sediments revealed by synthetic long-read sequencing (LoopSeq)

John E. Hinkle[1], Jeffrey P. Chanton[2], Molly A. Moynihan[3,4], S. Emil Ruff[3,4], Andreas Teske[1*]

[1]Department of Earth, Marine and Environmental Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
[2]Department of Earth, Ocean and Atmospheric Science, Florida State University, Tallahassee, FL
[3]The Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA
[4]The Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA

* Correspondence:
Andreas Teske
teske@email.unc.edu

Keywords: LoopSeq, 16S rRNA, sulfur-cycling bacteria, sulfur cycle, Guaymas Basin, hydrothermal sediments

Abstract. Hydrothermal sediments host phylogenetically diverse and physiologically complex

microbial communities. Previous studies of microbial community structure in hydrothermal

sediments have typically used short-read sequencing approaches. To improve on these

approaches, we use LoopSeq, a high-throughput synthetic long-read sequencing method that has

yielded promising results in analyses of microbial ecosystems, such as the human gut

microbiome. In this study, LoopSeq is used to obtain near-full length (approximately 1400 -

1500 nucleotides) bacterial 16S rRNA gene sequences from hydrothermal sediments in Guaymas

Basin. Based on these sequences, high-quality alignments and phylogenetic analyses provided

new insights into previously unrecognized taxonomic diversity of sulfur-cycling microorganisms

and their distribution along a lateral hydrothermal gradient. Detailed phylogenies for free-living

and syntrophic sulfur-cycling bacterial lineages identified well-supported monophyletic clusters

that have implications for the taxonomic classification of these groups. Particularly, we identify

clusters within *Candidatus* Desulfofervidus that represent unexplored physiological and genomic

diversity. In general, LoopSeq-derived 16S rRNA gene sequences aligned consistently with

reference sequences in GenBank; however, chimeras were prevalent in sequences as affiliated

with the thermophilic *Candidatus* Desulfofervidus and *Thermodesulfobacterium*, and in smaller

numbers within the sulfur-oxidizing family *Beggiatoaceae*. Our analysis of sediments along a

well-documented thermal and geochemical gradient show how sulfur-cycling bacteria of

different taxonomic groups persist as active catalysts of the sulfur cycle throughout surficial

hydrothermal sediments in the Guaymas Basin.

INTRODUCTION

The Guaymas Basin is a marginal rift basin in the Gulf of California that is characterized

by the coexistence of hydrothermal vents and cold seeps. Unlike many other regions with

hydrothermal vents, Guaymas Basin is subject to relatively high sediment accumulation rates

(Calvert, 1966). Basaltic sills, freshly emplaced into organic-rich sediments, drive the

hydrothermal circulation of hydrocarbons, sulfide, and nutrient-rich fluids (Einsele et al. 1980,

Lizarralde et al., 2011). Abundant energy and carbon sources sustain highly active and very

diverse microbial communities at the sediment surface (Teske et al., 2021). Consequently,

Guaymas Basin has been the subject of many sequencing studies throughout the past two

decades, starting with the use of near-full-length (~1500 nucleotides) cloned PCR products

(Teske et al., 2002), then progressing to high throughput pyrosequencing of very short (~80

nucleotides) 16S rRNA gene fragments (Ruff et al., 2015). More recent studies utilize high

throughput sequencing methods that produce longer (~400-500 nucleotides) 16S rRNA gene

fragments (Teske et al., 2019, Su et al., 2023). The abundance of sequence data obtained via

different methods makes the Guaymas Basin an ideal model system to assess the performance of

new next-generation sequencing technologies, such as LoopSeq.

LoopSeq is a high-throughput synthetic long-read sequencing method, which can

generate near-full length gene sequences (Callahan et al., 2021). Studies using clone library

construction and sequencing methods yield near full-length and high-quality sequences but are

greatly limited by throughput and sequencing depth. Clone library-based methods are time

consuming and only the most abundant microorganisms are usually detected. In turn, high

throughput sequencing methods can yield tens of thousands of sequence reads, allowing to detect

extremely rare organisms (Sogin et al., 2006). Yet, these reads are too short for high-resolution

phylogenetic analyses and can only be used for relatively basic taxonomic identification. Short

reads of 400-500 nucleotides can be used to construct phylogenies, but placement of unknown or

poorly resolved lineages remains problematic, due to the limited information provided by short

sequences and limited bootstrap support for phylogenetic branching patterns (Teske et al., 2019,

Ramírez et al., 2021).

LoopSeq bridges this gap and allows investigators to obtain nearly full-length 16S rRNA

gene sequences in high throughput, for costs comparable to short read amplicons. The

combination of high throughput and nearly full-length sequences allows for thorough surveys of

the diversity of known microbial lineages, as well as high-resolution phylogenetic placement of

rare or poorly resolved microbial lineages even for highly diverse ecosystems, such deep-sea

hydrothermal sediments. To test the capabilities and reliability of LoopSeq for environmental

samples, Guaymas Basin is ideal because i) the ecosystem is generally well understood, ii) the

hydrothermal sediments are very diverse, iii) the sediments are rich in organics that can be

problematic for sequencing, e.g. hydrocarbons, and iv) extensive 16S rRNA gene sequence

databases are available to verify LoopSeq sequences against publicly available short read and

full-length sequences.

METHODS

Sampling. Guaymas Basin sites were visited and sampled with R/V *Atlantis*, HOV *Alvin*,

and AUV *Sentry* during cruise AT37-06 (December 6-29, 2016). *Alvin* dives targeted previously

explored sampling areas in the southern axial valley of Guaymas Basin (Teske et al., 2016), or

newly identified sites found by AUV *Sentry*. Hydrothermal sediment samples (*Alvin* push cores)

used in this study were obtained during *Alvin* Dive 4872 (Dec. 24, 2016) in the Cathedral Hill

area (27.00.684 N/ 111.24.266 W) at 2014 m depth. The sample set covers a hydrothermal

gradient from hot hydrothermally charged sediments covered with orange and white

Beggiatoaceae mats to temperate bare sediment, within ~1 m distance (Fig. 1). Core 4872-01

represents bare sediment that was not covered by a *Beggiatoaceae* mat, whereas Core 4872-06

was covered by a white *Beggiatoaceae* mat, and Core 4872-14 was covered by an orange

Beggiatoaceae mat. Once the sediment cores were returned to the surface, porewater profiles

were collected using rhizons (Rhizosphere Research Products, Wageningen, The Netherlands).

Then, these cores were divided into 3 cm sections and methane samples were collected by

injecting 2 mL sediment plugs into 30 mL serum vials containing 2 ml of 1 M sodium hydroxide

solution each. The serum vials were immediately sealed with thick rubber stoppers, crimped with

aluminum seals, and stored at 4 ºC. The remaining sediment samples were frozen at -80ºC by the

shipboard science crew, and later used for DNA extraction shoreside.

Thermal profiles. Thermal profiles were measured in surficial sediments using *Alvin's*

50 cm heat flow probe (https://ndsf.whoi.edu/alvin/using-alvin/sampling-equipment/). The 50 cm

probe contains thermal sensors every 10 cm, starting 5 cm below the attached plastic disc (the

"puck") that limits probe penetration and rests on the seafloor once the probe was inserted. After

5 to 10 minutes of temperature reading stabilization, temperature readings were recorded.

101   **Geochemical analyses.** For sulfide analysis, 1 mL porewater subsamples were fixed with

0.1 mL of 0.1 M zinc acetate solution to preserve sulfide as zinc sulfide until analysis using the

methylene blue method (Cline, 1969). For sulfate measurements, porewater samples were diluted

in water (1:50) and analyzed using ion chromatography (Metrohm 930 Compact IC flex with

Metrosep A PCC HC/4.0 preconcentration column, and Metrosep A Supp 5 Guard/4.0

chromatography column). Sulfate for the white and orange cores was measured by Wetland

Biogeochemistry Analytical Services (WBAS) using ion chromatography on a Dionex ICS-1000

using a RFIC IonPac AS22 4×250 mm column and a RFIC IonPac AG22 Guard column 4×50

mm. The IC was calibrated using a 5-point standard curve and a minimum R^2 value of 0.999.

The detection limit measured at 0.10 ppm sulfate. A check standard was made from a Dionex 7

anion standard (1000 ppm) to a concentration of 15 ppm. An external Standard from HACH (100

ppm sulfate) was used to make a 20 ppm quality check standard. In general, quality checks were

analyzed for 10 % of the samples.

114   Porewaters from the Guaymas Basin were analyzed at WBAS for nitrate/nitrite (NO_x)

and ammonium (NH_4^+) concentrations colorimetrically, using a Flow Solutions IV segmented

flow Auto Analyzer from O.I Analytical, College Station, TX. The HCl-acidified samples were

neutralized with sodium hydroxide (NaOH) before analysis. NO_x was determined using the

cadmium reduction method and NH_4^+ was determined using the phenate method. Both nutrients

were diluted to get their concentrations within the linear range of the auto analyzer. Quality

control standards from certified stock standards (Environmental Research Associates), were

analyzed every 15-20 samples.

122   For combined concentration and $\delta^{13}C$ analysis of methane, 2 mL sediment subsamples

were added to 30 mL serum vials containing 2 mL of 1 M NaOH solution, sealed with thick

butyl rubber stoppers, crimped with aluminum seals, and stored at 4°C. Since cores were

retrieved unpressurized, outgassing may have particularly impacted the measurements of

methane concentrations near and above saturation. After the cruise, the methane samples were

analyzed by headspace gas chromatography-flame ionization detection (GC-FID) at Florida State

University (Magen et al., 2014). Additionally, the gas samples were analyzed for $\delta^{13}CH_4$ by

injecting 0.1 to 0.5 mL of sample into a gas chromatograph interfaced to a Finnigan MAT Delta

S Isotope Ratio Mass Spectrometer inlet system as previously described (Chanton and Liptay

2000). Values are reported in the per mil (‰) notation relative to Vienna Pee Dee Belemnite

(VPDB).

DNA extraction. DNA was extracted from Guaymas Basin sediment samples using a

modified version of the protocol developed by Zhou et al., 1996 (see supplementary material).

The extraction method was optimized for deep-sea sediments. The detailed protocol can be found

in the supplementary materials.

LoopSeq sequencing. Sequencing libraries were prepared from extracted genomic DNA

with the commercially available LoopSeq kits from Loop Genomics

(https://go.elementbiosciences.com/16s-loopseq-user-guide, also available in the supplementary

material). Synthetic long reads (SLRs) were constructed from the short-read sequencing reads

using the standard Loop Genomics informatics pipeline and processed as previously described

(Callahan et al., 2021). The method involves attaching two DNA tags: one Unique Molecular

Identifier (UMI) to each unique "parent" DNA molecule and one sample-specific tag (i.e., a

sample index) equally to all molecules in the same sample. Barcoded molecules are amplified,

multiplexed, and each UMI is distributed intramolecularly to a random position within each

parent molecule. Molecules are then fragmented into smaller units at the position of each UMI,

creating a library of UMI-tagged fragments with an average length of 400 bp compatible with an

Illumina sequencing platform run in PE150 mode (Callahan et al., 2021).

This protocol allows for the assembly of continuous 16S rRNA gene long reads from

individual DNA molecules, providing sequencing data like classical long read sequencing

methods but with much higher throughput. The terminal primers determine the extent of the 16S

rRNA gene sequence yielded by this approach (27F, 5'-AGAGTTTGATCMTGGCTCAG-3';

1492R, 5'-TACCTTGTTACGACTT-3'). LoopSeq can be performed on existing DNA

sequencing instruments, and due to the chemistry and the tagging of individual molecules the

method is semi-quantitative providing a major advantage towards other amplification-based

methods.

We note that LoopSeq reconstructs a single long read from several short reads. Therefore,

we define the resulting amplicon sequence variants (ASVs) as LoopSeq variants (L-ASVs) to

differentiate them from "traditional" ASVs that are essentially PCR amplicons.

DADA2 analysis. The statistical programming language R 4.2.1 (R Core Team, 2022)

and the DADA2 (v. 1.20.0) bioinformatics package (Callahan et al., 2016), along with a

LoopSeq suitable DAD2 workflow (Callahan et al., 2021) were used to analyze 16S rRNA gene

sequences obtained by LoopSeq sequencing. DADA2 creates ASVs (or LoopSeq variants (L-

ASVs) in our case) by grouping together amplicon reads of the same sequence.

165   The function "removePrimers" was used to select for 16S rRNA amplicon sequences

with both the forward and reverse primers and allowing a maximum mismatch of two base pairs,

as reads lacking either primer do not represent the complete gene. The following parameters

were used to filter the remaining sequences: a minimum length of 1000 bp (minLen=1000), a

maximum length of 2000 bp (maxLen=2000), a minimum quality score of 3 (minQ=3), a

maximum expected error rate of 1 (maxEE=1), and a maximum Ns of 0 (maxN = 0). These

parameters may be used to filter reads without selecting for reads with a forward and reverse

primer to capture long, but not complete genes. We found that 20.4% of reads that fulfilled the

previously stated parameters lacked the forward/reverse primer. Only full-length gene sequences

were kept in our analysis. The function "dereqFastq" was used to dereplicate reads. The function

"learnErrors" was used with a band size of 32 (BAND_SIZE=32) to learn error rates. The error

estimate function was set to the PacBio Error Function. DADA2 was ran with a p-value cutoff of

1^{-10} (OMEGA_A=1e-10), single detection (DETECT_SINGLETONS=TRUE), and a band size

of 32 (BAND_SIZE=32). The function "removeBimeraDenovo" (method = consensus) was used

to remove chimeras. An overview of raw read processing is depicted in Supplementary Table 2.

180   The function "assignTaxonomy" was used to assign taxonomy using the Silva reference

181   database (v138) (Quast et al., 2013). A total of 12,494 L-ASVs were identified. Figures depicting

182   taxonomic distribution, alpha diversity, and beta diversity (Fig. 3, Suppl. Figs 2 & 3) were

183   created using the R Phyloseq package (McMurdie & Holmes, 2013). All sequence reads were

184   submitted to the NCBI Sequence Read Archive (BioProject: PRJNA1105367).

185   **16S rRNA gene sequence phylogeny.** We compared our L-ASV sequences against

reference sequences obtained from the NCBI GenBank database. All sequences were aligned

using Multiple Sequence Comparison by Log-Expectation (MUSCLE) in MEGA11 (Stecher et

al., 2020). From these alignments, MEGA11 was used to construct minimum-evolution

phylogenetic trees checked by the aggregate of 1000 bootstrap replicates (Rzhetsky & Nei, 1992,

Felsenstein, 1985). Bootstrap values greater than 70% are shown and NCBI GenBank accession

numbers are provided on tip labels.

192   **16S rRNA gene sequence distance matrices.** Distance matrices corresponding to each

phylogenetic tree are shown in the Supplementary Data File. The number of base substitutions

between sequences are shown as a percentage (i.e., per 100 sites). Analyses were conducted

using the Maximum Composite Likelihood model (Tamura et al., 2004). Each analysis involved

all 16S rRNA gene sequences displayed in the corresponding phylogenetic tree. All ambiguous

positions were removed for each sequence pair (pairwise deletion option). Evolutionary analyses

were conducted in MEGA11 (Tamura et al., 2021, Stecher et al., 2020).

RESULTS AND DISCUSSION

Sediment characteristics. The color of Guaymas Basin *Beggiatoaceae* mats indicated

the relative temperature of the underlying sediment as measured using the *Alvin* heat flow probe;

orange mats covered hotter sediments and white mats covered comparatively cooler sediments

(McKay et al., 2012). Temperature varied greatly with increasing sediment depth at the three

sample sites (Figs. 1 & 2a). Temperatures at 40 centimeters below the seafloor (cmbsf) exceeded

100 ℃ below the orange mat, reached 75 ℃ below the white mat, and remained near 20 ℃ in

sediments without a *Beggiatoaceae* mat (Fig. 2a).

Temperature considerably impacts microbial population density, and therefore the

amount of DNA available for extraction. Measurable concentrations of DNA persisted the

deepest (40 cmbsf) in Core 1, from bare sediment neighboring the warm mats. In contrast, DNA

recovery is limited to surficial sediments (~ 20 cmbsf) in Core 14 from the hot center of the

orange mat (Fig. 2b.). Interestingly, the highest DNA yields are found within the upper ~6 cm of

sediment in all three cores, indicating that the impact of hydrothermal activity extends beyond

the *Beggiatoaceae* mats to the bare sediments of Core 1 (Fig. 2b). Previous research in Guaymas

Basin consistently reveals a very rapid decline in cell counts and 16S rRNA gene abundances

with increasing depth and temperature in hot hydrothermal sediments (Meyer et al., 2013, Møller

et al., 2018, Lagostina et al., 2021).

Micromolar concentrations of sulfide are observed throughout the bare sediment core,

whereas the white and orange mat cores reach concentrations of sulfide up to 2 millimolar (Fig.

2c). At the surface, sulfide concentrations are near zero for the bare sediment and white mat

cores. In contrast, the orange mat core has elevated (~1 millimolar) sulfide levels at the surface,

consistent with high hydrothermal flow in the mat center. Sulfate is abundant in all three cores,

with depletion only occurring at ~32 cm and below in the orange mat core, indicating that

hydrothermal circulation contributes to the replenishment of sulfate in surficial sediments (Fig.

2d).

Ammonium is observed throughout our white and orange mat sediment cores (Fig. 2e)

Consistent with previous research, ammonium concentrations increase as temperature increases

downcore (Ramírez et al., 2021). Ammonium in Guaymas Basin hydrothermal sediments is

primarily derived from the thermal degradation of organic matter (Von Damm et al., 1985). Deep

in the orange mat core, ammonium concentrations reach ~8 mM (Fig. 2e), close to the ~10-15

mM concentrations observed in Guaymas Basin hydrothermal fluids (Von Damm et al., 1985).

Elevated nitrate/nitrite concentrations are observed in surficial mat covered sediments

(Fig. 2f). Bottom water concentrations of nitrate in Guaymas Basin are up to 50 μM (Winkel et

al., 2014). Elevated nitrate levels (up to nearly 0.25 mM; Fig. 2f) in mat-covered surficial

sediments are at least in part due to damaged *Beggiatoaceae* mats leaking nitrate stored in their

vacuoles into surficial sediments (Schutte et al., 2018).

High concentrations of methane are observed in all three cores (Fig. 2g). A gradient of

peak methane concentrations (~1 mM in the bare sediment core, ~2 mM in the white mat core,

and ~8 mM in the orange mat core) illustrates the impact of hydrothermal fluids that are

supersaturated in methane. High methane concentrations (multiple millimolar) are a

characteristic feature of Guaymas Basin hydrothermal sediments, similar to those found at cold

seep sites (Welhan, 1988, McKay et al., 2016, Song et al., 2021).

Hydrothermal methane in Guaymas Basin unmodified by microbial activity has a $\delta^{13}CH_4$

value of approximately -42 ‰ (McKay et al. 2016, Song et al. 2021). The preferential oxidation

of $^{12}CH_4$ by methane-oxidizing archaea leads to the orange and white mat cores containing

methane with a heavier carbon isotopic signature that is less depleted in carbon-13, with values

ranging from -30 to -20 ‰ (Fig. 2h). The lighter carbon isotopic signature of methane

throughout the bare sediment core and near the surface of the orange and white mat cores,

indicates some contribution due to microbial methanogenesis (Fig. 2h). The methanogens in

these sediments are most likely methylotrophs, which unlike other types of methanogens are

capable of successfully performing methanogenesis in sulfate-replete sediments (Zhuang et al.,

2018).

Inferred Carbon Sources. Guaymas Basin sediments are rich in organic carbon (De la

Lanza-Espino & Soto, 1999) and harbor numerous heterotrophic microbial lineages (Pérez

Castro et al., 2021). Carbon sources are likely to shift throughout hydrothermal gradients in the

sediments. Aromatic hydrocarbons serve as widely available carbon sources for aromatic-

degrading bacteria throughout surficial Guaymas Basin sediments (Mara et al., 2022, Gutierrez

et al., 2015, Goetz & Jannasch, 1993). In contrast, alkanes and methane occur are more abundant

near hydrothermally active "hot spots" (Song et al., 2021, McKay et al., 2016). The abundance

of alkanes and methane in hot sediments contributes to the evolution of thermotolerance and

thermophily in several groups of alkane and methane degrading bacteria and archaea (Zehnle et

al., 2023, Benito Merino et al., 2022). The upper limit of hydrocarbon degradation near

approximately 80 ℃ is supported by the increased abundance of ^{13}C-depleted substrates, which

reflects the absence of microbial degradation (Song et al., 2021).

Sequencing Profile. LoopSeq recovered nearly full-length sequences (Suppl. Fig 1) that

affiliated with several phyla of heterotrophic bacteria, such as most Proteobacteria

Desulfobacterota, Bacteroidota, and Chloroflexi (Fig. 3). These phyla have been detected by

previous Guaymas Basin sequencing surveys using clone libraries of near full-length 16S rRNA

genes (Teske et al., 2002, Dowell et al., 2016, McKay et al., 2016). We analyzed the alpha

diversity of each mat site using five measures (Observed, Chao1, Shannon, Simpson, and Inverse

Simpson). Although we expected hydrothermal stress to influence alpha diversity, we found no

significant difference in alpha diversity across these three cores (Suppl. Fig. 2). To explore

potential differences in beta diversity, we created a non-metric multi-dimensional scaling

(NMDS) plot using the Bray-Curtis dissimilarity method. The bare sediment samples largely

cluster together, particularly the deeper samples collected below 21 cmbsf (Suppl. Fig. 3. We

also observe a cluster of surficial sediment samples (the top 6 cmbsf) from all three coring sites

(Suppl. Fig. 3).

Sulfur-cycling lineages that frequently interface with hydrocarbon metabolism are

particularly well represented in the sequencing profile and present throughout surficial

hydrothermal sediments in the Guaymas Basin (Fig. 3). Sulfur-oxidizing *Beggiatoaceae*

assimilate methane-derived carbon (McKay et al., 2012). Numerous sulfate-reducing lineages

exist in syntrophic assemblages with methane and alkane oxidizing archaea (Knittel et al., 2003),

while other sulfate-reducers perform terminal oxidation of hydrocarbons as free-living organisms

(Teske et al., 2019). We performed phylogenetic analyses on these sulfur-cycling lineages to

determine their intragroup diversity.

Beggiatoaceae. The detection of *Beggiatoaceae* is an interesting test of LoopSeq

capabilities. Previous efforts to generate full-length *Beggiatoaceae* sequences required collecting

large amounts of individually prepared and cleaned filaments (McKay et al., 2012). Due to the

size of individual *Beggiatoaceae* cells, they are outnumbered by other much smaller marine

bacteria, complicating sequencing efforts and requiring high-throughput methods to overcome

the ubiquitous presence of other marine bacterial groups. In our survey, all LoopSeq variants (L-

ASV) identified as *Beggiatoaceae* matched reference sequences of the white or orange

phylotype, with no additional phylotypes identified (Fig. 4, Suppl. Fig. 4). Light microscopy

reveals that past and present sequencing efforts do not capture a complete picture of

Beggiatoaceae diversity in Guaymas Basin (Salman et al., 2013). There is no evidence for

introns within the 16S rRNA genes of Guaymas Basin *Beggiatoaceae,* although they have been

observed in non-Guaymas Basin *Beggiatoaceae* (Salman et al., 2011). Recovered L-ASVs were

generally found to be associated with surficial sediments underlying white and orange

Beggiatoaceae mats. Some L-ASVs were found deeper below the surface, which can be

attributed to the "Smear effect", when the coring device pulls some of the long filaments down

into the sediment column.

***Candidatus* Desulfofervidus.** In contrast to *Beggiatoaceae*, *Candidatus* Desulfofervidus

sequences are often recovered in Guaymas Basin sequencing surveys (McKay et al., 2016,

Dowell et al., 2016). *Ca.* Desulfofervidus is a sulfate-reducing syntrophic partner of methane-

and alkane-oxidizing archaea (Laso-Pérez et al., 2016) with a thermal optimum of 50ºC (Holler

et al., 2011) to 60ºC (Kniemeyer et al., 2007, Wegener et al., 2015). Although often observed as

a syntroph, *Ca.* Desulfofervidus can also survive as a free-living hydrogenotroph without

methane- or alkane-oxidizing partners (Krukenberg et al., 2016). *Ca.* Desulfofervidus is a

phylogenetically narrow clade, as it is classified as its own family, order, and class-level lineage

within the Phylum *Desulfobacterota* (Waite et al., 2020). However, previous research has not

examined the phylogenetic diversity within *Ca.* Desulfofervidus.

Our phylogenies identified several monophyletic well-supported clusters, each with

nearly identical 16S rRNA gene sequences. Each cluster contained reference sequences from

50ºC (Holler et al., 2011) or 60ºC (Kniemeyer et al. 2007, Wegener et al., 2015) enrichments. All

reference sequences were from Guaymas Basin, with no close matches from other sites. The

close alignment of our sequences with established Guaymas Basin reference sequences attests to

their quality. Our phylogeny supports the previous assertion of Krukenberg et al. (2016) that all

Ca. Desulfofervidus lineages are closely related, since they form a 16S rRNA gene sequence

cluster of approximately 98.5 % sequence identity (Suppl. Data File 1, Sheet 2). The five

clusters, supported with bootstrap values ≥ 81 % (Fig. 5), imply the presence of unrecognized

species diversity within *Ca.* Desulfofervidus.

We found sequences in GenBank that formed two deeply branching clusters with strong

bootstrap support that are distinct from the major group of *Ca.* Desulfofervidus (Fig. 5). These

clusters were separated from the main group by sequence divergence up to 5.8% (Suppl. Data

File 1, Sheet 2). The recovery of these sequences across independent studies at different vent

sites suggests unrecognized microbial diversity related to *Ca.* Desulfofervidus. A previous

metagenomic survey of Guaymas Basin hydrothermal sediments detected metagenome-

assembles genomes (MAGs) branching between *Ca.* Desulfofervidus and

Thermodesulfobacteriaceae (Dombrowski et al., 2018). Detailed analyses of these MAGs

(labeled as B74_G16 and B4_G9) would reveal the genomic content and physiological potential

of Desulfofervidus-affiliated lineages.

SEEP-SRB2. Like *Ca.* Desulfofervidus, SEEP-SRB2 sequences are often recovered in

Guaymas Basin sequencing surveys (Dowell et al., 2016). Our survey recovered sequences from

slightly below the surface to approximately 15 cmbsf. Unlike *Ca.* Desulfofervidus, SEEP-SRB2

sequences are recovered from diverse non-hydrothermal environments; cold seeps, salt lakes,

coastal sediments, and estuaries (Kleindienst et al., 2012, Ruff et al., 2015). Despite its

widespread occurrence and likely ecological importance as a syntrophic partner of methane and

alkane oxidizers (Krukenberg et al., 2018), the internal phylogenetic structure of this group has

not been examined in any detail.

Our SEEP-SRB2 phylogeny is largely comprised of a well-supported major cluster with

~97.5% sequence identity (Suppl. Data File 1, Sheet 3, equivalent to *Genus* level; Schloss &

Handelsman, 2005). The closest cultured relatives of SEEP-SRB2, *Dissulfuribacter thermophilus*

and *Dissulfurimicrobium hydrothermale*, are thermophilic sulfur disproportionators isolated from

hydrothermal environments (Slobodkin et al., 2013, Slobodkin et al., 2016). These bacteria

constitute separate lineages but share a deep branching point with SEEP-SRB2 (Fig. 6). The

distinctiveness of these lineages is further supported by sequence distances of up to 11% (Suppl.

Data File 1, Sheet 3). Our 16S rRNA gene sequences indicate that SEEP-SRB2 would form a

stand-alone group within *Dissulfuribacteraceae* (Waite et al., 2020). Further phylogenetic

diversity that would fall between the pure culture examples (*Dissulfuribacter thermophilus* and

Dissulfurimicrobium hydrothermale) and SEEP-SRB2 remains to be explored. Previously, the

creation of a taxonomic family to comprise *Dissulfuribacter thermophilus* and

Dissulfurimicrobium hydrothermale has been suggested (Slobodkin et al., 2013). Our phylogeny

shows that such a group would also include SEEP-SRB2 (Fig. 6).

SEEP-SRB4. In contrast to the syntrophic symbionts SEEP-SRB1 and SEEP-SRB2,

SEEP-SRB4 has largely been overlooked since it was proposed 20 years ago (Knittel et al.,

2003). FISH surveys indicate that SEEP-SRB4 is not a syntroph, but occurs as individual free-

living cells (Knittel et al., 2003). The closest cultured relatives of SEEP-SRB4 are non-

thermophilic sulfur-disproportionating genera such as *Desulforhopalus* (Isaksen & Teske, 1996),

Desulfocapsa (Finster et al., 1998), *Desulfofustis* (Friedrich et al., 1996), and the sulfate-reducing

psychrophile *Desulfotalea* (Knoblauch et al., 1999). While SEEP-SRB2 has been enriched from

Guaymas Basin sediments at 37ºC (Krukenberg et al., 2018), there is currently no evidence that

SEEP-SRB4 is moderately thermotolerant. We seek to test this assumption in our survey,

checking the distribution of SEEP-SRB4 in surficial hydrothermal sediments. As with SEEP-

SRB2, we seek to investigate the previously unrecognized phylogenetic diversity of SEEP-

SRB4.

Our SEEP-SRB4 phylogeny depicts one major cluster with strong bootstrap support

(100 %) and > 98.9% sequence identity (Suppl. Data File 1, Sheet 4). This cluster contains

several well-supported, phylogenetically narrow clusters (Fig. 7). While the phylogeny shows a

potential Guaymas Basin-specific subcluster of four sequences, the other subclusters contain

globally distributed sequences from diverse cold seep sites and from Guaymas Basin.

The majority of L-ASVs for SEEP-SRB4 are retrieved from the top ~6 cm of the

sediment across the entire lateral hydrothermal gradient (Suppl. Table 3). The preference for

surficial sediments may be due to the abundance of elemental sulfur and polysulfides located

there (Teske et al., 2016). A similar preference for surficial sediments with coexisting sulfate and

sulfide was noted in the first study of SEEP-SRB4 (Knittel et al., 2003). SEEP-SRB4 bacteria

seem to be able to colonize surficial hydrothermal sediments as long as in-situ temperatures

remain relatively cool, similar to the strategy utilized by Guaymas Basin *Beggiatoaceae* (McKay

et al., 2012).

Desulfatiglans. *Desulfatiglandales* is an order-level lineage within the phylum

Desulfobacterota that contains polyaromatic degrading organisms, both isolated and uncultured

species. The order and family level taxonomy are named after *Desulfatiglans annilini*, the first

cultured representative (Suzuki et al., 2014). Sequences related to *Desulfatiglans* are often

recovered in Guaymas Basin sequencing surveys (Ramírez et al., 2021, Dowell et al., 2016),

including the shallow subsurface (Edgcomb et al., 2022). In contrast to the tightly clustered

groups of SEEP-SRB2 and SEEP-SRB4, *Desulfatiglans* exhibits a greater degree of phylogenetic

diversity.

Our Guaymas Basin sequences and almost all GenBank sequences were not closely

related to cultured isolates that degrade polyaromatic and substituted-aromatic compounds (Fig.

7). Even finding uncultured sequences that were closely related (>98% sequence identity) to

some of our Guaymas Basin ASVs proved to be difficult. This could imply an undersampling of

the natural diversity of *Desulfatiglans*, or their ongoing diversification and speciation.

Previously, *Desulfatiglans* has been classified into two groups distinguished by genomic and

physiological differences, Group 1 consisting of sulfate reducers and Group 2 consisting of

dehalogenating subsurface bacteria (Jochum et al., 2018). Most of our L-ASVs belong to Group

1 and a few belong to Group 2.

Outside of these two groups, we found two well-supported hydrothermal clusters (Fig. 8).

The first cluster contained L-ASV 190 (predominately recovered from the white mat site),

sequences from 37 °C and 50 °C enrichments of Guaymas Basin sediment (Kellermann et al.,

2012, Wegener et al., 2015) and an environmental sequence from Guaymas Basin hydrothermal

sediment (McKay et al., 2016). The second cluster contained L-ASV 207 (predominately

recovered from the orange mat site), a sequence from a *Desulfatiglans* MAG that was prevalent

in a 70 °C enrichment of Guaymas Basin sediment (Zehnle et al., 2023), and environmental

sequences from diverse hydrothermal sediments (Li et al., 2014, Li et al., 2020). We examined

these sequences via distance matrix analyses and find that these hydrothermal clusters are

generally separated from other sequences by 11-12% sequence divergence (Suppl. Data File 1,

Sheet 5).

In contrast to other sulfate reducers that occur in warm and hot Guaymas Basin

sediments, we observed most of our *Desulfatiglans* L-ASVs in cool sediments, suggesting that

they do not depend on methane and light hydrocarbons derived from hydrothermal fluids, and

instead utilize polyaromatics (Schnell & Schink, 1991, Zehnle et al., 2023). Polyaromatics from

hydrothermal oils are prevalent throughout Guaymas Basin sediments of diverse thermal regimes

(Kawka & Simoneit, 1990), and *Desulfatiglans* appears to have evolved to utilize polyaromatics

across the thermal spectrum, as suggested by its hydrothermal clusters.

Chimera Issues. DADA2 detected chimeras in approximately 4% of our reads from a

natural hydrothermal community. Previous research found structural errors (chimeras and

introgressions) in approximately 2% of reads from a Zymo mock community (Callahan et al.,

2021). Specifically, we found that LoopSeq produced chimeras in the majority of *Candidatus*

Desulfofervidus sequences we examined. Segments of these sequences were entered into

BLASTn, and we discovered that the chimeras consistently occurred in the first ~200 bp.

BLASTn searches of the chimeric segments revealed their relation to other marine sediment

bacteria such as *Chloroflexi*. The chimera breaks are located within the highly variable Helix 11

of the 16S rRNA gene (Van de Peer, et al. 1996), and therefore appear to be secondary structure

dependent. These issues with *Ca.* Desulfofervidus sequences were observed in samples across

the DNA concentration range, from very high (4240 ng/cm^3) to very low (0.5 ng/cm^3).

Automated chimera identifying programs can have sensitivity issues when used with uncultured

lineages that have poorly resolved phylogenies. We found that trimming off the first ~200 bp

remedied the chimera issue, resulting in sequences with high alignment quality and >97%

identity to previously archived *Ca.* Desulfofervidus sequences available in GenBank.

LoopSeq recovered eight sequences identified by the DADA2 annotation pipeline as

"*Thermodesulfobacterium*". Initial analysis of theses sequences showed that they were plagued

with chimera and artifact issues, a potential consequence of low DNA yields from deep, hot

samples. Two sequences were salvaged; one did not have chimera issues (L-ASV 12240), while

the other was trimmed to only include the first ~340 bp (L-ASV 11484). These two sequences

came from samples with low DNA concentrations (Sample 4872-06-08, 50 ng/cm^3 and Sample

4872-06-09, 12 ng/cm^3). These two sequences formed a Guaymas Basin specific cluster (Suppl.

Fig. 5) with the thermophilic ANME-1 syntroph *Thermodesulfobacterium torris* (Benito Merino

et al., 2022) and a nearly identical hydrothermal sediment clone (McKay et al., 2016).

Conclusions. Our sequencing survey has shown that LoopSeq is a promising option for

the high throughput long read sequencing of diverse microbial communities in hydrothermal

sediments. The huge quantity of sequences yielded by LoopSeq allowed us to further resolve the

fine scale phylogenetic structure of sulfur-cycling bacterial groups that are characteristic of

Guaymas Basin hydrothermal sediments. The five well-supported clusters revealed in our *Ca.*

Desulfofervidus phylogeny may represent unrecognized physiological and genomic diversity that

calls for further investigation beyond the initial description of this group (Krukenberg et al.,

2016). Investigating SEEP-SRB2, SEEP-SRB4, and *Ca.* Desulfofervidus showed that they are all

phylogenetically distinct groups without a spectrum of close relatives among environmental

clones as represented in NCBI GenBank (Figs. 5, 6, & 7). SEEP-SRB2, SEEP-SRB4, and *Ca.*

Desulfofervidus all contain considerable internal phylogenetic diversity within their respective

clades. Sulfur-disproportionating species and genera are among the closest cultured relatives of

SEEP-SRB2 and SEEP-SRB4 (Figs. 6 & 7). In our distance analyses we see sequences distances

of up to ~8% between sulfur-disproportionating bacteria and SEEP-SRB2 and up to ~5%

between sulfur-disproportionating bacteria and SEEP-SRB4 (Suppl. Data File 1, Sheets 3 & 4).

The near-full-length sequences generated by LoopSeq enabled these distance analyses.

Elemental sulfur and sulfur species of intermediate oxidation states are abundant at seep sites

(Jørgensen et al., 2019). This provides a favorable environment for sulfur-disproportionating

bacteria, and for the diversification of sulfur-oxidizing and sulfate-reducing bacterial lineages.

However, the chimera issues that we detected in sequences identified as *Ca.*

Desulfofervidus, *Thermodesulfobacterium*, and *Beggiatoaceae*, underscore the need for

phylogenetic analyses and alignment comparisons with high quality reference sequences to

confirm the quality and identity of LoopSeq sequences, and the reliability of microbial

community analyses. We also note that our DADA2 analysis of these sequences was not

sufficient to detect these chimeras. Without manual verification of sequence alignments and

scrutiny of isolated distal branches in phylogenetic clusters, problematic sequences can go

undetected, and create artificial diversity. Our observation of lineage-specific chimera issues

calls for caution when using LoopSeq alone to investigate extreme and unusual bacterial groups.

Since we submitted our samples for sequencing (early 2022), Element Biosciences has instituted

changes with the handling of low-biomass samples (such as our own) and has adapted the

chemistry of LoopSeq to improve DNA yield in low biomass samples (Element Biosciences

representative Dr. Caroline Obert, personal communication). Further protocol development of

LoopSeq should monitor this issue, and check sequence accuracy among a wide range of

microbial target groups.

LoopSeq has successfully enabled a survey of the diversity of sulfur-cycling bacteria

across a thermal and substrate gradient up to the current limit of hydrocarbon degradation (Fig.

9). Due to the scope of this study, the diversity of archaea in surficial Guaymas Basin sediments

was not investigated with LoopSeq. Further work may use LoopSeq to investigate potentially

unknown archaeal diversity in habitats with extreme conditions or along environmental gradients

(e.g., Guaymas Basin). Phylogenetic resolution and range of archaeal lineages is limited by

sequence coverage. Due to the limitations of popular primers, most available archaeal 16S rRNA

gene sequences are 800 bp in length or less (Teske & Sørensen, 2008). High quality near full-

length 16S rRNA gene sequence data, such as that provided by LoopSeq, would allow for an

improved view of archaeal diversity.

Data Availability Statement. All sequence reads were submitted to the NCBI Sequence

Read Archive (BioProject: PRJNA1105367). L-ASV sequences used in our phylogenetic trees

are available in Supplementary Data File 2 or via their NCBI GenBank accession number

provided in each phylogenetic tree.

Author Contributions. JEH analyzed the sequence data, created the figures, isolated the

DNA, and wrote the manuscript. JPC performed the methane concentration and stable isotope

analyses. MAM wrote the DADA2 script used to analyze the sequence data and prepared

samples for submission. SER prepared samples for submission and handled interactions with

Element Biosciences over time and advocated on the group's behalf. AT obtained the samples

during Alvin Dive 4872, guided the interpretation of the phylogenetic trees and geochemical

data, supported the development of the manuscript, and taught JEH the DNA extraction method.

All authors contributed to critical revisions of the manuscript.

Funding. This study was supported by an NSF Biological Oceanography grant to AT

(2048489), and by a grant from the Simons Foundation to SER (824763).

Acknowledgements. We thank the R/V *Atlantis* and HOV *Alvin* crews for their

exemplary support at sea, especially *Alvin* pilot Jefferson Grau for expert handling of the craft.

We thank Chris Chambers for conducting the sulfide analyses, and Catherine Crowley for clean-

up and quantification of the DNA. We thank Element Biosciences for their patience with our

complex environmental samples.

Conflict of Interest Statement. The authors declare that the research was conducted in

the absence of any commercial or financial relationships that could be construed as a potential

conflict of interest.

504
505 **Figure 1.** Seafloor image of sampling mat area at Dive 4872. Approximate location of push
506 coring sites are labeled in white. Image courtesy of HOV *Alvin* Dive Team.

507
Figure 2. Plots of in-situ temperatures and geochemical parameters for sediment cores. Sulfide,
sulfate, Ammonium, and Nitrite/Nitrate concentrations obtained from porewater. Methane
concentrations and δ¹³C values obtained from whole sediment. Ammonium and Nitrite/Nitrate

concentrations are not available for the bare sediment site (Core 4872-01). Data for the bare
sediment site (Core 4872-01) are plotted in brown. Data for the white mat site (Core 4872-06)
are plotted in blue. Data for the orange mat site (Core 4872-14) are plotted in orange. Plotted
data are available in Supplementary Table 1.

515
516   **Figure 3.** Stacked bar plots of A) phylum-level and B) order-level bacterial diversity. Phyla that

comprise more than 3% of total sequence reads and Orders comprising more than 10% of total
sequence reads are displayed. These percentage cutoffs were chosen to represent a balance
between taxonomic resolution and figure legibility. The X-axis represents depth below seafloor
(in centimeters) for the three sediments cores across the mat transect (bare sediment 4872-01,
white mat 4872-06, orange mat 4872-14).

Figure 4. 16S rRNA gene sequence based phylogenetic tree of *Beggiatoaceae* and related
lineages. Bootstrap values greater than 70% are shown. Two distinct clusters are highlighted, 1)
an orange filament cluster composed of L-ASVs from the orange mat site (4872-14) and a
reference orange filament sequence obtained from GenBank and 2) a cluster composed of L-

527   ASVs from the white mat site (4872-06) and a reference white filament sequence obtained from
528   GenBank. A version of this tree with expanded L-ASV clusters is available as Supplementary
529   Figure 2.

530
Figure 5. 16S rRNA gene sequence based phylogenetic tree of *Candidatus* Desulfofervidus and
related lineages. Bootstrap values greater than 70% are shown. NCBI GenBank accession
numbers are provided in each tip label. L-ASV tip labels are color coded according to the site
where the majority (2/3) of reads were recovered. This phylogenetic tree is based on shorter
sequences where the first ~200 bp were discarded due to chimera issues. **Brown** = bare
sediment, 4872-01; **Blue** = white mat site, 4872-06; **Orange** = orange mat, 4872-14; **Grey** = no
site had a 2/3+ share of reads.

Figure 6. 16S rRNA gene sequence based phylogenetic tree of SEEP-SRB2 and related lineages. Bootstrap values greater than 70% are shown. NCBI GenBank accession numbers are provided in each tip label. L-ASV tip labels are color coded according to the site where the majority (2/3) of reads were recovered. This phylogenetic tree is based on shorter sequences where the first ~200 bp were discarded due to chimera issues. **Brown** = bare sediment, 4872-01; **Blue** = white mat site, 4872-06; **Orange** = orange mat, 4872-14; **Grey** = no site had a 2/3+ share of reads. *Thermodesulfobacterium torris 16S rRNA sequence was obtained from an annotated genome

546   (BioSample ID: SAMN27514933).
547   *Thermodesulfobacterium syntrophicum 16S rRNA sequence was obtained from an annotated
548   genome (BioSample ID: SAMN29995626).

549
Figure 7. 16S rRNA gene sequence based phylogenetic tree of SEEP-SRB4 and related lineages.
Bootstrap values greater than 70% are shown. NCBI GenBank accession numbers are provided
in each tip label. L-ASV tip labels are color coded according to the site where the majority (2/3)
of reads were recovered. This phylogenetic tree is based on shorter sequences where the first
~200 bp were discarded due to chimera issues. **Brown** = bare sediment, 4872-01; **Blue** = white
mat site, 4872-06; **Orange** = orange mat, 4872-14; **Grey** = no site had a 2/3+ share of reads.

Figure 8. 16S rRNA gene sequence based phylogenetic tree of *Desulfatiglans* and related lineages. Bootstrap values greater than 70% are shown. NCBI GenBank accession numbers are provided in each tip label. L-ASV tip labels are color-coded according to the site where the majority (2/3) of reads were recovered. This phylogenetic tree is based on shorter sequences where the first ~200 bp were discarded due to chimera issues. **Brown** = bare sediment, 4872-01; **Blue** = white mat site, 4872-06; **Orange** = orange mat, 4872-14; **Grey** = no site had a 2/3+ share of reads. The Guaymas 70°C enrichment sequence, marked with an asterisk (*), is from MAG 9 (B70, Benzene enrichment at 70°C) under BioProject PRJNA1013425.

Figure 9. Conceptual figure of sulfur-cycling bacterial groups (with examples of preferred substrates) and their approximate distribution along a lateral hydrothermal gradient in surficial Guaymas Basin sediments. Examples of preferred substrates are shown for both free-living and syntrophic lifestyles, when applicable. Molecular structure images of the preferred substrates were obtained from chemspider.com. *Beggiatoaceae* mat and sediment close up images were captured with the bottom facing camera of HOV *Alvin* during Dive 4872.

References

1. Benito Merino, D., Zehnle, H., Teske, A. and Wegener, G., 2022. Deep-branching ANME-1c archaea grow at the upper temperature limit of anaerobic oxidation of methane. *Frontiers in Microbiology*, 13, 988871. doi: 10.3389/fmicb.2022.988871.

2. Callahan, B.J., Grinevich, D., Thakur, S., Balamotis, M.A. and Yehezkel, T.B., 2021. Ultra-accurate microbial amplicon sequencing with synthetic long reads. *Microbiome*, 9(1), 130. doi: 10.1186/s40168-021-01072-3.

3. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. and Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, 13(7), pp.581-583. doi: doi.org/10.1038/nmeth.3869.

4. Calvert, S.E., 1966. Origin of diatom-rich, varved sediments from the Gulf of California. *The Journal of Geology*, 74(5, Part 1), 546-565. doi: 10.1086/627188.

5. Cline, J.D., 1969. Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnology and Oceanography*, 14(3), 454-458.

6. De la Lanza-Espino, G. and Soto, L.A., 1999. Sedimentary geochemistry of hydrothermal vents in Guaymas Basin, Gulf of California, Mexico. *Applied Geochemistry*, 14(4), 499-510. doi: 10.1016/S0883-2927(98)00064-X.

7. Dombrowski, N., Teske, A.P. and Baker, B.J., 2018. Expansive microbial metabolic versatility and biodiversity in dynamic Guaymas Basin hydrothermal sediments. *Nature Communications*, 9, 4999. doi: 10.1038/s41467-018-07418-0.

8. Dowell, F., Cardman, Z., Dasarathy, S., Kellermann, M.Y., Lipp, J.S., Ruff, S.E., Biddle, J.F., McKay, L.J., MacGregor, B.J., Lloyd, K.G. and Albert, D.B., 2016. Microbial communities in methane-and short chain alkane-rich hydrothermal sediments of Guaymas Basin. *Frontiers in Microbiology*, 7. doi: 10.3389/fmicb.2016.00017

9. Edgcomb, V.P., Teske, A.P. and Mara, P., 2022. Microbial hydrocarbon degradation in Guaymas Basin—exploring the roles and potential interactions of fungi and sulfate-reducing bacteria. *Frontiers in Microbiology*, 13, 831828. doi: 10.3389/fmicb.2022.831828.

10. Einsele, G., Gieskes, J.M., Curray, J., Moore, D.M., Aguayo, E., Aubry, M.P., Fornari, D., Guerrero, J., Kastner, M., Kelts, K. and Lyle, M., 1980. Intrusion of basaltic sills into highly porous sediments and resulting hydrothermal activity. *Nature*, 283, 441-445. doi: 10.1038/283441a0.

11. Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 39(4), 783-791. doi: 10.1111/j.1558-5646.1985.tb00420.x.

12. Finster, K., Liesack, W. and Thamdrup, B.O., 1998. Elemental sulfur and thiosulfate disproportionation by *Desulfocapsa sulfoexigens* sp. nov., a new anaerobic bacterium isolated from marine surface sediment. *Applied and Environmental Microbiology*, 64(1), 119-125. doi: 10.1128/AEM.64.1.119-125.1998.

13. Goetz, F.E. and Jannasch, H.W., 1993. Aromatic hydrocarbon-degrading bacteria in the petroleum-rich sediments of the Guaymas Basin hydrothermal vent site: preference for aromatic carboxylic acids. *Geomicrobiology Journal*, 11(1), 1-18. doi: doi.org/10.1080/01490459309377928

14. Gutierrez, T., Biddle, J.F., Teske, A. and Aitken, M.D., 2015. Cultivation-dependent and cultivation-independent characterization of hydrocarbon-degrading bacteria in Guaymas Basin sediments. *Frontiers in Microbiology*, 6, 149000. doi: 10.3389/fmicb.2015.00695.

15. Holler, T., Widdel, F., Knittel, K., Amann, R., Kellermann, M.Y., Hinrichs, K.U., Teske, A., Boetius, A. and Wegener, G., 2011. Thermophilic anaerobic oxidation of methane by marine microbial consortia. *The ISME Journal*, 5(12), 1946-1956. doi: 10.1038/ismej.2011.77.

16. Isaksen, M.F. and Teske, A., 1996. *Desulforhopalus vacuolatus* gen. nov., sp. nov., a new moderately psychrophilic sulfate-reducing bacterium with gas vacuoles isolated from a temperate estuary. *Archives of Microbiology*, 166, 160-168. doi: 10.1007/s002030050371.

17. Jochum, L.M., Schreiber, L., Marshall, I.P., Jørgensen, B.B., Schramm, A. and Kjeldsen, K.U., 2018. Single-cell genomics reveals a diverse metabolic potential of uncultivated *Desulfatiglans*-related Deltaproteobacteria widely distributed in marine sediment. *Frontiers in Microbiology*, 9, 384358. doi: 10.3389/fmicb.2018.02038

18. Jørgensen, B.B., Findlay, A.J. and Pellerin, A., 2019. The biogeochemical sulfur cycle of marine sediments. *Frontiers in Microbiology*, 10, 436320. doi: 10.3389/fmicb.2019.00849.

19. Kawka, O.E. and Simoneit, B.R., 1990. Polycyclic aromatic hydrocarbons in hydrothermal petroleums from the Guaymas Basin spreading center. *Applied Geochemistry*, 5(1-2), 17-27. doi: 10.1016/0883-2927(90)90032-Z.

20. Kellermann, M.Y., Wegener, G., Elvert, M., Yoshinaga, M.Y., Lin, Y.S., Holler, T., Mollar, X.P., Knittel, K. and Hinrichs, K.U., 2012. Autotrophy as a predominant mode of carbon fixation in anaerobic methane-oxidizing microbial communities. *Proceedings of the National Academy of Sciences*, 109(47), 19321-19326. doi: 10.1073/pnas.1208795109.

21. Kleindienst, S., Herbst, F.A., Stagars, M., Von Netzer, F., Von Bergen, M., Seifert, J., Peplies, J., Amann, R., Musat, F., Lueders, T. and Knittel, K., 2014. Diverse sulfate-reducing bacteria of the *Desulfosarcina/Desulfococcus* clade are the key alkane degraders at marine seeps. *The ISME Journal*, 8(10), 2029-2044. doi: 10.1038/ismej.2014.51.

22. Kniemeyer, O., Musat, F., Sievert, S.M., Knittel, K., Wilkes, H., Blumenberg, M., Michaelis, W., Classen, A., Bolm, C., Joye, S.B. and Widdel, F., 2007. Anaerobic oxidation of short-chain hydrocarbons by marine sulphate-reducing bacteria. *Nature*, 449, 898-901. doi: 10.1038/nature06200.

23. Knittel, K., Boetius, A., Lemke, A., Eilers, H., Lochte, K., Pfannkuche, O., Linke, P. and Amann, R., 2003. Activity, distribution, and diversity of sulfate reducers and other bacteria in sediments above gas hydrate (Cascadia Margin, Oregon). *Geomicrobiology Journal*, 20(4) 269-294. doi: 10.1080/01490450303896.

24. Knoblauch, C., Sahm, K. and Jørgensen, B.B., 1999. Psychrophilic sulfate-reducing bacteria isolated from permanently cold Arctic marine sediments: description of *Desulfofrigus oceanense* gen. nov., sp. nov., *Desulfofrigus fragile* sp. nov., *Desulfofaba gelida* gen. nov., sp. nov., *Desulfotalea psychrophila* gen. nov., sp. nov. and *Desulfotalea arctica* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 49(4), 1631-1643. doi: 10.1099/00207713-49-4-1631.

25. Krukenberg, V., Harding, K., Richter, M., Glöckner, F.O., Gruber-Vodicka, H.R., Adam, B., Berg, J.S., Knittel, K., Tegetmeyer, H.E., Boetius, A. and Wegener, G., 2016. *Candidatus* Desulfofervidus auxilii, a hydrogenotrophic sulfate-reducing bacterium involved in the thermophilic anaerobic oxidation of methane. *Environmental Microbiology*, 18(9), 3073-3091. doi: 10.1111/1462-2920.13283.

26. Krukenberg, V., Riedel, D., Gruber‑Vodicka, H.R., Buttigieg, P.L., Tegetmeyer, H.E., Boetius, A. and Wegener, G., 2018. Gene expression and ultrastructure of meso- and thermophilic methanotrophic consortia. *Environmental Microbiology*, 20(5), 1651-1666. doi: 10.1111/1462-2920.14077.

27. Lagostina, L., Frandsen, S., MacGregor, B.J., Glombitza, C., Deng, L., Fiskal, A., Li, J., Doll, M., Geilert, S., Schmidt, M. and Scholz, F., 2021. Interactions between temperature and energy supply drive microbial communities in hydrothermal sediment. *Communications Biology*, 4(1), 1006. doi: 10.1038/s42003-021-02507-1.

28. Laso-Pérez, R., Wegener, G., Knittel, K., Widdel, F., Harding, K.J., Krukenberg, V., Meier, D.V., Richter, M., Tegetmeyer, H.E., Riedel, D. and Richnow, H.H., 2016. Thermophilic archaea activate butane via alkyl-coenzyme M formation. *Nature*, 539, 396-401. doi: 10.1038/nature20152.

29. Li, J., Su, L., Wang, F., Yang, J., Gu, L., Sun, M., Li, Q., Zhou, H. and Fang, J., 2020. Elucidating the biomineralization of low-temperature hydrothermal precipitates with varying Fe, Si contents: Indication from ultrastructure and microbiological analyses. *Deep Sea Research Part I: Oceanographic Research Papers*, 157, 103208. doi: 10.1016/j.dsr.2019.103208.

30. Li, J., Zhou, H., Fang, J., Sun, Y. and Dasgupta, S., 2014. Microbial distribution in different spatial positions within the walls of a black sulfide hydrothermal chimney. *Marine Ecology Progress Series*, 508, 67-85. doi: 10.3354/meps10841.

31. Lizarralde, D., Soule, S.A., Seewald, J.S. and Proskurowski, G., 2011. Carbon release by off-axis magmatism in a young sedimented spreading centre. *Nature Geoscience*, 4(1), 50-54. doi: 10.1038/ngeo1006.

32. Mara, P., Nelson, R.K., Reddy, C.M., Teske, A. and Edgcomb, V.P., 2022. Sterane and hopane biomarkers capture microbial transformations of complex hydrocarbons in young hydrothermal Guaymas Basin sediments. *Communications Earth & Environment*, 3(1), 250. doi: 10.1038/s43247-022-00582-8.

33. McKay, L., Klokman, V.W., Mendlovitz, H.P., LaRowe, D.E., Hoer, D.R., Albert, D., Amend, J.P. and Teske, A., 2016. Thermal and geochemical influences on microbial biogeography in the hydrothermal sediments of Guaymas Basin, Gulf of California. *Environmental Microbiology Reports*, 8(1), 150-161. doi: 10.1111/1758-2229.12365.

34. McKay, L.J., MacGregor, B.J., Biddle, J.F., Albert, D.B., Mendlovitz, H.P., Hoer, D.R., Lipp, J.S., Lloyd, K.G. and Teske, A.P., 2012. Spatial heterogeneity and underlying geochemistry of phylogenetically diverse orange and white Beggiatoa mats in Guaymas Basin hydrothermal sediments. *Deep Sea Research Part I: Oceanographic Research Papers*, 67, 21-31. doi: 10.1016/j.dsr.2012.04.011.

35. McMurdie, P.J. and Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS One*, 8(4). doi: 10.1371/journal.pone.0061217.

36. Meyer, S., Wegener, G., Lloyd, K.G., Teske, A., Boetius, A. and Ramette, A., 2013. Microbial habitat connectivity across spatial scales and hydrothermal temperature gradients at Guaymas Basin. *Frontiers in Microbiology*, 4, 51282. doi: 10.3389/fmicb.2013.00207.

37. Møller, M.H., Glombitza, C., Lever, M.A., Deng, L., Morono, Y., Inagaki, F., Doll, M., Su, C.C. and Lomstein, B.A., 2018. D: L-amino acid modeling reveals fast microbial

turnover of days to months in the subsurface hydrothermal sediment of Guaymas Basin. *Frontiers in Microbiology*, 9, e967. doi: 10.3389/fmicb.2018.00967.

38. Pérez Castro, S., Borton, M.A., Regan, K., Hrabe de Angelis, I., Wrighton, K.C., Teske, A.P., Strous, M. and Ruff, S.E., 2021. Degradation of biological macromolecules supports uncultured microbial populations in Guaymas Basin hydrothermal sediments. *The ISME Journal*, 15(12), 3480-3497. doi: 10.1038/s41396-021-01026-5.

39. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590-D596. doi: 10.1093/nar/gks1219.

40. R Core Team, 2022. *R: A Language and Environment for Statistical Computing*, Vienna, Austria. Available at: https://www.R-project.org/

41. Ramírez, G.A., Mara, P., Sehein, T., Wegener, G., Chambers, C.R., Joye, S.B., Peterson, R.N., Philippe, A., Burgaud, G., Edgcomb, V.P. and Teske, A.P., 2021. Environmental factors shaping bacterial, archaeal and fungal community structure in hydrothermal sediments of Guaymas Basin, Gulf of California. *PLoS One*, *16*(9). doi: 10.1371/journal.pone.0256321.

42. Ruff, S.E., Biddle, J.F., Teske, A.P., Knittel, K., Boetius, A. and Ramette, A., 2015. Global dispersion and local diversification of the methane seep microbiome. *Proceedings of the National Academy of Sciences*, 112(13), 4015-4020. doi: 10.1073/pnas.1421865112.

43. Rzhetsky, A. and Nei, M., 1995. Tests of applicability of several substitution models for DNA sequence data. *Molecular Biology and Evolution*, 12(1), 131-151. doi: 10.1093/oxfordjournals.molbev.a040182.

44. Salman, V., Amann, R., Girnth, A.C., Polerecky, L., Bailey, J.V., Høgslund, S., Jessen, G., Pantoja, S. and Schulz-Vogt, H.N., 2011. A single-cell sequencing approach to the classification of large, vacuolated sulfur bacteria. *Systematic and Applied Microbiology*, 34(4), 243-259. doi: 10.1016/j.syapm.2011.02.001.

45. Schloss, P.D. and Handelsman, J., 2005. Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Applied and Environmental Microbiology*, 71(3), 1501-1506. doi: 10.1128/aem.71.3.1501-1506.2005.

46. Schnell, S. and Schink, B., 1991. Anaerobic aniline degradation via reductive deamination of 4-aminobenzoyl-CoA in *Desulfobacterium anilini*. *Archives of Microbiology*, 155, 183-190. doi: 10.1007/BF00248615.

47. Schutte, C.A., Teske, A., MacGregor, B.J., Salman-Carvalho, V., Lavik, G., Hach, P. and de Beer, D., 2018. Filamentous giant *Beggiatoaceae* from the Guaymas Basin are capable of both denitrification and dissimilatory nitrate reduction to ammonium. *Applied and Environmental Microbiology*, 84(15). doi: 10.1128/AEM.02860-17.

48. Slobodkin, A.I., Reysenbach, A.L., Slobodkina, G.B., Kolganova, T.V., Kostrikina, N.A. and Bonch-Osmolovskaya, E.A., 2013. *Dissulfuribacter thermophilus* gen. nov., sp. nov., a thermophilic, autotrophic, sulfur-disproportionating, deeply branching deltaproteobacterium from a deep-sea hydrothermal vent. *International Journal of Systematic and Evolutionary Microbiology*, 63(6), 1967-1971. doi: 10.1099/ijs.0.046938-0.

49. Slobodkin, A.I., Slobodkina, G.B., Panteleeva, A.N., Chernyh, N.A., Novikov, A.A. and Bonch-Osmolovskaya, E.A., 2016. *Dissulfurimicrobium hydrothermale* gen. nov., sp.

nov., a thermophilic, autotrophic, sulfur-disproportionating deltaproteobacterium isolated from a hydrothermal pond. *International Journal of Systematic and Evolutionary Microbiology*, 66(2), 1022-1026. doi: 10.1099/ijsem.0.000828.

50. Sogin, M.L., Morrison, H.G., Huber, J.A., Welch, D.M., Huse, S.M., Neal, P.R., Arrieta, J.M. and Herndl, G.J., 2006. Microbial diversity in the deep sea and the underexplored "rare biosphere". *Proceedings of the National Academy of Sciences*, 103(32), 12115-12120. doi: 10.1073/pnas.0605127103.

51. Song, M., Schubotz, F., Kellermann, M.Y., Hansen, C.T., Bach, W., Teske, A.P. and Hinrichs, K.U., 2021. Formation of ethane and propane via abiotic reductive conversion of acetic acid in hydrothermal sediments. *Proceedings of the National Academy of Sciences*, 118(47). doi: 10.1073/pnas.2005219118.

52. Stecher, G., Tamura, K. and Kumar, S., 2020. Molecular evolutionary genetics analysis (MEGA) for macOS. *Molecular Biology and Evolution*, 37(4), 1237-1239. doi: 10.1093/molbev/msz312.

53. Su, L., Teske, A.P., MacGregor, B.J., McKay, L.J., Mendlovitz, H., Albert, D., Ma, Z. and Li, J., 2023. Thermal Selection of Microbial Communities and Preservation of Microbial Function in Guaymas Basin Hydrothermal Sediments. *Applied and Environmental Microbiology*, 89(3). doi: 10.1128/aem.00018-23.

54. Suzuki, D., Li, Z., Cui, X., Zhang, C. and Katayama, A., 2014. Reclassification of *Desulfobacterium anilini* as *Desulfatiglans anilini* comb. nov. within *Desulfatiglans* gen. nov., and description of a 4-chlorophenol-degrading sulfate-reducing bacterium, *Desulfatiglans parachlorophenolica* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 64(9), 3081-3086. doi: 10.1099/ijs.0.064360-0.

55. Tamura, K., Nei, M. and Kumar, S., 2004. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences*, 101(30), 11030-11035. doi: 10.1073/pnas.0404206101.

56. Tamura, K., Stecher, G. and Kumar, S., 2021. MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38(7), 3022-3027. doi: 10.1093/molbev/msab120.

57. Teske, A. and Sørensen, K.B., 2008. Uncultured archaea in deep marine subsurface sediments: have we caught them all?. *The ISME Journal*, 2(1), 3-18. doi: 10.1038/ismej.2007.90.

58. Teske, A., Hinrichs, K.U., Edgcomb, V., de Vera Gomez, A., Kysela, D., Sylva, S.P., Sogin, M.L. and Jannasch, H.W., 2002. Microbial diversity of hydrothermal sediments in the Guaymas Basin: evidence for anaerobic methanotrophic communities. *Applied and Environmental Microbiology*, 68(4), 1994-2007. doi: 10.1128/AEM.68.4.1994-2007.2002.

59. Teske, A., McKay, L.J., Ravelo, A.C., Aiello, I., Mortera, C., Núñez-Useche, F., Canet, C., Chanton, J.P., Brunner, B., Hensen, C. and Ramírez, G.A., 2019. Characteristics and evolution of sill-driven off-axis hydrothermalism in Guaymas Basin–the Ringvent site. *Scientific Reports*, 9(1), 13847. doi: 10.1038/s41598-019-50200-5.

60. Teske, A., Wegener, G., Chanton, J.P., Hoer, D., de Beer, D., Zhuang, G., Saxton, M.A., Joye, S.B., Soule, S.A. and Ruff, S.E., 2021. Microbial communities under distinct thermal and geochemical regimes in axial and off-axis sediments of Guaymas Basin. *Frontiers in Microbiology*, 12, 633649. doi: 10.3389/fmicb.2021.633649.

61. Van de Peer, Y., Chapelle, S. and De Wachter, R., 1996. A quantitative map of nucleotide substitution rates in bacterial rRNA. *Nucleic Acids Research*, 24(17) 3381-3391. doi: 10.1093/nar/24.17.3381.

62. Von Damm, K.L., Edmond, J.T., Measures, C.I. and Grant, B., 1985. Chemistry of submarine hydrothermal solutions at Guaymas Basin, Gulf of California. *Geochimica et Cosmochimica Acta*, 49(11) 2221-2237. doi: 10.1016/0016-7037(85)90223-6.

63. Waite, D.W., Chuvochina, M., Pelikan, C., Parks, D.H., Yilmaz, P., Wagner, M., Loy, A., Naganuma, T., Nakai, R., Whitman, W.B. and Hahn, M.W., 2020. Proposal to reclassify the proteobacterial classes Deltaproteobacteria and Oligoflexia, and the phylum Thermodesulfobacteria into four phyla reflecting major functional capabilities. *International Journal of Systematic and Evolutionary Microbiology*, 70(11), 5972-6016. doi: 10.1099/ijsem.0.004213.

64. Wegener, G., Krukenberg, V., Riedel, D., Tegetmeyer, H.E. and Boetius, A., 2015. Intercellular wiring enables electron transfer between methanotrophic archaea and bacteria. *Nature*, 526, 587-590. doi: 10.1038/nature15733.

65. Welhan, J.A., 1988. Origins of methane in hydrothermal systems. Chemical Geology, 71(1-3), 183-198. doi: 10.1016/0009-2541(88)90114-3.

66. Winkel, M., de Beer, D., Lavik, G., Peplies, J. and Mußmann, M., 2014. Close association of active nitrifiers with Beggiatoa mats covering deep‐sea hydrothermal sediments. *Environmental Microbiology*, 16(6), 1612-1626. doi: 10.1111/1462-2920.12316.

67. Zehnle, H., Otersen, C., Benito Merino, D. and Wegener, G., 2023. Potential for the anaerobic oxidation of benzene and naphthalene in thermophilic microorganisms from the Guaymas Basin. *Frontiers in Microbiology*, 14, 1279865. doi: 10.3389/fmicb.2023.1279865.

68. Zhou, J., Bruns, M.A. and Tiedje, J.M., 1996. DNA recovery from soils of diverse composition. *Applied and Environmental Microbiology*, 62(2), 316-322. doi: 10.1128/aem.62.2.316-322.1996.

69. Zhuang, G.C., Heuer, V.B., Lazar, C.S., Goldhammer, T., Wendt, J., Samarkin, V.A., Elvert, M., Teske, A.P., Joye, S.B. and Hinrichs, K.U., 2018. Relative importance of methylotrophic methanogenesis in sediments of the Western Mediterranean Sea. *Geochimica et Cosmochimica Acta*, 224, 171-186. doi: 10.1016/j.gca.2017.12.024.