1

Metastasis of colon cancer requires Dickkopf-2 to generate cancer cells with Paneth cell properties

Jae Hun Shin[1†*], Jooyoung Park[2†], Jaechul Lim[3], Jaekwang Jeong[4], Ravi K. Dinesh[5], Stephen E. Maher[6], Jun Young Hong[7], John Wysolmerski[4] Jungmin Choi[2] and Alfred L. M. Bothwell[8,9*]

[1]Integrative Science and Engineering Division, Underwood International College, Yonsei University, Incheon 21983, Korea.

[2]Department of Biomedical Sciences, Korea University College of Medicine, Seoul 02841, Korea.

[3]College of Veterinary Medicine, Seoul National University, Seoul 08826, Korea.

[4]Internal Medicine, Yale University School of Medicine, New Haven, Connecticut 06520, U.S.

[5]Department of Pathology, Stanford University, Stanford, California 94305, U.S.

[6]Department of Urology, Yale University School of Medicine, New Haven, Connecticut 06520, U.S.

[7]Department of Systems Biology, College of Life Science and Biotechnology, Yonsei University, Seoul, South Korea.

[8]Dept. of Pathology, Microbiology and Immunology, University of Nebraska Medical Center, 505 S. 45[th] Street., Omaha, NE 68198, U.S.

[9]Dept. of Immunobiology, Yale University School of Medicine, New Haven, Connecticut 06520, U.S.

[†]These authors are equally contributed.

[*]Correspondence to

Alfred L. M. Bothwell, Dept. of Pathology, Microbiology and Immunology, University of Nebraska Medical Center, 986842 Nebraska Medical Center, Fred & Pamela Buffett Cancer Center, 505 S. 45[th] Street, Omaha, NE 68198, 402-559-8050, albothwell@unmc.edu

Jae Hun Shin, Integrative Science and Engineering Division, Underwood International College, Yonsei University, Incheon 21983, South Korea, 82-10-2676-0525, jaehun.shin@yonsei.ac.kr

Running title

Formation of lysozyme positive colon cancer cells by DKK2

Keywords

Abstract

Metastasis is the leading cause of cancer-related mortality. Paneth cells provide stem cell

niche factors in homeostatic conditions, but the underlying mechanisms of cancer stem cell niche

development are unclear. Here we report that Dickkopf-2 (DKK2) is essential for the generation of

cancer cells with Paneth cell properties during colon cancer metastasis. Splenic injection of *Dkk2*-

knockout (KO) cancer organoids into C57BL/6 mice resulted in a significant reduction of liver

metastases. Transcriptome analysis showed reduction of Paneth cell markers such as lysozymes

in KO organoids. Single cell RNA sequencing analyses of murine metastasized colon cancer cells

and patient samples identified the presence of lysozyme positive cells with Paneth cell properties

including enhanced glycolysis. Further analyses of transcriptome and chromatin accessibility

suggested Hepatocyte nuclear factor 4-alpha (HNF4A) as a downstream target of DKK2.

Chromatin immunoprecipitation followed by sequencing analysis revealed that HNF4A binds to

the promoter region of *Sox9*, a well-known transcription factor for Paneth cell differentiation. In

the liver metastatic foci, DKK2 knockout rescued HNF4A protein levels followed by reduction of

lysozyme positive cancer cells. Taken together, DKK2-mediated reduction of HNF4A protein

promotes the generation of lysozyme positive cancer cells with Paneth cell properties in the

metastasized colon cancers.

Introduction

Metastasis is the main cause of cancer-related death (1). Mutations in the *Apc* gene in

intestinal epithelial cells cause adenoma formation by dys-regulation of Wnt signaling (2).

Accumulating oncogenic mutations in *Kras*, *Tp53* and *Smad4* genes facilitate colorectal

carcinogenesis and metastasis (3-5). Recent studies have identified that *Lgr5* positive cells are

required for metastasis in the murine models of colorectal cancer (6, 7). *Lgr5* positive cells

include stem cells, transit amplifying cells and Paneth cells (8). Ablation of *Lgr5* positive cells in

the primary tumor in mice restricted metastatic progression of colon cancer cells (6). Time-lapse

live imaging analysis of metastatic cancer cells in the murine primary colon tumors revealed that

both *Lgr5* positive and negative cancer cells are able to escape from the primary tumors and

circulate in the blood stream (7). Importantly, generation or presence of *Lgr5* positive cells is

necessary to form metastases suggesting that cancer stem cells and their niche formation upon

metastases seeding is a prerequisite of metastatic tumor growth.

The stem cell niche provides so called 'stem cell niche factors' in order to regulate

proliferation and differentiation of stem cells (9, 10). In the homeostatic condition, Paneth cells

function as stem cell niche cells by providing Wnt3, EGF, Notch ligand, and Dll4 to intestinal stem

cells (11). Paneth cell-mediated glucose conversion into lactate promotes proliferation of

intestinal stem cells that lack glucose metabolism (12). Stem cell niche factors not only maintain

stem cells but also generate new stem cells by dedifferentiation from the progenitors when the

original stem cells are damaged by intestinal injury or inflammation (13-15). Regeneration of stem

cells by niche factors also occurs in the context of colorectal cancer (6, 16, 17). However, the

underlying mechanisms of cancer stem cell niche cell development in the primary and metastatic

colon tumors are largely unknown.

In normal mucosa, the balance between Wnt and Notch signaling determines the fate of

intestinal stem cells differentiating to secretory or absorptive precursors (18-20). *Atoh1*

expression in the secretory precursors directs their differentiation to Paneth or goblet cells (21).

Sox9 expression then induces Paneth cell differentiation (22, 23) in the small intestine. Recent

findings has identified the presence of Paneth-like cells in the large intestine (24-26). However,

the existence of Paneth or Paneth-like cells in colon cancers remains unknown. Using colon

cancer organoids carrying mutations in *Apc*, *Kras* and *Tp53* genes, we showed here that *Dkk2*

knockout disrupted Lysozyme (LYZ) positive cell formation in colon cancer organoids as well as

metastatic nodules *in vivo*. Single cell RNA sequencing analyses for mouse and human

metastatic colon cancer samples have identified the existence of LYZ+ cells exhibiting Paneth

cell properties including metabolic support for stem cells (27). Mechanistically, DKK2 protein

deficiency recovered protein levels of Hepatocyte Nuclear Factor 4 alpha (HNF4A) in colon

cancer cells (28). HNF4A binding on the *Sox9* promoter inhibited the formation of LYZ+ cancer

cells exhibiting Paneth cell characteristics. The loss of LYZ+ cells by *Dkk2* knockout rescued

mice from the liver metastasis of colorectal cancer induced by cancer organoid transplantation.

Our findings suggest that DKK2 promotes LYZ+ cell formation exhibiting Paneth cell properties to

develop cancer stem cell niches for outgrowth of metastasized colon cancers.

Materials and Methods

Splenic injection

Generation of AKP, *Dkk2* knockout AKP, *HNF4α1* knockdown and *Dkk2* knockout AKP organoids

and their culture protocols have been published in the previous study (28). We used the

previously reported splenic injection methodology to develop organoid-derived liver metastasis in

mice (29). Cultured organoids were prepared as single cells following 10-30 min of digestion in

TrypLE solution (Thermo, 12605010) at 37°C, followed by centrifugation with advanced DMEM-

F12 (Thermo, 12634010). The isolated single cells were re-suspended with PBS as 300,000 cells

per 100 μl of PBS. Surgical instruments were sterilized by steam autoclave (250 ° F., 15 p.s.i. for

30 minutes). 8-12 weeks old C57BL/6 mice were anesthetized with Ketamine (100 mg/kg) plus

Xylazine (10 mg/kg) via intraperitoneal injection. Puralube Vet ointment was applied to the eyes

to prevent drying while surgery. Spleen was divided by two Horizon medium size ligating clips

(Teleflex, 002200) in the center of the spleen and one hemi-spleen was returned into the

peritoneum. Using a 26-½ Gauge needle, 100-200 μl volume of cells was injected underneath of

the splenic capsule over the course of 30 s. A small Horizon clip (Teleflex, 001200) was applied

to ligate the pancreas and splenic vessels and the injected hemi-spleen was removed. The

incision was closed with a 5-0 running stitch using Vicryl (Ethicon, J493G) and 2-3 skin clips.

Meloxicam (0.3 mg/kg) was injected subcutaneously to alleviate postsurgical pain at 24-72 h

post-surgery.

In vivo **imaging**

Noninvasive *in vivo* imaging was performed using a Microfocus X-ray imaging source (Kodak).

Mice were anesthetized by isoflurane inhalation and their abdominal hair was shaved. Tandem

repeat Tomato (tdTomato) fluorescence in liver metastatic tumors was recorded for 30 sec.

tdTomato signals were statistically analyzed using the Bruker image analysis system. X-ray

image was captured by 10 sec exposure and merged with the tdTomato image.

Data availability

All data are included in the article and SI Appendix. All the sequencing data reported in this study

have been deposited in the Gene Expression Omnibus as follows; Bulk RNA-seq: GSE157531,

ATAC-seq: GSE157529, Chip-seq: will be available before publication. scRNA-seq: GSE157645.

Results

DKK2 is indispensable for liver metastasis of colorectal cancer

Metastatic potential of *Lgr5*-expressing cells has been reported in colorectal cancer (30).

The ablation of *Lgr5*-expressing cells in primary colon tumors inhibited metastasis whereas the

growth of primary tumors quickly recovered by dedifferentiation of non-stem cells into *Lgr5*-

expressing stem cells (6). Moreover, *Fumagalli et al.* have shown that *Lgr5*-expressing cells are

indispensable for the growth of metastases (7). We have recently reported that DKK2 enhanced

Lgr5 expression in colon cancers through activation of c-Src (28). Enhanced activation of c-Src

has been reported in metastatic colon cancers as well (31). Based on these observations, we

queried whether DKK2 is required for metastasis of colorectal cancer. We developed colon

cancer organoids carrying oncogenic mutations in *Apc*, *Kras* and *Tp53* genes (AKP) (28). Splenic

injection of the *Dkk2* knockout AKP (KO) organoids into wild type C57BL/6 mice resulted in

significantly less liver metastasis and mortality compared to the control (AKP) organoids (Fig. 1A-

D). Quantitative gene expression analyses revealed that the levels of *Dkk2* and *Lgr5* were still

decreased in KO-derived liver metastasized cells compared to AKP-derived cells (Fig. 1E). Lgr5

protein expression and c-Src phosphorylation were reduced in KO tumor samples (Fig. 1F and

G). The percentage of *Lgr5*-expressing cells with c-Src phosphorylation in KO tumor tissues was

decreased about 2-fold compared to AKP tissues (Fig. 1H). These data suggest the necessity of

DKK2 in the growth of *Lgr5*-expressing cancer stem cells in liver metastasized nodules originated

from the splenic injection of cancer organoids.

DKK2 is required for lysozyme positive cell formation in colon cancer organoids

145   To understand the underlying mechanisms of the DKK2-mediated increase of *Lgr5*-

expressing stem cells in liver metastases, we performed RNA sequencing (RNA-seq) analysis on

AKP and *Dkk2* knockout (KO) organoids. KO organoids were generated using the CRISPR

technique and DKK2 reconstitution was followed using recombinant mouse DKK2 protein (Fig.

2A). Consistent with our previous report using a colitis-induced tumor model, expression of stem

cell marker genes including *Lgr5* was reduced in KO organoids (Fig. S1A) (28). In particular, the

expression of Paneth cell marker genes such as *Lyz1*, *Lyz2*, and alpha defensins were

significantly reduced in KO organoids (Fig. 2B and S1B). Paneth cells are derived from *Lgr5*-

expressing stem cells by asymmetric division and maintain their *Lgr5* expression (15). These cells

produce stem cell niche factors such as Wnt3A in order to facilitate stem cell proliferation and

differentiation into other epithelial lineage cells (11). Paneth cells are also crucial for the glycolysis

of intestinal stem cells that lack of glucose metabolism (12). Our bulk RNA-seq data showed that

Paneth cell-derived Wnt3a as well as *Fgfr3*, a marker of Paneth cell-induced expansion of stem

cells were reduced by *Dkk2* knockout, indicating that the formation of stem cell niche might be

disrupted in KO organoids (Fig. S1C) (32). Indeed, formation of Lysozyme positive (LYZ+) cells

was disrupted by *Dkk2* knockout during organoid culture (Fig. 2C). LYZ+ cells were detected in

AKP organoids from day 2 after plating a single or multiple cells of enzyme-digested organoids

whereas those cells were less frequently observed in KO organoids. It correlates to the

expression pattern of *Lgr5* in our previous report (28). A significant increase of *Lgr5* expression

was observed between day 2 and day 4 and peaked at day 4 to day 6 in AKP organoid culture.

Lgr5 expression was reduced about 3-fold in KO organoids compared to AKP organoids at day 8

(28). Likewise, the markers of Paneth cells, *Lyz1* and *Lyz2* were reduced in KO organoids at day

8 and recombinant mouse DKK2 protein treatment into KO organoids rescued it (Fig. 2D). These

data indicate that DKK2 is necessary for LYZ+ cell formation in colon cancer organoids that might

be required for the cancer stem cell niche formation.

LYZ+ cancer cells exhibit Paneth cell properties in both mouse and human systems.

172   To investigate the necessity of DKK2 in the formation of LYZ+ cancer cells during colon

cancer metastasis, we employed the murine *in vivo* model of liver metastasis shown in Figure 1.

In order to analyze transcriptome changes by *Dkk2* knockout in each cell type of metastasized

tumors, we performed single-cell RNA sequencing (scRNA-seq) analysis on liver metastasized

AKP and KO tumor cells. The Louvain method revealed 4 clusters in our dataset. A cell type was

assigned to each cluster on the basis of gene expression patterns and visualized in a Uniform

manifold approximation and projection (UMAP) representation (Fig. 3A). Epithelial tumor cells

were sub-clustered as 7 clusters and the cluster 6 has been identified as cancer cells with Paneth

cell properties based on the expression of Paneth marker genes using the Panglao database

(Fig. 3A, S2A-D). Total of 17 cells in cluster 6 were further defined as cancer cells with Paneth

cell properties (Paneth+) or goblet cell properties (Goblet+) using the relevant marker genes

expression, *Lyz1*, *Muc2* and *Atoh1* (Fig. S2E and S2F). The calculated ratio between Paneth+

cells and Goblet+ cells in KO metastasized tumors was 0.43, reduced as half compared to the

ratio in the AKP control, 1 (Fig. S2F). This suggests a reduction of LYZ+ cells in metastasized

colon tumors by knockout a *Dkk2* gene, which was confirmed by flow cytometry analysis of LYZ+

cells in metastasized mouse colon cancers in the liver (Fig. 6H and 6I).

188   To analyze the consequence of the loss of LYZ+ cells by *Dkk2* knockout in cancer stem

cell niche formation, we enriched *Lgr5* positive cells that consist of stem cells, stem cell niche

cells and transit amplifying cells in the homeostatic condition (Fig. S3A and S3B) (8). Elevated

expression of stem cell marker genes represents cancer stemness in *Lgr5* positive cells

compared to *Lgr5* negative cells (Fig. S3C). The UMAP plot of *Lgr5* positive cells displayed 2

clusters (Fig. S3D). Interestingly, expression of Noggin (*Nog*), one of the stem cell niche factors

was detected mostly in cluster 1 where only 1 cell was counted in KO (Fig. S3E-S3G). Noggin is

an inhibitor of BMP signaling, which restricts stemness of *Lgr5* positive stem cells in the gut (33,

34). Following the reduction of *Nog*, *Bmp4* expression was reversed in KO tumor epithelial cells in

the scRNA-seq data (Fig. S3H). This has been confirmed by quantitative gene expression

analysis of *Bmp4* in KO organoids (Fig. S3I). The correlative expression between *Dkk2* and the

stem cell marker gene, *Lgr5* has been also confirmed in the scRNA-seq data (Fig. 3B).

Furthermore, pathways enrichment analysis of the scRNA-seq data using IPA suggested that the

glycolysis I pathway was significantly inhibited in KO *Lgr5* positive tumor epithelial cells (Fig. 3C).

Expression of glycolysis-related genes such as *Aldoa*, *Aldoc* and *Fbp2* was reduced by *Dkk2*

knockout in *Lgr5* positive cells (Fig. 3D) (35). Previous studies have defined that Paneth cells

participate in the regulation of energy metabolism in intestinal stem cells (12, 36). Since stem

cells are not able to generate lactate, Paneth cells instead process glucose in order to provide

lactate to stem cells (12). Therefore, reduction of the glycolysis I pathway in the absence of DKK2

indicates the impaired stem cell niche formation by the loss of LYZ+ cells carrying Paneth cell

properties.

Paneth cells constitute the stem cell niche in the small intestine (21). Recent studies

have identified the presence of Paneth-like cells in the colon (25, 26). However, the presence of

Paneth cells or Paneth-like cells within the tumor microenvironment is elusive. To discern the

existence of cancer cells exhibiting Paneth cell properties in humans, we conducted an analysis

of scRNA-seq data obtained from colorectal cancer patients (37). Our analysis focused on CD45-

EpCAM+ epithelial cells, excluding hematopoietic cells. UMAP plotted 31 clusters of single cells

derived from normal tissue, primary tumor and metastasis samples (Fig. 4A and B). Employing

lysozyme expression and Paneth cell module scores (Pangalo database) as criteria, we have

identified LYZ+ cancer cells harboring Paneth cell properties across 5 clusters (Fig. 4C-H, Fig. S4

and S5). LYZ+ cancer cell population is distinct in metastasis samples (Fig. 4E and 4F). LYZ

expression over 2.0 (3[rd] Quantile, round 1) and Paneth module scores over 0.2 (3[rd] Quantile,

round 2 to avoid getting 0) single cells were identified as LYZ+ cancer cells harboring Paneth cell

properties, 1%, 13% and 24% in normal, primary tumor and metastasis samples, respectively.

This delineates the distribution of LYZ+ cancer cells with Paneth cell characteristics in different

stages of colorectal cancer progression.

To unravel the functional significance of LYZ+ cancer cells, we performed gene set

enrichment analysis for the scRNA-seq data obtained from colon cancer patients. In correlation

with our murine scRNA-seq data IPA results, glycolysis emerged as the 2nd-ranked enriched

pathway in LYZ+ cancer cells compared to LYZ- cancer cells (Fig 5A). Notably, key players in

glycolysis such as *ALDOB* and *KDELR3* genes are increased in LYZ+ cancer cells (Fig. 5B).

Also, the hallmark of protein secretion was the top-ranked enriched pathway, which is essential

for Paneth cell activity including secretion of anti-microbial peptides (Fig. 5A). These results

suggest the existence of a distinct population of LYZ+ colon cancer cells endowed with Paneth

cell properties. Collectively, DKK2 is required for the generation of LYZ+ cells to form the cancer

stem cell niche, a prerequisite for the outgrowth of metastasized colorectal cancer cells.

HNF4A mediates the formation of Lysozyme positive colon cancer cells by DKK2.

Ingenuity pathway analysis (IPA) of the human scRNA-seq data from primary tumor and

metastasis samples suggested hepatocyte nuclear factor 4 alpha (HNF4A) as an upstream

regulator of LYZ+ cancer cells compared to LYZ- cancer cells (Fig. 5C). In line with this, IPA of

the bulk RNA-seq data using mouse colon cancer organoids suggested that HNF4A is activated

in KO organoids (Fig. 6A). This is consistent with our previous report that DKK2 enhanced *Lgr5*

expression in colitis-induced cancer cells via HNF4α1, an isoform of HNF4A (28). Significant

enrichment of the HNF4A binding motif was also detected in the open chromatin regions of KO

organoids compared to AKP controls determined by assay for transposase-associated chromatin

using sequencing (ATAC-seq) (Fig. 6B). In comparison with normal colonic organoids, the

HNF4A motif was listed as the most reduced motif in AKP indicating significant reduction of

HNF4A-regulated transcription whereas it might be rescued by *Dkk2* knockout (Fig. 6C). Notably,

HNF4A was the only molecule suggested by all of these multi-omics data analyses in both mouse

and human. These results suggest HNF4A as a key transcription factor in downstream signaling

of DKK2 in colon cancer cells.

HNF4A is necessary for maturation of fetal intestine and barrier functions of intestinal

epithelium in the homeostatic condition (38, 39). Two *Hnf4a* isoforms—*Hnf4α1* and *Hnf4α7*— are

involved in intestinal homeostasis and regeneration (40). Intestinal inflammation induces

expression of the HNF4α1 isoform in the crypt epithelial cells to regenerate the epithelium

whereas loss of the HNF4α1 isoform occurs in colitis-induced tumor cells (40). Loss of the

HNF4α1 isoform was also observed in AKP organoids while *Dkk2* knockout restored the

presence of HNF4α1 protein in the nucleus of AKP organoid cells (28). This allowed us to

investigate the binding regions of HNF4α1 in colon cancer cells using KO organoids. Chromatin

immunoprecipitation followed by sequencing (Chip-seq) analysis using the Chip-seq grade anti-

HNF4A antibody revealed that HNF4A binds to the promoter region of *Sox9* in KO organoid cells

(Fig. 6D). Sox9 is a transcription factor required for the secretory lineage of epithelial cell

differentiation including Paneth cells (23). Quantitative gene expression analysis showed that

Dkk2 knockout reduced *Sox9* expression in AKP organoids, which was rescued by recombinant

DKK2 protein treatment (Fig. 6E). Reduced *Sox9* expression in KO organoids was also reversed

by HNF4α1 knockdown (Fig. 6F). These data indicate that DKK2-driven loss of HNF4α1 protein

leads to *Sox9* expression followed by formation of LYZ+ colon cancer cells. Confocal microscopy

analysis confirmed that HNF4α1 knockdown rescued the presence of LYZ+ cancer cells in KO

organoids (Fig. 6G). In order to quantify LYZ+ cancer cells in the mouse model of liver

metastasis, tdTomato positive cells in the liver were identified as the implanted organoid-derived

cancer cells then, LYZ+ cells were calculated (Fig. 6H). The percentile of LYZ+ cells in total

cancer cells was decreased about 3-fold in *Dkk2* knockout metastasized cells (KO) compared to

the AKP control cells in liver (Fig. 6I). Simultaneously, HNF4α1 positive cells were tripled to about

60 percent in KO (Fig. 6H and 6I). These findings suggest that DKK2 is indispensable for the

generation of LYZ+ cancer cells in liver metastasized nodules by reducing protein levels of

HNF4α1. Further analyses for human colorectal cancer scRNA-seq data revealed that HNF4A

regulation activity is reduced in LYZ+ human colon cancer cells compared to the LYZ- cancer

cells while SOX9 regulation activity is significantly increased (Fig. 6J-6K and S5) suggesting the

key roles of HNF4A and SOX9 in LYZ+ cancer cell formation in human. Taken together, DKK2-

driven loss of HNF4α1 protein enhances Sox9 expression in colon cancer cells to generate LYZ+

cells with Paneth cell properties.

Discussion

282   We have shown that DKK2 is indispensable for metastatic tumor growth of colorectal

283 cancer in the murine model developed by transplantation of colon cancer organoids carrying

284 mutation in the *Apc*, *Kras* and *Tp53* genes. Bulk RNA-seq analysis of DKK2-deficient colon

285 cancer organoids showed reduction of the marker genes of Paneth cells, which play a role in the

286 stem cell niche development. Single cell RNA-seq analyses for mouse and human metastatic

287 colon cancers have identified the existence of LYZ+ cancer cells harboring Paneth cell properties,

288 in particular, glycolysis for stem cells. In the absence of DKK2, both pathway enrichment analysis

289 of the RNA-seq and ATAC-seq data suggested that HNF4A is activated. This is consistent with

290 our previous report in the murine model of colitis-induced tumorigenesis (28). ChIP-seq analysis

291 using an anti-HNF4A antibody revealed that HNF4A directly binds to the promoter region of *Sox9*,

292 a transcriptional factor required for epithelial differentiation into Paneth cells (22, 23). Notably,

293 DKK2 induces the loss of HNF4α1 isoform in colon cancer cells followed by elevation of *Sox9*

294 expression lead to the formation of LYZ+ cancer cells exhibiting Paneth cell properties. DKK2-

295 deficient colon cancer cells possessed high levels of HNF4α1 protein and failed to generate LYZ+

296 cancer cells *in vitro* and *in vivo*. DKK2 deficiency resulted in reduced glycolysis in mouse liver

297 metastasized colon cancer cells. These results demonstrate the significant reduction of liver

298 metastasis in DKK2 knockout (KO) colon cancer cells infused mice.

299   *Fumagalli et al.* have recently shown that the generation of *Lgr5* positive cells from the

300 metastasized *Lgr5* negative cancer cells is necessary for their outgrowth using *in vivo* time-lapse

301 imaging and organoid culture of metastatic cancer cells that escaped from the primary colon

302 cancer tissues (7). This finding implies the necessity of both cancer stem cells and their niche for

303 metastatic outgrowth. Indeed, Paneth cell-derived stem cell niche factors such as Wnt3, EGF and

304 Dll4 are essential for the maintenance of stem cells (11). Dietary induced mouse sporadic

305 intestinal cancer showed elevation of Paneth cell marker genes in the intestinal epithelium (41).

306 IDO1+ Paneth cells contribute to the immune evasion of colon cancer (42). Depletion of Paneth

307 cells or Paneth cell-derived Wnt3 in *Apc^{Min}* mice impaired intestinal adenoma formation (43). The

plasticity of Paneth cells allows dedifferentiation of Paneth cells into intestinal stem cells (15, 44).

These reports indicate the necessity of the Paneth cells during the development of colon cancer.

However, the existence of Paneth cells in large intestine or tumor tissues remains unclear. In

2016, Reg4+ secretory cells have been suggested as Paneth-like cells in the colon (24). Single

cell transcriptome analysis showed the presence of Paneth-like cells in human colon (25). Recent

study further supported the presence of Paneth-like cells within the colonic stem cell niche during

bifidobacterium-induced intestinal stem cell regeneration (26). Our study suggests the presence

of LYZ+ cells exhibiting Paneth cell properties in the context of colon cancer and metastasis. This

extends our knowledge of colorectal cancer progression that DKK2 is required for the generation

of LYZ+ cells forming cancer stem cell niches.

Further investigation will be needed to characterize LYZ+ cancer cells with Paneth cell

properties observed in colon cancers carrying mutations in *Apc*, *Kras* and *Tp53* genes compared

to normal Paneth cells in order to identify their roles in cancer stem cell niche development.

Recent studies have reported stem cell niche remodeling in the primary tumor in the presence of

oncogenic mutations. Cancer stem cells outcompete their neighboring normal stem cells during

intestinal tumorigenesis, which is known as clonal fixation (45). *Apc*-mutant cancer cells secrete

several Wnt antagonists such as NOTUM to inhibit proliferation of normal intestinal stem cells and

facilitate their differentiation (46). Other oncogenic mutations in *Kras* or *Pik3ca* genes lead to

paracrine secretion of BMP ligands, remodeling the stem cell niche that is detrimental for

maintaining normal stem cells (47). These changes are beneficial for the outgrowth of cancer

stem cells in the crypt where the stem cell niche had been developed prior to tumorigenesis. In

the context of metastasis, the formation of stem cell niches is required upon metastases seeding

in the liver. We have shown that DKK2 is required for the formation of LYZ+ cancer cells carrying

Paneth cell properties. DKK2 expression is induced by *Apc* knockout and further increased by

additional oncogenic mutations in *Kras* and *Tp53* genes (28). Characterization of DKK2-induced

LYZ+ cancer cells will provide better understanding of the biology of cancer stem cell niches,

particularly in liver metastases of colorectal cancer.

DKK2 expression in colon cancer promotes LYZ+ cancer cell generation through the

regulation of HNF4α1, which suppresses expression of *Sox9,* the transcription factor for Paneth

cell differentiation. It has been recently reported that HNF4α2 isoform acts as an upstream

regulator of Wnt3 and Paneth cell differentiation implying the roles of HNF4A for Paneth cell

differentiation in the homeostatic condition (48). Using *Villin^{Cre}-Apc^{floxed}* mice, Suzuki et al. have

confirmed that HNF4A proteins are completely absent in cancer cells with high β-catenin activity

along with increased *Sox9* expression (49). *Sox9* regulates cell proliferation and is required for

Paneth cell differentiation (22, 23). Bi-allelic inactivation of the *Apc* gene induces *Sox9* expression

in colon cancer (50). Likewise, *Apc* inactivation induces *Dkk2* expression (28). Here we showed

that DKK2-mediated HNF4α1 protein degradation enhanced *Sox9* expression in colon cancer

(Fig. 7). Notably, *Apc* mutation-driven *Sox9* expression blocks intestinal differentiation and

activates stem cell-like program (50). This is consistent with our previous report that DKK2

enhances *Lgr5* expression in colon cancer (28). In sum, our findings suggest that a Wng ligand,

DKK2 is an upstream regulator of SOX9 expression for enhanced stem cell activity via the

formation of LYZ+ cells with Paneth cell characteristics in colorectal cancers.

In this study, we have shown that DKK2 promotes liver metastasis of colon cancer. DKK2

is required for the formation of LYZ+ colon cancer cells exhibiting Paneth cell properties such as

glycolysis for stem cells. Glycolysis in *Lgr5* expressing cancer cells was reduced along with the

loss of LYZ+ cells by *Dkk2* knockout. As a result, metastatic growth of colon cancer cells was

significantly inhibited *in vivo*. These findings suggest the necessity of DKK2 in the development of

cancer stem cell niches for developing metastases. Mechanistically, DKK2-mediated degradation

of HNF4α1 protein elevated *Sox9* expression followed by LYZ+ cell formation. Given that DKK2

expression is intestinal tumor specific, inhibition of DKK2 may have promising therapeutic

outcomes in the treatment of metastatic colon cancer.

Acknowledgements

The authors would like to thank Gouzel Tokmulina for the FACS sorting.

Author contributions

J.H.S. and A.L.M.B. designed research; J.H.S., J.L., J.J., R.K.D, S.E.M., and J.Y.H. performed

research; J.H.S., J.L., J.J., J.P., R.K.D., S.E.M., J.Y.H., and A.L.M.B. contributed new

reagents/analytic tools; J.H.S., J.J., J.L., J.P., J.C., J.W. and A.L.M.B. analyzed data; and J.H.S.,

J.J., J.L., J.C. and A.L.M.B. wrote the paper.

This work was supported by NIH RO1 CA168670-01 (awarded to A.L.M.B) and Yonsei University

Future-Leading Research Initiative 2023-22-0438 (awarded to J.H.S.).

Conflict of interest

The authors declare no competing interests.

References

1.   Dillekås H, Rogers MS, Straume O. Are 90% of deaths from cancer caused by metastases? Cancer Med. 2019;8(12):5574-6.

2.   Rowan AJ, Lamlum H, Ilyas M, Wheeler J, Straub J, Papadopoulou A, et al. APC mutations in sporadic colorectal tumors: A mutational "hotspot" and interdependence of the "two hits". Proc Natl Acad Sci U S A. 2000;97(7):3352-7.

3.   van Houdt WJ, Hoogwater FJ, de Bruijn MT, Emmink BL, Nijkamp MW, Raats DA, et al. Oncogenic KRAS Desensitizes Colorectal Tumor Cells to Epidermal Growth Factor Receptor Inhibition and Activation. Neoplasia. 2010;12(6):443-IN2.

4.   Drost J, van Jaarsveld RH, Ponsioen B, Zimberlin C, van Boxtel R, Buijs A, et al. Sequential cancer mutations in cultured human intestinal stem cells. Nature. 2015;521(7550):43-7.

5.   Fumagalli A, Drost J, Suijkervuijk SJ, van Boxtel R, de Ligt J, Offerhaus GJ, et al. Genetic dissection of colorectal cancer progression by orthotopic transplantation of engineered cancer organoids. Proc Natl Acad Sci U S A. 2017:E2357-E64.

6.   de Sousa e Melo F, Kurtova AV, Harnoss JM, Kljavin N, Hoeck JD, Hung J, et al. A distinct role for Lgr5(+) stem cells in primary and metastatic colon cancer. Nature. 2017;543(7647):676-80.

7.   Fumagalli A, Oost KC, Kester L, Morgner J, Bornes L, Bruens L, et al. Plasticity of Lgr5-Negative Cancer Cells Drives Metastasis in Colorectal Cancer. Cell Stem Cell. 2020;26(4):569-78.e7.

8.   Sato T, Vries RG, Snippert HJ, van de Wetering M, Barker N, Stange DE, et al. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. Nature. 2009;459(7244):262-5.

9.   McCarthy N, Kraiczy J, Shivdasani RA. Cellular and molecular architecture of the intestinal stem cell niche. Nature Cell Biology. 2020;22(9):1033-41.

10.   Santos AJM, Lo Y-H, Mah AT, Kuo CJ. The Intestinal Stem Cell Niche:
Homeostasis and Adaptations. Trends in Cell Biology. 2018;28(12):1062-78.
11.   Sato T, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M, et al.
Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. Nature.
2011;469(7330):415-8.
12.   Rodriguez-Colman MJ, Schewe M, Meerlo M, Stigter E, Gerrits J, Pras-Raves M,
et al. Interplay between metabolic identities in the intestinal crypt supports stem
cell function. Nature. 2017;543(7645):424-7.
13.   Tetteh Paul W, Basak O, Farin Henner F, Wiebrands K, Kretzschmar K,
Begthel H, et al. Replacement of Lost Lgr5-Positive Stem Cells through Plasticity of
Their Enterocyte-Lineage Daughters. Cell Stem Cell. 2016;18(2):203-13.
14.   van Es JH, Sato T, van de Wetering M, Lyubimova A, Yee Nee AN, Gregorieff A,
et al. Dll1+ secretory progenitor cells revert to stem cells upon crypt damage. Nat
Cell Biol. 2012;14(10):1099-104.
15.   Buczacki SJ, Zecchini HI, Nicholson AM, Russell R, Vermeulen L, Kemp R, et al.
Intestinal label-retaining cells are secretory precursors expressing Lgr5. Nature.
2013;495(7439):65-9.
16.   Batlle E, Clevers H. Cancer stem cells revisited. Nature Medicine.
2017;23(10):1124-34.
17.   Hilkens J, Timmer NC, Boer M, Ikink GJ, Schewe M, Sacchetti A, et al. RSPO3
expands intestinal stem cell and niche compartments and drives tumorigenesis. Gut.
2017;66(6):1095-105.
18.   van Es JH, Jay P, Gregorieff A, van Gijn ME, Jonkheer S, Hatzis P, et al. Wnt
signalling induces maturation of Paneth cells in intestinal crypts. Nature Cell
Biology. 2005;7(4):381-6.
19.   Suzuki K, Fukui H, Kayahara T, Sawada M, Seno H, Hiai H, et al. Hes1-deficient
mice show precocious differentiation of Paneth cells in the small intestine.
Biochemical and Biophysical Research Communications. 2005;328(1):348-52.
20.   Andreu P, Peignon G, Slomianny C, Taketo MM, Colnot S, Robine S, et al. A
genetic study of the role of the Wnt/β-catenin signalling in Paneth cell
differentiation. Developmental Biology. 2008;324(2):288-96.
21.   Lueschow SR, McElroy SJ. The Paneth Cell: The Curator and Defender of the
Immature Small Intestine. Frontiers in Immunology. 2020;11.
22.   Bastide P, Darido C, Pannequin J, Kist R, Robine S, Marty-Double C, et al. Sox9
regulates cell proliferation and is required for Paneth cell differentiation in the
intestinal epithelium. J Cell Biol. 2007;178(4):635-48.
23.   Mori–Akiyama Y, van den Born M, van Es JH, Hamilton SR, Adams HP, Zhang
J, et al. SOX9 Is Required for the Differentiation of Paneth Cells in the Intestinal
Epithelium. Gastroenterology. 2007;133(2):539-46.
24.   Sasaki N, Sachs N, Wiebrands K, Ellenbroek SI, Fumagalli A, Lyubimova A, et
al. Reg4+ deep crypt secretory cells function as epithelial niche for Lgr5+ stem cells
in colon. Proc Natl Acad Sci U S A. 2016;113(37):E5399-407.
25.   Wang Y, Song W, Wang J, Wang T, Xiong X, Qi Z, et al. Single-cell
transcriptome analysis reveals differential nutrient absorption functions in human
intestine. J Exp Med. 2020;217(2).

26.   Qi Y, He J, Zhang Y, Ge Q, Wang Q, Chen L, et al. Heat-inactivated
Bifidobacterium adolescentis ameliorates colon senescence through Paneth-like-
cell-mediated stem cell activation. Nat Commun. 2023;14(1):6121.
27.   Dayton TL, Clevers H. Beyond growth signaling: Paneth cells metabolically
support ISCs. Cell Res. 2017;27(7):851-2.
28.   Shin JH, Jeong J, Choi J, Lim J, Dinesh RK, Braverman J, et al. Dickkopf-2
regulates the stem cell marker LGR5 in colorectal cancer via HNF4α1. iScience.
2021;24(5):102411-.
29.   Shin JH, Jeong J, Maher SE, Lee H-W, Lim J, Bothwell ALM. Colon cancer cells
acquire immune regulatory molecules from tumor-infiltrating lymphocytes by
trogocytosis. Proceedings of the National Academy of Sciences. 2021;118(48).
30.   Li XB, Yang G, Zhu L, Tang YL, Zhang C, Ju Z, et al. Gastric Lgr5(+) stem cells
are the cellular origin of invasive intestinal-type gastric cancer in mice. Cell Res.
2016;26(7):838-49.
31.   Chen J, Elfiky A, Han M, Chen C, Saif MW. The role of Src in colon cancer and
its therapeutic implications. Clin Colorectal Cancer. 2014;13(1):5-13.
32.   Vidrich A, Buzan JM, Brodrick B, Ilo C, Bradley L, Fendig KS, et al. Fibroblast
growth factor receptor-3 regulates Paneth cell lineage allocation and accrual of
epithelial stem cells during murine intestinal development. American Journal of
Physiology-Gastrointestinal and Liver Physiology. 2009;297(1):G168-G78.
33.   Haramis A-PG, Begthel H, Born Mvd, Es Jv, Jonkheer S, Offerhaus GJA, et al. De
Novo Crypt Formation and Juvenile Polyposis on BMP Inhibition in Mouse Intestine.
Science. 2004;303(5664):1684-6.
34.   Qi Z, Li Y, Zhao B, Xu C, Liu Y, Li H, et al. BMP restricts stemness of intestinal
Lgr5+ stem cells by directly suppressing their signature genes. Nature
Communications. 2017;8(1):13824.
35.   Cascone T, McKenzie JA, Mbofung RM, Punt S, Wang Z, Xu C, et al. Increased
Tumor Glycolysis Characterizes Immune Resistance to Adoptive T Cell Therapy. Cell
Metabolism. 2018;27(5):977-87.e4.
36.   Yilmaz ÖH, Katajisto P, Lamming DW, Gültekin Y, Bauer-Rowe KE, Sengupta
S, et al. mTORC1 in the Paneth cell niche couples intestinal stem-cell function to
calorie intake. Nature. 2012;486(7404):490-5.
37.   Joanito I, Wirapati P, Zhao N, Nawaz Z, Yeo G, Lee F, et al. Single-cell and bulk
transcriptome sequencing identifies two epithelial tumor cell states and refines the
consensus molecular classification of colorectal cancer. Nat Genet. 2022;54(7):963-
75.
38.   Chen L, Toke NH, Luo S, Vasoya RP, Aita R, Parthasarathy A, et al. HNF4
factors control chromatin accessibility and are redundantly required for maturation
of the fetal intestine. Development. 2019;146(19):dev179432.
39.   Cattin AL, Le Beyec J, Barreau F, Saint-Just S, Houllier A, Gonzalez FJ, et al.
Hepatocyte nuclear factor 4alpha, a key factor for homeostasis, cell architecture, and
barrier function of the adult intestinal epithelium. Mol Cell Biol. 2009;29(23):6294-
308.
40.   Chellappa K, Deol P, Evans JR, Vuong LM, Chen G, Briancon N, et al. Opposing
roles of nuclear receptor HNF4alpha isoforms in colitis and colitis-associated colon
cancer. Elife. 2016;5.

41.   Wang D, Peregrina K, Dhima E, Lin EY, Mariadason JM, Augenlicht LH. Paneth
cell marker expression in intestinal villi and colon crypts characterizes dietary
induced risk for mouse sporadic intestinal cancer. Proceedings of the National
Academy of Sciences. 2011;108(25):10272-7.
42.   Pflügler S, Svinka J, Scharf I, Crncec I, Filipits M, Charoentong P, et al. IDO1+
Paneth cells promote immune escape of colorectal cancer. Communications Biology.
2020;3(1).
43.   Chen Q, Suzuki K, Sifuentes-Dominguez L, Miyata N, Song J, Lopez A, et al.
Paneth cell–derived growth factors support tumorigenesis in the small intestine.
Life Science Alliance. 2021;4.
44.   Clevers H. The intestinal crypt, a prototype stem cell compartment. Cell.
2013;154(2):274-84.
45.   Vermeulen L, Morrissey E, Heijden Mvd, Nicholson AM, Sottoriva A, Buczacki
S, et al. Defining Stem Cell Dynamics in Models of Intestinal Tumor Initiation.
Science. 2013;342(6161):995-8.
46.   Flanagan DJ, Pentinmikko N, Luopajärvi K, Willis NJ, Gilroy K, Raven AP, et al.
NOTUM from Apc-mutant cells biases clonal competition to initiate cancer. Nature.
2021;594(7863):430-5.
47.   Yum MK, Han S, Fink J, Wu S-HS, Dabrowska C, Trendafilova T, et al. Tracing
oncogene-driven remodelling of the intestinal stem cell niche. Nature.
2021;594(7863):442-7.
48.   Jones C, Avino M, Giroux V, Boudreau F. HNF4alpha Acts as Upstream
Functional Regulator of Intestinal Wnt3 and Paneth Cell Fate. Cell Mol Gastroenterol
Hepatol. 2023;15(3):593-612.
49.   Suzuki T, Kilbey A, Casa-Rodriguez N, Lawlor A, Georgakopoulou A, Hayman
H, et al. beta-Catenin Drives Butyrophilin-like Molecule Loss and gammadelta T-cell
Exclusion in Colon Cancer. Cancer Immunol Res. 2023;11(8):1137-55.
50.   Feng Y, Sentani K, Wiese A, Sands E, Green M, Bommer GT, et al. Sox9
Induction, Ectopic Paneth Cells, and Mitotic Spindle Axis Defects in Mouse Colon
Adenomatous Epithelium Arising From Conditional Biallelic Apc
Inactivation. The American Journal of Pathology. 2013;183(2):493-503.
522
523

Figure Legends

Figure 1. DKK2 is required for Lgr5 positive stem cells-driven liver metastasis of colorectal cancer. (A) Isolated cells from AKP tumor organoids expressing the *tdTomato* reporter gene were dissociated and injected into the spleen of wild type C57BL/6 mice. 8 weeks after the injection, metastatic tumor growth was measured by in vivo imaging analysis. **(B)** Representative pictures of *in vivo* imaging analysis. Ctrl: AKP control organoids transduced with scrambled guide RNA, KO: *Dkk2* knockout AKP organoids. **(C-D)** Statistic analysis of liver metastasis (C) and survival (D). Ctrl (*n*=10), KO (*n*=15), *n* represents the number of mice. **(E)** Quantitative gene expression analysis of *Dkk2, Lgr5* and *Hnf4α1* in liver metastasized colon cancer cells. Ctrl (*n*=11), KO (*n*=12), *n* represents the number of isolated cancer nodules. **(F)** Representative flow cytometry analysis data of c-Src phosphorylation (pSrc) and *Lgr5* expression in metastasized colon cancer cells. **(G)** Statistic analysis of *Lgr5* expression and pSrc in (E). The average of mean fluorescence intensity in control samples was set as 1 and relative fluorescence intensity was calculated. **(H)** Statistic analysis of the percentile of Lgr5 high (Lgr5 and pSrc double positive) cells in metastasized colon cancer in (F). Each *symbol* represents an individually isolated cancer nodule. n.s = not significant, **P<0.01, ****P<0.0001; two-tailed Welch's t-test (C, E, G, H). Error bars indicate mean ± s.d. Log-rank test (D). Results are representative of three independent experiments.

Figure 2. DKK2 is indispensable for the generation of cancer cells with Paneth cell properties in colon cancer organoids. (A) A schematic diagram of the generation of KO organoids using CRISPR technique and the reconstitution of DKK2 in organoids by recombinant mouse DKK2 protein (rmDKK2) treatment. Lysozyme (LYZ) expression is highlighted for the followings. **(B)** A volcano plot of RNA-seq analysis comparing KO versus AKP organoids. Paneth cell marker genes are highlighted as blue circles (AKP=3 and KO=5 biological replicates were analyzed). **(C)** Confocal microscopy analysis of LYZ positive cells in AKP or KO organoids in a time-dependent manner using anti-Lysozyme (LYZ) antibody. **(D)** Quantitative real time PCR analyses of *Lyz1* and *Lyz2* in 8 days cultured colon cancer organoids. KO organoids were

cultured in the presence of 1 μg/ml of recombinant mouse DKK2 protein. *P<0.05, **P<0.01,

***P<0.001; two-tailed Welch's t-test. Error bars indicate mean ± s.d. Results are representative

of three biological replicates.

Figure 3. Cancer cells harboring Paneth cell properties were reduced in *Dkk2* knockout

metastasized colon cancer tissues in mice.

Control AKP or KO colon cancer organoids were transplanted via splenic vein as described in

Figure 1. 3 weeks after transplantation, mice were sacrificed to analyze metastatic tumor growth

in liver. **(A)** Single cell RNA sequencing analysis (scRNA-seq) of liver metastasized colon cancer

tissues. The uniform manifold approximation and projection (UMAP) plot clustered epithelial,

endothelial, liver and immune cells in metastasized cancers based on transcriptome analysis. The

cancer epithelial cell cluster was sub-clustered to identify cells with Paneth cell properties (cluster

6, Paneth positive). **(B)** The correlation between *Dkk2* and *Lgr5* expression in the cluster of

epithelial cells by Pearson r test. **(C)** Ingenuity pathway analysis (IPA)-suggested top 5 canonical

pathways of the scRNA-seq data of *Lgr5* positive epithelial cells in KO compared to AKP. z-

Scores indicate activation or inhibition of the suggested pathways. The significance values for the

pathways are calculated by the right-tailed Fisher's exact test. **(D)** Box plots show expressions of

the genes involved in the glycolysis I pathway in (C). ns = not significant, *<P<0.05, **P<0.01,

***P<0.001, ****P<0.0001; Wilcoxon signed-rank test.

Figure 4. Identification of colon cancer cells harboring Paneth cell properties in humans.

Published colorectal cancer patient scRNA-seq data (REF) was analyzed to identify the presence

of cancer cells with Paneth cell properties. **(A)** The UMAP plot of total 31 clusters. **(B)** Normal

cells, primary tumor cells and liver metastasized cells (Metastasis) are shown in the UMAP plot

clusters. **(C-D)** Expression levels of lysozyme (LYZ) and Paneth cell module scores are displayed

in the UMPA plot. **(G)** Based on the analysis in (C) and (D), cancer cells harboring Paneth cell

properties are indicated as red dots (LYZ+ cancer cells). **(E)** Lysozyme expression and Paneth

cell module scores of single cells in Normal, Primary Tumor and Metastasis samples are

presented by dot plots. **(F)** Dot plots in (E) are overlayed. **(H)** The percentiles of cancer cells with

Paneth cell properties (LYZ+ cancer cells) are shown.

Figure 5. Colon cancer cells with Paneth cell properties contribute to glycolysis.

Colorectal cancer patient scRNA-seq data were further analyzed by Gene set enrichment

analysis (GSEA) and Ingenuity Pathway Analysis (IPA) **(A)** GSEA of cancer cells with Paneth cell

properties (LYZ+ cancer cells) compared to all other cancer cells (LYZ- cancer cells), shown in

Fig.4 (G). **(B)** Representative gene expressions in the hallmark of glycolysis pathway are shown.

(C) Upstream regulators suggested by IPA in Primary Tumor and Metastasis are presented.

Activation z-scores are indicated in the bar. IPA predicted activation or inhibition of the upstream

regulators are colored as red and cyan, respectively.

Figure 6. DKK2-driven reduction of HNF4α1 protein in murine colon cancer organoids

promotes cancer cells with Paneth cell properties.

(A) Ingenuity pathway analysis (IPA)-suggested upstream regulators of the RNA-seq data shown

in Figure 2A. Z-scores are presented in the bar. IPA predicted activation or inhibition of the

upstream regulators are colored as red and cyan, respectively. **(B-C)** List of top 5 transcriptional

factors (TF) in the motif enrichment analysis of ATAC-seq data comparing normal colonic

organoids, AKP and KO cancer organoids. * indicates possible false positive. **(D)** Chromatin

immunoprecipitation sequencing (Chip-seq) analysis of KO organoids using an anti-HNF4A

antibody. **(E)** Quantitative expression of *Sox9* in colon cancer organoids in the presence or

absence of DKK2 (rmDKK2: recombinant mouse DKK2 protein added in organoid culture). **(F)**

Analysis of Sox9 expression after knockdown *HNF4α1* in KO colon cancer organoids (KO

HNF4α1 KD). ns = not significant, *P<0.05, **P<0.01, ***P<0.001; two-tailed Welch's t-test. Error

bars indicate mean ± s.d. **(G)** Representative images of confocal microscopy analysis of Lyz-

stained cancer cells with Paneth cell properties in DKK2 KO HNF4α1 KD organoids. **(H-I)** Control

AKP or KO colon cancer organoids were transplanted via splenic vein as described in Figure 1. 3

weeks after transplantation, mice were sacrificed to analyze metastatic tumor growth in liver.

Quantification of cancer cells with Paneth cell properties in metastasized tumor nodules by flow

cytometry for Lysozyme (LYZ) and HNF4α1. Tumor cells were initially gated by the tdTomato

reporter expression. Representative images of flow cytometry are shown (H). Statistic analyses of

the percentiles of LYZ positive cells (% of upper left) and HNF4α1 positive cells (% of lower right)

in tumor nodules (I). **$P<0.01$, ****$P<0.0001$; two-tailed Welch's t-test. Error bars indicate mean ±

s.d. Data are representative of two independent experiments. **(J- K)** Reduced HNF4A regulation

activity with enhanced SOX9 regulation activity in LYZ+ cancer cells in human colorectal cancer

scRNA-seq data. Box plots represent the regulon activity of HNF4A and SOX9 in LYZ+ cancer

cells. ****$P<0.0001$; Wilcoxon signed-rank test (J). Z-scaled regulon activities of HNF4A and

SOX9 in human colon cancer cells is displayed by heatmap (K).

Figure 7. The suggested mechanism of DKK2 in the formation of colon cancer cells with

Paneth cell properties.

In the absence of DKK2, Sox9 expression is inhibited by HNF4A. Our previous report has shown

that DKK2 activates Src followed by degradation of HNF4A protein (28). HNF4A deficiency leads

to Sox9 expression in colon cancer cells that induces Paneth cell properties including the

expression of lysozymes and defensins. Formation of cancer cells with Paneth cell properties by

DKK2 contributes the outgrowth of metastasized colon cancer cells in the liver.

A

sgRNA targeting a Dkk2 gene

Cas9

AKP Cancer organoid
(*Apc, Kras, Tp53* mutations)

1) Knockdown *Dkk2* using LentiviralCRISPRv2
2) Selection with Puromycin
3) Sorting for Dkk2 knockout single stem cells
4) Generation of organoids

LYZ+ cells

LYZ+ ↓

DKK2 KO organoid
(*Apc, Kras, Tp53* mutations
+ *Dkk2* gene knockout)

rmDKK2

LYZ+ ↑

**Reconstitution of Lyz expression in
DKK2 KO organoid by rmDKK2 treatment**

B

-log₁₀(adjusted P-value) vs Log2 fold change (KO/AKP)

Defa24, Lyz1, Defa21, Lyz2, Defa36, Defa30, Defa3

n = 12,813

C

	LYZ	tdTomato	DAPI	Merge
AKP Day 2				
KO Day 2				
AKP Day 4				
KO Day 4				
AKP Day 8				
KO Day 8				

D

Lyz1

Relative Expression to *Hprt*

AKP, KO, rmDKK2

*** * **

Lyz2

Relative Expression to *Hprt*

AKP, KO, rmDKK2

*** ** ***

A

B
Normal | Primary Tumor | Metastasis

C
LYZ

D
Paneth cell Module Score

G
● LYZ+ cancer cells
● LYZ− cancer cells

E
Normal | Primary Tumor | Metastasis

LYZ expression

Paneth cell Module Score

F
LYZ expression

Paneth cell Module Score

H

LYZ+ cancer cells with Paneth cell properties (LYZ+ cancer cells)
: LYZ > 2, Paneth cell Module Score > 0.2

	LYZ+ cancer cells	LYZ− cancer cells
Normal	33 (1%)	3540 (99%)
Tumor	5545 (13%)	37579 (87%)
Metastasis	234 (24%)	751 (76%)

In the absence of DKK2

In the presence of DKK2